

# $L$ -Functions of Hypergeometric Motives

Henri Cohen

Institut de Mathématiques de Bordeaux

July 18, 2011, Oberwolfach

# Hypergeometry I

Joint with Fernando Rodriguez-Villegas.

Hypergeometric data :

$$\alpha = (\alpha_1, \dots, \alpha_r), \quad \beta = (\beta_1, \dots, \beta_r),$$

Pochhammer symbol  $(a)_m = a(a+1)\cdots(a+m-1)$ , (generalized)  
hypergeometric function :

$$F(\alpha, \beta; t) = \sum_{m \geq 0} \frac{\prod_{1 \leq j \leq r} (\alpha_j)_m}{\prod_{1 \leq j \leq r} (\beta_j)_m} t^m$$

(classical case  $\beta_r = 1$ , so  $(\beta_r)_m = m!$ ).

## Hypergeometry II

Satisfies a linear differential equation with regular singular points at  $0$ ,  $1$ , and  $\infty$ .

Monodromy at  $1$  is a complex reflection. Characteristic polynomials of the monodromy at  $0$  and  $\infty$  :

$$P_0(T) = \prod_j (T - e^{2i\pi\beta_j}), \quad P_\infty(T) = \prod_j (T - e^{2i\pi\alpha_j}).$$

## Hypergeometry III

**Hypergeometric Assumption** : we will assume that these characteristic polynomials are products of **coprime** and **cyclotomic** polynomials. Several equivalent formulations. The simplest is :  $\alpha_j$  and  $\beta_j$  rational and

$$P_\infty(T)/P_0(T) = \prod_{\nu \geq 1} (T^\nu - 1)^{\gamma_\nu}$$

for some  $\gamma_\nu \in \mathbb{Z}$  such that  $\sum_\nu \nu \gamma_\nu = 0$ . We set  $\gamma(T) = \sum_\nu \gamma_\nu T^\nu$ .

More complicated but more concrete equivalent statement : for any integer  $A$ , let  $R(A)$  be the set of  $\phi(A)$  rational numbers  $a/A$  such that  $(a, A) = 1$  and  $0 \leq a < A$ . The assumption means that there exist integers  $A_i$  and  $B_i$  such that  $\alpha = \bigcup_j R(A_j)$  and  $\beta = \bigcup_j R(B_j)$  (concatenation and not union since the  $A_i$  or  $B_i$  are not necessarily distinct). It is clear that we may assume that the  $A_i$  are distinct from the  $B_i$ , and that  $r = \sum_j \phi(A_j) = \sum_j \phi(B_j)$ .

**Hypergeometric Assumption** : we will assume that these characteristic polynomials are products of **coprime** and **cyclotomic** polynomials. Several equivalent formulations. The simplest is :  $\alpha_j$  and  $\beta_j$  rational and

$$P_\infty(T)/P_0(T) = \prod_{\nu \geq 1} (T^\nu - 1)^{\gamma_\nu}$$

for some  $\gamma_\nu \in \mathbb{Z}$  such that  $\sum_\nu \nu \gamma_\nu = 0$ . We set  $\gamma(T) = \sum_\nu \gamma_\nu T^\nu$ .

More complicated but more concrete equivalent statement : for any integer  $A$ , let  $R(A)$  be the set of  $\phi(A)$  rational numbers  $a/A$  such that  $(a, A) = 1$  and  $0 \leq a < A$ . The assumption means that there exist integers  $A_j$  and  $B_j$  such that  $\alpha = \bigcup_j R(A_j)$  and  $\beta = \bigcup_j R(B_j)$  (concatenation and not union since the  $A_j$  or  $B_j$  are not necessarily distinct). It is clear that we may assume that the  $A_j$  are distinct from the  $B_j$ , and that  $r = \sum_j \phi(A_j) = \sum_j \phi(B_j)$ .

## Examples :

For  $r = 1$ , only possibility  $\alpha = (1/2)$ ,  $\beta = (0)$ . For  $r = 2$  exactly 13 possibilities (enumeration given below), for instance  $\alpha = (1/3, 2/3)$ ,  $\beta = (0, 0)$ .

Examples for  $r = 4$  :

$$\alpha = (1/2, 1/2, 1/3, 2/3) , \quad \beta = (1/6, 1/6, 5/6, 5/6) ,$$

or

$$\alpha = (1/5, 2/5, 3/5, 4/5) , \quad \beta = (0, 0, 0, 0) .$$

In this example,

$$P_0(T) = (T - 1)^4 , \quad P_\infty(T) = T^4 + T^3 + T^2 + T + 1 ,$$

$$P_\infty(T)/P_0(T) = (T^5 - 1)/(T - 1)^5, \text{ hence } \gamma(T) = T^5 - 5T.$$

## Associated Motives I

**Theorem** (essentially N. Katz) : If  $(\alpha, \beta)$  is hypergeometric data satisfying the fundamental hypergeometric assumption above, for each  $t \in \mathbb{P}^1(\mathbb{Q}) \setminus \{0, 1, \infty\}$  there exists a **motive**  $H(\alpha, \beta; t)$ , which is defined over  $\mathbb{Q}$ , of rank  $r$ , and pure with a certain weight  $w$ , explicitly given in terms of data.

**Conjecture** : there must therefore exist a global  $L$ -function  $\Lambda(s)$ , holomorphic sur  $\mathbb{C}$ , with functional equation  $\Lambda(w + 1 - s) = \pm \Lambda(s)$ , and with an Euler product

$$\Lambda(s) = N^{s/2} L_\infty(s) L(s) \quad \text{with} \quad L(s) = \prod_p L_p(p^{-s}) = \sum_{n \geq 1} \frac{a(n)}{n^s}, \quad \text{where}$$

$$L_p(T) = \prod_{j=1}^r (1 - \xi_j T)^{-1}, \quad \text{with} \quad |\xi_j| = p^{w/2}$$

for almost all  $p$ .

## Associated Motives I

**Theorem** (essentially N. Katz) : If  $(\alpha, \beta)$  is hypergeometric data satisfying the fundamental hypergeometric assumption above, for each  $t \in \mathbb{P}^1(\mathbb{Q}) \setminus \{0, 1, \infty\}$  there exists a **motive**  $H(\alpha, \beta; t)$ , which is defined over  $\mathbb{Q}$ , of rank  $r$ , and pure with a certain weight  $w$ , explicitly given in terms of data.

**Conjecture** : there must therefore exist a global  $L$ -function  $\Lambda(s)$ , holomorphic sur  $\mathbb{C}$ , with functional equation  $\Lambda(w + 1 - s) = \pm \Lambda(s)$ , and with an Euler product

$$\Lambda(s) = N^{s/2} L_\infty(s) L(s) \quad \text{with} \quad L(s) = \prod_p L_p(p^{-s}) = \sum_{n \geq 1} \frac{a(n)}{n^s}, \quad \text{where}$$

$$L_p(T) = \prod_{j=1}^r (1 - \xi_j T)^{-1}, \quad \text{with} \quad |\xi_j| = p^{w/2}$$

for almost all  $p$ .



## Associated Motives II

Katz's theorem gives precise recipes for the weight  $w$ , the archimedean factor  $L_\infty$ , and the factors  $L_p$  for the **good** primes  $p$ . On the other hand, not for the conductor  $N$  (of course divisible only by the bad primes), nor for the  $L_p$  for bad primes, nor for the sign  $\pm$  of the f.e. Goal of our work : numerical check of the conjecture, and deduce conjectures for the unknown quantities ( $N$ ,  $L_p$  for  $p$  bad, and  $\pm$ ).

We have treated hundreds of examples, in degree  $r = 2$ ,  $r = 4$ , and  $r = 6$ . One of the best-known is the case

$$\alpha = (1/5, 2/5, 3/5, 4/5), \quad \beta = (0, 0, 0, 0),$$

which in fact corresponds to the theory of **mirror symmetry** on the **Calabi–Yau quintic** threefold

$$x_1^5 + x_2^5 + x_3^5 + x_4^5 + x_5^5 - 5tx_1x_2x_3x_4x_5.$$

## Associated Motives II

Katz's theorem gives precise recipes for the weight  $w$ , the archimedean factor  $L_\infty$ , and the factors  $L_p$  for the **good** primes  $p$ . On the other hand, not for the conductor  $N$  (of course divisible only by the bad primes), nor for the  $L_p$  for bad primes, nor for the sign  $\pm$  of the f.e. Goal of our work : numerical check of the conjecture, and deduce conjectures for the unknown quantities ( $N$ ,  $L_p$  for  $p$  bad, and  $\pm$ ).

We have treated hundreds of examples, in degree  $r = 2$ ,  $r = 4$ , and  $r = 6$ . One of the best-known is the case

$$\alpha = (1/5, 2/5, 3/5, 4/5), \quad \beta = (0, 0, 0, 0),$$

which in fact corresponds to the theory of **mirror symmetry** on the **Calabi–Yau quintic** threefold

$$x_1^5 + x_2^5 + x_3^5 + x_4^5 + x_5^5 - 5tx_1x_2x_3x_4x_5.$$

## Details on the Motive : The function $\mathcal{L}(x)$ I

Recall that if  $P_0$  and  $P_\infty$  are the characteristic polynomials of the monodromies at  $0$  and  $\infty$  of the hypergeometric diff. eq. we set

$$P_\infty(T)/P_0(T) = \prod_{\nu \geq 1} (T^\nu - 1)^{\gamma_\nu}$$

for some  $\gamma_\nu \in \mathbb{Z}$  such that  $\sum_\nu \nu \gamma_\nu = 0$ , and  $\gamma(T) = \sum_\nu \gamma_\nu T^\nu$ .

We define

$$\mathcal{L}(x) := \sum_{\nu \geq 1} \gamma_\nu (\{ \nu x \} - 1/2),$$

where  $\{z\}$  is the fractional part of  $z$ . Then  $\mathcal{L}$  is locally constant, right continuous, periodic of period 1, and such that

$$\mathcal{L}^+(-x) = -\mathcal{L}^-(x) \quad \text{where} \quad \mathcal{L}^\pm(x) := \lim_{y \rightarrow x^\pm} \mathcal{L}(y).$$

Note that by right continuity we have  $\mathcal{L}^+(x) = \mathcal{L}(x)$ .

## Details on the Motive : The function $\mathcal{L}(x)$ I

Recall that if  $P_0$  and  $P_\infty$  are the characteristic polynomials of the monodromies at  $0$  and  $\infty$  of the hypergeometric diff. eq. we set

$$P_\infty(T)/P_0(T) = \prod_{\nu \geq 1} (T^\nu - 1)^{\gamma_\nu}$$

for some  $\gamma_\nu \in \mathbb{Z}$  such that  $\sum_\nu \nu \gamma_\nu = 0$ , and  $\gamma(T) = \sum_\nu \gamma_\nu T^\nu$ .

We define

$$\mathcal{L}(x) := \sum_{\nu \geq 1} \gamma_\nu (\{\nu x\} - 1/2),$$

where  $\{z\}$  is the fractional part of  $z$ . Then  $\mathcal{L}$  is locally constant, right continuous, periodic of period  $1$ , and such that

$$\mathcal{L}^+(-x) = -\mathcal{L}^-(x) \quad \text{where} \quad \mathcal{L}^\pm(x) := \lim_{y \rightarrow x^\pm} \mathcal{L}(y).$$

Note that by right continuity we have  $\mathcal{L}^+(x) = \mathcal{L}(x)$ .

## The function $\mathcal{L}(x)$ II

We define the **weight**  $w$  of the motive as

$$w = \max_{x \in [0,1[} \mathcal{L}(x) - \min_{x \in [0,1[} \mathcal{L}(x) - 1 .$$

It is a nonnegative integer,

$$\max_{x \in [0,1[} \mathcal{L}(x) = - \min_{x \in [0,1[} \mathcal{L}(x) = \frac{w + 1}{2} ,$$

so that  $\mathcal{L}^{\pm}(x) + (w + 1)/2$  is **integral-valued**.

Since

$$\mathcal{L}^{-}(1) = \sum_{\nu \geq 1} \gamma_{\nu}(\nu - 1/2) = -(1/2) \sum_{\nu \geq 1} \gamma_{\nu}$$

and  $\mathcal{L}^{-}(x) + (w + 1)/2$  is integral valued, it follows that

$$w + 1 \equiv \sum_{\nu \geq 1} \gamma_{\nu} \pmod{2} .$$

## The function $\mathcal{L}(x)$ II

We define the **weight**  $w$  of the motive as

$$w = \max_{x \in [0,1[} \mathcal{L}(x) - \min_{x \in [0,1[} \mathcal{L}(x) - 1 .$$

It is a nonnegative integer,

$$\max_{x \in [0,1[} \mathcal{L}(x) = - \min_{x \in [0,1[} \mathcal{L}(x) = \frac{w + 1}{2} ,$$

so that  $\mathcal{L}^{\pm}(x) + (w + 1)/2$  is **integral-valued**.

Since

$$\mathcal{L}^{-}(1) = \sum_{\nu \geq 1} \gamma_{\nu}(\nu - 1/2) = -(1/2) \sum_{\nu \geq 1} \gamma_{\nu}$$

and  $\mathcal{L}^{-}(x) + (w + 1)/2$  is integral valued, it follows that

$$w + 1 \equiv \sum_{\nu \geq 1} \gamma_{\nu} \pmod{2} .$$

# The Hodge Polynomial I

Set

$$\ell(x) := \mathcal{L}^+(x) - \mathcal{L}^-(x),$$

which measures the jumps at discontinuities. It is integral valued and in fact

$$\ell(x) = |\{i \mid \alpha_i = x\}| - |\{j \mid \beta_j = x\}|.$$

We also have  $\ell(-x) = \ell(x)$ .

We define the **Hodge polynomial**  $h(T)$  as

$$h(T) := \sum_{\ell(x) > 0} T^{\mathcal{L}^-(x) + (w+1)/2} [\ell(x)],$$

where

$$[\ell] := 1 + T + \dots + T^{\ell-1}.$$

# The Hodge Polynomial I

Set

$$\ell(x) := \mathcal{L}^+(x) - \mathcal{L}^-(x),$$

which measures the jumps at discontinuities. It is integral valued and in fact

$$\ell(x) = |\{i \mid \alpha_i = x\}| - |\{j \mid \beta_j = x\}|.$$

We also have  $\ell(-x) = \ell(x)$ .

We define the **Hodge polynomial**  $h(T)$  as

$$h(T) := \sum_{\ell(x) > 0} T^{\mathcal{L}^-(x) + (w+1)/2} [\ell(x)],$$

where

$$[\ell] := 1 + T + \cdots + T^{\ell-1}.$$



# The Hodge Polynomial II

Easy to show that :

①  $h$  is reciprocal of degree  $w$  ( $T^w h(1/T) = h(T)$ ) and has nonnegative integer coefficients.

②

$$h(T) = \sum_{\ell(x) < 0} T^{\mathcal{L}^+(x) + (w+1)/2} [-\ell(x)] .$$

**Conjecture** (Corti–Golyshév) : The Hodge numbers of the motive  $H(\alpha, \beta; t)$  are the coefficients of  $h(T)$  :

$$h(T) = \sum_{p+q=w} h^{p,q} T^p .$$

As already mentioned this gives the gamma factors of  $\Lambda(s)$  at least when  $w$  is odd (otherwise must look at action of complex conjugation on  $(w/2, w/2)$  piece) :

$$L_\infty(s) = \prod_{p < q} \Gamma_{\mathbb{C}}(s - p)^{h_{p,q}} .$$

## The Hodge Polynomial II

Easy to show that :

①  $h$  is reciprocal of degree  $w$  ( $T^w h(1/T) = h(T)$ ) and has nonnegative integer coefficients.

②

$$h(T) = \sum_{\ell(x) < 0} T^{\mathcal{L}^+(x) + (w+1)/2} [-\ell(x)] .$$

**Conjecture (Corti–Golyushev)** : The Hodge numbers of the motive  $H(\alpha, \beta; t)$  are the coefficients of  $h(T)$  :

$$h(T) = \sum_{p+q=w} h^{p,q} T^p .$$

As already mentioned this gives the gamma factors of  $\Lambda(s)$  at least when  $w$  is odd (otherwise must look at action of complex conjugation on  $(w/2, w/2)$  piece) :

$$L_\infty(s) = \prod_{p < q} \Gamma_{\mathbb{C}}(s - p)^{h_{p,q}} .$$

## The Hodge Polynomial II

Easy to show that :

①  $h$  is reciprocal of degree  $w$  ( $T^w h(1/T) = h(T)$ ) and has nonnegative integer coefficients.

②

$$h(T) = \sum_{\ell(x) < 0} T^{\mathcal{L}^+(x) + (w+1)/2} [-\ell(x)] .$$

**Conjecture (Corti–Golyshev)** : The Hodge numbers of the motive  $H(\alpha, \beta; t)$  are the coefficients of  $h(T)$  :

$$h(T) = \sum_{p+q=w} h^{p,q} T^p .$$

As already mentioned this gives the gamma factors of  $\Lambda(s)$  at least when  $w$  is odd (otherwise must look at action of complex conjugation on  $(w/2, w/2)$  piece) :

$$L_\infty(s) = \prod_{p < q} \Gamma_{\mathbb{C}}(s - p)^{h_{p,q}} .$$

## Recipe for the $L_p$ for good primes $p$ I

We will define below the trace of Frobenius  $a_q(t) = \text{Tr}(\text{Frob}_q)$ , and we will then set as usual

$$L_p(T) = \exp\left(\sum_{f \geq 1} \frac{a_{p^f}(t)}{f} T^f\right).$$

The fundamental quantity which occurs in the definition of  $a_q(t)$  is the following : for any multiplicative character  $\chi$  of  $\mathbb{F}_q^*$  we set

$$Q(\chi) = \prod_{\nu} g(\chi^{\nu})^{\gamma_{\nu}},$$

where  $\gamma_{\nu}$  is as above and  $g(\chi^{\nu})$  is the corresponding **Gauss sum**. Using Möbius inversion and elementary formulas for Gauss sums, for suitable easily computed integers  $a$  and  $b$  we also have

$$Q(\chi) = (-1)^a q^b \frac{\prod_i \prod_{d|A_i} g(\chi^{-d})^{\mu(A_i/d)}}{\prod_i \prod_{d|B_i} g(\chi^{-d})^{\mu(B_i/d)}},$$

where the  $A_i$  and  $B_i$  are as above.

## Recipe for the $L_p$ for good primes $p \nmid l$

The **hypergeometric assumptions** imply that  $Q(\chi)$  can be expressed only in terms of **Jacobi** sums (which belong to a smaller number field), see below.

Finally, for  $t \neq 0, 1$ , or  $\infty$  we set

$$a_q(t) = \frac{q^d}{1-q} \left( 1 + \sum_{\chi \neq \chi_0} \chi(Mt) Q(\chi) \right),$$

where  $\chi$  ranges over all nontrivial characters of  $\mathbb{F}_q$ , for certain constants  $d$  and  $M$  which I do not define here (of course  $M$  is only a normalizing factor, could change  $Mt$  into  $t$ , but less clean). By Galois theory this is in  $\mathbb{Q}$ , and it is easy to show that it is in fact in  $\mathbb{Z}$ , as it should if it is indeed a trace.

## Recipe for the $L_p$ for good primes $p \nmid l$

The **hypergeometric assumptions** imply that  $Q(\chi)$  can be expressed only in terms of **Jacobi** sums (which belong to a smaller number field), see below.

Finally, for  $t \neq 0, 1$ , or  $\infty$  we set

$$a_q(t) = \frac{q^d}{1-q} \left( 1 + \sum_{\chi \neq \chi_0} \chi(Mt) Q(\chi) \right),$$

where  $\chi$  ranges over all nontrivial characters of  $\mathbb{F}_q$ , for certain constants  $d$  and  $M$  which I do not define here (of course  $M$  is only a normalizing factor, could change  $Mt$  into  $t$ , but less clean). By Galois theory this is in  $\mathbb{Q}$ , and it is easy to show that it is in fact in  $\mathbb{Z}$ , as it should if it is indeed a trace.

## Computation of the $L_p$ for good primes $p \mid$

In the case  $(\alpha, \beta) = (1/5, 2/5, 3/5, 4/5), (0, 0, 0, 0)$  mentioned above, we have the precise formula :

$$a_q(t) = \frac{1}{1-q} \left( 1 + \sum_{\chi \neq \chi_0} \chi(5^5 t) J(\chi, \chi, \chi, \chi) \right),$$

where  $q = p^f$  and  $J$  is the generalized **Jacobi sum** :

$$J(\chi_1, \dots, \chi_r) = \sum_{x_1 + \dots + x_r = 1} \chi_1(x_1) \cdots \chi_r(x_r).$$

**Elementary idea 1** : the local factors  $L_p$  have degree 4 and are Weil-symmetrical. Therefore only need to compute  $a_p$  and  $a_{p^2}$ .

**Elementary idea 2** : To obtain  $B$  terms of the Dirichlet series, need of course  $a_p$  for  $p \leq B$ , but  $a_{p^2}$  only for  $p \leq B^{1/2}$ .

## Computation of the $L_p$ for good primes $p \mid$

In the case  $(\alpha, \beta) = (1/5, 2/5, 3/5, 4/5), (0, 0, 0, 0)$  mentioned above, we have the precise formula :

$$a_q(t) = \frac{1}{1-q} \left( 1 + \sum_{\chi \neq \chi_0} \chi(5^5 t) J(\chi, \chi, \chi, \chi) \right),$$

where  $q = p^f$  and  $J$  is the generalized **Jacobi sum** :

$$J(\chi_1, \dots, \chi_r) = \sum_{x_1 + \dots + x_r = 1} \chi_1(x_1) \cdots \chi_r(x_r).$$

**Elementary idea 1** : the local factors  $L_p$  have degree 4 and are Weil-symmetrical. Therefore only need to compute  $a_p$  and  $a_{p^2}$ .

**Elementary idea 2** : To obtain  $B$  terms of the Dirichlet series, need of course  $a_p$  for  $p \leq B$ , but  $a_{p^2}$  only for  $p \leq B^{1/2}$ .



## Computation of the $L_p$ for good primes $p$ I

In the case  $(\alpha, \beta) = (1/5, 2/5, 3/5, 4/5), (0, 0, 0, 0)$  mentioned above, we have the precise formula :

$$a_q(t) = \frac{1}{1-q} \left( 1 + \sum_{\chi \neq \chi_0} \chi(5^5 t) J(\chi, \chi, \chi, \chi) \right),$$

where  $q = p^f$  and  $J$  is the generalized **Jacobi sum** :

$$J(\chi_1, \dots, \chi_r) = \sum_{x_1 + \dots + x_r = 1} \chi_1(x_1) \cdots \chi_r(x_r).$$

**Elementary idea 1** : the local factors  $L_p$  have degree 4 and are Weil-symmetrical. Therefore only need to compute  $a_p$  and  $a_{p^2}$ .

**Elementary idea 2** : To obtain  $B$  terms of the Dirichlet series, need of course  $a_p$  for  $p \leq B$ , but  $a_{p^2}$  only for  $p \leq B^{1/2}$ .

## Computation of the $L_p$ for good primes $p$ II

Main computational task is the computation of the Jacobi sums.

### (1) : Direct method.

By elementary properties of Jacobi sums, an order  $r$  Jacobi sum as above can be expressed as a product of  $r - 1$  usual Jacobi sums of order 2, of the form :

$$J(\chi_1, \chi_2) = \sum_{x \in \mathbb{F}_q} \chi_1(x) \chi_2(1 - x).$$

The direct computation of these sums requires essentially  $O(q)$  operations, neglecting the computation of the  $\chi_i$  values, and since we need  $q - 2$  Jacobi sums, the total cost of the computation of  $a_q(t)$  is of the order of  $O((r - 1)q^2)$ .

## (2). The use of Gauss sums.

Recall that

$$g(\chi) = \sum_{x \in \mathbb{F}_q^*} \chi(x) \exp(2i\pi \operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(x)/p).$$

It is well-known that

$$J(\chi_1, \dots, \chi_r) = \frac{\prod_j g(\chi_j)}{g(\prod_j \chi_j)}$$

when all the characters are nontrivial (and simpler formulas otherwise). A priori, why use Gauss sums, more complicated than Jacobi sums, and belong to a larger number field than Jacobi sums? A posteriori, they are in fact useful.

## Computation of the $L_p$ for good primes $p$ IV

Let  $\omega$  be a generator of the cyclic group  $\widehat{\mathbb{F}_q^*}$ . Taking our favorite example, we thus have

$$a_q(t) = \frac{1}{1-q} \left( 1 + \sum_{1 \leq r \leq q-2} \omega^r(5^5 t) J(\omega^r, \omega^r, \omega^r, \omega^r) \right).$$

If we compute once and for all the  $g(\omega^r)$  for  $1 \leq r \leq q-2$ , the above computation requires time  $O(q)$ , for a total time of  $O(q^2)$ , hence gain of a factor  $r-1$ .

### (3). Use of $\Theta$ functions.

Idea of S. Louboutin : when  $q = p$  is prime (by far the largest part of the computation), Gauss sums are directly linked to the **root numbers** in the functional equation of  $L$ -functions, but even better, of  $\Theta$ -functions, associated to the characters. These can be computed in time  $O(q^{1/2+\epsilon})$ , so a considerable gain : combined with the preceding ideal, total cost  $O(q^{3/2+\epsilon})$ .

## Computation of the $L_p$ for good primes $p$ IV

Let  $\omega$  be a generator of the cyclic group  $\widehat{\mathbb{F}}_q^*$ . Taking our favorite example, we thus have

$$a_q(t) = \frac{1}{1-q} \left( 1 + \sum_{1 \leq r \leq q-2} \omega^r(5^5 t) J(\omega^r, \omega^r, \omega^r, \omega^r) \right).$$

If we compute once and for all the  $g(\omega^r)$  for  $1 \leq r \leq q-2$ , the above computation requires time  $O(q)$ , for a total time of  $O(q^2)$ , hence gain of a factor  $r-1$ .

### (3). Use of $\Theta$ functions.

Idea of S. Louboutin : when  $q = p$  is prime (by far the largest part of the computation), Gauss sums are directly linked to the **root numbers** in the functional equation of  $L$ -functions, but even better, of  $\Theta$ -functions, associated to the characters. These can be computed in time  $O(q^{1/2+\epsilon})$ , so a considerable gain : combined with the preceding ideal, total cost  $O(q^{3/2+\epsilon})$ .

## Computation of the $L_p$ for good primes $p \nmid V$

### (4). Use of $p$ -adic methods.

The marvelous formula of **Gross–Koblitz** allows us to express Gauss sums (for all  $q$ , not only for  $q = p$ ) in terms of the **Morita  $p$ -adic gamma function**  $\Gamma_p$ . There exist efficient methods to compute this function, time  $O(p^{1+\epsilon})$ , so in principle no more efficient, even less than using theta functions.

The main interest of the method is that we only need to compute values modulo  $p$  or  $p^2$ , since we know that  $a_q(t)$  is an **integer** and that we have Weil–Deligne bounds. Although this is a  $O(q^2)$  method, it is the best available when  $q = p^2$ , and even when  $q = p$ , since we can work mod  $p$  and the implicit constant of  $O()$  is **very small**, it is quite competitive in practice ( $p \leq 10^4$  for instance).

### (4). Use of $p$ -adic methods.

The marvelous formula of **Gross–Koblitz** allows us to express Gauss sums (for all  $q$ , not only for  $q = p$ ) in terms of the **Morita  $p$ -adic gamma function**  $\Gamma_p$ . There exist efficient methods to compute this function, time  $O(p^{1+\epsilon})$ , so in principle no more efficient, even less than using theta functions.

The main interest of the method is that we only need to compute values modulo  $p$  or  $p^2$ , since we know that  $a_q(t)$  is an **integer** and that we have Weil–Deligne bounds. Although this is a  $O(q^2)$  method, it is the best available when  $q = p^2$ , and even when  $q = p$ , since we can work mod  $p$  and the implicit constant of  $O()$  is **very small**, it is quite competitive in practice ( $p \leq 10^4$  for instance).

## Verification of the Functional Equation I

(Also Implemented in several computer packages such as **T. Dokschitser** and `lcalc` of **M. Rubinstein**).

Even assuming that we know  $N$  and  $L_p$  for  $p$  bad, to check the functional equation  $\Lambda(w+1-s) = \pm \Lambda(s)$  need to compute  $\Lambda(s)$ . For this, efficient general recipes :

Let  $K(x)$  be the inverse Mellin transform of  $L_\infty(s)$ , i.e.,

$$\int_0^\infty t^{s-1} K(t) dt = L_\infty(s) ,$$

and set

$$\gamma(s, x) = \int_x^\infty t^{s-1} K(t) dt .$$

For any  $t_0 > 0$  we have

$$L_\infty(s)L(s) = \sum_{n \geq 1} \frac{a(n)}{n^s} \gamma(s, nt_0/N^{1/2}) \pm \sum_{n \geq 1} \frac{a(n)}{n^{w+1-s}} \gamma(w+1-s, n/(t_0 N^{1/2})) .$$

This series converges **exponentially fast**, but slows down if  $N$  is large.



## Verification of the Functional Equation II

The “method” is thus as follows : we guess values for  $N$  and the bad Euler factors  $L_p$  (a lot of information is known about them), and we test if the above expression is independent of  $t_0$ , by choosing for instance  $t_0 = 1$  and  $t_0 = 1.1$ .

## Verification of the Functional Equation III

In the above computation, must do **three** costly computations. First, compute the inverse Mellin transform  $K(x)$  (see below). Second, compute the partial Mellin transforms  $\int_x^\infty t^{s-1} K(t) dt$ . Finally, the third consists in computing the two series of the form.

$$\sum_{n \geq 1} (a(n)/n^s) \gamma(s, nt_0/N^{1/2}).$$

**Elementary idea 3** : to check the functional equation, there is no need to come back to the Dirichlet series by computing  $\int_x^\infty t^{s-1} K(t)$ . Indeed, we can write

$$\Lambda(s) = \int_0^\infty t^{s-1} F(N^{1/2}t) dt ,$$

so the f.eq. is in our case equivalent to

$$F(1/t) = \pm t^{w+1} F(t) .$$

Thus we only check this, and so only need to compute  $K(x)$  and series sums.

## Verification of the Functional Equation III

In the above computation, must do **three** costly computations. First, compute the inverse Mellin transform  $K(x)$  (see below). Second, compute the partial Mellin transforms  $\int_x^\infty t^{s-1} K(t) dt$ . Finally, the third consists in computing the two series of the form.

$$\sum_{n \geq 1} (a(n)/n^s) \gamma(s, nt_0/N^{1/2}).$$

**Elementary idea 3** : to check the functional equation, there is no need to come back to the Dirichlet series by computing  $\int_x^\infty t^{s-1} K(t)$ . Indeed, we can write

$$\Lambda(s) = \int_0^\infty t^{s-1} F(N^{1/2}t) dt ,$$

so the f.eq. is in our case equivalent to

$$F(1/t) = \pm t^{w+1} F(t) .$$

Thus we only check this, and so only need to compute  $K(x)$  and series sums.

## Verification of the Functional Equation IV

The archimedean factor  $L_\infty(s)$  is up to exponential factors  $Q^s$ , of the form  $\Gamma(s)$ ,  $\Gamma(s)\Gamma(s-1)$ ,  $\Gamma(s)\Gamma(s-1)\Gamma(s-2)$  for instance (can of course also have  $\Gamma(s/2)$ , etc...) The precise recipe for this factor in a very general context is due to Serre around 1970. It depends on the **Hodge numbers**  $h^{p,q}$  of the motive, which can easily be computed from the hypergeometric data, as can the weight  $w$ . More precisely : if as usual  $\Gamma_{\mathbb{C}}(s) = (2\pi)^{-s}\Gamma(s)$  and  $\Gamma_{\mathbb{R}}(s) = \pi^{-s/2}\Gamma(s/2)$  then

$$L_\infty(s) = N^{s/2} \prod_{\substack{p < q \\ p+q=w}} \Gamma_{\mathbb{C}}(s-p)^{h_{p,q}} \prod_{p=q=w/2} \Gamma_{\mathbb{R}}(s-p)^{h_{p,p}^+} \Gamma_{\mathbb{R}}(s-p+1)^{h_{p,p}^-},$$

where the second product of course occurs only in even weight and  $h_{p,p}^\pm$  are the dimensions of the  $\pm$ -eigenspaces of complex conjugation on  $H^{p,p}$  of the motive.

## Verification of the Functional Equation V

The computation of the inverse Mellin transform of the archimedean factor is a fundamental problem for which many solutions exist, and of course have been implemented.

Recall that the inverse Mellin transform is given by

$$K(x) = \frac{1}{2i\pi} \int_{\sigma-i\infty}^{\sigma+i\infty} \gamma(s)x^{-s} ds ,$$

for  $\sigma$  sufficiently large. We move the line of integration to the left until  $-\infty$ , and we catch all the residues along the way. This allows us to express  $K(x)$  as a power series, together with finitely many logarithmic terms or negative powers of  $x$ .

## Verification of the Functional Equation VI

It is easy to show that these series converge like the series

$$\sum_{n \geq 0} (-1)^n \frac{x^n}{(n!)^{r/2}},$$

where  $r$  is the degree of the motive, in other words the number of  $\alpha$  or of  $\beta$ , or equivalently the number of  $\Gamma(s/2)$  factors, recalling that  $\Gamma(s-p)$  counts for 2.

In the case  $r = 2$ , the factor is  $\Gamma(s)$  whose inverse Mellin transform is  $e^{-x}$ , nothing to add. For  $r = 4$ , the inverse Mellin transform is essentially the well-known  $K$ -Bessel function  $K_1(x)$ . For  $r = 6$  the convergence is in  $1/n!^3$ .

Although some cancellation exists, it is no problem to use the above series even when  $x$  is large, for  $r \geq 2$ , and even better for  $r \geq 4$ . Otherwise, there exist asymptotic expansions for  $x \rightarrow \infty$ .

## Verification of the Functional Equation VI

It is easy to show that these series converge like the series

$$\sum_{n \geq 0} (-1)^n \frac{x^n}{(n!)^{r/2}},$$

where  $r$  is the degree of the motive, in other words the number of  $\alpha$  or of  $\beta$ , or equivalently the number of  $\Gamma(s/2)$  factors, recalling that  $\Gamma(s - p)$  counts for 2.

In the case  $r = 2$ , the factor is  $\Gamma(s)$  whose inverse Mellin transform is  $e^{-x}$ , nothing to add. For  $r = 4$ , the inverse Mellin transform is essentially the well-known  $K$ -Bessel function  $K_1(x)$ . For  $r = 6$  the convergence is in  $1/n!^3$ .

Although some cancellation exists, it is no problem to use the above series even when  $x$  is large, for  $r \geq 2$ , and even better for  $r \geq 4$ . Otherwise, there exist asymptotic expansions for  $x \rightarrow \infty$ .

## Conjectures I

Three types of prime numbers :

- (1) **Wild** : those which divide the denominator of one of the  $\alpha_j$  or  $\beta_j$ .
- (2) **Tame** : not wild, and such that either  $v_p(t) > 0$ ,  $v_p(1/t) > 0$ , or  $v_p(t-1) > 0$ .
- (3) **Good** : all the other primes.

We have given the recipe for  $L_p(T)$  when  $p$  is good, and in that case  $p \nmid N$ , the conductor. When  $p$  is tame, we have been able to obtain precise (but complicated) conjectures, both for  $L_p(T)$  and for  $v_p(N)$ . They should not be too difficult to prove.



## Conjectures I

Three types of prime numbers :

(1) **Wild** : those which divide the denominator of one of the  $\alpha_j$  or  $\beta_j$ .

(2) **Tame** : not wild, and such that either  $v_p(t) > 0$ ,  $v_p(1/t) > 0$ , or  $v_p(t-1) > 0$ .

(3) **Good** : all the other primes.

We have given the recipe for  $L_p(T)$  when  $p$  is good, and in that case  $p \nmid N$ , the conductor. When  $p$  is tame, we have been able to obtain precise (but complicated) conjectures, both for  $L_p(T)$  and for  $v_p(N)$ . They should not be too difficult to prove.

## Conjectures II

When  $p$  is **wild**, a lot of experimental data and conjectures, very little precise conjectures. Sample phenomenon, only partly understood : if  $p$  is wild and  $v_p(t)$  tends to  $-\infty$ , then  $v_p(N)$  has a behavior of the form (almost random numbers) :

12, 8, 9, 6, 5, 6, 4, 3, 0, 2, 2, 0, 2, 2, 0, 2, 2, 0

for  $v_p(t) = -1, -2, -3, \dots$ . In other words it starts to “decrease”, with possible local increases, then **it reaches 0** for a certain **critical value** of  $v_p(t)$ , and then it is periodic with small amplitude. We have a precise conjecture for the critical value.

Example in dimension 0 : the **Belyi polynomials** which occur for suitable hypergeometric data : let  $a \geq 2$  and  $b \geq 2$  be coprime (not absolutely necessary) and let

$$P_t(X) = X^a(1 - X)^b - \frac{a^a b^b}{(a + b)^{a+b} t}.$$

Then the **discriminant**  $N$  of the number field generated by  $P_t$  has similar behavior for  $p \mid ab(a + b)$  (i.e., wild).

## Conjectures II

When  $p$  is **wild**, a lot of experimental data and conjectures, very little precise conjectures. Sample phenomenon, only partly understood : if  $p$  is wild and  $v_p(t)$  tends to  $-\infty$ , then  $v_p(N)$  has a behavior of the form (almost random numbers) :

12, 8, 9, 6, 5, 6, 4, 3, 0, 2, 2, 0, 2, 2, 0, 2, 2, 0

for  $v_p(t) = -1, -2, -3, \dots$ . In other words it starts to “decrease”, with possible local increases, then **it reaches 0** for a certain **critical value** of  $v_p(t)$ , and then it is periodic with small amplitude. We have a precise conjecture for the critical value.

Example in dimension 0 : the **Belyi polynomials** which occur for suitable hypergeometric data : let  $a \geq 2$  and  $b \geq 2$  be coprime (not absolutely necessary) and let

$$P_t(X) = X^a(1 - X)^b - \frac{a^a b^b}{(a + b)^{a+b} t}.$$

Then the **discriminant**  $N$  of the number field generated by  $P_t$  has similar behavior for  $p \mid ab(a + b)$  (i.e., wild).

## Examples

We give examples for  $r = 2$ . As mentioned, there are exactly 13 pairs  $(\alpha, \beta)$  of possible hypergeometric data. Three have motivic weight 0 (hence functional equation  $s \mapsto 1 - s$ ), corresponding to three families of number fields. The other ten have motivic weight 1 (func. eq.  $s \mapsto 2 - s$ ), and correspond to families of elliptic curves, hence by Wiles to families of modular cusp forms of weight 2.

Even though very explicit and general, we do **not** know how to prove modularity of these ten families without Wiles, and a dream is that perhaps it could be possible, since the data is purely of a combinatorial (as opposed to geometric) nature.

## Case (1)

$$\alpha = \{1/2, 1/2\}, \quad \beta = (0, 0).$$

$$a_p(t) = \frac{1}{1-p} \left( 1 + \sum_{1 \leq r \leq p-2} J(r, r)^2 \omega^r(16t) \right).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - 3 - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $x(x-1)y(y-1) = a$ , with  $a = 1/(16t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 + XY = X(X-a)^2.$$

## Case (1)

$$\alpha = \{1/2, 1/2\}, \quad \beta = (0, 0).$$

$$a_p(t) = \frac{1}{1-p} \left( 1 + \sum_{1 \leq r \leq p-2} J(r, r)^2 \omega^r(16t) \right).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - 3 - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $x(x-1)y(y-1) = a$ , with  $a = 1/(16t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 + XY = X(X-a)^2.$$

## Case (1)

$$\alpha = \{1/2, 1/2\}, \quad \beta = (0, 0).$$

$$a_p(t) = \frac{1}{1-p} \left( 1 + \sum_{1 \leq r \leq p-2} J(r, r)^2 \omega^r(16t) \right).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - 3 - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $x(x-1)y(y-1) = a$ , with  $a = 1/(16t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 + XY = X(X-a)^2.$$

## Case (2)

$$\alpha = \{1/3, 2/3\}, \quad \beta = (0, 0).$$

$$a_p(t) = \frac{1}{1-p} \left( 1 + \sum_{1 \leq r \leq p-2} J(r, r, r) \omega^r(27t) \right).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - 2 - N_p(t),$$

$N_p(t)$  number of affine  $\mathbb{F}_p$ -points on  $xy(1-x-y) = a$ , with  $a = 1/(27t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of projective points on

$$Y^2 + XY + aY = X^3.$$



## Case (2)

$$\alpha = \{1/3, 2/3\}, \quad \beta = (0, 0).$$

$$a_p(t) = \frac{1}{1-p} \left( 1 + \sum_{1 \leq r \leq p-2} J(r, r, r) \omega^r(27t) \right).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - 2 - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $xy(1-x-y) = a$ , with  $a = 1/(27t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 + XY + aY = X^3.$$

## Case (2)

$$\alpha = \{1/3, 2/3\}, \quad \beta = (0, 0).$$

$$a_p(t) = \frac{1}{1-p} \left( 1 + \sum_{1 \leq r \leq p-2} J(r, r, r) \omega^r(27t) \right).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - 2 - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $xy(1-x-y) = a$ , with  $a = 1/(27t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 + XY + aY = X^3.$$

### Case (3)

$$\alpha = \{1/4, 3/4\}, \quad \beta = (0, 0).$$

$$a_p(t) = \frac{1}{1-p} \left( 1 + \left(\frac{-t}{p}\right)p + \sum_{1 \leq r \leq p-2} J(r, r, 2r) \omega^r(64t) \right).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - 2 - \left(\frac{-t}{p}\right) - N_p(t),$$

$N_p(t)$  number of affine  $\mathbb{F}_p$ -points on  $x^2y(1-x-y) = a$ , with  $a = 1/(64t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of projective points on

$$Y^2 + XY = X^3 + aX; .$$

### Case (3)

$$\alpha = \{1/4, 3/4\}, \quad \beta = (0, 0).$$

$$a_p(t) = \frac{1}{1-p} \left( 1 + \left(\frac{-t}{p}\right)p + \sum_{1 \leq r \leq p-2} J(r, r, 2r)\omega^r(64t) \right).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - 2 - \left(\frac{-t}{p}\right) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $x^2y(1-x-y) = a$ , with  $a = 1/(64t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 + XY = X^3 + aX; .$$

### Case (3)

$$\alpha = \{1/4, 3/4\}, \quad \beta = (0, 0).$$

$$a_p(t) = \frac{1}{1-p} \left( 1 + \left(\frac{-t}{p}\right)p + \sum_{1 \leq r \leq p-2} J(r, r, 2r)\omega^r(64t) \right).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - 2 - \left(\frac{-t}{p}\right) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $x^2y(1-x-y) = a$ , with  $a = 1/(64t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 + XY = X^3 + aX; .$$

## Case (4)

$$\alpha = \{1/6, 5/6\}, \quad \beta = (0, 0).$$

$$a_p(t) = \frac{1}{1-p} \left( 1 + \left( \left( \frac{-3t}{p} \right) + \left( \frac{2t}{p} \right)_3 + \left( \frac{2t}{p} \right)_3^{-1} \right) p \right. \\ \left. + \sum_{1 \leq r \leq p-2} J(r, 2r, 3r) \omega^r(432t) \right).$$

If  $v_p(t) = 0$ :

$$a_p(t) = p - 2 - \left( \left( \frac{-3t}{p} \right) + \left( \frac{2t}{p} \right)_3 + \left( \frac{2t}{p} \right)_3^{-1} \right) - N_p(t),$$

$N_p(t)$  number of affine  $\mathbb{F}_p$ -points on  $x^3 y^2 (1 - x - y) = a$ , with  $a = 1/(432t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of projective points on

$$Y^2 + XY = X^3 - a.$$

## Case (4)

$$\alpha = \{1/6, 5/6\}, \quad \beta = (0, 0).$$

$$a_p(t) = \frac{1}{1-p} \left( 1 + \left( \left( \frac{-3t}{p} \right) + \left( \frac{2t}{p} \right)_3 + \left( \frac{2t}{p} \right)_3^{-1} \right) p \right. \\ \left. + \sum_{1 \leq r \leq p-2} J(r, 2r, 3r) \omega^r(432t) \right).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - 2 - \left( \left( \frac{-3t}{p} \right) + \left( \frac{2t}{p} \right)_3 + \left( \frac{2t}{p} \right)_3^{-1} \right) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $x^3 y^2 (1 - x - y) = a$ , with  $a = 1/(432t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 + XY = X^3 - a.$$

## Case (4)

$$\alpha = \{1/6, 5/6\}, \quad \beta = (0, 0).$$

$$a_p(t) = \frac{1}{1-p} \left( 1 + \left( \left( \frac{-3t}{p} \right) + \left( \frac{2t}{p} \right)_3 + \left( \frac{2t}{p} \right)_3^{-1} \right) p \right. \\ \left. + \sum_{1 \leq r \leq p-2} J(r, 2r, 3r) \omega^r(432t) \right).$$

If  $v_p(t) = 0$ :

$$a_p(t) = p - 2 - \left( \left( \frac{-3t}{p} \right) + \left( \frac{2t}{p} \right)_3 + \left( \frac{2t}{p} \right)_3^{-1} \right) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $x^3 y^2 (1 - x - y) = a$ , with  $a = 1/(432t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 + XY = X^3 - a.$$



## Case (8)

$$\alpha = \{1/2, 1/2\}, \quad \beta = (1/3, 2/3).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - \left( \left( \frac{-3t}{p} \right) + \left( \frac{-3t}{p} \right)_4 + \left( \frac{-3t}{p} \right)_4^{-1} \right) - N_p(t),$$

$N_p(t)$  number of affine  $\mathbb{F}_p$ -points on  $y^2(1-y)^2 = x^3(1-x)/a$ , with  $a = 27/(16t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of projective points on

$$Y^2 - aY = X^3 - aX^2.$$

## Case (8)

$$\alpha = \{1/2, 1/2\}, \quad \beta = (1/3, 2/3).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - \left( \left( \frac{-3t}{p} \right) + \left( \frac{-3t}{p} \right)_4 + \left( \frac{-3t}{p} \right)_4^{-1} \right) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $y^2(1-y)^2 = x^3(1-x)/a$ , with  $a = 27/(16t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 - aY = X^3 - aX^2.$$

## Case (8)

$$\alpha = \{1/2, 1/2\}, \quad \beta = (1/3, 2/3).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - \left( \binom{-3t}{p} + \binom{-3t}{p}_4 + \binom{-3t}{p}_4^{-1} \right) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $y^2(1-y)^2 = x^3(1-x)/a$ , with  $a = 27/(16t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 - aY = X^3 - aX^2.$$

## Case (9)

$$\alpha = \{1/2, 1/2\}, \quad \beta = (1/4, 3/4).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - \left(\frac{-t}{p}\right) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $y^2(1-y)^2 = x(1-x)/a$ , with  $a = 4/t$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 = X^3 - aX^2 + aX.$$

## Case (9)

$$\alpha = \{1/2, 1/2\}, \quad \beta = (1/4, 3/4).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - \left(\frac{-t}{p}\right) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $y^2(1-y)^2 = x(1-x)/a$ , with  $a = 4/t$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 = X^3 - aX^2 + aX.$$

## Case (9)

$$\alpha = \{1/2, 1/2\}, \quad \beta = (1/4, 3/4).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - \left(\frac{-t}{p}\right) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $y^2(1-y)^2 = x(1-x)/a$ , with  $a = 4/t$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 = X^3 - aX^2 + aX.$$

## Case (10)

$$\alpha = \{1/2, 1/2\}, \quad \beta = (1/6, 5/6).$$

If  $v_p(t) = 0$  :

$$pa_p(t) = p(p^2 + 6p - 12) - \left( \left( \frac{-3t}{p} \right) + \left( \frac{t}{p} \right)_3 J(\rho_3, \rho_3)^2 + \left( \frac{t}{p} \right)_3^{-1} J(\rho_3^{-1}, \rho_3^{-1})^2 \right) -$$

$N_p(t)$  number of affine  $\mathbb{F}_p$ -points on the threefold :  
 $x^3 y^2 (1 - x - y) = z(1 - z)w(1 - w)/a$ , with  $a = 27/t$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of projective points on

$$Y^2 = X^3 - (a/4)(X - 1)^2.$$

## Case (10)

$$\alpha = \{1/2, 1/2\}, \quad \beta = (1/6, 5/6).$$

If  $v_p(t) = 0$  :

$$pa_p(t) = p(p^2 + 6p - 12) - \left( \left( \frac{-3t}{p} \right) + \left( \frac{t}{p} \right)_3 J(\rho_3, \rho_3)^2 + \left( \frac{t}{p} \right)_3^{-1} J(\rho_3^{-1}, \rho_3^{-1})^2 \right) -$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on the **threefold** :  
 $x^3 y^2 (1 - x - y) = z(1 - z)w(1 - w)/a$ , with  $a = 27/t$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 = X^3 - (a/4)(X - 1)^2.$$



## Case (10)

$$\alpha = \{1/2, 1/2\}, \quad \beta = (1/6, 5/6).$$

If  $v_p(t) = 0$  :

$$pa_p(t) = p(p^2 + 6p - 12) - \left( \left( \frac{-3t}{p} \right) + \left( \frac{t}{p} \right)_3 J(\rho_3, \rho_3)^2 + \left( \frac{t}{p} \right)_3^{-1} J(\rho_3^{-1}, \rho_3^{-1})^2 \right) -$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on the **threefold** :  
 $x^3 y^2 (1 - x - y) = z(1 - z)w(1 - w)/a$ , with  $a = 27/t$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 = X^3 - (a/4)(X - 1)^2.$$

## Case (11)

$$\alpha = \{1/3, 2/3\}, \quad \beta = (1/4, 3/4).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - \left(\frac{3t}{p}\right) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $y^3(1-y) = x(1-x)/a$ , with  $a = 64/(27t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 - aY = X^3 - aX.$$

## Case (11)

$$\alpha = \{1/3, 2/3\}, \quad \beta = (1/4, 3/4).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - \left(\frac{3t}{p}\right) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $y^3(1-y) = x(1-x)/a$ , with  $a = 64/(27t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 - aY = X^3 - aX.$$

## Case (11)

$$\alpha = \{1/3, 2/3\}, \quad \beta = (1/4, 3/4).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - \left(\frac{3t}{p}\right) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $y^3(1-y) = x(1-x)/a$ , with  $a = 64/(27t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 - aY = X^3 - aX.$$

## Case (12)

$$\alpha = \{1/3, 2/3\}, \quad \beta = (1/6, 5/6).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - \left(\frac{t}{p}\right) + \left(\frac{4t}{p}\right)_3 J(\rho_3, \rho_3) + \left(\frac{4t}{p}\right)_3^{-1} J(\rho_3^{-1}, \rho_3^{-1}) - N_p(t),$$

$N_p(t)$  number of affine  $\mathbb{F}_p$ -points on the genus 2 curve :  
 $y^3(1-y)^3 = x(1-x)/a$ , with  $a = 16/t$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of projective points on

$$Y^2 = X^3 - 3aX + 4a + a^2/4.$$

## Case (12)

$$\alpha = \{1/3, 2/3\}, \quad \beta = (1/6, 5/6).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - \left(\frac{t}{p}\right) + \left(\frac{4t}{p}\right)_3 J(\rho_3, \rho_3) + \left(\frac{4t}{p}\right)_3^{-1} J(\rho_3^{-1}, \rho_3^{-1}) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on the **genus 2 curve** :  
 $y^3(1-y)^3 = x(1-x)/a$ , with  $a = 16/t$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 = X^3 - 3aX + 4a + a^2/4.$$

## Case (12)

$$\alpha = \{1/3, 2/3\}, \quad \beta = (1/6, 5/6).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - \left(\frac{t}{p}\right) + \left(\frac{4t}{p}\right)_3 J(\rho_3, \rho_3) + \left(\frac{4t}{p}\right)_3^{-1} J(\rho_3^{-1}, \rho_3^{-1}) - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on the **genus 2 curve** :  
 $y^3(1-y)^3 = x(1-x)/a$ , with  $a = 16/t$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 = X^3 - 3aX + 4a + a^2/4.$$

## Case (13)

$$\alpha = \{1/4, 3/4\}, \quad \beta = (1/6, 5/6) .$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - 2 \left( \frac{3t}{p} \right) - \left( \frac{4t}{p} \right)_3 - \left( \frac{4t}{p} \right)_3^{-1} - N_p(t) ,$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $y^4(1-y)^2 = x^2(1-x)/a$ , with  $a = 27/(4t)$ .

$$a_p(t) = p + 1 - M_p(t) ,$$

$M_p(t)$  number of **projective** points on

$$Y^2 = X^3 - aX + a .$$



## Case (13)

$$\alpha = \{1/4, 3/4\}, \quad \beta = (1/6, 5/6).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - 2\left(\frac{3t}{p}\right) - \left(\frac{4t}{p}\right)_3 - \left(\frac{4t}{p}\right)_3^{-1} - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $y^4(1-y)^2 = x^2(1-x)/a$ , with  $a = 27/(4t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 = X^3 - aX + a.$$

## Case (13)

$$\alpha = \{1/4, 3/4\}, \quad \beta = (1/6, 5/6).$$

If  $v_p(t) = 0$  :

$$a_p(t) = p - 2\left(\frac{3t}{p}\right) - \left(\frac{4t}{p}\right)_3 - \left(\frac{4t}{p}\right)_3^{-1} - N_p(t),$$

$N_p(t)$  number of **affine**  $\mathbb{F}_p$ -points on  $y^4(1-y)^2 = x^2(1-x)/a$ , with  $a = 27/(4t)$ .

$$a_p(t) = p + 1 - M_p(t),$$

$M_p(t)$  number of **projective** points on

$$Y^2 = X^3 - aX + a.$$