

Chaotic D-Term Inflation

Teruhiko Kawano

*Department of Physics, University of Tokyo
Hongo, Tokyo 113-0033, Japan*

A simple model for chaotic inflation in supergravity is proposed. The model is $\mathcal{N} = 1$ supersymmetric massive U(1) gauge theory via the Stuckelberg superfield and gives rise to D-term inflation with a quadratic term of inflaton in the potential. The Fayet-Iliopoulos field plays a role of the inflaton.

December, 2007

Cosmological inflation is a very attractive idea to solve the horizon problem and the flatness problem in the early universe [1], and it also explains the density fluctuations in the cosmic microwave background. Thus, it would be interesting to incorporate the idea into viable models of high energy physics. Since supersymmetry is one of the solutions to the hierarchy problem between the electoweak scale and the Planck scale, it is natural to consider the inflation scenario in supergravity.

Among many inflation scenarios, chaotic inflation doesn't suffer from the initial condition problems and gives rise to a large tensor-to-scalar ratio, which would be observed by the Planck satellite observation. However, it doesn't seem an easy task to find chaotic inflation models in supergravity [2]. This is because the F-term potential in supergravity have the exponential factor of the Kähler potential in general to prevent inflaton fields from getting the large fluctuation required by chaotic inflation. One thus is led to make use of the D-term potential to give rise to chaotic inflation.

In fact, the paper [3] is the first to give a chaotic inflation model by using the D-term potential, where it was shown that the quartic term of an inflaton field could be obtained from the D-term potential. Therefore, it would be interesting to obtain the quadratic term of inflaton fields from the D-term potential. Since in general, the D-term potential provides the quartic term of charged fields transforming linearly under the gauge group, it seems difficult to obtain the quadratic term of an inflaton field. However, the Fayet-Iliopoulos parameter of a U(1) gauge group enters the D-term potential quadratically. Thus, if one can replace the Fayet-Iliopoulos parameter by a dynamical field, one can expect that the field could play a role of an inflaton field. In this paper, it will be shown that this is indeed the case.

Let us consider $\mathcal{N} = 1$ supersymmetric massive U(1) gauge theory with the Stuckelburg superfield S given in terms of component fields by

$$S = \frac{1}{\sqrt{2}} (\rho + i\zeta) + \sqrt{2}\theta\psi + \theta^2 F, \quad (1)$$

where ρ and ζ are real scalar fields and play roles as the Stuckelburg field and the Fayet-Iliopoulos field, respectively, as will be seen. The U(1) gauge vector multiplet is given in the Wess-Zumino gauge by

$$V = -i\theta\sigma^\mu\bar{\theta}v_\mu - i\bar{\theta}^2\theta\lambda + i\theta^2\bar{\theta}\bar{\lambda} + \frac{1}{2}\theta^2\bar{\theta}^2 D. \quad (2)$$

The Lagrangian density L is given by

$$L = -\frac{1}{2} \int d^2\theta d^2\bar{\theta} \left(S - \bar{S} + i\sqrt{2}M_p V \right)^2 + \text{Re} \left[\int d^2\theta \frac{1}{2e^2} W^\alpha W_\alpha \right], \quad (3)$$

where M_p is a mass scale and the gaugino superfield W_α is given by

$$W_\alpha = -\frac{1}{4} \bar{D}^2 D_\alpha V. \quad (4)$$

The action is invariant under the U(1) gauge transformation

$$V \rightarrow V + i(\Lambda - \bar{\Lambda}), \quad S \rightarrow S + \sqrt{2}M_p \Lambda, \quad \bar{S} \rightarrow \bar{S} + \sqrt{2}M_p \bar{\Lambda}. \quad (5)$$

Note here that the non-linear transformation of the chiral superfield S prohibits it from appearing in the gauge kinetic function. Rescaling the vector superfield V to obtain the canonical kinetic term for it, in component fields one finds the Lagrangian density

$$L = -\frac{1}{2} (\partial_\mu \zeta)^2 - \frac{1}{2} (\partial_\mu \rho - M v_\mu)^2 - \frac{1}{4} F_{\mu\nu}^2 - i\bar{\psi} \bar{\sigma}^\mu \partial_\mu \psi - i\bar{\lambda} \bar{\sigma}^\mu \partial_\mu \lambda \\ + M\zeta D - M\psi\lambda - M\bar{\psi}\bar{\lambda} + |F|^2 + \frac{1}{2} D^2, \quad (6)$$

where M is defined as $M = eM_p$. It is obvious from (6) that the field ρ plays a role of the Stueckelberg field and gives mass to the gauge field v_μ . The field ζ plays a role of the Fayet-Iliopoulos parameter, but here it is a real scalar field. Changing the variable $MA_\mu = Mv_\mu - \partial_\mu \rho$ and solving the equation of motions of the auxiliary fields D and F , one obtains the Lagrangian density

$$L = -\frac{1}{2} (\partial_\mu \zeta)^2 - \frac{1}{2} M^2 \zeta^2 - \frac{1}{4} F_{\mu\nu}^2 - \frac{1}{2} M^2 A_\mu^2 - i\bar{\psi} \bar{\sigma}^\mu \partial_\mu \psi - i\bar{\lambda} \bar{\sigma}^\mu \partial_\mu \lambda - M\psi\lambda - M\bar{\psi}\bar{\lambda}. \quad (7)$$

This is the Lagrangian density of a free massive U(1) vector multiplet. The Fayet-Iliopoulos field ζ plays a role of inflaton. Therefore, one expects that the mass M of the inflaton field ζ is of order $10^{13} \sim 10^{14}$ GeV. Thus, if the mass scale M_p is set to the reduced Planck scale $\sim 10^{18}$ GeV, the U(1) gauge coupling may be of order $10^{-4} \sim 10^{-5}$ to give rise to the observed density fluctuations. However, one can see that the reheating doesn't occur without any modification in the model. Since the model is very simple, we hope that a simple modification of it could lead to a full-fledged inflation model.

Acknowledgements

The author would like to thank Koichi Hamaguchi, Yuuki Shinbara, Taizan Watari, and Masahide Yamaguchi for helpful discussions and reading the manuscript. The work was supported in part by a Grant-in-Aid (#19540268) from the MEXT of Japan.

References

- [1] See, for example, A. D. Linde, *Particle Physics and Inflationary Cosmology*, (Harwood, Chur, Switzerland, 1990);
A. R. Liddle and D. H. Lyth, *Cosmological Inflation and Large Scale Structure*, (Cambridge University Press, Cambridge, England, 2000);
D. H. Lyth and A. Riotto, “Particle Physics Models of Inflation and the Cosmological Density Perturbation,” *Phys. Rept.* **314**, 1 (1999), [hep-ph/9807278](#).
- [2] A. B. Goncharov and A. D. Linde, “Chaotic Inflation in Supergravity,” *Phys. Lett. B* **139**, 27 (1984); “A Simple Realization of the Inflationary Universe Scenario in SU(1,1) Supergravity,” *Class. Quant. Grav.* **1**, L75 (1984);
H. Murayama, H. Suzuki, T. Yanagida and J. Yokoyama, “Chaotic Inflation and Baryogenesis in Supergravity,” *Phys. Rev. D* **50**, 2356 (1994), [hep-ph/9311326](#);
M. Kawasaki, M. Yamaguchi and T. Yanagida, “Natural Chaotic Inflation in Supergravity,” *Phys. Rev. Lett.* **85**, 3572 (2000), [hep-ph/0004243](#);
J. R. Ellis, Z. Lalak, S. Pokorski and K. Turzynski, “The Price of WMAP Inflation in Supergravity,” *JCAP* **0610**, 005 (2006), [hep-th/0606133](#).
- [3] K. Kadota and M. Yamaguchi, “D-term Chaotic Inflation in Supergravity,” *Phys. Rev. D* **76**, 103522 (2007), [arXiv:0706.2676 \[hep-ph\]](#).