

# Comments on "Slip coefficient in nanoscale pore flow"

Zotin K.-H. Chu

*3/F, 24, 260th. Lane, First Section, Muzha Road,  
Wen-Shan District, Taipei, Taiwan 116, China*

## Abstract

We make some remarks on Sokhan and Quirke's [*Phys. Rev. E* 78, 015301(R) (2008)] paper. Sokhan and Quirke mentioned that, considering their main result, the slip coefficient is independent of the external force (flux) which is not consistent with previous measurements and approaches. We also discuss the sudden changes of the slip coefficient for larger Knudsen numbers or smaller nanopores.

PACS numbers: 47.15.gm, 05.20.Jj, 47.11.Mn, 66.20.-d

Sokhan and Quirke just presented an expression for the slip coefficient ( $l_s$ ) applicable to flows in nanoscale pores, which, as they claimed [1], has been verified by nonequilibrium molecular-dynamics simulation. Their results showed that the slip coefficient depends strongly on the pore width for small pores tending to a constant value for pores of width ( $H$ )  $> 20$  molecular diameters ( $\sigma$ ) for their systems, in contrast to the linear scaling predicted by Maxwell's theory of slip. Note that  $\mathbf{u}_s = l_s \nabla \mathbf{u}$  in [1] is of doubt as the left-hand-side (the slip velocity) is a vector while the right-hand-side is generally a tensor ( $\nabla \mathbf{u}$  : the shear rate). To the best knowledge of the present author, the slip velocity is normally defined in one-dimensional sense [2-3].

The present author, however, based on his previous experiences [2] (the gravity-driven case in [1] is similar to that of electric field-driven case in [2] considering the external force), would like to make some remarks about their presentation [1]. To begin with, we like to recall the recent critical review : '*The continuum approximation seems to break down below 10 nm in case of water in nanotubes. Smooth liquid-gas interface (meniscus) disappears in tubes with the diameter less than 8-10 nm and an anomalous behavior of water is observed in 1-7 nm carbon nanotubes*' [4]. The main results for a *nanoscale* pore flow claimed in [1] (cf. Eq. (8) therein) :

$$l_s = \frac{\tau\eta}{\rho h} - \frac{H}{6}, \quad (1)$$

where  $\tau$  is the relaxation time,  $\eta$  : the interfacial shear viscosity,  $\rho$  : mass density of the fluid and  $h = H/2$ , by Sokhan and Quirke are still mainly based on the continuum approximation : Equations (2),(3),(6) in [1] which contradict to the summarized results just mentioned above [4] and other researchers' claim [5]. There is no wonder that the authors of [1] made remarks : *Deviation of the order of 10% from the bulk values for the  $H = 5\sigma$  (cf. the detailed explanation of the notations in [1]) pore is due to the overlap of adsorbed layers at two surfaces and to inaccuracy in determination of the reference bulk density. At lower densities, when  $Kn \geq 1$ , the viscosity in the pore deviates markedly from bulk values.* To be concise, the continuum approximation breaks down for  $Kn \geq 1$  [2,6]. On the other hand, the present author argues that Sokhan and Quirke should illustrate the realistic range of  $H$  and  $\sigma$  in terms of dimensional units for their calculations in [1]. What are the exactly valid minimum dimensions for  $H$  and  $\sigma$  considering their numerical approaches [1]? Note that Mattia and Gogotsi raised the question : Whether the nonslip boundary condition is applicable at very small scales remains open [4].

Secondly, it was mentioned in [1] (cf. statements near Eq. (8) in [1]), saying that the main result of [1], that the slip coefficient (or slip length) is independent of the external force (flux), but depends nonlinearly on the pore width, both directly and indirectly through the relaxation time (cf. Eq. (1) above). This result is quite different from those presented by Majumder *et al.* [7] (cf. Table 1 in [7] especially for those different external-forcing measurements or calculations for the slip length about water). At least, considering Majumder *et al.*'s data [7], for the same fluid (water) : increasing the external forcing will increase the slip coefficient (or slip length). Meanwhile this result is also different from those presented in [2] or [8] where the external forcing can adjust the (averaged) velocity profiles (not to mention the slip velocity near the walls or interfaces). To be specific, the present author likes to argue that there is a possibility that *grazing* collisions between particles (atoms and molecules) and surfaces (or boundaries) [9] occurred for those particles approaching to as well as reflecting from or along the nanopore walls. The latter could be neglected by Sokhan and Quirke in [1]. In essence, the grazing collisions between particles and boundaries are mainly tuned by the external forcing [9].

Thirdly, Sokhan and Quirke claimed [1] for the systems with the same Knudsen number ( $\text{Kn}=\lambda$  (mean free path)/ $H$  (pore width)), the slip coefficient increases with the pore width (cf. Fig. 2 in [1]). This might be trivial in microdomains (but not in nanodomains) from previous kinetic theory as the Knudsen number being fixed (once  $H$  increases then  $\lambda$  should also increase so that  $\text{Kn}$  is kept to be the same; considering Sokhan and Quirke's derivation :

$$l_s = \lambda \frac{2}{\alpha} - \frac{H}{6}, \quad (2)$$

compared to the Maxwell's relation :

$$l_s = \lambda \left( \frac{2}{\alpha} - 1 \right) \quad (3)$$

cf. Eqs. (9) and (1) in [1]; here  $\alpha$  defines a fraction of specularly reflected molecules or  $\alpha$  defines the fraction of the flux of tangential momentum transmitted in collisions and is called the 'accommodation coefficient' (TMAC) [1]) then in transitional flow regime ( $\text{Kn} \approx \text{O}(1)$ ) due to the increasing of the mean free path, the slip velocity (proportional to the slip coefficient or slip length) will also increase (cf. Fig. 3 in [2]). On the other hand, to check the trend illustrated in Fig. 3 of [1], we have, using Eq. (9) in [1] and  $\rho = 0.125m/\sigma^3$  (from

$n^* = 0.125 = \rho\sigma^3/m$ , please see the first line of the right column at page 4 of [1]),

$$l_s = \frac{32\eta\sigma^3}{m\bar{c}\alpha} - \frac{H}{6} \equiv l_s|_{SQ} \quad (4)$$

which is related to the main result derived by Sokhan and Quirke in [1] and

$$l_s = \frac{32\eta\sigma^3}{m\bar{c}\alpha} - \frac{16\eta\sigma^3}{m\bar{c}} \equiv l_s|_{Max} \quad (5)$$

considering the Maxwell's expression (i.e., Eq. (1)) in [1], where  $\bar{c}$  is the mean speed of the molecules. The difference of above two expressions then give us the understanding of those shown in Fig. 3 of [1]:

$$l_s|_{Max} - l_s|_{SQ} = \frac{16\eta\sigma^3}{m\bar{c}} - \frac{H}{6} \equiv \lambda - \frac{H}{6}. \quad (6)$$

Unfortunately, even we normalized this difference with respect to  $\sigma$  (the same as that of Fig. 3 of [1]), the obtained difference is still dependent upon  $\eta$ ,  $m$ , and  $\bar{c}$  (or  $\lambda$ ). However, we cannot get the values of  $\eta$ ,  $m$ , and  $\bar{c}$  (or  $\lambda$ ) from [1] to evaluate the further detailed difference. Can Sokhan and Quirke clarify this issue? What is the Kn for calculations illustrated in Fig. 3 of [1]? Meanwhile for  $H \sim \sigma$ , what is the realistic *mean free path* for particles or molecules in confined nanodomains? This is crucial to our judgement whether Fig. 3 in [1] is universal?

Finally, the present author, based on the sudden decrease of the slip coefficient of case  $H = 5\sigma$  (w.r.t.  $H = 10\sigma$ ) for the larger Kn (say,  $\text{Kn} > 3.0$ ) in Fig. 2 of [1], also wonders that once  $H < 5\sigma$ , the curves of  $l_s/H$  vs. Kn will become nonmonotonic as the decreasing of  $l_s/H$  for smaller  $H$  (less than  $5\sigma$ ) will continue? We argue that the possible cause is due to the estimation from the relaxation times and bulk viscosities at densities equal to that in the middle of the pore considering Sokhan and Quirke's calculations [1]. It seems to us the latter treatment should be refined in much more narrow nanopores considering the other realistic physical influences due to confinement (e.g., is the mean free path definition still be valid in confined nanodomains?).

---

[1] V.P. Sokhan and N. Quirke, Phys. Rev. E **78**, 015301(R) (2008).

[2] P.H. Paul, M.G. Garguilo and D.J. Rakestraw, Anal. Chem. **70**, 2459 (1998). A. K.-H. Chu, Phys. Rev. E **67**, 041106 (2003).

- [3] A.V. Latyshev, V.N. Popov and A.A. Yushkanov, *High Temperature* **42**, 134 (2004).
- [4] D. Mattia and Y. Gogotsi, *Microfluid Nanofluid* **5**, 289 (2008).
- [5] R.E. Tuzun, D.W Noid, B.G. Sumpter and R.C. Merkle, *Nanotechnology* **7**, 241 (1996). R.E. Tuzun, D.W Noid, B.G. Sumpter and R.C. Merkle, *Nanotechnology* **8**, 112 (1997).
- [6] C. Cercignani, *Mathematical Methods in Kinetic Theory* (Plenum, New York, 1969).
- [7] M. Majumder, N. Chopra, R. Andrews and B.J. Hinds, *Nature* **438**, 44 (2005). [Correction] *Nature* **438**, 930 (2005).
- [8] J.K. Holt, H.G. Park, Y.m. Wang, M. Stadermann, A.B. Artyukhin, C.P. Grigoropoulos, A. Noy and O. Bakajin, *Science* **312**, 1034 (2006).
- [9] K.-H. W. Chu, *J. Phys. A : Math. Gen.* **33**, 7103 (2000).