

On a possible observation of “dihelion” in dissociation of relativistic ${}^9\text{C}$ nuclei

D. A. Artemenkov, N. K. Kornegrutsa, D. O. Krivenkov, R. Stanoeva, and P. I. Zarubin*

Joint Insitute for Nuclear Research, Dubna, Russia

(Dated: September 5, 2021)

Abstract

Nuclear track emulsion is exposed to $1.2 \text{ A } {}^9\text{C}$ GeV nuclei. Pairs of 2^3He nuclei having unusually narrow opening angles are observed in channel ${}^9\text{C} \rightarrow 3^3\text{He}$ pointing to the possible 2^3He resonance near the production threshold.

PACS numbers: 21.45.+v, 23.60+e, 25.10.+s

arXiv:1105.3813v1 [nucl-ex] 19 May 2011

*Electronic address: zarubin@lhe.jinr.ru; URL: <http://becquerel.jinr.ru>

I. INTRODUCTION

The BECQUEREL Collaboration explores the peripheral dissociation of light relativistic nuclei in nuclear emulsion [1]. The structural features of a projectile nucleus are most clearly manifested in the coherent dissociation of nuclei not accompanied by the formation of the target nucleus fragments and mesons (or in “white” stars). Nucleon clustering in stable and radioactive Be, B, C and N isotopes are studied in this approach with a complete observation of the fragments at a record angular resolution [2–8]. Macro photos of such interactions are assembled in the collection [1].

In particular, nuclear track emulsion is irradiated in a ${}^9\text{C}$ nucleus beam produced at the JINR Nuclotron by the fragmentation of 1.2 A GeV ${}^{12}\text{C}$ nuclei [8]. Dominance of the ${}^9\text{C}$ isotope in the secondary beam is confirmed by emulsion measurements of the ionization of the secondary beam nuclei, features of the charge topology of their fragmentation, as well as momentum measurements of the accompanying the ${}^3\text{He}$ nuclei. As is established, the ${}^8\text{B} + p$, ${}^7\text{Be} + 2p$, $2\text{He} + 2p$ and $\text{He} + 2\text{H} + 2p$ channels are leading ones in the ${}^9\text{C}$ coherent dissociation [7]. The two latter are related to the ${}^7\text{Be}$ core dissociation.

In addition, population of the three ${}^3\text{He}$ nucleus state is observed in the ${}^9\text{C}$ coherent dissociation with 14% probability. Its origin can be caused by a virtual regrouping of a neutron from the ${}^4\text{He}$ cluster to form ${}^3\text{He}$ clusters as well as the presence of the $3{}^3\text{He}$ component in the ${}^9\text{C}$ ground state. In the second variant, the probability of the $3{}^3\text{He}$ ensemble production points to the significance of this deeply bound configuration in the wave function of the ${}^9\text{C}$ ground state. The mechanism of the ${}^9\text{C}$ coherent dissociation in the channels with nucleon separation and the $3{}^3\text{He}$ channel is a nuclear diffractive interaction which is established on the basis of measurements of the total transverse momentum transferred to the fragment ensemble.

II. NARROW PAIRS IN COHERENT DISSOCIATION ${}^9\text{C} \rightarrow 3{}^3\text{He}$

Few pairs of ${}^3\text{He}$ nuclei with opening angles $\Theta_{2\text{He}}$ less than 10^{-2} rad are detected in the ${}^9\text{C} \rightarrow 3{}^3\text{He}$ channel. The macro photo of one identified event is shown in figure 1. This observation indicates to a possible existence of a narrow $2{}^3\text{He}$ resonant state with the decay energy near the $2{}^3\text{He}$ mass threshold (or “dihelion”). In the same way the formation



FIG. 1: Macro photo of “white” star of ${}^9\text{C}$ dissociation to $3{}^3\text{He}$ nuclei in nuclear track emulsion; the interaction vertex IV is shown by the arrow.




FIG. 2: Total distribution of opening angles $\Theta_{2\text{He}}$ between the relativistic He fragments in the “white” stars ${}^9\text{C} \rightarrow 3{}^3\text{He}$ and in events ${}^8\text{B} \rightarrow 2\text{He} + \text{H}$ with the formation of target nucleus fragments or meson; dotted line indicates the “white” stars contribution.

of ${}^8\text{Be}$ nuclei is reliably manifested in the production of ${}^4\text{He}$ pairs with extremely small opening angles in the relativistic dissociation of ${}^9\text{Be} \rightarrow 2{}^4\text{He}$ [3] and ${}^{10}\text{C} \rightarrow 2{}^4\text{He} + 2p$ [4, 5]. Significant probability of coherent dissociation ${}^9\text{C} \rightarrow 3{}^3\text{He}$ makes it an efficient source for search for an analog of the unbound ${}^8\text{Be}$ nucleus among the ${}^3\text{He}$ pairs. In what follows, a rather unexpected and potentially important feature of the spectrum $\Theta_{2\text{He}}$ of ${}^3\text{He}$ pairs produced in the dissociation of ${}^9\text{C}$ [7] and ${}^8\text{B}$ [8] nuclei, is given.

In figure 2 the dotted line shows the distribution $\Theta_{2\text{He}}$ for “white” stars ${}^9\text{C} \rightarrow 3{}^3\text{He}$. Its main part, corresponding to 30 pairs, is described by a Gaussian distribution with mean value $\langle \Theta_{2\text{He}} \rangle = (46 \pm 3) \times 10^{-3} \text{ rad}$ (RMS $16 \times 10^{-3} \text{ rad}$). In addition, thanks to excellent spatial resolution eight ${}^3\text{He}$ pairs within $\Theta_{2\text{He}} < 10^{-2} \text{ rad}$ are reliably observed. These

pairs form a special group with a mean value $\langle Q(2^3He) \rangle = (6 \pm 1) \times 10^{-3}$ rad and RMS 3×10^{-3} rad, which is obviously beyond the previous description. These values correspond to the average relative energy $\langle Q(2^3He) \rangle = (142 \pm 35)$ keV (RMS 100 keV). The parameter $Q(2^3He)$ was defined as the difference between the invariant mass of the pair and the double 3He mass assuming that the fragments conserve the 9C momentum per nucleon.

III. NARROW PAIRS IN PERIPHERAL DISSOCIATION $^8B \rightarrow 2He + H$

Additional search for the resonance 2^3He is carried out in the events of 1.2 A GeV peripheral dissociation $^8B \rightarrow 2He + H$. In this case, inelastic interactions with target nucleus fragments or produced mesons are selected in order to enhance the “dihelion” effect. This condition provides the selection of interactions with knocking of a neutron out of the 4He cluster in the 8B nucleus. Thus, the resulting distribution Θ_{2^3He} in the figure also includes a separate group of narrow pairs with $\langle \Theta_{2^3He} \rangle = (4.5 \pm 0.5) \times 10^{-3}$ rad (RMS 1.5×10^{-3} rad), corresponding to the case of “white” stars $^9C \rightarrow 3^3He$.

The total distribution for both the (figure 2) makes the indication to the existence of the 2^3He resonance more reliable. Moreover, the question arises about the nature of the broad peak with a maximum of Θ_{2^3He} about $(40 - 50) \times 10^{-3}$ rad. It is possible that in this Θ_{2^3He} region the 2^3He system shows its similarity with the first excited 2^+ state of the 8Be nucleus.

IV. CONCLUSIONS

Obviously, the “dihelion” indication should be reviewed using a significantly larger statistics. The BECQUEREL Collaboration performed irradiation of nuclear track emulsion in a mixed beam of 1.2 A GeV ^{12}N , ^{10}C and 7Be nuclei [5]. Thus, there are new opportunities with regard to the issue of “dihelion” based on the analysis of the found 400 “non-white” stars $^7Be \rightarrow 2^3He$ with knocking out of a neutron and the formation of fragments of target nuclei or mesons, as in the case of $^8B \rightarrow 2He + H$.

However, it is possible that the “dihelion” formation is due to the presence of the 2^3He component in the 9C and 8B structures. In principle, in a lighter 7Be nucleus such a component can be suppressed, this means that the “dihelion” formation can be suppressed as well. Therefore, it is important to search for the 2^3He resonance with high statistics exactly

in low energy ${}^9\text{C}$ and ${}^8\text{B}$ beams. At the same time, pointing to the existence of “dihelion”, our observation motivates the search for a mirror state of a pair of nuclei ${}^3\text{H}$ - “ditriton”.

Acknowledgments

This work was supported by grants from 96-1596423, 02-02-164-12a, 03-02-16134, 03-02-17079, 04-02-17151. 04-02-16593, and 09-02-9126 CT-a RFBR. The authors thank S. P. Kharlamov (FIAN, Moscow), and A. I. Malakhov (JINR, Dubna) for discussions and friendly support.

-
- [1] The BECQUEREL Project, <http://becquerel.jinr.ru/>
- [2] Peresadko N G et al. 2007 Phys. At. Nucl. 70, 1226 (Preprint arXiv:nucl-ex/0605014)
- [3] Artemenkov D A, Krivenkov D O, Shchedrina T V, Stanoeva R and Zarubin P I 2008 Few Body Syst. 44, 273
- [4] Stanoeva R et al. 2009 Phys. At. Nucl. 72, 690 (Preprint arXiv:0906.4220)
- [5] Kattabekov R R, Mamatkulov K Z et al. 2010 Phys. At. Nucl. 73, 2110 (Preprint arXiv:1104.5320)
- [6] Artemenkov D A, Alikulov S S, Kattabekov R R, Mamatkulov K Z, Kornegrutsa N K, Krivenkov D O and Zarubin P I 2011 Few Body Syst. 50, 259 (Preprint arXiv: 1105.2438)
- [7] Krivenkov D O et al. 2010 Phys. At. Nucl. 73, 2103 (Preprint arXiv:1104.2439)
- [8] Rukoyatkin P A et al. 2008 Eur. Phys. J. ST 162, 267