

C^* -ALGEBRAS OF LABELLED GRAPHS III - K -THEORY COMPUTATIONS

TERESA BATES, TOKE MEIER CARLSEN, AND DAVID PASK

ABSTRACT. In this paper we give a formula for the K -theory of the C^* -algebra of a weakly left-resolving labelled space. This is done by realising the C^* -algebra of a weakly left-resolving labelled space as the Cuntz-Pimsner algebra of a C^* -correspondence. As a corollary we get a gauge invariant uniqueness theorem for the C^* -algebra of any weakly left-resolving labelled space. In order to achieve this we must modify the definition of the C^* -algebra of a weakly left-resolving labelled space.

We also establish strong connections between the various classes of C^* -algebras which are associated with shift spaces and labelled graph algebras. Hence, by computing the K -theory of a labelled graph algebra we are providing a common framework for computing the K -theory of graph algebras, ultragraph algebras, Exel-Laca algebras, Matsumoto algebras and the C^* -algebras of Carlsen.

We provide an inductive limit approach for computing the K -groups of an important class of labelled graph algebras, and give examples.

1. INTRODUCTION

The purpose of this paper is to provide a formula for and a practical method of computing the K -theory of the C^* -algebra $C^*(E, \mathcal{L}, \mathcal{B})$ of a weakly left-resolving labelled space $(E, \mathcal{L}, \mathcal{B})$ (see Theorem 4.4 and Theorem A.9). To do this we first realise $C^*(E, \mathcal{L}, \mathcal{B})$ as a Cuntz-Pimsner algebra (this is done in Theorem 3.8, see also Remark A.7) and then use the results of [20] to identify the K -groups in question as the kernel and cokernel of a certain map. Then, for an important class of labelled graph C^* -algebras, we give a procedure for computing these groups through an inductive limit process (see Theorem 6.5). Our construction of $C^*(E, \mathcal{L}, \mathcal{B})$ as a Cuntz-Pimsner algebra also allows us to derive a gauge invariant uniqueness theorem for $C^*(E, \mathcal{L}, \mathcal{B})$ (see Corollaries 3.10 and A.8) and a proof that $C^*(E, \mathcal{L}, \mathcal{B})$ is nuclear (see Corollary 3.9 and Remark A.7), and, under a suitable countability condition, satisfies the Universal Coefficient Theorem of [42] (see Corollary 3.11 and Remark A.7).

As was already pointed out in [5], the original definition of $C^*(E, \mathcal{L}, \mathcal{B})$ has to be modified in order to avoid degeneracy of certain projections when E has sinks. Furthermore, in [6] it was pointed out that the Gauge Invariant Uniqueness Theorem stated in [4] was incorrect and a new (and correct) gauge invariant uniqueness theorem for $C^*(E, \mathcal{L}, \mathcal{B})$ when $(E, \mathcal{L}, \mathcal{B})$, in addition to being weakly left-resolving, is so-called set-finite (every set of vertices in \mathcal{B} emits a finite number of different labels) and \mathcal{B} is closed under relative complements was proved. In order to get a gauge invariant uniqueness theorem for the C^* -algebra of any weakly left-resolving labelled space it is necessary to change the definition of $C^*(E, \mathcal{L}, \mathcal{B})$ in the case where \mathcal{B} is not closed under relative complements (see the appendix for details). Since this makes things a bit more complicated, we have chosen to only work with weakly left-resolving labelled spaces $(E, \mathcal{L}, \mathcal{B})$ where \mathcal{B} is closed under relative complements (we call such labelled spaces *normal*,

Date: September 25, 2018.

This research was supported by the Australian Research Council, The Research Council of Norway and The Danish Natural Science Research Council.

see Subsection 2) in the main part of the paper, but in the appendix we show how to deal with the general case.

The motivation for our K -theory computations is to investigate the relationship between certain dynamical invariants of shift spaces and the K -theoretical invariants of C^* -algebras associated to these shift spaces. This connection was first brought to light in the work of Cuntz and Krieger in [12] where they showed how to associate a C^* -algebra \mathcal{O}_A to a finite 0-1 matrix A with no zero rows or columns, provided that the matrix satisfied a certain condition called (I). In [13, Proposition 3.1] it was shown that the K -groups of a Cuntz-Krieger algebra are isomorphic to the Bowen-Franks groups of the shift of finite type X_A associated to A (see [7]). Thus, a deep connection was established between the combinatorially-defined C^* -algebra \mathcal{O}_A and the (one-sided) shift of finite type X_A . Several generalisations of Cuntz-Krieger algebras have now been widely studied.

Combining the universal algebra approach of [1] and the graphical approach to Cuntz-Krieger algebras begun in [15], graph algebras were introduced in [25]. Graph algebras were originally defined for graphs satisfying a finiteness condition – the need for this condition was removed by Fowler and Raeburn in [18] (see also [17]). Using a different approach, Exel and Laca showed how to associate a C^* -algebra to an infinite 0-1 matrix with no zero rows or columns in [16]. A link between graph algebras and Exel-Laca algebras was provided by the ultragraph algebras introduced by Tomforde (see [39, 40]).

Motivated by the symbolic dynamical data contained in a Cuntz-Krieger algebra, Matsumoto provided a generalisation of Cuntz-Krieger algebras by associating to an arbitrary two-sided shift space Λ over a finite alphabet a C^* -algebra \mathcal{O}_Λ (see [27, 28, 29, 30, 32, 33]). Later Carlsen and Matsumoto modified this construction (see [9, 35]), and Carlsen further modified the definition of \mathcal{O}_Λ and extended it to one-sided shift spaces in [8] (see [10] for a discussion of the relationship between the three different definitions of \mathcal{O}_Λ and for an alternative construction of the C^* -algebra constructed in [8]). The results of [28, 29, 31, 34, 35] give connections between certain dynamical invariants of Λ and K -theoretical invariants of \mathcal{O}_Λ in the same spirit as [13].

By adapting the left-Krieger cover construction given in [23], any shift space over a countable alphabet may be presented by a left-resolving labelled graph. Hence, the labelled graph algebras introduced in [4] provide a method for associating a C^* -algebra to a shift space over a countable alphabet. As we shall see in Section 5, the class of labelled graph algebras contains the class of graph algebras and all of the classes of C^* -algebras discussed above. By computing the K -theory of a labelled graph algebra we will therefore be providing a unified approach to computing the K -theory of this wide collection of C^* -algebras.

The paper is organised as follows: Section 2 contains some background on labelled graphs, labelled spaces and their C^* -algebras. In Section 3 we give our key result, Theorem 3.8, which shows that the C^* -algebra of a weakly left-resolving normal labelled space may be realised as a Cuntz-Pimsner algebra in the sense of [20], and we also show a gauge invariant uniqueness theorem for it (Corollary 3.10) and that it is nuclear (Corollary 3.9), and, under a suitable countability condition, satisfies the Universal Coefficient Theorem of [42] (Corollary 3.11). In Section 4 we use the general results of [20] to provide a formula for the K -theory of the C^* -algebra of a weakly left-resolving normal labelled space (see Theorem 4.4). Then in Section 5 we show how our K -theory formulas reduce to those for graph algebras, ultragraph algebras and Matsumoto algebras in the appropriate cases. Furthermore in Proposition 5.6 we realise the Carlsen algebra $C^*(X)$ associated to a one-sided shift space X as a labelled graph algebra and hence compute its K -theory (see Corollary 5.7). In Section 6 we show how to compute the K -theory of certain labelled graph algebras as inductive limits; this is done in analogy with the computations of [36], and in Section 7 we provide a few examples of the computations outlined in Section 6. Finally, in the appendix we give an example that illustrates the necessity of

changing the definition of the C^* -algebra of a weakly left-resolving labelled space which is not normal in order to obtain a gauge invariant uniqueness theorem. We state this new definition and generalise the main results of Sections 3 and 4 to the non-normal case.

Acknowledgements. The authors wish to thank the Fields Institute and the Banff International Research Station for providing the facilities for them to discuss various aspects of this paper.

2. LABELLED SPACES AND THEIR C^* -ALGEBRAS

In this section we will briefly review the definitions of labelled spaces. We also introduce the notion of a *normal* labelled space and associate a C^* -algebras with any weakly left-resolving normal labelled space.

Directed graphs. A directed graph E consists of a quadruple (E^0, E^1, r, s) where E^0 and E^1 are sets of vertices and edges, respectively and $r, s : E^1 \rightarrow E^0$ are maps giving the direction of each edge. A path $\lambda = \lambda_1 \dots \lambda_n$ is a sequence of edges $\lambda_i \in E^1$ such that $r(\lambda_i) = s(\lambda_{i+1})$ for $i = 1, \dots, n-1$; we define $s(\lambda) = s(\lambda_1)$ and $r(\lambda) = r(\lambda_n)$. The collection of paths of length n in E is denoted by E^n and the collection of all finite paths in E by E^* , so that $E^* = \bigcup_{n \geq 1} E^n$.

A *loop* in E is a path which begins and ends at the same vertex, that is $\lambda \in E^*$ with $s(\lambda) = r(\lambda)$. We say that E is *row-finite* if every vertex emits finitely many edges. We denote the collection of all infinite paths in E by E^∞ . A vertex $v \in E^0$ is a *sink* if $s^{-1}(v) = \emptyset$ and we define E_{sink}^0 to be the set of all sinks in E^0 .

Labelled graphs. A *labelled graph* (E, \mathcal{L}) over a countable alphabet \mathcal{A} consists of a directed graph E together with a labelling map $\mathcal{L} : E^1 \rightarrow \mathcal{A}$. By replacing \mathcal{A} with $\mathcal{L}(E^1)$ (if necessary) we may assume that the map \mathcal{L} is onto.

Let \mathcal{A}^* be the collection of all *words* in the symbols of \mathcal{A} . The map \mathcal{L} extends naturally to a map $\mathcal{L} : E^n \rightarrow \mathcal{A}^*$, where $n \geq 1$. For $\lambda = e_1 \dots e_n \in E^n$ we set $\mathcal{L}(\lambda) = \mathcal{L}(e_1) \dots \mathcal{L}(e_n)$. In this case the path $\lambda \in E^n$ is said to be a *representative* of the *labelled path* $\mathcal{L}(e_1) \dots \mathcal{L}(e_n)$. Let $\mathcal{L}(E^n)$ denote the collection of all labelled paths in (E, \mathcal{L}) of length n where we write $|\alpha| = n$ if $\alpha \in \mathcal{L}(E^n)$. The set $\mathcal{L}^*(E) = \bigcup_{n \geq 1} \mathcal{L}(E^n)$ is the collection of all labelled paths in the labelled graph (E, \mathcal{L}) .

The labelled graph (E, \mathcal{L}) is *left-resolving* if for all $v \in E^0$ the map $\mathcal{L} : r^{-1}(v) \rightarrow \mathcal{A}$ is injective. The left-resolving condition ensures that for all $v \in E^0$ the labels of the incoming edges to v are all different. For $\alpha \in \mathcal{L}^*(E)$ we put

$$s_{\mathcal{L}}(\alpha) = \{s(\lambda) \in E^0 : \mathcal{L}(\lambda) = \alpha\} \text{ and } r_{\mathcal{L}}(\alpha) = \{r(\lambda) \in E^0 : \mathcal{L}(\lambda) = \alpha\},$$

so that $r_{\mathcal{L}}, s_{\mathcal{L}} : \mathcal{L}^*(E) \rightarrow 2^{E^0}$, where 2^{E^0} denotes the set of subsets of E^0 . We shall drop the subscript on $r_{\mathcal{L}}$ and $s_{\mathcal{L}}$ if the context in which it is being used is clear.

For $A \subseteq E^0$ and $\alpha \in \mathcal{L}^*(E)$ the *relative range of α with respect to A* is defined to be

$$r_{\mathcal{L}}(A, \alpha) = \{r(\lambda) : \lambda \in E^*, \mathcal{L}(\lambda) = \alpha, s(\lambda) \in A\}.$$

A collection $\mathcal{B} \subseteq 2^{E^0}$ of subsets of E^0 is said to be *closed under relative ranges for (E, \mathcal{L})* if for all $A \in \mathcal{B}$ and $\alpha \in \mathcal{L}^*(E)$ we have $r_{\mathcal{L}}(A, \alpha) \in \mathcal{B}$. If \mathcal{B} is closed under relative ranges for (E, \mathcal{L}) , contains $r_{\mathcal{L}}(\alpha)$ for all $\alpha \in \mathcal{L}^*(E)$ and is also closed under finite intersections and unions, then we say that \mathcal{B} is *accommodating* for (E, \mathcal{L}) . If \mathcal{B} in addition is closed under relative complements, then we say that it is *non-degenerate*.

We are particularly interested in the sets of vertices which are the ranges of words in $\mathcal{L}^*(E)$, so we form

$$\mathcal{E}^- = \{\{v\} : v \in E_{\text{sink}}^0\} \cup \{r(\alpha) : \alpha \in \mathcal{L}^*(E)\}.$$

We then define $\mathcal{E}^{0,-}$ to be the smallest subset of 2^{E^0} which contains \mathcal{E}^- and is accommodating and non-degenerate for (E, \mathcal{L}) . Of course, 2^{E^0} is the largest accommodating and non-degenerate collection of subsets for (E, \mathcal{L}) .

Labelled spaces. A *labelled space* consists of a triple $(E, \mathcal{L}, \mathcal{B})$ where (E, \mathcal{L}) is a labelled graph and \mathcal{B} is accommodating for (E, \mathcal{L}) . If, in addition, \mathcal{B} is non-degenerate, then we say that the labelled space $(E, \mathcal{L}, \mathcal{B})$ is *normal*.

A labelled space $(E, \mathcal{L}, \mathcal{B})$ is *weakly left-resolving* if for every $A, B \in \mathcal{B}$ and every $\alpha \in \mathcal{L}^*(E)$ we have $r(A, \alpha) \cap r(B, \alpha) = r(A \cap B, \alpha)$. If (E, \mathcal{L}) is left-resolving then $(E, \mathcal{L}, \mathcal{B})$ is weakly left-resolving for any accommodating $\mathcal{B} \subseteq 2^{E^0}$.

Let (E, \mathcal{L}) be a labelled graph. For $A \subseteq E^0$ let $\mathcal{L}(AE^1) = \{\mathcal{L}(e) : e \in E^1, s(e) \in A\}$ (the set $\mathcal{L}(AE^1)$ was denoted L_A^1 in [4]).

C^* -algebras of normal labelled spaces. We will now define representations of weakly left-resolving normal labelled spaces, and the C^* -algebra associated with a weakly left-resolving normal labelled space $(E, \mathcal{L}, \mathcal{B})$. Notice that unlike [4], we require $(E, \mathcal{L}, \mathcal{B})$ to be normal. The reason is that if we do not assume that $(E, \mathcal{L}, \mathcal{B})$ is normal, then the C^* -algebra $C^*(E, \mathcal{L}, \mathcal{B})$ associated with $(E, \mathcal{L}, \mathcal{B})$ might not satisfy the Gauge Invariant Uniqueness Theorem as stated in [4, Theorem 5.3] (see [6, Remark 2.5]). In the appendix we will give an example of what can go wrong if $(E, \mathcal{L}, \mathcal{B})$ is not normal and explain how the definition of a representation of $(E, \mathcal{L}, \mathcal{B})$ must be changed in order to obtain a gauge invariant uniqueness theorem for $C^*(E, \mathcal{L}, \mathcal{B})$ when $(E, \mathcal{L}, \mathcal{B})$ is not normal.

Definition 2.1. Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving normal labelled space. A *representation* of $(E, \mathcal{L}, \mathcal{B})$ in a C^* -algebra consists of projections $\{p_A : A \in \mathcal{B}\}$ and partial isometries $\{s_a : a \in \mathcal{A}\}$ with the properties that:

- (i) If $A, B \in \mathcal{B}$, then $p_A p_B = p_{A \cap B}$ and $p_{A \cup B} = p_A + p_B - p_{A \cap B}$, where $p_\emptyset = 0$.
- (ii) If $a \in \mathcal{A}$ and $A \in \mathcal{B}$, then $p_A s_a = s_a p_{r(A,a)}$.
- (iii) If $a, b \in \mathcal{A}$, then $s_a^* s_a = p_{r(a)}$ and $s_a^* s_b = 0$ unless $a = b$.
- (iv) For $A \in \mathcal{B}$ with $\mathcal{L}(AE^1)$ finite and $A \cap B = \emptyset$ for all $B \in \mathcal{B}$ satisfying $B \subseteq E_{\text{sink}}^0$, we have

$$p_A = \sum_{a \in \mathcal{L}(AE^1)} s_a p_{r(A,a)} s_a^*.$$

Remark 2.2. Notice that if the directed graph E contains sinks, then condition (iv) is different from condition (iv) of the original definition [4, Definition 4.1]. The original definition [4, Definition 4.1] was in error since it would lead to degeneracy of the vertex projections for sinks.

Remark 2.3. Notice also that if A is any set in \mathcal{B} with $\mathcal{L}(AE^1)$ finite and $A \cap E_{\text{sink}}^0 \in \mathcal{B}$, then combining (i) and (iv) we obtain the relation

$$p_A = p_{A \cap E_{\text{sink}}^0} + \sum_{a \in \mathcal{L}(AE^1)} s_a p_{r(A,a)} s_a^*,$$

cf. [5, Remark 3.2].

Remark 2.4. Suppose that $\{v\} \in \mathcal{B}$ for all $v \in E_{\text{sink}}^0$. Then for any $A \in \mathcal{B}$, the condition

$$A \cap B = \emptyset \text{ for all } B \in \mathcal{B} \text{ satisfying } B \subseteq E_{\text{sink}}^0$$

that appears in (iv) is equivalent to the condition

$$A \cap E_{\text{sink}}^0 = \emptyset.$$

Definition 2.5. Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving normal labelled space. Then $C^*(E, \mathcal{L}, \mathcal{B})$ is the C^* -algebra generated by a universal representation of $(E, \mathcal{L}, \mathcal{B})$.

The existence of $C^*(E, \mathcal{L}, \mathcal{B})$ is shown by an argument similar to the one given in [4, Theorem 4.5]. The existence of $C^*(E, \mathcal{L}, \mathcal{B})$ will also follow from Theorem 3.8. The universal property of $C^*(E, \mathcal{L}, \mathcal{B})$ allows us to define a strongly continuous action γ of \mathbf{T} on $C^*(E, \mathcal{L}, \mathcal{B})$ called the *gauge action* (see [4, Section 5]).

Remark 2.6. Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving normal labelled space, and let $\{p_A, s_a : A \in \mathcal{B}, a \in \mathcal{A}\}$ be the universal representation of $(E, \mathcal{L}, \mathcal{B})$ that generates $C^*(E, \mathcal{L}, \mathcal{B})$. For $\alpha = \alpha_1 \dots \alpha_n \in \mathcal{L}(E^n)$ (where $n \geq 2$), we let $s_\alpha = s_{\alpha_1} \dots s_{\alpha_n}$. Let $\mathcal{L}^\#(E) = \mathcal{L}^*(E) \cup \{\epsilon\}$ where ϵ is a symbol not belonging to $\mathcal{L}^*(E)$ (ϵ denotes the empty word), and let s_ϵ denote the unit of the multiplier algebra of $C^*(E, \mathcal{L}, \mathcal{B})$. It then follows from [4, Lemma 4.4] that we have

$$C^*(E, \mathcal{L}, \mathcal{B}) = \overline{\text{span}}\{s_\alpha p_A s_\beta^* : \alpha, \beta \in \mathcal{L}^\#(E), A \in \mathcal{B}\}.$$

3. VIEWING LABELLED GRAPH C^* -ALGEBRAS AS C^* -ALGEBRAS OF C^* -CORRESPONDENCES

Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving normal labelled space. In this section we will show how to construct a C^* -correspondence $X(E, \mathcal{L}, \mathcal{B})$ whose Cuntz-Pimsner algebra (see [20] and [37]) is isomorphic to $C^*(E, \mathcal{L}, \mathcal{B})$.

It should be noticed that this construction also works if $(E, \mathcal{L}, \mathcal{B})$ is not normal (see the appendix).

Remark 3.1. In [38] it is shown that for every weakly left-resolving labelled space $(E', \mathcal{L}', \mathcal{B}')$ satisfying $\{\{v\} : v \in (E')_{\text{sink}}^0\} \subseteq \mathcal{B}'$, there exists a C^* -correspondence for which the corresponding Cuntz-Pimsner algebra is isomorphic to $C^*(E', \mathcal{L}', \mathcal{B}')$. Although the construction of the C^* -correspondence given in [38] is similar to the one given below, the coefficient algebra of the C^* -correspondence constructed in [38] does not seem to be useful for obtaining a formula for the K -theory of the C^* -algebra of the labelled space as is done in Section 4.

For each $A \in 2^{E^0}$ we let χ_A denote the function defined on E^0 by

$$\chi_A(v) = \begin{cases} 1 & \text{if } v \in A, \\ 0 & \text{if } v \notin A. \end{cases}$$

We will regard χ_A as an element of the C^* -algebra of bounded functions on E^0 (when we calculate the K -theory we will regard χ_A as an element of the group of functions from E^0 to \mathbb{Z}). Let $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ denote the C^* -subalgebra of the C^* -algebra of bounded functions on E^0 generated by $\{\chi_A : A \in \mathcal{B}\}$.

Lemma 3.2. *Let $\{p_A : A \in \mathcal{B}\}$ be a family of projections in a C^* -algebra \mathcal{X} such that $p_{A \cap B} = p_A p_B$, $p_{A \cup B} = p_A + p_B - p_{A \cap B}$ for all $A, B \in \mathcal{B}$ and $p_\emptyset = 0$. Then there is a unique $*$ -homomorphism $\phi : \mathcal{A}(E, \mathcal{L}, \mathcal{B}) \rightarrow \mathcal{X}$ such that $\phi(\chi_A) = p_A$ for all $A \in \mathcal{B}$.*

Proof. Since $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ is generated by $\{\chi_A : A \in \mathcal{B}\}$, there can be at most one $*$ -homomorphism $\phi : \mathcal{A}(E, \mathcal{L}, \mathcal{B}) \rightarrow \mathcal{X}$ satisfying $\phi(\chi_A) = p_A$ for all $A \in \mathcal{B}$. We show that such a $*$ -homomorphism exists.

Let \mathfrak{F} denote the collection of finite subsets of \mathcal{B} which are closed under relative complements, intersections and unions. Then $\bigcup_{B' \in \mathfrak{F}} \text{span}\{\chi_A : A \in B'\}$ is dense in $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ and so it is enough to prove that for every $B' \in \mathfrak{F}$ there is a $*$ -homomorphism $\phi_{B'} : \text{span}\{\chi_A : A \in B'\} \rightarrow \mathcal{X}$ satisfying $\phi_{B'}(\chi_A) = p_A$ for every $A \in B'$.

If $B' \in \mathfrak{F}$, then, since B' is closed under relative complements, intersections and unions, there is a collection F of mutually disjoint elements of B' such that $\text{span}\{\chi_A : A \in B'\} = \text{span}\{\chi_A :$

$A \in F$ }. It follows that the map $\phi_{\mathcal{B}'} : \text{span}\{\chi_A : A \in \mathcal{B}'\} \rightarrow \mathcal{X}$ defined by

$$\phi_{\mathcal{B}'} \left(\sum_{A \in F} c_A \chi_A \right) = \sum_{A \in F} c_A p_A$$

is a well-defined $*$ -homomorphism which maps χ_A to p_A for every $A \in \mathcal{B}'$. Our result follows. \square

Lemma 3.3. *Let \mathcal{I} be a closed two-sided ideal of $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$. Then we have*

$$\mathcal{I} = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}, \chi_A \in \mathcal{I}\}.$$

Proof. We certainly have $\overline{\text{span}}\{\chi_A : A \in \mathcal{B}, \chi_A \in \mathcal{I}\} \subseteq \mathcal{I}$. To prove the reverse inclusion, suppose that $f \in \mathcal{I}$ and let $\epsilon > 0$. Then there is a finite collection F of mutually disjoint elements of \mathcal{B} and a family $(c_A)_{A \in F}$ of complex numbers such that

$$\left\| f - \sum_{A \in F} c_A \chi_A \right\| < \frac{\epsilon}{2}.$$

Let $q_{\mathcal{I}} : \mathcal{A}(E, \mathcal{L}, \mathcal{B}) \rightarrow \mathcal{A}(E, \mathcal{L}, \mathcal{B})/\mathcal{I}$ be the quotient map. Then

$$\left\| \sum_{A \in F, \chi_A \notin \mathcal{I}} c_A q_{\mathcal{I}}(\chi_A) \right\| = \left\| q_{\mathcal{I}} \left(f - \sum_{A \in F} c_A \chi_A \right) \right\| < \frac{\epsilon}{2},$$

and since $(q_{\mathcal{I}}(\chi_A))_{A \in F, \chi_A \notin \mathcal{I}}$ is a family of mutually orthogonal non-zero projections we must have $|c_A| < \frac{\epsilon}{2}$ for all $A \in F$ such that $\chi_A \notin \mathcal{I}$. It then follows that

$$\left\| f - \sum_{A \in F, \chi_A \in \mathcal{I}} c_A \chi_A \right\| < \epsilon$$

from which we may deduce that $f \in \overline{\text{span}}\{\chi_A : A \in \mathcal{B}, \chi_A \in \mathcal{I}\}$. Our result follows. \square

For each $a \in \mathcal{A}$, let X_a be the ideal of $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ generated by $\chi_{r(a)}$ so that $f \in X_a$ if and only if $f(v) = 0$ for all $v \in E^0 \setminus r(a)$. Since X_a is an ideal, it is straightforward to see that X_a is a right Hilbert $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ -module with inner product defined by $\langle f, g \rangle = f^*g$ and right action given by the usual multiplication in $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$.

We let $X(E, \mathcal{L}, \mathcal{B})$ be the right Hilbert $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ -module $\bigoplus_{a \in \mathcal{A}} X_a$. To turn $X(E, \mathcal{L}, \mathcal{B})$ into a C^* -correspondence we need to specify a left action of $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ on $X(E, \mathcal{L}, \mathcal{B})$, that is a $*$ -homomorphism $\phi : \mathcal{A}(E, \mathcal{L}, \mathcal{B}) \rightarrow \mathcal{L}(X(E, \mathcal{L}, \mathcal{B}))$ where $\mathcal{L}(X(E, \mathcal{L}, \mathcal{B}))$ denotes the C^* -algebra of adjointable operators on $X(E, \mathcal{L}, \mathcal{B})$ (see, for example, [20]).

Lemma 3.4. *For each $a \in \mathcal{A}$ there is a unique $*$ -homomorphism $\phi_a : \mathcal{A}(E, \mathcal{L}, \mathcal{B}) \rightarrow X_a$ satisfying $\phi_a(\chi_A) = \chi_{r(A,a)}$ for every $A \in \mathcal{B}$.*

Proof. This follows from Lemma 3.2 with $\mathcal{X} = X_a$ and $p_A = \chi_{r(A,a)}$ for $A \in \mathcal{B}$. \square

Remark 3.5. If the labelled graph (E, \mathcal{L}) is left-resolving, then we have

$$\phi_a(f)(v) = \begin{cases} f(w) & \text{if } s(r^{-1}(v) \cap \mathcal{L}^{-1}(a)) = \{w\} \\ 0 & \text{if } s(r^{-1}(v) \cap \mathcal{L}^{-1}(a)) = \emptyset \end{cases}$$

for all $a \in \mathcal{A}$, $v \in E^0$ and $f \in \mathcal{A}(E, \mathcal{L}, \mathcal{B})$.

Lemma 3.6. *Let $X(E, \mathcal{L}, \mathcal{B})$ be the right Hilbert $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ -module $\bigoplus_{a \in \mathcal{A}} X_a$.*

(i) For each $f \in \mathcal{A}(E, \mathcal{L}, \mathcal{B})$ the map

$$(1) \quad \varphi(f) : (x_a)_{a \in \mathcal{A}} \mapsto (\phi_a(f)x_a)_{a \in \mathcal{A}}$$

is an adjointable operator on $X(E, \mathcal{L}, \mathcal{B})$. The formula (1) defines a $*$ -homomorphism $\varphi : \mathcal{A}(E, \mathcal{L}, \mathcal{B}) \rightarrow \mathcal{L}(X(E, \mathcal{L}, \mathcal{B}))$. Hence $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$ is a C^* -correspondence over $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$.

(ii) For $a \in \mathcal{A}$ let $e_a = (\delta_{a,b}\chi_{r(a)})_{b \in \mathcal{A}} \in X(E, \mathcal{L}, \mathcal{B})$, where $\delta_{a,b}$ is the Kronecker delta function. Then we have

$$X(E, \mathcal{L}, \mathcal{B}) = \overline{\text{span}}_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})} \{e_a : a \in \mathcal{A}\}.$$

Proof. Since the effect of $\varphi(f)$ is to multiply each term in $X(E, \mathcal{L}, \mathcal{B})$ by $\phi_a(f)$, one checks that (1) defines an adjointable operator with $\varphi(f)^* = \varphi(f^*)$. It follows that $f \mapsto \varphi(f)$ defines a $*$ -homomorphism φ from $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ to $\mathcal{L}(X(E, \mathcal{L}, \mathcal{B}))$.

The second statement holds since, for each $x = (x_a)_{a \in \mathcal{A}} \in X(E, \mathcal{L}, \mathcal{B})$ and each $\varepsilon > 0$, we have $\|x - \sum_{a \in F} e_a x_a\| < \varepsilon$ for some finite subset F of \mathcal{A} . \square

Let $\mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))$ denote the ideal of $\mathcal{L}(X(E, \mathcal{L}, \mathcal{B}))$ consisting of generalized compact operators (see, for example, [20]). In order to define the Cuntz-Pimsner algebra $\mathcal{O}_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}$ associated with $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$, we must first characterise the ideal

$$J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)} = \varphi^{-1}(\mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))) \cap \ker(\varphi)^\perp$$

of $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ (cf. [20, Definition 3.2]). To do this, we let

$$(2) \quad \mathcal{B}_J := \{A \in \mathcal{B} : \mathcal{L}(AE^1) \text{ is finite and } A \cap B = \emptyset \text{ for all } B \in \mathcal{B} \text{ satisfying } B \subseteq E_{\text{sink}}^0\}.$$

Lemma 3.7. *The ideal $J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}$ of $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ is given by*

$$J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)} = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}_J\}.$$

Proof. By Lemma 3.3 it is enough to prove that for $A \in \mathcal{B}$, we have $\chi_A \in J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}$ if and only if $A \in \mathcal{B}_J$. Since $\ker(\varphi) = \bigcap_{a \in \mathcal{A}} \ker(\phi_a)$, it follows from Lemma 3.3 that

$$\ker(\varphi) = \overline{\text{span}}\{\chi_B : B \in \mathcal{B}, r(B, a) = \emptyset \text{ for all } a \in \mathcal{A}\}.$$

Since $r(B, a) = \emptyset$ for all $a \in \mathcal{A}$, if and only if $B \subseteq E_{\text{sink}}^0$, it follows that

$$\ker(\varphi)^\perp = \overline{\text{span}}\{A \in \mathcal{B} : A \cap B = \emptyset \text{ for all } B \in \mathcal{B} \text{ satisfying } B \subseteq E_{\text{sink}}^0\}.$$

We claim that for each $\eta \in \mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))$ the set $\{a \in \mathcal{A} : \|\langle e_a, \eta(e_a) \rangle\| \geq 1\}$ is finite. It suffices to check our claim for $\eta = \theta_{x,y}$ since these elements form a spanning set for $\mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))$. For each $a_0 \in \mathcal{A}$ and $x = (x_a), y = (y_a) \in X(E, \mathcal{L}, \mathcal{B})$ one checks that $\langle e_{a_0}, \theta_{x,y} e_{a_0} \rangle = x_{a_0}^* y_{a_0}$, and since $\|x_a^* y_a\| \geq 1$ for only a finite number of a 's, our claim follows. So, if $A \in \mathcal{B}$ and $\mathcal{L}(AE^1)$ is infinite, then $\varphi(\chi_A) \notin \mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))$.

Conversely, if $A \in \mathcal{B}$ and $\mathcal{L}(AE^1)$ is finite, then we have

$$(3) \quad \varphi(\chi_A) = \sum_{a \in \mathcal{L}(AE^1)} \theta_{e_a, e_a \chi_{r(A,a)}} \in \mathcal{K}(X(E, \mathcal{L}, \mathcal{B})).$$

Thus, if $A \in \mathcal{B}$, then $\varphi(\chi_A) \in \mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))$ if and only if $\mathcal{L}(AE^1)$ is finite. The result follows by definition of $J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}$ and \mathcal{B}_J . \square

Recall from [20, Definition 3.4] that a representation of the C^* -correspondence $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$ on a C^* -algebra \mathcal{X} consists of a pair (π, t) where $\pi : \mathcal{A}(E, \mathcal{L}, \mathcal{B}) \rightarrow \mathcal{X}$ is a $*$ -homomorphism and $t : X(E, \mathcal{L}, \mathcal{B}) \rightarrow \mathcal{X}$ is a linear map satisfying

- (1) $t(x)^* t(y) = \pi(\langle x, y \rangle)$ for $x, y \in X(E, \mathcal{L}, \mathcal{B})$ and
- (2) $\pi(f) t(x) = t(\varphi(f)x)$ for $f \in \mathcal{A}(E, \mathcal{L}, \mathcal{B})$ and $x \in X(E, \mathcal{L}, \mathcal{B})$.

Following [20, Definition 2.3] we define a $*$ -homomorphism $\psi_t : \mathcal{K}(X(E, \mathcal{L}, \mathcal{B})) \rightarrow \mathcal{X}$ by

$$\psi_t(\theta_{x,y}) = t(x)t(y)^* \text{ for } x, y \in X(E, \mathcal{L}, \mathcal{B}).$$

Moreover, such a representation is called *covariant* if, in addition, we have $\pi(f) = \psi_t(\varphi(f))$ for all $f \in J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}$. The *Cuntz-Pimsner algebra* $\mathcal{O}_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}$ of $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$ is defined in [20] (see also [37]) to be the C^* -algebra generated by a universal covariant representations of $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$.

Theorem 3.8. *Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving normal labelled space. Then there is a one-to-one correspondence between covariant representations of $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$ and representations of $(E, \mathcal{L}, \mathcal{B})$ that takes a covariant representation (π, t) of $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$ to the representation $\{\pi(\chi_A), t(e_a) : A \in \mathcal{B}, a \in \mathcal{A}\}$ of $(E, \mathcal{L}, \mathcal{B})$, and so $\mathcal{O}_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)} \cong C^*(E, \mathcal{L}, \mathcal{B})$.*

Proof. Let (π, t) be a covariant representation of $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$. We claim that

$$\{\pi(\chi_A), t(e_a) : A \in \mathcal{B}, a \in \mathcal{A}\}$$

is a representation of $(E, \mathcal{L}, \mathcal{B})$.

It is straightforward to check that $\{\pi(\chi_A), t(e_a) : A \in \mathcal{B}, a \in \mathcal{A}\}$ satisfies (i)–(iii) of Definition 2.1. We will now show that $\{\pi(\chi_A), t(e_a) : A \in \mathcal{B}, a \in \mathcal{A}\}$ also satisfies (iv) of Definition 2.1. Assume $A \in \mathcal{B}$, $\mathcal{L}(AE^1)$ is finite and that $A \cap B = \emptyset$ for any $B \in \mathcal{B}$ satisfying $B \subseteq E_{\text{sink}}^0$. It follows from Lemma 3.7 that $\chi_A \in J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}$. By covariance and Equation (3) we have

$$\pi(\chi_A) = \psi_t(\phi(\chi_A)) = \psi_t\left(\sum_{a \in \mathcal{L}(AE^1)} \theta_{e_a, e_a \chi_{r(A,a)}}\right) = \sum_{a \in \mathcal{L}(AE^1)} t(e_a)\pi(\chi_{r(A,a)})t(e_a)^*.$$

Thus $\{\pi(\chi_A), t(e_a) : A \in \mathcal{B}, a \in \mathcal{A}\}$ satisfies (iv) of Definition 2.1, and is therefore a representation of $(E, \mathcal{L}, \mathcal{B})$.

On the other hand, suppose that $\{P_A, S_a : A \in \mathcal{B}, a \in \mathcal{A}\}$ is a representation of $(E, \mathcal{L}, \mathcal{B})$ in a C^* -algebra \mathcal{X} . By definition $\mathcal{A}(E, \mathcal{L}, \mathcal{B}) = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}\}$ and by Lemma 3.6, we have $X(E, \mathcal{L}, \mathcal{B}) = \overline{\text{span}}_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})}\{e_a : a \in \mathcal{L}(E^1)\}$. Hence there can at most be one representation (π, t) of $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$ such that $\pi(\chi_A) = P_A$ for $A \in \mathcal{B}$ and $t(e_a) = S_a$ for $a \in \mathcal{A}$. We now construct such a representation.

By Lemma 3.2 there is a unique $*$ -homomorphism $\pi : \mathcal{A}(E, \mathcal{L}, \mathcal{B}) \rightarrow \mathcal{X}$ with $\pi(\chi_A) = p_A$ for $A \in \mathcal{B}$. We now construct $t : (X(E, \mathcal{L}, \mathcal{B}), \varphi) \rightarrow \mathcal{X}$. Given a finite subset F of \mathcal{A} and a family $(f_a)_{a \in F}$ of functions with $f_a \in X_a$ for each $a \in F$, define

$$t\left(\sum_{a \in F} e_a f_a\right) = \sum_{a \in F} S_a \pi(f_a) \in \mathcal{X}.$$

If $(f_a)_{a \in F}$ and $(g_a)_{a \in F}$ are families of functions with $f_a, g_a \in X_a$ for each $a \in F$, then

$$t\left(\sum_{a \in F} e_a f_a\right)^* t\left(\sum_{a \in F} e_a g_a\right) = \left(\sum_{a \in F} S_a \pi(f_a)\right)^* \left(\sum_{a \in F} S_a \pi(g_a)\right) = \sum_{a \in F} \pi(f_a)^* P_{r(a)} \pi(g_a)$$

by (iii) of Definition 2.1. Since $P_{r(a)} = \pi(\chi_{r(a)})$ we have

$$t\left(\sum_{a \in F} e_a f_a\right)^* t\left(\sum_{a \in F} e_a g_a\right) = \sum_{a \in F} \pi(f_a^* g_a) = \pi\left(\left\langle \sum_{a \in F} e_a f_a, \sum_{a \in F} e_a g_a \right\rangle\right).$$

Thus t extends to a linear map from $X(E, \mathcal{L}, \mathcal{B})$ to \mathcal{X} which satisfies $t(x)^*t(y) = \pi(\langle x, y \rangle)$ for $x, y \in X(E, \mathcal{L}, \mathcal{B})$. We claim that for $x \in X(E, \mathcal{L}, \mathcal{B})$ and $f \in \mathcal{A}(E, \mathcal{L}, \mathcal{B})$ we have $\pi(a)t(x) = t(\varphi(a)x)$.

Let $A \in \mathcal{B}$ and $a \in \mathcal{A}$, then

$$(4) \quad \pi(\chi_A)S_a = P_A S_a = S_a P_{r(A,a)} = S_a \pi(\chi_{r(A,a)}) = S_a \pi(\phi_a(\chi_A)).$$

Since $\mathcal{A}(E, \mathcal{L}, \mathcal{B}) = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}\}$, it follows from Equation (4) that

$$\pi(f)t(e_a) = \pi(f)S_a = S_a \pi(\phi_a(f)) = t(e_a)\pi(\phi_a(f)) = t(\varphi(f)e_a)$$

for all $f \in \mathcal{A}(E, \mathcal{L}, \mathcal{B})$ and all $a \in \mathcal{A}$. Our claim then follows from Lemma 3.6 (ii). Hence we have constructed a representation (π, t) of $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$. It remains to check covariance.

Let $A \in \mathcal{B}_J$, then by Equation (3) and (4) we have

$$\begin{aligned} \pi(\chi_A) &= P_A = \sum_{a \in \mathcal{L}(AE^1)} S_a P_{r(A,a)} S_a^* = \sum_{a \in \mathcal{L}(AE^1)} t(e_a)\pi(\chi_{r(A,a)})t(e_a)^* \\ &= \sum_{a \in \mathcal{L}(AE^1)} \psi_t(\theta_{e_a, e_a \phi_a(\chi_A)}) = \psi_t(\varphi(\chi_A)). \end{aligned}$$

By Lemma 3.7 we have $J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)} = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}_J\}$ and it follows that $\pi(f) = \psi_t(\varphi(f))$ for all $f \in J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}$. Thus (π, t) is a covariant representation of $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$.

The final statement follows by the universal nature of both algebras. \square

In the appendix, it is explained how Theorem 3.8 can be generalised to the case where $(E, \mathcal{L}, \mathcal{B})$ is not normal (see Remark A.7).

From Theorem 3.8 we may deduce a number of corollaries. The second of these is a generalization of the Gauge Invariant Uniqueness Theorem found in [6].

Corollary 3.9. *Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving normal labelled space. Then there exists an injective $*$ -homomorphism $\iota_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})} : \mathcal{A}(E, \mathcal{L}, \mathcal{B}) \rightarrow C^*(E, \mathcal{L}, \mathcal{B})$ such that $\iota_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})}(\chi_A) = p_A$ for every $A \in \mathcal{B}$.*

Proof. Follows from Theorem 3.8 and [20, Proposition 4.11]. \square

In the appendix, it is explained how Corollary 3.9 can be generalised to the case where $(E, \mathcal{L}, \mathcal{B})$ is not normal (see Remark A.7).

Corollary 3.10. *Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving normal labelled space. Let $\{p_A, s_a : A \in \mathcal{B}, a \in \mathcal{A}\}$ be the universal representation of $(E, \mathcal{L}, \mathcal{B})$ that generates $C^*(E, \mathcal{L}, \mathcal{B})$, let $\{q_A, t_a : A \in \mathcal{B}, a \in \mathcal{A}\}$ be a representation of $(E, \mathcal{L}, \mathcal{B})$ in a C^* -algebra \mathcal{X} and let π be the unique $*$ -homomorphism from $C^*(E, \mathcal{L}, \mathcal{B})$ to \mathcal{X} that maps each p_A to q_A and each s_a to t_a . Then π is injective if and only if q_A is non-zero whenever $A \neq \emptyset$, and for each $z \in \mathbb{T}$ there exists a $*$ -homomorphism $\beta_z : C^*(\{q_A, t_a : A \in \mathcal{B}, a \in \mathcal{A}\}) \rightarrow C^*(\{q_A, t_a : A \in \mathcal{B}, a \in \mathcal{A}\})$ such that $\beta_z(q_A) = q_A$ and $\beta_z(t_a) = zt_a$ for $A \in \mathcal{B}$ and $a \in \mathcal{A}$.*

Proof. Follows from Lemma 3.3, Theorem 3.8, and [20, Theorem 6.4]. \square

In the appendix, Corollary 3.10 is generalised to the case where $(E, \mathcal{L}, \mathcal{B})$ is not normal (see Corollary A.8).

Corollary 3.11. *Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving normal labelled space. Then $C^*(E, \mathcal{L}, \mathcal{B})$ is nuclear. If, in addition, \mathcal{B} and \mathcal{A} are countable, then $C^*(E, \mathcal{L}, \mathcal{B})$ satisfies the Universal Coefficient Theorem of [42].*

Proof. The first part follows from [20, Corollary 7.4] and the fact that $J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}$ and $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ are commutative and thus nuclear. If, in addition, \mathcal{B} and \mathcal{A} are countable, then $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ and $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$ are separable. It then follows from [20, Proposition 8.8] that $C^*(E, \mathcal{L}, \mathcal{B})$ satisfies the Universal Coefficient Theorem of [42]. \square

In the appendix, it is explained how Corollary 3.11 can be generalised to the case where $(E, \mathcal{L}, \mathcal{B})$ is not normal (see Remark A.7).

4. K -THEORY

We now compute the K -theory of $C^*(E, \mathcal{L}, \mathcal{B})$ when $(E, \mathcal{L}, \mathcal{B})$ is a weakly left-resolving normal labelled space (see the appendix for a procedure for computing the K -theory when $(E, \mathcal{L}, \mathcal{B})$ is not normal). Our approach is to use [20, Theorem 8.6] which involves detailed knowledge of the map $[X] : K_*(J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}) \rightarrow K_*(\mathcal{A}(E, \mathcal{L}, \mathcal{B}))$. Since we will need to work explicitly with this map, we will now recall its definition in detail (cf. [20, Definition 8.3]).

Let $D_{X(E, \mathcal{L}, \mathcal{B})}$ denote the linking algebra $\mathcal{K}(X(E, \mathcal{L}, \mathcal{B}) \oplus \mathcal{A}(E, \mathcal{L}, \mathcal{B}))$. Following [20, Definition B.1] we denote the natural embedding of $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ into $D_{X(E, \mathcal{L}, \mathcal{B})}$ by $\iota_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})}$ and the natural embedding of $\mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))$ into $D_{X(E, \mathcal{L}, \mathcal{B})}$ by $\iota_{\mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))}$. It is shown in [20, Proposition B.3] that the map $(\iota_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})})_* : K_*(\mathcal{A}(E, \mathcal{L}, \mathcal{B})) \rightarrow K_*(D_{X(E, \mathcal{L}, \mathcal{B})})$ induced by $\iota_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})}$ is an isomorphism. The map $[X] : K_*(J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}) \rightarrow K_*(\mathcal{A}(E, \mathcal{L}, \mathcal{B}))$ is then defined to be $(\iota_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})})_*^{-1} \circ (\iota_{\mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))})_* \circ (\varphi|_{J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}})_*$.

Lemma 4.1. *Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving normal labelled space. Let $a \in \mathcal{A}$ and $A \in \mathcal{B}$ such that $A \subseteq r(a)$. Then for $[\chi_A]_0 \in K_0(\mathcal{A}(E, \mathcal{L}, \mathcal{B}))$ and $[\theta_{e_a, e_a \chi_A}]_0 \in K_0(\mathcal{K}(X(E, \mathcal{L}, \mathcal{B})))$ we have*

$$(\iota_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})})_*([\chi_A]_0) = (\iota_{\mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))})_*([\theta_{e_a, e_a \chi_A}]_0)$$

in $K_0(D_{X(E, \mathcal{L}, \mathcal{B})})$.

Proof. Let v be the function that maps $(\eta, f) \in X(E, \mathcal{L}, \mathcal{B}) \times \mathcal{A}(E, \mathcal{L}, \mathcal{B})$ to $(e_a \chi_A f, 0) \in X(E, \mathcal{L}, \mathcal{B}) \times \mathcal{A}(E, \mathcal{L}, \mathcal{B})$. One checks that $v \in \mathcal{K}(X(E, \mathcal{L}, \mathcal{B}) \oplus \mathcal{A}(E, \mathcal{L}, \mathcal{B}))$, that $vv^* = \iota_{\mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))}(\theta_{e_a, e_a \chi_A})$ and $v^*v = \iota_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})}(\chi_A)$. It follows that the elements $\iota_{\mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))}(\theta_{e_a, e_a \chi_A})$ and $\iota_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})}(\chi_A)$ are equivalent in $K_0(D_{X(E, \mathcal{L}, \mathcal{B})})$. \square

Since $\mathcal{A}(E, \mathcal{L}, \mathcal{B}) = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}\}$ and $J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)} = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}_J\}$, it follows that there is an isomorphism from $\text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}\}$ to $K_0(\mathcal{A}(E, \mathcal{L}, \mathcal{B}))$ which for each $A \in \mathcal{B}$ takes χ_A to $[\chi_A]_0$, and an isomorphism from $\text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}_J\}$ to $K_0(J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)})$ which for each $A \in \mathcal{B}_J$ takes χ_A to $[\chi_A]_0$. We will simply identify $K_0(\mathcal{A}(E, \mathcal{L}, \mathcal{B}))$ with $\text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}\}$, $K_0(J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)})$ with $\text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}_J\}$ and each $[\chi_A]_0$ with χ_A .

Lemma 4.2. *Identifying $K_0(\mathcal{A}(E, \mathcal{L}, \mathcal{B}))$ with $\text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}\}$ and $K_0(J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)})$ with $\text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}_J\}$, the homomorphism $[X] : K_0(J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}) \rightarrow K_0(\mathcal{A}(E, \mathcal{L}, \mathcal{B}))$ induces the map $\Phi : \text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}_J\} \rightarrow \text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}\}$ determined by*

$$(5) \quad \chi_A \mapsto \sum_{a \in \mathcal{L}(AE^1)} \chi_{r(A, a)} \text{ for } A \in \mathcal{B}_J.$$

Proof. Let $A \in \mathcal{B}_J$. Since $\phi_a(\chi_A) = \chi_{r(A, a)}$ is a projection in $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$, we have $\varphi(\chi_A) = \sum_{a \in \mathcal{L}(AE^1)} \theta_{e_a, e_a \chi_{r(A, a)}}$ from Equation (3). It follows from Lemma 4.1 that

$$\begin{aligned} (\iota_{\mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))})_* \circ (\varphi|_{J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}})_*([\chi_A]_0) &= (\iota_{\mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))})_* \left(\left[\sum_{a \in \mathcal{L}(AE^1)} \theta_{e_a, e_a \chi_{r(A, a)}} \right]_0 \right) \\ &= \sum_{a \in \mathcal{L}(AE^1)} (\iota_{\mathcal{K}(X(E, \mathcal{L}, \mathcal{B}))})_*([\theta_{e_a, e_a \chi_{r(A, a)}}]_0) \\ &= \sum_{a \in \mathcal{L}(AE^1)} (\iota_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})})_*([\chi_{r(A, a)}]_0), \end{aligned}$$

from which Equation (5) follows using the given identifications and the definition of $[X]$. \square

If $\mathcal{B} \subseteq 2^{E^0}$ is uncountable, then $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ is not separable. However, it is still locally finite dimensional. Such nonseparable AF algebras were considered in [21].

Lemma 4.3. *Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving normal labelled space. Then $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ and $J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}$ are locally finite dimensional algebras with $K_1(\mathcal{A}(E, \mathcal{L}, \mathcal{B})) = K_1(J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}) = 0$.*

Proof. We prove the result for $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ and note that the proof for $J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}$ is similar. The first statement follows by the argument given in the proof of [4, Theorem 5.3]. To prove that $K_1(\mathcal{A}(E, \mathcal{L}, \mathcal{B})) = 0$, we argue as follows¹. By adding E^0 to \mathcal{B} (if necessary) we may assume that $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ is unital. Let u be a unitary in $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$, then given $0 < \epsilon < 1$ there is a unitary u' in $\text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}\}$ such that $\|u - u'\| < \epsilon$. Using the local finite dimensionality of $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ one can show that u' belongs to a finite dimensional subalgebra U of $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ which contains the identity $1_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})}$. Since the unitary group of a finite dimensional C^* -algebra is connected there is a continuous path of unitaries from u' to $1_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})} = \chi_{E^0}$ which remains continuous when we consider U as a subalgebra of $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$. Since $\|u - u'\| < \epsilon < 1$ there is a continuous path of unitaries from u to u' and hence from u to $1_{\mathcal{A}(E, \mathcal{L}, \mathcal{B})}$. The result now follows. \square

Theorem 4.4. *Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving normal labelled space. Let $(1 - \Phi)$ be the linear map from $\text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}_J\}$ to $\text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}\}$ given by*

$$(1 - \Phi)(\chi_A) = \chi_A - \sum_{a \in \mathcal{L}(AE^1)} \chi_{r(A, a)}, \quad A \in \mathcal{B}_J.$$

Then $K_1(C^(E, \mathcal{L}, \mathcal{B})) \cong \ker(1 - \Phi)$, and $K_0(C^*(E, \mathcal{L}, \mathcal{B})) \cong \text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}\} / \text{Im}(1 - \Phi)$ via $[p_A]_0 \mapsto \chi_A + \text{Im}(1 - \Phi)$ for $A \in \mathcal{B}$.*

Proof. The result follows by [20, Theorem 8.6], Theorem 3.8, Lemma 4.2 and the fact that $K_1(\mathcal{A}(E, \mathcal{L}, \mathcal{B})) = K_1(J_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}) = 0$ from Lemma 4.3. \square

In the appendix, Theorem 4.4 is generalises to the case where $(E, \mathcal{L}, \mathcal{B})$ is not normal (see Theorem A.9).

5. EXAMPLES

In this section we show that Theorem 4.4 unifies the K -theory formulas for several interesting classes of C^* -algebras.

Example 1 – Graph algebras. Let E be a directed graph. Following [4, Examples 3.3 and 4.3 (i)] we may realise $C^*(E)$ as the C^* -algebra of a left-resolving normal labelled space in the following manner. We take the trivial labelling $\mathcal{L}_t : E^1 \rightarrow E^1$ defined for $e \in E^1$ by $\mathcal{L}_t(e) = e$. Then (E, \mathcal{L}) is a left-resolving labelled graph and $\mathcal{E}^{0,-}$ is the set of all finite subsets of E^0 . Since (E, \mathcal{L}) is left-resolving, it follows that the labelled space $(E, \mathcal{L}_t, \mathcal{E}^{0,-})$, in addition to being normal, is weakly left-resolving. By [4, Proposition 5.1 (i)] there is an isomorphism $\phi : C^*(E) \rightarrow C^*(E, \mathcal{L}_t, \mathcal{E}^{0,-})$ that maps P_v to $p_{\{v\}}$ and S_e to s_e where $\{S_e, P_v : e \in E^1, v \in E^0\}$ and $\{s_e, p_A : e \in E^1, A \in \mathcal{E}^{0,-}\}$ are the canonical generators of $C^*(E)$ and $C^*(E, \mathcal{L}_t, \mathcal{E}^{0,-})$, respectively.

Let $E_{ns}^0 = \{v \in E^0 : 0 < |s^{-1}(v)| < \infty\}$ denote the set of nonsingular vertices in E^0 . For each $v \in E_{ns}^0$ we let $e_v \in \bigoplus_{w \in E_{ns}^0} \mathbb{Z}$ be defined by $e_v = (\delta_{v,w})_{w \in E_{ns}^0}$. We similarly define $e_v \in \bigoplus_{v \in E^0} \mathbb{Z}$ for each $v \in E^0$.

Note that $\mathcal{E}_J^{0,-}$ is precisely the collection of all finite subsets of E_{ns}^0 . One checks that there are isomorphisms $\psi : \text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}\} \rightarrow \bigoplus_{v \in E^0} \mathbb{Z}$ and $\psi_J : \text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{E}_J^{0,-}\} \rightarrow \bigoplus_{v \in E_{ns}^0} \mathbb{Z}$ such that for all $v \in E^0$ (respectively $v \in E_{ns}^0$) we have $\psi(\chi_v) = e_v$ (respectively $\psi_J(\chi_v) = e_v$).

¹Thanks to Iain Raeburn for showing us how to do this.

For $v \in E_{ns}^0$ we have $\Phi(\chi_v) = \sum_{f \in s^{-1}\{v\}} \chi_{r(f)}$ and we may now deduce the following Corollary from Theorem 4.4.

Corollary 5.1 (cf. [14, Theorem 3.1]). *Let E be a directed graph. With notation as above, define a linear map $(1 - \Phi) : \bigoplus_{v \in E_{ns}^0} \mathbb{Z} \rightarrow \bigoplus_{v \in E^0} \mathbb{Z}$ by*

$$(1 - \Phi)(e_v) = e_v - \sum_{f \in s^{-1}\{v\}} e_{r(f)}, \quad v \in E_{ns}^0.$$

Then $K_1(C^(E))$ is isomorphic to $\ker(1 - \Phi)$ and there exists an isomorphism from $K_0(C^*(E))$ to $\text{coker}(1 - \Phi)$ which maps $[P_v]_0$ to $e_v + \text{Im}(1 - \Phi)$ for each $v \in E^0$.*

Example 2 – Ultragraph algebras. For the definition and properties of an ultragraph see [39]. Following [4, Examples 3.3 and 4.3 (ii)] we may realise an ultragraph as a labelled graph as follows: Let $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ be an ultragraph. We define a directed graph $E_{\mathcal{G}} = (E_{\mathcal{G}}^0, E_{\mathcal{G}}^1, r', s')$ by putting $E_{\mathcal{G}}^0 = G^0$, $E_{\mathcal{G}}^1 = \{(e, w) : e \in \mathcal{G}^1, w \in r(e)\}$, $r'(e, w) = w$ and $s'(e, w) = s(e)$. Setting $\mathcal{A} = \mathcal{G}^1$ we may define a labelling map $\mathcal{L}_{\mathcal{G}} : E^1 \rightarrow \mathcal{A}$ by $\mathcal{L}_{\mathcal{G}}(e, w) = e$. If we let \mathcal{G}^0 be as defined in [22, Definition 2.4], then $(E_{\mathcal{G}}, \mathcal{L}_{\mathcal{G}}, \mathcal{G}^0)$ is a normal weakly left-resolving labelled space. As in [4, Proposition 5.1 (ii)], using the gauge invariant uniqueness theorem [39, Theorem 6.8] there is an isomorphism $\phi : C^*(\mathcal{G}) \rightarrow C^*(E_{\mathcal{G}}, \mathcal{L}_{\mathcal{G}}, \mathcal{G}^0)$ that maps P_A to p_A and S_e to s_e where $\{S_e, P_A\}$ and $\{s_e, p_A\}$ are the canonical generators of $C^*(\mathcal{G})$ and $C^*(E_{\mathcal{G}}, \mathcal{L}_{\mathcal{G}}, \mathcal{G}^0)$, respectively.

The following definitions are taken from [22]. Let $\mathfrak{A}_{\mathcal{G}}$ denote the C^* -subalgebra of $\ell^\infty(G^0)$ generated by the point masses $\{\delta_v : v \in G^0\}$ and the characteristic functions $\{\chi_{r(e)} : e \in \mathcal{G}^1\}$ and let $Z_{\mathcal{G}}$ denote the algebraic subalgebra of $\ell^\infty(G^0, \mathbb{Z})$ generated by these functions. Let $G_{rg}^0 \subseteq G^0$ denote the set of vertices $v \in G^0$ satisfying $0 < |s^{-1}(v)| < \infty$. For $v \in G_{rg}^0$ let $e_v \in \bigoplus_{v \in G_{rg}^0} \mathbb{Z}$ be defined by $e_v = (\delta_{v,w})_{w \in G_{rg}^0}$ and define $e_v \in Z_{\mathcal{G}} \subseteq \ell^\infty(G_0, \mathbb{Z})$ similarly for $v \in E^0$.

It follows from [22, Proposition 2.6 and 2.20] that $Z_{\mathcal{G}} = K_0(\mathfrak{A}_{\mathcal{G}}) = \text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{G}^0\}$. One checks that $A \in \mathcal{G}^0$ belongs to \mathcal{G}_j^0 if and only if A is a finite subset of G_{rg}^0 . It then follows that there is an isomorphism from $\bigoplus_{v \in G_{rg}^0} \mathbb{Z}$ to $\text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{G}_j^0\}$ which maps e_v to $\chi_{\{v\}}$ for every $v \in G_{rg}^0$. If $v \in G_{rg}^0$, then $\Phi(\chi_{\{v\}}) = \sum_{f \in s^{-1}\{v\}} \chi_{r(f)}$, and we may deduce the following Corollary to Theorem 4.4.

Corollary 5.2 (cf. [22, Theorem 5.4]). *Let $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ be an ultragraph. With notation as above, define a linear map $(1 - \Phi) : \bigoplus_{v \in G_{rg}^0} \mathbb{Z} \rightarrow Z_{\mathcal{G}}$ by*

$$(1 - \Phi)(e_v) = e_v - \sum_{f \in s^{-1}\{v\}} \chi_{r(f)}, \quad v \in G_{rg}^0.$$

Then $K_1(C^(\mathcal{G}))$ is isomorphic to $\ker(1 - \Phi)$ and there is an isomorphism from $K_0(C^*(\mathcal{G}))$ to $\text{coker}(1 - \Phi)$ which maps $[P_A]_0$ to $\chi_A + \text{Im}(1 - \Phi)$ for each $A \in \mathcal{G}^0$.*

Example 3 – Matsumoto algebras. We are interested in two C^* -algebras which Matsumoto associates to a (two-sided) shift space Λ over a finite alphabet \mathcal{A} . The first, which we denote \mathcal{O}_{Λ^*} , was described in [27], [28], [29], [30], [32] and [33]. It follows from an argument similar to the one used in [4, Theorem 6.3] (but using Corollary 3.10 instead of [4, Theorem 5.3]) that $\mathcal{O}_{\Lambda^*} \cong C^*(E_{\Lambda^*}, \mathcal{L}_{\Lambda^*}, \mathcal{E}_{\Lambda^*}^{0,-})$ where $(E_{\Lambda^*}, \mathcal{L}_{\Lambda^*})$ is the predecessor graph (or past set cover) of Λ , as defined in [4, Examples 3.3] (vii) (see also [2, Definition 5.2]).

The second, which we denote \mathcal{O}_{Λ} was described in [9] and [35]. Under a technical hypothesis, analogous to Cuntz and Krieger's condition (I), it follows from an argument similar to the one used in [4, Theorem 6.3] (again using Corollary 3.10 instead of [4, Theorem 5.3]) that $\mathcal{O}_{\Lambda} \cong C^*(E_{\Lambda}, \mathcal{L}_{\Lambda}, \mathcal{E}_{\Lambda}^{0,-})$ where $(E_{\Lambda}, \mathcal{L}_{\Lambda})$ is the left Krieger cover of Λ , as defined in [23, Section 2] (see also [4, Examples 3.3 (vi)] and [2, Definition 2.11]).

Let $\mathcal{A}^* \setminus \{\epsilon\}$ denote the set of non-empty finite words over \mathcal{A} . Let $\{S_a : a \in \mathcal{A}\}$ be the generators of \mathcal{O}_Λ (respectively \mathcal{O}_{Λ^*}). For $u = u_1u_2 \dots u_m \in \mathcal{A}^* \setminus \{\epsilon\}$, let $S_u := S_{u_1}S_{u_2} \dots S_{u_m}$.

Corollary 5.3 (cf. [28, Theorem 4.9] and [34, Lemma 2.7]). *Let Λ be a (two-sided) shift space over a finite alphabet \mathcal{A} . Let $(1 - \Phi) : \text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{E}_{\Lambda^*}^{0,-}\} \rightarrow \text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{E}_{\Lambda^*}^{0,-}\}$ be the linear map given by*

$$(1 - \Phi)(\chi_A) = \chi_A - \sum_{a \in \mathcal{A}} \chi_{r(A,a)} \quad A \in \mathcal{E}_{\Lambda^*}^{0,-}.$$

Then $K_1(\mathcal{O}_{\Lambda^})$ is isomorphic to $\ker(1 - \Phi)$ and there exists an isomorphism from $K_0(\mathcal{O}_{\Lambda^*})$ to $\text{coker}(1 - \Phi)$ which maps $[S_u^*S_u]_0$ to $\chi_{r(u)} + \text{Im}(1 - \Phi)$ for each $u \in \mathcal{A}^* \setminus \{\epsilon\}$.*

If Λ satisfies condition (I) of [29], then we have similar formulas for $K_0(\mathcal{O}_\Lambda)$ and $K_1(\mathcal{O}_\Lambda)$.

Proof. The labelled spaces $(E_\Lambda, \mathcal{L}_\Lambda, \mathcal{E}_\Lambda^{0,-})$ and $(E_{\Lambda^*}, \mathcal{L}_{\Lambda^*}, \mathcal{E}_{\Lambda^*}^{0,-})$ are weakly left-resolving and normal. Since $\mathcal{E}_\Lambda^{0,-} = (\mathcal{E}_{\Lambda^*}^{0,-})_J$ and $\mathcal{E}_{\Lambda^*}^{0,-} = (\mathcal{E}_\Lambda^{0,-})_J$, the result follows from [4, Theorem 6.3] and Theorem 4.4. \square

Example 4 – Cuntz-Pimsner algebras associated with subshifts. Let X be a one-sided shift space over a finite alphabet \mathcal{A} . In [8] a C^* -algebra was associated with X . An alternative construction of this C^* -algebra is given in [10] and its relationship with the algebras \mathcal{O}_Λ and \mathcal{O}_{Λ^*} is explored. We will denote this C^* -algebra by $C^*(X)$ (it is denoted by \mathcal{O}_X in [8] and by $\mathcal{D}_X \rtimes_{\alpha, \mathcal{L}} \mathbb{N}$ in [10]).

We show how one may realise $C^*(X)$ as the C^* -algebra of a labelled space. We let E_X be the directed graph (E_X^0, E_X^1, r, s) where $E_X^0 = X$, $E_X^1 = \{(x, a, y) \in X \times \mathcal{A} \times X : x = ay\}$ and $r, s : E_X^1 \rightarrow E_X^0$ are defined by $s(x, a, y) = x$ and $r(x, a, y) = y$. We let $\mathcal{L}_X : E_X^1 \rightarrow \mathcal{A}$ be the labeling given by $\mathcal{L}_X(x, a, y) = a$.

For $u, v \in \mathcal{L}_X^\#(E_X)$, let $C(u, v) = \{vx \in X : ux \in X\}$. Let \mathcal{B}_X be the Boolean algebra generated by $\{C(u, v) : u, v \in \mathcal{L}_X^\#(E_X)\}$. That is, \mathcal{B}_X is the smallest subset of 2^X which is closed under finite intersections, finite unions, relative complements and which contains $C(u, v)$ for every $u, v \in \mathcal{L}_X^\#(E_X)$.

Lemma 5.4. *With the above definitions $(E_X, \mathcal{L}_X, \mathcal{B}_X)$ is a normal weakly left-resolving labelled space.*

Proof. We will first prove that \mathcal{B}_X is accommodating for (E_X, \mathcal{L}_X) . Let $\alpha \in \mathcal{L}_X^*(E_X)$. Then we have that $r(\alpha) = C(\alpha, \epsilon)$, from which it follows that \mathcal{B}_X contains $r(\alpha)$. It remains to show that \mathcal{B}_X is closed under relative ranges.

To check that \mathcal{B}_X is closed under relative ranges it suffices to check that $r(C(u, v), a) \in \mathcal{B}_X$ for all $u, v \in \mathcal{L}_X^\#(E_X)$ and $a \in \mathcal{A}$. If $u, v \in \mathcal{L}_X^\#(E_X)$ are such that $v = v_1v_2 \dots v_n \neq \epsilon$, then we claim that

$$(6) \quad r(C(u, v), a) = \begin{cases} C(a, \epsilon) \cap C(u, v_2v_3 \dots v_n) & \text{if } a = v_1 \\ \emptyset & \text{otherwise.} \end{cases}$$

To see this observe that every vertex in $C(u, v)$ emits exactly one edge, labelled v_1 , and so $r(C(u, v), a) = \emptyset$ if $v_1 \neq a$. If $v_1 = a$ and $x \in C(a, \epsilon) \cap C(u, v_2v_3 \dots v_n)$, then $ax \in C(u, v)$ and so $x \in r(C(u, v), a)$. If $v_1 = a$ and $x \in r(C(u, v), a)$, then $ax \in C(u, v)$ and so $x \in C(a, \epsilon) \cap C(u, v_2v_3 \dots v_n)$. Thus Equation (6) holds. If $u \in \mathcal{L}_X^\#(E_X)$ and $a \in \mathcal{A}$, then it is easy to check that $r(C(u, \epsilon), a) = C(ua, \epsilon)$. It follows that \mathcal{B}_X is accommodating for (E_X, \mathcal{L}_X) . Moreover, \mathcal{B}_X is non-degenerate by definition.

Since a vertex $x \in E_X^0$ can only receive an edge labelled a from the vertex ax we see that (E_X, \mathcal{L}_X) is left-resolving. Thus $(E_X, \mathcal{L}_X, \mathcal{B}_X)$ is weakly left-resolving and normal. \square

As in [8], we let $\widetilde{\mathcal{D}}_X$ denote the C^* -algebra generated by $\{\chi_{C(u,v)} : u, v \in \mathcal{L}_X^*(E_X)\}$. We then have that $\widetilde{\mathcal{D}}_X = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}_X\} = \mathcal{A}(E_X, \mathcal{L}_X, \mathcal{B}_X)$.

According to [8, Remark 7.3], $C^*(X)$ is the universal C^* -algebra generated by a family $\{S_a : a \in \mathcal{A}\}$ of partial isometries such that there exists a $*$ -homomorphism from $\iota_X : \widetilde{\mathcal{D}}_X \rightarrow C^*(X)$ given by

$$(7) \quad \iota_X : \chi_{C(u,v)} \mapsto (S_{v_1} S_{v_2} \dots S_{v_m})(S_{u_1} S_{u_2} \dots S_{u_n})^*(S_{u_1} S_{u_2} \dots S_{u_n})(S_{v_1} S_{v_2} \dots S_{v_m})^*$$

where $u = u_1 u_2 \dots u_n$ and $v = v_1 v_2 \dots v_m \in \mathcal{L}_X^\#(E_X)$. Note that if u (respectively v) is the empty word, then the product $(S_{u_1} S_{u_2} \dots S_{u_n})$ (respectively $(S_{v_1} S_{v_2} \dots S_{v_m})$) is the identity of $C^*(X)$. If $u = u_1 u_2 \dots u_n \in \mathcal{L}_X^*(E_X)$, then we will denote the product $S_{u_1} S_{u_2} \dots S_{u_n}$ by S_u . If $u = \epsilon$, then we let S_u denote the unit of $C^*(X)$.

Lemma 5.5. *Let $\iota_X : \widetilde{\mathcal{D}}_X \rightarrow C^*(X)$ be defined in (7) above. Then for every $x \in \widetilde{\mathcal{D}}_X$ and $a \in \mathcal{A}$ we have $\iota_X(\phi_a(x)) = S_a^* \iota_X(x) S_a$ where $\phi_a : \widetilde{\mathcal{D}}_X \rightarrow \widetilde{\mathcal{D}}_X$ is given by $\phi_a(\chi_A) = \chi_{r(A,a)}$ for $A \in \mathcal{B}_X$ (see Lemma 3.4).*

Proof. Let $u, v \in \mathcal{L}_X^\#(E_X)$ with $v = v_1 v_2 \dots v_m \neq \epsilon$. If $a = v_1$ then

$$\begin{aligned} S_a^* \iota_X(\chi_{C(u,v)}) S_a &= S_a^* S_v S_u^* S_u S_v^* S_a = S_a^* S_a S_{v_2} S_{v_3} \dots S_{v_m} S_u^* S_u (S_{v_2} S_{v_3} \dots S_{v_m})^* \\ &= \iota_X(\chi_{C(a,\epsilon)} \chi_{C(u,v_2 v_3 \dots v_m)}), \end{aligned}$$

otherwise it is zero. Thus $S_a^* \iota_X(\chi_{C(u,v)}) S_a = \iota_X(\chi_{r(C(u,v),a)}) = \iota_X(\phi_a(\chi_{C(u,v)}))$ by Equation (6). One can similarly show that if $u \in \mathcal{L}_X^\#(E_X)$, then $S_a^* \iota_X(\chi_{C(u,\epsilon)}) S_a = \iota_X(\phi_a(\chi_{C(u,\epsilon)}))$. Since $\widetilde{\mathcal{D}}_X$ is generated by $\{\chi_{C(u,v)} : u, v \in \mathcal{L}_X^\#(E_X)\}$, it follows that $\iota_X(\phi_a(x)) = S_a^* \iota_X(x) S_a$ for every $x \in \widetilde{\mathcal{D}}_X$. \square

Proposition 5.6. *Let X be a one-sided shift space over a finite alphabet \mathcal{A} . Let $\{p_A, s_a : A \in \mathcal{B}_X, a \in \mathcal{A}\}$ and $\{S_a : a \in \mathcal{A}\}$ be the canonical generators of $C^*(E_X, \mathcal{L}_X, \mathcal{B}_X)$ and $C^*(X)$, respectively. Then the map $S_a \mapsto s_a$ for $a \in \mathcal{A}$ induces an isomorphism from $C^*(X)$ to $C^*(E_X, \mathcal{L}_X, \mathcal{B}_X)$.*

Proof. Let ι_X denote (the unique) $*$ -homomorphism from $\widetilde{\mathcal{D}}_X$ to $C^*(X)$ mapping $\chi_{C(u,v)}$ to

$$(S_{v_1} S_{v_2} \dots S_{v_m})(S_{u_1} S_{u_2} \dots S_{u_n})^*(S_{u_1} S_{u_2} \dots S_{u_n})(S_{v_1} S_{v_2} \dots S_{v_m})^*$$

for $u, v \in \mathcal{L}_X^\#(E_X)$. We will show that $\{\iota_X(\chi_A), S_a : A \in \mathcal{B}_X, a \in \mathcal{A}\}$ is a representation of $(E_X, \mathcal{L}_X, \mathcal{B}_X)$. First, one checks that $\{\iota_X(\chi_A) : A \in \mathcal{B}_X\}$ satisfies condition (i) of Definition 2.1. Second, if $a \in \mathcal{A}$ and $A \in \mathcal{B}_X$, then it follows from Lemma 5.5 that

$$\iota_X(\chi_A) S_a = \iota_X(\chi_A) S_a S_a^* S_a = S_a S_a^* \iota_X(\chi_A) S_a = S_a \iota_X(\phi_a(\chi_A)) = S_a \iota_X(\chi_{r(A,a)})$$

which shows that $\{\iota_X(\chi_A), S_a : A \in \mathcal{B}_X, a \in \mathcal{A}\}$ satisfies condition (ii) of Definition 2.1. If $a \in \mathcal{A}$, then $S_a^* S_a = \iota_X(\chi_{C(a,\epsilon)}) = \iota_X(\chi_{r(a)})$. If $a, b \in \mathcal{A}$ and $a \neq b$, then $S_a^* S_b = S_a^* \iota_X(\chi_{C(\epsilon,a)} \chi_{C(\epsilon,b)}) S_b = 0$. Thus $\{\iota_X(\chi_A), S_a : A \in \mathcal{B}_X, a \in \mathcal{A}\}$ satisfies condition (iii) of Definition 2.1. We have that $\bigcup_{a \in \mathcal{A}} C(\epsilon, a) = E_X^0$ and $C(\epsilon, a) \cap C(\epsilon, b) = \emptyset$ for $a, b \in \mathcal{A}$ with $a \neq b$ and so $\chi_{E_X^0} = \sum_{a \in \mathcal{A}} \chi_{C(\epsilon,a)}$ in $\widetilde{\mathcal{D}}_X$. Since $\iota_X(\chi_{E_X^0}) = 1_{C^*(X)}$ it follows from Lemma 5.5 that if $A \in \mathcal{B}_X$, then

$$\begin{aligned} \iota_X(\chi_A) &= \sum_{a \in \mathcal{L}_X(E_X^1)} \iota_X(\chi_{C(\epsilon,a)} \chi_A \chi_{C(\epsilon,a)}) = \sum_{a \in \mathcal{L}_X(AE_X^1)} S_a S_a^* \iota_X(\chi_A) S_a S_a^* \\ &= \sum_{a \in \mathcal{L}_X(AE_X^1)} S_a \iota_X(\phi_a(\chi_A)) S_a^* = \sum_{a \in \mathcal{L}_X(AE_X^1)} S_a \iota_X(\chi_{r(A,a)}) S_a^* \end{aligned}$$

which shows that $\{\iota_X(\chi_A), S_a : A \in \mathcal{B}_X, a \in \mathcal{A}\}$ satisfies condition (iv) of Definition 2.1. Hence $\{\iota_X(\chi_A), S_a : A \in \mathcal{B}_X, a \in \mathcal{A}\}$ is a representation of $(E_X, \mathcal{L}_X, \mathcal{B}_X)$. It follows from the universal

property of $C^*(E_X, \mathcal{L}_X, \mathcal{B}_X)$ that there is a $*$ -homomorphism $\pi_{s,p} : C^*(E_X, \mathcal{L}_X, \mathcal{B}_X) \rightarrow C^*(X)$ which sends p_A to χ_A and s_a to S_a for all $A \in \mathcal{B}_X$ and $a \in \mathcal{A}$.

It follows from Lemma 3.2 that there exists a $*$ -homomorphism from $\tilde{D}_X = \mathcal{A}(E_X, \mathcal{L}_X, \mathcal{B}_X)$ to $C^*(E_X, \mathcal{L}_X, \mathcal{B}_X)$ which maps χ_A to p_A for every $A \in \mathcal{B}_X$. In particular, if $u, v \in \mathcal{L}_X^\#(E_X)$, then $\phi(\chi_{C(u,v)}) = p_{C(u,v)}$. Repeated applications of conditions (iii) and (iv) of Definition 2.1 show that if $u \in \mathcal{L}_X^\#(E_X)$ then $s_u^* s_u = p_{r(u)} = p_{C(u,\epsilon)}$ (if $u = \epsilon$, then $r(u) = C(u, \epsilon) = X$). For $v = v_1 v_2 \dots v_m \in \mathcal{L}_X^*(E_X)$, condition (iv) of Definition 2.1 implies that

$$\begin{aligned} p_{C(u,v)} &= \sum_{a \in \mathcal{L}_X(C(u,v)E_X^1)} s_a p_{r(C(u,v),a)} s_a^* = s_{v_1} p_{C(u,v_2 v_3 \dots v_m)} s_{v_1}^* \text{ by Equation (6)} \\ &= s_{v_1} \left(\sum_{a \in \mathcal{L}_X(C(u,v_2 v_3 \dots v_m)E_X^1)} s_a p_{r(C(u,v_2 v_3 \dots v_m),a)} s_a^* \right) s_{v_1}^* = s_{v_1} s_{v_2} p_{C(u,v_3 \dots v_m)} s_{v_2}^* s_{v_1}^* \\ &= \dots = s_{v_1} s_{v_2} \dots s_{v_m} p_{C(u,\epsilon)} s_{v_m}^* \dots s_{v_2}^* s_{v_1}^* = s_v s_u^* s_u s_v^*. \end{aligned}$$

Thus, it follows from the universal property of $C^*(X)$ that there is a $*$ -homomorphism $\pi_{S,P} : C^*(X) \rightarrow C^*(E_X, \mathcal{L}_X, \mathcal{B}_X)$ which sends S_a to s_a and $\iota_X(\chi_A)$ to p_A for all $a \in \mathcal{A}$ and $A \in \mathcal{B}_X$.

Since $\pi_{s,p}$ and $\pi_{S,P}$ are evidently inverses, the result follows. \square

Corollary 5.7 (cf. [11, Theorem 1.1]). *Let X be a one-sided shift space over a finite alphabet \mathcal{A} . Let $(1 - \Phi) : \text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}_X\} \rightarrow \text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}_X\}$ be the linear map given by*

$$(1 - \Phi)(\chi_A) = \chi_A - \sum_{a \in \mathcal{L}_X(AE_X^1)} \chi_{r(A,a)}, \quad A \in \mathcal{B}_X.$$

Then $K_1(C^(X))$ is isomorphic to $\ker(1 - \Phi)$ and there exists an isomorphism from $K_0(C^*(X))$ to $\text{coker}(1 - \Phi)$ which maps $[S_u^* S_u]_0$ to $\chi_{r(u)} + \text{Im}(1 - \Phi)$ for each $u \in \mathcal{L}_X^*(E_X)$.*

Proof. Notice that $(\mathcal{B}_X)_J = \mathcal{B}_X$. The result now follows from Theorem 4.4 and Proposition 5.6. \square

6. COMPUTING K -THEORY FOR $C^*(E, \mathcal{L}, \mathcal{E}^{0,-})$

Recall from [5] that a labelled space $(E, \mathcal{L}, \mathcal{B})$ is *set-finite* if $\mathcal{L}(AE^1)$ is finite for all $A \in \mathcal{B}$, and *receiver set-finite* if for all $A \in \mathcal{B}$ and all $\ell \geq 1$ the set $\{\mathcal{L}(\lambda) : \lambda \in E^\ell, r(\lambda) \in A\}$ is finite.

In this section we give a practical method for computing the K -theory of labelled graph algebras of the form $C^*(E, \mathcal{L}, \mathcal{E}^{0,-})$ for left-resolving labelled graphs (E, \mathcal{L}) with no sources and sinks such that $(E, \mathcal{L}, \mathcal{E}^{0,-})$ is a set-finite and receiver set-finite labelled space. We use Theorem 4.4 to give a description of the K -theory of $C^*(E, \mathcal{L}, \mathcal{E}^{0,-})$ as an inductive limit in analogy with the computations of [36].

Standing Assumption: Throughout this section (E, \mathcal{L}) shall be a left-resolving labelled graph with no sources and sinks such that $(E, \mathcal{L}, \mathcal{E}^{0,-})$ is a set-finite and receiver set-finite (weakly left-resolving normal) labelled space.

Before we begin, let us recall some notation from [5]: For $v \in E^0$ and $\ell \geq 1$ let

$$\Lambda_\ell(v) = \{\alpha \in \mathcal{L}(E^k) : k \leq \ell, v \in r(\alpha)\}$$

denote the set of labelled paths of length at most ℓ which have a representative terminating at v . Define the relation \sim_ℓ on E^0 by $v \sim_\ell w$ if and only if $\Lambda_\ell(v) = \Lambda_\ell(w)$; hence $v \sim_\ell w$ if v and w receive exactly the same labelled paths of length at most ℓ . Evidently \sim_ℓ is an equivalence relation and we use $[v]_\ell$ to denote the equivalence class of $v \in E^0$. We call the $[v]_\ell$ *generalised vertices* as they play the same role in labelled spaces as vertices in a directed graph.

Set $\Omega_\ell = E^0 / \sim_\ell = \{[v]_\ell : v \in E^0\}$ and let

$$\Omega = \bigcup_{\ell=1}^{\infty} \Omega_\ell.$$

It follows from [5, Proposition 2.4] that if $v \in E^0$ and $\ell \geq 1$, then

$$(8) \quad [v]_\ell = \bigcup_{i=1}^n [w_i]_{\ell+1} \text{ for some } w_1, \dots, w_n \in [v]_\ell.$$

We then have that

$$(9) \quad r([v]_\ell, a) = \bigcup_{j=1}^m r([w_j]_{\ell+1}, a) \text{ for all } a \in \mathcal{A}.$$

It also follows from [5, Proposition 2.4] that $\mathcal{E}^{0,-}$ is the smallest subset of 2^{E^0} that contains Ω and is closed under finite unions and relative complements.

We let $\mathbb{Z}(\Omega)$ denote the subgroup $\text{span}_{\mathbb{Z}}\{\chi_{[v]_\ell} : [v]_\ell \in \Omega\}$ of the group of functions from E^0 to \mathbb{Z} .

Corollary 6.1. *Let $(1 - \Phi) : \mathbb{Z}(\Omega) \rightarrow \mathbb{Z}(\Omega)$ be the linear map defined by*

$$(1 - \Phi)(\chi_{[v]_\ell}) = \chi_{[v]_\ell} - \sum_{a \in \mathcal{L}([v]_\ell E^1)} \chi_{r([v]_\ell, a)} \text{ for } [v]_\ell \in \Omega.$$

Then

$$\begin{aligned} K_1(C^*(E, \mathcal{L}, \mathcal{E}^{0,-})) &\cong \ker(1 - \Phi) \text{ and} \\ K_0(C^*(E, \mathcal{L}, \mathcal{E}^{0,-})) &\cong \text{coker}(1 - \Phi) \end{aligned}$$

via $[p_{[v]_\ell}]_0 \mapsto \chi_{[v]_\ell} + \text{Im}(1 - \Phi)$ for $[v]_\ell \in \Omega$.

Proof. Since $\text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{E}^{0,-}\} = \text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{E}_j^{0,-}\} = \mathbb{Z}(\Omega)$, the result follows from Theorem 4.4. \square

In Theorem 6.5 we shall show how to compute the kernel and cokernel of $(1 - \Phi)$ in terms of a certain inductive limit. For $\ell \geq 1$ define $\mathbb{Z}(\Omega_\ell) = \text{span}_{\mathbb{Z}}\{\chi_{[v]_\ell} : [v]_\ell \in \Omega_\ell\}$. Note that since for every $v \in E^0$ and $\ell \geq 1$ there are vertices $w_1, \dots, w_m \in [v]_\ell$ such that $[v]_\ell = \bigcup_{j=1}^m [w_j]_{\ell+1}$ we have a linear map $i_\ell : \mathbb{Z}(\Omega_\ell) \rightarrow \mathbb{Z}(\Omega_{\ell+1})$ defined by

$$i_\ell(\chi_{[v]_\ell}) = \sum_{j=1}^m \chi_{[w_j]_{\ell+1}}.$$

Therefore we have an inductive system $\varinjlim (\mathbb{Z}(\Omega_\ell), i_\ell)$. Note that since, for fixed $\ell \geq 1$, the sets $[v]_\ell$ are disjoint, the functions $\chi_{[v]_\ell}$ form a basis for $\mathbb{Z}(\Omega_\ell)$.

Proposition 6.2. *There is an isomorphism*

$$\Psi_\infty : \varinjlim (\mathbb{Z}(\Omega_\ell), i_\ell) \rightarrow \mathbb{Z}(\Omega)$$

such that $\Psi_\infty(\chi_{[v]_\ell}) = \chi_{[v]_\ell}$ for all $[v]_\ell \in \Omega$.

Proof. For each $\ell \geq 1$ define $\Psi_\ell : \mathbb{Z}(\Omega_\ell) \rightarrow \mathbb{Z}(\Omega)$ by $\Psi_\ell(\chi_{[v]_\ell}) = \chi_{[v]_\ell}$. One checks that $\Psi_\ell \circ (i_\ell \circ \dots \circ i_k) = \Psi_k$ for every $\ell < k$ and so there is a map $\Psi_\infty : \varinjlim (\mathbb{Z}(\Omega_\ell), i_\ell) \rightarrow \mathbb{Z}(\Omega)$. Since $\mathbb{Z}(\Omega) = \bigcup_{\ell \geq 1} \Psi_\ell(\mathbb{Z}(\Omega_\ell))$ and each Ψ_ℓ is injective, it follows that Ψ_∞ is an isomorphism and our result is established. \square

Equations (8) and (9) allow us to make the following definitions.

Definition 6.3. For $\ell \geq 1$ define a linear map $(1 - \Phi)_\ell : \mathbb{Z}(\Omega_\ell) \rightarrow \mathbb{Z}(\Omega_{\ell+1})$ by

$$(1 - \Phi)_\ell (\chi_{[v]_\ell}) = i_\ell (\chi_{[v]_\ell}) - \sum_{a \in \mathcal{L}([v]_\ell E^1)} \chi_{r([v]_\ell, a)},$$

and define a linear map $(1 - \Phi) : \mathbb{Z}(\Omega) \rightarrow \mathbb{Z}(\Omega)$ by

$$(1 - \Phi) (\chi_{[v]_\ell}) = \chi_{[v]_\ell} - \sum_{a \in \mathcal{L}([v]_\ell E^1)} \chi_{r([v]_\ell, a)}.$$

Lemma 6.4. For all $\ell \geq 1$, we have $(1 - \Phi)_{\ell+1} \circ i_\ell = i_{\ell+1} \circ (1 - \Phi)_\ell$. Therefore the $(1 - \Phi)_\ell$'s induce a map $(1 - \Phi)_\infty : \varinjlim (\mathbb{Z}(\Omega_\ell), i_\ell) \rightarrow \varinjlim (\mathbb{Z}(\Omega_\ell), i_\ell)$ which satisfies $(1 - \Phi) \circ \Psi_\infty = \Psi_\infty \circ (1 - \Phi)_\infty$.

Proof. Let $\ell \geq 1$. Then for $[v]_\ell \in \Omega_\ell$ with $[v]_\ell = \bigcup_{j=1}^m [w_j]_{\ell+1}$ we have

$$\begin{aligned} (1 - \Phi)_{\ell+1} (i_\ell (\chi_{[v]_\ell})) &= (1 - \Phi)_{\ell+1} \left(\sum_{j=1}^m \chi_{[w_j]_{\ell+1}} \right) \\ &= \sum_{j=1}^m \left(i_{\ell+1} (\chi_{[w_j]_{\ell+1}}) - \sum_{a \in \mathcal{L}([w_j]_{\ell+1} E^1)} \chi_{r([w_j]_{\ell+1}, a)} \right). \end{aligned}$$

Also,

$$\begin{aligned} i_{\ell+1} ((1 - \Phi)_\ell (\chi_{[v]_\ell})) &= i_{\ell+1} \left(i_\ell (\chi_{[v]_\ell}) - \sum_{a \in \mathcal{L}([v]_\ell E^1)} \chi_{r([v]_\ell, a)} \right) \\ &= i_{\ell+1} \left(\sum_{j=1}^m \chi_{[w_j]_{\ell+1}} - \sum_{j=1}^m \sum_{a \in \mathcal{L}([w_j]_{\ell+1} E^1)} \chi_{r([w_j]_{\ell+1}, a)} \right) \end{aligned}$$

since $\chi_{r([v]_\ell, a)} = \sum_{j=1}^m \chi_{r([w_j]_{\ell+1}, a)}$,

$$= \sum_{j=1}^m \left(i_{\ell+1} (\chi_{[w_j]_{\ell+1}}) - \sum_{a \in \mathcal{L}([w_j]_{\ell+1} E^1)} \chi_{r([w_j]_{\ell+1}, a)} \right)$$

as $i_{\ell+1}$ is really just the identity map on $\Omega_{\ell+1}$ and so $(1 - \Phi)_{\ell+1} \circ i_\ell = i_{\ell+1} \circ (1 - \Phi)_\ell$.

Also for $\ell \geq 1$ we have

$$(1 - \Phi)(\Psi_\infty(\chi_{[v]_\ell})) = (1 - \Phi)(\chi_{[v]_\ell}) = i_\ell(\chi_{[v]_\ell}) - \sum_{a \in \mathcal{L}([v]_\ell E^1)} \chi_{r([v]_\ell, a)}$$

and

$$\begin{aligned} \Psi_\infty((1 - \Phi)_\infty(\chi_{[v]_\ell})) &= \Psi_\infty((1 - \Phi)_\ell(\chi_{[v]_\ell})) = \Psi_\infty \left(i_\ell(\chi_{[v]_\ell}) - \sum_{a \in \mathcal{L}([v]_\ell E^1)} \chi_{r([v]_\ell, a)} \right) \\ &= i_\ell(\chi_{[v]_\ell}) - \sum_{a \in \mathcal{L}([v]_\ell E^1)} \chi_{r([v]_\ell, a)} \end{aligned}$$

and so $(1 - \Phi) \circ \Psi_\infty = \Psi_\infty \circ (1 - \Phi)_\infty$. □

We have established the following commuting diagram:

$$\begin{array}{ccccccc}
 \mathbb{Z}(\Omega_1) & \xrightarrow{i_1} & \mathbb{Z}(\Omega_2) & \xrightarrow{i_2} & \mathbb{Z}(\Omega_2) & \xrightarrow{i_3} & \dots & \lim_{\rightarrow} (\mathbb{Z}(\Omega_\ell), i_\ell) & \xrightarrow{\Psi_\infty} & \mathbb{Z}(\Omega) \\
 & & \searrow (1-\Phi)_1 & & \searrow (1-\Phi)_2 & & & \downarrow (1-\Phi)_\infty & & \downarrow 1-\Phi \\
 \mathbb{Z}(\Omega_1) & \xrightarrow{i_1} & \mathbb{Z}(\Omega_2) & \xrightarrow{i_2} & \mathbb{Z}(\Omega_2) & \xrightarrow{i_3} & \dots & \lim_{\rightarrow} (\mathbb{Z}(\Omega_\ell), i_\ell) & \xrightarrow{\Psi_\infty} & \mathbb{Z}(\Omega)
 \end{array}$$

Let $\ell \geq 1$. It is easy to check that $i_\ell(\ker(1-\Phi)_\ell) \subseteq \ker(1-\Phi)_{\ell+1}$ and $i_\ell(\text{Im}(1-\Phi)_\ell) \subseteq \text{Im}(1-\Phi)_{\ell+1}$. It follows that $i_\ell : \mathbb{Z}(\Omega_\ell) \rightarrow \mathbb{Z}(\Omega_{\ell+1})$ induces maps $(i_\ell)_* : \ker(1-\Phi)_\ell \rightarrow \ker(1-\Phi)_{\ell+1}$ and $\tilde{i}_{\ell+1} : \text{coker}(1-\Phi)_\ell \rightarrow \text{coker}(1-\Phi)_{\ell+1}$.

Theorem 6.5. *With the above notation we have*

$$(10) \quad \ker(1-\Phi) \cong \lim_{\rightarrow} (\ker(1-\Phi)_\ell, (i_\ell)_*),$$

$$(11) \quad \text{coker}(1-\Phi) \cong \lim_{\rightarrow} (\text{coker}(1-\Phi)_\ell, \tilde{i}_\ell).$$

Proof. By definition of $(1-\Phi)_\infty$ we have

$$\ker(1-\Phi)_\infty \cong \lim_{\rightarrow} (\ker(1-\Phi)_\ell, (i_\ell)_*).$$

It follows from Lemma 6.4 that $\ker(1-\Phi) \cong \ker(1-\Phi)_\infty$ and Equation (10) follows. We claim that

$$\text{coker}(1-\Phi)_\infty \cong \lim_{\rightarrow} (\text{coker}(1-\Phi)_\ell, \tilde{i}_\ell).$$

Define $\eta_\ell : \text{coker}(1-\Phi)_\ell \rightarrow \text{coker}(1-\Phi)_\infty$ by

$$\eta_\ell (\chi_{[v]_{\ell+1}} + \text{Im}(1-\Phi)_\ell) = \chi_{[v]_\ell} + \text{Im}(1-\Phi)_\infty.$$

Note that η_ℓ is well-defined since $\text{Im}(1-\Phi)_\ell \subseteq \text{Im}(1-\Phi)_\infty$ for all $\ell \geq 1$. We claim that $\eta_{\ell+1} \circ \tilde{i}_\ell = \eta_\ell$ for all $\ell \geq 1$. Since

$$\begin{aligned}
 \eta_{\ell+1} \tilde{i}_\ell (\chi_{[v]_{\ell+1}} + \text{Im}(1-\Phi)_\ell) &= \eta_{\ell+1} (i_{\ell+1} \chi_{[v]_{\ell+1}} + \text{Im}(1-\Phi)_{\ell+1}) \\
 &= i_{\ell+1} (\chi_{[v]_{\ell+1}}) + \text{Im}(1-\Phi)_\infty \\
 &= \chi_{[v]_{\ell+1}} + \text{Im}(1-\Phi)_\infty \\
 &= \eta_\ell (\chi_{[v]_{\ell+1}} + \text{Im}(1-\Phi)_\ell),
 \end{aligned}$$

this establishes our claim. By the universal property of the inductive limit the η_ℓ 's induce a map $\eta_\infty : \lim_{\rightarrow} (\text{coker}(1-\Phi)_\ell, \tilde{i}_\ell) \rightarrow \text{coker}(1-\Phi)_\infty$ which is injective since each η_ℓ is injective, and is surjective as $\text{coker}(1-\Phi)_\infty = \bigcup_{\ell \geq 1} \eta_\ell(\text{coker}(1-\Phi)_\ell)$. Hence η_∞ is an isomorphism which establishes our claim. Finally, it follows from Lemma 6.4 that $\text{coker}(1-\Phi)_\infty \cong \text{coker}(1-\Phi)$ and Equation (11) follows. \square

Remark 6.6. Recall from [5] that to a left-resolving normal labelled graph (E, \mathcal{L}) with no sources and sinks over a finite alphabet we may associate an essential symbolic matrix system $(M(E), I(E))$, which by [31, Proposition 2.1] determines a unique λ -graph system $\mathfrak{L}_{M(E), I(E)}$.

By [5, Proposition 3.6] we know that $C^*(E, \mathcal{L}, \mathcal{E}^{0,-})$ is isomorphic to $\mathcal{O}_{\mathfrak{L}_{M(E), I(E)}}$. We should therefore expect some similarities between our computation of the K -theory of $C^*(E, \mathcal{L}, \mathcal{E}^{0,-})$ and the computation of the K -theory of $\mathcal{O}_{\mathfrak{L}_{M(E), I(E)}}$ outlined in [31, Section 9] (see also [28, 34]). Indeed, this is the case.

To an essential symbolic matrix system $(M, I) = (M_{\ell, \ell+1}, I_{\ell, \ell+1})_{\ell \geq 1}$ we may associate a λ -graph system $\mathfrak{L}_{M, I}$ as in [31, Section 2]. Following [31, Section 9] we see that

$$K_*(\mathcal{O}_{\mathfrak{L}_{M, I}}) = K_*(M, I) = \varinjlim (K_*(M, I), i_*^\ell)$$

for $* = 0, 1$ where

$$\begin{aligned} K_0^\ell(M, I) &= \mathbb{Z}^{m(\ell+1)} / (I_{\ell, \ell+1}^t - M_{\ell, \ell+1}^t) \mathbb{Z}^{m(\ell)} \\ K_1^\ell(M, I) &= \ker(I_{\ell, \ell+1}^t - M_{\ell, \ell+1}^t) \text{ in } \mathbb{Z}^{m(\ell)} \end{aligned}$$

and $i_*^\ell : K_*^\ell(M, I) \rightarrow K_*^{\ell+1}(M, I)$ is induced by the map $I_{\ell, \ell+1}^t : \mathbb{Z}^{m(\ell)} \rightarrow \mathbb{Z}^{m(\ell+1)}$.


Under our identification of the labelled graph (E, \mathcal{L}) with the essential symbolic matrix system $(M(E), I(E))$ we have $|\Omega_\ell| = m(\ell)$, and so we may identify the group $\mathbb{Z}(\Omega_\ell)$ with $\mathbb{Z}^{m(\ell)}$ and the map $(1 - \Phi)_\ell : \mathbb{Z}(\Omega_\ell) \rightarrow \mathbb{Z}(\Omega_{\ell+1})$ with the map $(I(E)_{\ell, \ell+1}^t - M(E)_{\ell, \ell+1}^t) : \mathbb{Z}^{m(\ell)} \rightarrow \mathbb{Z}^{m(\ell+1)}$. Hence $\text{coker}(1 - \Phi)_\ell$ may be identified with $K_0^\ell(M(E), I(E))$ and $\ker(1 - \Phi)_\ell$ may be identified with $K_1^\ell(M(E), I(E))$. Since the map $(i_\ell)_* : \ker(1 - \Phi)_\ell \rightarrow \ker(1 - \Phi)_{\ell+1}$ corresponds to $i_0^\ell : K_0^\ell(M(E), I(E)) \rightarrow K_0^{\ell+1}(M(E), I(E))$ and the map $\tilde{i}_\ell : \text{coker}(1 - \Phi)_\ell \rightarrow \text{coker}(1 - \Phi)_{\ell+1}$ corresponds to $i_1^\ell : K_1^\ell(M(E), I(E)) \rightarrow K_1^{\ell+1}(M(E), I(E))$, we may identify $\text{coker}(1 - \Phi)$ with $K_0(M(E), I(E))$ and $\ker(1 - \Phi)$ with $K_1(M(E), I(E))$.

7. COMPUTATIONS

In this section we compute the K -theory of certain weakly left-resolving normal labelled graphs using the techniques outlined in Section 6.

Example 7.1. Let (E_F, \mathcal{L}_F) be the left Fischer cover (see [2, Example 2.15]) of the even shift. Then $\Omega_\ell = E_F^0$ for all $\ell \geq 1$ and so $\mathbb{Z}(\Omega) = \mathbb{Z}^2$ and the matrix of $(1 - \Phi)$ is $\begin{pmatrix} 0 & -1 \\ -1 & 1 \end{pmatrix}$. Hence the K -theory of the labelled graph is the same as the K -theory of the underlying graph, that is $K_0 = K_1 = \{0\}$. This should not be a surprise in the light of [4, Theorem 6.6]. We obtain similar conclusions for the left Krieger cover (E_K, \mathcal{L}_K) of the even shift, though in this case $\Omega_\ell = E_K^0$ for $\ell \geq 2$.

Example 7.2. Now consider the labelled graph (E, \mathcal{L}) shown below



which was discussed in [5, Section 7.2] and [19]. We label the vertices of E by the integers, where $0 = r(a)$.

Fix $\ell \geq 1$. We set $\text{rest}_\ell = \{\pm\ell, \pm(\ell+1), \dots\}$. Then $[n]_\ell = \{n\}$ for $|n| \leq \ell - 1$ and $[n]_\ell = \text{rest}_\ell$ for $|n| \geq \ell$. Hence, if we identify each vertex n with the singleton $\{n\}$, then we have that

$$\Omega_1 = \{0, \text{rest}_1\}, \Omega_2 = \{0, \pm 1, \text{rest}_2\}, \dots, \Omega_\ell = \{0, \pm 1, \pm 2, \dots, \pm(\ell - 1), \text{rest}_\ell\}, \dots$$

By Definition 6.3 we have $(1 - \Phi)_1(\chi_0) = -\chi_1 - \chi_{-1}$ and $(1 - \Phi)_1(\chi_{\text{rest}_1}) = -2\chi_0 - \chi_{\text{rest}_2}$. It is straightforward to show that $\ker(1 - \Phi)_1 = \{0\}$ and that every element of $\mathbb{Z}(\Omega_2)$ can be written as $a\chi_0 + b\chi_1 + w$ where $a, b, \in \mathbb{Z}$ and $w \in \text{Im}(1 - \Phi)_1$. Hence $\text{coker}(1 - \Phi)_1 \cong \mathbb{Z}^2$. For $\ell \geq 2$ the maps $(1 - \Phi)_\ell : \mathbb{Z}(\Omega_\ell) \rightarrow \mathbb{Z}(\Omega_{\ell+1})$ are given by

$$(12) \quad (1 - \Phi)_\ell(\chi_n) = \chi_n - \chi_{n-1} - \chi_{n+1} \text{ if } n \neq \text{rest}_\ell, 0$$

$$(13) \quad (1 - \Phi)_\ell(\chi_0) = -\chi_1 - \chi_{-1}$$

$$(14) \quad (1 - \Phi)_\ell(\chi_{\text{rest}_\ell}) = -\chi_{\ell-1} - \chi_{-\ell+1} - \chi_{\text{rest}_{\ell+1}}.$$

Through systematic use of (14), then (12) for $n = -\ell + 1, \dots, -1$, followed by (13) and then (12) for $n = \ell - 1, \dots, 1$, one checks that $\{(1 - \Phi)_\ell(\chi_i) : i \in \Omega_\ell\}$ is linearly independent in $\mathbb{Z}(\Omega_{\ell+1})$, and so $\ker(1 - \Phi)_\ell = \{0\}$ for $\ell \geq 2$. A similar procedure allows one to show that every element of $\mathbb{Z}(\Omega_\ell)$ can be written in the form $a\chi_0 + b\chi_1 + w$ where $a, b, \in \mathbb{Z}$ and $w \in \text{Im}(1 - \Phi)_\ell$. Hence for $\ell \geq 2$ $\text{coker}(1 - \Phi)_\ell = \mathbb{Z}(\Omega_{\ell+1})/\text{Im}(1 - \Phi)_\ell$ is freely generated by

$$\chi_0 + \text{Im}(1 - \Phi)_\ell \text{ and } \chi_1 + \text{Im}(1 - \Phi)_\ell.$$

It follows that $\text{coker}(1 - \Phi)_\ell \cong \mathbb{Z}^2$ for all $\ell \geq 1$.

For $\ell \geq 1$ the maps $i_\ell : \mathbb{Z}(\Omega_\ell) \rightarrow \mathbb{Z}(\Omega_{\ell+1})$ are given by


$$i_\ell(\chi_i) = \chi_i \text{ for } 0 \leq |i| \leq \ell - 1 \text{ and}$$

$$i_\ell(\chi_{\text{rest}_\ell}) = \chi_\ell + \chi_{-\ell} + \chi_{\text{rest}_{\ell+1}}$$

and so for $\ell \geq 2$ the maps \tilde{i}_ℓ are the identity map. Thus, it follows from Theorem 6.5 that

$$K_0(C^*(E, \mathcal{L}, \mathcal{E}^{0,-})) \cong \mathbb{Z}^2 \text{ and } K_1(C^*(E, \mathcal{L}, \mathcal{E}^{0,-})) \cong \{0\}.$$

Example 7.3. Consider the Dyck shift D_2 which has labelled graph presentation (E_2, \mathcal{L}_2) below:



Since every vertex receives an edge labelled β_1, β_2 , the vertices in Ω_ℓ are distinguished by which labelled paths of length ℓ involving the symbols α_1, α_2 they receive.

For a word w in the symbols α_1, α_2 , we abuse our notation to denote the set of vertices comprising $r(w)$ by w . We then have

$$\begin{aligned} \Omega_1 &= \{\alpha_1, \alpha_2\} \\ \Omega_2 &= \{\alpha_1\alpha_1, \alpha_1\alpha_2, \alpha_2\alpha_1, \alpha_2\alpha_2\} \\ \Omega_3 &= \{\alpha_1\alpha_1\alpha_1, \alpha_1\alpha_1\alpha_2, \alpha_1\alpha_2\alpha_1, \alpha_1\alpha_2\alpha_2, \alpha_2\alpha_1\alpha_1, \alpha_2\alpha_1\alpha_2, \alpha_2\alpha_2\alpha_1, \alpha_2\alpha_2\alpha_2\} \\ &\vdots \\ \Omega_\ell &= \{\alpha_1^\ell, \alpha_1^{\ell-1}\alpha_2, \alpha_1^{\ell-2}\alpha_2\alpha_1, \alpha_1^{\ell-2}\alpha_2\alpha_2, \dots, \alpha_2^\ell\}. \end{aligned}$$

For $\ell = 1$ we have

$$(1 - \Phi)_1(\chi_{\alpha_1}) = -\chi_{\alpha_1\alpha_1} - 2\chi_{\alpha_1\alpha_2} - \chi_{\alpha_2\alpha_2}$$

$$(1 - \Phi)_1(\chi_{\alpha_2}) = -\chi_{\alpha_1\alpha_1} - 2\chi_{\alpha_2\alpha_1} - \chi_{\alpha_2\alpha_2}.$$

It is straightforward to see that $\ker(1 - \Phi)_1 = \{0\}$. Let

$$x_{11} = \chi_{\alpha_1\alpha_1} + \text{Im}(1 - \Phi)_1, x_{12} = \chi_{\alpha_1\alpha_2} + \text{Im}(1 - \Phi)_1, x_{21} = \chi_{\alpha_2\alpha_1} + \text{Im}(1 - \Phi)_1, x_{22} = \chi_{\alpha_2\alpha_2} + \text{Im}(1 - \Phi)_1.$$

Then $2(x_{11} + x_{12} + x_{21} + x_{22}) = 0$ since

$$(1 - \Phi)_1(\chi_{\alpha_1} + \chi_{\alpha_2}) = 2(\chi_{\alpha_1\alpha_1} + \chi_{\alpha_1\alpha_2} + \chi_{\alpha_2\alpha_1} + \chi_{\alpha_2\alpha_2}).$$

It follows that $\text{coker}(1 - \Phi)_1 \cong \mathbb{Z}^2 \times (\mathbb{Z}/2\mathbb{Z})$, generated by $x_{12}, x_{22}, x_{11} + x_{12} + x_{21} + x_{22}$.

As every vertex emits an edge labelled α_1 and an edge labelled α_2 , we have $r(w, \alpha_i) = w\alpha_i$ for $i = 1, 2$ and all words w in the symbols α_1, α_2 . Furthermore $r(w\alpha_i, \beta_j) = w$ if $i = j$ and is empty if $i \neq j$. Since $[w]_\ell = [\alpha_1 w]_{\ell+1} \cup [\alpha_2 w]_{\ell+1}$ for all words $w \in \Omega_\ell$, we have $i_\ell(\chi_w) = \chi_{\alpha_1 w} + \chi_{\alpha_2 w}$, and so

$$\begin{aligned} (1 - \Phi)_2(\chi_{\alpha_1 \alpha_1}) &= -\chi_{\alpha_1 \alpha_1 \alpha_1} - \chi_{\alpha_1 \alpha_1 \alpha_2} - \chi_{\alpha_1 \alpha_2 \alpha_1} - \chi_{\alpha_2 \alpha_2 \alpha_1} \\ (1 - \Phi)_2(\chi_{\alpha_1 \alpha_2}) &= -\chi_{\alpha_1 \alpha_1 \alpha_1} + \chi_{\alpha_1 \alpha_1 \alpha_2} - 2\chi_{\alpha_1 \alpha_2 \alpha_1} - \chi_{\alpha_1 \alpha_2 \alpha_2} - \chi_{\alpha_2 \alpha_1 \alpha_1} + \chi_{\alpha_2 \alpha_1 \alpha_2} - \chi_{\alpha_2 \alpha_2 \alpha_1} \\ (1 - \Phi)_2(\chi_{\alpha_2 \alpha_1}) &= -\chi_{\alpha_1 \alpha_1 \alpha_2} + \chi_{\alpha_1 \alpha_2 \alpha_1} - \chi_{\alpha_1 \alpha_2 \alpha_2} - \chi_{\alpha_2 \alpha_1 \alpha_1} - 2\chi_{\alpha_2 \alpha_1 \alpha_2} + \chi_{\alpha_2 \alpha_2 \alpha_1} - \chi_{\alpha_2 \alpha_2 \alpha_2} \\ (1 - \Phi)_2(\chi_{\alpha_2 \alpha_2}) &= -\chi_{\alpha_1 \alpha_1 \alpha_2} - \chi_{\alpha_2 \alpha_1 \alpha_2} - \chi_{\alpha_2 \alpha_2 \alpha_1} - \chi_{\alpha_2 \alpha_2 \alpha_2}. \end{aligned}$$

One checks that $\ker(1 - \Phi)_2 = \{0\}$. For a word $w \in \Omega_3$, let $x_w = \chi_w + \text{Im}(1 - \Phi)_2$. Then $2 \sum_{w \in \Omega_3} x_w = 0$, because

$$(1 - \Phi)_2(\chi_{\alpha_1 \alpha_2} + \chi_{\alpha_2 \alpha_2} + \chi_{\alpha_1 \alpha_1} + \chi_{\alpha_2 \alpha_2}) = 2 \sum_{w \in \Omega_3} \chi_w.$$

It follows that $\text{coker}(1 - \Phi)_2 \cong \mathbb{Z}^4 \times (\mathbb{Z}/2\mathbb{Z})$, generated by $x_{112}, x_{122}, x_{212}, x_{222}, \sum_{w \in \Omega_3} x_w$.

Since $\tilde{i}_2(x_{12}) = x_{112} + x_{212}$, $\tilde{i}_2(x_{22}) = x_{122} + x_{222}$ and $\tilde{i}_2(x_{11} + x_{12} + x_{21} + x_{22}) = \sum_{w \in \Omega_3} x_w$ we see that $\tilde{i}_2 : \mathbb{Z}^2 \times (\mathbb{Z}/2\mathbb{Z}) \rightarrow \mathbb{Z}^4 \times (\mathbb{Z}/2\mathbb{Z})$ is the identity on the cyclic group of order 2 and an injection on the free abelian part.

In general we have for a word w of length n in Ω_n that

$$(1 - \Phi)_n(\chi_w) = \chi_{\alpha_1 w} + \chi_{\alpha_2 w} - \chi_{w\alpha_1} - \chi_{w\alpha_2} - \chi_{\alpha_1 \alpha_1 w_{n-1}} - \chi_{\alpha_2 \alpha_1 w_{n-1}} - \chi_{\alpha_1 \alpha_2 w_{n-1}} - \chi_{\alpha_2 \alpha_2 w_{n-1}}$$

where w_{n-1} represents the first $n - 1$ symbols of w . Again, a short calculation shows that $\ker(1 - \Phi)_n = \{0\}$. For a word $w \in \Omega_{n+1}$, let $x_w = \chi_w + \text{Im}(1 - \Phi)_n$. Then $2 \sum_{w \in \Omega_{n+1}} x_w = 0$, because

$$(1 - \Phi)_n \left(\sum_{w' \in \Omega_n} \chi_{w'} \right) = 2 \sum_{w \in \Omega_{n+1}} \chi_w.$$


Using this, one can show that $\text{coker}(1 - \Phi)_n \cong \mathbb{Z}^{2^n} \times (\mathbb{Z}/2\mathbb{Z})$, generated by $\{x_{w'2} : w' \in \Omega_n\} \cup \{\sum_{w \in \Omega_{n+1}} x_w\}$.

Observe that \tilde{i}_n is induced by the map i_n which only adds symbols to the beginning of words. It follows that $\tilde{i}_n : \mathbb{Z}^{2^n} \times (\mathbb{Z}/2\mathbb{Z}) \rightarrow \mathbb{Z}^{2^{n+1}} \times (\mathbb{Z}/2\mathbb{Z})$ is the identity on the cyclic group of order 2 and an injection on the free abelian part. From Theorem 6.5 we have that

$$K_0(C^*(E_2, \mathcal{L}_2, \mathcal{E}_2^{0,-})) \cong \left(\prod_{i=1}^{\infty} \mathbb{Z} \right) \times (\mathbb{Z}/2\mathbb{Z}) \text{ and } K_1(C^*(E_2, \mathcal{L}_2, \mathcal{E}_2^{0,-})) \cong 0.$$

This confirms the calculations done in [24, §3]. It is worth noting that $C^*(E_2, \mathcal{L}_2, \mathcal{E}_2^{0,-})$ is unital (the unit is $p_{r(\alpha_1)} + p_{r(\alpha_2)}$) and has a K_0 -group which is not finitely generated, and so cannot be isomorphic to a graph algebra.

Example 7.4. Consider the following left-resolving labelled graph (E, \mathcal{L}) over the infinite alphabet $\{0, 1\} \times \mathbb{Z}$:



We write $E^0 = \{(v, i), (w, i) : i \in \mathbb{Z}\}$, and label the edges as shown. Since (v, i) is the only vertex which receives an edge labelled $(0, i - 1)$, we have $[(v, i)]_\ell = \{(v, i)\}$ for all $i \in \mathbb{Z}$ and $\ell \geq 1$. Since (w, i) is the only vertex which receives an edge with label $(1, i - 1)$, but does not receive any edge with label $(0, j)$ for any $j \in \mathbb{Z}$, we have $[(w, i)]_\ell = \{(w, i)\}$ for all $i \in \mathbb{Z}$ and $\ell \geq 1$. We will simply identify $[(v, i)]_\ell = \{(v, i)\}$ with (v, i) and $[(w, i)]_\ell = \{(w, i)\}$ with (w, i) for all $i \in \mathbb{Z}$. We then have that $\Omega_\ell = E^0$ for all $\ell \geq 1$.

Since $\Omega_\ell = E^0$ for each $\ell \geq 1$, and E^0 is infinite, we have

$$\mathbb{Z}(\Omega_\ell) = \bigoplus_{E^0} \mathbb{Z} = \text{span}_{\mathbb{Z}}\{\chi_{(v,i)}, \chi_{(w,i)} : i \in \mathbb{Z}\}.$$

Since (v, i) only connects to $(v, i + 1)$ and $(w, i + 1)$ and (w, i) only connects to $(v, i + 1)$ for all $i \in \mathbb{Z}$, we have

$$(15) \quad (1 - \Phi)_\ell(\chi_{(v,i)}) = \chi_{(v,i)} - \chi_{(v,i+1)} - \chi_{(w,i+1)} \text{ and}$$

$$(16) \quad (1 - \Phi)_\ell(\chi_{(w,i)}) = \chi_{(w,i)} - \chi_{(v,i+1)}.$$

We claim that $\ker(1 - \Phi)_\ell = \{0\}$ for all $\ell \geq 1$.

Suppose, for contradiction, that $b \in \ker((1 - \Phi)_\ell) \subseteq \bigoplus_{E^0} \mathbb{Z}$ is a nonzero vector. Since $b = \sum_{i \in \mathbb{Z}} b_{(v,i)} \chi_{(v,i)} + b_{(w,i)} \chi_{(w,i)}$ is in the direct sum there must be a maximum i such that not both $b_{(v,i)}$ and $b_{(w,i)}$ are 0. Then by (15) we have $b_{(v,i)} = b_{(v,i+1)} + b_{(w,i+1)} = 0$, and by (16) we have $b_{(w,i)} = b_{(v,i+1)} = 0$ which is a contradiction. Hence $\ker(1 - \Phi)_\ell = \{0\}$ for all $\ell \geq 1$.

Now we compute the cokernel of $(1 - \Phi)_\ell$. From Equations (15) and (16) we see that

$$(17) \quad (1 - \Phi)_\ell(\chi_{(v,i)} - \chi_{(w,i)}) = \chi_{(v,i)} - \chi_{(w,i)} - \chi_{(w,i+1)}.$$

Repeated use of Equation (17) shows that every vector in $\mathbb{Z}(\Omega_\ell)$ is equivalent, using vectors in $\text{Im}(1 - \Phi)_\ell$, to a vector which has all (v, i) -coordinates zero. From Equations (15) and (16) we also see that

$$(18) \quad (1 - \Phi)_\ell(\chi_{(v,i)} + \chi_{(w,i-1)} - \chi_{(w,i)}) = \chi_{(w,i-1)} - \chi_{(w,i)} - \chi_{(w,i+1)}.$$

Repeated use of Equation (18) shows that every vector in $\mathbb{Z}(\Omega_\ell)$ is equivalent, using vectors in $\text{Im}(1 - \Phi)_\ell$, to a vector which has all (v, i) -coordinates zero and all (w, i) -coordinates zero except for $(w, 0)$ and $(w, 1)$, with no relation between them. Hence $\text{coker}(1 - \Phi)_\ell$ is generated by $\chi_{(w,0)} + \text{Im}(1 - \Phi)_\ell$ and $\chi_{(w,1)} + \text{Im}(1 - \Phi)_\ell$ for all $\ell \geq 1$ and so we may conclude that $\text{coker}(1 - \Phi)_\ell \cong \mathbb{Z}^2$ for all $\ell \geq 1$.

Since \tilde{i}_ℓ is the identity map for all $\ell \geq 1$, we have


$$K_0(C^*(E, \mathcal{L}, \mathcal{E}^{0,-})) \cong \mathbb{Z}^2 \text{ and } K_1(C^*(E, \mathcal{L}, \mathcal{E}^{0,-})) \cong \{0\}.$$

APPENDIX A. C^* -ALGEBRAS OF NON-NORMAL LABELLED SPACES

In this appendix we give an example showing that if $(E, \mathcal{L}, \mathcal{B})$ is a weakly left-resolving labelled space which is not normal, and we define a representation of $(E, \mathcal{L}, \mathcal{B})$ as in Definition 2.1 and $C^*(E, \mathcal{L}, \mathcal{B})$ as in Definition 2.5, then $C^*(E, \mathcal{L}, \mathcal{B})$ might not satisfy the gauge invariant uniqueness theorem (Corollary 3.10). We will then describe how to modify the definition of a representation of a non-normal weakly left-resolving labelled space, and consequently of $C^*(E, \mathcal{L}, \mathcal{B})$, in order for $C^*(E, \mathcal{L}, \mathcal{B})$ to satisfy the gauge invariant uniqueness theorem as well as Theorem 3.8, Corollary 3.9, and Corollary 3.11.

Example A.1. Let E be the graph with $E^0 = \{v_0, v_1\}$, $E^1 = \{e_n : n \in \mathbb{N}\}$, $r(e_0) = s(e_0) = v_0$, and $r(e_n) = s(e_n) = v_1$ for $n > 0$. Define a labelling map $\mathcal{L} : E^1 \rightarrow \{a_n : n \in \{1, 2, \dots\}\}$ by

$\mathcal{L}(e_0) = a_1$ and $\mathcal{L}(e_n) = a_n$ for $n > 0$.



Then (E, \mathcal{L}) is a left-resolving labelled graph. Let $\mathcal{B} = \{E^0, \{v_1\}\}$. Then $(E, \mathcal{L}, \mathcal{B})$ is a weakly left-resolving labelled space, but it is not normal.

We will now construct two representations of $(E, \mathcal{L}, \mathcal{B})$. Let H_1 be a Hilbert space with an orthonormal basis $(e_{(x,n)})_{(x,n) \in (\mathbb{N} \cup \mathbb{N}^{\mathbb{N}}) \times \mathbb{Z}}$ indexed by $(\mathbb{N} \cup \mathbb{N}^{\mathbb{N}}) \times \mathbb{Z}$ (we can, for example, let H_1 be $l^2((\mathbb{N} \cup \mathbb{N}^{\mathbb{N}}) \times \mathbb{Z})$) and define bounded operators $q_{E^0}, q_{\{v_1\}}, t_{a_1}, t_{a_2}, \dots$ on H_1 in the following way:

$$q_{E^0}e_{(x,n)} = e_{(x,n)} \text{ for all } x \in \mathbb{N} \cup \mathbb{N}^{\mathbb{N}}, \quad q_{\{v_1\}}e_{(x,n)} = \begin{cases} 0 & \text{if } x \in \mathbb{N}, \\ e_{(x,n)} & \text{if } x \in \mathbb{N}^{\mathbb{N}}, \end{cases}$$

$$t_{a_1}e_{(x,n)} = \begin{cases} e_{(x+1,n)} & \text{if } x \in \mathbb{N}, \\ e_{(1x,n)} & \text{if } x \in \mathbb{N}^{\mathbb{N}}, \end{cases} \quad t_{a_i}e_{(x,n)} = \begin{cases} 0 & \text{if } x \in \mathbb{N}, \\ e_{(ix,n)} & \text{if } x \in \mathbb{N}^{\mathbb{N}}, \end{cases} \text{ for } i \geq 2.$$

Then q_{E^0} and $q_{\{v_1\}}$ are projections and t_{a_1}, t_{a_2}, \dots are partial isometries satisfying

- (i) If $A, B \in \mathcal{B}$, then $q_A q_B = q_{A \cap B}$ and $q_{A \cup B} = q_A + q_B - q_{A \cap B}$, where $q_\emptyset = 0$.
- (ii) If $a \in \mathcal{A}$ and $A \in \mathcal{B}$, then $q_A t_a = t_a q_{r(A,a)}$.
- (iii) If $a, b \in \mathcal{A}$, then $t_a^* t_a = p_{r(a)}$ and $t_a^* t_b = 0$ unless $a = b$.
- (iv) For $A \in \mathcal{B}$ with $\mathcal{L}(AE^1)$ finite and $A \cap B = \emptyset$ for all $B \in \mathcal{B}$ satisfying $B \subseteq E_{\text{sink}}^0$, we have

$$q_A = \sum_{a \in \mathcal{L}(AE^1)} t_a q_{r(A,a)} t_a^*$$

(condition (iv) is satisfied because there are no $A \in \mathcal{B}$ with $\mathcal{L}(AE^1)$ finite). For each $z \in \mathcal{T}$ there is an automorphism β_z on $B(H_1)$ given by $\beta_z(x) = u_z x u_z^*$ for $x \in B(H_1)$ where u_z is the unitary operator given by $u_z e_{(x,n)} = e_{(x,n+1)}$ for $(x,n) \in (\mathbb{N} \cup \mathbb{N}^{\mathbb{N}}) \times \mathbb{Z}$. Then $\beta_z(q_A) = q_A$ for all $A \in \mathcal{B}$, and $\beta_z(t_a) = z t_a$ for all $a \in \mathcal{A}$.

Similarly, let H_2 be a Hilbert space with an orthonormal basis $(f_y)_{y \in \mathbb{Z} \cup \{\mathbb{N}^{\mathbb{N}}\}}$ indexed by $(\mathbb{Z} \cup \mathbb{N}^{\mathbb{N}}) \times \mathbb{Z}$ and define bounded operators $r_{E^0}, r_{\{v_1\}}, u_{a_1}, u_{a_2}, \dots$ on H_2 in the following way:

$$r_{E^0}f_{(y,n)} = f_{(y,n)} \text{ for all } y \in \mathbb{Z} \cup \mathbb{N}^{\mathbb{N}}, \quad r_{\{v_1\}}f_{(y,n)} = \begin{cases} 0 & \text{if } y \in \mathbb{Z}, \\ f_{(y,n)} & \text{if } y \in \mathbb{N}^{\mathbb{N}}, \end{cases}$$

$$u_{a_1}f_{(y,n)} = \begin{cases} f_{(y+1,n)} & \text{if } y \in \mathbb{Z}, \\ f_{(1y,n)} & \text{if } y \in \mathbb{N}^{\mathbb{N}}, \end{cases} \quad u_{a_i}f_{(y,n)} = \begin{cases} 0 & \text{if } y \in \mathbb{Z}, \\ f_{(iy,n)} & \text{if } y \in \mathbb{N}^{\mathbb{N}}, \end{cases} \text{ for } i \geq 2.$$

Then r_{E^0} and $r_{\{v_1\}}$ are projections and u_{a_1}, u_{a_2}, \dots are partial isometries satisfying

- (i) If $A, B \in \mathcal{B}$, then $r_A r_B = r_{A \cap B}$ and $r_{A \cup B} = r_A + r_B - r_{A \cap B}$, where $r_\emptyset = 0$.
- (ii) If $a \in \mathcal{A}$ and $A \in \mathcal{B}$, then $r_A u_a = u_a r_{r(A,a)}$.
- (iii) If $a, b \in \mathcal{A}$, then $u_a^* u_a = r_{r(a)}$ and $u_a^* u_b = 0$ unless $a = b$.

(iv) For $A \in \mathcal{B}$ with $\mathcal{L}(AE^1)$ finite and $A \cap B = \emptyset$ for all $B \in \mathcal{B}$ satisfying $B \subseteq E_{\text{sink}}^0$, we have

$$r_A = \sum_{a \in \mathcal{L}(AE^1)} u_a r_{r(A,a)} u_a^*$$

For each $z \in \mathbf{T}$ there is an automorphism γ_z on $B(H_2)$ given by $\gamma_z(x) = v_z x v_z^*$ for $x \in B(H_2)$ where v_z is the unitary operator given by $v_z e_{(y,n)} = e_{(y,n+1)}$ for $(y,n) \in (\mathbb{Z} \cup \mathbb{N}^{\mathbb{N}}) \times \mathbb{Z}$. Then $\gamma_z(r_A) = r_A$ for all $A \in \mathcal{B}$, and $\gamma_z(u_a) = z u_a$ for all $a \in \mathcal{A}$.

Suppose we define a representation of $(E, \mathcal{L}, \mathcal{B})$ as in Definition 2.1 and let $C^*(E, \mathcal{L}, \mathcal{B})$ be the C^* -algebra generated by a universal representation of $(E, \mathcal{L}, \mathcal{B})$. Then $C^*(E, \mathcal{L}, \mathcal{B})$ cannot satisfy a gauge invariant uniqueness theorem because that would imply that there is a $*$ -isomorphism from the C^* -algebra generated by $\{q_{E^0}, q_{\{v_1\}}, t_{a_1}, t_{a_2}, \dots\}$ to the C^* -algebra generated by $\{r_{E^0}, r_{\{v_1\}}, u_{a_1}, u_{a_2}, \dots\}$ mapping q_{E^0} to r_{E^0} , $q_{\{v_1\}}$ to $r_{\{v_1\}}$, and t_{a_i} to u_{a_i} for each i . This cannot be the case since $q_{E^0} - q_{\{v_1\}} \neq (q_{E^0} - q_{\{v_1\}})t_{a_1}t_{a_1}^*$ whereas $r_{E^0} - r_{\{v_1\}} = (r_{E^0} - r_{\{v_1\}})u_{a_1}u_{a_1}^*$.

Remark A.2. Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving labelled space. Even if $(E, \mathcal{L}, \mathcal{B})$ is not normal the C^* -algebra $\mathcal{A}(E, \mathcal{L}, \mathcal{B})$ and the C^* -correspondence $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$ can still be defined as in Section 3. Let

$$\mathcal{P} = \{(A, B) : A \in \mathcal{B}, B \in \mathcal{B} \cup \emptyset, B \subseteq A\}$$

and suppose $(A, B) \in \mathcal{P}$. Notice that $r(B, a) \subseteq r(A, a)$ for all $a \in \mathcal{A}$ (because $B \subseteq A$). Let

$$\mathcal{A}_{(A,B)} = \{a \in \mathcal{A} : r(B, a) \neq r(A, a)\}$$

and let

$$\mathcal{N} = \{A \setminus B : (A, B) \in \mathcal{P}, \mathcal{A}_{(A,B)} = \emptyset\}.$$

An analysis of $(X(E, \mathcal{L}, \mathcal{B}), \varphi)$ similar to the one given in Section 3 shows that $\mathcal{O}_{(X(E, \mathcal{L}, \mathcal{B}), \varphi)}$ is the universal C^* -algebra generated by a family $\{p_A : A \in \mathcal{B}\}$ of projections and a family $\{s_a : a \in \mathcal{A}\}$ of partial isometries satisfying

- (i') If $A, B \in \mathcal{B}$, then $p_A p_B = p_{A \cap B}$ and $p_{A \cup B} = p_A + p_B - p_{A \cap B}$, where $p_\emptyset = 0$.
- (ii') If $a \in \mathcal{A}$ and $A \in \mathcal{B}$, then $p_A s_a = s_a p_{r(A,a)}$.
- (iii') If $a, b \in \mathcal{A}$, then $s_a^* s_a = p_{r(a)}$ and $s_a^* s_b = 0$ unless $a = b$.
- (iv') For $(A, B) \in \mathcal{P}$ with $\mathcal{A}_{(A,B)}$ finite and $A \cap C = B \cap C$ for all $C \in \mathcal{N}$ we have

$$p_A = p_B + \sum_{a \in \mathcal{A}_{(A,B)}} s_a (p_{r(A,a)} - p_{r(B,a)}) s_a^*.$$

We therefore define a representation of $(E, \mathcal{L}, \mathcal{B})$ and $C^*(E, \mathcal{L}, \mathcal{B})$ in the following way.

Definition A.3. Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving labelled space. A *representation* of $(E, \mathcal{L}, \mathcal{B})$ in a C^* -algebra consists of projections $\{p_A : A \in \mathcal{B}\}$ and partial isometries $\{s_a : a \in \mathcal{A}\}$ satisfying the conditions (i')–(iv') above.

Definition A.4. Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving labelled space. Then $C^*(E, \mathcal{L}, \mathcal{B})$ is the C^* -algebra generated by a universal representation of $(E, \mathcal{L}, \mathcal{B})$.

Remark A.5. If $(E, \mathcal{L}, \mathcal{B})$ is a weakly left-resolving normal labelled space, then the definition of a representation of $(E, \mathcal{L}, \mathcal{B})$ given in Definition A.3 agrees with the definition of a representation of $(E, \mathcal{L}, \mathcal{B})$ given in Definition 2.1.

Remark A.6. If $(E, \mathcal{L}, \mathcal{B})$ is a weakly left-resolving labelled space such that $r(A \setminus B, a) = r(A, a) \setminus r(B, a)$ for all $(A, B) \in \mathcal{P}$ and all $a \in \mathcal{A}$ (this will always be the case if (E, \mathcal{L}) is left-resolving), and we let $\tilde{\mathcal{B}}$ be the collection of subsets of E^0 which can be written as a

finite disjoint union of sets of the form $A \setminus B$ where $(A, B) \in \mathcal{P}$, then $(E, \mathcal{L}, \tilde{\mathcal{B}})$ is a weakly left-resolving normal labelled space and $C^*(E, \mathcal{L}, \mathcal{B}) \cong C^*(E, \mathcal{L}, \tilde{\mathcal{B}})$.

Remark A.7. It follows from Remark A.2 that with the definition of a representation of a weakly left-resolving labelled space $(E, \mathcal{L}, \mathcal{B})$ given in Definition A.3 and the definition of the C^* -algebra $C^*(E, \mathcal{L}, \mathcal{B})$ given in Definition A.4 that Theorem 3.8, and therefore also Corollary 3.9 and Corollary 3.11, remain true if the assumption that $(E, \mathcal{L}, \mathcal{B})$ is normal is dropped.

We also get the following gauge-invariant uniqueness Theorem for $C^*(E, \mathcal{L}, \mathcal{B})$.

Corollary A.8. *Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving labelled space. Let $\{p_A, s_a : A \in \mathcal{B}, a \in \mathcal{A}\}$ be the universal representation of $(E, \mathcal{L}, \mathcal{B})$ that generates $C^*(E, \mathcal{L}, \mathcal{B})$, let $\{q_A, t_a : A \in \mathcal{B}, a \in \mathcal{A}\}$ be a representation of $(E, \mathcal{L}, \mathcal{B})$ in a C^* -algebra \mathcal{X} and let π be the unique $*$ -homomorphism from $C^*(E, \mathcal{L}, \mathcal{B})$ to \mathcal{X} that maps each p_A to q_A and each s_a to t_a . Then π is injective if and only if $q_A = q_B$ implies that $A = B$ for all $(A, B) \in \mathcal{P}$, and for each $z \in \mathbb{T}$ there exists a $*$ -homomorphism $\beta_z : C^*(\{q_A, t_a : A \in \mathcal{B}, a \in \mathcal{A}\}) \rightarrow C^*(\{q_A, t_a : A \in \mathcal{B}, a \in \mathcal{A}\})$ such that $\beta_z(q_A) = q_A$ and $\beta_z(t_a) = zt_a$ for $A \in \mathcal{B}$ and $a \in \mathcal{A}$.*

The K -theory of $C^*(E, \mathcal{L}, \mathcal{B})$ can be determined by the following adaptation of Theorem 4.4.

Theorem A.9. *Let $(E, \mathcal{L}, \mathcal{B})$ be a weakly left-resolving labelled space. Let*

$$\mathcal{P}_J = \{(A, B) \in \mathcal{P} : \mathcal{A}_{(A,B)} \text{ is finite, and } A \cap C = B \cap C \text{ for all } C \in \mathcal{N}\},$$

let

$$\mathcal{B}_J = \{A \setminus B : (A, B) \in \mathcal{P}_J\},$$

and let $(1 - \Phi)$ be the linear map from $\text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}_J\}$ to $\text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}\}$ given by

$$(1 - \Phi)(\chi_{A \setminus B}) = \chi_{A \setminus B} - \sum_{a \in \mathcal{A}_{(A,B)}} \chi_{r(A,a) \setminus r(B,a)}, \quad (A, B) \in \mathcal{P}_J.$$

Then $K_1(C^*(E, \mathcal{L}, \mathcal{B})) \cong \ker(1 - \Phi)$, and $K_0(C^*(E, \mathcal{L}, \mathcal{B})) \cong \text{span}_{\mathbb{Z}}\{\chi_A : A \in \mathcal{B}\} / \text{Im}(1 - \Phi)$ via $[p_A]_0 \mapsto \chi_A + \text{Im}(1 - \Phi)$ for $A \in \mathcal{B}$.

If $(E, \mathcal{L}, \mathcal{B})$ is normal, then the definitions of \mathcal{B}_J and $(1 - \Phi)$ given in the theorem above agree with the definition of \mathcal{B}_J given by Equation (2) and the definition of $(1 - \Phi)$ given in Theorem 4.4, so the theorem above reduces to Theorem 4.4 in this case.

REFERENCES

- [1] Astrid an Huef and Iain Raeburn. The ideal structure of Cuntz–Krieger algebras. *Ergodic Theory & Dynam. Systems* **17** (1997), 611–624.
- [2] Teresa Bates, Søren Eilers and David Pask. Reducibility of covers of AFT shifts. *Israel J. Math.*, **185** (2011), 207–234.
- [3] Teresa Bates and David Pask. Flow equivalence of graph algebras. *Ergodic Theory & Dynam. Systems* **24** (2004), 367–382.
- [4] Teresa Bates and David Pask. C^* -algebras of labelled graphs. *J. Operator Theory* **57** (2007), 207–226.
- [5] Teresa Bates and David Pask. C^* -algebras of labelled graphs II - simplicity results. *Math. Scand.* **104** (2009), 249–274.
- [6] Teresa Bates, David Pask and Paulette Willis. Group actions on labeled graphs and their C^* -algebras *Illinois J. Math.* **56** (2012), 1149–1168.
- [7] Rufus Bowen and John Franks. Homology for zero-dimensional nonwandering sets. *Ann. Math.* **106** (1977), 73–92.
- [8] Toke Meier Carlsen. Cuntz-Pimsner C^* -algebras associated with subshifts. *Internat. J. Math.* **19** (2008), 47–70.
- [9] Toke Meier Carlsen and Kengo Matsumoto. Some remarks on the C^* -algebras associated with subshifts. *Math. Scand.* **95** (2004), 145–160.

- [10] Toke Meier Carlsen and Sergei Silvestrov. C*-crossed products and shift spaces. *Expo. Math.* **25** (2007), 275–307.
- [11] Toke Meier Carlsen and Sergei Silvestrov. On the K-theory of the C*-algebra associated to a one-sided shift space. *Proc. Est. Acad. Sci.* **59** (2010), 272–279.
- [12] Joachim Cuntz and Wolfgang Krieger. A class of C*-algebras and topological Markov chains. *Invent. Math.* **56** (1980), 251–268.
- [13] Joachim Cuntz. A class of C*-Algebras and Topological Markov Chains: Reducible Chains and the Ext-functor for C*-Algebras. *Invent. Math.* **63** (1981), 25–40.
- [14] Doug Drinen and Mark Tomforde. Computing K-theory and Ext for graph C*-algebras. *Illinois J. Math.* **46** (2002), 81–91.
- [15] Masatoshi Enomoto, Yasuo Watatani. A graph theory for C*-algebras. *Math. Japon.* **25** (1980), 435–442.
- [16] Ruy Exel and Marcelo Laca. Cuntz-Krieger algebras for infinite matrices. *J. Reine Angew. Math.* **512** (1999), 119–172.
- [17] Neal Fowler, Marcelo Laca, and Iain Raeburn. The C*-algebras of infinite graphs. *Proc. Amer. Math. Soc.* **128** (2000), 2319–2327.
- [18] Neal Fowler and Iain Raeburn. The Toeplitz algebra of a Hilbert bimodule. *Indiana Univ. Math. J.* **48** (1999), 155–181.
- [19] Ja Jeong and Sun Kim. On simple labelled graph C*-algebras. *J. Math. Anal. Appl.* **386** (2012), 631–640.
- [20] Takeshi Katsura. On C*-algebras associated with C*-correspondences. *J. Funct. Anal.* **217** (2004), 366–401.
- [21] Takeshi Katsura. Nonseparable AF algebras. *Operator Algebras: The Abel Symposium 2004*, 165–173, Abel Symp., **1**, Springer, Berlin, 2006.
- [22] Takeshi Katsura, Paul S. Muhly, Aidan Sims, and Mark Tomforde. Ultragraph C*-algebras via topological quivers. *Studia Math.* **187** (2008), 137–155.
- [23] Wolfgang Krieger. On sofic systems I. *Israel J. Math.* **48** (1984), 305–330.
- [24] Wolfgang Krieger and Kengo Matsumoto. A Lambda-Graph system for the Dyck shift and its K-Groups. *Doc. Math.* **8** (2003), 79–96.
- [25] Alex Kumjian, David Pask and Iain Raeburn. Cuntz–Krieger algebras of directed graphs, *Pacific. J. Math.* **184** (1998), 161–174.
- [26] Doug Lind and Brian Marcus. An Introduction to Symbolic Dynamics and Coding. Cambridge University Press, Cambridge, 1995.
- [27] Kengo Matsumoto. On C*-algebras associated with subshifts. *Internat. J. Math.* **8** (1997), 357–374.
- [28] Kengo Matsumoto. K-theory for C*-algebras associated with subshifts. *Math. Scand.* **82** (1998), 237–255.
- [29] Kengo Matsumoto. Dimension groups for subshifts and simplicity of the associated C*-algebras. *J. Math. Soc. Japan* **51** (1999), 679–698.
- [30] Kengo Matsumoto. Relations among generators of C*-algebras associated with subshifts. *Internat. J. Math.* **10** (1999), 385–405.
- [31] Kengo Matsumoto. Presentations of subshifts and their topological conjugacy invariants. *Doc. Math.* **4** (1999), 285–340.
- [32] Kengo Matsumoto. On automorphisms of C*-algebras associated with subshifts. *J. Operator Theory* **44** (2000), 91–112.
- [33] Kengo Matsumoto. Stabilized C*-algebras constructed from symbolic dynamical systems. *Ergodic Theory & Dynam. Systems* **20** (2000), 821–841.
- [34] Kengo Matsumoto. Bowen-Franks groups as an invariant for flow equivalence of subshifts. *Ergodic Theory & Dynam. Systems* **21** (2001), 1831–1842.
- [35] Kengo Matsumoto. Bowen-Franks groups for subshifts and Ext-groups for C*-algebras. *K-Theory* **23** (2001), 67–104.
- [36] Kengo Matsumoto. C*-algebras associated with presentations of subshifts. *Doc. Math.* **7** (2002), 1–30.
- [37] Michael V. Pimsner. A class of C*-algebras generalizing both Cuntz-Krieger algebras and crossed products by \mathbf{Z} *Free probability theory* (Waterloo, ON, 1995), 189–212, Fields Inst. Commun., **12**, Amer. Math. Soc., Providence, RI, 1997.
- [38] David Robertson and Wojciech Szymański. C*-algebras associated to C*-correspondences and applications to mirror quantum spheres. *Illinois J. Math.* **55** (2011), 845–870.
- [39] Mark Tomforde. A unified approach to Exel-Laca algebras and C*-algebras associated to graphs. *J. Operator Theory* **50** (2003), 345–368.
- [40] Mark Tomforde. Simplicity of ultragraph algebras. *Indiana Univ. Math. J.* **52** (2003), 901–926.
- [41] Iain Raeburn and Wojciech Szymański. Cuntz-Krieger algebras of infinite graphs and matrices. *Trans. Amer. Math. Soc.* **356** (2003), 39–59.

- [42] Jonathan Rosenberg and Claude Schochet. The Künneth theorem and the universal coefficient theorem for Kasparov's generalized K -functor. *Duke Math. J.* **55** (1987), 431–474.

TERESA BATES, SCHOOL OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF NSW, UNSW SYDNEY 2052, AUSTRALIA

E-mail address: `teresa@unsw.edu.au`

TOKE MEIER CARLSEN, DEPARTMENT OF MATHEMATICAL SCIENCES, NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY (NTNU), 7491 TRONDHEIM, NORWAY

E-mail address: `Toke.Meier.Carlsen@math.ntnu.no`

DAVID PASK, SCHOOL OF MATHEMATICS AND APPLIED STATISTICS, AUSTIN KEANE BUILDING (15), UNIVERSITY OF WOLLONGONG, NSW 2522, AUSTRALIA

E-mail address: `dpask@uow.edu.au`