## Flavor-Tuned 125 GeV SUSY Higgs Boson at the LHC : MSSM and NATURAL SUSY TESTS

Vernon Barger\*, Peisi Huang\*, Muneyuki Ishida<sup>†</sup>, and Wai-Yee Keung<sup>‡</sup>

<sup>∗</sup> Department of Physics, University of Wisconsin, Madison, WI 53706, USA

† Department of Physics, Meisei University, Hino, Tokyo 191-8506, Japan

‡ Department of Physics, University of Illinois at Chicago, IL 60607, USA

(Dated: October 26, 2018)

## Abstract

We show that an enhanced two-photon signal of the Higgs boson, h, observed with 125 GeV mass by the ATLAS and CMS collaborations, can be obtained if it is identified principally with the neutral  $H_u^0$  of the two Higgs doublets of minimal Supersymmetry. We focus on sparticles and the pseudoscalar Higgs A at the TeV scale. The off-diagonal element of the  $(H_u^0, H_d^0)$  mass matrix in the flavor basis must be suppressed, and this requires both a large Higgsino mass parameter,  $\mu \sim TeV$ , and large tan $\beta$ . A MSSM sum rule is derived that relates  $\gamma\gamma$  and  $b\bar{b}$  rates, and a  $\gamma\gamma$  enhancement relative to the SM predicts the  $b\bar{b}$  reduction. On the contrary, Natural SUSY requires  $|\mu| \leq 0.5$  TeV, for which  $\gamma\gamma$  is reduced and  $b\bar{b}$  is enhanced. This conclusion is independent of the  $m_A$ -value and the SUSY quantum correction  $\Delta_b$ . Relative  $\tau\bar{\tau}$  to  $b\bar{b}$  rates are sensitive to  $\Delta_b$ .

A  $\gamma\gamma$  enhancement of the 125 GeV Higgs boson signal relative to the Standard Model (SM) expectation has been reported by the ATLAS and CMS experiments at the LHC[\[1](#page-7-0), [2\]](#page-7-1). We investigate this in the Minimal Supersymmetric Standard Model (MSSM) in the region of large  $m_A \sim \text{TeV}$  by flavor-tuning of the mixing angle  $\alpha$  between two neutral CP-even Higgs flavor states  $H_u^0$  and  $H_d^0$ , with the 125 GeV Higgs signal identified principally with  $H_u^0$ . Then, the  $b\bar{b}$  decay, which is predicted to be dominant decay of the SM Higgs boson, is reduced and the cross sections of the other channels are correspondingly enhanced, except possibly the  $\tau\tau$  channel. We relate the cross-section enhancements/suppressions in  $\gamma\gamma/bb/\tau\tau$ channels compared with those of the SM Higgs boson. We also consider the consequences for Natural SUSY[\[3](#page-7-2)]. Our focus is on a heavy pseudoscalar A and large  $\tan \beta \equiv \langle H_u^0 \rangle / \langle H_d^0 \rangle$ , a region that has not yet been constrained by LHC experiments[\[5\]](#page-9-0). Light stau[\[6](#page-9-1), [7\]](#page-9-2) and light stop[\[8\]](#page-9-3) scenarios that have been considered are outside of our purview.

Ratios of the SUSY Higgs couplings to those of the SM Higgs The SUSY Higgs mechanism is based on the two Higgs doublet model of type II[\[9](#page-9-4)[–11](#page-9-5)] with the  $H_u$  doublet coupled to uptype quarks and the  $H_d$  doublet coupled to down-type quarks. After spontaneous symmetry breaking, the physical Higgs states are two CP-even neutral Higgs  $h, H$ , one CP-odd neutral pseudo-scalar A and the charged Higgs  $H^{\pm}$ .

We focus on the CP-even neutral Higgs boson  $h$  and  $H$ , which are related to the flavor eigenstates  $H_u^0$  and  $H_d^0$  by

$$
\frac{h}{\sqrt{2}} = c_{\alpha}H_u^0 - s_{\alpha}H_d^0 \ , \quad \frac{H}{\sqrt{2}} = s_{\alpha}H_u^0 + c_{\alpha}H_d^0 \ , \tag{1}
$$

where  $H_{u,d}^0$  is the shorthand for the real part of  $H_{u,d}^0 - \langle H_{u,d}^0 \rangle$ . We use the notation  $s_\alpha = \sin \alpha$ ,  $c_{\alpha} = \cos \alpha$ , and  $t_{\alpha} = \tan \alpha$ . Our interest is in large  $\tan \beta$ ,  $\tan \beta \stackrel{>}{\sim} 20$ , and in the decoupling regime with large  $m_A$  for which  $\alpha \simeq \beta - \frac{\pi}{2}$  $\frac{\pi}{2}$ .

The ratios of the h and H couplings to those of the SM Higgs  $h_{SM}$ , denoted as  $r_{PP}^{h,H}$  ( $\equiv$  $g_{h,H}$   $_{P\bar{P}}/g_{h_{SM}P\bar{P}}$ , are given by

<span id="page-1-0"></span>
$$
r_{VV}^h = s_{\beta-\alpha}, \quad r_t^h = r_{cc}^h = \frac{c_{\alpha}}{s_{\beta}}, \quad r_{\tau\tau}^h = \frac{-s_{\alpha}}{c_{\beta}}, \quad r_{bb}^h = \frac{-s_{\alpha}}{c_{\beta}} \left[ 1 - \frac{\Delta_b}{1 + \Delta_b} (1 + \frac{1}{t_{\alpha} t_{\beta}}) \right]
$$

$$
r_{VV}^H = c_{\beta-\alpha}, \quad r_t^H = r_{cc}^H = \frac{s_{\alpha}}{s_{\beta}}, \quad r_{\tau\tau}^H = \frac{c_{\alpha}}{c_{\beta}}, \quad r_{bb}^H = \frac{c_{\alpha}}{c_{\beta}} \left[ 1 - \frac{\Delta_b}{1 + \Delta_b} (1 - \frac{t_{\alpha}}{t_{\beta}}) \right]
$$
(2)

where we include the 1-loop contribution  $\Delta_b$  to the  $b\bar{b}$  coupling. It is the b-quark mass

correction factor [\[12](#page-9-6), [13\]](#page-9-7), which may be sizable, especially if both  $\mu$  and  $\tan\beta$  are large.

$$
\Delta_b = \bar{\mu} t_\beta \left[ \frac{2\alpha_s}{3\pi} \hat{m}_{\tilde{g}} I(\hat{m}_{\tilde{g}}^2, \hat{m}_{\tilde{b}_1}^2, \hat{m}_{\tilde{b}_2}^2) + \frac{h_t^2}{16\pi^2} a_t I(\bar{\mu}^2, \hat{m}_{\tilde{t}_1}^2, \hat{m}_{\tilde{t}_2}^2) \right]
$$
(3)  

$$
I(x, y, z) = -\frac{xy \ln x/y + yz \ln y/z + zx \ln z/x}{(x - y)(y - z)(z - x)}
$$
  

$$
I(x, y, z = y) = -\left[ x - y + x \log \frac{y}{x} \right] / (x - y)^2, \quad I(x, x, x) = \frac{1}{2x}.
$$
(4)

The first(second) term of  $\Delta_b$  is due to the sbottom-gluino(stop-chargino) loop. We take  $M_{\text{susy}} = 1$  TeV and sparticle masses  $\hat{m}$  in units of  $M_{\text{susy}}$ . The top Yukawa coupling is  $h_t = \bar{m}_t/v_u = \bar{m}_t/(v \ s_\beta)$  and  $\bar{m}_t = m_t(\bar{m}_t) = 163.5 \ \text{GeV}$  is the running top quark mass[\[14\]](#page-9-8). We consider  $m_Q = m_U = m_D = M_{\text{susy}}$  for the squark masses in the third generation.

The off-diagonal element of the stop squared mass matrix is  $\bar{m}_t X_t$  where the stop mixing parameter  $X_t$  is given by  $X_t = A_t - \mu/t_\beta$ . These quantities are also defined in units of  $M_{\text{susy}}$ as  $a_t \equiv A_t/M_{\text{susy}}, \bar{\mu} \equiv \mu/M_{\text{susy}}, \text{ and } x_t \equiv X_t/M_{\text{susy}} = a_t - \bar{\mu}/t_{\beta}$ . Our sign convention for  $\mu$  and  $A_t$  is the same as [\[15\]](#page-9-9), which are opposite sign convention of [\[16\]](#page-9-10). We fix  $\hat{m}_{\tilde{g}} = 2$ , well above the current LHC reach,  $\hat{m}_{\tilde{b}_1} = \hat{m}_{\tilde{b}_2} = 1$ , and  $\hat{m}_{\tilde{t}_1} = 0.8$ ,  $\hat{m}_{\tilde{t}_2} = 1.2$ . A stop mass difference  $m_{\tilde{t}_2} - m_{\tilde{t}_1} \geq 0.4$  TeV is chosen in accord with Natural SUSY prediction[\[24](#page-10-0)]. Then  $\Delta_b$  is well approximated numerically by

<span id="page-2-2"></span>
$$
\Delta_b \simeq \bar{\mu} \frac{t_\beta}{20} \left[ 0.26 + \left( \frac{0.09}{|\bar{\mu}| + 0.6} - 0.003 \right) a_t \right] , \qquad (5)
$$

where the first and the second terms in the square bracket are the numerical values from the gluino and the chargino contributions respectively.

The chargino and neutralino masses have no special role except possibly in  $b \to s\gamma$  decay, but consistency with Natural SUSY has been found there<sup>[\[4](#page-8-0)]</sup>. Large  $m_A$  implies large charged Higgs  $H^+$  mass that suppresses the  $H^+$  loop contribution to  $b \to s\gamma$ .

The gg,  $\gamma\gamma$  coupling ratios  $r^{\phi}_{gg,\gamma\gamma}$  for  $\phi = h, H, A$  relative to those of  $h_{SM}$  are [\[17\]](#page-9-11)

<span id="page-2-1"></span>
$$
r_{gg}^{\phi} = \frac{I_{tt}^{\phi} r_{tt}^h + I_{bb}^{\phi} r_{bb}^h}{I_{tt}^{\phi} + I_{bb}^{\phi}} , \qquad r_{\gamma\gamma}^{\phi} = \frac{\frac{7}{4} I_{WW}^{\phi} r_{VV}^h - \frac{4}{9} I_{tt}^{\phi} r_{tt}^h - \frac{1}{9} I_{bb}^{\phi} r_{bb}^h}{\frac{7}{4} I_{WW}^{\phi} - \frac{4}{9} I_{tt}^{\phi} - \frac{1}{9} I_{bb}^{\phi}} , \qquad (6)
$$

where  $I_{WW,tt,bb}^{\phi}$  represent the triangle-loop contributions to the amplitudes normalized to the  $m_h \rightarrow 0$  limit[\[18](#page-9-12)[–20\]](#page-9-13).

The  $XX \to h \to PP$  cross section ratios[\[17\]](#page-9-11) relative to  $h_{SM}$  are obtained from

<span id="page-2-0"></span>
$$
\sigma_P \equiv \frac{\sigma_{PP}}{\sigma_{\rm SM}} = \frac{\sigma_{XX \to PP}}{\sigma_{XX \to h_{SM} \to PP}} = \frac{|r_{XX}^h r_{PP}^h|^2}{R^h} \,,\tag{7}
$$

$$
R^{h} = \frac{\Gamma_{\text{tot}}^{h}}{\Gamma_{\text{tot}}^{h_{SM}}} = 0.57|r_{bb}^{h}|^{2} + 0.06|r_{\tau\tau}^{h}|^{2} + 0.25|r_{VV}^{h}|^{2} + 0.09|r_{gg}^{h}|^{2} + 0.03|r_{cc}^{h}|^{2},\tag{8}
$$

where  $R^h$  is the ratio of the h total width to that of  $h_{SM}$ ,  $\Gamma_{h_{SM}}^{\text{tot}} = 4.14 \text{ MeV}[21]$  $\Gamma_{h_{SM}}^{\text{tot}} = 4.14 \text{ MeV}[21]$  for  $m_h =$ 125.5 GeV. The coefficients in Eq. [\(8\)](#page-2-0) are the SM Higgs branching fractions. Here we have assumed no appreciable decays to dark matter.

Sum rule of cross-section ratios In the large  $m_A$  region close to the decoupling limit,  $\alpha$ takes a value

$$
\alpha = \beta - \frac{\pi}{2} + \epsilon \tag{9}
$$

with  $|\epsilon| < \frac{\pi}{2} - \beta$ . Then, the  $r_{XX}^h$  of Eq. [\(2\)](#page-1-0) are well approximated by

<span id="page-3-1"></span>
$$
r_{VV}^h = 1, \quad r_{tt,cc}^h = 1 + \epsilon/t_\beta, \quad r_{\tau\tau}^h \simeq 1 - \epsilon t_\beta, \quad r_{bb}^h \simeq 1 - \frac{1}{1 + \Delta_b} \epsilon t_\beta \; . \tag{10}
$$

through first order in  $\epsilon$ . The  $r_{tt,cc}^h$  are close to unity because those deviations from SM are  $t_\beta$  suppressed. Thus,

$$
r_{gg}^h \simeq r_{\gamma\gamma}^h \simeq 1 \tag{11}
$$

since the bottom triangle loop function  $I_{bb}^h$  is negligible in Eq. [\(6\)](#page-2-1). Only  $r_{bb}^h$ ,  $r_{\tau\tau}^h$  can deviate sizably from unity for large  $m_A$  and large tan $\beta$ . Following Eqs. [\(7\)](#page-2-0) and [\(8\)](#page-2-0), the  $\sigma_P \equiv$  $\sigma_{PP}/\sigma_{SM}$  of the other channels are commonly reduced(enhanced) in correspondence with  $r_{bb}^h > 1$  ( $r_{bb}^h < 1$ ). We predict the cross sections relative to their individual SM expectations

<span id="page-3-0"></span>
$$
\sigma_{\gamma} = \sigma_W = \sigma_Z = \frac{1}{0.6(r_{bb}^h)^2 + 0.4} \,, \tag{12}
$$

and

$$
0.4\sigma_{\gamma} + 0.6\sigma_{b} = 1\tag{13}
$$

where the SM  $b\bar{b}$  branching fraction [\[22](#page-9-15)] is approximated as 60%. Equation [\(12\)](#page-3-0) holds independently of the production process. Enhanced  $\sigma_{\gamma}$  implies reduced  $\sigma_{b}$ , as well as enhanced  $\sigma_W$  and  $\sigma_Z$ .

Flavor-Tuning of mixing angle  $\alpha$  Note that  $r_{bb,\tau\tau}^h = 1$  in the exact decoupling limit  $m_A \to$  $\infty$  for which  $\epsilon = 0$ . Flavor-tuning of  $\epsilon$  to be small but non-zero is necessary to obtain a significant variation of  $r_{bb}^h$  from unity. Positive(negative)  $\epsilon$  gives bb-reduction(enhancement).

The mixing angle  $\alpha$  is obtained by diagonalizing the squared-mass matrix of the neutral Higgs in the  $u, d$  basis. Their elements at tree-level are

$$
(M_{ij}^2)^{\text{tree}} = M_Z^2 s_\beta^2 + m_A^2 c_\beta^2; \quad M_Z^2 c_\beta^2 + m_A^2 s_\beta^2; \quad -(M_Z^2 + m_A^2) s_\beta c_\beta \tag{14}
$$

for  $ij = 11; 22; 12$ , respectively, which gives  $\epsilon < 0$  in all region of  $m_A$ . Thus, in order to get  $b\bar{b}$ -reduction, it is necessary to cancel  $(M_{12}^2)$ <sup>tree</sup> by higher order terms  $\Delta M_{ij}^2$ .

In the 2-loop leading-log(LL) approximation the  $\Delta M_{ij}^2$  are given [\[15,](#page-9-9) [23](#page-9-16)] by

<span id="page-4-2"></span>
$$
M_{ij}^2 = (M_{ij}^2)^{\text{tree}} + \Delta M_{ij}^2 \tag{15}
$$

where

<span id="page-4-1"></span>
$$
\Delta M_{11}^2 = F_3 \frac{3\bar{m}_t^4}{4\pi^2 v^2 s_\beta^2} \left[ t(1 - G_{\frac{15}{2}} t) + a_t x_t (1 - \frac{a_t x_t}{12}) (1 - 2G_{\frac{9}{2}} t) \right] - M_Z^2 s_\beta^2 (1 - F_3)
$$
  
\n
$$
\Delta M_{22}^2 = -F_{\frac{3}{2}} \frac{\bar{m}_t^4}{16\pi^2 v^2 s_\beta^2} \left[ (1 - 2G_{\frac{9}{2}} t)(x_t \bar{\mu})^2 \right]
$$
\n
$$
\Delta M_{12}^2 = -F_{\frac{9}{4}} \frac{3\bar{m}_t^4}{8\pi^2 v^2 s_\beta^2} \left[ (1 - 2G_{\frac{9}{2}} t)(x_t \bar{\mu}) (1 - \frac{a_t x_t}{6}) \right] + M_Z^2 s_\beta c_\beta (1 - F_{\frac{3}{2}})
$$
\n(16)

where  $F_l = 1/(1 + \frac{h_t^2}{8\pi^2}t)$  with  $l = 3, \frac{3}{2}$  $\frac{3}{2}, \frac{9}{4}$  $\frac{9}{4}$  and  $G_l = -\frac{1}{16\pi^2} (l h_t^2 - 32\pi \alpha_s)$  with  $l = \frac{15}{2}$  $\frac{15}{2}, \frac{9}{2}$  $\frac{9}{2}$ . The  $F_l$  are due to the wave function (WF) renormalization of the  $H_u$  field and the index l is numbers of  $H_u^0$  fields in the effective potential of the two Higgs doublet model.  $F_3\xi^4 \simeq$  $F_{\frac{9}{4}}\xi^3 \simeq F_{\frac{3}{2}}\xi^2 \simeq 1$  where  $\xi$  is defined by  $H_u(M_s) = H_u(\bar{m}_t)\xi$  where  $\xi = F_{\frac{3}{4}}^{-1}$ .

The parameter tan $\beta = v_u/v_d$  is defined in terms of the Higgs vacuum expectation values  $v_{u,d} = \langle H_{u,d}^0 \rangle$  at the minimum of the 1-loop effective potential at the weak scale  $\mu = \bar{m}_t$  and  $v = \sqrt{v_u^2 + v_d^2} \simeq 174$  GeV, while  $a_t, x_t, \bar{\mu}$  have scale  $\mu = M_{\text{susy}}$ . The relation  $\cot\beta(\bar{m}_t) = \cot\beta(M_s) \xi^{-1}$  will be used in the following calculation.

Numerically  $\alpha_s = \alpha_s(\bar{m}_t) = 0.109$  giving  $-32\pi\alpha_s = -10.9$ , while  $h_t = \bar{m}_t/v = 0.939$  is small.  $G_{\frac{15}{2},\frac{9}{2}} = 0.0274, 0.0442$  and  $t = \log(\frac{1 \text{ TeV}}{\bar{m}_t})^2 = 3.62$ ; thus,  $G_{\frac{15}{2}}t = 0.099$  and  $2G_{\frac{9}{2}}t =$ 0.320, and  $F_3 = 0.892$ .

In large  $m_A$  limit, the  $m_h^2$  expression is

<span id="page-4-0"></span>
$$
m_h^2 = M_Z^2 c_{2\beta}^2 + F_3 \frac{3\bar{m}_t^4}{4\pi^2 v^2} \left[ t(1 - G_{\frac{15}{2}}t) + (1 - 2G_{\frac{9}{2}}t)(x_t^2 - \frac{x_t^4}{12}) \right] - M_Z^2 [s_\beta^4 (1 - F_3) - 2s_\beta^2 c_\beta^2 (1 - F_{\frac{3}{2}})] \tag{17}
$$

where the Higgs WF renormalization factor  $\xi$  is retained in the denominator of  $F_3$ . This  $F_3$ factor is usually expanded to the numerator in 2LL approximation, and correspondingly  $G_{\frac{15}{2}}$ and  $G_{\frac{9}{2}}$  are replaced by  $G_{\frac{3}{2}}$ :  $m_h^2 = M_Z^2 c_{2\beta}^2 + \frac{3\bar{m}_t^4}{4\pi^2 v^2} [t(1-G_3t) + (1-2G_3t)(x_t^2 - \frac{x_t^4}{12})] - M_Z^2 s_{\beta}^4$  $\frac{3h_t^2}{8\pi^2}t.$ However, numerically Eq. [\(17\)](#page-4-0) significantly increases  $m_h$  at large  $M_{\text{susy}}$  as shown in Fig. [1:](#page-5-0) Eq. [\(17\)](#page-4-0) gives increasing  $m_h$  as  $M_{\text{susy}}$  increases up to ~ 7 TeV, while the usual formula with



<span id="page-5-0"></span>FIG. 1.  $M_{\text{susy}}$  dependence of Higgs mass  $m_h$  by the improved formula Eq. [\(17\)](#page-4-1) (solid black) in comparison with the one by the usual 2LL approximation (dashed blue) where the first  $F_3$  is removed and  $G_{\frac{15}{2}}$  and  $G_{\frac{9}{2}}$  are replaced by  $G_{\frac{3}{2}}$ . In this illustration  $x_t$  is taken to be  $\sqrt{6}$  following the "maximal-mixing" condition, and  $tan\beta = 20$ .

the expansion approximated for  $F_3$  gives decreasing  $m_h$  when  $M_{\text{susy}} > 1.3 \text{ TeV}$  and is not applicable at large  $M_{\text{susy}}$ .

The experimental  $m_h$  determinations from the LHC experiments are [\[1,](#page-7-0) [2\]](#page-7-1)

$$
m_h = 125.3 \pm 0.4 \pm 0.4, \quad 126.0 \pm 0.4 \pm 0.4 \text{ GeV}
$$
 (18)

It seems unlikely that the central  $m_h$  determination will change much with larger statistics because of the excellent mass resolution in the  $\gamma\gamma$  channel. The experimental  $m_h$  value is near the maximum possible value of  $m_h$  in Eq. [\(17\)](#page-4-0) and this constrains the value of  $x_t$  to  $|x_t| \simeq \sqrt{6}$ , to maximize the term  $x_t^2 - \frac{x_t^4}{12}$ . This is known as "maximal-mixing" in the stop mass-matrix[\[24\]](#page-10-0). In Eq. [\(17\)](#page-4-0) we require  $m_h \ge 124$  GeV. This implies

<span id="page-5-1"></span>
$$
1.95 (\equiv x_{\text{tmin}}) < |x_t| < 2.86 (\equiv x_{\text{tmax}}), \tag{19}
$$

where we should note that the positive  $x_t$  branch is favored by the SUSY renormalization group prediction[\[24](#page-10-0)].

By using Eq. [\(15\)](#page-4-2) the Higgs mixing angle  $\alpha$  is determined from

$$
t_{2\alpha} = \frac{2M_{12}^2}{M_{22}^2 - M_{11}^2} \simeq \frac{(m_A^2 + M_Z^2)s_{2\beta} - 2\Delta M_{12}^2}{(m_A^2 - M_Z^2)c_{2\beta} + (\Delta M_{11}^2 - \Delta M_{22}^2)},
$$
\n(20)

$$
\Delta M_{12}^2 \simeq -\frac{\bar{\mu}}{s_\beta^2} x_t (1 - \frac{x_t^2}{6}) 558 \text{GeV}^2 + 24 \cdot \frac{20}{\tan \beta} \text{GeV}^2 \,. \tag{21}
$$

Defining  $z (\equiv M_Z^2/m_A^2)$ ,  $\delta (\equiv \Delta M_{12}^2/m_A^2)$ , and  $\eta (\equiv \frac{1}{2})$  $\frac{1}{2}(\Delta M_{11}^2 - \Delta M_{22}^2)/m_A^2$ ,  $\epsilon$  is simply given in the first order of z,  $\delta$ , and  $\eta$  by

$$
\epsilon = -2\frac{z+\eta}{\tan\beta} + \delta \tag{22}
$$

We note that  $r_{bb}^h$  is related to  $\epsilon$  through Eq. [\(10\)](#page-3-1). With the  $x_t$  constraint in Eq. [\(19\)](#page-5-1), we can derive the allowed region of  $r_{bb}^h$  for each  $\bar{\mu}$ -value. Correspondingly, the allowed regions of  $\sigma_{\gamma} (=\sigma_{\gamma\gamma}/\sigma_{\rm SM} = \frac{1}{0.6(r^{h})^2}$  $\frac{1}{0.6(r_{bb}^{h})^2+0.4}),\,\sigma_b(=\sigma_{b\bar{b}}/\sigma_{\rm SM}\ =\frac{(r_{bb}^{h})^2}{0.6(r_{bb}^{h})^2}.$  $\frac{(r_{bb}^h)^2}{0.6(r_{bb}^h)^2+0.4})$  and  $\sigma_\tau (= \sigma_{\tau\tau}/\sigma_{\rm SM} = \frac{(r_{\tau\tau}^h)^2}{0.6(r_{bb}^h)^2+0.4}$  $\frac{(r_{\tau\tau}^{\alpha})^2}{0.6(r_{bb}^h)^2+0.4}$ are given respectively by the two curves in Fig. [2](#page-11-0) where we take  $tan \beta = 50$ .

The condition  $r_{bb}^h = 1$ , or equivalently  $\epsilon = 0$ ,  $t_{2\alpha} = t_{2\beta}$ , defines the boundary that separates the  $\gamma\gamma$  enhancement and suppression in the parameter space.

$$
r_{bb}^h = 1 \Leftrightarrow \epsilon = 0 \Leftrightarrow \Delta M_{12}^2 = M_Z^2 s_{2\beta} - \frac{\Delta M_{11}^2 - \Delta M_{22}^2}{2} t_{2\beta},\tag{23}
$$

This condition is independent of  $m_A$  and the quantum correction  $\Delta_b$ .

 $\Delta M_{12}^2 > M_Z^2 s_{2\beta} - \frac{\Delta M_{11}^2 - \Delta M_{22}^2}{2} t_{2\beta}$  gives  $b\bar{b}$  reduction. Flavor-tuning (FT) with small  $\alpha$  requires a cancellation of  $(M_{12}^2)$ <sup>tree</sup> by the loop-level  $\Delta M_{12}^2$  contribution, which requires rather large values of  $\bar{\mu}$  and tan $\beta$ . This possibility was raised in ref.[\[25\]](#page-10-1).

The region of  $\gamma\gamma$  enhancement does not overlap with the region  $|\bar{\mu}| < 0.5$  of Natural SUSY for any value of  $tan\beta$  from 20 to 60. For  $tan\beta = 20$ ,  $|\bar{\mu}| \stackrel{>}{\sim} 2$  is necessary for  $\gamma\gamma$ enhancement.

We give a benchmark point of the FT model in the MSSM (FT1) and two benchmark points of Natural SUSY (NFT1,NFT2).

$$
\bar{\mu} \quad \tan\beta \quad x_t \quad m_h(\text{GeV}) \quad \sigma_\gamma \quad \sigma_b \quad \sigma_\tau
$$
\n
$$
FT1 \quad -3 \quad 20 \quad -2.86 \quad 124 \quad 1.17 \quad 0.89 \quad 1.05
$$
\n
$$
NFT1 \quad -0.5 \quad 20 \quad 2.70 \quad 125 \quad 0.84 \quad 1.11 \quad 1.04
$$
\n
$$
NFT2 \quad -0.15 \quad 20 \quad 2.70 \quad 125 \quad 0.87 \quad 1.08 \quad 1.07
$$
\n(24)

where  $M_{\text{susy}} = 1$  TeV and  $m_A = 0.5$  TeV. The relevant sparticle masses are taken commonly with the values given above Eq.  $(5)$ . The  $m_h$  value is predicted by Eq.  $(17)$ . We also note that the predicted  $BF(B_s \to \mu^+\mu^-)$  values of these bench mark points are consistent with the experimental measurement[\[26](#page-10-2)]  $BF(B_s \to \mu^+ \mu^-) = \left(3.2 \frac{+1.4}{-1.2 \text{stat}} \frac{+0.5}{-0.3 \text{syst}}\right) \times 10^{-9}$  within  $2\sigma$ . Natural SUSY predictions Natural SUSY always predicts bb-enhancement and  $\gamma\gamma$  reduction.[\[29\]](#page-10-3)



Here we have taken  $|\mu| \leq 500 \text{GeV}$  and the other parameters are fixed with the values given above Eq.  $(5)$ .

## Concluding remarks

We have explored the  $\gamma\gamma$ ,  $b\bar{b}$  and  $\tau\tau$  signals in the MSSM, relative to SM, and also in Natural SUSY. In MSSM an enhancement in the diphoton signal of the 125 GeV Higgs boson relative to the SM Higgs can be obtained in a flavor-tuned model with  $h = H_u^0$  provided that | $|\mu|$  is large( TeV) and negative. A  $\gamma\gamma$  enhancement is principally due to the reduction of the  $b\bar{b}$  decay width compared to  $h_{SM}$ . The ratios of  $WW^*$  and  $ZZ^*$  to their SM values are predicted to be the same as that of  $\gamma\gamma$ . There is also a corresponding reduction of the h to  $\tau\tau$  signal. The Tevatron evidence of a Higgs to  $b\bar{b}$  signal in W + Higgs production [\[28\]](#page-10-4) does not favor much  $b\bar{b}$  reduction. The flavor-tuning of the neutral Higgs mixing angle  $\alpha$  requires a large  $\mu \sim TeV$  and large tan $\beta$ . For small  $|\mu| \stackrel{<}{\sim} 0.5$  TeV of Natural SUSY,  $\gamma\gamma$ -suppression relative to the SM is predicted. Thus, precision LHC measurements of the  $\gamma\gamma$ , W<sup>\*</sup>W, Z<sup>\*</sup>Z and  $b\bar{b}$  signals of the 125 GeV Higgs boson can test MSSM models.

## Acknowledgements

W.-Y. K. and M. I. thank the hospitality of National Center for Theoretical Sciences and Academia Sinica while this work was completed. M.I. is also grateful for hospitality at UW-Madison. This work was supported in part by the U.S. Department of Energy under grants No. DE-FG02-95ER40896 and DE-FG02-12ER41811.

- <span id="page-7-0"></span>[1] ATLAS Collaboration, Phys. Lett. B 716, 1 (2012) [\[arXiv:1207.7214\[](http://arxiv.org/abs/1207.7214)hep-ex]].
- <span id="page-7-1"></span>[2] CMS Collaboration, Phys. Lett. B 716, 30 (2012) [\[arXiv:1207.7235\[](http://arxiv.org/abs/1207.7235)hep-ex]].
- <span id="page-7-2"></span>[3] M. Dine, A. Kagan and S. Samuel, Phys. Lett. B 243, 250 (1990). S. Kelley, J. Lopez, D. Nanopoulos, H. Pois, and K. Yuan, Nucl. Phys. B398, 3 (1993).

V. Barger, M. Berger, P. Ohmann, Phys. Rev. D49, 4908 (1994).

S. Dimopoulos and G. Giudice, Phys. Lett. B357, 573 (1995) [\[arXiv:hep-ph/9507282](http://arxiv.org/abs/hep-ph/9507282) [hepph]].

F. Gabbiani, E. Gabrielli, A. Masiero, and L. Silvestrini, Nucl. Phys. B477, 321 (1996) [\[arXiv:hep-ph/9604387](http://arxiv.org/abs/hep-ph/9604387) [hep-ph]].

A. Cohen, D. Kaplan, and A. Nelson, Phys. Lett. B388, 588 (1996).

K. Chan, U. Chattopadhyay and P. Nath, Phys. Rev. D58, 096004 (1998).

J. L. Feng, K. T. Matchev, and T. Moroi, Phys. Rev. D61, 075005 (2000) [\[arXiv:hep-ph/9909334](http://arxiv.org/abs/hep-ph/9909334) [hep-ph]]; Phys. Rev. Lett. 84, 2322 (2000) [\[arXiv:hep-ph/9908309](http://arxiv.org/abs/hep-ph/9908309) [hep-ph]].

R. Kitano and Y. Nomura, Phys. Lett. B631, 58 (2005) [\[arXiv:hep-ph/0509039](http://arxiv.org/abs/hep-ph/0509039) [hep-ph]].

C Brust, A. Katz, S. Lawrence and R. Sundrum, JHEP 1203, 103 (2012) [\[arXiv:1110.6670](http://arxiv.org/abs/1110.6670) [hep-ph]].

S. Akula, M. Liu, P. Nath, and G. Peim, Phys. Lett. B709, 192 (2012) [\[arXiv:1111.4589](http://arxiv.org/abs/1111.4589) [hepph]].

R. Essig, E. Izaguirre, J. Kaplan and J. G. Wacker, JHEP 1201, 074 (2012) [\[arXiv:1110.6443](http://arxiv.org/abs/1110.6443) [hep-ph]].

M. Papucci, J. T. Ruderman and A. Weiler, JHEP 1209, 035 (2012) [\[arXiv:1110.6926](http://arxiv.org/abs/1110.6926) [hepph]].

L. Hall, D. Pinner, and J. Rudermn, JHEP 1204, 131(2012)

S. King, M. Muhlleitner and R. Nevzorov, Nucl. Phys. B860, 207 (2012) [\[arXiv:1201.2671](http://arxiv.org/abs/1201.2671) [hep-ph]].

- N. Arkani-Hamed, talk at WG2 meeting, Oct. 31, 2012, CERN, Geneva.
- H. Baer, V. Barger, P. Huang, JHEP 1111, 031 (2011) [\[arXiv:1107.5581](http://arxiv.org/abs/1107.5581) [hep-ph]].

H. Baer, V. Barger, P. Huang, X. Tata,in JHEP 1205, 109 (2012) [\[arXiv:1203.5539\[](http://arxiv.org/abs/1203.5539)hep-ph]].

- H. Baer, V. Barger, P. Huang, A. Mustafayev, X. Tata, [arXiv:1207.3343](http://arxiv.org/abs/1207.3343) [hep-ph].
- D. M. Ghilencea, H. M. Lee, and M. Park, JHEP 1207, 046 (2012) [\[arXiv:1203.0569](http://arxiv.org/abs/1203.0569) [hep-ph]].
- J. Feng and D. Sanford, Phys. Rev. D 86, 055015 (2012) [\[arXiv:1205.2372](http://arxiv.org/abs/1205.2372) [hep-ph]].

L. Randall, M. Reece, [arXiv:1206.6540](http://arxiv.org/abs/1206.6540) [hep-ph].

<span id="page-8-0"></span>[4] H. Baer, V. Barger, and A. Mustafayev, Phys. Rev. D85, 075010 (2012) [\[arXiv:1112.3017v](http://arxiv.org/abs/1112.3017)3[hep-ph]].

- <span id="page-9-1"></span><span id="page-9-0"></span>[5] ATLAS collaboration, ATLAS-CONF-2012-094.
- [6] M. Carena, S. Gori, N. R. Shah, C. E. M. Wagner, L.-T. Wang, JHEP 1207, 175 (2012) [\[arXiv:1205.5842](http://arxiv.org/abs/1205.5842) [hep-ph]].
- <span id="page-9-3"></span><span id="page-9-2"></span>[7] K. Hagiwara, J. S. Lee, and J. Nakamura, JHEP 1210, 002 (2012) [\[arXiv:1207.0802](http://arxiv.org/abs/1207.0802) [hep-ph]].
- [8] D. Carmi, A. Falkowski, E. Kuflik, T. Volansky, JHEP 1207, 136 (2012) [\[arXiv:1202.3144](http://arxiv.org/abs/1202.3144) [hep-ph]].
- <span id="page-9-4"></span>[9] V. Barger, H. E. Logan, G. Shaughnessy, Phys. Rev. D 79, 115018 (2009) [\[arXiv:0902.0170\[](http://arxiv.org/abs/0902.0170)hep-ph]].
- <span id="page-9-5"></span>[10] D. Chung et al., Phys. Rept. 407, 1-203 (2005). [arXiv:hep-ph/0312378](http://arxiv.org/abs/hep-ph/0312378)
- [11] John F. Gunion, Howard E. Haber, Gordon Kane, Sally Dawson, "The Higgs Hunter's Guide", (Perseus Books, Boulder Colorado 1990).
- <span id="page-9-6"></span>[12] L. Hall, R. Rattazzi, and U. Sarid, Phys. Rev. D 50, 7048 (1994). M. Carena, M. Olechowski, S. Pokorski, and C. E. M. Wagner, Nucl. Phys. B426, 269. D. Pierce, J. Bagger, K. Matchev, and R. Zhang, Nucl. Phys. B491, 3 (1997).
- <span id="page-9-8"></span><span id="page-9-7"></span>[13] A. Djouadi, Phys. Rep. 459, 1 (2008) [\[hep-ph/0503173v](http://arxiv.org/abs/hep-ph/0503173)2].
- [14] P. Langacker in "the Review of Particle Physics", J. Beringer et al. (Particle Data Group), Phys. Rev. D86, 010001 (2012)
- <span id="page-9-9"></span>[15] M. Carena et al., JHEP 1207, 175 (2012); JHEP 1203, 014 (2012).
- <span id="page-9-10"></span>[16] H. Baer and X. Tata, Weak Scale Supersymmetry: From Superfields to Scattering Events, (Cambridge University Press, Cambridge, England, 2006).
- <span id="page-9-11"></span>[17] V. Barger, M. Ishida, and W.-Y. Keung, Phys. Rev. Lett. 108, 261801 (2012) [\[arXiv:1203.3456\[](http://arxiv.org/abs/1203.3456)hep-ph]].
- <span id="page-9-12"></span>[18] "Collider Physics", by V. D. Barger and R. J. N. Phillips, (Perseus Book, Boulder, Cololado 1997).
- [19] V. Barger, M. Ishida, and W.-Y. Keung, Phys. Rev. D85, 015024 (2012).
- <span id="page-9-13"></span>[20] J. Chang, K. Cheung, Po-Yan Tseng, and Tzu-Chiang Yuan, JHEP 1212, 058 (2012) [\[arXiv:1206.5853\]](http://arxiv.org/abs/1206.5853).
- <span id="page-9-14"></span>[21] A. Denner, S. Heinemeyer, I. Puljak, D. Rebuzzi, and M. Spira, Eur. Phys. J. C71, 1753 (2011) [\[arXiv:1107.5909\[](http://arxiv.org/abs/1107.5909)hep-ph]].
- <span id="page-9-15"></span>[22] A. Djouadi, J. Kalinowski and M. Spira, Comput. Phys. Commun. bf108:56-74, (1998)
- <span id="page-9-16"></span>[23] H. E. Haber and R. Hempfling, Phys. Rev. D48, 4280 (1993) [arXiv: [hep-ph/9307201\]](http://arxiv.org/abs/hep-ph/9307201).
- <span id="page-10-0"></span>[24] V. Barger, P. Huang, M. Ishida, and W.-Y. Keung, Phys. Lett. B718, 1024 (2013) [\[arXiv:1206.1777\[](http://arxiv.org/abs/1206.1777)hep-ph]].
- <span id="page-10-1"></span>[25] M. Carena, S. Mrenna, and C. E. M. Wagner, Phys. Rev. D 62, 055008 (2000) [\[arXiv:hep-ph/9907422\]](http://arxiv.org/abs/hep-ph/9907422).
- <span id="page-10-5"></span><span id="page-10-2"></span>[26] R. Aaij et al. [LHCb Collaboration], [arXiv:1211.2674.](http://arxiv.org/abs/1211.2674)
- <span id="page-10-4"></span>[27] K. Blum, R. T. D'Agnolo, Phys. Lett. B714, 66 (2012) [\[arXiv:1202.2364\[](http://arxiv.org/abs/1202.2364)hep-ph]].
- [28] The TEVNPH Working Group for the CDF and D0 Collaboration, FERMILAB-CONF-12- 065-E . arXiv: 1207.0449[hep-ex].
- <span id="page-10-3"></span>[29] Here we note that as  $\bar{\mu} = 0$ , then  $r_{bb}^h = 1 + 2M_Z^2/m_A^2 \rightarrow \sigma_{\gamma\gamma}/\sigma_{\rm SM} = 1 - 2.4M_Z^2/m_A^2$ , independently of tan $\beta$ . The smaller  $m_A$  gives the larger suppression of  $\sigma_{\gamma}[27]$  $\sigma_{\gamma}[27]$  $\sigma_{\gamma}[27]$ . Then, from the  $\gamma\gamma$ deviation from unity, the  $CP$ -odd state mass  $m_A$  could be estimated.



<span id="page-11-0"></span>FIG. 2.  $\bar{\mu}$  dependence of  $\sigma_{\gamma} = \sigma_{\gamma\gamma}/\sigma_{\rm SM}(\text{upper panel})$ ,  $\sigma_b = \sigma_{b\bar{b}}/\sigma_{\rm SM}(\text{middle panel})$ , and  $\sigma_b = \sigma_{b\bar{b}}/\sigma_{\rm SM}$ (lower panel) for  $m_A = 500$  GeV: Their allowed values are between the solid red curve (corresponding to  $|x_t| = x_{\text{tmax}}$ ) and the dashed blue curve (corresponding to  $|x_t| = x_{\text{tmin}}$ ). Left(Right) panels show negative(positive)  $x_t$  region. Deviations from unity are enlarged for a large negative  $\bar{\mu}$ , but there the perturbative calculation is unreliable due to a large quantum correction.