ExoMol line lists II: The ro-vibrational spectrum of SiO

Emma J. Barton, Sergei N. Yurchenko and Jonathan Tennyson

Department of Physics and Astronomy, University College London, London WC1E 6BT, UK

Accepted XXXX. Received XXXX; in original form XXXX

ABSTRACT

Accurate rotation-vibration line lists are calculated for silicon monoxide. Line lists are presented for the main isotopologue, ${}^{28}\text{Si}{}^{16}\text{O}$, and for four monosubsituted isotopologues (${}^{29}\text{Si}{}^{16}\text{O}$, ${}^{30}\text{Si}{}^{16}\text{O}$, ${}^{28}\text{Si}{}^{18}\text{O}$ and ${}^{28}\text{Si}{}^{17}\text{O}$), in their ground electronic states. These line lists are suitable for high temperatures (up to 9000 K) including those relevant to exoplanetary atmospheres and cool stars. A combination of empirical and *ab initio* methods is used: the potential energy curves are determined to high accuracy by fitting to extensive data from the analysis of both laboratory and sunspot spectra; a high quality *ab initio* dipole moment curve is calculated at the large basis set, multi-reference configuration interaction level. A partition function plus full line lists of rotation-vibration transitions are made available in an electronic form as supplementary data to this article and at www.exomol.com.

Key words: molecular data; opacity; astronomical data bases: miscellaneous; planets and satellites: atmospheres; stars: low-mass

1 INTRODUCTION

Silicon monoxide (SiO) is a widely observed astronomical species and appears to be ubiquitous in our galaxy. Since its initial detection by Snyder & Buhl (1974), SiO has proved to be a key astrophysical maser and its very bright maser emissions are the subject of continued study (Cotton et al. 2011; Assaf et al. 2011; Nakashima et al. 2011; Vlemmings et al. 2011; Deguchi et al. 2011).

SiO was originally detected in the interstellar medium by Wilson et al. (1971). Its emissions were subsequently observed in the envelopes of oxygen-rich giant and super-giant stars (Kaifu et al. 1975) and the remnants of Supernova 1987A (Aitken et al. 1988). SiO was detected in absorption in cool giant stars (Rinsland & Wing 1982) including α Tau (Cohen et al. 1992). SiO absorptions are prominent in sunspots which have proved to be a fruitful source of spectroscopic data on highly excited states (Glenar et al. 1985; Campbell et al. 1995) which we exploit below. Furthermore recent detections of hot, dense exoplanets have led to the speculation that their atmospheres might contain significant quantities of SiO from vaporised silicates (Schaefer et al. 2012).

These astronomical applications, combined with technological uses of SiO spectra (Wooldridge et al. 2000; Motret et al. 2002), have motivated a number of laboratory studies. Tipping & Chackerian (1981),Mollaaghababa et al. (1991), Campbell et al. (1995), Cho & Saito (1998) and Sanz et al. (2003) have all produced molecular constants characterising SiO vibration-rotation states. Transition probabibilities or Einstein-A coefficients have also been provided by Tipping & Chackerian (1981), Langhoff & Bauschlicher (1993) and Drira et al. (1998). Transition line lists have been constructed using these data (Lovas et al. 1981; Glenar et al. 1985; Langhoff & Bauschlicher 1993; Drira et al. 1998), but none of them appears to be particularly complete. For example no single line list combines a comprehensive set of transition frequencies with an accurate model for the transition intensities. It is this that we aim to do here.

Tipping & Chackerian (1981) reported transition probabilities for a large number of transitions based on a semi-empirical dipole moment which they claim should give results accurate to about 10 %. Langhoff & Bauschlicher (1993) computed an accurate electric dipole moment function for the SiO molecule which not only reproduces accurate dipole moments, but also reproduces line strengths for vibrational-rotational transitions within the ground-state manifold. The resulting line lists are for ²⁸Si¹⁶O, ²⁹Si¹⁶O and ³⁰Si¹⁶O for $J \leq 250$ but v is limited to 15. Intensities were calculated using band intensities and Hönl-London factors. Drira et al. (1998) calculated an *ab initio* ²⁸Si¹⁶O line list for $J \leq 100$ and $v \leq 40$ although the

Reference	Method	Transitions	Temperature (K)	Frequency range (cm^{-1})	Uncertainty (cm^{-1})
Sanz et al. (2003)	Laboratory	$\Delta v = 0, J' \to J'' = 1 \to 0$ ²⁸ Si ¹⁶ O v = 0 - 45 ²⁹ Si ¹⁶ O v = 0 - 26 ²⁸ Si ¹⁸ O v = 0 - 44	1000 - 9000	0.13 - 1.67	6.7×10^{-8}
Campbell et al. (1995)	Sunspots	$\begin{array}{l} \Delta v = 1, \ \Delta J = \pm 1 \\ ^{28}\mathrm{Si}^{16}\mathrm{O} \ v = 0 - 13 \ J \leqslant 141 \\ ^{29}\mathrm{Si}^{16}\mathrm{O} \ v = 0 - 6 \ J \leqslant 107 \\ ^{30}\mathrm{Si}^{16}\mathrm{O} \ v = 0 - 6 \ J \leqslant 92 \end{array}$	3200	900 - 1300	0.006

Table 1. Summary of observations data used to determine the SiO potential energy curve. Temperatures are only approximate and the uncertainties are the estimates given in the cited papers.

publically available line list from the Strasbourg Data Centre contains only 503 transitions. Drira et al. (1998)'s intensities are significantly larger than those of Tipping & Chackerian (1981) and they estimate their intensities are only accurate to about 20 %.

The ExoMol project Tennyson & Yurchenko (2012) aims to provide line lists of spectroscopic transitions for key molecular species which are likely to be important in the atmospheres of extrasolar planets and cool stars; it aims, scope and methodology have been summarised by Tennyson & Yurchenko (2012). Line lists for ${}^{2}\Sigma^{+}$ XH molecules, X = Be, Mg, Ca, have already been published (Yadin et al. 2012). In the present paper, we present rotation-vibration transition lists and associated spectra for the five major isotopologues of SiO. These line lists are particulary comprehensive and should be valid for temperatures up to 9000 K.

2 METHOD

Rotation-vibration line lists for the ground electronic state of SiO were obtained by direct solution of the nuclear motion Schrödinger equation using the program LEVEL8.0 (Le Roy 2007). In principle the calculations were initiated using a potential energy curve (PEC) calculated *ab initio*. In practice, as detailed below, there are sufficient experimental data available for SiO that the PEC was actually characterised by fitting to these data using the program DPotFit 1.1 (Le Roy 2006).

2.1 Dipole moments

There appears to be no experimental measurements of any SiO transition dipoles and a single measurement of its permanent dipole moment as a function of the vibrational state by Raymonda et al. (1970). We determined a new dipole moment curve (DMC) using high level *ab initio* calculations. These are compared to previous, high-level *ab initio* determinations (Langhoff & Bauschlicher 1993; Drira et al. 1998; Chattopadhyaya et al. 2003) given below.

The *ab initio* calculations were performed using MOLPRO (Werner et al. 2010); we tested both the coupled cluster (CCSD(T)) and multi-reference configuration interaction (MRCI) methods with large basis sets. Our largest, and best, calculations used an aug-cc-pCV5Z basis set for the CCSD(T) study and aug-cc-pwCV5Z for MRCI; in both cases a correction due to the correlation of the core electrons was included but the corresponding correction due to relativistic effects was found to be very small and was neglected. These calculations give a value for the dipole at equilibrium of 3.10 D and 3.07 D, respectively, which bracket the experimentally determined value of 3.088 D (Raymonda et al. 1970). The *ab initio* DMC grid points were used directly in LEVEL.

Figure 1 compares the DMC's arising from our calculations with those obtained by Chattopadhyaya et al. (2003) (CCK) using an MRCI method. Other previous studies based on the use of similar CCSD(T) method gave curves similar to our CCSD(T) calculation. The CCSD(T) dipoles appear correct at short bondlengths but behave in an unphysical fashion for R > 2 Å. CCSD(T) is known to have problems as molecules are dissociated and so this model was not pursued. Again at short R our MRCI dipoles are similar to those of CCK. It is unclear why CCK's dipoles do not vary smoothly as R is increased but such behaviour is not consistent with obtaining reliable transition intensities. All calculations presented below therefore use our MRCI dipole moments.

2.2 Fitting the potential

The potential energy curve is essentially characterised by fitting to spectroscopic data. Measurements with very small estimated uncertainties were chosen for the fit such that any uncertainty in the fitted surface depends only on the accuracy of the fit




Figure 1. Ab initio dipole moment curves for SiO in its ground electronic state. MRCI and CCSD(T) are *ab initio* calculations performed as part of this work; CCK is the MRCI dipole of Chattopadhyaya et al. (2003).

Table 2. Fitting parameters used in the Extended Morse Oscillator potential, see eq. (1). (Uncertainties are given in parentheses in units of the last digit.)

N	ϕ_i
0	1.86869070(93)
1	-0.142883(30)
2	0.188883(98)
3	0.2255(14)
4	0.1959(37)
5	-0.471(16)
6	2.714(29)

and, for higher energies, the extrapolation beyond the experimental input values. Measurements for multiple isotopologues were used both to maximise the input data and to ensure that the resulting curve is applicable to all isotopologues, including the yet-to-be observed spectrum of ${}^{28}\text{Si}{}^{17}\text{O}$. The most comprehensive and accurate sets of available measurements are the infrared ro-vibrational sunspot lines detected by Campbell et al. (1995) and the microwave rotational laboratory lines recorded by Sanz et al. (2003).

The potential was expressed as an Extended Morse Oscillator potential function (EMOp):

$$V(r) = D_{\rm e} \left[1 - e^{-\phi(r)(r-r_{\rm e})} \right]^2, \tag{1}$$

where

$$\phi(r) = \sum_{i=0}^{N} \phi_i y_p(r, r_e)^i,$$
(2)
$$y_p(r, r_e) = \frac{r^p - r_e^p}{r^p + r_e^p}$$
(3)

and p was set to 2 and N to 6. Spectroscopically derived values for the dissociation energy, $D_e = 66620.0 \text{ cm}^{-1}$ (Campbell et al. 1995) and $r_e = 1.5097377$ Å(Sanz et al. 2003) were used.

Data for isotopologues were fitted simultaneously; attempts to include Born-Oppenheimer Breakdown (BOB) terms in the fit did not result in an improvement and were not pursued. Of the 1816 lines used in the fit, 1294 were ${}^{28}\text{Si}{}^{16}\text{O}$, 301 were ${}^{29}\text{Si}{}^{16}\text{O}$, 178 were ${}^{30}\text{Si}{}^{16}\text{O}$ and 43 were ${}^{28}\text{Si}{}^{18}\text{O}$. These data were used to determine seven constants with values for $r_{\rm e}$ and $D_{\rm e}$ held fixed. Fits were started from an *ab initio* PEC but were not sensitive to this starting point. The resulting fits reproduced the input experimental data within 0.01 cm⁻¹ and often, particularly for the pure rotational data, much better than this. Parameters resulting from the fit are given in table 2.

Table 3. Comparison of ²⁸Si¹⁶O partition functions.

$T(\mathbf{K})$	This work	Sauval & Tatum (1984)	Irwin (1981)
$I(\mathbf{R})$	THIS WOLK	Sauvai & Tatulli (1984)	II WIII (1981)
1000	1163.1	1169.2	1164.2
2000	3319.0	3316.1	3322.0
3000	6638.3	6618.3	6639.8
4000	11172.6	11000.2	11164.1
5000	16976.3	16629.7	16937.5
6000	24109.4	23833.9	24002.6
7000	32640.8	33080.1	32403.5
8000	42648.8	44989.4	42186.6
9000	54221.8	60371.5	53401.1

Table 4. Fitting parameters used to fit the partition functions, see eq. 4. Fits are valid for temperatures between 900 and 9000 K.

9066 -2.64536 692 4.920534		6 -4.29631977 7.19145985	$\begin{array}{r} 4 \\ 2.34443076 \end{array}$	-3.273169177
	106 7.19640021	7.19145985	9 94449076	
			2.54445070	12.27676929
0.251599	-1.35602888	-1.35513910	-12.16129465	3.08495699
629 -2.11202	-1.50490351	-1.49684190	9.00993487	-9.93550466
1.038226	0.90924988	9 0.905530086	-2.947037343	4.767331120
50025 -0.19750	04943 -0.18290280	96 -0.18230564	08 0.4708553438	-0.9089004600
6048 0.013508	35360 0.012911179	95 0.012878760	01 -0.029961405	7 0.0626337008

$\mathbf{2.3}$ Partition function

A partition function for ²⁸Si¹⁶O was calculated by summing all the calculated energy levels, see Table 7 below, using Excel. As we use all ro-vibrational energy levels there are no issues with convergence of this sum. Given the high accuracy of our energy levels, this determination should be more accurate than the previous determinations given by (Irwin 1981) and (Sauval & Tatum 1984). Table 3 compares with these previous studies and finds generally good agreement except the values of (Sauval & Tatum 1984) become too large at higher temperatures.

For ease of use, we fitted our partition function, Q, to a series expansion of the form used by Vidler & Tennyson (2000):

$$\log_{10} Q(T) = \sum_{n=0}^{5} a_n \left[\log T\right]^n$$
(4)

with the values given in Table 4.

Line list calculations $\mathbf{2.4}$

Line lists were calculated for the five isotopologues ²⁸Si¹⁶O, ²⁹Si¹⁶O, ³⁰Si¹⁶O, ²⁸Si¹⁸O and ²⁸Si¹⁷O. All rotation-vibration states were considered and transitions satisfying the dipole selection rule $\Delta J = \pm 1$. A summary of each line list is given in Table 7. These line lists in principle span frequencies up to 65 000 cm⁻¹; in practice, transitions above 10 000 cm⁻¹ are very weak.

The procedure described above was used to produce line lists, i.e. catalogues of transition frequencies $\tilde{\nu}_{ij}$ and Einstein coefficients A_{ij} , for five silicon oxide isotopologues ²⁸Si¹⁶O, ²⁹Si¹⁶O, ³⁰Si¹⁶O, ²⁸Si¹⁸O and ²⁸Si¹⁷O. The computed line lists for are given in the supplementary materials.

RESULTS 3

The line lists contain almost two million transitions each and, therefore, for compactness and ease of use, divided into separate energy levels and transitions file. This is done using the standard ExoMol format (Tennyson et al. 2013) which is based on a method originally developed for the BT2 line list (Barber et al. 2006). Extracts for the start of the ${}^{28}Si^{16}O$ files are given in tables 5 and 6. The full line list for each of these isotopologues can be downloaded from the CDS, via ftp://cdsarc.u-strasbg.fr/pub/cats/J/MNRAS/xxx/yy, or http://cdsarc.u-strasbg.fr/viz-bin/qcat?J/MNRAS//xxx/yy The line lists and partition function together with auxiliary data including the potential parameters and dipole moment functions, as well as the absorption spectrum given in cross section format (Hill et al. 2013), can all be obtained from there as well as at www.exomol.com.

Table 8 compares our SiO line lists with previous attempts to study this system: it only provides an assessment of the

Table 5. Extract from the state file for ²⁸Si¹⁶O. Full tables are available at http://cdsarc.u-strasbg.fr/cgi-bin/VizieR?-source=J/MNRAS/xxx/yy. The files contain 24 306 entries for ²⁸Si¹⁶O, 25 254 for ²⁸Si¹⁷O, 26 162 for ²⁸Si¹⁸O, 24 617 for ²⁹Si¹⁶O and 24 915 for ³⁰Si¹⁶O.

Ι	$ ilde{E}$	g	J	v
1	0.000000	1	0	0
2	1.448467	3	1	0
3	4.345384	5	2	0
4	8.690712	7	3	0
5	14.484267	9	4	0
6	21.726203	11	5	0

I: State counting number; \tilde{E} : State energy in cm⁻¹; g: State degeneracy; J: State rotational quantum number; v: State vibrational quantum number.

 28 Si 16 O. Full from Table 6. Extracts from the transitions file for tables are available http://cdsarc.u-strasbg.fr/cgi-bin/VizieR?-source=J/MNRAS/xxx/yy. The files contain 1 784 964 entries for ²⁸Si¹⁶O, 1 890 039 for ${}^{28}Si^{17}O$, 1 993 414 for ${}^{28}Si^{18}O$, 1 818 923 for ${}^{29}Si^{16}O$ and 1 852 656 for ${}^{30}Si^{16}O$.

Ι	F	A_{IF}
2	1	2.8438E-06
3	2	2.7301E-05
4	3	9.8720E-05
5	4	2.4265 E-04
6	5	4.8470 E-04
7	6	8.5037E-04

I: Upper state counting number; F: Lower state counting number; A_{IF} : Einstein A coefficient in s⁻¹.

quantity of data provided not its quality. However it is clear that our new line lists provide a much more comprehensive coverage of the problem. We believe they also represent a substantial improvement in accuracy. Figure 2 gives a room temperature comparison of our 28 Si¹⁶O line list with the most complete previous one due to Langhoff & Bauschlicher (1993). It is clear our line list is more complete.

Illustrative high resolution comparisons with ${}^{28}\text{Si}{}^{16}\text{O}$ observed sunspot transition frequencies are given in Table 9; as can be seen the comparison is excellent. Also given are the frequencies due to Langhoff & Bauschlicher (1993), who generated a PEC from the empirical parameters of Glenar et al. (1985), again the comparison is very good. Overall the error in Langhoff and Bauschlicher's frequencies, where available, is only about twice ours.

An important aim in fitting the PEC is to also predict spectroscopic data for higher vibrational states than used in the fit. To test this comparisons are made with the experimentally-derived vibrational frequencies of Shanker et al. (1976), which have an estimated uncertainty 1.0 cm⁻¹, and Bredohl et al. (1973), as re-assigned by Barrow & Stone (1975). Our predictions agree well with the experimentally-derived values; however those of Tipping & Chackerian (1981) (TC) disagree significantly for the higher vibrational states. Indeed by v = 35 our vibrational term value is 3.3 cm^{-1} lower than TC's. TC were limited in the vibrational states they could use in their fit, extending only to v = 5 from the spectra of Lovas et al. (1981), compared to the line frequencies for vibrational transitions used which extend to v = 13.

Since there are no measured intensities to compare with, we compare the results of our calculations with astronomical spectra. Sunspots display many SiO features over a wide range of wavelengths. High resolution sunspot spectra have been compiled by Wallace et al. (1996). Figure 3 compares small regions of these spectra with our line lists. Note that only the theoretical spectra, generated at T=3200 K, are given in the absolute units of cm/molecule). These spectra, which are chosen to cover regions where Δv is both 1 and 2, illustrate the excellent agreement between our line positions and the observed features. Our intensities are also well correlated with the observed ones. We note that the sunspot spectra show many other features in this figure. These features are either due to water (Polyansky et al. 1997) or so to some so-far unidentified species.

Figure 4 compares the spectra of four red giant Wallace & Hinkle (2002) with our simulated spectrum of ²⁸Si¹⁶O at a temperature of 3200 K. These spectra are for the $\Delta v = 2$ overtone spectrum. To fully reproduce the observed stellar spectrum

	$^{28}\mathrm{Si}^{16}\mathrm{O}$	$^{29}\mathrm{Si}^{16}\mathrm{O}$	$^{30}\mathrm{Si}^{16}\mathrm{O}$	$^{28}\mathrm{Si}^{18}\mathrm{O}$	$^{28}\mathrm{Si}^{17}\mathrm{O}$
Maximum v	95	95	96	98	97
Maximum J	408	410	413	423	416
Number of lines	1784964	1818923	1852656	1993414	1890039

Table 7.	Summary	of our	SiO	line l	ists.
----------	---------	--------	----------------------	--------	-------




Figure 2. Absorption spectra of ²⁸Si¹⁶O at T=296 K: ExoMol versus (Langhoff & Bauschlicher 1993).




Figure 3. Sunspot absorption spectra (Wallace et al. 1996), continuous line and our predicted 28 Si 16 O absorptions at 3200 K, vertical lines. The absolute intensity scale refers to the calculated spectra.

Table 8. Summary comparison of SiO rotation-vibration line lists. Given are the isotopologues considered, the maximum values for vibrational (v) and rotational (J) states considered, the maximum change in vibrational state (Δv) , whether intensity information and a partition function are provided.

Reference	Lovas et al. (1981)	Glenar et al. (1985)	Langhoff & Bauschlicher (1993)	Drira et al. (1997)	This work
Isotopes	1	3	3	1	5
maximum v	6	10	15	40	98
maximum J	98	86	250	1	423
maximum Δv	2	1	4	3	98
Intensities?	No	No	Yes	Yes	Yes
Partition Function?	No	No	No	No	Yes




Figure 4. Red giant spectra due to Wallace & Hinkle (2002) compared to a synthesized ${}^{28}Si^{16}O$ absorption spectrum using T = 3200 K (bottom panel).

would require running a stellar model which is beyond the scope of this paper. However it clear that both the line positions and general band structure predicted by our calculations are in very good agreement with the observations.

4 CONCLUSIONS

We present comprehensive line lists for the five most important isotopologues of SiO. These are based on the direct solution of the nuclear motion Schrödinger equation using a potential energy curve obtained by fitting to an extensive data set of measured transitions. These data are reproduced to near experimental accuracy resulting in high accuracy line positions. A new *ab initio* dipole moment is computed, which appears to behave more physically at large internuclear separations. This dipole is used to compute Einstein A coefficients for all possible dipole-allowed transitions within each SiO isotopologue. The result is a comprehensive line list for each species, including the first data for ²⁸Si¹⁷O. The line lists can be downloaded from the CDS, via ftp://cdsarc.u-strasbg.fr/pub/cats/J/MNRAS/, or from http://cdsarc.u-strasbg.fr/viz-bin/qcat?J/MNRAS/, or from www.exomol.com.

ACKNOWLEDGEMENT

This work is supported by ERC Advanced Investigator Project 267219.

© 2012 RAS, MNRAS **000**, 1–10

Table 9. Comparison of theoretically predicted ro-vibrational wavenumbers, in cm^{-1} , with the high-resolution sunspot observations of Campbell et al. (1995).

J'	J''	v'	$v^{\prime\prime}$	Obs.	This	work	Langhoff & Ba	auschlicher (1993)
					Calc.	ObsCalc.	Calc.	ObsCalc
9	10	1	0	1214.6810	1214.6829	-0.0019	1214.68	0.00
9	10	2	1	1202.8878	1202.8876	0.0002	1202.89	-0.00
9	10	3	2	1191.1305	1191.1291	0.0014	1191.13	0.00
9	10	4	3	1179.4093	1179.4070	0.0023	1179.41	-0.00
9	10	8	7	1173.4035 1132.8751	1173.4070 1132.8725	0.0025	1132.87	0.01
9	10	10	9	1102.8731 1109.8133	1132.8723 1109.8123	0.0020	1109.81	0.00
9 9	10	10					1098.33	0.00
			10	1098.3324	1098.3329	-0.0005		
9	10	13	12	1075.4744	1075.4740	0.0004	1075.48	-0.01
11	10	2	1	1232.8767	1232.8735	0.0032	1232.87	0.01
11	10	3	2	1220.9058	1220.9040	0.0018	1220.91	-0.00
11	10	4	3	1208.9734	1208.9711	0.0023	1208.97	0.00
11	10	5	4	1197.0759	1197.0742	0.0017	1197.08	-0.00
11	10	6	5	1185.2130	1185.2129	0.0001	1185.21	0.00
11	10	7	6	1173.3880	1173.3867	0.0013	1173.39	-0.00
11	10	8	7	1161.5964	1161.5954	0.0010	1161.60	-0.00
11	10	10	9	1138.1158	1138.1155	0.0003	1138.12	-0.00
10	11	1	0	1213.1327	1213.1350	-0.0023	1213.13	0.00
10	11	2	1	1201.3510	1201.3499	0.0011	1201.35	0.00
10	11	3	2	1189.6031	1189.6015	0.0016	1189.60	0.00
10	11	5	4	1166.2155	1166.2135	0.0020	1166.21	0.01
10	11	6	5	1154.5742	1154.5727	0.0015	1154.57	0.00
10	11	7	6	1142.9704	1142.9668	0.0036	1142.97	0.00
10	11	9	8	1119.8600	1119.8586	0.0014	1119.86	0.00
10	11	12	11	1085.4509	1085.4502	0.0007	1085.45	0.00
10	11	13	12	1074.0547	1074.0474	0.0073	1074.05	0.00
10		10		101110011	101 110 11 1	0.0010	1011100	0.00
125	126	2	1	965.1065	965.1051	0.0014	965.09	0.02
125	126	3	2	954.5828	954.5856	-0.0028	954.57	0.01
$125 \\ 125$	126	4	3	944.0942	944.0980	-0.0038	944.09	0.00
$120 \\ 126$	$120 \\ 127$	1	0	973.1300	973.1257	0.0043	973.11	0.02
$120 \\ 126$	127 127	2	1	962.5855	962.5854	0.0001	962.57	0.02
$120 \\ 126$	$127 \\ 127$	2 4	3	902.5855 941.5952	902.3834 941.6004	-0.0052	962.57 941.59	0.0
$120 \\ 127$	127			941.5952 970.5914				0.02
		$\frac{1}{2}$	0		970.5876	0.0038	970.57	
127	128		1	960.0589	960.0585	0.0004	960.04	0.02
127	128	3	2	949.5598	949.5612	-0.0014	949.55	0.0
127	128	4	3	939.0904	939.0956	-0.0052	939.08	0.0
128	129	1	0	968.0462	968.0425	0.0037	968.03	0.02
128	129	2	1	957.5253	957.5244	0.0009	957.51	0.02
128	129	3	2	947.0347	947.0382	-0.0035	947.02	0.0
129	130	2	1	954.9832	954.9833	-0.0001	954.97	0.03
130	131	1	0	962.9348	962.9306	0.0042	962.91	0.02
130	131	2	1	952.4351	952.4349	0.0002	952.42	0.02
130	131	3	2	941.9681	941.9710	-0.0029	941.95	0.02
131	132	1	0	960.3668	960.3641	0.0027	960.35	0.02
131	132	2	1	949.8787	949.8795	-0.0008	949.86	0.02
131	132	3	2	939.4210	939.4266	-0.0056	939.41	0.0
132	133	1	0	957.7940	957.7905	0.0035	957.77	0.02
132	133	2	1	947.3225	947.3171	0.0054	947.30	0.02
132	133	3	2	936.8721	936.8753	-0.0032	936.85	0.02
134	135	2	1	942.1713	942.1709	0.0002	942.15	0.02
$134 \\ 135$	136	2	1	939.5876	939.5874	0.0004	939.56	0.03
$135 \\ 136$	$130 \\ 137$	1	0	939.3870 947.4301	939.3874 947.4255	0.0046	947.40	0.03
$130 \\ 137$	137	1	0	947.4301 944.8216	947.4255 944.8167	0.0040	944.79	0.03
137	138	2	1	934.3987	934.3992	-0.0005	934.37	0.03
139	140	2	1	929.1778	929.1832	-0.0054	929.15	0.03
$140 \\ 140$	$\begin{array}{c} 141 \\ 141 \end{array}$	$\frac{1}{2}$	0	936.9546	936.9489	0.0057	936.92	0.03
		•)	1	926.5650	926.5649	0.0001	926.53	0.04

Table 10. Comparison of theoretically predicted vibrational spacings, in cm⁻¹, with the low-resolution experiments of Bredohl et al. (1973) for $13 \le v \le 17$, and Bredohl et al. (1973), as re-assigned by Barrow & Stone (1975), for $23 \le v \le 28$.

v' - v"	Obs.	Calc.	0	bserved – Calculated
		This work	This work	Tipping & Chackerian (1981)
14 - 13	1078.5	1077.72	0.8	0,8
15 - 14	1067.0	1066.27	0.7	0.7
16 - 15	1055.0	1054.86	0.1	0.1
17 - 16	1044.0	1043.48	0.5	0.4
18 - 17	1033.0	1032.13	0.9	0.8
24 - 23	964.69	964.70	-0.01	-0.36
25 - 24	953.55	953.58	-0.03	-0.04
26 - 25	942.28	942.48	-0.20	-0.03
27 - 26	931.37	931.41	-0.04	-0.52
28 - 27	920.22	920.38	-0.16	-0.74

REFERENCES

Aitken D. K., Smith C. H., James S. D., Roche P. F., Hyland A. R., McGregor P. J., 1988, MNRAS, 231, 7

- Assaf K. A., Diamond P. J., Richards A. M. S., Gray M. D., 2011, MNRAS, 415, 1083
- Barber R. J., Tennyson J., Harris G. J., Tolchenov R. N., 2006, MNRAS, 368, 1087

Barrow R. F., Stone T. J., 1975, J. Phys. B: At. Mol. Opt. Phys., 8, L13

Bredohl H., Cornet R., Dubois I., Remy F., 1973, Can. J. Phys., 51, 2332

Campbell J. M., Klapstein D., Dulick M., Bernath P. F., 1995, ApJS, 101, 237

Chattopadhyaya S., Chattopadhyay A., Das K. K., 2003, Theochem-J. Mol. Struct., 639, 177

Cho S. H., Saito S., 1998, ApJ, 496, L51

Cohen M., Witteborn F. C., Carbon D. F., Augason G., Wooden D., Bregman J., Goorvitch D., 1992, AJ., 104, 2045 Cotton W. D., Ragland S., Danchi W. C., 2011, ApJ, 736, 96

Deguchi S., Koike K., Kuno N., Matsunaga N., Nakashima J.-i., Takahashi S., 2011, Publ. Astron. Soc. Jpn., 63, 309

Drira I., Hure J. M., Spielfiedel A., Feautrier N., Roueff E., 1997, A&A, 319, 720

Drira I., Spielfiedel A., Edwards S., Feautrier N., 1998, J. Quant. Spectrosc. Radiat. Transf., 60, 1

Glenar D. A., Hill A. R., Jennings D. E., Brault J. W., 1985, J. Mol. Spectrosc., 111, 403

Hill C., Yurchenko S. N., Tennyson J., 2013, Icarus, (in press)

Irwin A. W., 1981, ApJS, 45, 621

Kaifu N., Buhl D., Snyder L., 1975, ApJ, 195, 359

Langhoff S. R., Bauschlicher C. W., 1993, Chem. Phys. Lett., 211, 305

Le Roy R. J., 2006, DPotFit 1.1 A Computer Program for Fitting Diatomic Molecule Spectral Data to Potential Energy Functions. University of Waterloo Chemical Physics Research Report CP-662R, http://leroy.uwaterloo.ca/programs/

Le Roy R. J., 2007, LEVEL 8.0 A Computer Program for Solving the Radial Schrödinger Equation for Bound and Quasibound

Levels. University of Waterloo Chemical Physics Research Report CP-663, http://leroy.uwaterloo.ca/programs/

Lovas F. J., Maki A. G., Olson W. B., 1981, J. Mol. Spectrosc., 87, 449

Mollaaghababa R., Gottlieb C. A., Vrtilek J. M., Thaddeus P., 1991, ApJ, 368, L19

Motret O., Coursimault F., Viladrosa R., Pouvesle J. M., 2002, High Temp. Mater. Process, 6, 305

Nakashima J.-I., Deguchi S., Imai H., Kemball A., Lewis B. M., 2011, ApJ, 728, 76

Polyansky O. L., Zobov N. F., Viti S., Tennyson J., Bernath P. F., Wallace L., 1997, Science, 277, 346

Raymonda J. W., Muenter J. S., Klemperer W. A., 1970, J. Chem. Phys., 52, 3458

Rinsland C. P., Wing R. F., 1982, ApJ, 262, 201

Sanz M. E., McCarthy M. C., Thaddeus P., 2003, J. Chem. Phys., 119, 11715

Sauval A. J., Tatum J. B., 1984, ApJS, 56, 193

Schaefer L., Lodders K., Fegley, Jr. B., 2012, ApJ, 755, 41

Shanker R., Linton C., Verma R. D., 1976, J. Mol. Spectrosc., 60, 197

Snyder L. E., Buhl D., 1974, ApJ, 189, L31

Tennyson J., Hill C., Yurchenko S. N., 2013, in , AIP Conference Proceedings, AIP, New York

Tennyson J., Yurchenko S. N., 2012, MNRAS, 425, 21

Tipping R. H., Chackerian C., 1981, J. Mol. Spectrosc., 88, 352

Vidler M., Tennyson J., 2000, J. Chem. Phys., 113, 9766

Vlemmings W. H. T., Humphreys E. M. L., Franco-Hernandez R., 2011, ApJ, 728, 149

Wallace L., Hinkle K., 2002, AJ., 124, 3393

Wallace L., Livingston W., Hinkle K., Bernath P., 1996, ApJS, 106, 165 Werner H. J., Knowles P. J., Lindh R., Manby F. R., Schütz M., 2010, MOLPRO, a package of ab initio programs. See http://www.molpro.net/

Wilson R. W., Penzias A. A., Jefferts K. B., Kutner M., Thaddeus P., 1971, ApJ, 167, L97

Wooldridge M. S., Danczyk S. A., Wu J. F., 2000, J. Quant. Spectrosc. Radiat. Transf., 64, 573

Yadin B., Vaness T., Conti P., Hill C., Yurchenko S. N., Tennyson J., 2012, MNRAS, 425, 34