

CERN-PH-EP-2013-136 LHCb-PAPER-2013-044 November 1, 2013

Observation of the decay $B_c^+\to B_s^0\pi^+$

The LHC_b collaboration^{[†](#page-0-0)}

Abstract

The result of a search for the decay $B_c^+ \to B_s^0 \pi^+$ is presented, using the $B_s^0 \to D_s^- \pi^+$ and $B_s^0 \to J/\psi \phi$ channels. The analysis is based on a data sample of pp collisions collected with the LHCb detector, corresponding to an integrated luminosity of 1 fb^{-1} taken at a center-of-mass energy of 7 TeV, and 2 fb^{-1} taken at 8 TeV. The decay $B_c^+ \to B_s^0 \pi^+$ is observed with significance in excess of five standard deviations independently in both decay channels. The measured product of the ratio of cross-sections and branching fraction is

$$
\frac{\sigma(B_c^+)}{\sigma(B_s^0)} \times \mathcal{B}(B_c^+ \to B_s^0 \pi^+) = \left(2.37 \pm 0.31 \, (\text{stat}) \, \pm 0.11 \, (\text{syst}) \, {}^{+0.17}_{-0.13} \, (\tau_{B_c^+})\right) \times 10^{-3},
$$

in the pseudorapidity range $2 < \eta(B) < 5$, where the first uncertainty is statistical, the second is systematic and the third is due to the uncertainty on the B_c^+ lifetime. This is the first observation of a B meson decaying to another B meson via the weak interaction.

> Published in Phys. Rev. Lett. 111, 181801 (2013) c CERN on behalf of the LHCb collaboration, license [CC-BY-3.0.](http://creativecommons.org/licenses/by/3.0/)

[†]Authors are listed on the following pages.

LHCb collaboration

R. Aaij⁴⁰, B. Adeva³⁶, M. Adinolfi⁴⁵, C. Adrover⁶, A. Affolder⁵¹, Z. Ajaltouni⁵, J. Albrecht⁹, F. Alessio³⁷, M. Alexander⁵⁰, S. Ali⁴⁰, G. Alkhazov²⁹, P. Alvarez Cartelle³⁶, A.A. Alves Jr^{24,37}, S. Amato², S. Amerio²¹, Y. Amhis⁷, L. Anderlini^{17,f}, J. Anderson³⁹, R. Andreassen⁵⁶, J.E. Andrews⁵⁷, R.B. Appleby⁵³, O. Aquines Gutierrez¹⁰, F. Archilli¹⁸, A. Artamonov³⁴, M. Artuso⁵⁸, E. Aslanides⁶, G. Auriemma^{24,*m*}, M. Baalouch⁵, S. Bachmann¹¹, J.J. Back⁴⁷, C. Baesso⁵⁹, V. Balagura³⁰, W. Baldini¹⁶, R.J. Barlow⁵³, C. Barschel³⁷, S. Barsuk⁷, W. Barter⁴⁶, Th. Bauer⁴⁰, A. Bay³⁸, J. Beddow⁵⁰, F. Bedeschi²², I. Bediaga¹, S. Belogurov³⁰, K. Belous³⁴, I. Belyaev³⁰, E. Ben-Haim⁸, G. Bencivenni¹⁸, S. Benson⁴⁹, J. Benton⁴⁵, A. Berezhnoy³¹, R. Bernet³⁹, M.-O. Bettler⁴⁶, M. van Beuzekom⁴⁰, A. Bien¹¹, S. Bifani⁴⁴, T. Bird⁵³, A. Bizzeti^{17,h}, P.M. Bjørnstad⁵³, T. Blake³⁷, F. Blanc³⁸, J. Blouw¹⁰, S. Blusk⁵⁸, V. Bocci²⁴, A. Bondar³³, N. Bondar²⁹, W. Bonivento¹⁵, S. Borghi⁵³, A. Borgia⁵⁸, T.J.V. Bowcock⁵¹, E. Bowen³⁹, C. Bozzi¹⁶, T. Brambach⁹, J. van den Brand⁴¹, J. Bressieux³⁸, D. Brett⁵³, M. Britsch¹⁰, T. Britton⁵⁸, N.H. Brook⁴⁵, H. Brown⁵¹, A. Bursche³⁹, G. Busetto^{21,q}, J. Buytaert³⁷, S. Cadeddu¹⁵, O. Callot⁷, M. Calvi^{20,j}, M. Calvo Gomez^{35,n}, A. Camboni³⁵, P. Campana^{18,37}, D. Campora Perez³⁷, A. Carbone^{14,c}, G. Carboni^{23,k}, R. Cardinale^{19,i}, A. Cardini¹⁵, H. Carranza-Mejia⁴⁹, L. Carson⁵², K. Carvalho Akiba², G. Casse⁵¹, L. Cassina¹, L. Castillo Garcia³⁷, M. Cattaneo³⁷, Ch. Cauet⁹, R. Cenci⁵⁷, M. Charles⁵⁴, Ph. Charpentier³⁷, P. Chen^{3,38}, S.-F. Cheung⁵⁴, N. Chiapolini³⁹, M. Chrzaszcz^{39,25}, K. Ciba³⁷, X. Cid Vidal³⁷, G. Ciezarek⁵², P.E.L. Clarke⁴⁹, M. Clemencic³⁷, H.V. Cliff⁴⁶, J. Closier³⁷, C. Coca²⁸, V. Coco⁴⁰, J. Cogan⁶, E. Cogneras⁵, P. Collins³⁷, A. Comerma-Montells³⁵, A. Contu^{15,37}, A. Cook⁴⁵, M. Coombes⁴⁵, S. Coquereau⁸, G. Corti³⁷, B. Couturier³⁷, G.A. Cowan⁴⁹, D.C. Craik⁴⁷, S. Cunliffe⁵², R. Currie⁴⁹, C. D'Ambrosio³⁷, P. David⁸, P.N.Y. David⁴⁰, A. Davis⁵⁶, I. De Bonis⁴, K. De Bruyn⁴⁰, S. De Capua⁵³, M. De Cian¹¹, J.M. De Miranda¹, L. De Paula², W. De Silva⁵⁶, P. De Simone¹⁸, D. Decamp⁴, M. Deckenhoff⁹, L. Del Buono⁸, N. Déléage⁴, D. Derkach⁵⁴, O. Deschamps⁵, F. Dettori⁴¹, A. Di Canto¹¹, H. Dijkstra³⁷, M. Dogaru²⁸, S. Donleavy⁵¹, F. Dordei¹¹, A. Dosil Suárez³⁶, D. Dossett⁴⁷, A. Dovbnya⁴², F. Dupertuis³⁸, P. Durante³⁷, R. Dzhelyadin³⁴, A. Dziurda²⁵, A. Dzyuba²⁹, S. Easo⁴⁸, U. Egede⁵², V. Egorychev³⁰, S. Eidelman³³, D. van Eijk⁴⁰, S. Eisenhardt⁴⁹, U. Eitschberger⁹, R. Ekelhof⁹, L. Eklund^{50,37}, I. El Rifai⁵, Ch. Elsasser³⁹, A. Falabella^{14,e}, C. Färber¹¹, C. Farinelli⁴⁰, S. Farry⁵¹, D. Ferguson⁴⁹, V. Fernandez Albor³⁶, F. Ferreira Rodrigues¹, M. Ferro-Luzzi³⁷, S. Filippov³², M. Fiore^{16,e}, C. Fitzpatrick³⁷, M. Fontana¹⁰, F. Fontanelli^{19,i}, R. Forty³⁷, O. Francisco², M. Frank³⁷, C. Frei³⁷, M. Frosini^{17,37,f}, E. Furfaro^{23,k}, A. Gallas Torreira³⁶, D. Galli^{14,c}, M. Gandelman², P. Gandini⁵⁸, Y. Gao³, J. Garofoli⁵⁸, P. Garosi⁵³, J. Garra Tico⁴⁶, L. Garrido³⁵, C. Gaspar³⁷, R. Gauld⁵⁴, E. Gersabeck¹¹, M. Gersabeck⁵³, T. Gershon⁴⁷, Ph. Ghez⁴, V. Gibson⁴⁶, L. Giubega²⁸, V.V. Gligorov³⁷, C. Göbel⁵⁹, D. Golubkov³⁰, A. Golutvin^{52,30,37}, A. Gomes², P. Gorbounov^{30,37}, H. Gordon³⁷, M. Grabalosa Gándara⁵, R. Graciani Diaz³⁵, L.A. Granado Cardoso³⁷, E. Graugés³⁵, G. Graziani¹⁷, A. Grecu²⁸, E. Greening⁵⁴, S. Gregson⁴⁶, P. Griffith⁴⁴, O. Grünberg⁶⁰, B. Gui⁵⁸, E. Gushchin³², Yu. Guz^{34,37}, T. Gys³⁷, C. Hadjivasiliou⁵⁸, G. Haefeli³⁸, C. Haen³⁷, S.C. Haines⁴⁶, S. Hall⁵², B. Hamilton⁵⁷, T. Hampson⁴⁵, S. Hansmann-Menzemer¹¹, N. Harnew⁵⁴, S.T. Harnew⁴⁵, J. Harrison⁵³, T. Hartmann⁶⁰, J. He³⁷, T. Head³⁷, V. Heijne⁴⁰, K. Hennessy⁵¹, P. Henrard⁵, J.A. Hernando Morata³⁶, E. van Herwijnen³⁷, M. Heß⁶⁰, A. Hicheur¹, E. Hicks⁵¹, D. Hill⁵⁴, M. Hoballah⁵, C. Hombach⁵³, W. Hulsbergen⁴⁰, P. Hunt⁵⁴, T. Huse⁵¹, N. Hussain⁵⁴, D. Hutchcroft⁵¹, D. Hynds⁵⁰, V. Iakovenko⁴³, M. Idzik²⁶, P. Ilten¹², R. Jacobsson³⁷, A. Jaeger¹¹,

E. Jans⁴⁰, P. Jaton³⁸, A. Jawahery⁵⁷, F. Jing³, M. John⁵⁴, D. Johnson⁵⁴, C.R. Jones⁴⁶, C. Joram³⁷, B. Jost³⁷, M. Kaballo⁹, S. Kandybei⁴², W. Kanso⁶, M. Karacson³⁷, T.M. Karbach³⁷, I.R. Kenyon⁴⁴, T. Ketel⁴¹, B. Khanji²⁰, O. Kochebina⁷, I. Komarov³⁸, R.F. Koopman⁴¹, P. Koppenburg⁴⁰, M. Korolev³¹, A. Kozlinskiy⁴⁰, L. Kravchuk³², K. Kreplin¹¹, M. Kreps⁴⁷, G. Krocker¹¹, P. Krokovny³³, F. Kruse⁹, M. Kucharczyk^{20,25,37,j}, V. Kudryavtsev³³, K. Kurek²⁷, T. Kvaratskheliya^{30,37}, V.N. La Thi³⁸, D. Lacarrere³⁷, G. Lafferty⁵³, A. Lai¹⁵, D. Lambert⁴⁹, R.W. Lambert⁴¹, E. Lanciotti³⁷, G. Lanfranchi¹⁸, C. Langenbruch³⁷, T. Latham⁴⁷, C. Lazzeroni⁴⁴, R. Le Gac⁶, J. van Leerdam⁴⁰, J.-P. Lees⁴, R. Lefèvre⁵, A. Leflat³¹, J. Lefrançois⁷, S. Leo²², O. Leroy⁶, T. Lesiak²⁵, B. Leverington¹¹, Y. Li³, L. Li Gioi⁵, M. Liles⁵¹, R. Lindner³⁷, C. Linn¹¹, B. Liu³, G. Liu³⁷, S. Lohn³⁷, I. Longstaff⁵⁰, J.H. Lopes², N. Lopez-March³⁸, H. Lu³, D. Lucchesi^{21,q}, J. Luisier³⁸, H. Luo⁴⁹, O. Lupton⁵⁴, F. Machefert⁷, I.V. Machikhiliyan^{4,30}, F. Maciuc²⁸, O. Maev^{29,37}, S. Malde⁵⁴, G. Manca^{15,d}, G. Mancinelli⁶, J. Maratas⁵, U. Marconi¹⁴, P. Marino^{22,s}, R. Märki³⁸, J. Marks¹¹, G. Martellotti²⁴, A. Martens⁸, A. Martín Sánchez⁷, M. Martinelli⁴⁰, D. Martinez Santos^{41,37}, D. Martins Tostes², A. Martynov³¹, A. Massafferri¹, R. Matev³⁷, Z. Mathe³⁷, C. Matteuzzi²⁰, E. Maurice⁶, A. Mazurov^{16,32,37,e}, J. McCarthy⁴⁴, A. McNab⁵³, R. McNulty¹², B. McSkelly⁵¹, B. Meadows^{56,54}, F. Meier⁹, M. Meissner¹¹, M. Merk⁴⁰, D.A. Milanes⁸, M.-N. Minard⁴, J. Molina Rodriguez⁵⁹, S. Monteil⁵, D. Moran⁵³, P. Morawski²⁵, A. Mordà⁶, M.J. Morello^{22,s}, R. Mountain⁵⁸, I. Mous⁴⁰, F. Muheim⁴⁹, K. Müller³⁹, R. Muresan²⁸, B. Muryn²⁶, B. Muster³⁸, P. Naik⁴⁵, T. Nakada³⁸, R. Nandakumar⁴⁸, I. Nasteva¹, M. Needham⁴⁹, S. Neubert³⁷, N. Neufeld 37 , A.D. Nguyen 38 , T.D. Nguyen 38 , C. Nguyen-Mau 38,o , M. Nicol⁷, V. Niess 5 , R. Niet⁹, N. Nikitin³¹, T. Nikodem¹¹, A. Nomerotski⁵⁴, A. Novoselov³⁴, A. Oblakowska-Mucha²⁶, V. Obraztsov³⁴, S. Oggero⁴⁰, S. Ogilvy⁵⁰, O. Okhrimenko⁴³, R. Oldeman^{15,d}, M. Orlandea²⁸, J.M. Otalora Goicochea², P. Owen⁵², A. Oyanguren³⁵, B.K. Pal⁵⁸, A. Palano^{13,b}, M. Palutan¹⁸, J. Panman³⁷, A. Papanestis⁴⁸, M. Pappagallo⁵⁰, C. Parkes⁵³, C.J. Parkinson⁵², G. Passaleva¹⁷, G.D. Patel⁵¹, M. Patel⁵², G.N. Patrick⁴⁸, C. Patrignani^{19,*i*}, C. Pavel-Nicorescu²⁸, A. Pazos Alvarez³⁶, A. Pearce⁵³, A. Pellegrino⁴⁰, G. Penso^{24,*l*}, M. Pepe Altarelli³⁷, S. Perazzini^{14,c}, E. Perez Trigo³⁶, A. Pérez-Calero Yzquierdo³⁵, P. Perret⁵, M. Perrin-Terrin⁶, L. Pescatore⁴⁴, E. Pesen⁶¹, G. Pessina²⁰, K. Petridis⁵², A. Petrolini^{19,*i*}, A. Phan⁵⁸, E. Picatoste Olloqui³⁵, B. Pietrzyk⁴, T. Pilař⁴⁷, D. Pinci²⁴, S. Playfer⁴⁹, M. Plo Casasus³⁶, F. Polci⁸, G. Polok²⁵, A. Poluektov^{47,33}, E. Polycarpo², A. Popov³⁴, D. Popov¹⁰, B. Popovici²⁸, C. Potterat³⁵, A. Powell⁵⁴, J. Prisciandaro³⁸, A. Pritchard⁵¹, C. Prouve⁷, V. Pugatch⁴³, A. Puig Navarro³⁸, G. Punzi^{22,r}, W. Qian⁴, J.H. Rademacker⁴⁵, B. Rakotomiaramanana³⁸, M.S. Rangel², I. Raniuk⁴², N. Rauschmayr³⁷, G. Raven⁴¹, S. Redford⁵⁴, M.M. Reid⁴⁷, A.C. dos Reis¹, S. Ricciardi⁴⁸, A. Richards⁵², K. Rinnert⁵¹, V. Rives Molina³⁵, D.A. Roa Romero⁵, P. Robbe⁷, D.A. Roberts⁵⁷, A.B. Rodrigues¹, E. Rodrigues⁵³, P. Rodriguez Perez³⁶, S. Roiser³⁷, V. Romanovsky³⁴, A. Romero Vidal³⁶, J. Rouvinet³⁸, T. Ruf³⁷, F. Ruffini²², H. Ruiz³⁵, P. Ruiz Valls³⁵, G. Sabatino^{24,k}, J.J. Saborido Silva³⁶, N. Sagidova²⁹, P. Sail⁵⁰, B. Saitta^{15,d}, V. Salustino Guimaraes², B. Sanmartin Sedes³⁶, R. Santacesaria²⁴, C. Santamarina Rios³⁶, E. Santovetti^{23,k}, M. Sapunov⁶, A. Sarti¹⁸, C. Satriano^{24,*m*}, A. Satta²³, M. Savrie^{16,e}, D. Savrina^{30,31}, M. Schiller⁴¹, H. Schindler³⁷, M. Schlupp⁹, M. Schmelling¹⁰, B. Schmidt³⁷, O. Schneider³⁸, A. Schopper³⁷, M.-H. Schune⁷, R. Schwemmer³⁷, B. Sciascia¹⁸, A. Sciubba²⁴, M. Seco³⁶, A. Semennikov³⁰, K. Senderowska²⁶, I. Sepp⁵², N. Serra³⁹, J. Serrano⁶, P. Seyfert¹¹, M. Shapkin³⁴, I. Shapoval^{16,42,e}, P. Shatalov³⁰, Y. Shcheglov²⁹, T. Shears⁵¹, L. Shekhtman³³, O. Shevchenko⁴², V. Shevchenko³⁰, A. Shires⁹, R. Silva Coutinho⁴⁷, M. Sirendi⁴⁶, N. Skidmore⁴⁵, T. Skwarnicki⁵⁸, N.A. Smith⁵¹, E. Smith^{54,48},

E. Smith⁵², J. Smith⁴⁶, M. Smith⁵³, M.D. Sokoloff⁵⁶, F.J.P. Soler⁵⁰, F. Soomro³⁸, D. Souza⁴⁵, B. Souza De Paula², B. Spaan⁹, A. Sparkes⁴⁹, P. Spradlin⁵⁰, F. Stagni³⁷, S. Stahl¹¹, O. Steinkamp³⁹, S. Stevenson⁵⁴, S. Stoica²⁸, S. Stone⁵⁸, B. Storaci³⁹, M. Straticiuc²⁸, U. Straumann³⁹, V.K. Subbiah³⁷, L. Sun⁵⁶, W. Sutcliffe⁵², S. Swientek⁹, V. Syropoulos⁴¹, M. Szczekowski²⁷, P. Szczypka^{38,37}, D. Szilard², T. Szumlak²⁶, S. T'Jampens⁴, M. Teklishyn⁷, E. Teodorescu²⁸, F. Teubert³⁷, C. Thomas⁵⁴, E. Thomas³⁷, J. van Tilburg¹¹, V. Tisserand⁴, M. Tobin³⁸, S. Tolk⁴¹, D. Tonelli³⁷, S. Topp-Joergensen⁵⁴, N. Torr⁵⁴, E. Tournefier^{4,52}, S. Tourneur³⁸, M.T. Tran³⁸, M. Tresch³⁹, A. Tsaregorodtsev⁶, P. Tsopelas⁴⁰, N. Tuning^{40,37}, M. Ubeda Garcia³⁷, A. Ukleja²⁷, A. Ustyuzhanin^{52,p}, U. Uwer¹¹, V. Vagnoni¹⁴, G. Valenti¹⁴, A. Vallier⁷, R. Vazquez Gomez¹⁸, P. Vazquez Regueiro³⁶, C. Vázquez Sierra³⁶, S. Vecchi¹⁶, J.J. Velthuis⁴⁵, M. Veltri^{17,9}, G. Veneziano³⁸, M. Vesterinen³⁷, B. Viaud⁷, D. Vieira², X. Vilasis-Cardona^{35,n}, A. Vollhardt³⁹, D. Volyanskyy¹⁰, D. Voong⁴⁵, A. Vorobyev²⁹, V. Vorobyev³³, C. Vo 6^{60} , H. Voss¹⁰, J.A. de Vries⁴⁰, R. Waldi⁶⁰, C. Wallace⁴⁷, R. Wallace¹², S. Wandernoth¹¹, J. Wang⁵⁸, D.R. Ward⁴⁶, N.K. Watson⁴⁴, A.D. Webber⁵³, D. Websdale⁵², M. Whitehead⁴⁷, J. Wicht³⁷, J. Wiechczynski²⁵, D. Wiedner¹¹, L. Wiggers⁴⁰, G. Wilkinson⁵⁴, M.P. Williams^{47,48}, M. Williams⁵⁵, F.F. Wilson⁴⁸, J. Wimberley⁵⁷, J. Wishahi⁹, W. Wislicki²⁷, M. Witek²⁵, S.A. Wotton⁴⁶, S. Wright⁴⁶, S. Wu³, K. Wyllie³⁷, Y. Xie^{49,37}, Z. Xing⁵⁸, Z. Yang³, X. Yuan³, O. Yushchenko³⁴, M. Zangoli¹⁴, M. Zavertyaev^{10,a}, F. Zhang³, L. Zhang⁵⁸, W.C. Zhang¹², Y. Zhang³, A. Zhelezov¹¹, A. Zhokhov³⁰, L. Zhong³, A. Zvyagin³⁷.

 1 Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³Center for High Energy Physics, Tsinghua University, Beijing, China

 $^{4}LAPP,$ Université de Savoie, CNRS/IN2P3, Annecy-Le-Vieux, France

 5 Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

 6 CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

 ${}^{7}LAL$, Université Paris-Sud, CNRS/IN2P3, Orsay, France

⁸LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

 9 Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

 10 Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

 11 Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

 12 School of Physics, University College Dublin, Dublin, Ireland

¹³Sezione INFN di Bari, Bari, Italy

¹⁴Sezione INFN di Bologna, Bologna, Italy

¹⁵Sezione INFN di Cagliari, Cagliari, Italy

¹⁶Sezione INFN di Ferrara, Ferrara, Italy

¹⁷Sezione INFN di Firenze, Firenze, Italy

¹⁸Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy

¹⁹Sezione INFN di Genova, Genova, Italy

²⁰Sezione INFN di Milano Bicocca, Milano, Italy

²¹Sezione INFN di Padova, Padova, Italy

²²Sezione INFN di Pisa, Pisa, Italy

²³Sezione INFN di Roma Tor Vergata, Roma, Italy

²⁴Sezione INFN di Roma La Sapienza, Roma, Italy

²⁵ Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland ^{26}AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland

²⁷National Center for Nuclear Research (NCBJ), Warsaw, Poland

²⁸Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania 29 Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia

³⁰Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia

³¹Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia

³²Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia

- ³³Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia
- ³⁴Institute for High Energy Physics (IHEP), Protvino, Russia
- ³⁵Universitat de Barcelona, Barcelona, Spain
- ³⁶Universidad de Santiago de Compostela, Santiago de Compostela, Spain

³⁷European Organization for Nuclear Research (CERN), Geneva, Switzerland

 38 Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

 39 Physik-Institut, Universität Zürich, Zürich, Switzerland

⁴⁰Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands

⁴¹ Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands

⁴²NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine

⁴³Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine

⁴⁴University of Birmingham, Birmingham, United Kingdom

⁴⁵H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom

 46 Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom

⁴⁷Department of Physics, University of Warwick, Coventry, United Kingdom

⁴⁸STFC Rutherford Appleton Laboratory, Didcot, United Kingdom

⁴⁹School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom

⁵⁰School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom

⁵¹Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom

⁵²Imperial College London, London, United Kingdom

⁵³ School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom

⁵⁴Department of Physics, University of Oxford, Oxford, United Kingdom

⁵⁵Massachusetts Institute of Technology, Cambridge, MA, United States

⁵⁶University of Cincinnati, Cincinnati, OH, United States

⁵⁷University of Maryland, College Park, MD, United States

⁵⁸Syracuse University, Syracuse, NY, United States

 59 Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ²

 60 Institut für Physik, Universität Rostock, Rostock, Germany, associated to 11

 61 Celal Bayar University, Manisa, Turkey, associated to 37

 $a_P.N.$ Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia

 b Università di Bari, Bari, Italy

 c Università di Bologna, Bologna, Italy

 d Università di Cagliari, Cagliari, Italy

 e Università di Ferrara, Ferrara, Italy

 f Università di Firenze, Firenze, Italy

 g Università di Urbino, Urbino, Italy

 h Università di Modena e Reggio Emilia, Modena, Italy

 i Università di Genova, Genova, Italy

 j Università di Milano Bicocca, Milano, Italy

 k Università di Roma Tor Vergata, Roma, Italy

 ι Università di Roma La Sapienza, Roma, Italy

 $^m Università$ della Basilicata, Potenza, Italy</sup>

 $ⁿ LIFAELS, La Salle, Universitat Ramon Llull, Barcelona. Snain$ </sup>

 \degree Hanoi University of Science, Hanoi, Viet Nam

p Institute of Physics and Technology, Moscow, Russia

^qUniversit`a di Padova, Padova, Italy

 r Università di Pisa, Pisa, Italy

^sScuola Normale Superiore, Pisa, Italy

The B_c^+ meson is the ground state of the $\bar{b}c$ system. As such it is unique as it is the only weakly decaying doubly heavy meson. All measurements of B_c^+ meson decays to date are decays where the constituent b quark decays weakly to a c quark $[1-8]$ $[1-8]$. The decay of the B_c^+ meson to another B meson, with the bottom quark acting as a spectator (see Fig. [1\)](#page-7-0), has not previously been observed. This will improve the understanding of theoretical predictions, and provide valuable information for the source of B_s^0 mesons at the LHC.

A wide range of predictions for the branching fraction $\mathcal{B}(B_c^+ \to B_s^0 \pi^+)$ exists, between 16.4 % and 2.5 %, based on e.g. QCD sum rules [\[9,](#page-14-1) [10\]](#page-14-2), or quark-potential models (see Refs. [\[11](#page-14-3)[–16\]](#page-14-4) and references therein). Experimental clarification is needed to shed light on the present theoretical status. Unlike most other B decays, the higher order corrections in the expansion of Heavy Quark Effective Theory within the framework of quantum chromodynamics (QCD) are relatively large. The expansion is described in powers of m_c/m_b rather than $\Lambda_{\rm QCD}/m_b$, due to the presence of two heavy quark constituents, where Λ_{QCD} is the QCD scale, and m_c (m_b) the charm (bottom) quark mass. In addition, the energy release in the decay is relatively small, leading to larger non-factorizable effects compared to decays with lighter daughter particles. Study of the decay $B_c^+ \to B_s^0 \pi^+$ allows these models to be tested. Knowledge of the production of B_s^0 mesons from B_c^+ decays is also useful for time-dependent analyses of B_s^0 decays, to understand any associated decay-time bias due to the incorrect estimate of the B_s^0 decay time if originating from a B_c^+ decay, or to take advantage of flavor tagging capabilities using the accompanying ("bachelor") pion.

The data used in this analysis were collected with the LHCb detector [\[17\]](#page-14-5) from pp The data used in this analysis were conected with the EriCb detector [17] from pp collisions at $\sqrt{s} = 7$ TeV and 8 TeV, corresponding to integrated luminosities of 1 fb⁻¹ and 2 fb^{-1} , respectively.

The decays $B_s^0 \to D_s^- \pi^+$ and $B_s^0 \to J/\psi \phi$ are used, with the subsequent decays $D_s^ \to K^+K^-\pi^-$, $J/\psi \to \mu^+\mu^-$ and $\phi \to K^+K^-$. The inclusion of charge conjugate modes is implied throughout. The event selection and fits to the B_s^0 invariant mass

Figure 1: Leading-order Feynman diagram of the decay $B_c^+ \rightarrow B_s^0 \pi^+$.

distributions follow previous LHCb analyses based on these B_s^0 decay modes [\[18,](#page-14-6) [19\]](#page-15-0). The two channels are analysed independently and the final results are combined. The strategy is to normalize the final number of $B_c^+ \to B_s^0 \pi^+$ decays to the number of B_s^0 decays, which gives a result for the $B_c^+ \to B_s^0 \pi^+$ branching fraction multiplied by the ratio of B_c^+ and B_s^0 production rates, $(\sigma(B_c^+)/\sigma(B_s^0)) \times \mathcal{B}(B_c^+ \to B_s^0 \pi^+)$. The B_c^+ signal region was not examined until the event selection was finalized. Since the ratio of production rates, $\sigma(B_c^+)/\sigma(B_s^0)$, may depend on the kinematics of the produced B meson, the result is quoted for B mesons produced in the pseudorapidity range $2 < \eta(B) < 5$, corresponding to the LHCb detector acceptance.

The LHCb detector is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, described in detail in Ref. [\[17\]](#page-14-5). The combined tracking system provides momentum measurement with relative uncertainty that varies from 0.4% at $5 \text{ GeV}/c$ to 0.6% at $100 \text{ GeV}/c$, and impact parameter resolution of $20 \mu m$ for tracks with high transverse momentum, p_T . The impact parameter (IP) is defined as the distance of closest approach between the track and a primary interaction. Charged hadrons are identified using two ring-imaging Cherenkov detectors. The charged pions from B_c^+ decays are selected with efficiency of 93 % while keeping the misidentification rate of kaons below 7 %. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers with a typical efficiency of 97% at $1-3\%$ pion to muon misidentification probability. The trigger [\[20\]](#page-15-1) consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction. The B_s^0 candidates with muons in the final state are required to pass the hardware trigger, which selects muons with a transverse momentum, $p_{\rm T} > 1.48 \,\text{GeV}/c$, whereas the B_s^0 candidates with only hadrons in the final state are selected by requiring a hadron in the calorimeter with $E_T > 3.6$ GeV/c.

Monte Carlo simulations, used to develop the B_c^+ candidate selection, are performed using BCVEGPY [\[21\]](#page-15-2), interfaced with PYTHIA 6.4 [\[22\]](#page-15-3) using a specific LHCb configura-tion [\[23\]](#page-15-4). Decays of hadronic particles are described by $EVTGEN$ [\[24\]](#page-15-5), in which final state radiation is generated using PHOTOS [\[25\]](#page-15-6). The interaction of the generated particles with the detector and its response are implemented using the GEANT4 toolkit [\[26\]](#page-15-7) as described in Ref. [\[27\]](#page-15-8).

The B_s^0 candidates are selected using the multi-variate analysis known as boosted decision tree (BDT) [\[28,](#page-15-9) [29\]](#page-15-10), to optimally discriminate between signal and background. In the training, simulated B_s^0 decays are used as signal, whereas candidates in the B_s^0 mass sideband in data are used as background. To avoid potential biases, only one sixth of the data is used in the training. It is verified that the distribution of the BDT discriminant is the same for the events used in the training, compared to those that were not. All events are used for the final result. The BDT training for the selection of $B_s^0 \to D_s^- \pi^+$ candidates uses only the upper sideband [5466, 5800] MeV/ c^2 , as the lower sideband contains a large amount of irreducible partially reconstructed B decays, while the training for $B_s^0 \to J/\psi \phi$ uses both lower sideband [5200, 5316] MeV/ c^2 and upper mass sideband [5416, 5550] MeV/ c^2 . The B_s^0 vertex quality (χ^2_{vtx}) , flight distance, momentum p and p_T are used to discriminate signal from background. For the $D_s^- \pi^+$ final state we use

Figure 2: Invariant mass distributions of (a) $B_s^0 \to D_s^- \pi^+$ and (b) $B_s^0 \to J/\psi \phi$ candidates. The different components are defined in the legend.

in addition the χ^2_{vtx} , flight distance, p and p_T of the D_s^- candidate and the p, p_T and χ^2_{IP} of the bachelor pion from the B_s^0 decay to suppress combinatorial background. The quantity χ_{IP}^2 is defined as the difference in χ^2 of a given primary vertex (PV) reconstructed with and without the considered track. The training for $J/\psi \phi$ candidates uses p, p_T , χ^2_{vtx} and χ_{IP}^2 of the J/ψ and ϕ candidates, and the p_T of the final state kaons and muons. In the selection of B_s^0 candidates from B_c^+ decays, variables that require the candidate to point to a primary vertex, such as the impact parameter of the B_s^0 candidate, are explicitly not included. The minimum value of the BDT discriminant is chosen by optimizing the B_s^0 included. The minimum value of the BDT discriminant is chosen by optimizing the B_s^s
signal significance $S/\sqrt{S+B}$, where S and B are the expected numbers of signal and combinatorial background events, respectively.

The total number of B_s^0 decays is obtained from extended unbinned maximum likelihood fits to the invariant mass distributions, using mass constraints for the J/ψ candidates [\[30\]](#page-15-11). and are shown in Fig. [2.](#page-9-0) The signal shapes are taken as double Crystal Ball functions [\[31\]](#page-15-12) with common peak value and with tails to either side of the peak, to account for final state radiation and detector resolution effects. The parameters that describe the tails are obtained from simulation and are fixed in the fits. The peak and width parameters of the signal are allowed to vary. The combinatorial backgrounds are modeled with exponential distributions. The $B_s^0 \to D_s^- \pi^+$ final state is contaminated by partially reconstructed B decays such as $B_s^0 \to D_s^{*-} \pi^+$ and $B_s^0 \to D_s^- \rho^+$ decays, where the soft photon or neutral pion is not reconstructed, and by decays where one of the final state particles is misidentified as a kaon, such as $B^0 \to D^-\pi^+$ or $\bar{A}_b^0 \to A_c^-\pi^+$ decays. The shapes of these backgrounds are fixed from simulation, following Ref. [\[18\]](#page-14-6). In total $103\,760 \pm 380$ $B_s^0 \to J/\psi \phi$ and 73700 $\pm 500 B_s^0 \to D_s^- \pi^+$ decays are found.

Selected B_s^0 candidates with masses consistent with the known B_s^0 mass are combined with tracks that satisfy loose pion identification requirements. Subsequently, B_c^+ candidates are selected with a second BDT algorithm. In the training of the second BDT, simulated candidates with masses consistent with the B_c^+ mass [\[32\]](#page-15-13) are used as signal, and candidates in the B_c^+ mass sideband region in data are used as background. For this, only the upper

mass sideband is used in the case of $B_s^0 \to D_s^- \pi^+$, while also the lower mass sideband is used in the case of $B_s^0 \to J/\psi \phi$, to further suppress the larger combinatorial background at smaller values of the mass. Only one sixth of the total data set is used in the training. The second BDT uses the following variables: the B_c^+ candidate p_T , decay time, χ^2_{vtx} , χ^2_{IP} and the B_c^+ pointing angle, *i.e.* the angle between the B_c^+ candidate momentum vector and the line joining the associated PV and the B_c^+ decay vertex. The B_s^0 polar angle (the angle between B_s^0 flight direction and the beam axis), decay time, decay length and pointing angle are also used. The p and p_T of the bachelor pion from the B_c^+ decay are the most discriminating observables in the second BDT. Differences between the analyses of the $D_s^- \pi^+$ and $J/\psi \phi$ final states are: the use of $\chi^2_{\rm IP}$ of the B_s^0 candidate and bachelor pion (from the B_c^+ decay), and B_s^0 and B_c^+ momentum for the former; the use of the B_c^+ and B_s^0 decay-length uncertainties for the latter. The optimal selections are defined by maximizing figures of merit for a target level of significance of three standard deviations, maximizing figures of merit for a target level of significance of three standard deviations $\epsilon/(3/2 + \sqrt{B})$ [\[33\]](#page-16-0), where ϵ is the signal efficiency for a given BDT criterion. The figure of merit displays a plateau, and the chosen value is at the lower end to allow to better constrain the shape of the combinatiorial background. The chosen selection is very close constrain the snape of the combinational background. The chosen selection is very close
to the optimal point for a target level of 5σ and for the expected significance $S/\sqrt{S+B}$. The trigger for $B_s^0 \to D_s^- \pi^+$ decays preferentially selects candidates with high p_T with respect to the trigger for $B_s^0 \to J/\psi \phi$ decays, which results in higher efficiency for the second BDT requirement for the $B_s^0 \to D_s^- \pi^+$ final state. The B_c^+ and B_s^0 candidates are required to be produced in the pseudorapidity range $2 < \eta(B) < 5$.

The invariant mass distributions for the $B_c^+ \to B_s^0 \pi^+$ candidates are shown in Fig. [3,](#page-10-0) together with the resulting fits. The decay $B_c^+ \to B_s^0 \pi^+$ has a Q-value of 770 MeV/ c^2 (with $Q \equiv m_{B_c^+} - m_{B_s^0} - m_{\pi^+}$), which results in a resolution of about 6 MeV/ c^2 when a B_s^0 mass constraint is applied. The signal shape is modeled as a double Crystal Ball function, with its parameters obtained from simulated events. The larger number of B_c^+ candidates in the $B_s^0 \to D_s^- \pi^+$ channel allows variation of the peak position and the width in the

Figure 3: B_c^+ mass fits for the combined 2011 and 2012 data sets for (a) $B_c^+ \to B_s^0 (\to D_s^- \pi^+) \pi^+$ and (b) $B_c^+ \to B_s^0 (\to J/\psi \phi) \pi^+$ candidates. The different components are indicated in the legends.

fit. The combinatorial background is primarily due to signal B_s^0 decays combined with a random pion from the primary vertex, and is modeled with an exponential function. Backgrounds due to $B_c^+ \to B_s^* \pi^+$ and $B_c^+ \to B_s^0 \rho^+$ decays, where the photon or neutral pion are not reconstructed, are simulated, and their shapes are modeled with Gaussian distributions, with parameters fixed in the fit, and yields allowed to vary. Statistical signal significances of 7.7 σ for $B_c^+ \to B_s^0 (\to D_s^- \pi^+) \pi^+$ and 6.1 σ for $B_c^+ \to B_s^0 (\to J/\psi \phi) \pi^+$ decays are obtained from the likelihood ratio of fits with and without the probability density function for the signal shape, $\sqrt{-2\ln(\mathcal{L}_{B}/\mathcal{L}_{S+B})}$, with 64 ± 10 and 35 ± 8 signal decays, respectively.

In Fig. [3a](#page-10-0), the structure around $6225 \text{ MeV}/c^2$ is consistent with originating from $B_c^+ \to B_s^* \pi^+$ decays. However, this contribution is not significant.

To obtain the value for the $B_c^+ \to B_s^0 \pi^+$ branching fraction, multiplied by the ratio of B_c^+ and B_s^0 production rates, the relative detection efficiency of B_s^0 decays compared to $B_c^+ \to B_s^0 \pi^+$ decays is determined from simulation. Requiring the bachelor pion to be inside the LHCb acceptance reduces the $B_c^+ \to B_s^0 \pi^+$ yield by about 19% with respect to the B_s^0 yield. The most significant reduction in the number of selected B_c^+ candidates comes from suppressing B_s^0 combinations with a random pion from the primary interaction, by means of the second BDT selection. The total relative detection efficiency of $B_c^+ \to B_s^0 \pi^+$ decays with respect to B_s^0 decays is estimated to be 15.2% for the $B_s^0 \to J/\psi \phi$ decay and 33.9% for the $B_s^0 \to D_s^- \pi^+$ final state. This difference in B_c^+ selection efficiencies is a consequence of the difference in B_s^0 trigger and selection requirements.

The sources of systematic uncertainty for the efficiency-corrected ratio of B_c^+ and B_s^0 yields are listed in Table [1.](#page-11-0) The uncertainty on the B_s^0 yield in the $D_s^- \pi^+$ analysis is determined by varying the parameters that describe the tails of the signal mass distribution, and by reducing the exponent of the combinatorial background by a factor two. The

Table 1: Contributions of the various sources of (relative) systematic uncertainty on the efficiencycorrected ratio of event yields. The total systematic uncertainty is the quadratic sum of the individual contributions. The number of $B_c^+ \to B_s^0 (\to D_s^- \pi^+) \pi^+$ candidates is large enough that the peak position and width are freely varied in the fit, and hence the corresponding uncertainty is contained in the statistical uncertainty of the signal yield.

Source	$D_{s}^{-}\pi^{+}$	
$B_{\rm s}^0$ fit model	3.0	1.2
B_c^+ mean mass	.	2.0
B_{c}^{+} mass resolution	.	5.2
B_c^+ signal model	1.5	1.7
Combinatorial background model	1.8	0.3
Partially reconstructed bkgd.	1.8	1.7
Data-simulation difference	3.7	3.7
B_c^+ lifetime	$+6.8$ -3.5	7.4
Total	$+8.9$ -6.7	10.4

uncertainty on the $B_s^0 \to J/\psi \phi$ yield is obtained by comparing the fitted yield in simulated pseudo-experiments to the yield that was used as input to those experiments.

The uncertainty on the B_c^+ yield is quantified by varying the peak position and width in the fit to $B_c^+ \to B_s^0 (\to J/\psi \phi) \pi^+$ candidates. The signal model is validated using simulated pseudo-experiments in the $J/\psi \phi$ analysis, whereas the tail parameters are varied by $\pm 10\%$ in the $D_s^- \pi^+$ analysis. In addition, the combinatorial background shape is changed to a straight line, and the difference in the signal yield is taken as the associated systematic uncertainty. The effect of partially reconstructed $B_c^+ \rightarrow B_s^0 \rho^+$ decays is estimated by excluding candidates with mass less than $6150 \,\text{MeV}/c^2$ from the fit. The significance of the $B_c^+ \to B_s^0 \pi^+$ signal is reduced to 7.5 σ for $B_c^+ \to B_s^0 (\to D_s^- \pi^+) \pi^+$ and 5.5 σ for $B_c^+ \to B_s^0 (\to J/\psi \phi) \pi^+$ when the systematic uncertainties on the fit to the B_c^+ mass distribution are taken into account.

The relative detection efficiency of B_c^+ and B_s^0 events is determined from simulated events. The correspondence between data and simulation is quantified by varying the criterion on the BDT value, and by comparing the observed B_s^0 yield to the expected yield based on the change in efficiency as determined from simulation. The largest contribution is due to the 10% uncertainty on the B_c^+ lifetime [\[32\]](#page-15-13), which was recently improved by the CDF collaboration [\[34\]](#page-16-1). The change in selection efficiency when varying the B_c^+ lifetime by $\pm 10\%$ is assigned as systematic uncertainty. A longer (shorter) B_c^+ lifetime corresponds to a larger (smaller) efficiency and therefore a smaller (larger) ratio. As a cross-check, the effect of the choice of different sets of BDT input variables is investigated and the result is found to be stable.

The contribution from Cabibbo suppressed $B_c^+ \to B_s^0 K^+$ decays, the uncertainty on the efficiency of reconstructing the extra pion, and the uncertainty on the efficiency of the particle identification requirement on the bachelor pion all give small contributions $(< 1.0\%)$ to the total systematic uncertainty, and are not itemized in the summary in Table [1.](#page-11-0)

The B_s^0 and B_c^+ yields are corrected for the relative detection efficiencies, to obtain the efficiency-corrected ratios of $B_c^+ \to B_s^0 \pi^+$ over B_s^0 yields, $(2.54 \pm 0.40 \,\text{(stat)}_{-0.17}^{+0.23} \,\text{(syst)}) \times$ 10^{-3} and $(2.20 \pm 0.49 \text{ (stat)} \pm 0.23 \text{ (syst)}) \times 10^{-3}$ for the D_s^- π⁺ and $J/\psi \phi$ final states, respectively. The small fraction of B_s^0 candidates originating from B_c^+ decays is neglected. The uncertainty due to the uncertainty on the B_c^+ lifetime is correlated between the two measurements, and is accounted for in the combined result of the ratio of production rates multiplied with the branching fraction

$$
\frac{\sigma(B_c^+)}{\sigma(B_s^0)} \times \mathcal{B}(B_c^+ \to B_s^0 \pi^+) = (2.37 \pm 0.31 \, (\text{stat}) \pm 0.11 \, (\text{syst}) \, {}^{+0.17}_{-0.13} \, (\tau_{B_c^+}) \times 10^{-3},
$$

where the first uncertainty is statistical, the second is systematic and the third is due to the uncertainty on the B_c^+ lifetime. Since $\sigma(B_c^+)/\sigma(B_s^0)$ may depend on the kinematics of the produced B meson, the data are divided according to center-of-mass energy leading to $(1.27 \pm 0.42 \text{ (stat)} \pm 0.05 \text{ (syst)} \frac{+0.09}{-0.07} (\tau_{B_c^+}) \times 10^{-3} \text{ and}$ $(2.92 \pm 0.40 \text{ (stat)} \pm 0.12 \text{ (syst)} \pm 0.21 \text{ (r_s)} \times 10^{-3} \text{ for } \sqrt{s} = 7 \text{ and } 8 \text{ TeV } pp \text{ collisions},$ respectively. The lower value for the result of the 7 TeV data is attributed to a downward statistical fluctuation of the $B_c^+ \to B_s^0 (\to J/\psi \phi) \pi^+$ yield in the 2011 data set, with a p-value of 1.5% .

Assuming a value for $\mathcal{B}(B_c^+ \to J/\psi \pi^+)$ around 0.15 % [\[11\]](#page-14-3), combined with the results $(\sigma(B_c^+)/\sigma(B^+)) \times \mathcal{B}(B_c^+ \to J/\psi \pi^+)/\mathcal{B}(B^+ \to J/\psi K^+) = (0.68 \pm 0.10 \pm 0.03 \pm 0.05) \,\%$ [\[4\]](#page-14-7), and measurements of f_s/f_d [\[18\]](#page-14-6) and $\mathcal{B}(B^+ \to J/\psi K^+)$ [\[32\]](#page-15-13), results in a ratio of production rates of B_c^+ mesons over B_s^0 mesons of about 0.02. This leads to a branching fraction for $B_c^+ \to B_s^0 \pi^+$ of about 10%. Although precise quantification requires improved understanding of $\sigma(B_c^+)$ and $\mathcal{B}(B_c^+ \to J/\psi \pi^+)$, even taking the lower estimates for $\mathcal{B}(B_c^+\to J/\psi\,\pi^+)$ that are found in the literature [\[11\]](#page-14-3), leads to a value of $\mathcal{B}(B_c^+\to B_s^0\pi^+)$ which is the largest exclusive branching fraction of any known weak B meson decay.

In summary, the first observation of a weak decay of a B meson to another B meson is reported. This measurement will help to better understand flavor tagging and the decay time resolution in time-dependent B_s^0 analyses, and in addition will constrain models that predict branching fractions of B_c^+ decays.

Acknowledgements

We wish to thank A.K. Likhoded for useful discussions. We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 and Region Auvergne (France); BMBF, DFG, HGF and MPG (Germany); SFI (Ireland); INFN (Italy); FOM and NWO (The Netherlands); SCSR (Poland); MEN/IFA (Romania); MinES, Rosatom, RFBR and NRC "Kurchatov Institute" (Russia); MinECo, XuntaGal and GENCAT (Spain); SNSF and SER (Switzerland); NAS Ukraine (Ukraine); STFC (United Kingdom); NSF (USA). We also acknowledge the support received from the ERC under FP7. The Tier1 computing centres are supported by IN2P3 (France), KIT and BMBF (Germany), INFN (Italy), NWO and SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom). We are thankful for the computing resources put at our disposal by Yandex LLC (Russia), as well as to the communities behind the multiple open source software packages that we depend on.

References

- [1] CDF collaboration, F. Abe et al., Observation of the B_c meson in $p\bar{p}$ collisions at \sqrt{s} = 1.8 TeV, [Phys. Rev. Lett.](http://dx.doi.org/10.1103/PhysRevLett.81.2432) 81 (1998) 2432, [arXiv:hep-ex/9805034](http://arxiv.org/abs/hep-ex/9805034); CDF \sqrt{s} = 1.6 TeV, Thys. Nev. Lett. **61** (1998) 2452, **at ATV** hep-ext 9605054, CDF collaboration, F. Abe *et al., Observation of B_c mesons in pp̃ collisions at* \sqrt{s} = 1.8 TeV, Phys. Rev. D58 [\(1998\) 112004,](http://dx.doi.org/10.1103/PhysRevD.58.112004) [arXiv:hep-ex/9804014](http://arxiv.org/abs/hep-ex/9804014).
- [2] CDF collaboration, T. Aaltonen *et al.*, Observation of the decay $B_c^{\pm} \rightarrow J/\psi \pi^{\pm}$ and measurement of the B_c^{\pm} mass, [Phys. Rev. Lett.](http://dx.doi.org/10.1103/PhysRevLett.100.182002) **100** (2008) 182002, arXiv: 0712.1506.
- [3] D0 collaboration, V. M. Abazov et al., Observation of the B_c meson in the exclusive decay $B_c \to J/\psi \pi$, [Phys. Rev. Lett.](http://dx.doi.org/10.1103/PhysRevLett.101.012001) 101 (2008) 012001, [arXiv:0802.4258](http://arxiv.org/abs/0802.4258).
- [4] LHCb collaboration, R. Aaij et al., Measurements of B_c^+ production and mass with the $B_c^+ \to J/\psi \pi^+$ decay, [Phys. Rev. Lett.](http://dx.doi.org/10.1103/PhysRevLett.109.232001) **109** (2012) 232001, [arXiv:1209.5634](http://arxiv.org/abs/1209.5634).
- [5] LHCb collaboration, R. Aaij et al., First observation of the decay $B_c^+ \to J/\psi \pi^+ \pi^- \pi^+$, [Phys. Rev. Lett.](http://dx.doi.org/10.1103/PhysRevLett.108.251802) 108 (2012) 251802, [arXiv:1204.0079](http://arxiv.org/abs/1204.0079).
- [6] LHCb collaboration, R. Aaij et al., Observation of the decay $B_c^+ \to \psi(2S)\pi^+$, [Phys.](http://dx.doi.org/10.1103/PhysRevD.87.071103) Rev. D87 [\(2013\) 071103\(R\),](http://dx.doi.org/10.1103/PhysRevD.87.071103) [arXiv:1303.1737](http://arxiv.org/abs/1303.1737).
- [7] LHCb collaboration, R. Aaij et al., Observation of $B_c^+ \to J/\psi D_s^+$ and $B_c^+ \to J/\psi D_s^{*+}$ decays, Phys. Rev. D87 [\(2013\) 112012,](http://dx.doi.org/10.1103/PhysRevD.87.112012) [arXiv:1304.4530](http://arxiv.org/abs/1304.4530).
- [8] LHCb collaboration, R. Aaij et al., First observation of the decay $B_c^+ \to J/\psi K^+$, JHEP 09 [\(2013\) 075,](http://dx.doi.org/10.1007/JHEP09(2013)075) [arXiv:1306.6723](http://arxiv.org/abs/1306.6723).
- [9] V. Kiselev, A. Kovalsky, and A. Likhoded, B_c decays and lifetime in QCD sum rules, [Nucl. Phys.](http://dx.doi.org/10.1016/S0550-3213(00)00386-2) B585 (2000) 353, [arXiv:hep-ph/0002127](http://arxiv.org/abs/hep-ph/0002127).
- [10] I. Gouz et al., Prospects for the B_c studies at LHCb, [Phys. Atom. Nucl.](http://dx.doi.org/10.1134/1.1788046) 67 (2004) [1559,](http://dx.doi.org/10.1134/1.1788046) [arXiv:hep-ph/0211432](http://arxiv.org/abs/hep-ph/0211432).
- [11] M. A. Ivanov, J. G. Korner, and P. Santorelli, Exclusive semileptonic and nonleptonic decays of the B_c meson, Phys. Rev. D73 [\(2006\) 054024,](http://dx.doi.org/10.1103/PhysRevD.73.054024) [arXiv:hep-ph/0602050](http://arxiv.org/abs/hep-ph/0602050).
- [12] A. Y. Anisimov, I. Narodetsky, C. Semay, and B. Silvestre-Brac, The B_c meson lifetime in the light front constituent quark model, [Phys. Lett.](http://dx.doi.org/10.1016/S0370-2693(99)00273-7) B452 (1999) 129, [arXiv:hep-ph/9812514](http://arxiv.org/abs/hep-ph/9812514).
- [13] P. Colangelo and F. De Fazio, Using heavy quark spin symmetry in semileptonic B_c decays, Phys. Rev. $D61$ [\(2000\) 034012,](http://dx.doi.org/10.1103/PhysRevD.61.034012) $arXiv:hep-ph/9909423$.
- [14] D. Ebert, R. Faustov, and V. Galkin, Weak decays of the B_c meson to B_s and B mesons in the relativistic quark model, [Eur. Phys. J.](http://dx.doi.org/10.1140/epjc/s2003-01347-5) $C32$ (2003) 29, $arXiv:hep-ph/0308149$.
- [15] R. Dhir, N. Sharma, and R. Verma, Flavor dependence of B_c meson form factors and $B_c \to PP \ decays, J. Phys. G35 (2008) 085002.$ $B_c \to PP \ decays, J. Phys. G35 (2008) 085002.$
- [16] S. Naimuddin *et al., Nonleptonic two-body* B_c -meson decays, [Phys. Rev.](http://dx.doi.org/10.1103/PhysRevD.86.094028) **D86** (2012) [094028.](http://dx.doi.org/10.1103/PhysRevD.86.094028)
- [17] LHCb collaboration, A. A. Alves Jr. et al., The LHCb detector at the LHC, [JINST](http://dx.doi.org/10.1088/1748-0221/3/08/S08005) 3 [\(2008\) S08005.](http://dx.doi.org/10.1088/1748-0221/3/08/S08005)
- [18] LHCb collaboration, R. Aaij et al., Measurement of the fragmentation fraction ratio f_s/f_d and its dependence on B meson kinematics, JHEP 04 [\(2013\) 1,](http://dx.doi.org/10.1007/JHEP04(2013)001) [arXiv:1301.5286](http://arxiv.org/abs/1301.5286).
- [19] LHCb collaboration, R. Aaij et al., Measurement of CP-violation and the B_s^0 -meson decay width difference with $B_s^0 \to J/\psi K^+ K^-$ and $B_s^0 \to J/\psi \pi^+ \pi^-$ decays, [Phys. Rev.](http://dx.doi.org/10.1103/PhysRevD.87.112010) D87 [\(2013\) 112010,](http://dx.doi.org/10.1103/PhysRevD.87.112010) [arXiv:1304.2600](http://arxiv.org/abs/1304.2600).
- [20] R. Aaij et al., The LHCb trigger and its performance in 2011, JINST 8 [\(2013\) P04022,](http://dx.doi.org/10.1088/1748-0221/8/04/P04022) [arXiv:1211.3055](http://arxiv.org/abs/1211.3055).
- [21] C.-H. Chang, J.-X. Wang, and X.-G. Wu, BCVEGPY2.0: An upgraded version of the generator BCVEGPY with the addition of hadroproduction of the P-wave B_c^+ states, [Comput. Phys. Commun.](http://dx.doi.org/10.1016/j.cpc.2005.09.008) 174 (2006) 241, [arXiv:hep-ph/0504017](http://arxiv.org/abs/hep-ph/0504017).
- [22] T. Sjöstrand, S. Mrenna, and P. Skands, $PYTHIA$ 6.4 physics and manual, [JHEP](http://dx.doi.org/10.1088/1126-6708/2006/05/026) 05 [\(2006\) 026,](http://dx.doi.org/10.1088/1126-6708/2006/05/026) [arXiv:hep-ph/0603175](http://arxiv.org/abs/hep-ph/0603175).
- [23] I. Belyaev et al., Handling of the generation of primary events in Gauss, the LHCb simulation framework, [Nuclear Science Symposium Conference Record \(NSS/MIC\)](http://dx.doi.org/10.1109/NSSMIC.2010.5873949) IEEE [\(2010\) 1155.](http://dx.doi.org/10.1109/NSSMIC.2010.5873949)
- [24] D. J. Lange, The EvtGen particle decay simulation package, [Nucl. Instrum. Meth.](http://dx.doi.org/10.1016/S0168-9002(01)00089-4) A462 [\(2001\) 152.](http://dx.doi.org/10.1016/S0168-9002(01)00089-4)
- [25] P. Golonka and Z. Was, PHOTOS Monte Carlo: a precision tool for QED corrections in Z and W decays, [Eur. Phys. J.](http://dx.doi.org/10.1140/epjc/s2005-02396-4) $C45$ (2006) 97, [arXiv:hep-ph/0506026](http://arxiv.org/abs/hep-ph/0506026).
- [26] Geant4 collaboration, J. Allison et al., Geant4 developments and applications, [IEEE](http://dx.doi.org/10.1109/TNS.2006.869826) [Trans. Nucl. Sci.](http://dx.doi.org/10.1109/TNS.2006.869826) 53 (2006) 270; Geant4 collaboration, S. Agostinelli et al., Geant4: a simulation toolkit, [Nucl. Instrum. Meth.](http://dx.doi.org/10.1016/S0168-9002(03)01368-8) A506 (2003) 250.
- [27] M. Clemencic et al., The LHCb simulation application, GAUSS: design, evolution and experience, [J. Phys. Conf. Ser.](http://dx.doi.org/10.1088/1742-6596/331/3/032023) **331** (2011) 032023.
- [28] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, Classification and regression trees, Wadsworth international group, Belmont, California, USA, 1984.
- [29] R. E. Schapire and Y. Freund, A decision-theoretic generalization of on-line learning and an application to boosting, [Jour. Comp. and Syst. Sc.](http://dx.doi.org/10.1006/jcss.1997.1504) 55 (1997) 119.
- [30] W. D. Hulsbergen, Decay chain fitting with a Kalman filter, [Nucl. Instrum. Meth.](http://dx.doi.org/10.1016/j.nima.2005.06.078) A552 [\(2005\) 566,](http://dx.doi.org/10.1016/j.nima.2005.06.078) [arXiv:physics/0503191](http://arxiv.org/abs/physics/0503191).
- [31] T. Skwarnicki, A study of the radiative cascade transitions between the Upsilon-prime and Upsilon resonances, PhD thesis, Institute of Nuclear Physics, Krakow, 1986, [DESY-F31-86-02.](http://inspirehep.net/record/230779/files/230779.pdf)
- [32] Particle Data Group, J. Beringer *et al., [Review of particle physics](http://pdg.lbl.gov/)*, [Phys. Rev.](http://dx.doi.org/10.1103/PhysRevD.86.010001) **D86** [\(2012\) 010001.](http://dx.doi.org/10.1103/PhysRevD.86.010001)
- [33] G. Punzi, Sensitivity of searches for new signals and its optimization, in Statistical Problems in Particle Physics, Astrophysics, and Cosmology (L. Lyons, R. Mount, and R. Reitmeyer, eds.), p. 79, 2003. [arXiv:physics/0308063](http://arxiv.org/abs/physics/0308063).
- [34] CDF collaboration, T. Aaltonen et al., Measurement of the B_c^- meson lifetime in the decay B_c^- → J/ψ π^- , Phys. Rev. **D87** [\(2013\) 011101,](http://dx.doi.org/10.1103/PhysRevD.87.011101) [arXiv:1210.2366](http://arxiv.org/abs/1210.2366).