Characterizing some rings of finite order

Jutirekha Dutta, Dhiren K. Basnet and Rajat K. Nath[∗]

*Department of Mathematical Sciences, Tezpur University, Napaam-784028, Sonitpur, Assam, India. Emails: jutirekhadutta@yahoo.com, dbasnet@tezu.ernet.in and rajatkantinath@yahoo.com**

Abstract

In this paper, we compute the number of distinct centralizers of some classes of finite rings. We then characterize all finite rings with n distinct centralizers for any positive integer $n \leq 5$. Further we give some connections between the number of distinct centralizers of a finite ring and its commutativity degree.

Key words: finite ring, centralizer, commutativity degree. *2010 Mathematics Subject Classification:* 16U70.

1 Introduction

Finite abelian groups have been completely characterized up to isomorphism for a long time but finite rings have yet to be characterized. The problem of characterizing finite rings up to isomorphism has received considerable attention in recent years (see [\[2,](#page-12-0) [8,](#page-12-1) [9,](#page-12-2) [11,](#page-12-3) [12\]](#page-12-4)) starting from the works of Eldridge [\[10\]](#page-12-5) and Raghavendran [\[15\]](#page-12-6). In this paper we characterize finite rings in terms of their number of distinct centralizers. Given a ring R and

[∗]Corresponding author

an element $r \in R$, the subrings $C(r) = \{s \in R : rs = sr\}$ and $Z(R) =$ $\{s \in R : rs = sr \text{ for all } r \in R\}$ are known as centralizer of r in R and *center* of R respectively. We write $Cent(R)$ to denote the set of all centralizers in R. Firstly we compute the order of $Cent(R)$ for some classes of finite rings R. Motivated by the works of Belcastro and Sherman $\lbrack 3 \rbrack$ and Ashrafi [\[1\]](#page-11-0), we define *n*-centralizer ring for any positive integer *n*. A ring R is said to be *n*-centralizer ring if $|\text{Cent}(R)| = n$, for any positive integer n. We then characterize n-centralizer finite rings for all $n \leq 5$, adapting similar techniques that are used by Belcastro and Sherman [\[3\]](#page-12-7) in order to characterize *n*-centralizer finite groups for $n \leq 5$.

Further, we conclude the paper by noting some interesting connections between $d(R)$ and $|\text{Cent}(R)|$. Note that for any finite ring R, the ratio $d(R)$ = 1 $\frac{1}{|R|^2}\sum$ r∈R $|C(r)|$ is the probability that a randomly chosen pair of elements of R commute. This ratio is known as commutativity degree of finite ring R and it was introduced by MacHale [\[13\]](#page-12-8) in the year 1975. Some characterizations of finite rings in terms of commutativity degree can be found in [\[13,](#page-12-8) [5,](#page-12-9) [6\]](#page-12-10).

Throughout the paper R denotes a finite ring. For any subring S of R, R/S denotes the additive quotient group and $|R:S|$ denotes the index of the additive subgroup S in the additive group R . Note that the isomorphisms considered are the additive group isomorphisms. Also for any two non-empty subsets A and B of a ring R, we write $A + B = \{a + b : a \in A, b \in B\}$. We shall use the fact that for any non-commutative ring R , the additive group R $\frac{R}{Z(R)}$ is not a cyclic group (see [\[13,](#page-12-8) Lemma 1]).

2 Some computations of $|\text{Cent}(R)|$

In this section, we compute $|\text{Cent}(R)|$ for some classes of finite rings. However, first we prove some results which are useful for subsequent results as well as for the next sections.

Proposition 2.1. R is a commutative ring if and only if R is a 1-centralizer ring.

Proof. The proposition follows from the fact that a ring R is commutative if and only if $C(r) = R$ for each $r \in R$. □

Proposition 2.2. Let R , S be two rings, then

$$
Cent(R \times S) = Cent(R) \times Cent(S).
$$

Proof. It can be easily seen that $C((r, s)) = C(r) \times C(s)$ for any $r \in R$ and $s \in S$. This proves the proposition. \Box

The following lemmas play an important role in finding lower bound of $| Cent(R)|$ for any non-commutative ring R.

Lemma 2.3. Let R be a ring. Then $Z(R)$ is the intersection of all centralizers in R.

Proof. It is clear that $Z(R) \subseteq \bigcap_{r \in R} C(r)$. Now, for any $s \in \bigcap_{r \in R} C(r)$ we have $rs = sr$ for all $r \in R$. Therefore $s \in Z(R)$. Hence the lemma follows.

Lemma 2.4. If R is a ring, then R is the union of centralizers of all noncentral elements of R.

Proof. It is clear that $\bigcup_{r \in R-Z(R)} C(r) \subseteq R$. Again, for any $s \in Z(R)$, we have by Lemma [2.3,](#page-2-0) $s \in C(r)$ for all $r \in R$. So $s \in \bigcup_{r \in R-Z(R)} C(r)$. Also for any $s \in R - Z(R)$, we have $s \in C(s)$ and so $s \in \bigcup_{r \in R - Z(R)} C(r)$. Hence the lemma follows. \Box

Lemma 2.5. A ring R cannot be written as a union of two of its proper subrings.

Proof. The lemma follows from the well-known fact that a group can not be written as a union of two of its proper subgroups. \Box

Theorem 2.6. For any non-commutative ring R, $|\text{Cent}(R)| \geq 4$.

Proof. Since R is non-commutative, so $|\text{Cent}(R)| \geq 2$. If $|\text{Cent}(R)| = 2$, then, by Lemma [2.4,](#page-2-1) R is equal to a proper subset of itself, which is not possible. Also by Lemma [2.5,](#page-2-2) $|\text{Cent}(R)| \neq 3$. Hence the theorem follows. \Box

Note that the ring $R = \left\{ \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \right.$ $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$ $\begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}$ $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, where $0, 1 \in \mathbb{Z}_2$, has 4 distinct centralizers. So the above result is the best one possible.

At this point, the following question, similar to the question posed by Belcastro and Sherman [\[3,](#page-12-7) p. 371], arises naturally.

Question 2.7. Does there exist an *n*-centralizer ring for any positive integer $n \neq 2, 3$? Can we characterize an n-centralizer ring?

The following results show the existence of *n*-centralizer rings for some values of n.

Proposition 2.8. There exists a $(p+2)$ -centralizer ring for any prime p.

Proof. We consider the ring $R = \left\{ \begin{bmatrix} a & b \\ 0 & 0 \end{bmatrix} : a, b \in \mathbb{Z}_p \right\}$. For any element $\begin{bmatrix} x & y \\ 0 & 0 \end{bmatrix}$ of $C\left(\begin{bmatrix} a & b \\ 0 & 0 \end{bmatrix}\right)$ we have $xb - ay = 0$. Clearly, $C\left(\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}\right) = R$. Using simple calculations, we have for any $a\neq 0$ and $l\in \mathbb{Z}$ $C\left(\begin{bmatrix} a & 0 \ 0 & 0 \end{bmatrix}\right) = \left\{\begin{bmatrix} x & 0 \ 0 & 0 \end{bmatrix} \, : \, x \in \mathbb{Z}_p\right\}$ and $C\left(\begin{bmatrix}la & a \\ 0 & 0\end{bmatrix}\right) = \left\{\begin{bmatrix}lx & x \\ 0 & 0\end{bmatrix} \ : \ x \in \mathbb{Z}_p\right\}$. Hence $|\text{Cent}(R)| = p+2$. \Box

The above proposition is a particular case of the following theorem.

Theorem 2.9. Let R be a non-commutative ring of order p^2 , where p is a prime. Then $|\text{Cent}(R)| = p+2$.

Proof. For any $x \in R - Z(R)$, we consider $C(x)$. As $C(x)$ is an additive subgroup of R we have $|C(x)| = 1, p$ or p^2 . Clearly, $|C(x)| \neq 1, p^2$, as $x, 0_R \in C(x)$ and R is non-commutative, where 0_R is the additive identity in R. Hence $C(x)$ is additive cyclic group of order p and so $Z(R) = \{0_R\}.$

Let $x, y \in R - Z(R)$. If there exists an element $t \neq 0_R$ $\in C(x) \cap$ $C(y)$ then $C(x) = C(y)$, as $C(x)$, $C(y)$ are additive cyclic groups of order p. Thus for any $x, y \in R - Z(R)$ we have either $C(x) \cap C(y) = \{0_R\}$ or $C(x) = C(y)$. Therefore the number of centralizers of non-central elements p^2-1 $|R| - |Z(R)|$ is = $= p + 1.$ Hence $|\text{Cent}(R)| = p + 2.$ \Box $p-1$ $p-1$

Theorem 2.10. Let p be a prime number and R be a non-commutative ring of order p^3 with unity. Then $|\text{Cent}(R)| = p+2$.

Proof. Let x be an arbitrary element of $R - Z(R)$. Then $C(x)$ is an additive subgroup of R and so $|C(x)| = 1, p, p^2$ or p^3 . Here $|C(x)| \neq 1, p^3$ as $x, 0_R \in$ $C(x)$, where 0_R is the additive identity in R and R is non-commutative. If $|C(x)| = p$ then $|Z(R)| = 1$, which is not possible as $0_R, 1_R \in Z(R)$. So $|C(x)| = p^2$ and this gives $|Z(R)| = p$.

Now, we suppose that $y \in R - Z(R)$ and $y \in C(x)$. Let $z \in C(x)$ be an arbitrary element. We know that $Z(R) \subset Z(C(x))$ and so $|Z(C(x))| > 1$, therefore by Lemma 3 of [\[14\]](#page-12-11), $C(x)$ is commutative. Therefore $z \in C(y)$, as $y \in C(x)$. So $C(x) \subseteq C(y)$. Also $|C(x)| = |C(y)|$. Hence, $C(x) = C(y)$; and if $y \notin C(x)$ then $C(x) \cap C(y) = Z(R)$. Therefore the number of centralizers p^3-p $|R| - |Z(R)|$ of non-central elements of R is = $\frac{p}{p^2 - p} = p + 1$. Thus $|C(x)| - |Z(R)|$ $| \text{Cent}(R) | = p + 2.$ \Box

 $\int [a \; b]$ As an application of the above theorem, it follows that the ring $R =$ $0 \quad c$ $\Big\}$ | a, b, $c \in \mathbb{Z}_p$ having order p^3 is a $(p+2)$ -centralizer ring. The following theorem, which is generalization of Theorem [2.9,](#page-3-0) gives another class of $(p+2)$ -centralizer rings.

Theorem 2.11. Let R be a ring and $\frac{R}{Z(R)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$, where p is a prime. $Then |\text{Cent}(R)| = p+2.$

Proof. We write $Z := Z(R)$. Since $R/Z \cong \mathbb{Z}_p \times \mathbb{Z}_p$ we have

$$
\frac{R}{Z} = \langle Z + a, Z + b \; : \; p(Z + a) = p(Z + b) = Z; a, b \in R \rangle.
$$

If S/Z is additive non-trivial subgroup of R/Z then $|S/Z| = p$. Therefore any additive proper subgroup of R properly containing Z has p disjoint right cosets. Hence the proper additive subgroups of R properly containing Z are

$$
S_m = Z \cup (Z + (a + mb)) \cup (Z + 2(a + mb)) \cup \cdots \cup (Z + (p - 1)(a + mb)),
$$

where $1 \le m \le (p - 1)$,

$$
S_p = Z \cup (Z + a) \cup (Z + 2a) \cup \cdots \cup (Z + (p - 1)a)
$$
 and

$$
S_{p+1} = Z \cup (Z + b) \cup (Z + 2b) \cup \cdots \cup (Z + (p - 1)b).
$$

Now for any $x \in R - Z$, we have $Z + x$ is equal to $Z + k$ for some $k \in$ ${m a, mb, a + mb, 2(a + mb), \ldots, (p-1)(a + mb) : 1 \le m \le (p-1)}$. Therefore $C(x) = C(k)$. Again, let $y \in S_j - Z$ for some $j \in \{1, 2, ..., (p + 1)\},$ then $C(y) \neq S_q$, where $1 \leq q(\neq j) \leq (p+1)$. Thus $C(y) = S_j$. Hence $| \text{Cent}(R) | = p + 2.$ \Box

Further, we have the following theorem analogous to Lemma 2.7 of [\[1\]](#page-11-0).

Theorem 2.12. Let R be a non-commutative ring whose order is a power of a prime p. Then $|\text{Cent}(R)| \geq p+2$, and equality holds if and only if R $\frac{R}{Z(R)} \cong \mathbb{Z}_p \times \mathbb{Z}_p.$

Proof. Let R be a non-commutative ring whose order is a power of a prime p. Suppose $k = \lvert \text{Cent}(R) \rvert$. Let A_1, \ldots, A_k be the distinct centralizers of R such that $|A_1| \geq \cdots \geq |A_k|$ and $A_1 = R$. So $R = \bigcup_{i=2}^k A_i$ and by Cohn's theorem in [\[7\]](#page-12-12), we have $|R| \leq \sum_{k=1}^{k}$ $i=3$ | A_i | (as A_i 's are additive groups). Also $|A_i| \leq \frac{|R|}{p}$, where $i \neq 1$. Hence

$$
|R| \le \underbrace{\frac{|R|}{p} + \dots + \frac{|R|}{p}}_{(k-2)-\text{times}}
$$

which implies $|R| \leq (k-2)\frac{|R|}{p}$ and so $k \geq p+2$. That is $|\text{Cent}(R)| \geq p+2$.

For the equality, if $\frac{R}{Z(R)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$ then by Theorem [2.11,](#page-4-0) we have $|\text{Cent}(R)| = p + 2$. Conversely, we assume that $l = |\text{Cent}(R)| = p + 2$. Suppose A_1, A_2, \ldots, A_l are distinct centralizers of R such that $|A_1| \geq \cdots \geq$ |A_l| and $A_1 = R$. So $R = \bigcup_{i=2}^{l} A_i$ and by Cohn's theorem in [\[7\]](#page-12-12), we have $|R| \leq \sum^{l}$ $i=3$ | $|A_i|$. Also $|A_i| \leq \frac{|R|}{p}$, where $i \neq 1$. Suppose, there exists an A_i such that $|A_i| < \frac{|R|}{n}$ $\frac{n_1}{p}$ for $3 \leq i \leq l$ then

$$
|R| < \underbrace{\frac{|R|}{p} + \dots + \frac{|R|}{p}}_{(l-2)-\text{times}} = (l-2)\frac{|R|}{p} = |R|,
$$

a contradiction. Hence $|A_3| = \frac{|R|}{n}$ $\frac{R|}{p},\ldots,|A_l|=\frac{|R|}{p}$ $\frac{R_1}{p}$. Also $|A_2| \geq \cdots \geq |A_l|$, so $|A_i| = \frac{|R|}{n}$ $\frac{R}{p}$, where $2 \leq i \leq l$. Hence $\sum_{i=0}^{l}$ $i=3$ $|A_i| = (l-2)\frac{|R|}{p} = |R|$. Therefore $\sum_{i=1}^{l}$ $i=3$ $|A_i| = |R|$ if and only if $A_2 + A_m = R$, for all $m \neq 2$ and $A_k \cap A_l \subseteq A_2$ for all $k \neq l$ (By Cohn's Theorem in [\[7\]](#page-12-12)). Interchanging A_i 's we have $A_2 \cap A_3 =$ $Z(R)$. Thus

$$
|R| = |A_2 + A_3| = \frac{|A_2||A_3|}{|A_2 \cap A_3|} = \frac{|R|^2}{p^2 |Z(R)|}
$$

which gives $|R : Z(R)| = p^2$. Hence $\frac{R}{Z(R)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$, since R is noncommutative. This completes the proof. \Box

We conclude this section by the following result.

Proposition 2.13. There exists an 8-centralizer ring.

Proof. We consider the ring $R = \{a + bi + cj + dk : a, b, c, d \in \mathbb{Z}_p, i^2 = j^2 = j^2\}$ $k^2 = -1$, $ij = k$, $jk = i$, $ki = j$, $ji = -k$, $kj = -i$, $ik = -j$. If $b = c = d = 0$, then clearly $C(a) = R$. If $c = d = 0$ and $b \neq 0$, then $C(a + bi) = \{x + yi$: $x, y \in \mathbb{Z}_p$. If $b = d = 0$ and $c \neq 0$, then $C(a + cj) = \{x + zj : x, z \in \mathbb{Z}_p\}$. If $b = c = 0$ and $d \neq 0$, then $C(a + dk) = \{x + wk : x, w \in \mathbb{Z}_p\}$. If $d = 0$ and $b, c \neq 0$, then $C(a+bi+cj) = \{x+yi+zj : bz = cy, x, y, z \in \mathbb{Z}_p\}$. If $c = 0$ and $b, d \neq 0$, then $C(a+bi+dk) = \{x+yi+wk : bw = dy, x, y, w \in \mathbb{Z}_p\}.$ If $b = 0$ and $c, d \neq 0$, then $C(a + cj + dk) = \{x + zj + wk : cw = dz, x, z, w \in \mathbb{Z}_p\}.$ If $b, c, d \neq 0$, then $C(a + bi + cj + dk) = \{x + yi + zj + wk : bz = cy, dy =$ $bw, dz = cw, x, y, z, w \in \mathbb{Z}_p$. Hence $|\text{Cent}(R)| = 8$. \Box

3 4-centralizer rings

In this section, we give a characterization of finite 4-centralizer rings analogous to Theorem 2 of [\[3\]](#page-12-7). The following lemma which is useful in characterization of 4-centralizer rings.

Lemma 3.1. Let R be a 4-centralizer finite ring. Then at least one of the centralizers of non-central elements has index 2 in R.

Proof. Let A, B, C be the three proper centralizers of R. Suppose none of A, B, C has index 2, that is $|R: A| \geq 3, |R: B| \geq 3, |R: C| \geq 3$. Then as $R = A \cup B \cup C$, we have

$$
|R| \le |A| + |B| + |C| - 2|Z(R)| \le \frac{|R|}{3} + \frac{|R|}{3} + \frac{|R|}{3} - 2|Z(R)| < |R|,
$$

which is a contradiction. Hence the lemma follows.

$$
\Box
$$

We have the following characterization of finite 4-centralizer rings.

Theorem 3.2. Let R be a non-commutative finite ring. Then $|\text{Cent}(R)| = 4$ if and only if $\frac{R}{Z(R)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.

Proof. If $\frac{R}{Z(R)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ then by Theorem [2.11,](#page-4-0) we have $|\text{Cent}(R)| = 4$.

Conversely, let $|\text{Cent}(R)| = 4$ then R has exactly four distinct centralizers, say R, A, B, C where A, B, C are centralizers of three distinct non-central elements of R.

By Lemma [2.5,](#page-2-2) R cannot be written as the union of two of its proper subrings of R. Therefore we may choose $a \in A-(B\cup C), b \in B-(C\cup A), c \in$ $C - (A \cup B)$ respectively. It can be easily seen that $C(a) = A, C(b) =$ $B, C(c) = C$. By Lemma [3.1,](#page-6-0) at least one of the centralizers A, B, C, say A has index 2 in R, that is $|R : A| = 2$.

Now, let $x \in (A \cap B) - Z(R)$ then $C(x) \neq R$. If $C(x) = A$ then $a, b \in C(x)$. So, $C(x) \neq A$. Similarly it can be seen that $C(x) \neq B$. If $C(x) = C$ then $x \in A \cap B \cap C = Z(R)$ (using Lemma [2.3\)](#page-2-0), which is a contradiction. Therefore $|\text{Cent}(R)|$ must be at least 5, which is again a contradiction. So $A \cap B =$ $A \cap B \cap C = Z(R)$. Similarly it can be seen that $B \cap C = Z(R)$, $A \cap C = Z(R)$. Again A, B, C are additive subgroups of R , therefore

$$
|R| \ge |A + B| = \frac{|A||B|}{|A \cap B|} = \frac{|A||B|}{|Z(R)|}
$$

which gives $|B| \leq 2|Z(R)|$. Since $Z(R) \subset B$, so $\frac{|B|}{2} \leq |Z(R)| < |B|$. Hence $|B| = 2|Z(R)|$. Similarly $|C| = 2|Z(R)|$. Therefore

$$
|R| = |A| + |B| + |C| - 2|Z(R)| = \frac{|R|}{2} + 2|Z(R)|
$$

which gives $|R:Z(R)|=4$ and hence $\frac{R}{Z(R)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.

$$
\square
$$

4 5-centralizer rings

In this section, we give a characterization of finite 5-centralizer rings analogous to Theorem 4 of [\[3\]](#page-12-7). The following lemmas are useful in this regard.

Lemma 4.1. Let R be a ring and $R = A \cup B \cup C$, where A, B, C are the proper distinct subrings. We put $K = A \cap B \cap C, L = A \cap B - K, M = A \cap C - K$, $N = B \cap C - K$ and $A' = A - (B \cup C)$, $B' = B - (A \cup C)$, $C' = C - (A \cup B)$. Then

- (a) $L = M = N = \phi$,
- (b) $A' + B' \subseteq C', B' + C' \subseteq A'$ and $C' + A' \subseteq B'$,
- (c) $A' + A' \subseteq K$, $B' + B' \subseteq K$ and $C' + C' \subseteq K$,
- (d) $|R: K| = 4$.

Proof. (a) We consider $l \in L$ and $c' \in C'$. Then $c' + l \in A$ or B or C. If $c' + l \in A$ then $c' + l + (-l) = c' \in A$, a contradiction. If $c' + l \in B$ then $c'+l+(-l) = c' \in B$, a contradiction. If $c'+l \in C$ then $(-c')+c'+l = l \in C$, a contradiction. Since $C' \neq \phi$, we must have $L = \phi$. Similarly $M = N = \phi$.

(b) Let $a' \in A'$, then $a' \in A \Rightarrow -a' \in A \Rightarrow -a' \in K$ or A'. If $-a' \in K$ then $a' \in K$, a contradiction. Hence $-a' \in A'$. Similarly if $b' \in B'$ then $-b' \in B'$ and if $c' \in C'$ then $-c' \in C'$. Suppose $a' \in A', b' \in B'$ then $a' + b' \in K$ or A' or B' or C' . If $a' + b' \in A' \subseteq A$ then $b' = -a' + a' + b' \in A$, a contradiction. If $a' + b' \in B' \subseteq B$, then $a' = a' + b' + (-b') \in B$, a contradiction. If $a' + b' \in K$, then $a' + b' \in A$, a contradiction. Hence $a' + b' \in C'$. Thus $A' + B' \subseteq C'$. Similarly it can be seen that $B' + C' \subseteq A'$ and $C' + A' \subseteq B'$.

(c) Let $a', a_1' \in A' \subseteq A$. So $a' + a_1' \in A \Rightarrow a' + a_1' \in A'$ or K. Let $a' + a_1' \in A'$. We consider $b' + a' + a_1'$, for some $b' \in B'$. Then by second part we have $b' + (a' + a_1') \in C'$ and $(b' + a') + a_1' \in B'$. So $b' + a' + a_1' \in B' \cap C'$, a contradiction. Similarly we can show the other two.

(d) From part (a), we have $R = K \cup A' \cup B' \cup C'$. Let $k + a' \in K + a'$ where $k \in K$, $a' \in A'$ then $k + a' \in A = K \cup A'$. If $k + a' \in K$ then $a' \in K$, a contradiction. So $K + a' \subseteq A'$. Again $x' \in A'$ gives $x' + (-a') \in K$ (by part (c)). So, $x' \in K + a'$. Hence $K + a' = A'$. Similarly it can be seen that $K + b' = B', K + c' = C'$, where $b' \in B', c' \in C'$. Therefore $|R: K| = 4$. \Box

Lemma 4.2. Let R be a 5-centralizer finite ring and A, B, C, D be the four proper centralizers of R. Then

- (a) $|R| = |A| + |B| + |C| + |D| 3|Z(R)|.$
- (b) If S and T are distinct proper centralizers of R, then

$$
\frac{|S||T|}{|R|} \le |Z(R)| \le \frac{|R|}{6}.
$$

Proof. Let $a \in A - (B \cup C), b \in B - (A \cup C)$ and $c \in C - (A \cup B)$. Suppose there does not exist any $a \in A - (B \cup C)$ such that $C(a) = A$. Then $C(a) = D$ for all $a \in A - (B \cup C)$. Therefore $A - (B \cup C) \subseteq D - (B \cup C)$. Interchanging the roles of A and D we get $A - (B \cup C) = D - (B \cup C)$,

which gives $A \cup B \cup C = D \cup B \cup C = R$. Again, by Lemma [4.1](#page-7-0) (a), we have $B \cap C = C \cap D$ and so $Z(R) = A \cap B \cap C$. Therefore, by Lemma [4.1](#page-7-0) (d), we have $R/Z(R) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. This gives $|\text{Cent}(R)| = 4$, contradiction. Hence $C(a) = A$. Similarly $C(b) = B$ and $C(c) = C$.

(a) Let us assume without loss of generality that D is a subset of $A\cup B\cup C$. Then $R = A \cup B \cup C \cup D = A \cup B \cup C$. Now, by Lemma [4.1,](#page-7-0) we have $|R:K| = 4$ where $K = A \cap B \cap C = Z(R)$. Thus by Theorem [3.2,](#page-6-1) $|\text{Cent}(R)| = 4$, which is a contradiction. Therefore no one of A, B, C or D is contained in the union of the other three.

Let $r \in (A \cap B) - (C \cup D)$ then $r \in C(a) \cap C(b)$ which gives $a, b \in C(r)$. But $a \notin C(b)$, so $C(r) \neq A, B$. Again $r \notin C, D$; so $C(r) \neq C, D$. Also $C(r) \neq R$, since $r \in R - Z(R)$. Therefore $|\text{Cent}(R)|$ must be at least 6, a contradiction. Hence $(A \cap B) - (C \cup D) = \phi$. This shows that no element of R is in exactly two proper centralizers.

Let $r \in (A \cap B \cap C) - D$ then $r \in C(a) \cap C(b) \cap C(c)$. Therefore $a, b, c \in$ $C(r)$. But $b \notin C(a), c \notin C(b)$. So $C(r) \neq A, B, C$. Also $C(r) \neq D, R$; as $r \notin D$ and $r \notin Z(R)$. Therefore $|\text{Cent}(R)|$ must be at least 6, a contradiction. Hence $A \cap B \cap C - D = \phi$. Thus no element of R is in exactly three proper centralizers.

From above, it can be seen clearly that

$$
|R| = |A \cup B \cup C \cup D| = |A| + |B| + |C| + |D| - 3|Z(R)|.
$$

(b) Note that for any two proper centralizers S and T of R we have $S \cap T = Z(R)$, since no element of R is in exactly two as well as three proper centralizers. Also any proper centralizers of R are additive subgroups of R , so $\frac{|S||T|}{|S+T|} = |S \cap T| = |Z(R)|$. Since $S + T \subseteq R$ we have $|Z(R)| \ge \frac{|S||T|}{|R|}$.

Again by part (a),

$$
|R| = |A| + |B| + |C| + |D| - 3|Z(R)|
$$

\n
$$
\geq 2|Z(R)| + 2|Z(R)| + 2|Z(R)| + 2|Z(R)| - 3|Z(R)|.
$$

Thus $|R : Z(R)| \geq 5$. If $|R : Z(R)| = 5$ then $\frac{R}{Z(R)} \cong \mathbb{Z}_5$, a contradiction. Therefore $|Z(R)| \leq \frac{|R|}{6}$. So, $\frac{|S||T|}{|R|} \leq |Z(R)| \leq \frac{|R|}{6}$. \Box

We would like to mention here that the group theoretic analogues of Lemma [4.1](#page-7-0) and Lemma [4.2](#page-8-0) can be found in [\[4\]](#page-12-13) and [\[3\]](#page-12-7) respectively. Now we prove the main theorem of this section which characterizes finite 5-centralizer rings.

Theorem 4.3. Let R be a finite ring. Then $|\text{Cent}(R)| = 5$ if and only if R $\frac{R}{Z(R)} \cong \mathbb{Z}_3 \times \mathbb{Z}_3.$

Proof. Let $\frac{R}{Z(R)} \cong \mathbb{Z}_3 \times \mathbb{Z}_3$, then by Theorem [2.11,](#page-4-0) we get $|\text{Cent}(R)| = 5$.

Conversely, let $|\text{Cent}(R)| = 5$. Let A, B, C, D be the four proper cen-tralizers of R. Then by Lemma [4.2](#page-8-0) (b), $\frac{|A||B|}{|R|} \leq |Z(R)| \leq \frac{|R|}{6}$. Our aim is to get more near lower bound for $|Z(R)|$. We may assume without loss of generality that $|A| \geq |B| \geq |C| \geq |D|$. Suppose $|A| < \frac{|R|}{3}$, as $1 < |A| \leq \frac{|R|}{2}$. $\sum_{i=1}^{\infty}$ That is $|A| \leq \frac{|R|}{4}$. Now by Lemma [4.2](#page-8-0) (a), $|R| \leq |R| - 3|Z(R)| < |R|$, a contradiction. Hence $|A| = \frac{|R|}{2}$ $\frac{|R|}{2}$ or $|A| = \frac{|R|}{3}$ $\frac{|R|}{3}$. If $|A| = \frac{|R|}{2}$ $\frac{R_1}{2}$, then $|R| =$ $|A| + |B| + |C| + |D| - 3|Z(R)|$ gives $\frac{|R|}{2} < |B| + |C| + |D|$ and so $\frac{|R|}{6} < |B|$. Also, applying Lemma [4.2](#page-8-0) (b) on A and B we have $\frac{|R|}{6} < |B| \le \frac{|R|}{3}$. So |B| is one of $\frac{|R|}{3}, \frac{|R|}{4}$ $\frac{R}{4}$ or $\frac{|R|}{5}$ $\frac{R_1}{5}$. Reapplying Lemma [4.2](#page-8-0) (b) on A and B we have,

$$
\frac{|A||B|}{|R|} \le |Z(R)| \le \frac{|R|}{6}
$$

which gives $\frac{|R|}{10} \leq |Z(R)| \leq \frac{|R|}{6}$. Thus $|Z(R)|$ is one of $\frac{|R|}{6}, \frac{|R|}{7}$ $\frac{|R|}{7},\frac{|R|}{8}$ $\frac{|R|}{8},\frac{|R|}{9}$ $rac{R}{9}$ or $\frac{|R|}{10}$. Let $|Z(R)| = \frac{|R|}{7}$ $\frac{|R|}{7},\frac{|R|}{9}$ $\frac{\kappa_0}{9}$ then 2 divides 7 and 9, which is not possible. If $|Z(R)| = \frac{|R|}{6}$ $\frac{R}{6}$ then $\frac{R}{Z(R)} \cong \mathbb{Z}_6$, a contradiction. Let $|Z(R)| = \frac{|R|}{8}$ $\frac{|R|}{8}$ then $\frac{|R|}{8}$ divides |B|. If $|B| = \frac{|R|}{3}$ $\frac{|R|}{3},\frac{|R|}{5}$ $rac{\kappa_1}{5}$ then 3,5 divides 8, a contradiction. Therefore $|B| = \frac{|R|}{4}$ $\frac{|R|}{4}$. By Lemma [4.2](#page-8-0) (a), we have $\frac{5|R|}{8} = |C| + |D|$. Also $|B| \ge |C| \ge |D|$. So $|C|+|D| \leq \frac{|R|}{2} < \frac{5|R|}{8} = |C|+|D|$, a contradiction. If $|Z(R)| = \frac{|R|}{10}$, then $\frac{|R|}{10}$ divides |B|. If $|B| = \frac{|R|}{3}$ $\frac{|R|}{3},\frac{|R|}{4}$ $\frac{\kappa_1}{4}$ then 3,4 divides 10, a contradiction. Therefore $|B| = \frac{|R|}{5}$. Now Lemma [4.2\(](#page-8-0)a) gives, $|C| + |D| = \frac{6|R|}{10}$. Also $|B| \ge |C| \ge |D|$, therefore $|C| + |D| \le \frac{2|R|}{5} < \frac{6|R|}{10} = |C| + |D|$, a contradiction.

If $|A| = \frac{|R|}{3}$ $\frac{R}{3}$ then Lemma [4.2](#page-8-0) (a) gives, $\frac{2|R|}{3} < |B| + |C| + |D|$ which gives $\frac{2|R|}{3} < 3|B|$ and so $|B| \ge \frac{|R|}{4}$. Also $|A| \ge |B|$, so $|B| = \frac{|R|}{3}$ $rac{|R|}{3}$ or $rac{|R|}{4}$ $\frac{n_1}{4}$. Again, applying Lemma [4.2](#page-8-0) (b) on A and B we get,

$$
\frac{|A||B|}{|R|} \le |Z(R)| \le \frac{|R|}{6}
$$

which gives $\frac{|R|}{12} \leq |Z(R)| \leq \frac{|R|}{6}$. Therefore $|Z(R)|$ is one of $\frac{|R|}{6}$, $\frac{|R|}{7}$ $\frac{|R|}{7},\frac{|R|}{8}$ $\frac{|R|}{8},\frac{|R|}{9}$ which gives $\frac{|R|}{12} \leq |Z(R)| \leq \frac{|R|}{6}$. Therefore $|Z(R)|$ is one of $\frac{|R|}{6}, \frac{|R|}{7}, \frac{|R|}{8}, \frac{|R|}{10}, \frac{|R|}{10}$, $\frac{|R|}{11}$ or $\frac{|R|}{12}$. Now if $|Z(R)| = \frac{|R|}{7}, \frac{|R|}{8}, \frac{|R|}{10}, \frac{|R|}{11}$ then 3 divides 7, 8, 10, 11, $\frac{|R|}{7},\frac{|R|}{8}$ $\frac{R}{8}$, $\frac{|R|}{10}$, $\frac{|R|}{11}$ then 3 divides 7, 8, 10, 11, a contradiction. Let $|Z(R)| = \frac{|R|}{6}$ $\frac{\kappa_1}{6}$ then as above we get a contradiction. Let $|Z(R)| = \frac{|R|}{9}$ $\frac{R}{9}$ then $\frac{R}{Z(R)} \cong \mathbb{Z}_3 \times \mathbb{Z}_3$. Let $|Z(R)| = \frac{|R|}{12}$ and $|B| = \frac{|R|}{3}$ $\frac{n_1}{3}$ then applying Lemma [4.2](#page-8-0) (b) on A and B we have, $\frac{|R|}{9} \le \frac{|R|}{12}$, a contradiction. If $|B|=\frac{|R|}{4}$ $\frac{R}{4}$ then Lemma [4.2](#page-8-0) (a) gives, $|C| + |D| = \frac{4|R|}{6}$ $\frac{|R|}{6}$. Also $|C|, |D| \le \frac{|R|}{4}$, so $|C|+|D| \leq \frac{3|R|}{6} < \frac{4|R|}{6} = |C|+|D|$, which is not possible. Hence $\frac{R}{Z(R)} \cong$ $\mathbb{Z}_3\times\mathbb{Z}_3$.

5 Relation between $| Cent(R)|$ and $d(R)$

Note that $d(R) = 1$ if and only if R is commutative. Therefore, by Propo-sition [2.1,](#page-1-0) we have $|\text{Cent}(R)| = 1$ if and only if $d(R) = 1$. By Theorem [3.2](#page-6-1) and Theorem 1 of [\[13\]](#page-12-8), we have the following result.

Proposition 5.1. Let R be a non-commutative finite ring. Then $|\text{Cent}(R)| =$ 4 if and only if $d(R) = \frac{5}{8}$.

In [\[13\]](#page-12-8), MacHale also proved the following theorem:

Theorem 5.2. Let R be a non-commutative finite ring and p the smallest prime dividing the order of R. Then $d(R) \leq \frac{1}{n^2}$ $\frac{1}{p^3}(p^2+p-1)$, with equality if and only if $|R:Z(R)| = p^2$.

Now by Theorem [2.11](#page-4-0) and Theorem [5.2,](#page-11-1) we have the following interesting connection between $d(R)$ and $| Cent(R)|$.

Proposition 5.3. Let R be a non-commutative finite ring and p the smallest prime dividing the order of R. If $d(R) = \frac{1}{p^3}(p^2+p-1)$ then $|\text{Cent}(R)| = p+2$.

We conclude the paper by noting that the converse of Proposition [5.3](#page-11-2) holds for some finite non-commutative rings. In particular, by Theorem [2.12](#page-5-0) and Theorem [5.2,](#page-11-1) we have the following result.

Proposition 5.4. Let R be a non-commutative ring whose order is a power of a prime p. If $|\text{Cent}(R)| = p + 2$ then $d(R) = \frac{1}{p^3}(p^2 + p - 1)$.

References

[1] A. R. Ashrafi, On finite groups with a given number of centralizers, Alqebra Colloq., $7(2)(2000)$, 139-146.

- [2] M. Behboodi, R. Beyranvand, A. Hashemi and H. Khabazian, Classification of finite rings: theory and algorithm, *Czechoslovak Math. J.*, **64**(3) (2014), 641–658.
- [3] S. M. Belcastro and G. J. Sherman, Counting centralizers in finite groups, Math. Magazine, 67 (5) (1994), 366–374.
- [4] M. Bruckheimer, A. C. Bryan and A. Muir, Groups which are the union of three subgroups, Amer. Math. Monthly, 77 (1970), 52-57.
- [5] S. M. Buckley, D. MacHale and A. Ni Shé, Finite rings with many commuting pairs of elements, Preprint.
- [6] S. M. Buckley and D. MacHale, Contrasting the commuting probabilities of groups and rings, Preprint.
- [7] J. H. E. Cohn, On n-sum groups, *Math. Scand.*, **75** (1994), 44–58.
- [8] C. J. Chikunji, A Classification of a certain class of completely primary finite rings, Ring and Module Theory, Trends in Mathematics 2010, Springer Basel, pp 83–90.
- [9] J. B. Derr, G. F. Orr and P. S. Peck, Noncommutative rings of order p^4 , J. Pure Appl. Algebra, 97(2) (1994), 109–116.
- $[10]$ K. E. Eldridge, Orders for finite noncommutative rings with unity, Amer. Math. Monthly, 75 (1968), 512–514.
- [11] B. Fine, Classification of finite rings of order p^2 , Math. Magazine, 66(4) (1993), 248–252.
- [12] R.W. Goldbach and H.L. Claasen, Classification of not commutative rings with identity of order dividing p^4 , *Indag. Math.*, 6 (1995), 167–187.
- [13] D. MacHale, Commutativity in finite rings, Amer. Math. Monthly, 83(1976), 30–32.
- [14] G. R. Omidi and E. Vatandoost, On the commuting Graph of rings, J. Algebra and Appl., 10 3(2011), 521–527.
- [15] R. Raghavendran, A class of finite rings, Compositio Math., 22 (1970), 49–57.