# Measurement of the branching fraction for  $\psi(3770) \rightarrow \gamma \chi_{c0}$

M. Ablikim<sup>1</sup>, M. N. Achasov<sup>9,f</sup>, X. C. Ai<sup>1</sup>, O. Albayrak<sup>5</sup>, M. Albrecht<sup>4</sup>, D. J. Ambrose<sup>44</sup>, A. Amoroso<sup>49A,49C</sup>, F. F. An<sup>1</sup>, Q. An<sup>46,a</sup>, J. Z. Bai<sup>1</sup>, R. Baldini Ferroli<sup>204</sup>, Y. Ban<sup>31</sup>, D. W. Bennett<sup>19</sup>, J. V. Bennett<sup>5</sup>, M. Bertani<sup>204</sup>, D. Bettoni<sup>21A</sup>, J. M. Bian<sup>43</sup>, F. Bianchi<sup>49A,49C</sup>, E. Boger<sup>23,d</sup>, I. Boyko<sup>23</sup>, R. A. Briere<sup>5</sup>, H. Cai<sup>51</sup>, X. Cai<sup>1,a</sup>, O. Cakir<sup>40A,b</sup>, A. Calcaterra<sup>20A</sup>, G. F. Cao<sup>1</sup>, S. A. Cetin<sup>40B</sup>, J. F. Chang<sup>1,a</sup>, G. Chelkov<sup>23,d,e</sup>, G. Chen<sup>1</sup>, H. S. Chen<sup>1</sup>, H. Y. Chen<sup>2</sup>, J. C. Chen<sup>1</sup>, M. L. Chen<sup>1,a</sup>, S. J. Chen<sup>29</sup>, X. Chen<sup>1,a</sup>, X. R. Chen<sup>26</sup>, Y. B. Chen<sup>1,a</sup>, H. P. Cheng<sup>17</sup>, X. K. Chu<sup>31</sup>, G. Cibinetto<sup>21A</sup>, H. L. Dai<sup>1,a</sup>, J. P. Dai<sup>34</sup>, A. Dbeyssi<sup>14</sup>, D. Dedovich<sup>23</sup>, Z. Y. Deng<sup>1</sup>, A. Denig<sup>22</sup>, I. Denysenko<sup>23</sup>, M. Destefanis<sup>49A,49C</sup>, F. De Mori<sup>49A,49C</sup>, Y. Ding<sup>27</sup>, C. Dong<sup>30</sup>, J. Dong<sup>1,a</sup>, L. Y. Dong<sup>1</sup>, , M. Y. Dong<sup>1,a</sup>, Z. L. Dou<sup>29</sup>, S. X. Du<sup>53</sup>, P. F. Duan<sup>1</sup>, E. E. Eren<sup>40B</sup>, J. Z. Fan<sup>39</sup>, J. Fang<sup>1,a</sup>, S. S. Fang<sup>1</sup>, X. Fang<sup>46,a</sup>, Y. Fang<sup>1</sup>, R. Farinelli<sup>21A,21B</sup>, L. Fava<sup>49B,49C</sup>, O. Fedorov<sup>23</sup>, F. Feldbauer<sup>22</sup>, G. Felici<sup>20A</sup>, C. Q. Feng<sup>46,a</sup>, E. Fioravanti<sup>21A</sup>, M. Fritsch<sup>14,22</sup>, C. D. Fu<sup>1</sup>, Q. Gao<sup>1</sup>, X. L. Gao<sup>46,a</sup>, X. Y. Gao<sup>2</sup>, Y. Gao<sup>39</sup>, Z. Gao $^{46,a}$ , I. Garzia<sup>21A</sup>, K. Goetzen<sup>10</sup>, L. Gong<sup>30</sup>, W. X. Gong<sup>1,a</sup>, W. Gradl<sup>22</sup>, M. Greco<sup>49A,49C</sup>, M. H. Gu<sup>1,a</sup>, Y. T.  $\rm{Gu^{12}, Y. H.~Guan^{1}, A. Q.~Guo^{1}, L.~B.~Guo^{28}, Y.~Guo^{1}, Y.~P.~Guo^{22}, Z.~Haddadi^{25}, A.~Hafner^{22}, S.~Han^{51},}$  $X. Q. Hao<sup>15</sup>, F. A. Harris<sup>42</sup>, K. L. He<sup>1</sup>, T. Held<sup>4</sup>, Y. K. Heng<sup>1,a</sup>, Z. L. Hou<sup>1</sup>, C. Hu<sup>28</sup>, H. M. Hu<sup>1</sup>, J. F. Hu<sup>49A,49C</sup>,$ T.  $\rm{Hu}^{1,a}$ , Y.  $\rm{Hu}^1$ , G. S.  $\rm{Huang}^{46,a}$ , J. S.  $\rm{Huang}^{15}$ , X. T.  $\rm{Huang}^{33}$ , Y.  $\rm{Huang}^{29}$ , T.  $\rm{Hussain}^{48}$ , Q. Ji<sup>1</sup>, Q. P. Ji<sup>30</sup>, X. B. Ji<sup>1</sup>, X. L. Ji<sup>1,a</sup>, L. W. Jiang<sup>51</sup>, X. S. Jiang<sup>1,a</sup>, X. Y. Jiang<sup>30</sup>, J. B. Jiao<sup>33</sup>, Z. Jiao<sup>17</sup>, D. P. Jin<sup>1,a</sup>, S. Jin<sup>1</sup>, T. Johansson<sup>50</sup>, A. Julin<sup>43</sup>, N. Kalantar-Nayestanaki<sup>25</sup>, X. L. Kang<sup>1</sup>, X. S. Kang<sup>30</sup>, M. Kavatsyuk<sup>25</sup>, B. C. Ke<sup>5</sup>, P. Kiese<sup>22</sup>, R. Kliemt<sup>14</sup>, B. Kloss<sup>22</sup>, O. B. Kolcu<sup>40B,i</sup>, B. Kopf<sup>4</sup>, M. Kornicer<sup>42</sup>, W. Kuehn<sup>24</sup>, A. Kupsc<sup>50</sup>, J. S. Lange<sup>24,a</sup>, M. Lara<sup>19</sup>, P. Larin<sup>14</sup>, C. Leng<sup>49C</sup>, C. Li<sup>50</sup>, Cheng Li<sup>46,a</sup>, D. M. Li<sup>53</sup>, F. Li<sup>1,a</sup>, F. Y. Li<sup>31</sup>, G. Li<sup>1</sup>, H. B. Li<sup>1</sup>, J. C. Li<sup>1</sup>, Jin Li<sup>32</sup>, K. Li<sup>13</sup>, K. Li<sup>33</sup>, Lei Li<sup>3</sup>, P. R. Li<sup>41</sup>, Q. Y. Li<sup>33</sup>, T. Li<sup>33</sup>, W. D. Li<sup>1</sup>, W. G. Li<sup>1</sup>, X. L. Li<sup>33</sup>, X. M. Li<sup>12</sup>, X. N. Li<sup>1,a</sup>, X. Q. Li<sup>30</sup>, Z. B. Li<sup>38</sup>, H. Liang<sup>46,a</sup>, Y. F. Liang<sup>36</sup>, Y. T. Liang<sup>24</sup>, G. R. Liao<sup>11</sup> , D. X. Lin<sup>14</sup>, B. J. Liu<sup>1</sup>, C. X. Liu<sup>1</sup>, D. Liu<sup>46,a</sup>, F. H. Liu<sup>35</sup>, Fang Liu<sup>1</sup>, Feng Liu<sup>6</sup>, H. B. Liu<sup>12</sup>, H. H. Liu<sup>1</sup>, H. H. Liu $^{16}$ , H. M. Liu $^1$ , J. Liu $^1$ , J. B. Liu $^{46,a}$ , J. P. Liu $^{51}$ , J. Y. Liu $^1$ , K. Liu $^{39}$ , K. Y. Liu $^{27}$ , L. D. Liu $^{31}$ , P. L. Liu<sup>1,a</sup>, Q. Liu<sup>41</sup>, S. B. Liu<sup>46,a</sup>, X. Liu<sup>26</sup>, Y. B. Liu<sup>30</sup>, Z. A. Liu<sup>1,a</sup>, Zhiqing Liu<sup>22</sup>, H. Loehner<sup>25</sup>, X. C. Lou<sup>1,a,h</sup>, H. J. Lu<sup>17</sup>, J. G. Lu<sup>1,a</sup>, Y. Lu<sup>1</sup>, Y. P. Lu<sup>1,a</sup>, C. L. Luo<sup>28</sup>, M. X. Luo<sup>52</sup>, T. Luo<sup>42</sup>, X. L. Luo<sup>1,a</sup>,  $X.$  R. Lyu<sup>41</sup>, F. C. Ma<sup>27</sup>, H. L. Ma<sup>1</sup>, L. L. Ma<sup>33</sup>, Q. M. Ma<sup>1</sup>, T. Ma<sup>1</sup>, X. N. Ma<sup>30</sup>, X. Y. Ma<sup>1,a</sup>, Y. M. Ma<sup>33</sup>, F. E. Maas<sup>14</sup>, M. Maggiora<sup>49A,49C</sup>, Y. J. Mao<sup>31</sup>, Z. P. Mao<sup>1</sup>, S. Marcello<sup>49A,49C</sup>, J. G. Messchendorp<sup>25</sup>, J. Min<sup>1,a</sup>, R. E. Mitchell<sup>19</sup>, X. H. Mo<sup>1,a</sup>, Y. J. Mo<sup>6</sup>, C. Morales Morales<sup>14</sup>, N. Yu. Muchnoi<sup>9,f</sup>, H. Muramatsu<sup>43</sup>, Y. Nefedov<sup>23</sup>, F. Nerling<sup>14</sup>, I. B. Nikolaev<sup>9,f</sup>, Z. Ning<sup>1,a</sup>, S. Nisar<sup>8</sup>, S. L. Niu<sup>1,a</sup>, X. Y. Niu<sup>1</sup>, S. L. Olsen<sup>32</sup>, Q. Ouyang<sup>1,a</sup>, S. Pacetti<sup>20B</sup>, Y. Pan<sup>46,a</sup>, P. Patteri<sup>20A</sup>, M. Pelizaeus<sup>4</sup>, H. P. Peng<sup>46,a</sup>, K. Peters<sup>10</sup>, J. Pettersson<sup>50</sup>, J. L. Ping<sup>28</sup>, R. G. Ping<sup>1</sup>, R. Poling<sup>43</sup>, V. Prasad<sup>1</sup>, H. R. Qi<sup>2</sup>, M. Qi<sup>29</sup>, S. Qian<sup>1,a</sup>, C. F. Qiao<sup>41</sup>, L. Q. Qin<sup>33</sup>, N.  $\rm{Qin}^{51}$ , X. S.  $\rm{Qin}^{1}$ , Z. H.  $\rm{Qin}^{1,a}$ , J. F.  $\rm{Qiu}^{1}$ , K. H. Rashid<sup>48</sup>, C. F. Redmer<sup>22</sup>, M. Ripka<sup>22</sup>, G. Rong<sup>1</sup>, Ch. Rosner<sup>14</sup>, X. D. Ruan<sup>12</sup>, V. Santoro<sup>21A</sup>, A. Sarantsev<sup>23,g</sup>, M. Savrié<sup>21B</sup>, K. Schoenning<sup>50</sup>, S. Schumann<sup>22</sup>, W. Shan<sup>31</sup>, M. Shao<sup>46,a</sup>, C. P. Shen<sup>2</sup>, P. X. Shen<sup>30</sup>, X. Y. Shen<sup>1</sup>, H. Y. Sheng<sup>1</sup>, W. M. Song<sup>1</sup>, X. Y. Song<sup>1</sup>, S. Sosio<sup>49A,49C</sup>, S. Spataro<sup>49A,49C</sup>, G. X. Sun<sup>1</sup>, J. F. Sun<sup>15</sup>, S. S. Sun<sup>1</sup>, Y. J. Sun<sup>46,a</sup>, Y. Z. Sun<sup>1</sup>, Z. J. Sun<sup>1,a</sup>, Z. T. Sun<sup>19</sup>, C. J. Tang<sup>36</sup>, X. Tang<sup>1</sup>, I. Tapan<sup>40C</sup>, E. H. Thorndike<sup>44</sup>, M. Tiemens<sup>25</sup>, M. Ullrich<sup>24</sup>, I. Uman<sup>40D</sup>, G. S. Varner<sup>42</sup>, B. Wang<sup>30</sup>, B. L. Wang<sup>41</sup>, D. Wang<sup>31</sup>, D. Y. Wang<sup>31</sup>, K. Wang<sup>1,a</sup>, L. L. Wang<sup>1</sup>, L. S. Wang<sup>1</sup>, M.  $\text{Wang}^{33}$ , P.  $\text{Wang}^1$ , P. L.  $\text{Wang}^1$ , S. G.  $\text{Wang}^{31}$ , W.  $\text{Wang}^{1,a}$ , W. P.  $\text{Wang}^{46,a}$ , X. F.  $\text{Wang}^{39}$ , Y. D.  $\text{Wang}^{14}$ , Y. F. Wang<sup>1,a</sup>, Y. Q. Wang<sup>22</sup>, Z. Wang<sup>1,a</sup>, Z. G. Wang<sup>1,a</sup>, Z. H. Wang<sup>46,a</sup>, Z. Y. Wang<sup>1</sup>, T. Weber<sup>22</sup>, D. H. Wei<sup>11</sup>, J. B. Wei<sup>31</sup>, P. Weidenkaff<sup>22</sup>, S. P. Wen<sup>1</sup>, U. Wiedner<sup>4</sup>, M. Wolke<sup>50</sup>, L. H. Wu<sup>1</sup>, Z. Wu<sup>1,a</sup>, L. Xia<sup>46,a</sup>, L. G. Xia<sup>39</sup>, Y. Xia<sup>18</sup>, D. Xiao<sup>1</sup>, H. Xiao<sup>47</sup>, Z. J. Xiao<sup>28</sup>, Y. G. Xie<sup>1,a</sup>, Q. L. Xiu<sup>1,a</sup>, G. F. Xu<sup>1</sup>, L. Xu<sup>1</sup>, Q. J. Xu<sup>13</sup>, Q. N. Xu<sup>41</sup>, X. P. Xu<sup>37</sup>, L. Yan<sup>49A,49C</sup>, W. B. Yan<sup>46,a</sup>, W. C. Yan<sup>46,a</sup>, Y. H. Yan<sup>18</sup>, H. J. Yang<sup>34</sup>, H. X. Yang<sup>1</sup>, L. Yang<sup>51</sup> , Y. X. Yang<sup>11</sup>, M. Ye<sup>1,a</sup>, M. H. Ye<sup>7</sup>, J. H. Yin<sup>1</sup>, B. X. Yu<sup>1,a</sup>, C. X. Yu<sup>30</sup>, J. S. Yu<sup>26</sup>, C. Z. Yuan<sup>1</sup>, W. L. Yuan<sup>29</sup>, Y. Yuan<sup>1</sup>, A. Yuncu<sup>40B,c</sup>, A. A. Zafar<sup>48</sup>, A. Zallo<sup>20A</sup>, Y. Zeng<sup>18</sup>, Z. Zeng<sup>46,a</sup>, B. X. Zhang<sup>1</sup>, B. Y. Zhang<sup>1,a</sup>, C. Zhang<sup>29</sup>, C. C. Zhang<sup>1</sup>, D. H. Zhang<sup>1</sup>, H. H. Zhang<sup>38</sup>, H. Y. Zhang<sup>1, a</sup>, J. J. Zhang<sup>1</sup>, J. L. Zhang<sup>1</sup>, J. Q. Zhang<sup>1</sup>, J. W. Zhang<sup>1,a</sup>, J. Y. Zhang<sup>1</sup>, J. Z. Zhang<sup>1</sup>, K. Zhang<sup>1</sup>, L. Zhang<sup>1</sup>, X. Y. Zhang<sup>33</sup>, Y. Zhang<sup>1</sup>, Y. H. Zhang<sup>1,a</sup> , Y. N. Zhang<sup>41</sup>, Y. T. Zhang<sup>46,a</sup>, Yu Zhang<sup>41</sup>, Z. H. Zhang<sup>6</sup>, Z. P. Zhang<sup>46</sup>, Z. Y. Zhang<sup>51</sup>, G. Zhao<sup>1</sup>, J. W. Zhao<sup>1,a</sup>, J. Y. Zhao<sup>1</sup>, J. Z. Zhao<sup>1,a</sup>, Lei Zhao<sup>46,a</sup>, Ling Zhao<sup>1</sup>, M. G. Zhao<sup>30</sup>, Q. Zhao<sup>1</sup>, Q. W. Zhao<sup>1</sup>, S. J. Zhao<sup>53</sup>, T. C. Zhao<sup>1</sup>, Y. B. Zhao<sup>1,a</sup>, Z. G. Zhao<sup>46,a</sup>, A. Zhemchugov<sup>23,d</sup>, B. Zheng<sup>47</sup>, J. P. Zheng<sup>1,a</sup>, W. J. Zheng<sup>33</sup>, Y. H. Zheng<sup>41</sup>, B. Zhong<sup>28</sup>, L. Zhou<sup>1,a</sup>, X. Zhou<sup>51</sup>, X. K. Zhou<sup>46,a</sup>, X. R. Zhou<sup>46,a</sup>, X. Y. Zhou<sup>1</sup>, K. Zhu<sup>1</sup>, K. J. Zhu<sup>1,a</sup>, S. Zhu<sup>1</sup>, S. H. Zhu<sup>45</sup>, X. L. Zhu<sup>39</sup>, Y. C. Zhu<sup>46,a</sup>, Y. S. Zhu<sup>1</sup>, Z. A. Zhu<sup>1</sup>, J. Zhuang<sup>1,a</sup>,

L.  $Zotti^{49A,49C}$ , B. S.  $Zou<sup>1</sup>$ , J. H.  $Zou<sup>1</sup>$ 

(BESIII Collaboration)

<sup>1</sup> Institute of High Energy Physics, Beijing 100049, People's Republic of China

Beihang University, Beijing 100191, People's Republic of China

<sup>3</sup> Beijing Institute of Petrochemical Technology, Beijing 102617, People's Republic of China

 $4\overline{\hbox{Bochum Ruhr-University}}$ , D-44780 Bochum, Germany

Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

Central China Normal University, Wuhan 430079, People's Republic of China

China Center of Advanced Science and Technology, Beijing 100190, People's Republic of China

COMSATS Institute of Information Technology, Lahore, Defence Road, Off Raiwind Road, 54000 Lahore,

Pakistan

G.I. Budker Institute of Nuclear Physics SB RAS (BINP), Novosibirsk 630090, Russia

<sup>10</sup> GSI Helmholtzcentre for Heavy Ion Research GmbH, D-64291 Darmstadt, Germany

<sup>11</sup> Guangxi Normal University, Guilin  $5/100/4$ , People's Republic of China

GuangXi University, Nanning 530004, People's Republic of China

Hangzhou Normal University, Hangzhou 310036, People's Republic of China

Helmholtz Institute Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany

<sup>15</sup> Henan Normal University, Xinxiang 453007, People's Republic of China

<sup>16</sup> Henan University of Science and Technology, Luoyang 471003, People's Republic of China

Huangshan College, Huangshan 245000, People's Republic of China

Hunan University, Changsha 410082, People's Republic of China

Indiana University, Bloomington, Indiana 47405, USA

 $^{20}$  (A)INFN Laboratori Nazionali di Frascati, I-00044, Frascati, Italy; (B)INFN and University of Perugia,

I-06100, Perugia, Italy

<sup>21</sup> (A)INFN Sezione di Ferrara, I-44122, Ferrara, Italy; (B)University of Ferrara, I-44122, Ferrara, Italy

 $^{22}$  Johannes Gutenberg University of Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany

<sup>23</sup> Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia

Justus Liebig University Giessen, II. Physikalisches Institut, Heinrich-Buff-Ring 16, D-35392 Giessen, Germany

 $^{25}$  KVI-CART, University of Groningen, NL-9747 AA Groningen, The Netherlands

Lanzhou University, Lanzhou 730000, People's Republic of China

Liaoning University, Shenyang 110036, People's Republic of China

Nanjing Normal University, Nanjing 210023, People's Republic of China

Nanjing University, Nanjing 210093, People's Republic of China

Nankai University, Tianjin 300071, People's Republic of China<br> $31$  Petine University, Petities 100871, People's Penellis of China

Peking University, Beijing 100871, People's Republic of China

 $3^{2}$  Seoul National University, Seoul, 151-747 Korea

Shandong University, Jinan 250100, People's Republic of China

Shanghai Jiao Tong University, Shanghai 200240, People's Republic of China

Shanxi University, Taiyuan 030006, People's Republic of China

Sichuan University, Chengdu 610064, People's Republic of China

Soochow University, Suzhou 215006, People's Republic of China

Sun Yat-Sen University, Guangzhou 510275, People's Republic of China

Tsinghua University, Beijing 100084, People's Republic of China

(A)Istanbul Aydin University, 34295 Sefakoy, Istanbul, Turkey; (B)Istanbul Bilgi University, 34060 Eyup,

Istanbul, Turkey; (C)Uludag University, 16059 Bursa, Turkey; (D)Near East University, Nicosia, North Cyprus,

10, Mersin, Turkey

<sup>41</sup> University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China

University of Hawaii, Honolulu, Hawaii 96822, USA

University of Minnesota, Minneapolis, Minnesota 55455, USA

University of Rochester, Rochester, New York 14627, USA

University of Science and Technology Liaoning, Anshan 114051, People's Republic of China

University of Science and Technology of China, Hefei 230026, People's Republic of China

University of South China, Hengyang  $\mu$ 21001, People's Republic of China

University of the Punjab, Lahore-54590, Pakistan

 $^{49}$  (A)University of Turin, I-10125, Turin, Italy; (B)University of Eastern Piedmont, I-15121, Alessandria, Italy;

 $(C)$ INFN, I-10125, Turin, Italy

Uppsala University, Box 516, SE-75120 Uppsala, Sweden

Wuhan University, Wuhan 430072, People's Republic of China

<sup>52</sup> Zhejiang University, Hangzhou 310027, People's Republic of China <sup>53</sup> Zhengzhou University, Zhengzhou 450001, People's Republic of China

<sup>a</sup> Also at State Key Laboratory of Particle Detection and Electronics, Beijing 100049, Hefei 230026, People's

Republic of China

 $<sup>b</sup>$  Also at Ankara University,06100 Tandogan, Ankara, Turkey</sup>

 $c^c$  Also at Bogazici University, 34342 Istanbul, Turkey

<sup>d</sup> Also at the Moscow Institute of Physics and Technology, Moscow 141700, Russia

 $e$  Also at the Functional Electronics Laboratory, Tomsk State University, Tomsk, 634050, Russia

 $f$  Also at the Novosibirsk State University, Novosibirsk, 630090, Russia

<sup>g</sup> Also at the NRC "Kurchatov Institute", PNPI, 188300, Gatchina, Russia

<sup>h</sup> Also at University of Texas at Dallas, Richardson, Texas 75083, USA

<sup>i</sup> Also at Istanbul Arel University, 34295 Istanbul, Turkey

#### Abstract

By analyzing a data set of 2.92 fb<sup>-1</sup> of  $e^+e^-$  collision data taken at  $\sqrt{s} = 3.773$  GeV and  $106.41 \times 10^6$  $\psi(3686)$  decays taken at  $\sqrt{s} = 3.686$  GeV with the BESIII detector at the BEPCII collider, we measure the branching fraction and the partial decay width for  $\psi(3770) \to \gamma \chi_{c0}$  to be  $\mathcal{B}(\psi(3770) \to \gamma \chi_{c0})$  $(6.88 \pm 0.28 \pm 0.67) \times 10^{-3}$  and  $\Gamma[\psi(3770) \to \gamma \chi_{c0}] = (187 \pm 8 \pm 19)$  keV, respectively. These are the most precise measurements to date.

# 1. Introduction

Transitions between charmonium states can be used to shed light on various aspects of Quantum Chromodynamics (QCD), the theory of the strong interactions, in both the perturbative and non-perturbative regimes [\[1\]](#page-9-0). The  $\psi(3770)$  resonance is the lowest-mass charmonium state lying above the production threshold of open-charm  $D\overline{D}$  pairs. It is assumed to be the  $1^3D_1$  cc state with a small  $2^3S_1$  admixture. Based on this S-D mixing model, predictions have been made [\[2,](#page-10-0) [3,](#page-10-1) [4](#page-10-2), [5](#page-10-3), [6\]](#page-10-4) for the partial widths of the  $\psi(3770)$ electric-dipole  $(E1)$  radiative transitions. These predictions vary over a large range depending on the underlying model assumptions. One of the largest variations in predictions is for the partial width of  $\psi(3770) \rightarrow \gamma \chi_{c0}$ , with predictions ranging from 213 keV to 523 keV. A precise measurement of the partial width of  $\psi(3770) \to \gamma \chi_{c0}$  provides a stringent test of the various theoretical approaches, thereby providing a better understanding of  $\psi(3770)$  decays.

In 2006, the CLEO Collaboration reported the first observation of  $\psi(3770) \to \gamma \chi_{c0/1}$  and measured the partial widths [\[7,](#page-10-5) [8](#page-10-6)]. A comparison between their results and predictions of traditional theory models [\[2](#page-10-0), [3](#page-10-1), [4](#page-10-2), [5](#page-10-3)] indicates that relativistic and coupled-channel effects are necessary ingredients to describe the data. A similar conclusion has been drawn in  $\psi(3686) \to \gamma \chi_{cJ}$  decays [\[9\]](#page-10-7). The results of CLEO were normalized to the cross section of  $\psi(3770) \to DD$  to obtain the total number of  $\psi(3770)$  decays, which assumed the contribution of  $\psi(3770) \to \text{non-}DD$  decays is negligible [\[27](#page-10-8)]. Recently, the BESIII Collaboration presented an improved measurement of  $\psi(3770) \rightarrow \gamma \chi_{c1}$  [\[10\]](#page-10-9).

In this Letter, we report on an alternative and complementary measurement of the branching fraction and partial width of  $\psi(3770) \to \gamma \chi_{c0}$  using  $\chi_{c0} \to 2(\pi^+\pi^-)$ ,  $K^+K^-\pi^+\pi^-$ ,  $3(\pi^+\pi^-)$  and  $K^+K^-$  decays. The results of our measurements are obtained by taking the relative strength with respect to the well-known  $\psi(3686)$  radiative E1 transition [\[11\]](#page-10-10). In this way, the measurement will not depend on knowledge of the  $\chi_{cJ}$ branching fractions to light hadron final states, which have large uncertainties [\[7\]](#page-10-5). This measurement forms an independent and more precise benchmark that can be compared to the predictions of various theoretical models.

### 2. BESIII detector and Monte Carlo simulation

In this work, we use 2.92 fb<sup>-1</sup> of  $e^+e^-$  collision data taken at  $\sqrt{s} = 3.773$  GeV [\[12\]](#page-10-11), and 106.41 ×10<sup>6</sup>  $\psi(3686)$  decays taken at  $\sqrt{s} = 3.686$  GeV [\[13\]](#page-10-12) with the BESIII detector. These are labeled the  $\psi(3770)$  and  $\psi(3686)$  data samples, respectively, throughout this Letter.

The BESIII detector [\[14\]](#page-10-13) has a geometrical acceptance of 93% of  $4\pi$  and consists of four main components. In the following, we describe each detector component starting from the innermost (closest to the interaction region) to the most outside layer. The inner three components are immersed in the 1 T magnetic field of a superconducting solenoid. First, a small-cell, helium-based main drift chamber (MDC) with 43 layers provides charged particle tracking and measurement of ionization energy loss  $(dE/dx)$ . The average single wire resolution is 135  $\mu$ m, and the momentum resolution for 1 GeV electrons in a 1 T magnetic field is 0.5%. The next detector after the MDC is a time-of-flight system (TOF) used for particle identification. It is composed of a barrel part made of two layers of 88 plastic scintillators, each with 5 cm thickness and 2.4 m length; and two endcaps, each with 96 fan-shaped plastic scintillators of 5 cm thickness. The time resolution is 80 ps in the barrel, and 110 ps in the endcaps, corresponding to a  $K/\pi$  separation better than 2σ for momenta up to about 1.0 GeV. The third detector component is an electromagnetic calorimeter (EMC) made of 6240 CsI(Tl) crystals arranged in a cylindrical shape (barrel) plus two endcaps. For 1.0 GeV photons, the energy resolution is 2.5% in the barrel and 5% in the endcaps, and the position resolution is 6 mm in the barrel and 9 mm in the endcaps. Outside the EMC, a muon chamber system (MUC) is incorporated in the return iron of the superconducting magnet. It is made of  $1272 \text{ m}^2$  of resistive plate chambers arranged in 9 layers in the barrel and 8 layers in the endcaps. The position resolution is about 2 cm.

A GEANT4 [\[15\]](#page-10-14) based Monte Carlo (MC) simulation software package, which includes the geometric description of the detector and the detector response, is used to determine the detection efficiency of the signal process and to estimate the potential peaking backgrounds. Signal MC samples of  $\psi(3686)/\psi(3770) \rightarrow \gamma \chi_{cJ}$ are generated with the angular distribution that corresponds to an E1 transition, and the  $\chi_{cJ}$  decays to light hadron final states are generated according to a phase-space model. Particle decays are modeled using EvtGen [\[16](#page-10-15)], while the initial production is handled by the MC generator KKMC [\[17\]](#page-10-16), in which both initial state radiation (ISR) effects [\[18](#page-10-17)] and final state radiation (FSR) effects [\[19\]](#page-10-18) are considered. For the background studies of  $\psi(3686)$  decays,  $106\times10^6$  MC events of generic decays  $\psi(3686) \rightarrow$  anything are produced at  $\sqrt{s} = 3.686$  GeV. For the background studies of  $\psi(3770)$  decays, MC samples of  $\psi(3770) \rightarrow$  $D^0\bar{D}^0$ ,  $\psi(3770) \to D^+D^-$ ,  $\psi(3770) \to \text{non-}D\bar{D}$  decays, ISR production of  $\psi(3686)$  and  $J/\psi$ , QED, and  $q\bar{q}$  continuum processes are produced at  $\sqrt{s} = 3.773$  GeV. The known decay modes of the  $J/\psi$ ,  $\psi(3686)$ and  $\psi(3770)$  are generated with branching fractions taken from the PDG [\[11\]](#page-10-10), and the remaining events are generated with Lundcharm [\[20](#page-10-19)].

#### 3. Analysis

To select candidate events for  $\psi(3686)/\psi(3770) \to \gamma \chi_{cJ}$  with  $\chi_{cJ} \to 2(\pi^+\pi^-)/K^+K^-\pi^+\pi^-/3(\pi^+\pi^-)/$  $K^+K^-$ , we require at least  $4/4/6/2$  charged tracks to be reconstructed in the MDC, respectively. All charged tracks used in this analysis are required to be within a polar-angle  $(\theta)$  range of  $|\cos \theta|$  < 0.93. It is required that all charged tracks originate from the interaction region defined by  $|V_z| < 10$  cm and  $|V_{xy}| < 1$ cm, where  $|V_z|$  and  $|V_{xu}|$  are the distances of closest approach of the charged track to the collision point in the beam direction and in the plane perpendicular to the beam, respectively.

Charged particles are identified by confidence levels for kaon and pion hypotheses calculated using  $dE/dx$ and TOF measurements. To effectively separate pions and kaons, a track is identified as a pion (or kaon) only if the confidence level for the pion (or kaon) hypothesis is larger than the confidence level for the kaon (or pion) hypothesis.

Photons are selected by exploiting the information from the EMC. It is required that the shower time be within 700 ns of the event start time and the shower energy be greater than 25 (50) MeV in the barrel (endcap) region defined by  $|\cos \theta|$  < 0.80 (0.86 <  $|\cos \theta|$  < 0.92). Here,  $\theta$  is the photon polar angle with respect to the beam direction.

In the selection of  $\gamma 2(\pi^+\pi^-)$ , background events from radiative Bhabha events in which at least two radiative photons are produced and one of them converts into an  $e^+e^-$  pair are suppressed by requiring the opening angle of any  $\pi^+\pi^-$  combination be larger than 10°. For the selection of  $\gamma K^+K^-$ , the background events of  $e^+e^- \rightarrow \gamma e^+e^-$  are suppressed by requiring  $E_{EMC} < 1$  GeV and  $E_{EMC}/p_{MDC} < 0.8$  for each charged kaon, where  $E_{EMC}$  and  $p_{MDC}$  are the energy deposited in the EMC and the momentum measured by the MDC, respectively.

In each event, there may be several different charged and/or neutral track combinations which satisfy the selection criteria for each light hadron final state. Each combination is subjected to a 4C kinematic fit for the hypotheses of  $\psi(3686)/\psi(3770) \to \gamma(2\pi^+\pi^-), \gamma K^+K^-\pi^+\pi^-, \gamma(3\pi^+\pi^-)$  and  $\gamma K^+K^-$ . For each final state, if more than one combination satisfies the selection criteria, only the combination with the least  $\chi^2_{4C}$  is retained, where  $\chi^2_{4C}$  is the chi-square of the 4C kinematic fit. The final states with  $\chi^2_{4C}$  < 25 are kept for further analysis.

To identify the  $\chi_{cJ}$  decays, we examine the invariant mass spectra of the light hadron final states. Figure [1](#page-4-0) shows the corresponding mass spectra for the  $\psi(3686)$  data, in which clear  $\chi_{c0}$ ,  $\chi_{c1}$  and  $\chi_{c2}$  signals are observed. Since the  $\chi_{c1}$  cannot decay into two pseudoscalar mesons because of spin-parity conservation, the  $\chi_{c1}$  signal cannot be observed in the  $K^+K^-$  invariant mass spectrum. By fitting these spectra separately, we obtain the numbers of  $\chi_{cJ}$  observed from the  $\psi(3686)$  data,  $N_{\psi(3686)}$ , which are summarized in Table [1.](#page-8-0) In the fits, the  $\chi_{cJ}$  signals are described by the MC simulated line-shapes convoluted by Gaussian functions for the resolution. Backgrounds in the four channels are described by  $3/3/3/1$ -parameter polynomial functions. The parameters of the convoluted Gaussian functions and the Chebychev polynomial functions are all free.



<span id="page-4-0"></span>Figure 1: Invariant mass spectra of the (a)  $2(\pi^+\pi^-)$ , (b)  $K^+K^-\pi^+\pi^-$ , (c)  $3(\pi^+\pi^-)$  and (d)  $K^+K^-$  combinations for the  $\psi(3686)$  data. The dots with error bars are for data and the blue solid lines are the fit results. The red dashed lines are the fitted backgrounds. The red, pink and blue arrows show the  $\chi_{c0}$ ,  $\chi_{c1}$  and  $\chi_{c2}$  nominal masses, respectively.

Figure [2](#page-6-0) shows the corresponding mass spectra for the  $\psi(3770)$  data, in which clear peaks can be observed for the  $\chi_{c0}$  decays. Fitting to these spectra similarly, we obtain the number of  $\chi_{cJ}(J=0,1)$  decays observed from the  $\psi(3770)$  data,  $N_{\psi(3770)}$ , which are summarized in Table [1.](#page-8-0) Due to the limited statistics, the decay  $\psi(3770) \rightarrow \gamma \chi_{c2}$  is not further considered in this analysis. The means and widths of the convoluted Gaussian functions for the  $\chi_{c0}$  signals are left free. For the  $\chi_{c1}$ , the mean and width of the convoluted Gaussian functions are fixed at the values taken from the fits to the  $\psi(3686)$  data. Backgrounds in the four channels are described by 6/2/6/2-parameter polynomial functions.

The background events from  $e^+e^- \to (\gamma_{\rm ISR})\psi(3686)$  produced near  $\sqrt{s} = 3.773$  GeV have the same event topologies as those from  $\psi(3770)$  decays and are indistinguishable from  $\psi(3770)$  decays. In the fits to the  $\psi(3770)$  data, the size and line-shape of such backgrounds are fixed according to MC simulations, with the numbers of background events being determined by

<span id="page-5-0"></span>
$$
N_{\chi_{cJ}}^{\mathrm{b}} = \sigma_{\psi(3686)}^{\chi_{cJ}, LH, \mathrm{obs}} \cdot \mathcal{L}_{\psi(3770)} \cdot \eta,
$$
\n(1)

where  $\mathcal{L}_{\psi(3770)}$  is the integrated luminosity of the  $\psi(3770)$  data,  $\sigma_{\psi(3686)}^{\chi_{cJ},LH,obs}$  is the observed cross section of  $e^+e^- \to \psi(3686) \to \gamma \chi_{cJ}$  with  $\chi_{cJ} \to LH$ , in which LH denotes  $2(\pi^+\pi^-)$ ,  $K^+K^-\pi^+\pi^-$ ,  $3(\pi^+\pi^-)$  and  $K^+K^-$ . In this work, we assume that there is no other effect affecting the  $\psi(3686)$  and  $\psi(3770)$  production in the energy range from 3.73 to 3.89 GeV. The variable  $\eta$  represents the rate of misidentifying  $\psi(3686)$ decays as  $\psi(3770)$  decays, which is obtained by analyzing  $1.5 \times 10^6$  MC events of  $\psi(3686) \to \gamma \chi_{cJ}$  with  $\chi_{cJ} \to LH$  generated at  $\sqrt{s} = 3.773$  GeV. The observed cross section for  $\psi(3686) \to \gamma \chi_{cJ}$  with  $\chi_{cJ} \to LH$ at a center-of-mass energy of  $\sqrt{s}$  is given by

$$
\sigma_{\psi(3686)}^{\chi_{cJ},LH,obs} = \int \sigma_{\psi(3686)}^{\chi_{cJ},LH}(s')f(s')F(x,s)G(s,s'')ds''dx.
$$
\n(2)

where  $s' \equiv s(1-x)$  is the square of the actual center-of-mass energy of the  $e^+e^-$  after radiating photon(s), x is the fraction of the radiative energy to the beam energy;  $f(s')$  is the phase space factor,  $(E_\gamma(s')/E_\gamma^0)^3$ , in which  $E_{\gamma}(s')$  and  $E_{\gamma}^{0}$  are the photon energies in  $\psi(3686) \to \gamma \chi_{cJ}$  transition at  $\sqrt{s'}$  and at the  $\psi(3686)$  mass, respectively;  $F(x, s)$  is the sampling function describing the radiative photon energy fraction x at  $\sqrt{s}$  [\[18](#page-10-17)];  $G(s, s'')$  is a Gaussian function describing the distribution of the collision energy with an energy spread  $\sigma_E = 1.37$  MeV as achieved at BEPCII;  $\sigma_{\psi(3686)}^{XcJ, LH}(s')$  is the cross section described by the Breit-Wigner function

$$
\sigma_{\psi(3686)}^{\chi_{cJ},LH}(s') = \frac{12\pi\Gamma_{\psi(3686)}^{\text{ee}}\Gamma_{\psi(3686)}^{\text{tot}}B_{\psi(3686)}^{\chi_{cJ},LH}}{(s'^2 - M_{\psi(3686)}^2)^2 + (\Gamma_{\psi(3686)}^{\text{tot}}M_{\psi(3686)})^2},\tag{3}
$$

 $x, \bar{x}$ 

in which  $\Gamma^{ee}_{\psi(3686)}$  and  $\Gamma^{tot}_{\psi(3686)}$  are, respectively, the leptonic width and total width of the  $\psi(3686)$ ,  $M_{\psi(3686)}$ is the  $\psi(3686)$  mass,  $\mathcal{B}_{\psi(3686)}^{\chi_{cJ},LH}$  is the combined branching fraction of  $\psi(3686) \to \gamma \chi_{cJ}$  with  $\chi_{cJ} \to LH$ . Here, the upper limit of x is set at  $1 - m_{\chi_{cJ}}^2/s$ , where  $m_{\chi_{cJ}}$  is the  $\chi_{cJ}$  nominal mass. We determine the branching fraction  $\mathcal{B}_{\psi(3686)}^{\chi_{cJ},LH}$  by dividing the number of  $\chi_{cJ}$  decays of  $\psi(3686)$  by the total number of  $\psi(3686)$  and by the corresponding efficiency obtained in this work. The rates  $\eta$  of misidentifying  $\psi(3686) \to \gamma \chi_{c0/1/2}$  as  $\psi(3770) \rightarrow \gamma \chi_{c0/1/2}$  are estimated to be  $4.72/6.40/7.60 \times 10^{-4}$ ,  $4.40/6.27/7.57 \times 10^{-4}$ ,  $3.53/4.95/6.14 \times 10^{-4}$ and  $6.56/-/11.02 \times 10^{-4}$  for  $\chi_{c0/1/2} \to 2(\pi^+\pi^-)$ ,  $K^+K^-\pi^+\pi^-$ ,  $3(\pi^+\pi^-)$  and  $K^+K^-$ , respectively. These lead to the number of background events from  $e^+e^- \rightarrow (\gamma_{\rm ISR})\psi(3686)$  to be  $90.6 \pm 3.4/37.5 \pm 1.4/76.5 \pm 2.9$ ,  $70.0 \pm 2.7/23.5 \pm 0.9/51.0 \pm 1.9$ ,  $56.6 \pm 2.2/39.7 \pm 1.5/73.5 \pm 2.8$  and  $34.9 \pm 1.3/-/11.1 \pm 0.4$  for  $\psi(3770) \rightarrow$  $\gamma \chi_{c0/1/2}$  with  $\chi_{c0/1/2} \to 2(\pi^+\pi^-)$ ,  $K^+K^-\pi^+\pi^-$ ,  $3(\pi^+\pi^-)$  and  $K^+K^-$  decays, respectively. The errors arise from uncertainties in the  $\psi(3686)$  resonance parameters, the integrated luminosity of the  $\psi(3770)$  data  $\mathcal{L}_{\psi(3770)}$  and the misidentification rates  $\eta$ . In Eq. [\(1\)](#page-5-0), the number of background events depends on the ratio of the misidentification rate  $\eta$  over the efficiency  $\epsilon_{\psi(3686)}$  of reconstructing  $\psi(3686) \to \chi_{cJ}$ . Since  $\eta$ and  $\epsilon_{\psi(3686)}$  all contain the simulation of  $\chi_{cJ} \to LH$ , a possible systematic uncertainty from the simulation of  $\chi_{cJ} \to LH$  is canceled here.



<span id="page-6-0"></span>Figure 2: Invariant mass spectra of the (a)  $2(\pi^+\pi^-)$ , (b)  $K^+K^-\pi^+\pi^-$ , (c)  $3(\pi^+\pi^-)$  and (d)  $K^+K^-$  combinations for the  $\psi(3770)$  data. The dots with error bars are data and the blue solid lines are the fit results. the red solid lines are the fitted combinatorial backgrounds. The red dashed lines are the sums of the peaking and fitted combinatorial backgrounds. The red, pink and blue arrows show the  $\chi_{c0}$ ,  $\chi_{c1}$  and  $\chi_{c2}$  nominal masses, respectively.

# 4. Results

The ratio of the branching fraction for  $\psi(3770) \rightarrow \gamma \chi_{cJ}$  divided by the branching fraction for  $\psi(3686) \rightarrow$  $\gamma \chi_{cJ}$  is determined channel by channel as

$$
R_{cJ} = \frac{\mathcal{B}[\psi(3770) \to \gamma \chi_{cJ}]}{\mathcal{B}[\psi(3686) \to \gamma \chi_{cJ}]} = \frac{N_{\psi(3770)} \cdot N_{\psi(3686)}^{\text{tot}} \cdot \epsilon_{\psi(3686)}}{N_{\psi(3686)} \cdot N_{\psi(3770)}^{\text{tot}} \cdot \epsilon_{\psi(3770)}},
$$
(4)

where  $N_{\psi(3686)}$  and  $N_{\psi(3770)}$  are the numbers of  $\chi_{cJ}$  observed from the  $\psi(3686)$  and  $\psi(3770)$  data,  $N_{\psi(3686)}^{\text{tot}}$ and  $N_{\psi(3770)}^{\rm tot}$  are the total numbers of  $\psi(3686)$  and  $\psi(3770)$  decays,  $\epsilon_{\psi(3686)}$  and  $\epsilon_{\psi(3770)}$  are the efficiencies of reconstructing  $\psi(3686)$  and  $\psi(3770) \to \gamma \chi_{cJ}$  with  $\chi_{cJ} \to LH$  estimated by MC simulations, respectively. Here,  $N^{\text{tot}}_{\psi(3770)}$  is determined by  $\sigma^{\text{obs}}_{\psi(3770)}$ .  $\mathcal{L}_{\psi(3770)}$ , where  $\sigma^{\text{obs}}_{\psi(3770)} = (7.15 \pm 0.27 \pm 0.27)$  nb is the cross section for  $\psi(3770)$  production [\[21](#page-10-20), [22](#page-10-21), [23\]](#page-10-22) and  $\mathcal{L}_{\psi(3770)}$  is the integrated luminosity of the  $\psi(3770)$  data set [\[12](#page-10-11)].

Table [1](#page-8-0) summarizes the ratios  $R_{cJ}$  measured via the different channels. The results are consistent within statistical uncertainties. From these measurements, we obtain the statistical-weighted averages  $\bar{R}_{c0} =$  $(6.89 \pm 0.28 \pm 0.65)\%$  and  $\bar{R}_{c1} = (2.03 \pm 0.44 \pm 0.66)\%$ , where the first uncertainty is statistical and the second systematic.

In the measurements of  $\bar{R}_{c0/1}$ , the systematic uncertainty arises from the uncertainties in the total number  $(0.81\%)$  of  $\psi(3686)$  decays  $(N_{\psi(3686)}^{tot}$  [\[13\]](#page-10-12)); the integrated luminosity  $(1.0\%)$  of the  $\psi(3770)$  data  $(L_{\psi(3770)}$ [\[12](#page-10-11)]); the cross section (5.3%) for  $\psi(3770)$  ( $\sigma_{\psi(3770)}^{\text{obs}}$  [\[21](#page-10-20), [22,](#page-10-21) [23\]](#page-10-22)); the photon selection (1.4%), assigned based on 1.0% per photon [\[24\]](#page-10-23); the MDC tracking  $(2.6\%/4.0\%)$ ; the particle identification  $(2.6\%/4.0\%)$ ; the statistical uncertainty  $(1.0\%)$  of the efficiency due to the size of the simulated event sample; the 4C kinematic fit (1.0%), estimated by comparing the measurements with and without the kinematic fit correction; the fit to mass spectra  $(6.4\%/31.5\%)$ , estimated by comparing the measurements with alternative fit ranges ( $\pm 20$ )  $\text{MeV}/c^2$ ), signal shape (simple Breit-Wigner function) and background shapes ( $\pm 1$  order of the polynomial functions); and the subtraction of  $\psi(3686)$  peaking background  $(0.5\%/2.0\%)$ . The efficiencies of the MDC tracking and particle identification for  $K^+$  or  $\pi^+$  are examined by the doubly tagged hadronic  $D\bar{D}$  events. The difference between the efficiencies of data and MC is assigned as an uncertainty. Then, their effects on  $\bar{R}_{c0/1}$  are estimated to be [2](#page-8-1).6%/4.0%. Table 2 summarizes these uncertainties. Adding them in quadrature, we obtain the total systematic uncertainty for  $\bar{R}_{c0/1}$  to be 9.4%/32.6%.

Multiplying  $\bar{R}_{cJ}$  by the branching fraction  $\mathcal{B}[\psi(3686) \to \gamma \chi_{cJ}]$  (and the total width  $\Gamma_{\psi(3770)}^{\text{tot}}$ ) taken from the PDG [\[11\]](#page-10-10), we obtain the branching fractions (and the partial widths) for  $\psi(3770) \to \gamma \chi_{cJ}$ , which are summarized in Table [3,](#page-10-24) where the first uncertainty is statistical and the second systematic. In the measurement of  $\mathcal{B}[\psi(3770) \to \gamma \chi_{cJ}]$  (and  $\Gamma[\psi(3770) \to \gamma \chi_{cJ}]$ ), the systematic uncertainty arises from the uncertainties of  $\bar{R}_{c0/1}$  and the uncertainties of  $\mathcal{B}[\psi(3686) \to \gamma \chi_{c0/1}]$  of 2.7/3.2% (and the uncertainty of  $\Gamma_{\psi(3770)}^{\rm tot}$  of 3.7%).

$\chi_{cJ} \rightarrow LH\bar{l}$		$J=0$	$J=1$
	$N_{\psi(3770)}$	$756 \pm 51$	$80 \pm 26$
	$\epsilon_{\psi(3770)}$	$24.1 \pm 0.2$	$25.7 \pm 0.2$
$2(\pi^{+}\pi^{-})$	$N_{\psi(3686)}$	$59976 \pm 318$	$19712 \pm 175$
	$\epsilon_{\psi(3686)}$	$24.9 \pm 0.2$	$26.5 \pm 0.2$
	$R_{cJ}$	$6.64 \pm 0.45$	$2.13 \pm 0.69$
	$N_{\psi(3770)}$	$716 \pm 54$	$46 \pm 24$
	$\epsilon_{\psi(3770)}$	$24.0 \pm 0.2$	$25.4 \pm 0.2$
$K^+K^-\pi^+\pi^-$	$N_{\psi(3686)}$	$46929 \pm 240$	$11576 \pm 115$
	$\epsilon_{\psi(3686)}$	$23.3 \pm 0.2$	$24.9 \pm 0.2$
	$R_{cJ}$	$7.56 \pm 0.57$	$2.00 \pm 1.04$
	$N_{\psi(3770)}$	$502 \pm 54$	$76 \pm 27$
	$\epsilon_{\psi(3770)}$	$18.5 \pm 0.2$	$20.0 \pm 0.2$
$3(\pi^{+}\pi^{-})$	$N_{\psi(3686)}$	$36536 \pm 237$	$19593 \pm 153$
	$\epsilon_{\psi(3686)}$	$18.1 \pm 0.2$	$19.6 \pm 0.2$
	$R_{cJ}$	$6.86 \pm 0.74$	$1.94 \pm 0.69$
$K^+K^-$	$N_{\psi(3770)}$	$283 \pm 24$	
	$\epsilon_{\psi(3770)}$	$32.5 \pm 0.2$	
	$N_{\psi(3686)}$	$21452 \pm 154$	
	$\epsilon_{\psi(3686)}$	$32.1 \pm 0.2$	
	$R_{cJ}$	$6.65 \pm 0.57$	
Averaged $R_{cJ}$		$6.89 \pm 0.28$	$2.03 \pm 0.44$

<span id="page-8-0"></span>Table 1: Measured  $R_{cJ}$  (%), where  $N_{\psi(3770)}$  and  $N_{\psi(3686)}$  are the (peaking background corrected) numbers of  $\chi_{cJ}$ observed from the  $\psi(3770)$  and  $\psi(3686)$  data,  $\epsilon_{\psi(3770)}$  and  $\epsilon_{\psi(3686)}$  are the detection efficiencies (%). The uncertainties are statistical only.

Table 2: Systematic uncertainties (%) in the measurements of  $\bar{R}_{cJ}$ .

<span id="page-8-1"></span>

	$R_{c0}$	$R_{c1}$
tot $\psi(3686)$ $\left[13\right]$	0.81	0.81
$\sigma^{\rm obs}_{\psi(3770)}$ [21, 22, 23]	5.3	5.3
$L_{\psi(3770)}$ [12]	1.0	1.0
MC statistics	1.0	1.0
Photon selection	1.4	1.4
MDC tracking	2.6	4.0
Particle identification	2.6	4.0
$4C$ kinematic fit	1.0	1.0
Fit to mass spectra	6.4	31.5
Background subtraction	0.5	2.0
Total	9.4	32.6

## 5. Summary

In summary, by analyzing 2.92 fb<sup>-1</sup> of  $e^+e^-$  collision data taken at  $\sqrt{s} = 3.773$  GeV and  $106.41 \times 10^6$ .  $\psi(3686)$  decays taken at  $\sqrt{s} = 3.686$  GeV with the BESIII detector at the BEPCII collider, we measure the branching fraction  $\mathcal{B}(\psi(3770) \to \gamma \chi_{c0}) = (6.88 \pm 0.28 \pm 0.67) \times 10^{-3}$  and the partial width  $\Gamma[\psi(3770) \to$  $\gamma \chi_{c0}$  = (187 ± 8 ± 19) keV. These are obtained by first measuring the ratio with respect to the accurately known branching fraction for  $\psi(3686) \rightarrow \gamma \chi_{cJ}$  decays. Our results are, thereby, not influenced by the uncertainties in the branching fractions of  $\chi_{cJ}$  decays to light hadrons as done in Ref. [\[7\]](#page-10-5). The branching fraction and partial width for  $\psi(3770) \to \gamma \chi_{c1}$  measured in this work are consistent with our previous measurement [\[10\]](#page-10-9) within errors. Table [3](#page-10-24) compares the  $\Gamma[\psi(3770) \to \gamma \chi_{c0/1}]$  measured at BESIII with those measured by CLEO  $[7, 8]$  $[7, 8]$ <sup>[1](#page-9-1)</sup> and the theoretical calculations from Refs.  $[2, 3, 4, 5, 6]$  $[2, 3, 4, 5, 6]$  $[2, 3, 4, 5, 6]$  $[2, 3, 4, 5, 6]$  $[2, 3, 4, 5, 6]$  $[2, 3, 4, 5, 6]$  $[2, 3, 4, 5, 6]$  $[2, 3, 4, 5, 6]$ . The partial width  $\Gamma[\psi(3770) \to \gamma \chi_{c0}]$  measured at BESIII is consistent within errors with the one measured by CLEO with an improved precision. These results underline the fact that the traditional models [\[3](#page-10-1), [4](#page-10-2), [5\]](#page-10-3) with a relativistic assumption or a coupled-channel correction agree quantitatively better with the experimental data than those [\[2,](#page-10-0) [3,](#page-10-1) [4,](#page-10-2) [5](#page-10-3)] based upon non-relativistic calculations. For these traditional models, the non-relativistic calculations clearly overestimate the partial width  $\Gamma[\psi(3770) \to \gamma \chi_{cJ}]$ . The experimental data also support the recent calculation based on the non-relativistic constituent quark model (NRCQM) [\[6\]](#page-10-4). Together with further theoretical developments, our results aim to contribute to a deeper understanding of the dynamics of charmonium decays above the open-charm threshold.

## 6. Acknowledgements

The BESIII collaboration thanks the staff of BEPCII and the IHEP computing center for their strong support. This work is supported in part by National Key Basic Research Program of China under Contract Nos. 2009CB825204 and 2015CB856700; National Natural Science Foundation of China (NSFC) under Contracts Nos. 10935007, 11125525, 11235011, 11305180, 11322544, 11335008, 11425524; the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; Joint Large-Scale Scientific Facility Funds of the NSFC and CAS under Contracts Nos. 11179007, U1232201, U1332201; CAS under Contracts Nos. KJCX2- YW-N29, KJCX2-YW-N45; 100 Talents Program of CAS; the CAS Center for Excellence in Particle Physics (CCEPP); INPAC and Shanghai Key Laboratory for Particle Physics and Cosmology; German Research Foundation DFG under Contract No. Collaborative Research Center CRC-1044; Istituto Nazionale di Fisica Nucleare, Italy; Ministry of Development of Turkey under Contract No. DPT2006K-120470; Russian Foundation for Basic Research under Contract No. 14-07-91152; U. S. Department of Energy under Contracts Nos. DE-FG02-04ER41291, DE-FG02-05ER41374, DE-FG02-94ER40823, DESC0010118; U.S. National Science Foundation; University of Groningen (RuG) and the Helmholtzzentrum fuer Schwerionenforschung GmbH (GSI), Darmstadt; WCU Program of National Research Foundation of Korea under Contract No. R32-2008-000-10155-0.

#### References

<span id="page-9-0"></span>[1] M. B. Voloshin, Prog. Part. Nucl. Phys. 61 (2008) 455; E. Eichten et al., Rev. Mod. Phys. 80 (2008) 1161; N. Brambilla et al., Eur. Phys. J. C 71 (2011) 1534.

<span id="page-9-1"></span><sup>&</sup>lt;sup>1</sup> The CLEO measurements were based on the total width  $\Gamma_{\psi(3770)}^{\text{tot}} = (23.6 \pm 2.7)$  MeV [\[25\]](#page-10-25) and the cross section  $\sigma_{\psi(3770)\to D\bar{D}}^{\rm obs} = (6.39 \pm 0.10_{-0.08}^{+0.17})$  nb for determining the total number of  $\psi(3770)$  decays, where the  $\psi(3770) \to \text{non-}D\bar{D}$ decays were neglected. In addition, CLEO [\[7](#page-10-5)] cited  $\mathcal{B}(\psi(3686) \to \gamma \chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$  and  $\mathcal{B}(\psi(3686) \to \gamma \chi_{c1}) =$  $(9.07 \pm 0.11 \pm 0.54)\%$  to determine  $\mathcal{B}(\psi(3770) \to \gamma \chi_{cJ})$  from Ref. [\[26\]](#page-10-26). Although CLEO determined the branching fraction of  $\psi(3770) \to \text{non-}D\bar{D}$  decays to be  $(-3.3 \pm 1.4^{+4.8}_{-6.6})\%$  and set an upper limit of 9% at 90% confidence level [\[27](#page-10-8)], the PDG value for  $\Gamma_{\psi(3770)}^{D\bar{D}}/\Gamma_{\psi(3770)}^{\text{tot}}$  [\[28](#page-10-27)] from four measurements at BESII [\[22,](#page-10-21) [29,](#page-10-28) [30](#page-10-29), [31](#page-10-30)] implied the branching fraction for  $\psi(3770) \to \text{non-}D\bar{D}$ decays to be  $(14.7 \pm 3.2)$ %. At present, the PDG value for  $\Gamma_{\psi(3770)}^{D\bar{D}}/\Gamma_{\psi(3770)}^{\text{tot}}$  gives the branching fraction of  $\psi(3770) \to \text{non-}D\bar{D}$ decays to be  $(7\frac{+8}{-9})\%$  [\[11](#page-10-10)]. Therefore, for better comparisons, we also list the corrected CLEO partial widths with the same input values as those used in our measurements, which are  $\sigma_{\psi(3770)}^{\text{obs}} = (7.15 \pm 0.27 \pm 0.27)$  nb [\[21](#page-10-20), [22](#page-10-21), [23](#page-10-22)],  $\Gamma_{\psi(3770)}^{\text{tot}} = (27.2 \pm 1.0)$ MeV,  $\mathcal{B}(\psi(3686) \to \gamma \chi_{c0}) = (9.99 \pm 0.27)\%$  and  $\mathcal{B}(\psi(3686) \to \gamma \chi_{c1}) = (9.55 \pm 0.31)\%$  [\[11](#page-10-10)].

<span id="page-10-24"></span>Table 3: Comparisons of the partial widths for  $\psi(3770) \to \gamma \chi_{cJ}$  (in keV), where B and Γ denote the branching fraction and the partial width for  $\psi(3770) \to \gamma \chi_{cJ}$ , respectively. For the BESIII results, the first uncertainty is statistical and the second systematic. Detailed explanations about the CLEO results can be found in footnote 1.

Experiments	$J=0$	$J=1$
$\mathcal{B}^{\text{BESIII}}(\times 10^{-3})$	$6.88 \pm 0.28 \pm 0.67$	$1.94 \pm 0.42 \pm 0.64$
$\mathcal{B}^{\text{BESIII}}(\times 10^{-3})$ [10]		$2.48 \pm 0.15 \pm 0.23$
$T$ BESIII	$187 \pm 8 \pm 19$	$53 \pm 12 \pm 18$
$\Gamma$ <sup>BESIII</sup> [10]		$67.5 \pm 4.1 \pm 6.7$
$\Gamma^{\mathrm{CLEO}}$ [7, 8]	$172 \pm 30$	$70 \pm 17$
$\Gamma^{\rm CLEO}$ corrected	$192 \pm 24$	$72 \pm 16$
Theories		
Rosner $[2]$ (non-relativistic)	$523 \pm 12$	$73 \pm 9$
$Ding-Qing-Chao$ [3]		
non-relativistic	312	95
relativistic	199	72
Eichten-Lane-Quigg [4]		
non-relativistic	254	183
with coupled channels corrections	225	59
Barnes-Godfrey-Swanson [5]		
non-relativistic	403	125
relativistic	213	77
$NRCQM$ [6]	218	70

- <span id="page-10-0"></span>[2] J. L. Rosner, Phys. Rev. D  $64$  (2001) 094002; Ann. Phys. (N.Y.)  $319$  (2005) 1.<br>[3] Y.-B. Ding. D.-H. Qin and K.-T. Chao. Phys. Rev. D  $44$  (1991) 3562.
- <span id="page-10-1"></span>Y.-B. Ding, D.-H. Qin and K.-T. Chao, Phys. Rev. D 44 (1991) 3562.
- <span id="page-10-2"></span>[4] E. J. Eichten, K. Lane and C. Quigg, Phys. Rev. D 69 (2004) 094019.
- <span id="page-10-3"></span>[5] T. Barnes, S. Godfrey and E. S. Swanson, Phys. Rev. D 72 (2005) 054026.
- <span id="page-10-4"></span>[6] W.-J. Deng, L.-Y. Xiao, L.-C. Gui, X.-H. Zhong, arXiv:1510.08269 [hep-ph].
- <span id="page-10-5"></span>[7] CLEO Collaboration, B. A. Briere *et al.*, Phys. Rev. D **74** (2006) 031106.
- <span id="page-10-6"></span>[8] CLEO Collaboration, T. E. Coans et al., Phys. Rev. Lett. **96** (2006) 182002.
- <span id="page-10-7"></span>[9] T. Skwarnicki, Int. J. Mod. Phys. A 19 (2004) 1030.
- <span id="page-10-9"></span>[10] BESIII Collaboration, M. Ablikim et al., Phys. Rev. D 91 (2015) 092009.
- <span id="page-10-10"></span>[11] Particle Data Group, K. A. Olive et al., Chinese Physics C  $38$  (2014) 090001.
- <span id="page-10-11"></span>[12] BESIII Collaboration, M. Ablikim et al., Chinese Physics C  $37$  (2013) 123001.
- <span id="page-10-12"></span>[13] BESIII Collaboration, M. Ablikim et al., Chinese Physics C 37 (2013) 063001.
- <span id="page-10-13"></span>[14] BESIII Collaboration, M. Ablikim et al., Nucl. Instrum. Meth. A 614 (2010) 345.
- <span id="page-10-14"></span>[15] GEANT4 Collaboration, S. Agostinelli et al., Nucl. Instrum. Meth. A 506 (2003) 250.
- <span id="page-10-15"></span>[16] D. J. Lange, Nucl. Instrum. Meth. A 462 (2001) 152; R. G. Ping, Chinese Physics C 32 (2008) 599.
- <span id="page-10-16"></span> $(17)$  S. Jadach, B. F. L. Ward and Z. Was, Comp. Phys. Commu. 130 (2000) 260; Phys. Rev. D 63 (2001) 113009.
- <span id="page-10-17"></span>[18] E. A. Kureav and V. S. Fadin, Sov. J. Nucl. Phys. 41 (1985) 466, Yad. Fiz. 41 (1985) 733.
- <span id="page-10-18"></span>[19] E. Barberio, Z. Was, Comput. Phys. Commun. 79 (1994) 291.
- <span id="page-10-19"></span>[20] J. C. Chen et al., Phys. Rev. D 62 (2000) 034003.
- <span id="page-10-20"></span>[21] BES Collaboration, M. Ablikim *et al.*, Phys. Lett. B  $650$  (2007) 111.<br>[22] BES Collaboration, M. Ablikim *et al.*, Phys. Rev. Lett. **97** (2006) 12
- <span id="page-10-21"></span>BES Collaboration, M. Ablikim et al., Phys. Rev. Lett. 97 (2006) 121801.
- <span id="page-10-22"></span>[23] BES Collaboration, M. Ablikim et al., Phys. Lett. B **652** (2007) 238.
- <span id="page-10-23"></span>[24] BESIII Collaboration, M. Ablikim et al., Chinese Physics C  $34$  (2010) 421; Phys. Rev. D  $81$  (2010) 052005; Phys. Rev. Lett. 105 (2010) 261801; Phys. Rev. D 83 (2011) 012003; Phys. Rev. D 83 (2011) 012006; Phys. Rev. D 83 (2011) 032003.
- <span id="page-10-25"></span>[25] Particle Data Group, S. Eidelman et al., Phys. Lett. B 592 (2004) 1.
- <span id="page-10-26"></span>[26] CLEO Collaboration, S. B. Athar et al., Phys. Rev. D **70** (2004) 112002.
- <span id="page-10-8"></span>[27] CLEO Collaboration, D. Besson et al., Phys. Rev. Lett.  $96$  (2006) 092002, 104 (2010) 159901E.
- <span id="page-10-27"></span>[28] Particle Data Group, C. Amsler *et al.*, Phys. Lett. B **661** (2008) 1.
- <span id="page-10-28"></span>[29] BES Collaboration, M. Ablikim et al., Phys. Lett. B **641** (2006) 145.
- <span id="page-10-29"></span>[30] BES Collaboration, M. Ablikim et al., Phys. Rev. D **76** (2007) 122002.
- <span id="page-10-30"></span>[31] BES Collaboration, M. Ablikim et al., Phys. Lett. B 659 (2008) 74.