

Integrable orbit equivalence rigidity for free groups

Lewis Bowen*
University of Texas at Austin

January 7, 2018

Abstract

It is shown that every accessible group which is integrable orbit equivalent to a free group is virtually free. Moreover, we also show that any integrable orbit-equivalence between finitely generated groups extends to their end compactifications.

Contents

1	Introduction	1
1.1	Accessible groups	2
1.2	Strict integrable embeddings	2
2	Preliminaries	3
3	The Space of Ends	4

1 Introduction

Measure equivalence (ME) is an equivalence relation on groups, introduced by M. Gromov [Gro93] as a measure-theoretic counterpart to quasi-isometry. Most of the research in this area has focussed on rigidity phenomena. For example, Furman proved [Fur99a, Fur99b] that any group ME to a lattice in a higher rank simple Lie group has a finite normal subgroup whose quotient is commensurable to a lattice in the same Lie group. See [Fur11] for a survey of further results.

Here we consider the class of free groups. This class is far from rigid: there is a large variety of groups measure-equivalent to a free group [Gab05] and we do not even have a conjectural classification of such groups. So it makes sense to consider the more restrictive

*supported in part by NSF grant DMS-1500389, NSF CAREER Award DMS-0954606

notion of measure-equivalence known as **integrable orbit equivalence** (IOE) or L^1 -orbit-equivalence (L^1 -OE) which takes into account the metric in addition to measure-theoretic structure. This notion is stricter than integrable measure equivalence (IME) (also called L^1 -measurable equivalence) that first appeared in [BFS13] where it was shown that any group that is L^1 -ME to a lattice in $SO(n, 1)$ ($n \geq 3$) has a finite normal subgroup whose quotient is a lattice in $SO(n, 1)$. Another milestone is T. Austin's work proving L^1 -ME rigidity for nilpotent groups [Aus13] (see also M. Cantrell's recent strengthening of Austin's results [Can15]).

The main result of this paper is that any finitely generated accessible group that admits a strict integrable embedding into a free group is virtually free. In particular, any finitely generated accessible group that is IOE to a free group is virtually free. These terms are defined next.

1.1 Accessible groups

According to Stallings' Ends Theorem [Sta68], if a finitely generated group Γ has more than one end then it splits as either a nontrivial free product with amalgamation or as an HNN extension over a finite subgroup. If such splittings cannot occur indefinitely, then the group is called *accessible*. C.T.C. Wall conjectured [Wal71] that all finitely generated groups are accessible. A counterexample was obtained by Dunwoody [Dun93]. However, Dunwoody showed that all finitely presented groups are accessible [Dun85].

1.2 Strict integrable embeddings

Let Γ, G be finitely generated groups. Intuitively, a strict integrable embedding of Γ into G is a random map from Γ into G that is 'Lipschitz on average' and has a bounded number of preimages. To be precise, fix a finite symmetric generating set S_G of G and define the **word length** of any $g \in G$ by $|g|_G := n$ where n is the smallest natural number such that there exist $s_1, \dots, s_n \in S_G$ with $g = s_1 \cdots s_n$.

Given an action of Γ on a set X , a **cocycle** into G is a map $\alpha : \Gamma \times X \rightarrow G$ such that

$$\alpha(g_1 g_2, x) = \alpha(g_1, g_2 x) \alpha(g_2, x) \quad \forall g_1, g_2 \in \Gamma, x \in X.$$

In the case of concern, X is endowed with a probability measure μ , the action $\Gamma \curvearrowright (X, \mu)$ is measure-preserving and α is measurable. Then we say that α is **integrable** if

$$\int |\alpha(g, x)|_G d\mu(x) < \infty$$

for every $g \in \Gamma$. While the precise value of $\int |\alpha(g, x)|_G d\mu(x)$ depends on the generating set S_G , its finiteness does not and therefore integrability of α does not depend on S_G .

We say that α is a **strict integrable embedding** if in addition to being integrable there is a constant $C > 0$ such that for every $h \in G$,

$$\#\{g \in \Gamma : \alpha(g, x) = h\} \leq C$$

for a.e. x . This notion is more restrictive than the notion of integrable embedding defined in the appendix of [Aus13].

Our main result is:

Theorem 1.1. *Let Γ be a finitely generated accessible group. If Γ admits a strict integrable embedding into a free group then Γ is virtually free.*

Definition 1. Two groups Γ, Λ are **integrably orbit equivalent** (IOE) if there exist probability measure-preserving essentially free ergodic actions $\Gamma \curvearrowright (X, \mu)$ and $\Lambda \curvearrowright (X, \mu)$ such that for a.e. $x \in X$, $\Gamma x = \Lambda x$ and the orbit cycles, defined by

$$\alpha : \Gamma \times X \rightarrow \Lambda, \quad \alpha(g, x)x = gx,$$

$$\beta : \Lambda \times X \rightarrow \Gamma, \quad \beta(h, x)x = hx$$

are integrable.

Clearly an IOE cocycle is a strict integrable embedding. So Theorem 1.1 implies that any finitely generated accessible group IOE to a free group is virtually free.

We do not know whether accessibility is a necessary condition nor whether ‘strict integrable embedding’ can be weakened to ‘integrable embedding’ or IOE weakened to IME.

Acknowledgements. This work was inspired from conversations with Tim Austin, Uri Bader, Alex Furman and Roman Sauer.

2 Preliminaries

Definition 2. If X is a connected locally connected σ -compact topological space then let

$$\text{End}(X) := \varprojlim_K \pi_0(X \setminus K).$$

To be precise, the inverse limit is over all compact subsets $K \subset X$ and $\pi_0(X \setminus K)$ denotes the set of noncompact connected components of $X \setminus K$. We give $\pi_0(X \setminus K)$ the discrete topology. If $K \subset L$ are compact subsets of X then there is a natural map from $\pi_0(X \setminus L)$ to $\pi_0(X \setminus K)$ and $\text{End}(X)$ is the inverse limit of this system (where the collection of compacts of X is ordered by inclusion). In particular, for every compact K , there is a natural map $\pi_K : \text{End}(X) \rightarrow \pi_0(X \setminus K)$.

The **end compactification of X** , denoted \overline{X} is the space $\overline{X} := X \cup \text{End}(X)$ with the following topology: every open subset of X is \overline{X} . Also, for every compact $K \subset X$ and $C \in \pi_0(X \setminus K)$, the set

$$C \cup \{\xi \in \text{End}(X) : \pi_K(\xi) \in C\} \subset \overline{X}$$

is open. These sets form a basis for the topology of \overline{X} .

Let Γ be a group with a finite generating set S . The **Cayley graph of** (Γ, S) , denoted $\text{Cay}(\Gamma, S)$, has vertex set Γ and edge set $\{(g, gs) : g \in \Gamma, s \in S\}$. We usually let $\bar{\Gamma}$ denote the end compactification $\overline{\text{Cay}(\Gamma, S)}$, leaving the generating set implicit.

It is well-known that any finitely generated group quasi-isometric to a free group is itself virtually free. This is usually attributed to Gromov via Stallings' Ends Theorem. Alternatively, it follows from Thomassen-Woess [TW93] that accessibility is a quasi-isometric-invariant and from Papasoglu-Whyte [PW02] that any accessible group quasi-isometric to a free group must be virtually free. Indeed, this implies more:

Theorem 2.1. *If Γ is the fundamental group of a finite graph of groups in which all vertex and edge groups are finite then Γ is virtually free.*

Proof. This follows from [PW02] although it may have been known earlier. \square

We also note:

Lemma 2.2. *If Γ is the fundamental group of a finite graph of groups in which all edge groups are finite and $\Gamma_v \leq \Gamma$ is a vertex subgroup then Γ_v is quasi-isometrically embedded in Γ .*

Finally, we introduce a notion of L^1 -embedding:

Definition 3. Let $\Gamma \curvearrowright (X, \mu)$ a probability measure-preserving action and $\alpha : \Gamma \times X \rightarrow G$ a measurable cocycle. We say α is an L^1 -embedding if

- α is L^1 : for every $g \in \Gamma$, $\int |\alpha(g, x)|_G d\mu(x) < \infty$ where $|\cdot|_G$ denotes length with respect to a fixed word metric on G ;
- there is a constant $C > 0$ such that for any $h \in G$,

$$\#\{g \in \Gamma : \alpha(g, x) = h\} \leq C$$

for a.e. x .

Remark 1. It is straightforward to check that a composition of L^1 -embeddings is an L^1 -embedding and that any cocycle arising from an L^1 -OE is an L^1 -embedding.

In order to prove Theorem 1.1, it now suffices to show the following. Let $\Gamma \curvearrowright (X, \mu)$ be a probability measure-preserving action and $\alpha : \Gamma \times X \rightarrow G$ an L^1 -embedding. Then Γ is not 1-ended. We will show this in the next section.

3 The Space of Ends

Theorem 3.1. *Let Γ, G be finitely generated groups, $\Gamma \curvearrowright (X, \mu)$ a probability measure-preserving action, $\alpha : \Gamma \times X \rightarrow G$ an L^1 -embedding. Define $\alpha' : \Gamma \times X \rightarrow G$ by*

$$\alpha'(g, x) = \alpha(g^{-1}, x)^{-1}.$$

Let $\bar{\Gamma}, \bar{G}$ denote the end-compactifications of Γ, G respectively with respect to fixed finite generating subsets. Then α' extends to a map, also denoted by α' from $\bar{\Gamma} \times X \rightarrow \bar{G}$ such that

- $\alpha'_{gx}(g\xi) = \alpha(g, x)\alpha'_x(\xi)$ (for $g \in G, \xi \in \bar{\Gamma}$ and a.e. $x \in X$);
- $\alpha'_x : \bar{\Gamma} \rightarrow \bar{G}$ is continuous for a.e. x .

Here, $\alpha'_x(g) := \alpha'(g, x)$.

Remark 2. The Theorem above implies that every finitely generated group is 1-taut relative to its space of ends, in the terminology of [BFS13]. We will not need this fact.

Theorem 3.1 follows immediately from the next two lemmas.

Lemma 3.2. *Let X, Y be connected locally connected σ -compact topological spaces. Let $\alpha : X \rightarrow Y$ be a continuous map. Assume that for every compact $K \subset Y$ there exists a compact $F \subset X$ such that $\alpha(X \setminus F) \subset Y \setminus K$ and α descends to a well defined map $\pi_0(X \setminus F) \rightarrow \pi_0(Y \setminus K)$. Then α extends continuously to $\bar{\alpha} : \bar{X} \rightarrow \bar{Y}$.*

Proof. For every $K \subset Y$ we have a map

$$\text{End}(X) \rightarrow \pi_0(X \setminus D) \rightarrow \pi_0(Y \setminus K),$$

so the lemma follows by the definition of the inverse limit $\varprojlim_D \pi_0(Y \setminus K)$. \square

Lemma 3.3. *For a.e. $x \in X$ and every finite set $K \subset G$ there exists a finite set $F \subset \Gamma$ (depending on x and K) such that $\alpha'_x(\Gamma \setminus F) \subset G \setminus K$ and α'_x descends to a map $\pi_0(\Gamma \setminus F) \rightarrow \pi_0(G \setminus K)$.*

Proof. Let S_Γ, S_G be finite generating sets for Γ, G respectively. Let $|\cdot|_\Gamma, |\cdot|_G$ denote word length on Γ, G respectively.

For each $h_1, h_2 \in G$, choose a geodesic segment $\gamma[h_1, h_2]$ from h_1 to h_2 . More precisely, for every integer $0 \leq n \leq |h_1^{-1}h_2|_G$, there is an element $\gamma[h_1, h_2](n) \in G$ so that

$$\gamma[h_1, h_2](n)^{-1}\gamma[h_1, h_2](n+1) \in S_G$$

if $n < |h_1^{-1}h_2|_G$ and $\gamma[h_1, h_2](0) = h_1, \gamma[h_1, h_2](|h_1^{-1}h_2|_G) = h_2$. Let us also require that this choice is left-invariant so that $h\gamma[h_1, h_2] = \gamma[hh_1, hh_2]$ for any $h, h_1, h_2 \in G$.

For each $x \in X, g \in \Gamma, s \in S_\Gamma$, we imagine an airplane flying from $\alpha'_x(g)$ to $\alpha'_x(gs)$. The path of the flight is the geodesic $\gamma[\alpha'_x(g), \alpha'_x(gs)]$. We call this an **s -flight**. For $k \in G$, we let $F_{s,k}(x)$ denote the set of elements $g \in \Gamma$ such that the s -flight from $\alpha'_x(g)$ to $\alpha'_x(gs)$ contains k . That is:

$$F_{s,k}(x) := \{g \in \Gamma : k \in \gamma[\alpha'_x(g), \alpha'_x(gs)]\}.$$

Claim 1. $F_{s,k}(x)$ is finite for a.e. x . In fact,

$$\int \#F_{s,k}(x) d\mu(x) \leq C \int |\alpha(s^{-1}, x)^{-1}|_G d\mu(x) < \infty$$

where $C > 0$ is the constant in the definition of L^1 -embedding.

Proof of Claim 1. It suffices to show that $\int \#F_{s,k}(x) d\mu(x) < \infty$. In order to prove this, let

$$L_{s,k} = \{(g, x) \in \Gamma \times X : g^{-1} \in F_{s,k}(x)\}.$$

Let c_Γ denote the counting measure on Γ . Then $\int \#F_{s,k} d\mu = c_\Gamma \times \mu(L_{s,k})$. Because the action $\Gamma \curvearrowright (X, \mu)$ is invariant,

$$c_\Gamma \times \mu(L_{s,k}) = c_\Gamma \times \mu(R_{s,k})$$

where $R_{s,k} = \{(g^{-1}, gx) : (g, x) \in L_{s,k}\}$. By definition

$$c_\Gamma \times \mu(R_{s,k}) = \int \#\{g \in \Gamma : (g, x) \in R_{s,k}\} d\mu(x).$$

However, $(g, x) \in R_{s,k}$ if and only if $(g^{-1}, gx) \in L_{s,k}$ if and only if $g \in F_{s,k}(gx)$ if and only if $k \in \gamma[\alpha'_{gx}(g), \alpha'_{gx}(gs)]$ if and only if

$$\alpha'_{gx}(g)^{-1}k \in \gamma[e, \alpha'_{gx}(g)^{-1}\alpha'_{gx}(gs)].$$

Let us now compute

$$\alpha'_{gx}(g)^{-1}\alpha'_{gx}(gs) = \alpha(g^{-1}, gx)\alpha(s^{-1}g^{-1}, gx)^{-1} = \alpha(s^{-1}, x)^{-1}$$

by the cocycle equation. So

$$\begin{aligned} \int \#F_{s,k} d\mu &= c_\Gamma \times \mu(R_{s,k}) = \int \#\{g \in \Gamma : \alpha'_{gx}(g)^{-1}k \in \gamma[e, \alpha(s^{-1}, x)^{-1}]\} d\mu(x) \\ &\leq C \int |\alpha(s^{-1}, x)^{-1}|_G d\mu(x) < \infty. \end{aligned}$$

□

Now let $K \subset G$ be finite and define

$$F_K(x) := \bigcup \{F_{s,k} : s \in S_\Gamma, k \in K\}.$$

To finish the proof of the lemma, it suffices to show that if $g_1, g_2 \in \Gamma$ are in the same connected component of $\Gamma \setminus F_K(x)$ then $\alpha'_x(g_1), \alpha'_x(g_2)$ are in the same connected component of $G \setminus K$. Because S_Γ is a generating set, we may assume that $g_2 = g_1s$ for some $s \in S_\Gamma$. Because $g_1 \notin F_K(x)$, it follows that

$$K \cap \gamma[\alpha'_x(g_1), \alpha'_x(g_1s)] = \emptyset.$$

So $\alpha'_x(g_1), \alpha'_x(g_1s)$ are in the same connected component of $G \setminus K$ as required. □

Definition 4. Suppose H is a finitely generated group and $S_H \subset H$ is a finite symmetric generating set. Let $\text{Cay}(H, S_H)$ be the associated Cayley graph. Given a subset $F \subset H$, let ∂F be the set of all edges e of $\text{Cay}(H, S_H)$ with one endpoint in F and one endpoint in $H \setminus F$.

Lemma 3.4. *Suppose H is a finitely generated group and $S_H \subset H$ is a finite symmetric generating set. Suppose there exists a constant $C > 0$ and finite subsets $F_n \subset H$ such that*

- $|\partial F_n| \leq C$ for all $n \in \mathbb{N}$,
- $\lim_{n \rightarrow \infty} |F_n| = \infty$.

Then H has at least 2 ends.

Proof. We identify each F_n with its induced subgraph in $\text{Cay}(H, S_H)$. We may assume without loss of generality that every connected component of the complement $\text{Cay}(H, S_H) \setminus F_n$ is infinite. This is because we may add all finite components of $\text{Cay}(H, S_H) \setminus F_n$ to F_n without increasing the size of its boundary.

Choose elements $g_n \in F_n$, $s_n \in S_H$ so that

- $(g_n, g_n s_n) \in \partial F_n$
- if $F_n^\circ \subset F_n$ is the connected component of F_n containing g_n then $\lim_{n \rightarrow \infty} |F_n^\circ| = +\infty$
- there exists an infinite path $p_n \subset \text{Cay}(H, S_H) \setminus F_n$ starting from $g_n s_n$.

Let $F'_n = g_n^{-1} F_n^\circ$ and $p'_n = g_n^{-1} p_n$. After passing to a subsequence if necessary, we may assume that F'_n converges to a limit F'_∞ and p'_n converges to a limit p'_∞ (in the topology of uniform convergence on compact subsets). We observe that F'_∞ is infinite, $p'_\infty \subset \text{Cay}(H, S_H) \setminus F'_\infty$ is an infinite path and $|\partial F'_\infty| \leq C$. Thus the compact set $K := \partial F'_\infty$ is such that there are at least two infinite components of $\text{Cay}(H, S_H) \setminus K$ (namely, the component containing p'_∞ and the component containing F'_∞). This proves that H has at least two ends. □

Proposition 3.5. *Suppose Γ is an infinite finitely generated group, $G = \mathbb{F}_r$ be a nonabelian free group, $\Gamma \curvearrowright (X, \mu)$ a probability measure-preserving action and $\alpha : \Gamma \times X \rightarrow G$ an L^1 -embedding. Then Γ has more than one end.*

Proof. We fix a free generating set of G from which we obtain a word metric and a Cayley graph (which is a regular tree since G is a free group). We also fix a finite generating set S_Γ for Γ .

To obtain a contradiction, we assume that $\text{End}(\Gamma) = \{\xi\}$ is a singleton. Define $\phi : X \rightarrow \text{End}(G)$ by $\phi(x) = \alpha'(\xi, x)$ where α' is as in Theorem 3.1. By Theorem 3.1,

$$\phi(hx) = \alpha(h, x)\phi(x).$$

For $n \in \mathbb{N}$, $x \in X$, let $G(n, x)$ be the set of all $g \in G$ such that $(g|\phi(x))_e \leq n$ where $(\cdot|\cdot)_e$ is the Gromov product. To be precise $(g|\phi(x))_e = d(e, m)$ where, if $g \neq e$, $m \in G$ is the ‘midpoint’ of the geodesic triangle with vertices $\{g, \phi(x), e\}$. That is, m is the unique element contained in all three geodesic sides of the triangle with vertices $\{g, \phi(x), e\}$. If $g = e$ then by definition $m = e$. Thus $G(n, x)$ is the set of all elements $g \in G$ such that the geodesic from g to $\phi(x)$ contains a point of distance no more than n from e . Let

$F(n, x) = \{h \in \Gamma : \alpha'(h, x) \in G(n, x)\}$.

Claim 1. $\int |\partial F(n, x)| d\mu(x) \leq C \sum_{s \in S_\Gamma} \int |\alpha(s, x)|_G d\mu(x) =: M$. Note M is independent of n .

Proof of Claim 1. Let $r_n(x) \in G$ be the unique element satisfying

$$d(e, r_n(x)) = n = (r_n(x)|\phi(x))_e.$$

In other words, $n \mapsto r_n(x)$ is the geodesic from e to $\phi(x)$. Observe that $\partial G(n, x)$ is the unique edge from $r_n(x)$ to $r_{n+1}(x)$.

By definition $\partial F(n, x)$ consists of all edges of the form (g, gs) such that $g \in F(n, x)$ and $gs \notin F(n, x)$ ($s \in S$). Equivalently, $\alpha'_x(g) \in G(n, x)$ and $\alpha'_x(gs) \notin G(n, x)$. Equivalently, the s -flight from $\alpha'_x(g)$ to $\alpha'_x(gs)$ flies over $r_n(x)$. The claim now follows as in the proof of Lemma 3.3, Claim 1. \square

Claim 2. For every $n \in \mathbb{N}$ and a.e. $x \in X$, $|F(n, x)| < \infty$.

Proof. The previous claim implies $\partial F(n, x)$ is finite for a.e. x . Because Γ is 1-ended, for a.e. $x \in X$ either $F(n, x)$ or $\Gamma \setminus F(n, x)$ is finite. Because $\alpha' : \bar{\Gamma} \times X \rightarrow \bar{G}$ is continuous and $\alpha'(\xi, x) \notin \bar{G}(n, x) = \alpha'(F(n, x), x)$, it follows that $\Gamma \setminus F(n, x)$ must be infinite and therefore $F(n, x)$ is finite. \square

Observe that $G(n, x) \subset G(n+1, x)$ and $\cup_{n \geq 0} G(n, x) = G$. Therefore $F(n, x) \subset F(n+1, x)$ for all n and $\cup_{n \geq 0} F(n, x) = \Gamma$ which in particular implies that $\lim_{n \rightarrow \infty} |F(n, x)| = +\infty$.

Because

$$\lim_{n \rightarrow \infty} \int |F(n, x)| d\mu(x) = +\infty, \quad \int |\partial F(n, x)| d\mu(x) \leq M,$$

we can choose $x_n \in X$ so that $|F(n, x_n)| \rightarrow \infty$ while $|\partial F(n, x_n)|$ stays bounded. Lemma 3.4 now implies that Γ has at least 2 ends, a contradiction. \square

Proof of Theorem 1.1. By assumption there exists an L^1 -embedding $\alpha : \Gamma \times X \rightarrow G$ and G is a free group. Since Γ is accessible, we may write it as the fundamental group of a finite graph of groups in which each edge group is finite and each vertex group has ≤ 1 end. By Lemma 2.2 each vertex group H quasi-isometrically embeds into Γ . So if we restrict α to $H \times X$, it is still an L^1 -embedding. So Proposition 3.5 implies that H is not 1-ended. So every vertex group and edge group in the graph of groups decomposition of Γ is finite. This implies that Γ is virtually free by Theorem 2.1. \square

References

- [Aus13] Tim Austin. Integrable measure equivalence for groups of polynomial growth with an appendix by Lewis Bowen. *arXiv preprint arXiv:1310.3216*, 2013.

- [BFS13] Uri Bader, Alex Furman, and Roman Sauer. Integrable measure equivalence and rigidity of hyperbolic lattices. *Invent. Math.*, 194(2):313–379, 2013.
- [Can15] Michael Cantrell. Differentiability of integrable measurable cocycles between nilpotent groups. *arXiv preprint arXiv:1509.08966*, 2015.
- [Dun85] M. J. Dunwoody. The accessibility of finitely presented groups. *Invent. Math.*, 81(3):449–457, 1985.
- [Dun93] Martin J. Dunwoody. An inaccessible group. In *Geometric group theory, Vol. 1 (Sussex, 1991)*, volume 181 of *London Math. Soc. Lecture Note Ser.*, pages 75–78. Cambridge Univ. Press, Cambridge, 1993.
- [Fur99a] Alex Furman. Gromov’s measure equivalence and rigidity of higher rank lattices. *Ann. of Math. (2)*, 150(3):1059–1081, 1999.
- [Fur99b] Alex Furman. Orbit equivalence rigidity. *Ann. of Math. (2)*, 150(3):1083–1108, 1999.
- [Fur11] Alex Furman. A survey of measured group theory. In *Geometry, rigidity, and group actions*, Chicago Lectures in Math., pages 296–374. Univ. Chicago Press, Chicago, IL, 2011.
- [Gab05] D. Gaboriau. Examples of groups that are measure equivalent to the free group. *Ergodic Theory Dynam. Systems*, 25(6):1809–1827, 2005.
- [Gro93] M. Gromov. Asymptotic invariants of infinite groups. In *Geometric group theory, Vol. 2 (Sussex, 1991)*, volume 182 of *London Math. Soc. Lecture Note Ser.*, pages 1–295. Cambridge Univ. Press, Cambridge, 1993.
- [PW02] Panos Papasoglu and Kevin Whyte. Quasi-isometries between groups with infinitely many ends. *Comment. Math. Helv.*, 77(1):133–144, 2002.
- [Sta68] John R. Stallings. On torsion-free groups with infinitely many ends. *Ann. of Math. (2)*, 88:312–334, 1968.
- [TW93] Carsten Thomassen and Wolfgang Woess. Vertex-transitive graphs and accessibility. *J. Combin. Theory Ser. B*, 58(2):248–268, 1993.
- [Wal71] C. T. C. Wall. Pairs of relative cohomological dimension one. *J. Pure Appl. Algebra*, 1(2):141–154, 1971.