
A Radio Detection Survey of Narrow-Line Seyfert 1 Galaxies using Very-Long-Baseline Interferometry at 22 GHz

Akihiro DOI^{1,2}, Tomoaki OYAMA³, Yusuke KONO⁴, Aya YAMAUCHI³,
Syunsaku, SUZUKI⁴, Naoko MATSUMOTO^{5,4}, and Fumie TAZAKI⁴

¹The Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency,
3-1-1 Yoshinodai, Sagami-hara, Kanagawa 229-8510

²Department of Space and Astronautical Science, SOKENDAI (The Graduate University for
Advanced Studies), 3-1-1 Yoshinodai, Sagami-hara, Kanagawa 229-8510

³Mizusawa VERA Observatory, National Astronomical Observatory of Japan, 2-12
Hoshigaoka, Mizusawa, Oshu, Iwate 023-0861

⁴Mizusawa VLBI Observatory, NAOJ, 2-21-1 Osawa, Mitaka, Tokyo 181-8588

⁵The Research Institute for Time Studies, Yamaguchi University, 1677-1 Yoshida, Yamaguchi,
Yamaguchi 753-8511

Received 2016 February 8; Accepted 2016 March 15

Abstract

We conducted a high-sensitivity radio detection survey for forty narrow-line Seyfert 1 (NLS1) galaxies using very-long-baseline interferometry (VLBI) at 22 GHz through phase-referencing long-time integration and using a newly developing recorder with a data rate of 8 Gbps, which is a candidate of the next generation VLBI data recording systems for the Japanese VLBI Network. The baseline sensitivity was typically a few mJy. The observations resulted in a detection rate of 12/40 for our radio-selected NLS1 sample; 11 out of the detected 12 NLS1s showed inverted radio spectra between 1.4 and 22 GHz on the basis of the Very Large Array flux densities and the VLBI detections. These high fractions suggest that a compact radio core with a high brightness temperature is frequently associated with NLS1 nuclei. On the other hand, at least half of the sample indicated apparently steep spectra even with the limited VLBI sensitivity. Both the inverted and steep spectrum radio sources are included in the NLS1 population.

Key words: galaxies: active — galaxies: jets — radio continuum: galaxies — galaxies: Seyfert — techniques: interferometric

1 Introduction

The radio natures of narrow-line Seyfert 1 (NLS1s) galaxies potentially provide us key understandings of outflowing mechanisms in the growing phase of active galactic nuclei (AGNs). NLS1s as a class are thought to be fed at high mass accretion rates onto relatively small-mass black holes, potentially connecting between stellar-mass and supermassive black hole systems in the mass function. The first systematic studies in ra-

dio bands by interferometric observations were carried out at arc-second resolutions using the Very Large Array (VLA), and indicated little difference between NLS1s and Seyfert galaxies: similar radio luminosities, steep radio spectra, and scarcely resolved radio morphology suggest the presence of weak nonthermal jets as a radio emitting source (Ulvestad et al. 1995; Moran 2000), although several optical/X-ray properties are clearly different. NLS1s were thought to be radio-quiet objects as a class

with only a few exceptions of known radio-loud objects (e.g., Grupe et al. 2000; Oshlack et al. 2001; Siebert et al. 1999); subsequent systematic studies based on large database, such as the VLA Faint Images of the Radio Sky at Twenty-centimeters (FIRST; Becker et al. 1995) and the NRAO VLA Sky Survey (NVSS; Condon et al. 1998) in radio and the Sloan Digital Sky Survey (SDSS) in optical, have revealed a lower fraction of radio-loud objects ($\sim 7\%$; Komossa et al. 2006, see also Zhou et al. 2006; Whalen et al. 2006; Yuan et al. 2008) compared to that of broad-line Seyfert galaxies.

Recently, gamma-ray detections by *Fermi Gamma-Ray Space Telescope* toward six NLS1s galaxies with high significance (Abdo et al. 2009a, 2009c; D’Ammando et al. 2012, 2015) have offered a new population of gamma-ray emitting AGNs other than blazars and radio galaxies. The radio observations at high angular resolutions using very-long-baseline interferometry (VLBI) for the first discovered γ -ray emitting NLS1 SDSS J094857.31+002225.4 (PMN J0948+0022) had revealed the presences of a one-sided pc-scale jet and a rapidly variable, very high brightness radio core showing an inverted spectrum at the nucleus (Doi et al. 2006). PMN J0948+0022 also shows a core-dominant structure with two-sided kpc-scale radio emissions (Doi et al. 2012). The combination of these radio/gamma-ray properties is reminiscent of blazars, which are characterized by Doppler-beaming on relativistic jets viewed from pole-on typically at parsec (pc) scales and by decelerated components at from kpc to Mpc scales. VLBI observations of the other several gamma-ray emitting NLS1s also indicate the presence of beamed jets at pc scales (D’Ammando et al. 2012, 2013b; Doi et al. 2011, 2013; Wajima et al. 2014); some of them also show two-sided kpc-scale radio structures (Antón et al. 2008; Doi et al. 2011, 2012). However, the radio jets are mildly relativistic and their powers are comparable to the least energetic blazars, on the basis of single-dish monitoring (Angelakis et al. 2015).

On the other hand, VLBI observations of radio-loud NLS1s including gamma-ray detections/non-detections have revealed that the radio-loud aspect is possibly attributed not only to a beaming effect but also an intrinsically large radio power. Intrinsic radio powers that are corrected for the beaming effect using estimated Doppler factors suggest that radio-loud NLS1s include both intrinsically radio-loud and intrinsically radio-quiet cases (Doi et al. 2011, 2012). An extended pc-scale jet showing a steep spectrum dominates a total radio power, which suggests being not so relativistic and/or sufficiently inclined with respect to our line of sight for the jet (Gu & Chen 2010; Gu et al. 2015). Seven out of known ten NLS1s with kpc-scale radio structures (Richards & Lister 2015, see also Gliozzi et al. 2010; Doi et al. 2012, 2015) are not still gamma-ray detected. This situation suggests that jet powers may be intrinsically large sufficient to escape the core regions of host galaxies,

but not so beamed because of relatively large viewing angles. For several nearby radio-quiet NLS1s, VLBI observations have also revealed pc-scale jet-like structures showing steep spectra, which are presumably generated through the same process as that of radio-loud NLS1s, but probably with an only difference in power or viewing angle for nonthermal jets (Giroletti et al. 2005; Doi et al. 2013, 2015). Thus, at least a fraction of NLS1 nuclei has an ability to generate nonthermal jets, which have a wide range of radio powers; in some cases, the signatures of beaming effect on relativistic jets viewed from pole-on are observed.

However, a limited number of NLS1s has still been investigated at milli-arcsecond (mas) resolutions so far, because NLS1s are relatively weak radio sources in comparison with the other AGN classes. Systematic VLBI observations have been started by the other authors (Gu et al. 2015), whose targets are sixteen radio-loud NLS1s selected from several parent samples with 105 sources; fourteen sources were detected at 5 and 6.7 GHz, which were relatively low frequencies and may tend to be biased to the contribution of an extended jet component. The present paper reports the result of a new VLBI detection survey as a systematic study for NLS1s in the radio band at a higher frequency (22 GHz), where an inner jet component is potentially focused on. Our observation was planned before the publication of Gu et al. (2015). Our sample selection was relatively similar to theirs; the differences are (1) the combination of parent samples, (2) including radio-quiet NLS1s as well, and (3) at a higher observing frequency. Section 2 presents the sample selection. In Sections 3 and 4, VLBI observations and the procedures of data reductions are described. These results are presented and its implications are briefly discussed in Section 5.

2 Sample

We selected NLS1 radio sources by position-matching between the catalog of the Very Large Array (VLA) 1.4-GHz Faint Images of the Radio Sky at Twenty cm (FIRST) survey (Becker et al. 1995) and source lists in the following NLS1 studies: (1) 64 sources in Véron-Cetty et al. (2001), (2) 2011 sources in Zhou et al. (2006), (3) 23 sources in Yuan et al. (2008), and (4) 62 sources in Whalen et al. (2006). We found several sources that were counted redundantly; the number of uniquely selected sources resulted in 233 sources ($734-0.8$ mJy beam $^{-1}$). We clipped these sources into 41 sources with an intensity higher than 10 mJy beam $^{-1}$. The NLS1 radio source catalogue that is a target list in the present study is listed in Table 1.

The sample selection criteria are similar to those of the previous systematic VLBI study by Gu et al. (2015), which was also based on the radio selected samples with flux densities of > 10 mJy in the FIRST. The parent samples were restricted to only radio-loud objects from Komossa et al. (2006); Zhou

et al. (2006); Yuan et al. (2008); Foschini (2011), which were slightly different from ours including radio-quiet ones as well. Six radio-quiet objects are included in our sample ($\log RL < 1$, see Col. (7) in Table 1).

3 Observation

The observations are performed as a part of the experimental observations to test a developing OCTAVE-DAS (Data Acquisition System, Oyama et al. 2012). It is a candidate of the next generation data recording system for OCTAVE (Optically Connected Array for VLBI Exploration, Kono et al. 2012), JVN (Japanese VLBI Network, Fujisawa et al. submitted) and other VLBI arrays.

OCTAVE-DAS consists of three key components. The first is a high speed analog-to-digital (A/D) converter at a sampling rate of 8 gigabits per second (Gbps) of 3 bit quantization. It is called OCTAD (OCTave A/D converter). It is able to sample signals at a radio frequency (RF) directly. In addition, it has digital base-band converter (DBBC) functions for the VLBI Global Observing System (VGOS) observation. The second key component is a media converter between single 10 GigE (Gigabit Ethernet) port and four VSI-H (Vlbi Standard Interface-Hardware) I/O ports. It is called OCTAVIA (OCTave Vlbi Interface Adapter) or OCTAVIA2 depending on its version. The last key component is a recorder. It is able to record the data stream at the rate from 4.5 Gbps up to 32 Gbps. It is called OCTADISK (OCTave DISK recorder), OCTADISK2 or VSREC (Vlbi Software Recorder) depending on its version. OCTAVE-DAS is being developed at the Mizusawa VLBI observatory, a branch of the National Astronomical Observatory of Japan.

The test observations using OCTAVE-DAS were conducted on April 21 and 28, 2014 using four radio telescopes of the VLBI Exploration of Radio Astrometry project (VERA; Kobayashi et al. 2003); Mizusawa, Ogasawara, Iriki, and Ishigaki stations participated in the experiment. The polarization is the left-circular-polarization. VERA has a dual-beam receiving system. The dual beams are called A- and B-beam. For the dual beam observation in this paper, A-beam was used for the target sources. B-beam was used for the reference sources.

The received radio signals of dual beams are converted to IF frequency, respectively. The bandwidth is 512 MHz. The IF signals are sampled by two A/D converters (ADS-1000s). The sampling rate is 1.024 GHz. The quantization bit number is two. The data streams are recorded by OCTADISK. The aggregated bit rate is 4.096 Gbps.

In addition to this current VERA observation system, we installed OCTAVE-DAS in order to expand the bandwidth. We also installed the analog signal converters with wider bandwidth. The new system was installed on A-beam for this test

observation to increase the sensitivity of target sources. The received signal of A-beam was divided to two signals for the current and the new system. The signal for the new systems is divided to four IF channels of which bandwidth is 512 MHz. The starting frequencies of the four IFs are 21.459, 21.971, 22.483, and 22.995 GHz. The four IF signals are A/D converted by ADS-3000+ which was developed by NICT (National Institute of Information and Communications Technology) instead of OCTAD for the observation in this paper. The sampled data stream was recorded by VSREC. The aggregated recording rate is 8.192 Gbps.

The dual beam systems with the current VERA system and OCTAVE-DAS can work simultaneously. We utilized “dual-beam phase referencing” in order to increase coherent integration time by elimination of the atmospheric phase fluctuation of the targets with the phases of the calibrators. For several targets (SBS 0846+513, SDSS J103727.44+003635.5, B3 1441+476, [HB89] 1519-065, [HB89] 1546+353), proper calibrators cannot be found within a separation angle of 2.1 degrees. We used a nodding style phase-referencing. A-beam pointing was switched between a target and a reference calibrator. On-source tracking was used for PMN J0948+0022, which was expected to be detected without phase-referencing. SDSS J124634.64+023809.0 was mis-allocated in our observation. The number of observed targets resulted in 40.

The A/D converter ADS1000 at Ishigaki station was unlocked to a reference signal during both of the two observations unfortunately. The solutions of phase reference from the B-beam were unavailable.

4 Data reduction

The correlations were processed with the software correlator OCTACOR2 (OCTave CORrelator), which was developed at the Mizusawa VLBI observatory and NICT (Oyama et al. 2012). The correlated data were integrated every one second. The frequency channel number per an IF channel is 512.

Data reduction procedures were performed using the Astronomical Image Processing System (AIPS; Greisen 2003). Amplitude calibration using a priori gain values together with the system noise temperatures measured during the observations were applied. The calibration accuracy had not been evaluated in this new system, probably less than 20%, which was inferred from the comparison of results of known strong sources in the data. The delay differences among the four IF channels were calibrated using the fringe-fitting solutions of a bright calibrator (“manual-pcal”). This solution allowed us to adopt a small fringe-finding window in subsequent fringe fitting procedures.

For dual-beam phase-referencing, the fringe-fitting solution in phase for the reference calibrator was obtained with the B-

beam data at first. Next, we applied the solution to the four channels of the A-beam. Calibrators for five targets were not detected. The difference between the center frequencies of two beams is 256 MHz ($\sim 1\%$ with respect to the radio frequency). In the case of residual phase variation of ~ 100 deg (typically observed during VERA dual-beam observations) after a fringe-rate removal, the coherence loss is expected to be only 0.02% when we apply fringe-phase solutions to data at a $\sim 1\%$ -different frequency. With the dual beam phase referencing, we were able to extend the fringe-fitting solution interval to 720 seconds. The interval is equivalent to a scan duration time of the antenna schedule. As a result, six out of 29 targets were detected at signal-to-noise ratios (SNR s) higher than 3 in this observing mode.

We also performed fringe-fitting on the five target that observed in the manner of nodding style phase-referencing with a cycle of 40 seconds. The solution interval was 720 seconds with a net accumulation of 240 seconds on a target. As a result, two out of the five targets were detected at $SNR > 3$ in this observing mode.

PMN J0948+0022, which was observed without phase-referencing, was detected with a solution interval of 120 sec. Fringe findings for five targets whose calibrators were not detected in B-beam were also attempted with a solution interval of 120 sec. As a result, three out of the five were detected at $SNR > 3$.

5 Result and Discussion

Correlated flux densities averaged through baselines with detection are listed in the last column of Table 1; the table includes the references for sources for which previous VLBI studies exist. Upper limits of correlated flux density, ranging from 7 mJy to 23 mJy depending on the observing mode (Sect. 4), were determined by $SNR = 3$ on the most sensitive baseline during observations. Twelve out of 40 observed targets were detected with VERAs' baselines ranging from ~ 1000 km to ~ 2300 km, which imply a brightness temperature of the order of 10^7 Kelvin or more for high-frequency radio emissions associated with these NLS1s. Even if we adopted a detection limit of 5σ instead of 3σ , only [HB89] 1546+353, which is the weakest detection in our sample, would get into non-detection.

The two sources, FBQS J114654.2+323652 and FBQS J1713+3523, are for the first detected with very long baselines. FBQS J114654.2+323652 was presumably detected with the MIZUSAWA–IRIKI baseline in the dual-beam phase reference mode; we also confirmed the fringe detection in a normal fringe fitting, without a calibrator, with a shorter integration (2 min). Around this source, there is no strong source potentially confusing to our observation. FBQS J1713+352 was positively detected with baselines of all the four antennas

in fringe fitting without a calibrator (because its calibrator was not detected in B-beam). Around this source, there is no strong source potentially confusing to our observation. Both the two sources are weak radio sources at 1.4 GHz in the FIRST (15.42 mJy and 11.24 mJy for FBQS J114654.2+323652 (1994.4) and FBQS J1713+352 (1994.5), respectively); on the other hand, they are relatively strong at 22 GHz in our observation (104 mJy and 138 mJy). These sources should be inverted spectrum sources with $\alpha = +0.7$ and $\alpha = +0.9$, where α is the spectral index ($F_\nu \propto \nu^\alpha$), assuming that the 1.4 GHz FIRST (and the 22 GHz) fluxes do not vary significantly in ~ 20 years. Such a weak source showing an inverted spectrum has also been previously reported by Doi et al. (2007) for FBQS J1629+4007, which is also in the list of the present study and has been detected (11.95 mJy at 1.4 GHz (1994.6) and 145 mJy at 22 GHz). In the first place, relatively weak radio emissions at 1.4 GHz and a limited baseline sensitivity at 22 GHz were supposed to strongly bias VLBI detections toward inverted spectrum sources. In fact, all the detected sources except for one (11/12) show flat or inverted spectra ($\alpha = -0.1$ – $+0.9$) between 1.4–22 GHz, if variability in the time interval between measurements at 1.4 and 22 GHz is assumed insignificant. Although this is a sort of artificial effect, it is noteworthy that not so small fraction of NLS1 radio sources (12/40) have been detected at such a high frequency even with very long baselines.

On the other hand, at least half of the sources indicate apparently steep spectra ($\alpha < -0.2$) between 1.4 GHz and 22 GHz on the basis of the FIRST flux densities and the VLBI upper limits. Both the inverted and steep spectrum radio sources are included in the NLS1 population, as previously pointed out by several authors (Doi et al. 2007; Gu & Chen 2010; Doi et al. 2011). Particularly, [HB89] 1044+476 is the strongest radio source (767.44 mJy at 1.4 GHz in the FIRST (1997.2)) in our sample but not detected with the VLBI at 22 GHz (< 7 mJy). Only a weak radio emission (19.4 mJy) with a core-jet structure in the east-west direction has been previously found in VLBI images at 5 GHz while VLA images at 8.4 and 22 GHz show significantly large radio flux densities; a bulk of total flux must be resolved out with very long baselines and originate in extended components of a compact steep spectrum source (Gu et al. 2015). B3 1702+457 is in the similar situation (118.64 mJy at FIRST 1.4 GHz (1997.2) and < 9 mJy at VLBI 22 GHz); the previous VLBI studies revealed an extended radio structure with a steep spectrum for B3 1702+457 (Doi et al. 2007, 2011). Hence, our negative detections with a limited sensitivity at 22 GHz are genuine even for these sources relatively strong at 1.4 GHz.

Consequently, the systematic study by our VLBI detection survey has revealed that NLS1s are one of AGN subclasses that can possess compact radio components with high bright-

Table 1. Samples and Results of Observations

NED name (1)	z (2)	RA. (J2000) (3)	Dec. (J2000) (4)	$I_{1.4\text{GHz}}^{\text{FIRST}}$ (5)	Sample (6)	$\log RL$ (7)	Ref. (8)	Mode (9)	Calibrator (10)	$F_{\text{VLBI}}^{\dagger}$ (11)
2MASX J03474022+0105143	0.031	03 47 40.195	+01 05 14.25	38.85	a	0.99		2B	J0352+0238	< 7
FBQS J0713+3820	0.123	07 13 40.291	+38 20 40.08	10.43	b	0.33		2B	J0709+3737	< 7
2MASS J07440228+5149175	0.46	07 44 02.242	+51 49 17.48	11.89	b	1.62		2B	J0733+5022	< 7
FBQS J075800.0+392029	0.096	07 58 00.047	+39 20 29.09	10.8	b	-0.19		2B	J0752+3730	< 7
SDSS J081432.11+560956.6	0.510361	08 14 32.135	+56 09 56.55	69.18	c	2.53	e	2B	J0824+5552	117
SBS 0846+513	0.584701	08 49 57.990	+51 08 28.83	344.09	c	3.16	f	ND	J0905+4850	454
SDSS J085001.16+462600.5	0.524316	08 50 01.171	+46 26 00.41	20.9	c	2.23	e	2B	J0847+4609	< 7
SDSS J090227.16+044309.5	0.533025	09 02 27.152	+04 43 09.40	152.59	c	3.02	e	2B	J0901+0448	244
PMN J0948+0022	0.585102	09 48 57.295	+00 22 25.60	107.53	c	2.55	ghi	1B	...	501
MRK 1239	0.019927	09 52 19.099	-01 36 43.63	58.64	a	1.33	jk	2B	J0945-0153	< 7
SDSS J095317.09+283601.4	0.65891	09 53 17.106	+28 36 01.63	44.58	c	2.71	e	2B	J1001+2911	< 7
SDSS J103123.73+423439.3	0.377167	10 31 23.728	+42 34 39.40	16.57	c	2.34		2B	J1038+4244	< 7
KUG 1031+398	0.042443	10 34 38.599	+39 38 28.17	23.98	a	1.67		2B	J1033+4116	< 7
SDSS J103727.44+003635.5	0.595596	10 37 27.454	+00 36 35.76	27.23	c	2.66	e	ND	J1048+0055	< 12
[HB89] 1044+476	0.79902	10 47 32.654	+47 25 32.24	734.02	c	3.87	e	2B	J1051+4644	< 7
SDSS J111005.03+365336.3	0.62995	11 10 05.034	+36 53 36.12	18.62	c	2.97	e	2B	J1104+3812	< 7
2MASX J11193404+5335181	0.105975	11 19 34.026	+53 35 18.45	15.49	d	1.96		2B	J1120+5404	< 7
SDSS J113824.54+365327.1	0.356743	11 38 24.545	+36 53 26.99	12.54	c	2.34	e	2B	J1130+3815	< 7
2MASX J11404788+4622046	0.11439	11 40 47.897	+46 22 04.82	78.85	b	1.36		2B	J1138+4745	< 7
FBQS J114654.2+323652	0.4658	11 46 54.298	+32 36 52.24	14.67	c	2.11		2B	J1152+3307	104
FBQS J1151+3822	0.334575	11 51 17.757	+38 22 21.75	10.93	b	0.54		2B	J1146+3958	< 7
2MASX J12022678-0129155	0.150694	12 02 26.806	-01 29 15.54	11.37	d	1.49		2B	J1207-0106	< 7
NGC 4051	0.002336	12 03 09.594	+44 31 52.52	12.3	a	0.47	lj	2B	J1203+4510†	< 17
NGC 4253	0.012929	12 18 26.516	+29 48 46.52	38.16	a	1.19		2B	J1217+3007	< 7
SDSS J123852.12+394227.8	0.622668	12 38 52.147	+39 42 27.59	10.36	c	2.23		2B	J1242+3751	< 7
SDSS J124634.64+023809.0	0.362629	12 46 34.683	+02 38 09.02	37.05	c	2.38	e	...	J1250+0216	...
MRK 0783	0.0672	13 02 58.925	+16 24 27.49	18.53	a	1.36	j	2B	J1300+141B	< 9
SDSS J130522.74+511640.2	0.787552	13 05 22.746	+51 16 39.55	83.87	c	2.34	e	2B	J1259+5140	< 9
FBQS J1421+2824	0.539978	14 21 14.075	+28 24 52.23	46.79	b	2.14	e	2B	J1419+2706	117
SDSS J143509.49+313147.8	0.502218	14 35 09.523	+31 31 48.30	39.26	c	2.87		2B	J1435+3012†	< 23
B3 1441+476	0.705472	14 43 18.578	+47 25 56.53	164.79	c	3.07	e	ND	J1452+4522	< 16
[HB89] 1502+036	0.407882	15 05 06.467	+03 26 30.83	365.39	c	3.19	mn	2B	J1458+0416†	697
[HB89] 1519-065	0.083121	15 22 28.758	-06 44 41.83	11.8	a	0.56††		ND	J1510-0543	< 16
[HB89] 1546+353	0.479014	15 48 17.924	+35 11 28.37	140.94	c	2.84	eo	ND	J1602+3326	22
IRAS 15462-0450	0.099792	15 48 56.806	-04 59 34.26	10.68	a	1.05††		2B	J1550-0538	< 9
FBQS J1629+4007	0.272486	16 29 01.315	+40 07 59.62	11.97	b	1.46	pqr	2B	J1623+3909†	145
2MASX J16332357+4718588	0.116054	16 33 23.585	+47 18 58.96	62.63	c	2.22	pqr	2B	J1637+4717	163
FBQS J1644+2619	0.145	16 44 42.536	+26 19 13.19	87.53	c	2.65	prs	2B	J1642+2523	62
B3 1702+457	0.0604	17 03 30.379	+45 40 47.09	115.44	a	2.17	pqr	2B	J1658+4737	< 9
FBQS J1713+3523	0.083	17 13 04.476	+35 23 33.43	11.13	b	1.05		2B	J1708+3346†	138
SDSS J172206.02+565451.6	0.425967	17 22 06.081	+56 54 52.00	36.91	c	2.37		2B	J1722+5856	< 9

Col. (1) Source Name in NED; Col. (2) redshift; Col. (3) right ascension; Col. (4) declination; Col. (5) peak intensity in mJy beam^{-1} in VLA FIRST; Col. (6) parent sample; Col. (7) radio loudness from reference listed in Col. (6), except for sources from Véron-Cetty et al. (2001) that includes no value for radio loudness. Note that the definition of radio loudness differs from reference to reference. We calculated radio loudness for sources in Véron-Cetty et al. (2001) by the ratio of FIRST 1.4 GHz radio to SDSS g -band PSF optical flux densities, according to Véron-Cetty et al. (2001). Sources denoted by †† represent the use of B -band magnitudes listed in Véron-Cetty et al. (2001) because their SDSS results are not available; Col. (8) reference of previous VLBI observation; Col. (9) observation mode in the present study. "ND" and "2B" represent nodding-style and VERA's dual-beam phase referencing, respectively. "1B" represents an on-source observation with a single beam. SDSS J124634.64+023809.0 was mis-allocated in our observation; Col. (10) phase-referencing calibrator used in the present study. Calibrators denoted by † represent non-detection in B-beam; Col. (11) correlated flux density in mJy measured in the present study.

References — a: Véron-Cetty et al. (2001), b: Whalen et al. (2006), c: Yuan et al. (2008), d: Zhou et al. (2006), e: Gu et al. (2015), f: D'Ammando et al. (2013a), g: Doi et al. (2006), h: Abdo et al. (2009b), i: Giroletti et al. (2011), j: Doi et al. (2013), k: Doi et al. (2015), l: Giroletti & Panessa (2009), m: Fey & Charlot (2000), n: D'Ammando et al. (2013b), o: Orienti et al. (2015), p: Doi et al. (2007), q: Gu & Chen (2010), r: Doi et al. (2011), s: Doi et al. (2012)

ness temperatures ($\gtrsim 10^7$ Kelvin) even at the high frequency (22 GHz). These compact components show inverted spectra in almost all the VLBI-detected cases, which account for a significant fraction in our radio-selected NLS1 sample. These properties may be related to NLS1s' blazer-like aspects such as gamma-ray detections in several NLS1s in contrast to normal broad-line Seyfert galaxies.

Acknowledgments

This study was partially supported by Grants-in-Aid for Scientific Research (B) (24340042, AD) and Grant-in-Aid for Scientific Research on Innovative Areas (26120537, AD) from the Japan Society for the Promotion of Science (JSPS). We are grateful to all the staffs and students involved in the development and operation of the Japanese VLBI network (JVN) and the VLBI Exploration of Radio Astrometry project (VERA). The JVN project is led by the National Astronomical Observatory of Japan (NAOJ), which is a branch of the National Institutes of Natural Sciences (NINS), Yamaguchi University, Hokkaido University, Gifu University, Kagoshima University, Tsukuba University, Osaka Prefecture University, and Ibaraki University, in collaboration with the Geographical Survey Institute (GSI), the Japan Aerospace Exploration Agency (JAXA), and the National Institute of Information and Communications Technology (NICT). The VERA is operated by the Mizusawa VLBI observatory, a branch of the National Astronomical Observatory of Japan. We used the US National Aeronautics and Space Administration's (NASA) Astrophysics Data System (ADS) abstract service and NASA/IPAC Extragalactic Database (NED), which is operated by the Jet Propulsion Laboratory (JPL). In addition, we used the Astronomical Image Processing System (AIPS) software developed at the National Radio Astronomy Observatory (NRAO), a facility of the US National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

References

- Abdo, A. A., et al. 2009a, *ApJ*, 699, 976
 —. 2009b, *ApJ*, 707, 727
 —. 2009c, *ApJL*, 707, L142
 Angelakis, E., et al. 2015, *A&A*, 575, A55
 Antón, S., Browne, I. W. A., & Marchã, M. J. 2008, *A&A*, 490, 583
 Becker, R. H., White, R. L., & Helfand, D. J. 1995, *ApJ*, 450, 559
 Condon, J. J., Cotton, W. D., Greisen, E. W., Yin, Q. F., Perley, R. A., Taylor, G. B., & Broderick, J. J. 1998, *AJ*, 115, 1693
 D'Ammando, F., Orienti, M., Larsson, J., & Giroletti, M. 2015, *MNRAS*, 452, 520
 D'Ammando, F., et al. 2012, *MNRAS*, 426, 317
 —. 2013a, *MNRAS*, 436, 191
 —. 2013b, *MNRAS*, 433, 952
 Doi, A., Asada, K., Fujisawa, K., Nagai, H., Hagiwara, Y., Wajima, K., & Inoue, M. 2013, *ApJ*, 765, 69
 Doi, A., Asada, K., & Nagai, H. 2011, *ApJ*, 738, 126
 Doi, A., Nagai, H., Asada, K., Kameno, S., Wajima, K., & Inoue, M. 2006, *PASJ*, 58, 829
 Doi, A., Nagira, H., Kawakatu, N., Kino, M., Nagai, H., & Asada, K. 2012, *ApJ*, 760, 41
 Doi, A., Wajima, K., Hagiwara, Y., & Inoue, M. 2015, *ApJL*, 798, L30
 Doi, A., et al. 2007, *PASJ*, 59, 703
 Fey, A. L., & Charlot, P. 2000, *ApJS*, 128, 17
 Foschini, L. 2011, in *Narrow-Line Seyfert 1 Galaxies and their Place in the Universe*
 Fujisawa et al., K. submitted, *PASJ*
 Giroletti, M., & Panessa, F. 2009, *ApJL*, 706, L260
 Giroletti, M., Taylor, G. B., & Giovannini, G. 2005, *ApJ*, 622, 178
 Giroletti, M., et al. 2011, *A&A*, 528, L11
 Gliozzi, M., Papadakis, I. E., Grupe, D., Brinkmann, W. P., Raeth, C., & Kedziora-Chudczer, L. 2010, *ApJ*, 717, 1243
 Greisen, E. W. 2003, *Information Handling in Astronomy - Historical Vistas*, 285, 109
 Grupe, D., Leighly, K. M., Thomas, H.-C., & Laurent-Muehleisen, S. A. 2000, *A&A*, 356, 11
 Gu, M., & Chen, Y. 2010, *AJ*, 139, 2612
 Gu, M., Chen, Y., Komossa, S., Yuan, W., Shen, Z., Wajima, K., Zhou, H., & Zensus, J. A. 2015, *ApJS*, 221, 3
 Kobayashi, H., et al. 2003, in *Astronomical Society of the Pacific Conference Series*, Vol. 306, *New technologies in VLBI*, ed. Y. C. Minh, 367
 Komossa, S., Voges, W., Xu, D., Mathur, S., Adorf, H.-M., Lemson, G., Duschl, W. J., & Grupe, D. 2006, *AJ*, 132, 531
 Kono, Y., et al. 2012, in *Seventh General Meeting (GM2012) of the international VLBI Service for Geodesy and Astrometry (IVS)*, held in Madrid, Spain, March 4-9, 2012, Eds: D. Behrend and K.D. Bayer, *National Aeronautics and Space Administration*, ed. D. Behrend & K. D. Bayer
 Moran, E. C. 2000, *New Astronomy Reviews*, 44, 527
 Orienti, M., D'Ammando, F., Larsson, J., Finke, J., Giroletti, M., Dallacasa, D., Isacsson, T., & Stoby Hoglund, J. 2015, *MNRAS*, 453, 4037
 Oshlack, A. Y. K. N., Webster, R. L., & Whiting, M. T. 2001, *ApJ*, 558, 578
 Oyama, T., et al. 2012, in *Seventh General Meeting (GM2012) of the international VLBI Service for Geodesy and Astrometry (IVS)*, held in Madrid, Spain, March 4-9, 2012, Eds: D. Behrend and K.D. Bayer, *National Aeronautics and Space Administration*, p. 91-95, ed. D. Behrend & K. D. Bayer, 91-95
 Richards, J. L., & Lister, M. L. 2015, *ApJL*, 800, L8
 Siebert, J., Leighly, K. M., Laurent-Muehleisen, S. A., Brinkmann, W., Boller, T., & Matsuoka, M. 1999, *A&A*, 348, 678
 Ulvestad, J. S., Antonucci, R. R. J., & Goodrich, R. W. 1995, *AJ*, 109, 81
 Véron-Cetty, M.-P., Véron, P., & Gonçalves, A. C. 2001, *A&A*, 372, 730
 Wajima, K., Fujisawa, K., Hayashida, M., Isobe, N., Ishida, T., & Yonekura, Y. 2014, *ApJ*, 781, 75
 Whalen, D. J., Laurent-Muehleisen, S. A., Moran, E. C., & Becker, R. H. 2006, *AJ*, 131, 1948
 Yuan, W., Zhou, H. Y., Komossa, S., Dong, X. B., Wang, T. G., Lu, H. L., & Bai, J. M. 2008, *ApJ*, 685, 801
 Zhou, H., Wang, T., Yuan, W., Lu, H., Dong, X., Wang, J., & Lu, Y. 2006, *ApJS*, 166, 128