

ALMA OBSERVATIONS OF THE ARCHETYPAL “HOT CORE” THAT ISN’T: ORION KL

M. T. OROZCO-AGUILERA

Instituto Nacional de Astrofísica, Óptica y Electrónica, Luis Enrique Erro 1, Tonantzintla, Puebla, México

LUIS A. ZAPATA

Instituto de Radioastronomía y Astrofísica, UNAM, Apdo. Postal 3-72 (Xangari), 58089
Morelia, Michoacán, México

TOMOYA HIROTA

Mizusawa VLBI Observatory, National Astronomical Observatory of Japan, Osawa 2-21-1, Mitaka-shi, Tokyo 181-8588, Japan

SHENG-LI, QIN

Department of Astronomy, Yunnan University, and Key Laboratory of Astroparticle Physics of Yunnan Province, Kunming 650091, China
0000-0003-2302-0613

MASQUÉ, JOSEP M

Departamento de Astronomía, Universidad de Guanajuato, Apdo. Postal 144, 36000 Guanajuato, México

(Received; Revised; Accepted)

Draft version November 10, 2018

ABSTRACT

We present sensitive high angular resolution ($\sim 0.1'' - 0.3''$) continuum ALMA (The Atacama Large Millimeter/Submillimeter Array) observations of the archetypal hot core located in Orion-KL. The observations were made in five different spectral bands (bands 3, 6, 7, 8, and 9) covering a very broad range of frequencies (149 – 658 GHz). Apart of the well-know millimeter emitting objects located in this region (Orion Source I and BN), we report the first submillimeter detection of three compact continuum sources (ALMA 1-3) in the vicinities of the Orion-KL hot molecular core. These three continuum objects have spectral indices between 1.47 to 1.56, and brightness temperatures between 100 to 200 K at 658 GHz suggesting that we are seeing moderate optically thick dust emission with possible grain growth. However, as these objects are not associated with warm molecular gas, and some of them are farther out from the molecular core, we thus conclude that they cannot heat the molecular core. This result favours the hypothesis that the hot molecular core in Orion-KL core is heated externally.

Subject headings: ISM: general; ISM: kinematics and dynamics; stars: formation

1. INTRODUCTION

Hot Molecular Cores (HMC) are dense ($\geq 10^6 \text{ cm}^{-3}$), warm ($\geq 100 \text{ K}$) and compact ($\leq 10^5 \text{ AU}$) dusty regions within molecular clouds that are thought to harbour massive young stars (Kurtz et al. 2000). These regions are characterized by a strong line emission from a large amount of molecules, but in particular, from complex organic molecules (COMs), as CH_3CN , CH_3OH , HCOOH , HCOOCH_3 , CH_3OCH_3 , $\text{CH}_3\text{CH}_2\text{CN}$, and CH_3COCH_3 . COMs are defined by molecules composed with more than six atoms containing C and H elements (Herbst & van Dishoeck 2009). One of the first identified HMC was the one located in the Orion Kleinmann-Low (KL) region (Ho et al. 1979). Ho et al. (1979) identified it as a compact source of hot ammonia emission embedded in a more extended ridge of dense material. Later interferometric observations revealed the peculiar and cumply “heart” or “U” morphology of the Orion-KL HMC (Genzel et al. 1982; Wilner et al. 1994; Wright et al. 1996; Favre et al. 2011a). Such morphology is not seen in the southern compact HMCs located in Orion South even observing at the same spatial scales (Zapata et al. 2007, 2010).

The nature of the Orion-KL HMC was first questioned by Blake et al. (1996) using continuum/line Owens Valley Radio Observatory (OVRO) observations, that found no evidence of a luminous internal heating source within the Orion-KL HMC. This result was more aggravated with Berkeley-Illinois-Maryland Association (BIMA) line observations of the formic acid (that also trace the HMC) and the position in the sky of water masers, which suggested that these molecules trace the interaction region between the outflow and the molecular gas at nearly systemic velocities (Liu et al. 2002). Moreover, using CO (carbon monoxide) BIMA observations, Chernin & Wright (1996) proposed that the biconical outflow in Orion-KL is partly truncated by the hot molecular core.

Recent studies have also confirmed that the HMC in Orion-KL is indeed externally heated possibly by an explosive outflow occurred some 500 yrs ago (Zapata et al. 2009, 2011a; Bally et al. 2017) or maybe by the Orion Source I (Goddi et al. 2011; Wright & Plambeck 2017) or the expanding bubble-like outflow (Zapata et al. 2011b). Some other works also confirming this externally heating hypothesis include: Favre et al. (2011b); Hirota et al. (2011); Peng et al. (2012, 2013); Bell et al. (2014); Gong et al.

TABLE 1
SUMMARY OF THE OBSERVATIONS

ALMA Band	Date	Center frequency ^a [GHz]	Total bandwidth [MHz]	uv range [k λ]	Number of Antennas	On-source Time [min.]	Synthesized Beam [arcsec]	PA [degree]	rms [mJy beam ⁻¹]
4	2015/09/04	149.5	4000	7 – 815	33	~ 8	0.36×0.34	-46	2.5
4	150 – 815	0.29×0.28	-67	1.5
6	2015/08/28	232.6	2342	200 – 1480	40	~ 46	0.21×0.17	+77	4.5
7	2014/07/26	348.4	7500	39 – 870	31	~ 25	0.29×0.26	-89	6.1
7	200 – 870	0.26×0.20	+66	4.6
8	2015/09/22	433.7	5624	200 – 3100	35	~ 25	0.10×0.08	+70	5.0
9	2014/08/05	658.5	7500	300 – 1400	35	~ 5	0.17×0.14	-42	34.0

^a Center frequency after averaging the four spectral windows.

(2015); Peng et al. (2017); Wright & Plambeck (2017). For example, Peng et al. (2017) found that the Orion-KL HMC emission peaks of vibrationally excited HC₃N lines move from south to northeast with increasing E_u , and that the HC₃N higher-energy lines have higher rotational temperatures and low column densities which appear to support that the hot core is externally heated.

The strong bursting observed in the water masers located in the direction HMC in Orion-KL could be caused by the interaction between the explosive outflow and the ambient quiescent gas (Hirota et al. 2011, 2014).

In this study, using the tremendous sensitivities, a better uv -plane coverage, and a very broad covering of frequency range at (sub)millimeter wavelengths offered by the recent operational ALMA (The Atacama Large Millimeter/Submillimeter Array) observatory, we carried out a search for compact and faint continuum sources (in five different continuum bands of ALMA) that are probably associated with (proto)stellar objects and with hot molecular gas within the HMC in Orion-KL. We report, in addition to Orion Source I and Orion BN, the detection at submillimeter wavelengths of three new compact continuum sources (ALMA1-3) that are located in the vicinities of the HMC, but with not hot molecular gas associated. Additionally, one of these compact sources seem to be associated with high-mass stars, but it does not account by the internal heating of the HMC. We thus conclude that the hot molecular core in Orion-KL is indeed heated externally as suggested by many observational works.

2. OBSERVATIONS

The observations were carried out with ALMA between 2014 and 2015 as part of the ALMA programs: 2013.1.01034.S (Band 4), 2012.1.00146.S (Band 6), 2013.1.00048.S (Band 8), and 2012.1.00123.S (Bands 7 and 9). The total bandwidth used to estimate the continuum emission in the ALMA observations were more than 2 GHz, see Table 1. However, as there are many lines detected in the spectral windows, we have probably some contamination from very faint lines. In order to search for compact millimeter sources within the hot molecular core in Orion-KL, we constrain the uv -range of the observations, see Table 1 and Figures 1, 2, and 3. We chose the uv -range based on a trade off between removing as much extended emission as possible and keeping enough visibilities to obtain a good map. The resulting synthesized beams are presented in Table 1. The number of antennas used during the observations varied between 31 to 40, see Table 1. Weather conditions were very good and stable, with an average precipitable water vapour of 6 mm (Band 3), 1 mm (Band 6), 0.3–0.7 mm (Band 7), 0.3

mm (Band 8), and 0.15 mm (Band 9).

The ALMA calibration includes simultaneous observations of the 183 GHz water line with water vapour radiometers that measure the water column in the antenna beam, that is used to reduce the atmospheric phase noise. Quasars J0529–0519, J0423–013, J0607–0834, and J0532–0307 were used to calibrate the bandpass, the amplitude and the gain fluctuations.

The data were calibrated, imaged, and analyzed using the Common Astronomy Software Applications (CASA: McMullin et al. 2007). The data presented in this paper were also analyzed using the karma software (Gooch 1996). We used ROBUST parameter of CLEAN equal to zero in the continuum maps presented in this study. This was made in order to obtain an optimal compromise between sensitivity and angular resolution. The resulting rms-noises for the continuum images and their respective angular resolutions are presented in Table 1. All the resulting ALMA images are corrected by the primary beam attenuation. We self-calibrate all images in phase. In average, we give about 1 to 2 rounds in phase for every observation. This helped to decrease the rms-noises (on a few mJy) and thus increase the S/N ration on the final images. We also tried to do self-calibration in amplitude and phase, but we did not obtained substantial improvements on the images.

3. RESULTS

The main results of this study are presented in Table 2, and Figures 1, 2, and 3.

In Figure 1, we present the results of the ALMA observations in band 4 (149.5 GHz). In this Figure, we include the observations of the HMC in Orion-KL using the full uv -range (see Table 1) and one map only using baselines that are sensitive to compact emission (> 150 k λ) or structures with angular sizes of less than 1.3". In the image using the full uv -range (upper panel), the HMC in Orion-KL is well detected together with the two strong compact objects (Orion Source I and BN) at these millimeter wavelengths. The ALMA band 4 observations revealed the “heart” or “U” shape of the HMC as already traced by many authors, see for example Goddi et al. (2011), where it is mapped by the NH₃ line emission. This “heart” structure is very well resolved, with its peculiar clumpy structure. On the other hand, the map with the restricted uv -range, revealed 6 compact sources above the 5- σ (7.5 mJy Beam⁻¹) noise level, see the lower panel of Figure 1. However, as some of these compact sources are not detected in other ALMA bands, we only count five real detections (Orion Source I, BN, ALMA1, ALMA2, and ALMA 3). We give the physical parameters of these new sources in Table 2. We note that the submillimeter source SMA1 found using Submillimeter Ar-

TABLE 2
PHYSICAL PARAMETERS OF THE COMPACT SOURCES^a

Name	Position ^b		Flux Density					Spectral Index
	α_{2000}	δ_{2000}	Band 4	Band 6	Band 7	Band 8	Band 9	
ALMA 1	05 35 14.225	-05 22 28.02	18±2	80±5	90±5	170±15	–	1.47±0.15
ALMA 2	05 35 14.428	-05 22 33.56	41±5	138±10	210±20	305±25	820±120	1.48±0.18
ALMA 3	05 35 14.802	-05 22 30.69	31±3	117±15	180±15	260±22	720±60	1.56±0.13
Source I	05 35 14.515	-05 22 30.60	150±14	590±80	905±90	1518±150	3950±320	1.74±0.15

^a These parameters were obtained from a Gaussian fit using the viewer in CASA.

^b Positional errors are $\sim 0.05''$ for R.A. and decl.




FIG. 1.— ALMA continuum band 4 (3.0 mm) contour and colour-scale images of the hot molecular core located in Orion-KL. **Upper:** Continuum image using the full uv -range from the ALMA observations. The contours are starting from 15% to 90% in steps of 5% of the intensity peak. The intensity peak is $0.150 \text{ Jy Beam}^{-1}$. The half-power contour of the synthesized beam of the image is shown in the bottom left corner. **Lower:** Continuum image using only part of the uv -range ($> 150 \text{ kl}$) from the continuum observations. The contours are starting from 15% to 90% in steps of 5% of the intensity peak. The intensity peak is $0.130 \text{ Jy Beam}^{-1}$. The synthesized beam size is shown in the bottom left corner. The position of the source SMA1 is indicated with a cross, see [Beuther et al. \(2005\)](#). Both images are corrected by the primary beam attenuation. The spatial scale bar is presented in bottom right corner.

ray (SMA) observations and reported by [Beuther et al. \(2005\)](#) does not have any counterpart, perhaps it is resolved out by




FIG. 2.— ALMA continuum band 7 (0.8 mm) contour and colour-scale images from the hot molecular core located in Orion-KL. **Upper:** Continuum image using the full uv -range from the ALMA observations. The contours are starting from 30% to 90% in steps of 5% of the intensity peak. The intensity peak is $0.561 \text{ Jy Beam}^{-1}$. The half-power contour of the synthesized beam of the image is shown in the bottom left corner. **Lower:** Continuum image using only part of the uv -range ($> 150 \text{ kl}$) from the continuum observations. The contours are starting from 10% to 90% in steps of 5% of the intensity peak. The intensity peak is $0.453 \text{ Jy Beam}^{-1}$. The synthesized beam size is shown in the bottom left corner. The position of the source SMA1 is indicated with a cross, see [Beuther et al. \(2005\)](#). Both images are corrected by the primary beam attenuation. The spatial scale bar is presented in bottom right corner.

the present ALMA observations.

In Figure 2, we present the results of the ALMA observa-




FIG. 3.— ALMA continuum bands 6 and 9 contour and colour-scale images of the hot molecular core located in Orion-KL. **Upper:** Band 6 continuum image using part uv -range from the ALMA observations (> 200 k λ). The contours are starting from 30% to 90% in steps of 5% of the intensity peak. The intensity peak is 0.461 Jy Beam $^{-1}$. The half-power contour of the synthesized beam of the image is shown in the bottom left corner. **Lower:** Band 6 continuum image using part uv -range from the ALMA observations (> 200 k λ). The contours are starting from 55% to 90% in steps of 5% of the intensity peak. The intensity peak is 1.533 Jy Beam $^{-1}$. The first contour level, in the lower panel is about $10\text{-}\sigma$ (45 mJy Beam $^{-1}$). The half-power contour of the synthesized beam of the image is shown in the bottom left corner. The position of the source SMA1 is indicated with a cross, see [Beuther et al. \(2005\)](#). Both images are corrected by the primary beam attenuation. The blue asterisks mark the position of the peaks of the hot molecular gas as traced by the $\text{NH}_3(12,12)$ with an upper level energy over the ground of 1456 K, see [Goddi et al. \(2011\)](#). The red asterisk marks the position of the peak of the hot molecular gas as traced by the $\text{CH}_3\text{CN}(12_9,11_9)$ with an upper level energy over the ground of 646 K, see [Zapata et al. \(2011b\)](#). The spatial scale bar is presented in bottom right corner of each panel.

tions in band 7 (348.4 GHz). In this Figure, we include the observations of the HMC using the full uv -range (see Table 1) and one map only using baselines that are sensitive to compact emission (> 150 k λ) or structures with angular sizes of less than $1.3''$. In the image using the full uv -range (upper panel), only the most compact structures of the HMC are detected. These compact and clumpy structures are located close to Orion Source I and form an arc-like structure. On the other hand, the map with the restricted uv -range, revealed 4 compact sources above the $5\text{-}\sigma$ (23 mJy Beam $^{-1}$) noise level (see the lower panel of figure 2). ALMA 1 is present in the map, even though its emission is very faint. Despite starting the contours at $10\text{-}\sigma$, we consider all sources below this level are spurious because they do not show well-defined counterparts in all other ALMA bands, contrary to the objects ALMA1-3.




FIG. 4.— Spectral energy distributions (SEDs) of the detected compact sources from the millimeter to submillimeter wavelengths. The dashed lines are a least-squares power-law fit (of the form $S_\nu \propto \nu^\alpha$ to the spectrum). The α -values of the fitting are shown in the panels.

We give the flux densities at this ALMA band and from these new sources in Table 2.

In Figure 3, we present the results of the ALMA observations in band 6 and 9 (232.6 and 658.5 GHz). The data from ALMA band 8 is very similar to that obtained in band 7 and it is not shown in this study. In Figure 3, we only include the observations of the HMC using baselines that are sensitive to compact emission (> 200 k λ for band 6, and > 300 k λ for band 9) or structures with angular sizes of less than $1''$ and $0.65''$, respectively. In these maps, the Orion BN object is far from the phase center and it is very attenuated, so that it is not shown in the images of Figure 3. However, we see clearly the rest of the objects (Orion Source I, ALMA1, ALMA2, and ALMA3). In this image, we have additionally included the positions of the peaks of the hot molecular gas as traced by the $\text{NH}_3(12,12)$ with a temperature in the upper level of 1456 K, see [Goddi et al. \(2011\)](#). The positions of the peak of the hot molecular gas as traced by the $\text{CH}_3\text{CN}(12_9,11_9)$ with a temperature in the upper level of 646 K is also included in Figure 3, see [Zapata et al. \(2011b\)](#). None of these positions coincide with the submillimeter compact sources reported in this paper. ALMA 1 is not detected in the band 9 observations because, possibly, the observations are too noisy and, in addition, this source is close to the edge of the primary beam.

In Figure 4, we present the spectral energy distributions (SEDs) of the detected compact sources from the millimeter

to submillimeter wavelengths.

4. DISCUSSION AND CONCLUSIONS

A discussion on the nature of the strong sources in the field, Orion Source I and BN at submillimeter wavelengths is outside of the scope of the paper, but we refer to the readers to the papers of [Beuther et al. \(2006\)](#); [Hirota et al. \(2015, 2016\)](#); [Plambeck & Wright \(2016\)](#), where a complete discussion is made using ALMA and SMA data.

As we have observations from 149.5 GHz up to 658.5 GHz of the Orion-KL HMC, we can construct the spectral indices ($S_\nu = \nu^\alpha$, where S_ν is the density flux at different frequencies, ν is the frequency and α , the spectral index) of the compact objects by use of the new ALMA multiple band observations (Figure 4). We estimated the values of the spectral indices using the least-squares power-law technique. The estimated values of spectral indices are presented in Table 2. They are relatively not so steep compared with those reported to be associated with optically thick thermal dust emission (about 2 – 3), but probably it is due to possible grain growth that flattens the spectral indices ([Draine 2006](#)). We thus conclude that possibly we are detecting moderate optically thick thermal dust emission. Taking the values of their flux densities at 658 GHz, we estimated brightness temperatures of about 100 to 200 K. This again suggests that we are seeing dust emission. An estimation of the mass from the dust emission for these submillimeter objects seems not to be a reliable as the emission is moderate optically thick. Finally, we also construct the spectral index for Source I and compared the resulting value (see Table 2) with those reported in the literature, see [Hirota et al. \(2015, 2016\)](#); [Plambeck & Wright \(2016\)](#). The value of 1.8 for spectral index for Source I reported in [Hirota et al. \(2015\)](#) is the closest one to that found in this study, which confirm that the derived spectral indices of the ALMA1-3 continuum sources in this work are reliable. We also have compared the flux densities of the ALMA compact sources reported here with the full and restricted uv-ranges, and find that these are very similar. Moreover, we restricted the images to similar uv-ranges for band 6 and 9 observations, and find not different values for the flux densities presented in Table 2.

We found that ALMA 1 is well coincident (within a 0.8'' error) with the millimeter source found in CARMA observations, FW2011-C14 ([Friedel & Widicus 2011](#)), and the mid-infrared extended object called [RLK73] IRc6E ([Shuping et al. 2004](#)). However, as the mid-infrared is very extended it is difficult to know if they are really connected. For ALMA 2 and ALMA 3, we also find that, within the position errors, these submillimeter objects are coincident with the CARMA millimeter objects FW2011-C14 and FW2011-C22 ([Friedel & Widicus 2011](#)), respectively. We found that ALMA 3 additionally is associated with the compact mid-infrared object [RLK73] IRc12 ([Shuping et al. 2004](#)).

[Robberto et al. \(2005\)](#) reported that [RLK73] IRc12 is a very luminous object with a bolometric luminosity of $4 \times 10^4 L_\odot$, between 7.7 and 12.4 μm . This luminosity corresponds to a high-mass star.

As mentioned before in Figure 3, we have included the positions of the peaks of the hot molecular gas as traced by the $\text{NH}_3(12,12)$ and $\text{CH}_3\text{CN}(12_9,11_9)$, and found that these positions do not coincide with any of the submillimeter objects reported in this study. Even the infrared object [RLK73] IRc12 associated with ALMA3 and that is maybe a high-mass (proto)star, is too far away from the hot molecular gas. Additionally the other IR sources and massive stars are also not associated with hot molecular gas. We thus concluded that the hot molecular gas in the Orion-KL core probably is heated externally.

If the HMC is heated internally, we expect young massive stars in middle of hot molecular gas, but if the heating is externally the core should be illuminated from the edges, this physical effect has been already traced by radio observations made by [Zapata et al. \(2011b\)](#); [Goddi et al. \(2011\)](#). According to the results reported in the present paper, and those from [Zapata et al. \(2011b\)](#); [Goddi et al. \(2011\)](#) an externally heating model seems to fit much better with the HMC in Orion-KL. We thus favour the hypothesis that the hot molecular core in Orion-KL core is heated externally.

This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. L.A.Z. is grateful to CONACyT, Mexico, and DGAPA, UNAM for their financial support. S.-L. Q. is supported by NSFC under grant No. 11373026, and Top Talents Program of Yunnan Province (2015SHA030). T.H. is supported by MEXT/JSPS KAKENHI grant Nos. 21224002, 24684011, and 25108005, and the ALMA Japan Research Grant of NAOJ Chile Observatory, NAOJ-ALMA-0006. We are very thankful for the thoughtful suggestions of the anonymous referee that helped to improve our manuscript. This paper makes use of the following ALMA data:

ADS/JAO.ALMA#2012.1.00146.S,
ADS/JAO.ALMA#2013.1.00048.S,
ADS/JAO.ALMA#2013.1.01034.S,
ADS/JAO.ALMA#2012.1.00123.S.

ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada) and NSC and ASIAA (Taiwan) and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ.

REFERENCES

- Bally, J., Ginsburg, A., Arce, H., et al. 2017, *ApJ*, 837, 60
 Bell, T. A., Cernicharo, J., Viti, S., et al. 2014, *A&A*, 564, A114
 Beuther, H., Zhang, Q., Greenhill, L. J., et al. 2005, *ApJ*, 632, 355
 Beuther, H., Zhang, Q., Reid, M. J., et al. 2006, *ApJ*, 636, 323
 Blake, G. A., Mundy, L. G., Carlstrom, J. E., et al. 1996, *ApJ*, 472, L49
 Chernin, L. M., & Wright, M. C. H. 1996, *ApJ*, 467, 676
 Draine, B. T. 2006, *ApJ*, 636, 1114
 Favre, C., Wootten, H. A., Remijan, A. J., et al. 2011, *ApJ*, 739, L12
 Favre, C., Despois, D., Brouillet, N., et al. 2011, *A&A*, 532, A32
 Friedel, D. N., & Widicus Weaver, S. L. 2011, *ApJ*, 742, 64
 Herbst, E., & van Dishoeck, E. F. 2009, *ARA&A*, 47, 427
 Hirota, T., Tsuboi, M., Fujisawa, K., et al. 2011, *ApJ*, 739, L59
 Hirota, T., Tsuboi, M., Kurono, Y., et al. 2014, *PASJ*, 66, 106
 Hirota, T., Kim, M. K., Kurono, Y., & Honma, M. 2015, *ApJ*, 801, 82
 Hirota, T., Machida, M. N., Matsushita, Y., et al. 2016, *ApJ*, 833, 238
 Ho, P. T. P., Barrett, A. H., Myers, P. C., et al. 1979, *ApJ*, 234, 912
 Genzel, R., Ho, P. T. P., Bieging, J., & Downes, D. 1982, *ApJ*, 259, L103
 Goddi, C., Greenhill, L. J., Humphreys, E. M. L., Chandler, C. J., & Matthews, L. D. 2011, *ApJ*, 739, L13
 Gong, Y., Henkel, C., Thorwirth, S., et al. 2015, *A&A*, 581, A48
 Gooch, R. 1996, *Astronomical Data Analysis Software and Systems V*, 101, 80

- Kurtz, S., Cesaroni, R., Churchwell, E., Hofner, P., & Walmsley, C. M. 2000, *Protostars and Planets IV*, 299
- Liu, S.-Y., Girart, J. M., Remijan, A., & Snyder, L. E. 2002, *ApJ*, 576, 255
- McMullin, J. P., Waters, B., Schiebel, D., Young, W., & Golap, K. 2007, *Astronomical Data Analysis Software and Systems XVI*, 376, 127
- Plambeck, R. L., & Wright, M. C. H. 2016, *ApJ*, 833, 219
- Peng, T.-C., Despois, D., Brouillet, N., Parise, B., & Baudry, A. 2012, *A&A*, 543, A152
- Peng, T.-C., Despois, D., Brouillet, N., et al. 2013, *A&A*, 554, A78
- Peng, Y., Qin, S.-L., Schilke, P., et al. 2017, *ApJ*, 837, 49
- Shuping, R. Y., Morris, M., & Bally, J. 2004, *AJ*, 128, 363
- Robberto, M., Beckwith, S. V. W., Panagia, N., et al. 2005, *AJ*, 129, 1534
- Wilner, D. J., Wright, M. C. H., & Plambeck, R. L. 1994, *ApJ*, 422, 642
- Wright, M. C. H., Plambeck, R. L., & Wilner, D. J. 1996, *ApJ*, 469, 216
- Wright, M. C. H., & Plambeck, R. L. 2017, arXiv:1705.03957
- Zapata, L. A., Ho, P. T. P., Rodríguez, L. F., Schilke, P., & Kurtz, S. 2007, *A&A*, 471, L59
- Zapata, L. A., Schilke, P., & Ho, P. T. P. 2010, *MNRAS*, 402, 2221
- Zapata, L. A., Schmid-Burgk, J., Ho, P. T. P., Rodríguez, L. F., & Menten, K. M. 2009, *ApJ*, 704, L45
- Zapata, L. A., Schmid-Burgk, J., & Menten, K. M. 2011, *A&A*, 529, A24
- Zapata, L. A., Loinard, L., Schmid-Burgk, J., et al. 2011, *ApJ*, 726, L12