Coherence constraints on physical parameters at bright radio sources and FRB emission mechanism

Maxim Lyutikov,¹ Mohammad Z. Rafat,²

Department of Physics and Astronomy, Purdue University, 525 Northwestern Avenue, West Lafayette, IN 47907-2036, USA; lyutikov@purdue.edu
²SIfA, School of Physics, The University of Sydney, NSW 2006, Australia; mohammad.rafat@sydney.edu.au

(Received xx; revised xx; accepted xx)

We discuss physical constrains that observations of high brightness temperature coherent radio emission, with brightness temperatures as high as $T_b \sim 10^{35}$ K, impose on the plasma parameters at relativistically moving astrophysical sources. High brightness temperatures imply a minimal plasma energy density at the source. Additional important constraints come from the fact that resonantly emitting particles lose most of their energy to non-resonant inverse Compton and synchrotron processes. Overall, we find that coherence constraints can be accommodated by plasma parameters in the magnetospheres of neutron stars.

We also interpret recent observations of high-to-low frequency drifting features in the spectra of repeating FRBs as analogues of type-III Solar radio bursts produced by reconnection plasma beams within magnetospheres of highly magnetized neutron stars.

1. Introduction

A number of astrophysical sources show high brightness temperatures, reaching, in case of pulsars and fast radio bursts (FRBs), values of $\sim 10^{35}$ K (e.g. Manchester & Taylor 1977; Melrose 2000; Lorimer et al. 2007), and as high as $\sim 10^{40}$ K in extreme cases (e.g. Soglasnov et al. 2004). These observations imply physical constraints at the emitting plasma that we discuss below.

Though high brightness bursts from Crab and other pulsars have been known for a long time (Staelin & Reifenstein 1968), recent observations of mysterious FRBs (Lorimer et al. 2007; Petroff et al. 2019; Cordes & Chatterjee 2019) impose new constraints on the physical properties at the source. One of the more constraining limitations come from recent identification of a repeating FRB (Spitler et al. 2016). The implied luminosity of $\sim 10^{40}$ erg s⁻¹, taken with short duration \sim few milliseconds and large distances ~ 1 Gpc implies very high energy density at the source. Lyutikov (2017) (see also Lyutikov et al. (2016)) argued that these constraints limits the *loci* of the radio emission generation to neutron stars' magnetospheres.

In this paper we discuss more thoroughly what limitation on the physical parameters at the relativistically moving source (e.g. density and magnetic field) can be obtained from the the observations of high brightness radio emission. In $\S 2$ we discuss general relations, without limiting ourselves to any particular physical location, while in $\S 4$ we apply these to the neutron stars. Finally, in $\S 5.2$ we discuss implications of recent observations of frequency drifts in FRBs.

2. Limits on physical parameters at the sources of high brightness radio emission

2.1. Radiation energy density

The extremely high brightness temperatures in astrophysical radio sources, most likely, involve relativistic plasma so that in the center of momentum frame the spread of the Lorentz factors γ is large, $\gamma \gg 1$. The plasma is likely to be in a magnetic field and additionally the center of momentum frame may move with respect to the observer with relativistic bulk streaming Lorentz factor $\gamma_s \gg 1$. Let us first neglect the effects of the bulk motion, to be added later in §2.3.

Consider a radiation source with observed spectral flux density S_{ν} at distance r from the observer. For an isotropic source we have, Appendix A.1,

$$S_{\nu} = \pi \frac{R^2}{d^2} I_{\nu},\tag{2.1}$$

where R is the source radius and I_{ν} is its spectral brightness. For an unresolved source we may estimate its radius from its variability time τ as $R \approx c\tau$, Appendix A.2, so that $S_{\nu} = \pi (c\tau)^2 I_{\nu}/d^2$. For an anisotropic source (2.1) is further modified, Appedix A.3, to $S_{\nu} = \pi \theta_b^2(c\tau)^2 I_{\nu}/d^2$ where θ_b is half-cone angle of emission region. We may then write, using (2.1) and (A 18),

$$u_r = \frac{\Delta \nu S_{\nu} d^2}{c^3 \tau^2} = \frac{S d^2}{c^3 \tau^2},\tag{2.2}$$

where u_r is the energy density, S is the total flux and $\Delta \nu$ is the frequency range corresponding to central frequency ν with $\Delta/\nu/\nu \ll 1$. The brightness temperature for unresolved unpolarized source is defined as

$$k_B T_b = \frac{c^2 I_{\nu}}{2\nu^2}.$$
 (2.3)

We may then relate the energy density of radiation at the source u_r to the observed brightness temperature, using (A 18), as

$$u_r = \left(\frac{\Delta\nu}{\nu}\theta_b^2\right) \frac{2\pi k_B T_b}{\lambda^3} \tag{2.4}$$

where $\lambda = \nu/c$ is the wavelength, $\Delta \nu/\nu$ is the fractional bandwidth of the receiver, and $\theta_{\rm b}$ is a typical beaming angle of the radiation at the source. The factor in parenthesis is smaller than unity, hence it *relaxes* constraints on the plasma parameters. This factor can in fact be much smaller than unity, with $\theta_{\rm b}^2 \sim 1/\gamma^2$.

To avoid complications about the unknown microphysics of the coherent emission, below we incorporate the factor in parenthesis into brightness temperature

$$\overline{T}_b \equiv \left(\frac{\Delta\nu}{\nu}\theta_b^2\right)T_b. \tag{2.5}$$

The observed brightness temperature is $T_{\rm b}$ while we use $\overline{T}_{\rm b}$ to infer intrinsic plasma parameters. Relation (2.4) then becomes

$$u_r = \frac{2\pi k_B \overline{T}}{\lambda^3} \tag{2.6}$$

Relation (2.6) is one of the key results of the paper: it connects the observed brightness temperature T_B , and the properties of the detector $\Delta\nu/\nu$ and λ to the radiation energy density at the source (parametrized by beaming angle θ_b .

2.2. Required plasma density and magnetic field

Radiation energy density at the source u_r should be a fraction $\eta_r \leq 1$ of emitting particles' energy density,

$$u_r = \eta_r \gamma n m_e c^2, \tag{2.7}$$

where γ is the random Lorentz factor of particles at the source. Values of γ and n refer only to the coherently emitting particles, not the total energy/density. Hence equating (2.4) and (2.7) implies

$$\gamma n = \frac{1}{\eta_r} \frac{2\pi k_B \overline{T}_b}{m_e c^2 \lambda^3} \geqslant \frac{2\pi k_B \overline{T}_b}{m_e c^2 \lambda^3},\tag{2.8}$$

where the inequality follows as $\eta_r \leq 1$. This is the requirement on the plasma energy density at the source given the observed brightness temperature.

Given the radiation flux and the radiation energy density (2.4) one can calculate the equipartition magnetic field B_{eq} (Lyutikov 2017; Lyutikov et al. 2016)) by equating u_r with the magnetic energy density $u_B = B_{eq}^2/8\pi$:

$$B_{eq} = \left(\frac{16\pi^3 k_B}{c^3} \nu^3 \overline{T}_b\right)^{1/2} \approx 5 \times 10^8 \left(\frac{\nu}{1 \text{ GHz}}\right)^{3/2} \left(\frac{\overline{T}_b}{10^{35} \text{ K}}\right)^{1/2} \text{ G}$$
 (2.9)

The equipartition magnetic field is the lower estimate on the magnetic field at the source, so that we can parametrize $B = b_{eq} B_{eq}$ with $b_{eq} \ge 1$.

The magnetic field (2.9) can occur, first of all, in the neutron star magnetospheres, but also in the magnetospheres of white dwarfs and stellar mass black holes. (See also §3 for another requirements on the magnetic fields at the source.)

2.3. Relativistically moving variable sources

If an unresolved source of emission is moving relativistically with Lorentz factor Γ (and the corresponding Doppler factor with respect to the observer δ), and varies with a time scale τ' in the plasma rest frame, then using Lorentz transformations (e.g. Ghisellini 2013)

$$F_{\nu} = \delta^{3} F_{\nu}'$$

$$\nu F_{\nu} = \delta^{4} (\nu F_{\nu})'$$

$$\nu \propto \delta \nu'$$

$$\tau = \delta^{-1} \tau'$$

$$n = \Gamma n'$$

$$T_{b} = \delta^{3} T_{b}'$$

$$u_{r}' = \frac{(\nu F_{\nu})' d^{2}}{c^{3} \tau'^{2}} \delta^{2}$$
(2.10)

(primes denote quantities measured in the plasma rest frame),

Hence

$$k_B \overline{T}_b = \eta_r \frac{\gamma \lambda^3 m_e c^2 n'}{2\pi} \delta^4 \tag{2.11}$$

Or

$$\gamma \delta^4 n' > 2\pi \frac{k_B \overline{T}_b}{m_e c^2 \lambda^3} \tag{2.12}$$

Expression (2.12) is a requirement on internal plasma parameters, γ and n', and bulk mo-

tion Doppler factor δ in terms of the observed brightness temperature T_b and wavelength λ .

Inverting (2.12), we can put a limit on the highest possible frequency where coherent emission of given brightness T_b can be observed:

$$\nu_c \leqslant \left(\frac{m_e n' c^5}{k_B \overline{T}_b}\right)^{1/3} \gamma^{1/3} \delta^{4/3}$$
 (2.13)

Using definition of the brightness temperature this translates to

$$\nu_c \leqslant \frac{n' m_e c^5 \tau^2}{d^2 F_{\nu}} \gamma \delta^4 \tag{2.14}$$

We stress that n' is the plasma (emitting leptons') density in the rest frame; in the observer frame $n = \Gamma n'$.

If the plasma is moving along magnetic field, the estimate (2.9) remains valid in the plasma rest frame. (Otherwise $B' = B/\Gamma$). This immediately implies that $\sim 10^{35} \rm K$ brightens temperatures must come from neutron stars' magnetospheres (Lyutikov et al. 2016).

Using parametrization of the magnetic field at the source (2.9) and the enthalpy density $\approx \gamma n' m_e c^2$, the magnetization parameter σ (Kennel & Coroniti 1984) is

$$\sigma = \frac{B^2}{4\pi \gamma n' m_e c^2} \approx \eta_r b_{eq}^2 \delta^4 \tag{2.15}$$

In a relativistic plasma we expect $\sigma \geq 1$. This requires (i) super-equipartition magnetic field $b_{eq} \geq 1$; (ii) relativistic Doppler motion $\delta \geq 1$; (iii) not highly matter dominated regime - η_r not too small.

3. Nonlinearity parameter a and "normal" radiative losses

A particle emitting coherent radio emission will also experience "normal" losses due to synchrotron and/or Inverse Compton processes. Under certain conditions these normal losses may dominate over losses to coherent emission, further containing the properties of the emission region.

The conventional electromagnetic wave intensity parameter (e.g. Landau & Lifshitz 1975) evaluates to

$$a \equiv \frac{eE'}{2\pi m_e c\nu'} = \frac{e\sqrt{\nu k_B \overline{T}_b}}{m_e c^{5/2} \delta} = 3 \times 10^5 \nu_{GHz}^{1/2} T_{b,35}^{1/2} \delta^{-1}$$

$$\frac{E'^{,2}}{4\pi} = u'_r$$
(3.1)

(E' is the electric field of the wave in the rest frame). Conventionally, in the absence of strong external magnetic field, parameter a is a dimensionless transverse momentum of a particle in the electromagnetic wave, $a \sim p_{\perp}/(m_e c)$.

If a particle oscillates in the electromagnetic fields of the coherent wave with amplitude a, then in addition to the energy losses to the emission of coherent waves it will also suffer Inverse Compton (IC) and synchrotron losses (if magnetic field is present). Typically those "normal" losses will dominated over the "coherent" losses. For example, if an external magnetic field is present, the particle will also emit synchrotron radiation. The synchrotron radiation decay times become shorter than pulse duration in the plasma

frame for

$$\overline{T}_b \geqslant \frac{1}{k_B} \frac{m_e^{8/3} c^7}{e^{10/3} \nu^{7/3} \tau^{2/3}} = 10^{29} b_{eq}^{-4/3} \nu_{GHz}^{-7/3} \tau_{-3}^{-2/3} \,\mathrm{K}$$
(3.2)

In case of IC losses, the estimate (3.2) remains valid with b_{eq} set to unity.

Thus, under "normal" circumstances, for brightness temperatures exceeding (3.2) the coherent electromagnetic wave cannot be sustained: normal losses will drain particles' energy before it has time to emit a coherent wave.

There is an important caveat to the above statement. In large magnetic fields, with $\omega_B \geqslant 2\pi\nu'$ where $\omega_B = eB/(m_ec)$ and ν' is the wave frequency in the source frame, the nature of the particle's motion in the field of the electromagnetic wave changes: instead of oscillations in the electric field of the wave with the dimensionless momentum a a particle experiences slow drift with velocity $v_d/c \sim E'/B$ (here E' is the electric field of the wave, while B is the external magnetic field). This condition translates to

$$B \geqslant \frac{m_e c^2}{e \lambda \delta} \approx 100 \nu_{GHz} \delta^{-1} \,\mathrm{G}$$
 (3.3)

Thus, the presence of high magnetic fields in high brightness relativistic sources is required to avoid large radiative losses of coherently emitting particles to IC and synchrotron processes.

4. Neutron stars' magnetospheres

The relations derived above involve the source number density n' and Lorentz factors γ and γ_s . To proceed further we can parametrize the number density to the expected ones, particularly in the case of neutron star magnetospheres. Two parameterizations are possible: (i) pulsar-like, normalizing the rest frame density to the GJ density (Goldreich & Julian 1969)

$$n'_{pulsar} = \kappa \frac{\Omega B}{2\pi e c \gamma_s} \tag{4.1}$$

where $\kappa \sim 10^3-10^6$ is the observer frame multiplicity (Fawley et al. 1977; Timokhin 2010) and Ω is the neutron star spin; and (ii) magnetar-like (Thompson et al. 2002)

$$n'_{magnetar} = \Delta \phi \frac{B}{2\pi e r \gamma_{\circ}} \tag{4.2}$$

(the last comes from equating curl $B \sim \Delta \phi B/r$ to $(4\pi/c)2nec$, $\Delta \phi$ is a typical twist angle in the magnetar magnetosphere.)

For pulsar-like parametrization (4.1) the condition (2.12) gives

$$\eta_r = \frac{r_e k_B T}{\kappa \gamma \lambda^3 \Omega \omega_B m_e \gamma_s^3}$$

$$\kappa \gamma \gamma_s^3 > \frac{r_e k_B T}{\lambda^3 \Omega \omega_B m_e} = 5 \times 10^{18} \nu_{GHz}^3 P_0^4 T_{b,35} \left(\frac{B}{B_Q}\right)^{-1} \left(\frac{r}{R_{LC}}\right)^3 \tag{4.3}$$

where $B_q = 4 \times 10^{13}$ Gauss is the quantum critical magnetic field and we estimated $\delta \sim 2\gamma_s$ and normalized to period P_0 = one second; $R_{LC} = c/\Omega$ is the light cylinder radius. In the extreme case, a millisecond pulsar with quantum magnetic field producing radiation near the light cylinder, it is required that

$$\kappa \gamma \gamma_s^3 > 6 \times 10^6 T_{b,35} \tag{4.4}$$

This is not too constraining, since we expect $\kappa \sim 10^4 - 10^6$ and $\gamma \sim \gamma_s \sim 10^3 - 10^4$ (e.g. Hibschman & Arons 2001; Timokhin 2010).

Similarly, for Crab giant pulses

$$\kappa \gamma \gamma_s^3 > 3 \times 10^{13} \nu_{GHz}^3 \left(\frac{r}{R_{LC}}\right)^3 T_{b,35},$$
 (4.5)

which can be accommodated with the assumed multiplicity and Lorentz factors even at the light cylinder.

For (millisecond) magnetar-like parametrization (4.2), the condition (2.12) gives

$$\gamma \gamma_s^3 \geqslant \frac{e \nu^3 r k_B T}{m_e c^5 B \Delta \phi} = 10^7 \nu_{GHz}^3 \left(\frac{P}{10^{-3} \text{sec}}\right)^4 \frac{T_{b,35}}{\Delta \phi} \left(\frac{B}{B_O}\right)^{-1} \left(\frac{r}{R_{LC}}\right)^4$$
 (4.6)

Which can also be accommodated with $\gamma \sim \gamma_s \sim$ few hundred.

Finally, the ratio of the local magnetic field to the equipartition magnetic field (2.9) evaluates to

$$b_{eq} = \frac{B_{NS}(r/R_{LC})^3}{B_{eq}} = 3 \times 10^5 \left(\frac{B_{NS}}{B_q}\right) \nu_{GHz}^{-3/2} \left(\frac{r}{R_{LC}}\right)^{-3} T_{b,35}^{-1/2}$$
(4.7)

Though Eq. (4.7) has a number of unknown parameters, there is a large region of allowed $b_{eq} \ge 1$.

5. Origin of FRBs

5.1. Not Giant Pulses from young energetic pulsars

Having narrowed down the most likely location of the FRBs' emission to neutron stars magnetospheres, there are two possible energy sources: rotation and magnetic field. Lyutikov et al. (2016) argued that if the FRBs are analogues of giant pulses (GPs) but coming from young (ages tens to hundreds years) pulsars with Crab-like magnetic field, then the required initial periods need to be in a few msec range - a reasonable assumption for $D \leq$ few hundreds Mpc. Identification of the FRB host with a galaxy at D=1 Gpc makes this possibility unlikely, as we discuss next.

The localization of the Repeating FRB at ~ 1 Gpc (Chatterjee et al. 2017), an order of magnitude further away than what was a fiducial model in Lyutikov et al. (2016), combined with a very steady value of DM (if DM was coming from the newly ejecta SN material), virtually excludes rotationally-powered FRBs, e.g. as analogues of Crab giant pulses (Lyutikov 2017).

To reiterate the argument, the observed radio flux can be parametrized as a fraction $\eta \leq 1$ of the spin-down luminosity L_{sd}

$$\nu F_{\nu} = \eta \frac{L_{sd}}{4\pi d^2},\tag{5.1}$$

The longest possible spin-down time is then

$$\tau_{SD} = \eta \frac{\pi I_{NS}}{2d^2 \nu F_{\nu} P_{min}^2} \approx 600 \,\eta \,\text{yrs}$$

$$(5.2)$$

for $F_{\nu} = 1$ Jy and the minimal period of $P_{min} = 1$ msec. (For a given FRB luminosity L_{FRB} , scaled with spin-down power, the longest spin-down time is for shortest periods and, correspondingly, smallest magnetic fields). The longest possible spin-down time scale (5.2) is barely consistent with constant value of the properties of the Repeating FRB over

the period of few years - that would require an unrealistically high conversion efficiency $\eta \to 1$.

Rotationally-powered radio emission - originating on the open field lines - is also disfavored due to a lack of periodicities in the Repeater(s) bursts (Katz 2018).

5.2. Magnetar magnetospheres?

Repeating FRBs show similar features in their dynamic spectra: downward frequency drifts (Hessels et al. 2019; The CHIME/FRB Collaboration et al. 2019b,a; Josephy et al. 2019). Lyutikov (2020) interpreted these drifts as FRBs' analogues of radius-to-frequency mapping in pulsars and Solar type-III radio burst (but not in a sense of a particular emission mechanism). (Alternatively, spectral drifts are expected in the lensing scenarios (Cordes et al. 2017), though in that case both upward and downward drifts are expected.)

First, we interpret similar frequency behavior as an indication of a some kind of a stiff confining structures - most likely the magnetic field. Narrow spectral features could then be related to the local plasma parameters (e.g. plasma and cyclotron frequencies) or changing resonance conditions. In both case frequency drift then reflects the propagation of the emitting region in changing magnetospheric conditions, similar to what is called "radius-to-frequency mapping" in pulsar research (e.g. Manchester & Taylor 1977; Phillips 1992).

The frequency drift in FRBs are also reminiscent of type-III Solar radio bursts, whereby narrow frequency features show high-to-low temporal evolution (e.g. Fainberg & Stone 1974). Reconnection-driven beams in magnetars magnetospheres though may have different physical conditions than on the open field lines of pulsars' magnetospheres. First, the density is not limited to the Goldreich-Julian value (4.1) and can be much higher (4.2). Magnetic field at the source may be higher than in pulsars (both estimates of plasma densities increase with magnetic field). Finally, the bulk Lorentz factor may be smaller than on the open field lines of pulsar magnetospheres.

If the growth rate is on plasma frequency in the plasma rest frame, $\sim \omega_p'/\sqrt{\gamma}$, the condition of fast growth in the lab frame $\omega_p'/\gamma_s \geqslant \Omega$ translates to

$$\kappa \geqslant \gamma \gamma_s^3 \frac{\Omega}{\omega_B} \approx 10^{-15} \gamma \gamma_s^3 \left(\frac{B_{NS}}{B_Q}\right)^{-1} \left(\frac{P}{10^{-3} \text{sec}}\right)^2 \left(\frac{r}{r_{LC}}\right)^3 \quad \text{, for scaling (4.1)}$$

$$\Delta \phi \geqslant \gamma \gamma_s^3 \frac{\Omega^2 r}{\omega_B c} \approx 10^{-15} \gamma \gamma_s^3 \left(\frac{B_{NS}}{B_Q}\right)^{-1} \left(\frac{P}{10^{-3} \text{sec}}\right)^2 \left(\frac{r}{r_{LC}}\right)^4 \quad \text{, for scaling (4.2)} \quad (5.3)$$

Both these constraints can generally be satisfied in neutron stars magnetospheres. (Same numerical factor in front of expressions in (5.3) is due to the fact that at millisecond periods both estimates of density (4.1) and (4.2) coincide.)

A merger of two Langmuir waves with frequency $\sim \omega_p'/\sqrt{\gamma}$ in the plasma frame will produce observed radiation at

$$\omega \sim \gamma_s \frac{\omega_p'}{\sqrt{\gamma}} = \begin{cases} \left(\kappa \Omega \omega_B \frac{\gamma_s}{\gamma}\right)^{1/2} = 3 \times 10^{11} \left(\kappa \frac{\gamma_s}{\gamma}\right)^{1/2} \left(\frac{r}{r_{LC}}\right)^{-3/2} \text{rad s}^{-1} &, \text{for scaling (4.1)} \\ \left(\Delta \phi \frac{\omega_B c}{r} \frac{\gamma_s}{\gamma}\right)^{1/2} = 3 \times 10^{11} \left(\Delta \phi \frac{\gamma_s}{\gamma}\right)^{1/2} \left(\frac{r}{r_{LC}}\right)^{-2} \text{rad s}^{-1} &, \text{for scaling (4.2)} \end{cases}$$

$$(5.4)$$

(in both cases the numerical estimates are scaled with $(B_{NS}/B_Q)^{1/2} (P/(10^{-3}\text{sec}))^{-2}$.) Both these estimates can generally produce emission at the observed radio wavelengths.

The emission frequencies (5.4) demonstrate downward frequency drift as an emitting entity propagates up in the neutron star magnetosphere. For dipolar magnetic field $\propto r^{-3}$, the scalings are $\omega \propto t^{-3/2}$ and $\omega \propto t^{-2}$ for the two cases (assuming constant Doppler factor; for time-varying Doppler factor $t \to t/\delta(t)^2$).

Let us next list a few observational arguments for and against associating FRBs with magnetar flares. Coherent radio emission can be produced at the initial stage of a "reconnection flare", whereby coherent "kinetic jets" of particles are generated, like the ones in the studies of Crab flares (e.g. Cerutti et al. 2014; Lyutikov et al. 2017, 2018). But there are observational constraints: (i) the SGR 1806 - 20 flare had peak power of 10^{47} erg s⁻¹ (Palmer et al. 2005) but was not seen by Parkes radio telescope (Tendulkar et al. 2016); that puts an upper limit on radio-to-high -energy efficiency $\leq 10^{-6}$. For the Repeater, the first Repeater, the implied high energy luminosity would be then $\geq 10^{47}$ erg s⁻¹. On the other hand, if the Repeater was in our Galaxy the corresponding fluxes would be in GigaJansky, which are clearly not seen. Also in case of PSR J1119-6127 magnetar-like X-ray bursts seem to suppress radio emission (Archibald et al. 2017), but this is probably related to rotationally-driven radio emission, not reconnection-driven. Thus, only some special types of magnetars can produce FRBs.

Finally, associating FRBs with (special kinds of) magnetar flares may resolve the lack of periodicities in the appearance of the FRBs from the Repeater(s): magnetospheric reconnection events appear randomly on closed field lines. This, combined with short, millisecond-like periods, will likely erase the signatures of the rotational period in the observed time sequence of the bursts.

6. Discussion

In this paper we discuss the constraints on the properties of coherent emitting astrophysical sources, having in mind relativistic objects like pulsars and, presumably, FRBs. Several lines of reasoning point to neutron stars' magnetospheres as the origin of the high brightness emission (this is of course known for pulsars, but is important for FRBs).

An important point, besides the estimates of plasma parameters at the source for given observed brightness, is the estimate of non-coherent energy losses, §3. Let us discuss this important argument. Particles at the source lose energy to emission of coherent (resonant) waves and, in addition, to non-resonant interactions (e.g. IC and synchrotron). Typically, the part of the energy that goes to coherent (low frequency - radio) emission is much smaller than the one going to non-resonant interactions (often in optical and Xrays). This was not much of a problem before the identification of FRBs at cosmological distances - the radio emission was always energetically unimportant, subdominant part. For example, even the brightest and rarest giant pulses from Crab reach only 10^{-2} of the total spin-down luminosity. Observations of FRBs, raise the bar, so to say. The implied radio luminosity is some five to nine orders of magnitude larger that is seen in pulsars (In Crab the average radio power is $\sim 10^{32} \text{ erg s}^{-1}$, the peak is $\sim 10^{36} \text{ erg s}^{-1}$; the FRB Repeater is at $\sim 10^{41} \text{ erg s}^{-1}$). These are macroscopically (in astrophysical sense) important powers (Lyutikov 2017) and thus do offer, for the first time from radio observations, a meaningful physical constraints on the plasma parameters at the source. We demonstrated that though these constraints are important, they can be realistically satisfied.

In conclusion, FRB emission properties point to magnetospheres of neutron stars as the origin. Two types of mechanisms can be at work - rotationally or magnetically powered. Rotationally-powered FRB emission mechanisms (e.g. as analogues of Crab giant pulses Lyutikov et al. 2016) are excluded by the localization of the Repeating FRB at ~ 1 Gpc (Spitler et al. 2016), as discussed by Lyutikov (2017) and §5.1. Magnetically-powered emission has some observational constraints, as discussed at the end of §5.2, but remains theoretically viable.

Acknowledgments

This work had been supported by DoE grant DE-SC0016369 and NASA grant 80NSSC17K0757. We would like to thank organizers of SRitp FRB workshop for hospitality, and particularly discussions with Jim Cordes, David Eichler, Victoria Kaspi, Jonathan Katz, Amir Levinson, Yuri Lyubarski and Ue-Li Pen. We would like to thank Maxim Barkov for discussions.

REFERENCES

- ARCHIBALD, R. F., BURGAY, M., LYUTIKOV, M., KASPI, V. M., ESPOSITO, P., ISRAEL, G., KERR, M., POSSENTI, A., REA, N., SARKISSIAN, J., SCHOLZ, P. & TENDULKAR, S. P. 2017 Magnetar-like X-Ray Bursts Suppress Pulsar Radio Emission. *Astrophys. J. Lett.* 849, L20, arXiv: 1710.03718.
- CERUTTI, B., WERNER, G. R., UZDENSKY, D. A. & BEGELMAN, M. C. 2014 Gamma-ray flares in the Crab Nebula: A case of relativistic reconnection?a). *Physics of Plasmas* **21** (5), 056501, arXiv: 1401.3016.
- Chatterjee, S., Law, C. J., Wharton, R. S., Burke-Spolaor, S., Hessels, J. W. T., Bower, G. C., Cordes, J. M., Tendulkar, S. P., Bassa, C. G., Demorest, P., Butler, B. J., Seymour, A., Scholz, P., Abruzzo, M. W., Bogdanov, S., Kaspi, V. M., Keimpema, A., Lazio, T. J. W., Marcote, B., McLaughlin, M. A., Paragi, Z., Ransom, S. M., Rupen, M., Spitler, L. G. & van Langevelde, H. J. 2017 A direct localization of a fast radio burst and its host. *Nature* 541 (7635), 58–61, arXiv: 1701.01098.
- CORDES, JAMES M. & CHATTERJEE, SHAMI 2019 Fast Radio Bursts: An Extragalactic Enigma. arXiv e-prints p. arXiv:1906.05878, arXiv: 1906.05878.
- CORDES, J. M., WASSERMAN, I., HESSELS, J. W. T., LAZIO, T. J. W., CHATTERJEE, S. & WHARTON, R. S. 2017 Lensing of Fast Radio Bursts by Plasma Structures in Host Galaxies. *Astrophys. J.* **842**, 35, arXiv: 1703.06580.
- FAINBERG, J. & STONE, R. G. 1974 Satellite Observations of Type III Solar Radio Bursts at Low Frequencies. *Space Sci. Rev.* 16, 145–188.
- FAWLEY, W. M., Arons, J. & Scharlemann, E. T. 1977 Potential drops above pulsar polar caps Acceleration of nonneutral beams from the stellar surface. *Astrophys. J.* **217**, 227–243.
- GHISELLINI, G., ed. 2013 Radiative Processes in High Energy Astrophysics, Lecture Notes in Physics, Berlin Springer Verlag, vol. 873, arXiv: 1202.5949.
- Goldreich, P. & Julian, W. H. 1969 Pulsar Electrodynamics. Astrophys. J. 157, 869-+.
- Hessels, J. W. T., Spitler, L. G., Seymour, A. D., Cordes, J. M., Michilli, D., Lynch, R. S., Gourdji, K., Archibald, A. M., Bassa, C. G., Bower, G. C., Chatterjee, S., Connor, L., Crawford, F., Deneva, J. S., Gajjar, V., Kaspi, V. M., Keimpema, A., Law, C. J., Marcote, B., McLaughlin, M. A., Paragi, Z., Petroff, E., Ransom, S. M., Scholz, P., Stappers, B. W. & Tendulkar, S. P. 2019 FRB 121102 Bursts Show Complex Time? Frequency Structure. *Astrophys. J. Lett.* 876 (2), L23, arXiv: 1811.10748.
- HIBSCHMAN, J. A. & Arons, J. 2001 Pair Production Multiplicities in Rotation-powered Pulsars. Astrophys. J. 560, 871–884, arXiv: arXiv:astro-ph/0107209.
- Josephy, A., Chawla, P., Fonseca, E., Ng, C., Patel, C., Pleunis, Z., Scholz, P., Andersen, B. C., Bandura, K., Bhardwaj, M., Boyce, M. M., Boyle, P. J., Brar, C., Cubranic, D., Dobbs, M., Gaensler, B. M., Gill, A., Giri, U., Good, D. C., Halpern, M., Hinshaw, G., Kaspi, V. M., Landecker, T. L., Lang, D. A., Lin, H. H., Masui, K. W., Mckinven, R., Mena-Parra, J., Merryfield, M., Michilli, D., Milutinovic, N., Naidu, A., Pen, U., Rafiel-Ravand I, M., Rahman, M., Ransom, S. M., Renard, A., Siegel, S. R., Smith, K. M., Stairs, I. H., Tendulkar, S. P., Vanderlinde, K., Yadav, P. & Zwaniga, A. V. 2019 Chime/Frb Detection of the Original Repeating Fast Radio Burst Source Frb 121102. arXiv e-prints p. arXiv:1906.11305, arXiv: 1906.11305.

- KATZ, J. I. 2018 FRB Sky Distribution, Rarity, Energetics, Magnetic Reconnection and Sources. $arXiv\ e\text{-}prints$, arXiv: 1811.10755.
- KENNEL, C. F. & CORONITI, F. V. 1984 Confinement of the Crab pulsar's wind by its supernova remnant. *Astrophys. J.* **283**, 694–709.
- LANDAU, L. D. & LIFSHITZ, E. M. 1975 The classical theory of fields.
- LORIMER, D. R., BAILES, M., McLaughlin, M. A., Narkevic, D. J. & Crawford, F. 2007 A Bright Millisecond Radio Burst of Extragalactic Origin. *Science* **318**, 777, arXiv: 0709.4301.
- LYUTIKOV, M. 2006 Did Swift measure gamma-ray burst prompt emission radii? MNRAS 369, L5–L8, arXiv: arXiv:astro-ph/0601557.
- LYUTIKOV, M. 2017 Fast Radio Bursts' Emission Mechanism: Implication from Localization. Astrophys. J. Lett. 838, L13, arXiv: 1701.02003.
- LYUTIKOV, MAXIM 2020 Radius-to-frequency Mapping and FRB Frequency Drifts. *Astrophys. J.* **889** (2), 135, arXiv: 1909.10409.
- LYUTIKOV, M., BURZAWA, L. & POPOV, S. B. 2016 Fast radio bursts as giant pulses from young rapidly rotating pulsars. *Mon. Not. Roy. Astron. Soc.* 462, 941–950, arXiv: 1603.02891.
- LYUTIKOV, M., KOMISSAROV, S. & SIRONI, L. 2018 Particle acceleration in explosive relativistic reconnection events and Crab Nebula gamma-ray flares. *Journal of Plasma Physics* 84 (2), 635840201, arXiv: 1804.10291.
- LYUTIKOV, M., SIRONI, L., KOMISSAROV, S. S. & PORTH, O. 2017 Explosive X-point collapse in relativistic magnetically dominated plasma. *Journal of Plasma Physics* 83 (6), 635830601, arXiv: 1805.05233.
- Manchester, R. N. & Taylor, J. H. 1977 Pulsars.
- MELROSE, D. B. 2000 The Status of Pulsar Emission Theory. In IAU Colloq. 177: Pulsar Astronomy 2000 and Beyond (ed. M. Kramer, N. Wex & R. Wielebinski), Astronomical Society of the Pacific Conference Series, vol. 202, pp. 721—+.
- Palmer, D. M., Barthelmy, S., Gehrels, N., Kippen, R. M., Cayton, T., Kouveliotou, C., Eichler, D., Wijers, R. A. M. J., Woods, P. M., Granot, J., Lyubarsky, Y. E., Ramirez-Ruiz, E., Barbier, L., Chester, M., Cummings, J., Fenimore, E. E., Finger, M. H., Gaensler, B. M., Hullinger, D., Krimm, H., Markwardt, C. B., Nousek, J. A., Parsons, A., Patel, S., Sakamoto, T., Sato, G., Suzuki, M. & Tueller, J. 2005 A giant γ -ray flare from the magnetar SGR 1806 20. Nature 434, 1107–1109, arXiv: astro-ph/0503030.
- Petroff, E., Hessels, J. W. T. & Lorimer, D. R. 2019 Fast radio bursts. Astron. Astrophys. Rev. 27 (1), 4, arXiv: 1904.07947.
- Phillips, J. A. 1992 Radio emission altitudes in the pulsar magnetosphere. Astrophys. J. 385, 282–287.
- Soglasnov, V. A., Popov, M. V., Bartel, N., Cannon, W., Novikov, A. Y., Kondratiev, V. I. & Altunin, V. I. 2004 Giant Pulses from PSR B1937+21 with Widths 15 Nanoseconds and $T \geqslant 5 \times 10^{39}$ K, the Highest Brightness Temperature Observed in the Universe. Astrophys. J. **616**, 439–451, arXiv: astro-ph/0408285.
- SPITLER, L. G., SCHOLZ, P., HESSELS, J. W. T., BOGDANOV, S., BRAZIER, A., CAMILO, F., CHATTERJEE, S., CORDES, J. M., CRAWFORD, F., DENEVA, J., FERDMAN, R. D., FREIRE, P. C. C., KASPI, V. M., LAZARUS, P., LYNCH, R., MADSEN, E. C., MCLAUGHLIN, M. A., PATEL, C., RANSOM, S. M., SEYMOUR, A., STAIRS, I. H., STAPPERS, B. W., VAN LEEUWEN, J. & ZHU, W. W. 2016 A repeating fast radio burst. Nature 531, 202–205, arXiv: 1603.00581.
- STAELIN, D. H. & REIFENSTEIN, III, E. C. 1968 Pulsating Radio Sources near the Crab Nebula. Science ${\bf 162},\,1481-1483.$
- Tendulkar, S. P., Kaspi, V. M. & Patel, C. 2016 Radio Nondetection of the SGR 1806–20 Giant Flare and Implications for Fast Radio Bursts. *Astrophys. J.* **827**, 59, arXiv: 1602.02188.
- The CHIME/FRB Collaboration, :, Andersen, B. C., Band ura, K., Bhardwaj, M., Boubel, P., Boyce, M. M., Boyle, P. J., Brar, C., Cassanelli, T., Chawla, P., Cubranic, D., Deng, M., Dobbs, M., Fandino, M., Fonseca, E., Gaensler, B. M., Gilbert, A. J., Giri, U., Good, D. C., Halpern, M., Höfer, C., Hill, A. S., Hinshaw, G., Josephy, A., Kaspi, V. M., Kothes, R., Landecker, T. L., Lang,

D. A., LI, D. Z., LIN, H. H., MASUI, K. W., MENA-PARRA, J., MERRYFIELD, M., MCKINVEN, R., MICHILLI, D., MILUTINOVIC, N., NAIDU, A., NEWBURGH, L. B., NG, C., PATEL, C., PEN, U., PINSONNEAULT-MAROTTE, T., PLEUNIS, Z., RAFIEI-RAVANDI, M., RAHMAN, M., RANSOM, S. M., RENARD, A., SCHOLZ, P., SIEGEL, S. R., SINGH, S., SMITH, K. M., STAIRS, I. H., TENDULKAR, S. P., TRETYAKOV, I., VANDERLINDE, K., YADAV, P. & ZWANIGA, A. V. 2019a CHIME/FRB Detection of Eight New Repeating Fast Radio Burst Sources. arXiv e-prints p. arXiv:1908.03507, arXiv: 1908.03507.

The CHIME/FRB Collaboration, Amiri, M., Bandura, K., Bhardwaj, M., Boubel, P., Boyce, M. M., Boyle, P. J., Brar, C., Burhanpurkar, M., Cassanelli, T., Chawla, P., Cliche, J. F., Cubranic, D., Deng, M., Denman, N., Dobbs, M., Fandino, M., Fonseca, E., Gaensler, B. M., Gilbert, A. J., Gill, A., Giri, U., Good, D. C., Halpern, M., Hanna, D. S., Hill, A. S., Hinshaw, G., Höfer, C., Josephy, A., Kaspi, V. M., Landecker, T. L., Lang, D. A., Lin, H. H., Masui, K. W., Mckinven, R., Mena-Parra, J., Merryfield, M., Michilli, D., Milutinovic, N., Moatti, C., Naidu, A., Newburgh, L. B., Ng, C., Patel, C., Pen, U., Pinsonneault-Marotte, T., Pleunis, Z., Rafiei-Ravandi, M., Rahman, M., Ransom, S. M., Renard, A., Scholz, P., Shaw, J. R., Siegel, S. R., Smith, K. M., Stairs, I. H., Tendulkar, S. P., Tretyakov, I., Vanderlinde, K. & Yadav, P. 2019b A second source of repeating fast radio bursts. Nature 566 (7743), 235–238, arXiv: 1901.04525.

Thompson, C., Lyutikov, M. & Kulkarni, S. R. 2002 Electrodynamics of Magnetars: Implications for the Persistent X-Ray Emission and Spin-down of the Soft Gamma Repeaters and Anomalous X-Ray Pulsars. *Astrophys. J.* **574**, 332–355, arXiv: arXiv:astro-ph/0110677.

Timokhin, A. N. 2010 Time-dependent pair cascades in magnetospheres of neutron stars - I. Dynamics of the polar cap cascade with no particle supply from the neutron star surface. *Mon. Not. Roy. Astron. Soc.* 408, 2092–2114, arXiv: 1006.2384.

Appendix A. Brightness Temperature and radiation energy-density

A.1. Isotropic source

The energy dE from within a solid angle $d\Omega$ passing through a projected area dA in time dt and in a narrow frequency range $d\nu$ is

$$dE = I_{\nu} \, dA \, dt \, d\Omega \, d\nu, \tag{A 1}$$

where I_{ν} is the specific intensity (or spectral brightness). The intensity I_{ν} is a conserved quantity along the ray path.

The spectral flux density F_{ν} , power emitted per unit area per unit frequency, is given by

$$F_{\nu} = \int I_{\nu} \cos \theta \, d\Omega, \quad \text{with} \quad d\Omega = \sin \theta \, d\theta \, d\phi,$$
 (A 2)

using the effective area $\cos \theta \, dA$ in (A 1).

The flux density S_{ν} , the spectral power received per unit area per unit frequency by a detector from a discrete source (i.e. one for which there is a well-defined solid angle) subtending a solid angle Ω_{source} is

$$S_{\nu} = \int_{\Omega_{\text{course}}} I_{\nu} \cos \theta \, d\Omega, \tag{A 3}$$

For an isotropic source of radius R (i.e. a sphere of uniform spectral brightness) located at a distance d from the observer we obtain

$$S_{\nu} = I_{\nu} \int_{0}^{2\pi} d\phi \int_{0}^{\theta_{c}} d\theta \cos\theta \sin\theta = \pi I_{\nu} \sin^{2}\theta_{c} = \pi I_{\nu} \frac{R^{2}}{d^{2}}, \tag{A 4}$$

where $\theta_c = \sin^{-1}(R/d)$. In particular $S_{\nu} = \pi I_{\nu}$ for d = R.

We comment that I_{ν} cannot be obtained for unresolved sources since $\theta_{\rm c}$ cannot be measured. If we know the distance and variability time, then we can estimate $\theta_{\rm c}$.

Planck's law states that for blackbody radiation $I_{\nu} = B_{\nu}$ with

$$B_{\nu} = \frac{2h\nu^3/c^2}{\exp(h\nu/k_BT) - 1} \approx \frac{2k_BT\nu^2}{c^2},$$
 (A 5)

where the approximation applies when $h\nu/k_BT \ll 1$. With a substitution $T \to T_b$ Eq. (A 5) defines the brightness temperature.

For an opaque (can receive radiation from only one hemisphere) isotropic source with spectral brightness I_{ν} , the spectral flux density F_{ν} is

$$F_{\nu} = I_{\nu} \int_0^{2\pi} d\phi \int_0^{\pi/2} d\theta \cos\theta \sin\theta = \pi I_{\nu}. \tag{A 6}$$

Comparing (A 4) and (A 6) we note that for an isotropic source $F_{\nu} = S_{\nu}$ on the surface of the source; they are not equal otherwise.

For isotropic blackbody radiation this implies that

$$F_{\nu} = \pi I_{\nu} = \pi B_{\nu}, \quad S_{\nu} = \pi I_{\nu} \frac{R^2}{d^2} = \pi B_{\nu} \frac{R^2}{d^2}, \quad \text{and} \quad F_{\nu} = S_{\nu} \frac{d^2}{R^2}.$$
 (A 7)

We may write dE in (A 1) as

$$dE = cu_{\nu}(\Omega) \, dA \, dt \, d\Omega \, d\nu, \tag{A 8}$$

where $u_{\nu}(\Omega)$ is the spectral energy density per unit solid angle per unit frequency. Using (A 1) and (A 8) we obtain the relationship

$$I_{\nu} = cu_{\nu}(\Omega). \tag{A 9}$$

The spectral energy density u_{ν} , energy per unit volume per unit frequency, is obtained by integrating over solid angle

$$u_{\nu} = \int u_{\nu}(\Omega) d\Omega = \frac{1}{c} \int I_{\nu} d\Omega$$
, implying $u_{\nu} = \frac{4\pi}{c} I_{\nu}$ (A 10)

for an isotropic source.

The brightness temperature of an isotropic source may be defined using (A5), (A7) and (A10) as

$$k_B T_b = \frac{c^2 I_\nu}{2\nu^2} = \frac{c^2 F_\nu}{2\pi\nu^2} = \frac{c^2 S_\nu}{2\pi\nu^2} \frac{r^2}{R^2} = \frac{c^3 u_\nu}{8\pi\nu^2}.$$
 (A 11)

We may integrate I_{ν} , F_{ν} , S_{ν} and u_{ν} over the observation frequency range $\nu_1 \leqslant \nu \leqslant \nu_2$ to obtain, respectively,

$$I = \frac{2k_B T_b \nu^3}{c^2} \frac{\Delta \nu}{\nu}, \quad F = \pi I, \quad S = \pi \frac{R^2}{r^2} I, \quad u_r = \frac{4}{c} I,$$
 (A 12)

where we define $\nu = (\nu_1 + \nu_2)/2$ and $\Delta \nu = \nu_2 - \nu_1$, and assume that $\Delta \nu / \nu \ll 1$ so that $\nu_2^3 - \nu_1^3 \approx 3\nu^3 (\Delta \nu / \nu)$. The brightness temperature may then be expressed as

$$k_B T_b = \frac{c^2 I}{2\nu^3} \frac{\nu}{\Delta \nu} = \frac{c^2 F}{2\pi \nu^3} \frac{\nu}{\Delta \nu} = \frac{c^2 S}{2\pi \nu^3} \frac{\nu}{\Delta \nu} \frac{r^2}{R^2} = \frac{c^3 u_r}{8\pi \nu^3} \frac{\nu}{\Delta \nu}.$$
 (A 13)

A.2. Unresolved source

The emitted spectral density, F_{ν} , and received spectral density, S_{ν} , are defined in terms of specific intensity, I_{ν} , but with the integral over solid angle performed over different

domain sizes. For a resolved isotropic sources relations (A 7) apply. For an unresolved source F_{ν} and S_{ν} cannot be related through I_{ν} as $\theta_{\rm c}$ cannot defined. However, if we are able to infer the size of the source through another mean then it is possible to relate S_{ν} and F_{ν} . Suppose an isolated (i.e. radiation arriving at the detector is only from the source), unresolved, isotropic source has an inferred radius of $R_{\rm th}$ (e.g. through variability time scale $R_{\rm th} \sim \tau c$) and is at a distance d from the observer. Then its angular size is given by $\theta_{\rm c,th} = \sin^{-1}(R_{\rm th}/r)$. For an unresolved source we make the replacement $\theta_{\rm c} \to \theta_{\rm c,th}$ in (A 11) and (A 13). Of course we have $\theta_{\rm c} = \theta_{\rm c,th}$ for resolved sources.

A.3. Anisotropic source

Pulsars' (and possibly FRBs) emission is beamed and hence anisotropic at the source. Let us assume that within the beam, there are no anisotropies in frequency. We may approximate the anisotropy due to beaming by restricting the emission region to $0 \le \theta \le \theta_b$: emission is from a circular patch of the source surface with angular extent θ_b . For blackbody radiation we may write this as $I_{\nu}(\theta) = 2k_BT(\theta)\nu^2/c^2$ with

$$T_b(\theta) = \begin{cases} T_b, & \text{for } \theta \leqslant \theta_b, \\ 0, & \text{otherwise,} \end{cases}$$
 (A 14)

which implies that

$$F_{\nu} = \frac{2k_B T \nu^2}{c^2} \int_0^{2\pi} d\phi \int_0^{\theta_b} d\theta \cos\theta \sin\theta \approx \pi \theta_b^2 I_{\nu}, \tag{A 15}$$

where the final expression is valid for $\theta_b \ll 1$, and $I_{\nu} = B_{\nu}$ is given by (A 5). Similarly,

$$u_{\nu} = \frac{2\pi}{c} (1 - \cos \theta_{\rm b}) I_{\nu} \approx \frac{\pi \theta_{\rm b}^2}{c} I_{\nu}, \quad \text{and} \quad S_{\nu} = \pi \theta_{\rm b}^2 \frac{R_{\rm th}^2}{r^2} I_{\nu},$$
 (A 16)

we approximate $\theta_{\rm c,th}$ using $\tan \theta_{\rm c,th} = R_{\rm th} \sin \theta_{\rm b}/r$ to obtain $\theta_{\rm c,th} \approx (R_{\rm th}/r)\theta_{\rm b}$, where we assume $\{\theta_{\rm c,th}, \theta_{\rm b}\} \ll 1$. Therefore, for an anisotropic source described by (A 14) the brightness temperature as given by (A 11) and (A 13) is modified to

$$k_B T_b = \frac{c^2 I_{\nu}}{2\nu^2} = \frac{c^2 F_{\nu}}{2\pi \theta_{\rm b}^2 \nu^2} = \frac{c^2 S_{\nu}}{2\pi \theta_{\rm b}^2 \nu^2} \frac{r^2}{R_{\rm tb}^2} = \frac{c^3 u_{\nu}}{2\pi \theta_{\rm b}^2 \nu^2},\tag{A 17}$$

and

$$k_B T_b = \frac{c^2 I}{2\nu^3} \frac{\nu}{\Delta \nu} = \frac{c^2 F}{2\pi \theta_b^2 \nu^3} \frac{\nu}{\Delta \nu} = \frac{c^2 S}{2\pi \theta_b^2 \nu^3} \frac{\nu}{\Delta \nu} \frac{r^2}{R_{tb}^2} = \frac{c^3 u_r}{2\pi \theta_b^2 \nu^3} \frac{\nu}{\Delta \nu}.$$
 (A 18)

Appendix B. Coordinate and observer times, and mini-jets

Let us clarify here the often confusing notions of coordinate times in two systems of references and the laboratory observer time. Relation $\tau = \delta^{-1}\tau'$ is between the *observer* time τ in our, the laboratory, frame and the coordinate time τ' in the plasma rest frame (which is the same as the observer time in the plasma frame). The coordinate time in the plasma rest frame is related to the coordinate time in the laboratory frame τ_{coord} as $\tau' = \tau_{coord}/\gamma_s$. Thus, $\tau = \tau_{coord}/(\delta\gamma_s) \approx \tau_{coord}/(2\gamma_s^2)$, where the last relation approximates $\delta \approx 2\gamma_s$ for relativistic motion along the line of sight. Thus, the relations (2.10) do include the commonly used notion that a relativistically moving emitter that is active for (coordinate time) τ_{coord} produces a pulse in observer time which is $2\gamma_s^2$ shorter.

Along the similar lines of thought, the coherently emitting entity can be a type of mini-jet propagating within the emission regions with internal Lorentz factor γ_i . The

causally connected region, that can in principle supply energy to the jet with duration τ' in the frame of the blob is then $\sim \pi (c\tau')^3 \gamma_j^2$ (Lyutikov 2006). In the observer frame the pulse will still have a duration $\tau = \delta^{-1} \tau'$.

Hence relation (2.11) will be modified as

$$k_B T_b = \eta_r \gamma \lambda^3 m_e c^2 n' \delta^4 \gamma_j^2 \tag{B1}$$

Thus, minijets will further reduce demands on the intrinsic energy density at the source, by $\sim \gamma_j^2$.