## Universal behavior of p-wave proton-proton fusion near threshold

Bijaya Acharya, 1, 2, \* Lucas Platter, 2, 3, † and Gautam Rupak 4, ‡

1 Institut für Kernphysik and PRISMA+ Cluster of Excellence,
Johannes Gutenberg-Universität Mainz, 55128 Mainz, Germany

2 Department of Physics and Astronomy, University of Tennessee, Knoxville, TN 37996, USA

3 Physics Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA

4 Department of Physics & Astronomy and HPC2 Center for Computational Sciences,
Mississippi State University, Mississippi State, MS 39762, USA

We calculate the p-wave contribution to the proton-proton fusion S factor and its energy derivative in pionless effective field theory (EFT) up to next-to-leading order. The leading contributions are given by a recoil piece from the Gamow-Teller and Fermi operators, and from relativistic 1/m suppressed weak interaction operators. We obtain the value of  $(2.5\pm0.3)\times10^{-28}$  MeV fm² for the S factor and  $(2.2\pm0.2)\times10^{-26}$  fm² for its energy derivative at threshold. These are smaller than the results of a prior study that employed chiral EFT by several orders of magnitude. We conclude that, contrary to what has been previously reported, the p-wave contribution does not need to be considered in a high-precision determination of the S factor at astrophysical energies. Combined with the chiral EFT calculation of Acharya  $et\ al.$  [Phys. Lett. B  $\mathbf{760}$ , 584 (2016)] for the s-wave channel, this gives a total threshold S factor of  $S(0)=(4.047^{+0.024}_{-0.032})\times10^{-23}$  MeV fm².

The Sun is powered by nuclear burning of hydrogen, the most abundant element in the universe, into helium. The elementary proton-proton (pp) fusion process that results in a deuteron, a positron and a neutrino is the first step in the chain of reactions producing heavier elements in stellar environments [1]. Solar models for quantities such as core temperatures and neutrino flux are sensitive to the pp fusion cross section. At the relevant solar core temperatures ( $T \approx 1.5 \times 10^7$  K), the Coulomb repulsion and the slow weak process result in a very small cross section. Thus, experimental measurements are prohibitive and non-existent. Theoretical calculations with well-justified uncertainty estimates are essential for providing critical input data for stellar models [2–4]. Inference of solar neutrino masses from terrestrial measurements depends crucially on the pp fusion rate. This reaction involves all the fundamental interactions except gravity. It is important in the field of astro, nuclear and particle physics, and there is an active effort to calculate the cross section with ever higher accuracy and precision [see Reference [3] for an extensive review of the existing literature].

The reaction cross section  $\sigma(E)$  at center-of-mass (c.m.) kinetic energy E is conventionally expressed in terms of the S factor  $S(E) = E \exp(2\pi\eta_p)\sigma(E)$ . The Sommerfeld parameter  $\eta_p = \sqrt{m_p/E} \,\alpha/2$  with proton mass  $m_p = 938.28$  MeV and fine structure constant  $\alpha = 1/137$ . Reference [3] provides the best estimates of  $S(0) = (4.01 \pm 0.04) \times 10^{-23}$  MeV fm<sup>2</sup> at threshold, and  $S'(0)/S(0) = (11.2 \pm 0.1)$  MeV<sup>-1</sup> for the logarithmic derivative. Reference [3] also estimated the contribution of the S''(0) term to be  $\approx 1\%$  at the solar core temperature and recommended that a modern calculation be undertaken. The threshold S factor and its energy derivatives have since been calculated in pionless [5] and chiral [6, 7] effective field theories (EFTs). Refer-

ence [7], the only study so far to have included capture from the p-wave, has claimed that this channel makes a significant contribution to S(E), of roughly the same size as the s-wave S''(0) term, in the astrophysically relevant  $E \approx 10$  keV region. An independent calculation of the p-wave contribution is therefore imperative, especially since the s-wave S(E) has now been constrained to subpercentage precision [6].

EFTs provide a description of interacting particles in terms of only those degrees of freedom that are relevant below a breakdown momentum scale,  $\Lambda$ . Low-energy observables are then calculated as expansions in powers of  $Q/\Lambda$ , where Q is the characteristic momentum of the process under study. Such approaches have been widely used in nuclear physics. They provide a clear guidance on how to systematically construct the nuclear Hamiltonian and couplings to external electroweak sources as perturbation in  $Q/\Lambda$ . They also enable us to use the convergence of the expansion to estimate the uncertainty in theoretical calculations. The pp fusion process at solar energies  $E \lesssim 100 \text{ keV}$  is peripheral, and thus can be accurately described in terms of the incoming pp s-wave phase shift and the outgoing deuteron bound state wave function to within 10% model-independently [1]. Thus the characteristic momentum scale  $Q \approx p, \gamma, 1/a_{pp}, \alpha m_p \ll m_{\pi}$ , where  $p = \sqrt{m_p E} \lesssim 10 \text{ MeV}$ ,  $a_{pp} \approx 25 \text{ MeV}$  is the pp s-wave scattering length,  $\gamma = 45.701$  MeV the deuteron

<sup>&</sup>lt;sup>1</sup> After the authors of Reference [7] were notified about this Rapid Communication, they revisited their calculation and published an Erratum [8] whose p-wave result, albeit much closer, is still not in agreement with this work within our uncertainty estimate. More importantly, as we will later discuss, the result of Reference [8] for the total S factor, with s and p waves included, does not agree with the value we quote here due to basis-truncation issues in their calculation.

binding momentum,  $m_{\pi} \approx 140$  MeV the pion mass. It is therefore appropriate to employ Pionless EFT (#EFT) for the calculation of the pp fusion S factor. This is an EFT with non-relativistic nucleons that interact through short-ranged forces without an explicit pion degree of freedom [9, 10]. Its breakdown scale is  $\Lambda \approx m_{\pi}$  and the perturbative expansion is therefore in  $Q/\Lambda \lesssim 1/3$ . #EFT provides a simple description of pp fusion, to about 10% precision, in terms of nucleon-nucleon observables [11]. Calculation of the fusion rate to a few percent precision requires contribution from two-body currents that represent short-distance physics not constrained by elastic-channel nucleon-nucleon phase shifts [12].

In this Rapid Communication, we present the first calculation of the p-wave contributions to pp fusion in  $\not$ EFT. The results are expressed in terms of model-independent parameters, and, therefore, universal. It provides an important constraint on the precise determination of the solar pp fusion rate and provides insights into further steps that are needed to reduce uncertainties in the future.

Pionless effective field theory: The cross section calculation depends on the strong interaction, the Coulomb repulsion between the two protons, and the weak interaction. The dominant p-wave contribution requires the strong interaction only in the outgoing deuteron ( ${}^3S_1$ ) channel, which is given by [9, 13–15]

$$\mathcal{L}_S = d_i^{\dagger} \left[ \Delta - \left( i \partial_0 + \frac{\nabla^2}{4m} \right) \right] d_i + g_0 \left[ d_i^{\dagger} (N^T P_i N) + \text{h.c.} \right], \tag{1}$$

where m = 938.92 MeV is the isospin-averaged nucleon mass, N represents a nucleon and the vector  $d_i$  represents the deuteron.  $P_i = \sigma_2 \sigma_i \tau_2 / \sqrt{8}$ , where the Pauli matrices  $\sigma$  and  $\tau$  respectively act on spins and isospins, projects the nucleons onto the spin-triplet isospin-singlet  ${}^{3}S_{1}$  (deuteron) channel. The two couplings  $\Delta$ ,  $g_{0}$  are fixed by requiring that the deuteron bound state wave function has the correct exponential decay and normalization constant. In #EFT, this corresponds to ensuring the  ${}^{3}S_{1}$  elastic scattering amplitude has a pole at  $p^{*}=i\gamma$ , and has the correct residue at the said pole. While these depend only on  $\gamma$  at leading order (LO), the contributions of the effective range  $\rho = 1.764$  fm to the residue, which enter at next-to-leading order (NLO), can be expressed in terms of the deuteron wave function renormalization constant,  $Z_d$ , and treated exactly using the zed parameterization [16].

We include the Coulomb interaction between the protons using the t-matrix  $-it_C(E; \boldsymbol{q}, \boldsymbol{p})$  for incoming (outgoing) momentum  $\boldsymbol{p}(\boldsymbol{q})$ . It can be expressed in closed form using the momentum-space Coulomb wave function  $\chi_{\boldsymbol{p}}^{(+)}(\boldsymbol{q})$  as:  $t_C(E; \boldsymbol{q}, \boldsymbol{p}) = (E - q^2/m_p + i0^+)\chi_{\boldsymbol{p}}^{(+)}(\boldsymbol{q})$ . Coulomb amplitude  $t_C$  includes non-perturbative resummation of Coulomb photon exchanges.

The capture from pp p-wave initial state receives contribution from two sets of weak interactions. The first set constitutes the usual Fermi and Gamow-Teller interactions:

$$\mathcal{L}_{W}^{(\text{FGT})} = -\frac{G_{V}}{\sqrt{2}} \left( l_{+}^{0} N^{\dagger} \tau^{-} N + g_{A} \boldsymbol{l}_{+} \cdot N^{\dagger} \boldsymbol{\sigma} \tau^{-} N \right) , \quad (2)$$

where  $G_V$  and  $g_A$  are the vector and axial coupling constants, for which we use the latest Particle Data Group's [17] values of 1.1363(3)  $\times$  10<sup>-11</sup> MeV<sup>-2</sup> and 1.2724(23), respectively.  $l_+^{\mu}$  is the leptonic Dirac current, and  $\tau^- = (\tau_1 - i\tau_2)/2$  is the isospin lowering operator.

The second set of interactions constitutes relativistic p/m effects:

$$\mathcal{L}_{W}^{(\text{rel})} = \frac{G_{V}}{\sqrt{2}} \left[ g_{A} l_{+}^{0} N^{\dagger} \boldsymbol{\sigma} \cdot i \overleftarrow{\nabla}_{2m} \tau^{-} N + \boldsymbol{l}_{+} \cdot N^{\dagger} \left( i \overleftarrow{\nabla}_{2m} \tau^{-} - \mu_{V} \boldsymbol{\sigma} \times \frac{\overline{\nabla}_{2m}}{2m} \tau^{-} \right) N \right], \quad (3)$$

where  $\mu_V = (\mu_p - \mu_n)/2$  denotes the isovector magnetic moment,  $\overrightarrow{\nabla} = \overleftarrow{\nabla} - \overrightarrow{\nabla}$  and  $\overrightarrow{\nabla} = \overleftarrow{\nabla} + \overrightarrow{\nabla}$ .




FIG. 1. Feynman diagram for pp fusion: solid lines nucleons, short-dashed line positron  $e^+$ , dashed line neutrino  $\nu_e$  and double line deuteron. The blob represents Coulomb amplitude  $t_c$ , " $\otimes$ " a weak vertex, " $\blacksquare$ " a strong interaction vertex.

The Feynman diagrams in Fig. 1 provide the dominant p-wave contribution to pp fusion. A straightforward calculation shows that p-wave capture from the weak interaction vertex in Eq. (2) comes from the deuteron recoil momentum k. Thus this p-wave contribution scales as  $kp/Q^2$  compared to the LO s-wave amplitude in #EFT [11, 18]. We name this recoil contribution  $T_{\text{FGT}}$ . The weak interaction vertex generated by the  $\overrightarrow{\nabla}$  terms in Eq. (3) contribute even in the zero-recoil limit. Relative to the LO s-wave amplitude, it is suppressed by a factor of p/m and we name this relativistic contribution  $T_{\rm rel}$ . The contribution from the  $\mu_V \overline{\nabla}$  term is suppressed by  $k^2$  and we do not include it. Compared to the LO s-wave amplitude, at momentum  $p \approx \gamma \approx Q$ , the recoil contribution  $kp/Q^2$  and the relativistic contribution p/m are similar scaling as  $Q^3/\Lambda^3 \approx 0.04$ . We use this estimate for  $T_{\mathrm{FGT}} \approx |T_{\mathrm{rel}}|$  that holds up to  $p \lesssim \gamma$  to keep the EFT analysis simple. Empirically, at solar energies  $E \leq 100 \text{ keV}$ , p/m is small but  $pk/\gamma^2$  is smaller.

Thus  $T_{\rm rel}$  contribution is larger making  $p/m \lesssim 0.01$  to be a better estimate for the relative contribution of the p-wave amplitude. Furthermore, the cross section (and therefore the S factor) can be decomposed into a partial wave expansion as  $\sigma(E) = \sigma_0(E) + \sigma_1(E) + \ldots$ , where the subscript l refers to the l-th pp partial wave. We threfore anticipate the p-wave cross section (and S factor) to be smaller by a factor of  $p^2/m^2 \lesssim 10^{-4}$  compared to the s-wave value. We include the NLO correction from the effective range  $\rho$ . Initial state p-wave strong interactions are suppressed by relative powers of  $Q^3/\Lambda^3$ . Higher order corrections to the weak interactions are suppressed by at least  $Q^2/\Lambda^2 \approx 0.01$ , and constitute a 10% uncertainty in the p-wave cross section.

The p-wave cross section: The p-wave amplitude is

$$i\mathcal{M}_{1} = i8 \frac{G_{v}}{\sqrt{2}} \epsilon_{i}^{d^{*}} u_{N}(-\boldsymbol{p}) \mathbb{P}_{j} u_{N}(\boldsymbol{p})$$

$$\times \left\{ \left( l_{+}^{0} T_{\text{FGT}} - \boldsymbol{l}_{+} \cdot \boldsymbol{T}_{\text{rel}} \right) \operatorname{Tr} \left[ P_{i} \tau^{-} \mathbb{P}_{j}^{\dagger} \right] + g_{A} \left( l_{+}^{k} T_{\text{FGT}} - l_{+}^{0} T_{\text{rel}}^{k} \right) \operatorname{Tr} \left[ P_{i} \sigma_{k} \tau^{-} \mathbb{P}_{j}^{\dagger} \right] \right\}, \quad (4)$$

where  $\mathbb{P}_j$  is the spin-triplet isospin-triplet projector,  $i\sigma_2\sigma_j(1+\tau_3)/4$ . The non-relativistic two component nucleon spinor fields  $u_N(\boldsymbol{p})$  are normalized as  $[u_N(\boldsymbol{p})]_{\alpha}[u_N^*(\boldsymbol{p})]_{\beta}=\delta_{\alpha\beta}$  when summed over polarizations. The amplitudes from the loop integrals are

$$T_{\text{FGT}} = g_0 \sqrt{Z_d} m \int \frac{d^3 q}{(2\pi)^3} \chi_{\mathbf{p}}^{(+)}(\mathbf{q}) \frac{\mathbf{q} \cdot \mathbf{k}}{(\gamma^2 + q^2)^2}, \quad (5)$$

and

$$T_{\rm rel} = g_0 \sqrt{Z_d} m \int \frac{d^3 q}{(2\pi)^3} \chi_{\mathbf{p}}^{(+)}(\mathbf{q}) \frac{\mathbf{q}}{m} \frac{1}{\gamma^2 + q^2}.$$
 (6)

The solid angle integral of  $q\chi_{p}^{(+)}(q)$  picks out the vector direction p and constitutes the l=1 partial wave contribution. The c.m. deuteron momentum k is related to the positron/neutrino pair momenta  $p_{e,\nu}$  from momentum conservation as  $k = -(p_e + p_{\nu})$ . The expressions for  $T_{\text{FGT}} \propto e^{i\delta_1}$  and  $T_{\text{rel}} \propto e^{i\delta_1}$  are derived further below. Since  $T_{\text{FGT}}T_{\text{rel}}^* = T_{\text{FGT}}^*T_{\text{rel}}$ , Eq. (4) gives

$$\overline{|\mathcal{M}_{1}|^{2}} = 8 \left(\frac{G_{v}}{\sqrt{2}}\right)^{2} \left\{ \left(E_{\nu}E_{e} + \boldsymbol{p}_{\nu} \cdot \boldsymbol{p}_{e}\right) \times \left(3|T_{\text{FGT}}|^{2} + 2g_{A}^{2}\boldsymbol{T}_{\text{rel}} \cdot \boldsymbol{T}_{\text{rel}}^{*}\right) + 6\left(\boldsymbol{p}_{\nu} \cdot \boldsymbol{T}_{\text{rel}}\right)\left(\boldsymbol{p}_{e} \cdot \boldsymbol{T}_{\text{rel}}^{*}\right) + 3\left(E_{\nu}E_{e} - \boldsymbol{p}_{\nu} \cdot \boldsymbol{p}_{e}\right)\boldsymbol{T}_{\text{rel}} \cdot \boldsymbol{T}_{\text{rel}}^{*} - \left(6 + 4g_{A}^{2}\right)\left(E_{\nu}\boldsymbol{p}_{e} + E_{e}\boldsymbol{p}_{\nu}\right) \cdot \boldsymbol{T}_{\text{rel}}T_{\text{FGT}}^{*} + 2g_{A}^{2}\left(3E_{\nu}E_{e} - \boldsymbol{p}_{\nu} \cdot \boldsymbol{p}_{e}\right)|T_{\text{FGT}}|^{2} \right\}, \quad (7)$$

where we used the polarization sum over the leptonic currents  $l_{+}^{\mu}l_{-}^{\nu}$ .

The spin averaged cross section for non-relativistic fields is given by Fermi's Golden Rule as

$$\sigma_1(E) = \int \frac{\mathrm{d}^3 p_e}{(2\pi)^3} \frac{\mathrm{d}^3 p_\nu}{(2\pi)^3} \frac{1}{4E_e E_\nu} \frac{1}{v_{rel}} \overline{|\mathcal{M}_1|^2} \times 2\pi \,\delta \left(\delta m + E - \frac{k^2}{2M_d} - E_e - E_\nu\right) , \quad (8)$$

where  $\delta m = 2m_p - M_d = m_p - m_n + \gamma^2 m/(m_p m_n)$ . The integral can be reduced to 4-dimensions. The magnitude  $p_{\nu}$  is constrained from the Dirac  $\delta$ -function. We are free to choose the spin quantization axis ( $\hat{z}$  axis) along p direction. Azimuthal symmetry of the total lepton momentum  $p_e + p_{\nu} = -k$  implies dependence only on the difference in the azimuthal angle  $\phi = \phi_e - \phi_{\nu}$  of the pair  $p_{e,\nu}$ . The integral in Eq. (8) can then be written as

$$\sigma_1(E) = \frac{1}{(2\pi)^4} \int_0^{p_e^{\text{max}}} dp_e p_e^2 \int_{-1}^1 dx_e \int_{-1}^1 dx_\nu \times \int_0^{2\pi} d\phi \frac{p_\nu^2}{|1 + \frac{p_\nu}{M_d} + \frac{p_e}{M_d} x_{e\nu}|} \frac{1}{4E_e E_\nu} \frac{1}{v_{rel}} \overline{|\mathcal{M}_1|^2}, \quad (9)$$

where  $x_{e,\nu} = \hat{\mathbf{p}} \cdot \hat{\mathbf{p}}_{e,\nu}$  and  $x_{e\nu} = \hat{\mathbf{p}}_e \cdot \hat{\mathbf{p}}_{\nu} = x_e x_{\nu} + \sqrt{1 - x_e^2} \sqrt{1 - x_{\nu}^2} \cos \phi$ . The neutrino momentum magnitude is given by

$$p_{\nu} = -M_d - p_e x_{e\nu} + [(M_d + p_e x_{e\nu})^2 + 2M_d (2m_p - M_d + E - E_e) - p_e^2]^{1/2}, \quad (10)$$

and the maximal positron momentum is

$$p_e^{\text{max}} = \left\{ \left( 2 - \frac{2m_p + E}{M_d} \right) \times \left( [2m_p - M_d + E]^2 - m_e^2 \right) \right\}^{1/2}.$$
 (11)

**Results:** The cross section  $\sigma_1(E)$  in Eq. (9) is evaluated by numerical integration using analytic expressions for  $T_{\text{FGT}}$  and  $T_{\text{rel}}$ . These can be derived from the coordinate space wavefunction

$$\chi_{\mathbf{p}}^{(+)}(\mathbf{r}) \equiv \int \frac{d^3q}{(2\pi)^3} e^{i\mathbf{q}\cdot\mathbf{r}} \chi_{\mathbf{p}}^{(+)}(\mathbf{q})$$

$$= \sum_{l=0}^{\infty} (2l+1)i^l e^{i\delta_l} P_l(\hat{\mathbf{r}}\cdot\hat{\mathbf{p}}) \frac{F_l(\eta_p; pr)}{pr} , \qquad (12)$$

where  $\delta_l = \arg \Gamma(l+1+i\eta_p)$  is the Coulomb phase shift and

$$F_{l}(\eta_{p}; \rho) = \frac{2^{l} e^{-\pi \eta_{p}/2} |\Gamma(l+1+i\eta_{p})|}{\Gamma(2l+2)} \rho^{l+1} e^{-i\rho} \times M(l+1-i\eta_{p}, 2l+2, 2i\rho),$$
(13)

is the regular Coulomb wave function expressed in terms of the conventionally defined Kummer's function M(a, b, z). Equation (5) can then be written as

$$T_{\text{FGT}} = \frac{1}{6} g_0 \sqrt{Z_d} m(\boldsymbol{p} \cdot \boldsymbol{k}) e^{i\delta_1} e^{-\pi \eta_p/2} |\Gamma(2 + i\eta_p)|$$

$$\times \int_0^\infty dr r^3 e^{-(\gamma + ip)r} M(2 - i\eta_p, 4, i2pr)$$

$$= -\sqrt{\frac{8\pi \gamma}{1 - \rho \gamma}} e^{i\delta_1} e^{-\pi \eta_p/2} (\boldsymbol{p}_\nu + \boldsymbol{p}_e) \cdot \boldsymbol{p}$$

$$\times |\Gamma(2 + i\eta_p)| \frac{1}{(\gamma^2 + p^2)^2} e^{2\eta_p \arctan p/\gamma}, \quad (14)$$

where we have used the NLO relation  $g_0\sqrt{Z_d}m = \sqrt{8\pi\gamma/(1-\rho\gamma)}$ . Similarly, Eq. (6) can be written as

$$T_{\rm rel} = \frac{1}{3} g_0 \sqrt{Z_d} m e^{i\delta_1} e^{-\pi \eta_p/2} |\Gamma(2+i\eta_p)| \frac{\mathbf{p}}{m}$$

$$\times \int_0^\infty dr r (1+\gamma r) e^{-(\gamma+ip)r} M(2-i\eta_p, 4, i2pr)$$

$$= \sqrt{\frac{8\pi\gamma}{1-\rho\gamma}} e^{i\delta_1} e^{-\pi\eta_p/2} \frac{\mathbf{p}}{m} |\Gamma(2+i\eta_p)|$$

$$\times \frac{1}{2p^2 + 2p^2\eta_p^2}$$

$$\left[1 + \frac{p^2 + 2p\eta\gamma - \gamma^2}{\gamma^2 + p^2} e^{2\eta_p \arctan p/\gamma}\right]. \tag{15}$$

In Fig. 2 we show the result for the S-factor  $S_1(E)$ . We perform a polynomial fit to the results shown in Fig. 2 and use it to extrapolate the S factor and its derivative to zero energy. We obtain

$$S_1(0) = (2.47 \pm 0.25 \pm 0.01) \times 10^{-28} \text{ MeV fm}^2,$$
  
 $S_1'(0) = (2.16 \pm 0.22 \pm 0.01) \times 10^{-26} \text{ fm}^2,$  (16)

where the first errors indicate EFT uncertainties and the second ones are numerical errors from polynomial fits to S(E).

Our result for  $S_1(0)$  agrees with the tentative estimates we made earlier based on the power counting, but does not agree with the value of  $S_1(0) = 2.0 \times 10^{-25} \text{ MeV fm}^2$ claimed in Reference [7]. In fact, the p-wave contribution is much smaller than the  $\approx 1\%$  contribution obtained by Reference [7] in the entire energy region in which they perform their calculations. We, therefore, disagree with the findings of Marcucci et al. in Reference [7] and claim that the p-wave contributions need not be considered in the calculation of the pp S factor at astrophysically relevant energies since these are much smaller than the precision of the s-wave calculation [see Reference [6] for a state-of-the-art uncertainty analysis]. Furthermore, Refs. [6, 19] have found that basis truncation errors accounted for a reduction in Reference [7]'s s-wave S factor by about 0.7 %. Since Marcucci et al. only addressed the error in the p-wave calculation in their




FIG. 2. The p-wave S-factor  $S_1(E)$ .

Erratum [8], their revised value for the S factor, with combined s and p waves, is still incorrect, and does not agree with value  $S(0) = (4.047^{+0.024}_{-0.032}) \times 10^{-23} \text{ MeV fm}^2$  calulated by Reference [6] within the uncertainty band, which remains unmodified upon inclusion of the p-wave contribution calculated in this Rapid Communication. Finally, we emphasize that, even though the p-wave numbers we calculated are negligible given the large uncertainty in the s-wave value, the correction we make to Reference [7]'s results is at least as important as all sources of uncertainty combined.

**Conclusion:** We calculated for the first time the contribution of p-wave pp configuration to the fusion rate in #EFT. This analysis was motivated by a recent calculation with chiral potentials that suggested that the leading p-wave contributions are comparable to the next-to-next-to-leading s-wave contributions.

We determined the dominant Feynman diagrams contributing to the p-wave S factor and calculated their contribution at low energies. The NLO calculation includes the recoil contributions from the Gamow-Teller and Fermi operators as well as the relativistic 1/m suppressed weak interaction operators. We found that the p-wave contribution to the pp fusion S factor is smaller than the value obtained in Reference [7] by several orders of magnitude, and that the effect of p-wave fusion is therefore negligible for a high-precision determination.

Our analytic results for the p-wave fusion matrix element can serve as a benchmark for numerical calculations of chiral effective theory matrix elements. They

<sup>&</sup>lt;sup>2</sup> The relationship between the chiral EFT counterterms  $c_D$  and  $d_R$  have since been updated [20]. This correction makes a negligible modification in the S-factor value compared to the uncertainty band [21].

are expressed in terms of the weak couplings constants  $g_A$  and  $G_F$  and two observables from the strong sector, the deuteron binding energy and wave function renormalization and therefore do not suffer from any short-distance model ambiguities. We note that the input observables needed to predict the p-wave fusion rate are s-wave observables. Our results illustrate furthemore the importance of calculating astrophysically relevant three-nucleon capture reactions in pionless effective field theory to reduce currently accepted uncertainties and to explore the importance of recoil corrections in these nuclei.

## ACKNOWLEDGEMENTS

This work was supported by the U.S. National Science Foundation under Grants PHY-1555030 and PHY-1615092, by the Office of Nuclear Physics, U.S. Department of Energy under Contract No. DE-AC05-00OR22725, and by the Deutsche Forschungsgemeinschaft through The Low-Energy Frontier of the Standard Model CRC (SFB 1044) and through the PRISMA<sup>+</sup> Cluster of Excellence. GR acknowledges support from the JINPA at Oak Ridge National Laboratory and The University of Tennessee, during his sabbatical where part of this research was completed. LP and GR thank the GSI-funded EMMI RRTF workshop "ER15-02: Systematic Treatment of the Coulomb Interaction in Few-Body Systems" participants for valuable discussions and hosts for their hospitality.

- \* acharya@uni-mainz.de
- $^{\dagger}$ lplatter@utk.edu
- <sup>‡</sup> grupak@ccs.msstate.edu
- H. A. Bethe and C. L. Critchfield, Phys. Rev. 54, 248 (1938).

- [2] E. G. Adelberger et al., Rev. Mod. Phys. 70, 1265 (1998), arXiv:astro-ph/9805121 [astro-ph].
- [3] E. G. Adelberger et al., Rev. Mod. Phys. 83, 195 (2011), arXiv:1004.2318 [nucl-ex].
- [4] N. Vinyoles, A. M. Serenelli, F. L. Villante, S. Basu, J. Bergstrm, M. C. Gonzalez-Garcia, M. Maltoni, C. Pea-Garay, and N. Song, Astrophys. J. 835, 202 (2017), arXiv:1611.09867 [astro-ph.SR].
- [5] J.-W. Chen, C. P. Liu, and S.-H. Yu, Phys. Lett. B720, 385 (2013), arXiv:1209.2552 [nucl-th].
- [6] B. Acharya, B. D. Carlsson, A. Ekström, C. Forssén, and L. Platter, Phys. Lett. B760, 584 (2016), arXiv:1603.01593 [nucl-th].
- [7] L. E. Marcucci, R. Schiavilla, and M. Viviani, Phys. Rev. Lett. 110, 192503 (2013), arXiv:1303.3124 [nucl-th].
- [8] L. E. Marcucci, R. Schiavilla, and M. Viviani, Phys. Rev. Lett. **123**, 019901(E) (2019).
- [9] U. van Kolck, Nucl. Phys. A645, 273 (1999), arXiv:nuclth/9808007.
- [10] J.-W. Chen, G. Rupak, and M. J. Savage, Nuclear Physics A 653, 386412 (1999).
- [11] X. Kong and F. Ravndal, Nucl. Phys. A656, 421 (1999), arXiv:nucl-th/9902064 [nucl-th].
- [12] M. Butler and J.-W. Chen, Phys. Lett. B520, 87 (2001), arXiv:nucl-th/0101017 [nucl-th].
- [13] D. B. Kaplan, M. J. Savage, and M. B. Wise, Nucl. Phys. B 534, 329 (1998), nucl-th/9802075.
- [14] D. B. Kaplan, M. J. Savage, and M. B. Wise, Phys. Lett. B424, 390 (1998), arXiv:nucl-th/9801034.
- [15] D. B. Kaplan, M. J. Savage, and M. B. Wise, Phys. Rev. C59, 617 (1999), arXiv:nucl-th/9804032.
- [16] D. R. Phillips, G. Rupak, and M. J. Savage, Phys. Lett. B473, 209 (2000), arXiv:nucl-th/9908054.
- [17] M. Tanabashi *et al.* (Particle Data Group), Phys. Rev. D 98, 030001 (2018).
- [18] X. Kong and F. Ravndal, Phys. Rev. C64, 044002 (2001), arXiv:nucl-th/0004038 [nucl-th].
- [19] B. Acharya, A. Ekström, D. Odell, T. Papenbrock, and L. Platter, Phys. Rev. C95, 031301(R) (2017), arXiv:1608.04699 [nucl-th].
- [20] D. Gazit, S. Quaglioni, and P. Navratil, Phys. Rev. Lett. 103, 102502 (2009); 122, 029901(E) (2019).
- [21] B. Acharya, A. Ekström, and L. Platter, Phys. Rev. C98, 065506 (2018), arXiv:1806.09481 [nucl-th].