

ALMA OBSERVATIONS OF THE EXTRAORDINARY CARINA PILLARS: HH 901/902

GEOVANNI CORTES-RANGEL, LUIS A. ZAPATA, JESÚS A. TOALÁ
Instituto de Radioastronomía y Astrofísica, Universidad Nacional Autónoma de México,
P.O. Box 3-72, 58090, Morelia, Michoacán, México.

PAUL T. P. HO
Academia Sinica Institute of Astronomy and Astrophysics, PO Box 23-141, Taipei, 10617, Taiwan.
East Asian Observatory, 666 N. A'ohoku Place, Hilo, Hawaii 96720, USA.

SATOKO TAKAHASHI
Joint ALMA Observatory, Alonso de Cordova 3108, Vitacura, Santiago, Chile.
NAOJ Chile Observatory, Alonso de Cordova 3108, Vitacura, Santiago, Chile.
Department of Astronomical Science, School of Physical Sciences, SOKENDAI, Mitaka, Tokyo 181-8588, Japan.

ADAL MESA-DELGADO
Instituto de Astrofísica, Facultad de Física, Pontificia Universidad Católica de Chile,
Av. Vicuña Mackenna 4860, 782-0436 Macul, Santiago, Chile.

JOSEP M. MASQUÉ
Departamento de Astronomía, Universidad de Guanajuato, Apdo. Postal 144, 36000 Guanajuato, México.

ABSTRACT

We present Atacama Large Millimeter/Submillimeter Array (ALMA) 1.3 mm continuum and C¹⁸O(2–1), N₂D⁺(3–2), ¹³CS(5–4), and ¹²CO(2–1) line sensitive and high angular resolution ($\sim 0.3''$) observations of the famous carina pillars and protostellar objects HH 901/902. Our observations reveal for the first time, the bipolar CO outflows and the dusty disks (plus envelopes) that are energizing the extended and irradiated HH objects far from the pillars. We find that the masses of the disks+envelopes are about 0.1 M_⊙ and of the bipolar outflows are between 10^{−3} - 10^{−4} M_⊙, which suggests that they could be low- or maybe intermediate- mass protostars. Moreover, we suggest that these young low-mass stars are likely embedded Class 0/I protostars with high-accretion rates. We also show the kinematics of the gas in the pillars together with their respective gas masses (0.1 – 0.2 M_⊙). We estimate that the pillars will be photo-evaporated in 10⁴ to 10⁵ years by the massive and luminous stars located in the Trumpler 14 cluster. Finally, given the short photo-evaporated timescales and that the protostars in these pillars are still very embedded, we suggest that the disks inside of the pillars will be quickly affected by the radiation of the massive stars, forming proplyds, like those observed in Orion.

Keywords: molecular data – techniques: interferometric – ISM: individual objects (HH 901, HH 902)
– ISM: jets and outflows

1. INTRODUCTION

The Carina Nebula is one most active star forming regions in our Galaxy. This region contains nearly one hundred O-type stars and tens of thousands of lower-mass young stars (Walborn 1973, 1995; Massey, & John-

son 1993; Smith 2006; Povich et al. 2011; Preibisch et al. 2011; Feigelson et al. 2011), with the Trumpler (Tr) 14 and 16 massive clusters being one the most luminous regions within Carina. The clusters Tr 14 and 16 have ultraviolet (UV) luminosities about 20 and 60 times higher

than θ 1 Ori C located in the Orion Nebula (Smith & Brooks 2008). Hence, the Carina Nebula is an ideal laboratory to study the time-scale of the photo-evaporation of disks, envelopes, and molecular pillars located especially in the core of the massive clusters.

As a clear case of irradiated/photo-evaporating objects, using APEX FLASH+ and CHAMP+ line observations, Sahai et al. (2012) reported an irradiated object with a cometary-shaped (CN 105–600) in the Carina star-forming region. They found that this externally irradiated object is not a classical protoplanetary disk as those reported in the Orion Nebula (O’Dell et al. 1993, 2008; Henney & O’Dell 1999), this object is instead a large and massive molecular globule ($0.1 - 0.3 M_{\odot}$) in which very compact disks and outflows are embedded. The young star inside of this globule has a mass of about $0.5 M_{\odot}$ and total luminosity of about $10 L_{\odot}$ (Sahai et al. 2012), corresponding to a low-mass young star. Haikala et al. (2017) also reported APEX observations of several globulettes in Carina, these globulettes are compact and denser than objects of similar mass in the Rosette nebula.

There are however many more well studied irradiated globules in the Carina Nebula, they include HH 666, HH 1066, HH 900, and HH 1010 (Reiter & Smith 2013). But any of them include interferometric millimeter observations. Given that different young massive star forming regions have a very varied morphology, ages, and number of massive stars, these regions hold distinct kinds of irradiated interfaces (or globulettes) and their study is important because this will allow us to better understand the evolution of the young stars. A clear example is found in the Carina region (Hartigan et al. 2012), which contains a rich array of walls, pillar and globules and in the older Cyg OB2 association is found mostly isolated globules.

Very recently, Mesa-Delgado et al. (2016) using ALMA (Atacama Large Millimeter/Submillimeter Array) Band 6 Long Baseline continuum observations reported the first direct imaging of protoplanetary disks in the star-forming region of Carina and studied their possible external strong photo-evaporation. They detected a few protoplanetary disks (CN104-593 and CN105-600) in the highly obscured globules within the Carina Nebula and found no signs of disks near to the massive clusters Tr 14 and Tr 16. The disks have an average size of 120 au and total masses of 30 and $50 M_{Jup}$. The non-detection of millimeter emission above the 4σ threshold ($\sim 7M_{Jup}$) in the core of the massive cluster Tr 14, suggest evidence for rapid photo-evaporative disk destruction in the cluster’s harsh radiation field.

Bipolar molecular outflows and collimated optical jets are one of the phenomena associated with star formation (Bally 2016). The outflows are composed of swept-

up molecular shells energized by neutral and atomic jets containing molecules. As these molecular outflows escape from their parent molecular clouds to scales of some parsecs, they become mostly atomic or ionized and traced by Herbig-Haro (HH) objects revealed at optical wavelengths.

The HH 901/902 were first reported by Smith et al. (2010) using *Hubble Space Telescope (HST)*/Advanced Camera for Surveys H α observations. They catalogued the HH objects as irradiated bipolar jets. These objects are extremely bright and clear in their *HST* images (see their Figure 5). The dusty pillars from which HH 901 and 902 emerge are between Tr 14 and 16 clusters. Both HH objects axis are clearly perpendicular to the dusty pillars and are located to a distance of about 1.5 arcmin. to the south of the Tr 14 and about 7 arcmin. from Tr 16. These flows consist of a highly collimated (with opening angles of only a few degrees) chain of H α -emitting knots with an extent of 16 arcsec (0.18 pc) for the HH 901 and an extent of 38 arcsec (0.42 pc) for the HH 902. Both flows are strongly irradiated by the massive clusters Tr 14 and 16. HH 901 and HH 902 both lie close to plane of the sky with tilt angles of about 20° (Reiter & Smith 2014). Reiter & Smith (2013, 2014) suggested that these two jets are energized by intermediate-mass stars ($2-8 M_{\odot}$). However, these studies could not corroborate this interpretation due to the fact that the authors reported that the driving sources were not detected in the IR, probably because they were too embedded and because of inadequate angular resolution. Given these problems, the fits to the IR SEDs (Spectral Energy Distributions) become very poor and the estimated luminosity is uncertain.

The HH 901/HH 902 objects have also been detected in H $_2$ ($2.2 \mu m$) and Br γ spectral lines by Hartigan et al. (2015). These observations represented the first detection of the molecular material associated with the HH objects. In particular, the H $_2$ emission is tracing the innermost parts of the outflow as can be seen in Figures 19 and 20 of Hartigan et al. (2015). Molecular material far from the exciting objects is probably destroyed by the UV radiation.

In this study, we have carried out new ALMA observations of the HH 901/HH 902 objects in order to better characterize their optical jets, disks, and molecular pillars. We reveal their compact molecular outflows, dusty disk, and envelopes, together with the kinematics of the pillars. We describe these results in more detail in the next sections.

2. OBSERVATIONS

The observations of the HH 901/902 protostellar objects were carried out with the ALMA at Band 6 in 2018 January 1st and 4th (C43-6), and April 20th (C43-




Figure 1. ALMA CO(2–1) moment zero (blue and red colors) and millimeter continuum (contours) images of the HH 901 object overlaid in a *HST*(H α) optical image (green colors). The blue and red color represent the blue-shifted and red-shifted CO emission, respectively, from the HH 901 outflow. The contours range from 50% to 90% of the peak emission, in steps of 20%. The peak of the millimeter continuum emission is $2.8 \text{ mJy Beam}^{-1}$. Here, we are only contouring the most compact 1.3 mm emission from our observations revealing the envelope and the disk. There is still some extended emission at low levels. The half-power contour of the synthesized beam of the line image is shown in the bottom-left corner. We integrate in radial velocities from $+6.4$ to -1.2 km s^{-1} for the redshifted emission, and from -8.2 to -14.6 km s^{-1} for the blueshifted emission for the CO outflow in the HH 901 object. The LSR systemic velocity of the entire pillar associated with the object HH 901 is about -5.0 km s^{-1} .

3) as part of the Cycle 5 program 2017.1.00912.S. The observations used 43 antennas with a diameter of 12 m, yielding baselines with projected lengths from 15 – 2516 m (11.5 – 1935 k λ) for C43-6 and 15 – 500 m (11.5 – 384.6 k λ) for C43-3. The primary beam at this frequency has a full width at half-maximum (FWHM) of about $25''$, so that each object was covered with a single pointing at the sky position $\alpha(J2000) = 10^h 44^m 1^s.7$; $\delta(J2000) = -59^{\circ} 30' 32.0''$ for the HH 902, and $\alpha(J2000) = 10^h 44^m 3^s.5$; $\delta(J2000) = -59^{\circ} 31' 2.0''$ for the HH 901. The integration time on source (HH 901 and 902) for the C43-6 configuration was about 9 min.

and for the C43-3 configuration is about 3.5 min.

The continuum images were obtained by averaging line-free spectral channels of four spectral windows centered at: 218.014 GHz(spw0), 219.578 GHz(spw1), 231.239 GHz(spw2), and 230.556 GHz(spw3). The total bandwidth for the continuum is about 4.3 GHz. These four spectral windows were centered to observe different molecular lines as the C 18 O(2–1) ($\nu_{\text{rest}} = 219.56035 \text{ GHz}$), $^{13}\text{CS}(5-4)$ ($\nu_{\text{rest}} = 231.22068 \text{ GHz}$), N $_2$ D $^+(3-2)$ ($\nu_{\text{rest}} = 231.32186 \text{ GHz}$), and $^{12}\text{CO}(2-1)$ ($\nu_{\text{rest}} = 230.53800 \text{ GHz}$). We detected emission only from the lines C 18 O, N $_2$ D $^+$, and ^{12}CO with a channel spacing




Figure 2. ALMA CO(2–1) moment zero (contours) and one (colors) maps of the pillar and outflow HH 901. The contours range from 10% to 90% of the peak emission, in steps of 10%. The peak of the millimeter line CO emission is $1.8 \text{ Jy Beam}^{-1} \text{ km s}^{-1}$. The half-power contour of the synthesized beam of the line image is shown in the bottom-left corner. The LSR radial velocity scale-bar is shown at the right. The brown circle marks the position of the disk revealed in these ALMA observations for the HH 901 object. The red and blue arrows trace the orientation of the outflow. For the CO moment zero image, we integrate in radial velocities from -15.9 to $+6.4 \text{ km s}^{-1}$. The LSR systemic velocity of the entire pillar associated with the object HH 901 is about -5.0 km s^{-1} .

of 0.63 km s^{-1} . The thermal emission from these species can be seen in Figures 1, 2, 3, 4, 5, and 6.

The weather conditions were very good for these observations and stable with an average precipitable water vapor between 1.3 and 2.2 mm and an average system temperature around 90 K. The ALMA calibration included simultaneous observations of the 183 GHz water line with water vapor radiometers, used to reduce atmospheric phase fluctuations. Quasars, J0904-5735 and J1107-4449 were used for the bandpass and flux calibrations. J1032-5917 was used for correcting the gain fluctuations.

The data were calibrated, imaged, and analyzed using the Common Astronomy Software Applications (CASA) Version 5.1. Imaging of the calibrated visibilities was done using the CLEAN and TCLEAN tasks. We concatenated the data from both dates for the C43-6 and

C43-3 configurations with the CONCAT task. We used the ROBUST parameter of CLEAN in CASA set to natural. The resulting image rms noise for the continuum was $50 \mu\text{Jy beam}^{-1}$ at a angular resolution of $0.32'' \times 0.27''$ with a PA of -65° for both images. The ALMA theoretical rms noise for this configuration, integration time, and frequency is about $43 \mu\text{Jy beam}^{-1}$, which is very close to the value we obtain in the continuum images. For the line image rms noise we obtained a value of $6.5 \text{ mJy beam}^{-1} \text{ km s}^{-1}$ at a angular resolution of $0.31'' \times 0.26''$ with a PA of -64° . The ROBUST parameter was set also to natural. The ALMA theoretical rms noise for this configuration, integration time, and frequency is about $4.3 \text{ mJy beam}^{-1}$, which is very close to the value we obtain in the line images. Phase self-calibration was attempted on the continuum images but the resulting maps did not improved significantly, so at this point we did not used the self-calibrated maps. We did not include the channel maps because they do not add extra information from that obtained from the moments maps.

3. RESULTS AND DISCUSSION

3.1. HH 901

In Figure 1 we present the *HST/ACS* image from Smith et al. (2010) of the optical jet HH 901 overlaid with the ALMA high velocity CO(2–1) and the 1.3 mm continuum emission. The CO emission reveals a bipolar and collimated outflow that is tracing the most inner part of the optical jet in an east-west orientation (with an approximate size of 5000 au). This map shows that the blue-shifted emission is located toward an eastern orientation, while red-shifted emission is to its western side. We integrate in radial velocities from $+6.4$ to -1.2 km s^{-1} for the redshifted emission, and from -8.2 to -14.6 km s^{-1} for the blueshifted emission for the CO outflow in the HH 901 object. The CO emission between -1.2 km s^{-1} to -8.2 km s^{-1} is associated with the dusty pillar as is shown in Figure 2. The LSR systemic velocity of the entire pillar associated with the object HH 901 is about -5.0 km s^{-1} (Figure 3). The radial velocities of the innermost CO outflow are in agreement to those presented in Reiter & Smith (2014). They found, using slit positions of the optical [SII] and $\text{H}\alpha$ toward the HH 901, red-shifted emission in the west side of the flow with radial velocities about $+25.0 \text{ km s}^{-1}$.

The 1.3 mm continuum image traces the disk and possibly part of the envelope at slightly larger scales. From a gaussian fitting to the continuum source from a map obtained using ROBUST parameter equal to 0.5 (yielding a better angular resolution), we obtained a deconvolved size of $0.49'' \pm 0.03'' \times 0.33'' \pm 0.03''$ with a PA of $100^\circ \pm 7^\circ$ and an integrated flux of $6.8 \pm 0.4 \text{ mJy}$




Figure 3. ALMA $\text{C}^{18}\text{O}(2-1)$ (left panel) and $\text{N}_2\text{D}^+(3-2)$ (right panel) moment one maps overlaid with a contour image of the $\text{CO}(2-1)$ moment zero of the pillar and outflow HH 901. The contours range from 10% to 90% of the peak emission, in steps of 10%. The peak of the millimeter line CO emission is $1.8 \text{ Jy Beam}^{-1} \text{ km s}^{-1}$. The half-power contour of the synthesized beam of the line image is shown in the bottom-left corner of each panel. The LSR radial velocity scale-bar is shown at the right. The brown circle marks the position of the disk+envelope revealed in these ALMA observations for the HH 901 object.

Beam^{-1} . The corresponding physical sizes of these deconvolved values are about 700 au, which suggests that we are also seeing part of the envelope and disk. Assuming that the dust emission is optically thin and isothermal, the dust mass (M_d) is directly proportional to the flux density (S_ν) integrated over the source, as:

$$M_d = \frac{D^2 S_\nu}{\kappa_\nu B_\nu(T_d)}, \quad (1)$$

where D is the distance to Carina Nebula ($2.3 \pm 0.1 \text{ kpc}$, [Smith & Brooks 2008](#)), κ_ν the dust mass opacity, and $B_\nu(T_d)$ the Planck function for the dust temperature T_d . In reality, thermal dust emission is probably not optically thin, hence the estimated mass is considered to be the lower limit. Assuming a dust mass opacity (κ_ν) of $0.015 \text{ cm}^2 \text{ g}^{-1}$ (taking a dust-to-gas ratio of 100) appropriate for these wavelengths (1.3 mm) ([Ossenkopf et al. 1994](#)), a typical opacity power-law index $\beta = 1.2$ (obtained from our ALMA data), as well as a characteristic dust temperature of 50 K, we estimated a lower limit for the mass of the most compact part (*i.e.*, not interferometrically filtered) of the disk and envelope system associated with HH 901 of about $0.1 M_\odot$. The mass uncertainty could be very large (a factor of up to 3 or 4) given the uncertainty in the opacity and in the dust

temperature.

We can also estimate the volumetric and column density following the procedure described in [Hernández-Hernández et al. \(2014\)](#). We obtained a lower limit for the column density of 10^{24} cm^{-2} and a volumetric density of 10^8 cm^{-3} for the dusty source associated with the HH 901 object. These values are observed in very dense regions as compact envelopes and circumstellar disks toward the nearest GMC, Orion ([Teixeira et al. 2016](#); [Takahashi et al. 2013](#)).

If we assume that the continuum emission at these millimeter wavelengths is mostly arising from the disk, with a very small contribution of the envelope, the $0.1 M_\odot$ could be linked to the disk. If we further adopt a value $M_{\text{star}}/M_{\text{disk}}$ between 1 and 10 ([Bate 2018](#)), we can estimate that the protostar in the middle of the HH 901 object should be a low-mass star. Furthermore, if the envelope is even more massive and the disk mass is smaller, we will have a protostar of lower mass. However, given the uncertainties on the mass estimation for the disks (a factor of 3 or 4), it could also be possible that the star in the middle is an intermediate-mass star. This is in agreement with recent observations ([Reiter & Smith 2013, 2014](#)).

In Figures 2 and 3 show the kinematics and distribu-




Figure 4. ALMA CO(2–1) moment zero (blue and red colors) and millimeter continuum (contours) images of the HH 902 object overlaid in a *HST* optical image (green colors). The blue and red color represent the blue-shifted and red-shifted emission, respectively, from the HH 902 outflow. The contours range from 50% to 80% of the peak emission, in steps of 10%. The peak of the millimeter continuum emission is $5.1 \text{ mJy Beam}^{-1}$. Here, we are only contouring the most compact 1.3 mm emission from our observations revealing the envelope and the disk. There is still some extended emission at low levels. The half-power contour of the synthesized beam of the continuum image is shown in the bottom-left corner. We integrate in radial velocities from -3.7 to -5.6 km s^{-1} for the redshifted emission, and from -10.8 to -15.2 km s^{-1} for the blueshifted emission for the CO outflow. The LSR systemic velocity of the entire pillar associated with the object HH 902 is about -8.5 km s^{-1} .

tion of the molecular gas using the C^{18}O , N_2D^+ , and ^{12}CO spectral lines. These Figures present the moment zero emission from ^{12}CO together with the moment one of the C^{18}O , N_2D^+ , and ^{12}CO lines. In order to compute these moments we integrate in radial velocities from -14.6 to $+6.4 \text{ km s}^{-1}$, which includes the emission from the pillars and the outflows. The molecule that better traces the pillars and the outflows is the ^{12}CO , see Figure 1. From this Figure, one can see how the pillar is very well defined and with a clear east-west velocity gradient going from the blueshifted to redshifted, respectively. In the southern most side of the pillar, the bipolar outflow HH 901 is evident. The bipolar outflow is only observed in the area traced by the pillar probably because the critical density for the CO(2–1) ($\sim 10^4 \text{ cm}^{-3}$) decreases drastically outside of the pillar and the outflow emission becomes too faint at larger spatial scales to be detected. Both lines, the C^{18}O and N_2D^+ confirmed the east-west velocity gradient clearly revealed by the CO, as seen in Figure 3. However, for the case of N_2D^+ , this molecule is only observed in a small solid angle coincident with the 1.3 mm continuum image, probably

tracing the densest parts of the pillar where the star formation is taking place. The critical density of N_2D^+ $\text{J}=3-2$ is about $3 \times 10^6 \text{ cm}^{-3}$, which explains the fact that the N_2D^+ line has better spatial correlation with the 1.3 mm dust continuum emission. Matsushita et al. (2019) reported that the N_2D^+ line is strongly depleted towards the dusty massive core MM5/OMC-3 and this is explained in terms of chemical evolution. Once CO evaporates in the gas phase, molecules like N_2H^+ or N_2D^+ will not have an efficient formation process. This is, however, not observed in our maps probably because it is needed a better angular resolution or sensitivity.

As the CO(2–1) is probably optically thick, we can use the C^{18}O to estimate the opacity (τ_0) and then to estimate the mass of the pillar and the outflow. Assuming Local Thermodynamic Equilibrium, we can estimate the masses using the equations presented in Scoville et al. (1986); Palau et al. (2007) for the $\text{J}=2-1$ transition, obtaining:

$$\left[\frac{M_{\text{H}_2}}{M_{\odot}} \right] = 1.2 \times 10^{-15} T_{\text{ex}} e^{\frac{16.59}{T_{\text{ex}}}} \times \frac{H_2}{\text{CO}} \left[\frac{\int I_{\nu} dv}{\text{Jy km s}^{-1}} \right] \left[\frac{\theta_{\text{maj}} \theta_{\text{min}}}{\text{arcsec}^2} \right] \left[\frac{D}{\text{pc}} \right]^2. \quad (2)$$




Figure 5. ALMA CO(2–1) moment zero (contours) and one (colors) maps of the pillar and outflow HH 902. The contours range from 10% to 90% of the peak emission, in steps of 10%. The peak of the millimeter line CO emission is $1.6 \text{ Jy Beam}^{-1} \text{ km s}^{-1}$. The half-power contour of the synthesized beam of the line image is shown in the bottom-left corner. The LSR radial velocity scale-bar is shown at the right. The brown circle marks the position of the disk revealed in these ALMA observations for the HH 902 object. For the CO moment zero image, we integrate in radial velocities from -15.2 to $+3.7 \text{ km s}^{-1}$. The LSR systemic velocity of the entire pillar associated with the object HH 902 is about -8.5 km s^{-1} .




Figure 6. ALMA $\text{C}^{18}\text{O}(2-1)$ (left panel) and $\text{N}_2\text{D}^+(3-2)$ (right panel) moment one maps overlaid with a contour image of the CO(2–1) moment zero of the pillar and outflow HH 902. The contours range from 10% to 90% of the peak emission, in steps of 10%. The peak of the millimeter line CO emission is $1.8 \text{ Jy Beam}^{-1} \text{ km s}^{-1}$. The half-power contour of the synthesized beam of the line image is shown in the bottom-left corner of each panel. The LSR radial velocity scale-bar is shown at the right. The brown circle marks the position of the disk+envelope revealed in these ALMA observations for the HH 902 object.

We used 2.8 as mean molecular weight, an abundance ratio between the molecular hydrogen and the carbon monoxide ($X_{\frac{H_2}{CO}}$) of $\sim 10^4$ (Scoville et al. 1986), an excitation temperature (T_{ex}) of 50 K, an average intensity (I_ν) of 0.01 Jy Beam $^{-1}$ for the outflow and 0.06 Jy Beam $^{-1}$ for the pillar, we take a velocity range (dv) of 5.5 km s $^{-1}$ for the outflow and 2.5 km s $^{-1}$ for the pillar, and a distance (D) to the source of 2.3 ± 0.1 kpc. We then estimate a gas mass for the outflow of $\sim 5 \times 10^{-4} M_\odot$ and for the pillar of $\sim 0.1 M_\odot$. For the mass of the outflow, we can correct this value for the inclination, therefore the true velocity range is $dv_{\text{real}} = \frac{dv}{\sin \theta}$, where θ is the inclination angle taken to be 20° (Reiter & Smith 2013, 2014), so the $dv_{\text{real}} = 16$ km s $^{-1}$ and the mass is $10^{-3} M_\odot$.

The uncertainty in the values of the mass arises mainly from the error in the distance, which is only 4%, the flux density which is about 10% and the excitation temperature. Given that the C 18 O intensity line is approximately five times weaker than the 12 CO line, we estimate an average opacity (τ_0) of 0.2, so we corrected the mass value by a factor of $\left[\frac{\tau_0}{1 - \exp(-\tau_0)} \right]$. Furthermore, this mass estimation is in agreement with the mass obtained for the HH 901 pillar using a volumetric density of 10^5 cm $^{-3}$ and a physical volume of $10'' \times 2'' \times 2''$ obtained in Reiter & Smith (2013) and with a value of about $0.1 M_\odot$.

For the outflow, the resulting kinetic energy ($E_k = \frac{1}{2}mv^2$) is about 10^{42} ergs and the momentum ($p=mv$) is $1 \times 10^{-2} M_\odot$ km s $^{-1}$. These values are slightly higher than those estimated by Lumbreras & Zapata (2014); Zapata et al. (2015, 2018) for outflows associated with low-mass protostars, but possibly consistent if the correction for outflow inclination is taken into account in these studies. If we estimate the mechanical force for the outflow F_{CO} , we obtain a value of $1.5 \times 10^{-5} M_\odot$ yr $^{-1}$ km s $^{-1}$. This value is low compared from those obtained for intermediate-mass young stars reported in Table 7 of van Kempen et al. (2016), again suggesting that this outflow is associated with a low-mass or in the limit of an intermediate star. If we add the momentum and energy from the optical counterpart could be an intermediate-mass star. Here, it is important to mention that the outflows in Carina are strongly irradiated, and maybe most of the energy budget is already in the optical irradiated counterparts, thus that a direct comparison with van Kempen's outflows could be not so straight.

As already mentioned in Section 1, Reiter & Smith (2013, 2014) suggested that the HH 901 jet is energized by a relatively young intermediate-mass star (2-8 M_\odot). However, this estimation of the mass (and evolutionary stage) is very poorly constrained because they have not

direct detection of the IR flux from the driving source of HH 901 (Reiter & Smith 2014). This assumption is only inferred by the large mass-loss rates of a few $10^{-6} M_\odot$ yr $^{-1}$ (these are about of two orders of magnitude larger than those observed in T-Tauri stars) obtained from the optical ionized lines. These large mass-loss rates can be later transformed to mass-accretion rates and then to luminosity of the exciting star. However, the possibility that the powering source of the HH 901 is of intermediate-mass is in agreement with our ALMA data, but this intermediate-mass star must be on the low range of mass as suggested by the outflow energy.

Given the high the mass-lost rate reported in the HH 901 object ($10^{-6} M_\odot$ yr $^{-1}$), the protostar in the middle could be Class 0/I object, so this should be very obscured even in the IR wavelengths. This is in agreement with the fact that the protostar is still surrounded by an envelope traced by the 1.3 mm and N $_2$ D $^+$ emission.

Following Mesa-Delgado et al. (2016), we can estimate the photo-evaporation time-scale for the pillar or globulette associated with the HH 901 object. Taken the Equation 2 (already discussed in that paper,) and which allow us to estimate the theoretical value of the globule/disk mass photo-evaporation rate \dot{M} by Extreme Ultraviolet radiation (EUV):

$$\dot{M}_{\text{EUV}} \simeq 4 \times 10^{-9} \left(\frac{F_{\text{EUV}}}{10^{10} \text{ s}^{-1} \text{ cm}^2} \right)^{1/2} \left(\frac{R}{100 \text{ AU}} \right)^{3/2} M_\odot \text{ year}^{-1}, \quad (3)$$

where F_{EUV} is EUV flux, and R is the radius of the globule or disk. As the massive cluster Tr 14 is the closest (1.0 pc) to the HH 901 object, we ignored at this point the radiation for the massive cluster located also in this region Tr 16. The Tr 14 cluster has an UV luminosity (Q_H) of 2×10^{50} photon s $^{-1}$ (Smith & Brooks 2008), implying $F_{\text{EUV}} = 10^{12}$ photon s $^{-1}$ cm $^{-2}$ at the distance of the HH 901 object. For a radius of 1000 au, and using the Equation 3, we obtain a mass photo-evaporation rate \dot{M} equal to $1 \times 10^{-6} M_\odot$ year $^{-1}$. Therefore, taking a mass of $0.1 M_\odot$ for the pillar HH 901, we obtain that the pillar will be photo-evaporated in around 10^5 years. This timescale is shorter to that estimated, for example, for the massive EGG or globulette 105-600 (Mesa-Delgado et al. 2016) of about 10^6 years. However, the 105-600 globulette is farther away from the Tr 14 massive cluster (17 pc), and is more massive (0.1 to 0.3 M_\odot), which explains the large difference.

3.2. HH 902

In Figure 4 we present the *Hubble Space Telescope*/Advanced Camera for Surveys image from Smith et al. (2010) of the optical jet HH 902 overlaid with the ALMA high velocity CO(2-1) and the 1.3 mm continuum emission. As in the HH 901 object (Figure 1), the CO emission also reveals a bipolar and collimated out-

flow that is tracing the most inner part of the optical jet in a north-west and south-east orientation. This map reveals that the blue-shifted emission is located toward an eastern orientation, while red-shifted emission is to its western side, similar to the HH 901 object. We integrate in radial velocities from $+3.7$ to -5.6 km s $^{-1}$ for the redshifted emission, and from -10.8 to -15.2 km s $^{-1}$ for the blueshifted emission for the CO outflow in the HH 902 object. The CO emission between -5.5 km s $^{-1}$ to -10.7 km s $^{-1}$ is associated with the dusty pillar as is shown in Figure 2. The LSR systemic velocity of the entire pillar associated with the object HH 902 is about -8.5 km s $^{-1}$. The radial velocities of the innermost CO outflow are in agreement to those presented in Reiter & Smith (2014).

The 1.3 mm continuum emission in Figure 4 is also tracing, as in Figure 1, the disk+envelope associated with the HH 902 object. However, in this Figure two objects, named A and B, are revealed. The component A is associated with HH 902 object, while the object B is to its north-west. The source B is not associated with any molecular outflow, maybe because it is still in its pre-stellar phase. This source is extended with a size of about 2000 au and has a flux density of 38 ± 2.0 mJy Beam $^{-1}$. Assuming a dust temperature of 30 K, this flux density corresponds to a mass of $1.5 M_{\odot}$, which is likely associated with an extended envelope. For the millimeter component A, from a gaussian fitting to the continuum source from a map obtained using ROBUST equals to 0.5 (with a better angular resolution) we obtained a deconvolved size of $0.34'' \pm 0.02'' \times 0.28'' \pm 0.02''$ with a PA of $107^{\circ} \pm 13^{\circ}$ and an integrated flux of 6.5 ± 0.7 mJy Beam $^{-1}$. The corresponding physical sizes of these deconvolved values are about 700 au, which again suggests that we are also seeing part of the envelope and disk.

Following the Equation 1 and taking similar values to the ones assumed for the object HH 901, we estimated a lower limit (because again the thermal dust emission is probably not optically thin) for the mass of the most compact part of the disk and envelope system (of source A) associated with the HH 902 of about $0.1 M_{\odot}$. In a similar way as for the HH 901 continuum object, this mass for the dust for the HH 902 object corresponds to a low-mass protostar.

In Figures 5 and 6, the kinematics and distribution of the molecular gas using the C 18 O, N $_2$ D $^+$, and 12 CO spectral lines are presented. These Figures present the moment zero emission from 12 CO together with the moment one of the C 18 O, N $_2$ D $^+$, and 12 CO lines. Similar to Figures 3 and 4, in order to compute these moments we integrate in radial velocities from -15.2 to $+3.7$ km s $^{-1}$, which includes the emission from the pillars and the outflow. The molecule that better traces the pillars and

the outflow is again the 12 CO, as seen in Figure 5, but now for the HH 902 object. From this Figure, one can see how the pillar is very well defined and with a clear east-west velocity gradient going from the blueshifted to redshifted, respectively. In the most southern side of the pillar, the bipolar outflow HH 902 is evident. Both lines, the C 18 O and N $_2$ D $^+$ confirm the east-west velocity gradient clearly revealed by the CO, as seen in Figure 5. However, for the case of N $_2$ D $^+$, this molecule again is only observed in small solid angle coincident with the 1.3 mm continuum image, probably tracing the densest parts of the pillar, as in the HH 901 object.

Using the Equation 2 and assuming similar values for the excitation temperature, an abundance ratio, and distance, we can obtain the mass for the outflow and the pillar associated with the HH 902 object. We take an average intensity (I_{ν}) of 0.015 Jy Beam $^{-1}$ for the outflow and 0.07 Jy Beam $^{-1}$ for the pillar, and we also take a velocity range (dv) of 5.0 km s $^{-1}$ for the outflow and 3.0 km s $^{-1}$ for the pillar. For these values, we estimate a gas mass for the outflow of $7 \times 10^{-4} M_{\odot}$ and for the pillar of $\sim 0.2 M_{\odot}$. We also corrected these mass values by the opacity. Moreover, correcting these values by the inclination, we obtain a mass for the outflow of $10^{-3} M_{\odot}$.

Taking again the Equation 3 and similar values for the EUV radiation ($F_{\text{EUV}} = 10^{12}$ photon s $^{-1}$ cm $^{-2}$) at the distance of the HH 902 object and a radius of 4×10^3 au, we obtain a mass photo-evaporation rate \dot{M} equal to $1 \times 10^{-5} M_{\odot}$ year $^{-1}$. Therefore, taking a mass of $0.2 M_{\odot}$ for the pillar HH 902, we obtain that the pillar will be photo-evaporated in around 2×10^4 years, which is a lower value to the timescale of the HH 901.

Given the short photo-evaporated timescales of the molecular pillars and that the protostars in these pillars are still very embebed, we suggest that that the disks inside of the pillars will be quickly affected by the radiation of the massive stars (in about $10^4 - 10^5$ years), forming proplyds like those observed in Orion (O'Dell et al. 1993, 2008; Henney & O'Dell 1999).

4. CONCLUSIONS

In this study, we present sensitive and high angular resolution observations of the famous HH 901 and HH 902 objects located in the Carina Nebula using ALMA. The main results of our work can be summarized as follows.

- Our observations reveal for the first time, the bipolar CO collimated outflows, and the dusty disks (plus envelopes) that are energizing the extended and irradiated HH 901/902 objects far from the pillars.
- We find that the masses of the disks+envelopes

are about $0.1 M_{\odot}$ and that of the bipolar outflows are between $10^{-3} - 10^{-4} M_{\odot}$, which suggests that the outflows could be energized by low- or maybe intermediate- mass protostars.

- We also reveal the kinematics of the gas in the pillars together with their respective gas masses ($0.1 - 0.2 M_{\odot}$).
- We estimate that the pillars will be photo-evaporated in around of 10^4 to 10^5 years by the massive and luminous stars located in the Tr 14 cluster.

We are very thankful for the thoughtful suggestions of the anonymous referee that helped to improve our

manuscript. This paper makes use of the following ALMA data: ADS/JAO.ALMA#2017.1.00912.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), MOST and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. L.A.Z. and J. A. T. are grateful to CONACyT, México, and DGAPA, UNAM for the financial support.

Facilities: HST(STIS), ALMA

Software: CASA(McMullin et al. 2007), KARMA(Gooch 1996)

REFERENCES

- Bally, J. 2016, ARA&A, 54, 491
 Bate, M. R. 2018, MNRAS, 475, 5618
 Feigelson, E. D., Getman, K. V., Townsley, L. K., et al. 2011, ApJS, 194, 9
 Gooch, R. 1996, Astronomical Data Analysis Software and Systems V, 101, 80
 Haikala, L. K., Gahm, G. F., Grenman, T., et al. 2017, A&A, 602, A61
 Hartigan, P., Reiter, M., Smith, N., et al. 2015, AJ, 149, 101
 Hartigan, P., Palmer, J., & Cleaves, L. I. 2012, High Energy Density Physics, 8, 313
 Henney, W. J., & O'Dell, C. R. 1999, AJ, 118, 2350
 Hernández-Hernández, V., Zapata, L., Kurtz, S., & Garay, G. 2014, ApJ, 786, 38
 Lumbreras, A. M., & Zapata, L. A. 2014, AJ, 147, 72
 McMullin, J. P., Waters, B., Schiebel, D., Young, W., & Golap, K. 2007, Astronomical Data Analysis Software and Systems XVI, 376, 127
 Matsushita, Y., Takahashi, S., Machida, M. N., et al. 2019, ApJ, 871, 221
 Mesa-Delgado, A., Zapata, L., Henney, W. J., Puzia, T. H., & Tsamis, Y. G. 2016, ApJL, 825, L16
 Massey, P., & Johnson, J. 1993, AJ, 105, 980
 O'Dell, C. R., Wen, Z., & Hu, X. 1993, ApJ, 410, 696
 O'Dell, C. R., Muench, A., Smith, N., & Zapata, L. 2008, Handbook of Star Forming Regions, Volume I, 4, 544
 Ossenkopf, V., & Henning, T. 1994, A&A, 291, 943
 Palau, A., Estalella, R., Ho, P. T. P., Beuther, H., & Beltrán, M. T. 2007, A&A, 474, 911
 Povich, M. S., Smith, N., Majewski, S. R., et al. 2011, ApJS, 194, 14
 Preibisch, T., Hodgkin, S., Irwin, M., et al. 2011, ApJS, 194, 10
 Takahashi, S., Ho, P. T. P., Teixeira, P. S., et al. 2013, ApJ, 763, 57
 Teixeira, P. S., Takahashi, S., Zapata, L. A., & Ho, P. T. P. 2016, A&A, 587, A47
 Sahai, R., Güsten, R., & Morris, M. R. 2012, ApJL, 761, L21
 Scoville, N. Z., Sargent, A. I., Sanders, D. B., et al. 1986, ApJ, 303, 416
 Smith, N. 2006, MNRAS, 367, 763
 Smith, N., & Brooks, K. J. 2008, Handbook of Star Forming Regions, Volume II, 5, 138
 Smith, N., Bally, J., & Walborn, N. R. 2010, MNRAS, 405, 1153
 Reiter, M., & Smith, N. 2013, MNRAS, 433, 2226
 Reiter, M., & Smith, N. 2014, MNRAS, 445, 3939
 van Kempen, T. A., Hogerheijde, M. R., van Dishoeck, E. F., et al. 2016, A&A, 587, A17
 Walborn, N. R. 1973, ApJ, 179, 517
 Walborn, N. R. 1995, Revista Mexicana De Astronomia Y Astrofisica Conference Series, 51
 Zapata, L. A., Fernández-López, M., Rodríguez, L. F., et al. 2018, AJ, 156, 239
 Zapata, L. A., Lizano, S., Rodríguez, L. F., et al. 2015, ApJ, 798, 131