

A note on factorial \mathbb{A}^1 -forms with retractions

Prosenjit Das

*Stat-Math Unit, Indian Statistical Institute,
203, B.T. Road, Kolkata 700 108, India.*

Abstract

Let k be a field. In this paper we will show that any factorial \mathbb{A}^1 -form A over any k -algebra R is trivial if A has a retraction to R .

Keywords: \mathbb{A}^1 -form, Factorial domain, Retraction

AMS Subject classifications (2000). Primary 13B25; Secondary 12F05, 13B10

Throughout this paper, by “ring”, we shall mean “commutative ring with unity”.

Definitions. Let k be a field, R a k -algebra and A an R -algebra. A is said to be an \mathbb{A}^1 -form over R (with respect to k) if there exists an algebraic field extension $k'|_k$ such that $A \otimes_k k' \cong (R \otimes_k k')[X]$. An \mathbb{A}^1 -form is called *purely inseparable* (resp. *separable*), if we can take the extension $k'|_k$ to be a purely inseparable (resp. a separable) extension.

A ring homomorphism $\Phi : A \rightarrow R$ is said to be a *retraction* if Φ is R -linear. If a retraction $\Phi : A \rightarrow R$ exists, we say R is a *retract* of A .

It is well known that any separable \mathbb{A}^1 -form over any field is trivial. More generally, it has been shown that a separable \mathbb{A}^1 -form over an arbitrary commutative algebra is trivial ([Dut00], Theorem 7), i.e.,

Theorem 1. *Let k be a field, L a separable field extension of k , R a k -algebra and A an R -algebra such that $A \otimes_k L \cong \text{Sym}_{(R \otimes_k L)}(P')$ for a finitely generated rank one projective module P' over $R \otimes_k L$. Then $A \cong \text{Sym}_R(P)$ for a finitely generated rank one projective module P over R .*

If k is not perfect, there exist non-trivial purely inseparable \mathbb{A}^1 -forms. Asanuma gave a complete structure theorem for purely inseparable \mathbb{A}^1 -forms over a field k of characteristic $p > 2$ ([Asa05], Theorem 8.1). However, from Asanuma’s results, it can be deduced that any factorial \mathbb{A}^1 -form over a field k with a k -rational point is trivial, i.e.,

Email address: prosenjit.das@gmail.com (Prosenjit Das)

Theorem 2. *Let k be a field and A an \mathbb{A}^1 -form over k such that*

- (1) *A is a UFD.*
- (2) *A has a k -rational point.*

Then $A \cong k[X]$.

In this paper we shall show that Theorem 2 has a generalization, in the spirit of Theorem 1, to \mathbb{A}^1 -forms over k -UFDs (see Corollary 4 for precise formulation). The generalization turns out to be a special case of our main theorem (Theorem 3) which may be envisaged as an example of “faithfully flat descent”.

Theorem 3. *Let R be a ring and A be an R -algebra such that*

- (1) *A is a UFD.*
- (2) *There is a retraction $\Phi : A \rightarrow R$.*
- (3) *There exists a faithfully flat ring homomorphism $\eta : R \rightarrow R'$ such that $A \otimes_R R' \cong R'[X]$.*

Then $A \cong R[X]$.

Proof. Let $A' = A \otimes_R R' = R'[f]$, $P = \text{Ker } \Phi$, $P' = PA' (= P \otimes_R R')$, and let $\Phi' = \Phi \otimes 1$ be the induced retraction from A' to R' . Then P is a prime ideal of A and we have a short exact sequence of R -modules

$$0 \longrightarrow P \longrightarrow A \xrightarrow{\Phi} R \longrightarrow 0$$

and hence a short exact sequence of R' -modules

$$0 \longrightarrow P' \longrightarrow A' (= R'[f]) \xrightarrow{\Phi'} R' \longrightarrow 0.$$

Then $P' = \text{Ker } \Phi' = (f - \Phi'(f))R'[f]$. Replacing $f - \Phi'(f)$ by f , we assume $P' = fA' = fR'[f]$. Since A' is faithfully flat over A , going-down theorem holds between A and A' ([Mat89], Pg. 68, Theorem 9.5) and also $P' \cap A = P$ ([Mat89], Pg. 49, Theorem 7.5). As $ht(P') = 1$, it follows that $ht(P) = 1$.

Now, since A is a UFD, there exists $g \in A$ such that $P = gA$. Thus we get $gA' = P' = fA'$. Since f is a non-zero divisor in $A' (= R'[f])$, it follows that $g = \lambda f$ for some unit λ in A' . Let

$$\lambda = a_0 + a_1f + a_2f^2 + \cdots + a_nf^n$$

where a_0 is a unit in R' and a_i is nilpotent in R' for $1 \leq i \leq n$. Let $I = (a_1, a_2, \cdots, a_n)R'$. Then I is a nilpotent ideal of R' . Let N be the least positive integer such that $I^N = (0)$. Since $g \equiv a_0f \pmod{I}$, we have

$$R'[f] = R'[g] + IR'[f] = \cdots = R'[g] + I^N R'[f] = R'[g].$$

Thus we have $R[g] \subseteq A$ and $R[g] \otimes_R R' = R'[g] = A \otimes_R R'$. Since R' is faithfully flat over R , it follows that $A = R[g] \cong R[X]$. \square

As an immediate consequence of Theorem 3, we get the following corollary:

Corollary 4. *Let k be a field, R a k -algebra and A an R -algebra such that*

- (1) *A is a UFD.*
- (2) *R is a retract of A .*
- (3) *A is an \mathbb{A}^1 -form over R .*

Then $A \cong R[X]$.

The following two well-known examples ([KMT74], Pg. 70–71, Remark 6.6(a), Examples (i) and (ii)) respectively show that in Theorem 2 (and hence in Theorem 3), the hypothesis on the existence of a retraction and the hypothesis “ A is a UFD” are necessary.

Example 1. Let \mathbb{F}_p be the prime field of characteristic p and let $k = \mathbb{F}_p(t, u)$ be a purely transcendental extension of \mathbb{F}_p with variables t and u . Then $A = k[X, Y]/(Y^p - t - X - uX^p)$ is a factorial non-trivial \mathbb{A}^1 -form over k which does not have a retraction to k .

Example 2. Let k be a field of characteristic $p \geq 2$ and $A = k[X, Y]/(Y^p - X - aX^p)$ where $a \in k \setminus k^p$. Then A is a non-trivial \mathbb{A}^1 -form over k with a retraction to k . Here A is not a UFD.

Acknowledgement:

In the earlier version of the article, Corollary 4 was presented as the main theorem. I thank the referee for his observation that the arguments give the more general statement Theorem 3.

I thank Amartya K. Dutta for suggesting the problem and for his useful comments and suggestions. I am grateful to T. Asanuma for his lectures on purely inseparable \mathbb{A}^1 -forms in Indian Statistical Institute, Kolkata. I also thank S. M. Bhatwadekar and N. Onoda for helpful discussions.

References

- [Asa05] Teruo Asanuma, *Purely inseparable k -forms of affine algebraic curves*, Affine algebraic geometry, Contemp. Math., vol. 369, Amer. Math. Soc., Providence, RI, 2005, pp. 31–46.
- [Dut00] Amartya K. Dutta, *On separable \mathbb{A}^1 -forms*, Nagoya Math. J. **159** (2000), 45–51.

- [KMT74] Tatsuji Kambayashi, Masayoshi Miyanishi, and Mitsuhiro Takeuchi, *Unipotent algebraic groups*, Lecture Notes in Mathematics, Vol. 414, Springer-Verlag, Berlin, 1974.
- [Mat89] Hideyuki Matsumura, *Commutative ring theory*, second ed., Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, Cambridge, 1989, Translated from the Japanese by M. Reid.