

AN ALMA SURVEY OF H₂CO IN PROTOPLANETARY DISKS

JAMILA PEGUES¹, KARIN I. ÖBERG¹, JENNIFER B. BERGNER^{2,1}, RYAN A. LOOMIS^{3,1}, CHUNHUA QI¹, ROMANE LE GAL¹, L. ILSEDORE CLEEVES⁴, VIVIANA V. GUZMÁN^{5,6}, JANE HUANG¹, JES K. JØRGENSEN⁷, SEAN M. ANDREWS¹, GEOFFREY A. BLAKE⁸, JOHN M. CARPENTER⁵, KAMBER R. SCHWARZ⁹, JONATHAN P. WILLIAMS¹⁰, DAVID J. WILNER¹

¹Center for Astrophysics | Harvard & Smithsonian, Cambridge, MA 02138, USA

²Department of Geophysical Sciences, University of Chicago, Chicago, IL 60637, USA

³National Radio Astronomy Observatory, Charlottesville, VA 22903, USA

⁴Department of Astronomy, University of Virginia, Charlottesville, VA 22904, USA

⁵Joint ALMA Observatory, Alonso de Córdova 3107 Vitacura, Santiago, Chile

⁶Instituto de Astrofísica, Pontificia Universidad Católica de Chile, Av. Vicuña Mackenna 4860, 7820436 Macul, Santiago, Chile

⁷Niels Bohr Institute & Centre for Star and Planet Formation, University of Copenhagen, Øster Voldgade 5-7, DK-1350 Copenhagen K., Denmark

⁸Division of Geological & Planetary Sciences, California Institute of Technology, Pasadena, CA 91125, USA

⁹Sagan Fellow at the Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ 85721, USA and

¹⁰Institute for Astronomy, University of Hawai'i at Mānoa, Honolulu, HI 96822, USA

Draft version March 2, 2020

ABSTRACT

H₂CO is one of the most abundant organic molecules in protoplanetary disks and can serve as a precursor to more complex organic chemistry. We present an ALMA survey of H₂CO towards 15 disks covering a range of stellar spectral types, stellar ages, and dust continuum morphologies. H₂CO is detected towards 13 disks and tentatively detected towards a 14th. We find both centrally-peaked and centrally-depressed emission morphologies, and half of the disks show ring-like structures at or beyond expected CO snowline locations. Together these morphologies suggest that H₂CO in disks is commonly produced through both gas-phase and CO-ice-regulated grain-surface chemistry. We extract disk-averaged and azimuthally-averaged H₂CO excitation temperatures and column densities for four disks with multiple H₂CO line detections. The temperatures are between 20-50K, with the exception of colder temperatures in the DM Tau disk. These temperatures suggest that H₂CO emission in disks is generally emerging from the warm molecular layer, with some contributions from the colder midplane. Applying the same H₂CO excitation temperatures to all disks in the survey, we find that H₂CO column densities span almost three orders of magnitude ($\sim 5 \times 10^{11} - 5 \times 10^{14} \text{ cm}^{-2}$). The column densities appear uncorrelated with disk size and stellar age, but Herbig Ae disks may have less H₂CO compared to T Tauri disks, possibly because of less CO freeze-out. More H₂CO observations towards Herbig Ae disks are needed to confirm this tentative trend, and to better constrain under which disk conditions H₂CO and other oxygen-bearing organics efficiently form during planet formation.

Subject headings: astrochemistry, protoplanetary disks, ISM: molecules, radio lines: ISM

1. INTRODUCTION

Protoplanetary disks are the precursors of planetary systems. These disks consist of gas and dust orbiting young stars. Over time, the dust grains collide and stick together to form pebbles, eventually growing into planetesimals and planets (e.g. Mordasini et al. 2008, and references therein). The organic compositions of pebbles, planetesimals, and planets are therefore shaped by the organic chemistry of the gas and dust in their ancestral disks. By studying this chemistry, we can ultimately model and predict the chemical formation environment of young planets (e.g. Cridland et al. 2016; Öberg & Bergin 2016; Cridland et al. 2017) and their potential ability to develop Earth-like life.

The small organic molecule formaldehyde (H₂CO) is expected to be important for the overall organic chemical budget of disks. H₂CO is commonly and abundantly detected in disks (e.g., Dutrey et al. 1997; Aikawa et al. 2003; Thi et al. 2004; Öberg et al. 2010, 2011). Based on cometary observations, H₂CO was also abundant in the solar nebula; in comets it is present at levels of ~ 0.1 -1% with respect to water (Mumma & Charnley 2011). There are several possible chemical origins

of H₂CO in disks. H₂CO can form through neutral-neutral gas-phase chemistry (e.g., Fockenberg & Preses 2002; Atkinson et al. 2006), and through grain-surface chemistry via CO ice hydrogenation (e.g., Hiraoka et al. 1994; Hiraoka et al. 2002; Watanabe & Kouchi 2002; Hidakata et al. 2004; Watanabe et al. 2004; Fuchs et al. 2009). Both processes can occur in disks, though likely in different locations, since neutral-neutral gas-phase chemical reactions increase with density and temperature towards the inner disk, while CO ice hydrogenation is only possible beyond the CO snowline. In addition, H₂CO has been detected in ices around young stellar objects (e.g., detections/tentative detections by Gibb et al. 2004; Pontoppidan et al. 2004; Boogert et al. 2008), and some of the H₂CO observed in disks might be inherited from this preceding evolutionary stage (e.g., Drozdovskaya et al. 2014, 2019).

Understanding H₂CO emission and its origins in disks is important for modeling more complex organic disk chemistry. When formed through CO-ice hydrogenation, H₂CO serves as a stepping-stone towards forming oxygen-bearing, complex organic molecules (known as COMs, and typically defined as unsaturated carbon-

bearing molecules with five or more atoms; e.g., Herbst & van Dishoeck 2009), especially methanol (CH_3OH ; e.g., Hiraoka et al. 1994; Watanabe & Kouchi 2002). COMs are expected to be rich in disks based on astrochemical disk models (Walsh et al. 2014). Unfortunately, without the help of rare heating phenomena such as stellar outbursts (e.g., van ’t Hoff et al. 2018; Lee et al. 2019), COM emission lines are predicted to be difficult to detect in disks (Henning & Semenov 2013; Walsh et al. 2014), due to their large partition functions and comparatively small abundances. To date, the largest COMs detected towards disks are cyanoacetylene (HC_3N , Chapillon et al. 2012), methyl-cyanide (CH_3CN , Öberg et al. 2015b), methanol (CH_3OH , Walsh et al. 2016), and formic acid (HCOOH , Favre et al. 2018), and only cyanoacetylene and methyl-cyanide have been detected in more than one disk (Bergner et al. 2018; Loomis et al. 2018). H_2CO , on the other hand, is readily detectable in disks. The presence/absence of H_2CO emission, originating specifically from grain-surface chemistry, can thus indicate whether the disk hosts a rich organic grain-surface chemistry or not (Walsh et al. 2014; Öberg et al. 2017).

The chemical origins of H_2CO in disks, and its potential to form more complex organic chemistry, can be addressed through spatially-resolved observations of H_2CO emission at millimeter and submillimeter wavelengths. To date, H_2CO has been spatially resolved with the Atacama Large Millimeter/submillimeter Array (ALMA) towards six disks: (1) the T Tauri disk Oph IRS 48 (van der Marel et al. 2014); (2) the T Tauri disk DM Tau (Loomis et al. 2015); (3) the Herbig Ae disk HD 163296 (Carney et al. 2017; Guzmán et al. 2018); (4) the T Tauri disk TW Hya (Öberg et al. 2017); (5) the T Tauri disk V4046 Sgr (Kastner et al. 2018); and (6) the T Tauri disk DG Tau (Podio et al. 2019). These six studies show a variety of H_2CO morphologies, from centrally peaked (indicative of gas-phase formation) to ring-like in the outer disk (indicative of formation through CO ice hydrogenation). Two of these studies (Carney et al. 2017; Guzmán et al. 2018) calculated H_2CO excitation temperatures of $>20\text{K}$ and 24K , which are consistent with both formation pathways. Three of these studies (Loomis et al. 2015; Carney et al. 2017; Öberg et al. 2017) determined that the detected H_2CO emission could be described by two distinct components of H_2CO : (1) a warm inner component produced through gas-phase chemistry, and (2) a cold outer component exterior to the CO snowline produced through grain-surface chemistry. A larger survey of H_2CO emission towards disks is necessary to explore the typical distribution and chemical origins of H_2CO in protoplanetary disks, as well as their connections to stellar and disk characteristics.

In this work, we present an ALMA survey of H_2CO towards 15 protoplanetary disks. We characterize the H_2CO morphologies, excitation temperatures, and column densities, evaluate their dependence on stellar and disk characteristics, and assess the chemical origins of H_2CO . In Section 2, we describe the disk sample, the ALMA observations, and the data reduction process. In Section 3, we present our methodology for visualizing and analyzing the imaged line emission. In Section 4, we report the detections of H_2CO , the H_2CO emission morphologies, excitation temperatures, and column den-




FIG. 1.— **Stellar Effective Temperature versus Stellar Age** for our disk sample (Table 1). Each point represents a disk, and the sizes denote the sizes of the submillimeter/millimeter dust continuum. An empty vs. filled point marks whether or not the disk shows a large inner cavity in the submillimeter/millimeter dust continuum. For disks where an age range rather than a single age is given in the literature, horizontal bars span that range.

sities, and any trends relating the H_2CO to stellar and disk characteristics. In Section 5, we discuss the results in the context of H_2CO formation chemistry and disk environment, and in Section 6 we summarize the key findings from our survey.

2. OBSERVATIONAL DETAILS

2.1. Disk Sample

Figure 1 and Table 1 summarize the stellar and disk characteristics of the 15 disks in our sample. The sample was assembled from three main ALMA observing projects, with H_2CO lines from other ALMA observing projects added in as available. The sample consists of well-studied T Tauri and Herbig Ae disks, and gas, dust, and temperature structures are available for most sources from previous studies. The host stars in the sample range in luminosity, mass, and age from $\sim 0.2\text{--}26 L_{\odot}$, $\sim 0.5\text{--}2.0 M_{\odot}$, and $\lesssim 1\text{--}15\text{--}15\text{ Myrs}$, respectively. All disks in the sample are relatively close by, $\sim 70\text{--}170\text{ pc}$ away. Six disks in the sample are associated with the Taurus-Auriga region, five with the Upper Scorpius region, and one each with the Lupus and Ophiuchus regions.

The disks vary in dust morphology and size, but most are large. Outer millimeter dust radii range from $<75\text{ AU}$ to $\sim 200\text{ AU}$ (where we define the outer radius as the radius where the azimuthally-averaged continuum emission drops below 5% of the peak). Five of the disks are “transition disks”, i.e. have large inner holes in their submillimeter/millimeter dust continuum: DM Tau (Marsh & Mahoney 1992), GM Aur (Jensen & Mathieu 1997), J1604-2130 (Mathews et al. 2012), LkCa 15 (Bergin et al. 2004), and V4046 Sgr (Jensen & Mathieu 1997).

2.2. ALMA Observations

The core of our study contains observations of the H_2CO $3_{03}\text{--}2_{02}$ and $4_{04}\text{--}3_{03}$ lines towards eight disks each

TABLE 1
Stellar and Disk Characteristics of the Sample.

Disk	Stellar Type	R.A. ^[0] [J2000]	Dec. ^[0] [J2000]	Dist [pc]	Age _* [Myr]	L _* [L _⊙]	M _* [M _⊙]	T _{eff} [K]
AS 209 ^[1]	K5	16:49:15.3	-14:22:09.0	121	1.0	1.41	0.83	4266
CI Tau ^[2]	K7	04:33:52.0	+22:50:29.8	159	0.7 (0.4-1.8)	1.20	0.66	4060
DM Tau ^[2]	M1	04:33:48.7	+18:10:9.7	145	4.0 (2.5-7.1)	0.24	0.53	3705
DO Tau ^[2]	M0	04:38:28.6	+26:10:49.1	140	0.4 (0.1-0.9)	1.40	0.45	3850
GM Aur ^[2]	K5.5 ^[3]	04:55:11.0	+30:21:59.0	159	2.6 (1.4-4.3)	1.60	1.30	4730
HD 143006 ^[1]	G7	15:58:36.9	-22:57:15.5	165	4.0	3.80	1.78	5623
HD 163296 ^[1]	A1	17:56:21.3	-21:57:22.5	101	13	17	2.04	9333
IM Lup ^[1]	K5	15:56:09.2	-37:56:06.5	158	0.5	2.57	0.89	4266
J16042165-2130284 ^{*[2]}	K2	16:04:21.6	-21:30:28.9	149	13.8 (7.4-32)	0.62	1.11	4900
J16090075-1908526 ^{*[2]}	K9	16:09:00.7	-19:08:53.1	137	4.8 (2.5-9.1)	0.32	0.68	3890
J16123916-1859284 ^{*[2]}	M0.5	16:12:39.2	-18:59:28.9	138	4.1 (1.8-8.3)	0.29	0.56	3775
J16142029-1906481 ^{*[2]}	M0	16:14:20.3	-19:06:48.5	143	2.1 (1.0-5.4)	0.46	0.60	3850
LkCa 15 ^[2]	K5	04:39:17.8	+22:21:03.1	158	2.0 (0.9-4.3)	1.04	1.03	4350
MWC 480 ^[2]	A4.5±2	04:58:46.3	+29:50:36.6	161	6.5	25	1.84	8330
V4046 Sgr ^{**[4]}	K5,K7 ^[5]	18:14:10.5	-32:47:35.3	73	13	0.86	1.75	4350, 4060

NOTE. — R.A. and Dec. (Columns 3 and 4) are from GAIA (e.g., Gaia Collaboration et al. 2016a,b, 2018). All other table values are taken from the literature. For AS 209, HD 143006, HD 163296, and IM Lup, we use values as updated by [2] using new GAIA distances. For disks denoted by [1], we use values calculated/scaled using the same methodology of [2] unless otherwise individually referenced. For V4046 Sgr, we take all values except for the stellar type from [4], which assumed a distance of 73pc. Stellar ages (Column 6) are given as ranges in parentheses (which account for error) when available. *: The disks J16042165-2130284, J16090075-1908526, J16123916-1859284, and J16142029-1906481 are referred to respectively as J1604-2130, J1609-1908, J1612-1859, and J1614-1906 in subsequent figures, tables, and text. **: V4046 Sgr is a protoplanetary disk orbiting a binary star system. The individual stellar effective temperatures and total stellar luminosity and mass are reported. [0] Gaia Collaboration et al. (2016a,b, 2018), [1] Andrews et al. (2018), [2] Calculated using the same methodology as Andrews et al. (2018), [3] Espaillat et al. (2010), [4] Rosenfeld et al. (2012), [5] Quast et al. (2000).

TABLE 2
Observed H₂CO Lines in the Sample.

H ₂ CO Line	Freq. [GHz]	A _{ul} [10 ⁻⁴ s ⁻¹]	E _u [K]	g _u	ALMA Project Code	Observed Disks
p-3 ₀₃ -2 ₀₂	218.222	2.818	20.956	7	2016.1.00627.S	CI Tau, DM Tau, DO Tau
p-3 ₂₂ -2 ₂₁	218.476	1.571	68.094	7	2015.1.00964.S	HD 143006, J1604-2130, J1609-1908, J1612-1859, J1614-1906
					2013.1.00226.S	HD 143006, J1604-2130, J1609-1908, J1612-1859, J1614-1906
p-3 ₂₁ -2 ₂₀	218.760	1.578	68.111	7	2015.1.00964.S	AS 209, HD 163296, IM Lup, LkCa 15, MWC 480, V4046 Sgr
					2013.1.00226.S	HD 143006, J1604-2130, J1609-1908, J1612-1859, J1614-1906
o-4 ₁₄ -3 ₁₃	281.527	5.884	45.570	27	2015.1.00657.S	LkCa 15, MWC 480
p-4 ₀₄ -3 ₀₃	290.623	6.903	34.904	9	2015.1.00678.S	AS 209, DM Tau, GM Aur, IM Lup, V4046 Sgr
					2012.1.00681.S	HD 163296
					2015.1.00657.S	LkCa 15, MWC 480
p-4 ₂₃ -3 ₂₂	291.238	5.211	82.071	9	2015.1.00657.S	LkCa 15, MWC 480
p-4 ₂₂ -3 ₂₁	291.948	5.249	82.122	9	2015.1.00657.S	LkCa 15, MWC 480
o-4 ₁₃ -3 ₁₂	300.837	7.178	47.887	27	2015.1.00657.S	LkCa 15, MWC 480
o-5 ₁₅ -4 ₁₄	351.769	12.02	62.452	33	2011.0.00629.S	DM Tau, MWC 480

NOTE. — ‘o-’ and ‘p-’ denote ortho and para lines, respectively. All frequencies, Einstein coefficients (A_{ul}), upper energy levels (E_u), and degeneracies (g_u) were obtained from the Cologne Database for Molecular Spectroscopy (CDMS; Endres et al. 2016).

(both lines are observed towards one disk, DM Tau). These observations are complemented by the H₂CO 5₁₅-4₁₄ line towards two disks; the H₂CO 3₂₂-2₂₁ and 3₂₁-2₂₀ lines towards 11 disks each; and the H₂CO 4₁₄-3₁₃, 4₁₃-3₁₂, 4₂₃-3₂₂, and 4₂₂-3₂₁ lines towards two disks each. When detected, these additional lines are used to determine H₂CO excitation temperatures and column densities.

In total, nine H₂CO lines were considered, providing a total of 48 H₂CO observations across the 15 disks. All H₂CO lines and their molecular characteristics, ALMA observing projects, and corresponding disks are listed in

Table 2. The ALMA observing projects are described and referenced briefly in Table 3.

We also consider complementary dust continuum and C¹⁸O observations towards all disks except GM Aur. All CO observations were taken from ALMA observing projects 2013.1.00226.S, 2015.1.00964.S, and 2016.1.00627.S.

2.3. Data Reduction

The ALMA/NAASC staff performed the initial calibration (including bandpass, flux, and phase calibration) for each observation. The flux calibration for HD

TABLE 3
ALMA Observing Project Details and References.

ALMA Project Code, Band	On-Source Int. Times [min]	Num. of Ant.	Baseline Ranges [m]	Largest Ang. Scale ["]	References
2016.1.00627.S, 6	22-26	47	15-704	4-5	Bergner et al. (2019)
2015.1.00964.S, 6	14-21	48	15-704	5-10	Bergner et al. (2019)
2015.1.00678.S, 7	20	41	20-640	2.6-4.2*	Kastner et al. (2018); Qi et al. (2019)
2015.1.00657.S, 7	23-61	39-44	15-650	5	Loomis et al. (subm.)
2013.1.00226.S, 6	12-21	31-37	20-784	6	Huang et al. (2017)
2012.1.00681.S, 7	45	32	18-650	2.9*	Qi et al. (2015)
2011.0.00629.S, 7	35	23	17.5-380	6	Loomis et al. (2015)

NOTE. — *: The $\text{H}_2\text{CO } 4_{04}\text{-}3_{03}$ line towards DM Tau, HD 163296, and V4046 Sgr was observed with largest angular scales of $2''.6$, $2''.9$, and $4''.2$, respectively. It is possible that we are missing extended H_2CO emission for these large disks.

163296 in ALMA observing projects 2012.1.00681.S and 2013.1.00226.S was further optimized by Qi et al. (2015) and Huang et al. (2017), respectively. As the datasets were observed at different times across the past decade, we note that this initial calibration was non-uniform across the sample. After initial calibration, we used the COMMON ASTRONOMY SOFTWARE APPLICATIONS package (CASA) version 4.7.2 to self-calibrate and image each targeted line. We self-calibrated using the continuum emission from all available spectral windows, after removing any strong lines. Solution intervals were 10-100 seconds. We used intervals within this range that maximized the number of solutions with signal-to-noise ratios ≥ 2 . We carried out 2 iterations of phase calibration and 1 iteration of amplitude calibration for each line, and we used a Briggs weighting scheme and a robust value of 0.5 during the self-calibration process for all disks. There were a few exceptions to this approach. For one observation of the $\text{H}_2\text{CO } 4_{13}\text{-}3_{12}$ line towards MWC 480, we used 3 iterations of phase calibration and 2 iterations of amplitude calibration to improve phase and amplitude solutions. We did not self-calibrate the $\text{H}_2\text{CO } 4_{04}\text{-}3_{03}$ line towards DM Tau, or the $\text{H}_2\text{CO } 3_{03}\text{-}2_{02}$ line towards J1612-1859, due to low continuum peaks for these sources.

After self-calibration, we subtracted out the continuum using CASA’s `uvcontsub` function and imaged each targeted line using CASA’s `clean` algorithm. We initially used a Briggs weighting scheme and a robust value of 0.5 for `cleaning` for all disks. We created the `cleaning` masks by hand to cover the emission and little else in each channel. For the weaker lines (e.g., $\text{H}_2\text{CO } 3_{22}\text{-}2_{21}$), we recycled `clean` masks from the stronger lines available. The weaker H_2CO lines that were tentatively-detected or non-detected with `robust=0.5` were reimaged with `robust=2` to maximize sensitivity.

We did not image the two weaker $\text{H}_2\text{CO } 3_{22}\text{-}2_{21}$ and $3_{21}\text{-}2_{20}$ lines for the disks J1609-1908, J1612-1859, and J1614-1906, because the emission from the much brighter $\text{H}_2\text{CO } 3_{03}\text{-}2_{02}$ line was already very faint for these three disks.

3. IMAGE ANALYSIS

Our basic image products were image cubes, which are visualized as channel maps. Figure 2 displays the channel map for the $\text{H}_2\text{CO } 3_{03}\text{-}2_{02}$ line towards J1604-2130; channel maps for all detected lines and sources are shown in

Figure Sets A1-A3 in the Appendix. We generated Keplerian masks (e.g., Rosenfeld et al. 2013; Yen et al. 2016; Salinas et al. 2017) to extract the line emission within each channel for each disk, ultimately reducing the noise incorporated into the extracted emission. We generated each set of Keplerian masks in four steps¹: (1) calculate each disk’s line-of-sight velocities within the disk’s coordinate system, (2) incorporate thermal and turbulent broadening using the approach of Yen et al. (2016), (3) mask the pixels within each velocity channel that fall within the broadened velocity bins, and (4) convolve the masks with the average beam size. For this survey, we empirically fit for the position and inclination angles using the masks, C^{18}O emission towards each disk, and a grid-search algorithm (^{12}CO emission was used for the three disks where C^{18}O was not detected, and H_2CO emission was used for the one disk where no CO was observed). The Keplerian masks calculated for the $\text{H}_2\text{CO } 3_{03}\text{-}2_{02}$ line towards J1604-2130 are overplotted in the channel map of Figure 2. All parameters used to generate the Keplerian masks for the sample are given in Table B1 in the Appendix.

We created a velocity-integrated emission map (a.k.a. a moment-0 map) for each line and disk pair by summing up the emission within the Keplerian masks across the velocity channels in the data cube. To estimate the noise of each velocity-integrated emission map, it is important to account for how each pixel in the image appears in a different number of Keplerian masks, depending on the pixel’s location. We therefore generated an ‘rms map’ for each line and disk, which calculates the rms at each pixel *through* all masks. Each rms map has the same dimensions as the velocity-integrated emission map and holds different rms values for each pixel. For each line and disk, we randomly sampled 1000 rms maps across channels with no emission, and then took the standard deviation at each pixel. We used the median of the standard deviations as the estimate of the line emission map’s noise (σ_{map}). See Bergner et al. (2018) for the full methodology for generating rms maps.

Velocity-integrated fluxes for each line and disk were calculated by summing over the velocity-integrated emission maps pixel-by-pixel. Noise was estimated similarly to the velocity-integrated maps, i.e. from the standard

¹ The Keplerian mask package is available on GitHub (<https://github.com/jpegues/kepmask>). Version 1.0.0 was used for this paper and is available at <https://doi.org/10.5281/zenodo.3382082>.




FIG. 2.— **H₂CO 3₀₃-2₀₂ Channel Map and Keplerian Masks for J1604-2130.** The colorbars have been clipped to exclude values below zero. The outlines of the calculated Keplerian masks are drawn with dashed lines.

deviation of 1000 random samples. To create a radial emission profile for each line, we azimuthally averaged each velocity-integrated emission map. In this process, we deprojected the disks and divided them into 0.1'' rings about the host star, and then averaged the emission within each ring. Following Bergner et al. (2018), we estimated the noise within each ring as (σ_{map}/\sqrt{N}) , where N is the number of independent measurements in this ring, taken to be the number of pixels in the ring divided by the beam area (in pixels). For rings within the beam's circumference, we fixed N to be the beam circumference (in pixels) divided by the beam area. Finally, we summed the emission in the Keplerian masks as a function of velocity channel to extract a spectrum for each line and disk (shown in Figure B1 in the Appendix).

4. RESULTS

4.1. Detection Statistics

We define a *detection* as a line that fulfills all of the following criteria:

1. Peak emission is $\geq 3\sigma$ in the velocity-integrated emission map
2. Emission within the Keplerian masks is $\geq 3\sigma$ in at least 4 velocity channels of the channel map

We define a *tentative detection* as a line that fulfills at least one of the above criteria, and a *non-detection* as a line that fails all criteria. Based on these criteria, we detect the H₂CO 3₀₃-2₀₂ line towards 6/8 disks. We tentatively detect this line towards the disk J1614-1906, and do not detect it towards the disk J1612-1859. We detect the H₂CO 4₀₄-3₀₃ line towards 8/8 disks. Altogether, we report detections of H₂CO towards 13/15 disks. Of these 13 disks, we detect one line towards 8 disks, two lines towards 1 disk, three lines towards 2 disks, four lines towards 1 disk, and six lines towards 1 disk. Table 4 lists the fluxes for all detected H₂CO lines and flux upper limits for all tentative and non-detected H₂CO lines. The fluxes for the associated continuum and C¹⁸O emission are in Tables C1 and C2, respectively, in the Appendix.

4.2. Spatial Distributions of H₂CO Emission

Figure 3 presents emission maps for all observed H₂CO 3₀₃-2₀₂ and 4₀₄-3₀₃ lines, associated dust continuum, and C¹⁸O line emission. Figure 4 summarizes the emission morphologies of the disks detected in H₂CO. Figure 5 shows the radial profiles of all H₂CO 3₀₃-2₀₂ and 4₀₄-3₀₃ lines. We see a variety of H₂CO spatial structures between the 13 detected disks. The majority (8/13) of the disks are centrally peaked in H₂CO. Three disks have central dips (central depressions that are $\sim 1-3\sigma_{peak}$

deep) in H₂CO, while the last two disks have central holes (central depressions that are $\gtrsim 3\sigma_{peak}$ deep) in H₂CO. About half (6/13) of the disks show outer rings or plateaus in H₂CO emission. DM Tau, for example, has rings in its H₂CO 4₀₄-3₀₃ emission and a plateau with suggestive ring-like substructure in its H₂CO 3₀₃-2₀₂ emission.

Figure 6 compares the H₂CO radial profiles with the dust continuum and C¹⁸O radial profiles. The majority (10/13) of the disks show H₂CO emission beyond the edges of the dust continuum (defined as the radial extent at which the azimuthally-averaged continuum emission is 5% of its peak). Of the five disks that show inner holes in the dust continuum at our resolution, only one disk, J1604-2130, shows a similar inner hole in the H₂CO emission. Dust and H₂CO emission substructure thus do not necessarily coincide. The C¹⁸O and H₂CO emission morphologies are more similar than the dust and H₂CO: seven disks have similar radial extents in H₂CO and C¹⁸O, and five have similar outer slopes in H₂CO and C¹⁸O emission. However, there are also distinct differences between the H₂CO and C¹⁸O emission morphologies, indicating that H₂CO substructure is set by chemistry as well as by the disk gas distribution.

A priori, one possible explanation for the central H₂CO depressions observed is dust opacity, i.e. optically thick dust blocking out molecular emission. C¹⁸O emission profiles provide an avenue to test this. We do see C¹⁸O depressions in three disks with H₂CO inner dips or holes: IM Lup, J1604-2130, and LkCa 15 (Figure 6). However, for two of these disks (J1604-2130 and LkCa 15) the C¹⁸O depressions coincide with central holes in the dust continuum and are therefore explained by holes in gas and dust rather than by dust opacity. This leaves IM Lup as the only case where the central H₂CO depression is likely due to dust opacity, at least in part. For all disks except IM Lup, optically thick dust is likely not a major cause for the H₂CO emission substructure in the sample.

4.3. Rotational Diagram Analysis

Three or more H₂CO lines are detected towards four disks: DM Tau, J1604-2130, LkCa 15, and MWC 480. The velocity-integrated emission maps are shown in Figure 7, while the radial profiles and spectra are shown in Figure 8. For these four disks, we fit for the area-averaged H₂CO excitation temperatures ($\overline{T_{ex}}$) and column densities ($\overline{N_{tot}}$) using the rotational diagram method (e.g. Blake et al. 1987; Goldsmith & Langer 1999). This method assumes that the lines are in local thermodynamic equilibrium (LTE), that all lines originate from the same disk environment, and that the emission fills the beam. LTE should be a reasonable approximation, since the H₂CO critical densities are lower than




FIG. 3.— **H₂CO 3₀₃-2₀₂ and 4₀₄-3₀₃ Velocity-Integrated Emission Maps.** The left and right groups of columns correspond to all detected/tentatively-detected H₂CO 3₀₃-2₀₂ (3-2) and 4₀₄-3₀₃ (4-3) lines, respectively. In each group, each row shows emission for a different disk. The first column in each group shows the dust continuum at wavelengths of either 1.3mm (left group) or 1.0mm (right group), the second shows the velocity-integrated emission maps above $2\sigma_{map}$ for C¹⁸O, and the third shows the H₂CO velocity-integrated emission maps above $2\sigma_{map}$. Colorbars are provided for the H₂CO emission maps to the right of each group. The contour lines for all subplots are $[3\sigma, 5\sigma, 10\sigma, 20\sigma\dots]$; for the continuum emission σ is the continuum rms, while for the integrated emission maps σ is the corresponding σ_{map} . Beams are drawn in the lower right corners. The H₂CO 3-2 line is tentatively detected towards J1614-1906. All other H₂CO 3-2 and 4-3 lines are detected. C¹⁸O is not observed towards GM Aur and is not detected towards J1609-1908 and J1614-1906.

TABLE 4
H₂CO Emission Flux Measurements.

Disk	H ₂ CO Line	Flux* [mJy × km s ⁻¹]	Peak Flux* [mJy beam ⁻¹ × km s ⁻¹]	Int. Vel. Range [km s ⁻¹]	Chan. Width [km s ⁻¹]	rms** [mJy beam ⁻¹]	Beam Size (P.A.)	Robust Value
AS 209	4 ₀₄ -3 ₀₃	580 ± 17	19 ± 2.6	0.9 - 8.1	0.4	2.9	0.31" x 0.22" (-69.84°)	0.5
	3 ₂₂ -2 ₂₁	<82	30 ± 7.2	0.9 - 8.1	0.4	7.5	0.61" x 0.56" (-69.47°)	2.0
	3 ₂₁ -2 ₂₀	<80	21 ± 6.4	0.9 - 8.1	0.4	6.6	0.61" x 0.56" (-70.17°)	2.0
CI Tau	3 ₀₃ -2 ₀₂	420 ± 12	28 ± 3.1	2.6 - 9.0	0.2	4.8	0.67" x 0.46" (30.53°)	0.5
DM Tau	4 ₀₄ -3 ₀₃	337 ± 28	20 ± 3.1	3.6 - 8.4	0.3	5.5	0.59" x 0.51" (-81.65°)†	0.5
	3 ₀₃ -2 ₀₂	462 ± 17	24 ± 2.7	3.6 - 8.6	0.2	4.9	0.66" x 0.50" (35.24°)†	0.5
	5 ₁₅ -4 ₁₄	256 ± 37	39 ± 6.6	3.6 - 8.6	0.2	13	0.77" x 0.60" (-25.76°)†	0.5
DO Tau	3 ₀₃ -2 ₀₂	85 ± 17	36 ± 3.7	3.4 - 8.6	0.4	3.4	0.90" x 0.58" (32.19°)	2.0
GM Aur	4 ₀₄ -3 ₀₃	1074 ± 17	44 ± 3.6	-0.1 - 11.3	0.3	3.1	0.29" x 0.19" (-2.69°)	0.5
HD 143006	3 ₀₃ -2 ₀₂	161 ± 9.4	32 ± 3.0	5.4 - 10.2	0.2	4.7	0.68" x 0.47" (-74.10°)	0.5
	3 ₂₂ -2 ₂₁	<30 [×]	8.2 ± 2.3	5.4 - 10.2	0.2	4.0	0.87" x 0.57" (-71.56°)	2.0
	3 ₂₁ -2 ₂₀	<32	17 ± 3.2	5.4 - 10.2	0.2	4.2	0.87" x 0.57" (-71.74°)	2.0
HD 163296	4 ₀₄ -3 ₀₃	942 ± 43	51 ± 5.9	-1.4 - 13.0	0.4	6.9	0.57" x 0.52" (86.68°)†	0.5
	3 ₂₂ -2 ₂₁	<132	<30	-1.4 - 13.0	0.4	3.8	0.69" x 0.59" (67.20°)	2.0
	3 ₂₁ -2 ₂₀	119 ± 40	47 ± 9.8	-1.4 - 13.0	0.4	3.4	0.72" x 0.59" (70.98°)	2.0
IM Lup	4 ₀₄ -3 ₀₃	1063 ± 25	45 ± 4.3	0.2 - 8.6	0.2	7.4	0.45" x 0.39" (70.99°)†	0.5
	3 ₂₂ -2 ₂₁	<55 [×]	12 ± 3.5	0.6 - 8.6	0.4	3.7	0.67" x 0.49" (-76.11°)	2.0
	3 ₂₁ -2 ₂₀	<45 [×]	9.1 ± 3.0	0.6 - 8.6	0.4	3.2	0.66" x 0.48" (-76.35°)	2.0
J1604-2130	3 ₀₃ -2 ₀₂	821 ± 6.4	60 ± 1.6	3.6 - 5.4	0.2	4.5	0.63" x 0.44" (-76.97°)	0.5
	3 ₂₂ -2 ₂₁	157 ± 7.1	18 ± 1.5	3.6 - 5.4	0.2	4.6	0.63" x 0.44" (-77.13°)	0.5
	3 ₂₁ -2 ₂₀	151 ± 6.6	18 ± 1.3	3.6 - 5.4	0.2	4.7	0.63" x 0.44" (-77.51°)	0.5
J1609-1908	3 ₀₃ -2 ₀₂	43 ± 13	36 ± 4.2	-1.6 - 9.2	0.4	3.3	0.81" x 0.53" (-72.33°)	2.0
J1612-1859	3 ₀₃ -2 ₀₂	<26 [×]	<12	-0.8 - 10.4	0.4	3.0	0.81" x 0.53" (-71.66°)	2.0
J1614-1906	3 ₀₃ -2 ₀₂	<30	25 ± 5.2	-3.0 - 10.6	0.4	3.3	0.80" x 0.53" (-72.36°)	2.0
LkCa 15	4 ₀₄ -3 ₀₃	662 ± 24	20 ± 2.8	2.1 - 10.2	0.3	3.3	0.26" x 0.20" (-14.57°)	0.5
	3 ₂₂ -2 ₂₁	<77	16 ± 3.3	2.1 - 10.2	0.3	4.6	0.64" x 0.51" (13.70°)	2.0
	3 ₂₁ -2 ₂₀	<52	15 ± 2.8	2.1 - 10.2	0.3	3.5	0.64" x 0.50" (14.42°)	2.0
MWC 480	4 ₁₄ -3 ₁₃	1199 ± 16	124 ± 3.2	1.9 - 10.7	1.1	2.2	0.82" x 0.47" (-36.05°)	0.5
	4 ₁₃ -3 ₁₂	1342 ± 33	169 ± 5.7	1.9 - 10.7	1.1	3.9	0.80" x 0.48" (-37.21°)	0.5
	4 ₂₃ -3 ₂₂	<62	29 ± 4.0	1.9 - 10.7	1.1	2.3	0.99" x 0.58" (-40.33°)	2.0
	4 ₂₂ -3 ₂₁	37 ± 20	32 ± 3.6	1.9 - 10.7	1.1	2.2	0.99" x 0.58" (-40.35°)	2.0
	4 ₀₄ -3 ₀₃	239 ± 21	74 ± 8.3	-1.4 - 11.8	1.1	3.5	1.02" x 0.83" (-3.58°)	0.5
	3 ₂₂ -2 ₂₁	36 ± 30	55 ± 9.8	-1.2 - 11.2	0.4	4.0	0.76" x 0.50" (13.92°)	2.0
	3 ₂₁ -2 ₂₀	55 ± 22	60 ± 8.3	-1.2 - 11.2	0.4	3.0	0.76" x 0.50" (14.31°)	2.0
	4 ₁₄ -3 ₁₃	297 ± 16	49 ± 4.9	-1.4 - 11.8	1.1	2.1	0.90" x 0.46" (-28.87°)	0.5
	4 ₁₃ -3 ₁₂	452 ± 32	86 ± 8.4	-1.4 - 11.8	1.1	3.6	0.87" x 0.47" (-28.96°)	0.5
	4 ₂₃ -3 ₂₂	<59 [×]	<16	-1.4 - 11.8	1.1	2.1	1.08" x 0.59" (-32.09°)	2.0
V4046 Sgr	4 ₂₂ -3 ₂₁	<61	30 ± 6.0	-1.4 - 11.8	1.1	2.1	1.08" x 0.59" (-32.10°)	2.0
	5 ₁₅ -4 ₁₄	396 ± 54	104 ± 13	-1.2 - 11.2	0.4	12	0.88" x 0.54" (-22.89°)†	0.5
	4 ₀₄ -3 ₀₃	1218 ± 40	33 ± 6.3	-4.2 - 10.2	0.3	5.8	0.54" x 0.45" (-67.39°)†	0.5
	3 ₂₂ -2 ₂₁	<69	<19	-4.2 - 10.2	0.3	3.9	0.86" x 0.54" (-84.12°)	2.0
	3 ₂₁ -2 ₂₀	<67	14 ± 3.6	-4.2 - 10.2	0.3	3.4	0.85" x 0.53" (-84.18°)	2.0

NOTE. — *: The velocity-integrated fluxes (Column 3) were measured within the radii of the Keplerian masks (Table B1 in the Appendix). The 3 σ upper limit is given for tentative and non-detections, and the upper limits of non-detections are additionally marked with a [×]. The peak fluxes (Column 4) are the peaks of the velocity-integrated emission maps; note the difference in unit compared to the velocity-integrated fluxes. The uncertainty in each peak flux is the standard deviation of the peaks across 1000 random samples. The 3 σ upper limit is given whenever the peak flux is less than 3 σ itself. Uncertainties do not include absolute flux uncertainties. **: The channel rms was estimated as the standard deviation of 1000 1"-by-1" random samples. †: The following UV tapers were applied to reduce the effects of small-scale noise in the image: 0.25" for the H₂CO 4₀₄-3₀₃ line towards IM Lup and V4046 Sgr, 0.40" for the 5₁₅-4₁₄ line towards MWC 480, and 0.50" for the 4₀₄-3₀₃ line towards HD 163296. UV tapers of 0.25", 0.50", and 0.60" were applied to the 3₀₃-2₀₂ (0.25"), 5₁₅-4₁₄ (0.50"), and 4₀₄-3₀₃ (0.60") lines towards DM Tau for the same reason, as well as to make the beam sizes comparable for these lines.




FIG. 4.— $\text{H}_2\text{CO } 3_{03-2_{02}}$ and $4_{04-3_{03}}$ Grid of Morphologies. The rows classify the H_2CO detections by the shape of the central emission at/inwards of the emission peak. The columns classify the detections based on the substructure beyond the central emission when it is significant in both the integrated emission maps and radial profiles.

typical disk densities in the midplane and warm molecular disk layers (see Figure D1 in the Appendix). It is a reasonable first approximation that the emission fills the beam, since the H_2CO emission is always resolved; however unresolved emission substructure may be present.

Following Loomis et al. (2018), we construct opacity-corrected rotational diagrams by relating $\overline{T_{ex}}$ and $\overline{N_{tot}}$ to the area-averaged column density $\overline{N_u}$ and degeneracy g_u of each line's upper state:

$$\begin{aligned}
 \ln\left(\frac{\overline{N_u}}{g_u}\right) &= \ln\left(\frac{\overline{N_u^{thin}} C_{\tau_{ul}}}{g_u}\right) \\
 &= \ln\left[\left(\frac{4\pi \int S dv}{A_{ul} \Omega_a hc}\right) \left(\frac{\tau_{ul}}{1 - e^{-\tau_{ul}}}\right)\right] \\
 &= \left[\ln\left(\frac{\overline{N_{tot}}}{Q\{\overline{T_{ex}}\}}\right) - \frac{E_u}{\overline{T_{ex}}}\right], \quad (1)
 \end{aligned}$$

where $\overline{N_u^{thin}}$ is the area-averaged column density if the line is actually optically thin, $\int S dv$ is the area-averaged velocity-integrated flux (in units of spectral flux density \times velocity), A_{ul} is the Einstein coefficient, and Ω_a is the solid angle of area over which $\int S dv$ is averaged. Next, h is the Planck constant, c is the speed of light, $Q\{T\}$ is the partition function at temperature T (interpolated from T and $Q\{T\}$ values from CDMS; Müller et al. 2005), and E_u is the upper energy level. Finally, $C_{\tau_{ul}} = \tau_{ul}/(1 - e^{-\tau_{ul}})$ is the opacity correction factor (e.g., Goldsmith & Langer 1999), where the opacity of the line τ_{ul} can be calculated as:

$$\begin{aligned}
 \tau_{ul} &= \left(\frac{A_{ul} c^3}{8\pi \nu^3 \Delta v_l}\right) N_u (e^{h\nu/(k_B T_{ex})} - 1) \\
 &= \left(\frac{A_{ul} c^3}{8\pi \nu^3 \Delta v_l}\right) N_u^{thin} C_{\tau_{ul}} (e^{h\nu/(k_B T_{ex})} - 1)
 \end{aligned}$$

$$\begin{aligned}
 (1 - e^{-\tau_{ul}}) &= \left(\frac{A_{ul} c^3}{8\pi \nu^3 \Delta v_l}\right) N_u^{thin} (e^{h\nu/(k_B T_{ex})} - 1) \\
 \tau_{ul} - \frac{\tau_{ul}^2}{2} + \frac{\tau_{ul}^3}{6} &\approx \left(\frac{A_{ul} c^3}{8\pi \nu^3 \Delta v_l}\right) N_u^{thin} (e^{h\nu/(k_B T_{ex})} - 1), \quad (2)
 \end{aligned}$$

where $(1 - e^{-\tau_{ul}}) \approx \tau_{ul} - \frac{\tau_{ul}^2}{2} + \frac{\tau_{ul}^3}{6}$, with accuracy within 99%, 95%, and 90% at τ_{ul} of 0.6, 1.0, and 1.25 respectively. ν is the line frequency, k_B is the Boltzmann constant, and Δv_l is the intrinsic line width, assumed to be a combination of thermal and turbulent broadening (e.g., Flaherty et al. 2015):

$$\Delta v_l \approx \sqrt{\left(\sqrt{\frac{2k_B T_{ex}}{m_{\text{H}_2\text{CO}}}}\right)^2 + \left(t_0 \sqrt{\frac{k_B T_{ex}}{\mu m_H}}\right)^2}, \quad (3)$$

where $m_{\text{H}_2\text{CO}}$ and m_H are the masses of H_2CO and H respectively, $\mu = 2.37$ is the assumed mean molecular weight of the disk, and $t_0 \approx 0.01$ is the assumed prefactor for the turbulent contribution (e.g., Flaherty et al. 2015).

Combining Equations 1 and 2, we initially solve for the opacity-uncorrected $\overline{T_{ex}}$ and $\overline{N_{tot}}$ using a linear fit assuming $C_{\tau_{ul}} = 1$, and then we correct the rotational diagram for the opacity using an iterative approach: (1) solve for each τ_{ul} using Equation 2, (2) recalculate each $\overline{N_u}/g_u$ using Equation 1, (3) redo the linear fit to the rotational diagram (the last line of Equation 1), and use the fit to (4) recalculate the values for $\overline{T_{ex}}$ and $\overline{N_{tot}}$ from the fit's slope and y-intercept, respectively. We repeat this process until we reach convergence, defined as when the difference in the τ_{ul} values and $\overline{T_{ex}}$ between two iterations is at most 0.001% (or, in the case of the uncertain spike in excitation temperature for LkCa 15, until 5000 iterations have passed).

Figure 9 displays the disk-averaged rotational diagrams. The disk area is measured out to where at least one H_2CO line has azimuthally-averaged intensities above 1σ . Table 5 lists the extracted disk-averaged excitation temperatures and column densities. The disk-averaged excitation temperatures span $\sim 11\text{-}37\text{K}$. The disk-averaged column densities span a little more than an order of magnitude: J1604-2130 has the largest column density at $2.1 \times 10^{13} \text{cm}^{-2}$, while MWC 480 has the lowest at $1.9 \times 10^{12} \text{cm}^{-2}$. Notably the energy ranges covered by the H_2CO lines for each disk are similar, implying that these results are not due to excitation effects from using different H_2CO lines.

For each disk we also construct azimuthally-averaged rotational diagrams. For each fit, we extract the integrated flux for each line from within the same narrow ($0.1''$) rings of the deprojected disk around the host star. We then construct a rotational diagram for each ring, assuming that each ring has uniform temperature and




FIG. 5.— **H₂CO 3₀₃-2₀₂ and 4₀₄-3₀₃ Radial Profiles.** H₂CO 3₀₃-2₀₂ (3-2) and 4₀₄-3₀₃ (4-3) lines are depicted in light and dark purple, respectively, above and below the black line. DM Tau appears twice because both lines are observed towards DM Tau. The H₂CO 3-2 line is tentatively detected towards J1614-1906. All other H₂CO 3-2 and 4-3 lines are detected. The shading represents the 1 σ uncertainties, which do not include absolute flux uncertainties. Proposed locations for the midplane CO snowlines of the disks, extracted from the literature and listed in Table 6 (Section 5.2), are overplotted with gray vertical shading. The light gray or dark gray regions indicate the snowline locations were derived from either disk temperature models or N₂H⁺ fits, respectively. The average half-power beam widths are represented by the horizontal bars in the lower right corners.

TABLE 5
Disk-Averaged H₂CO Excitation Temperatures and Column Densities.

Disk	$\overline{T_{ex}}$ [K]	$\overline{N_{tot}}$ [10 ¹² cm ⁻²]	τ_{ul} (H ₂ CO Line)
DM Tau	11 ± 0.39	2.4 ± 0.34	0.85 ± 0.20 (3 ₀₃ -2 ₀₂), 0.35 ± 0.08 (4 ₀₄ -3 ₀₃), 0.19 ± 0.06 (5 ₁₅ -4 ₁₄)
J1604-2130	37 ± 3.6	21 ± 3.2	0.44 ± 0.06 (3 ₀₃ -2 ₀₂), 0.07 ± 0.01 (3 ₂₁ -2 ₂₀), 0.07 ± 0.01 (3 ₂₂ -2 ₂₁)
LkCa 15	29 ± 8.8	3.1 ± 1.4	0.14 ± 0.04 (4 ₀₄ -3 ₀₃), 0.26 ± 0.08 (4 ₁₃ -3 ₁₂), 0.26 ± 0.07 (4 ₁₄ -3 ₁₃)
MWC 480	21 ± 2.2	1.9 ± 0.47	0.16 ± 0.03 (4 ₀₄ -3 ₀₃), 0.32 ± 0.07 (4 ₁₃ -3 ₁₂), 0.22 ± 0.04 (4 ₁₄ -3 ₁₃), 0.19 ± 0.05 (5 ₁₅ -4 ₁₄)

NOTE. — Values were derived from the linear fits of Figure 9. Errors and fits include 10% absolute flux uncertainties, added in quadrature to the velocity-integrated flux uncertainties.

column density (see Loomis et al. (2018) for more details). Figure 10 displays the azimuthally-averaged excitation temperatures. In all disks except J1604-2130, the radially-resolved temperature profiles are close to consistent with a constant temperature. The excitation temperature increase towards J1604-2130 coincides with the inner dust edge, and may be due to direct radiation from the host star. Otherwise, the H₂CO excitation temperature across disks are ~20-50~K, with the exception of

DM Tau, where most H₂CO appears to be at ~10-15~K. We note that DM Tau is an M-star with a massive disk, and it is therefore perhaps unsurprising that it hosts the coldest H₂CO.

Figure 10 also displays the azimuthally-averaged H₂CO column densities. The column density profiles generally vary between the four disks, closely following their H₂CO line emission profiles. Notably the observed H₂CO line emission plateau towards DM Tau seems to




FIG. 6.— Same as Figure 5, Normalized with C^{18}O (orange) and Dust Continuum (gray) Radial Profiles for Comparison. C^{18}O is not detected towards J1609-1908 and J1614-1906, and we have no CO observations for GM Aur.

be due to a near-flat column density profile beyond ~ 100 AU in this disk. Interestingly, H_2CO has the smallest column density towards MWC 480, the only disk of these four that is irradiated by a Herbig Ae star.

4.4. Radial H_2CO Column Density Profiles across the Sample

For each disk where a single H_2CO line was detected, we estimate a range of azimuthally-averaged column density profiles using Equation 1 for H_2CO excitation temperatures between 20-50K. Figure 11 displays the resulting H_2CO column density profiles, demonstrating that they are quite insensitive to excitation temperature assumptions in this range. Across the disks, estimated column densities span nearly three orders of magnitude, from $\sim 5 - 5000 \times 10^{11} \text{cm}^{-2}$. The highest column densities are seen at about 25 AU and 100 AU for the disks GM Aur and J1604-2130, respectively; notably, these locations are near the inner edges of the holes in the disks' dust continuum. The lowest column densities are seen for the disks HD 163296, MWC 480, and V4046 Sgr, which present low column densities across their entire disks. Notably HD 163296 and MWC 480 are the only Herbig Ae disks in the sample, and V4046 Sgr is a very old disk.

Figure 12 plots the median radial H_2CO column density profiles from Figure 11, grouped by stellar age on the

left and stellar type (T Tauri or Herbig Ae) on the right. The right-hand panel of Figure 12 shows that the median Herbig Ae H_2CO column densities are generally lower than the T Tauri medians. Note that this trend is tentative, since there are only two Herbig Ae disks in the sample, and there is overlap between the T Tauri and Herbig Ae distributions. The left-hand panel shows that the median H_2CO column densities for the old disks ($\geq 5\text{Myr}$) are similar to the Herbig Ae medians, because both Herbig Ae disks are $\geq 5\text{Myr}$. However, given that the $\geq 5\text{Myr}$ disk J1604-2130 hosts some of the largest H_2CO column densities in the survey, it seems more likely that the tentative trend is explained by stellar type rather than stellar age. There is no trend in H_2CO column density with dust continuum size (not shown).

5. DISCUSSION

5.1. H_2CO Formation Chemistry

Interpretations of our observed H_2CO morphologies and profiles rely on an understanding of the possible astrochemical origins of H_2CO in disks, which we briefly review here.

Astrochemical laboratory experiments and models have shown that H_2CO can form efficiently from both gas-phase and grain-surface chemistry. In the gas phase, H_2CO can form through multiple pathways, including




FIG. 7.— **Velocity-Integrated Emission Maps for Disks with Multiple H₂CO Detections.** Each row plots emission for a different disk, and each subplot gives the velocity-integrated emission map above $2\sigma_{map}$ of a different line of H₂CO. Each subplot has an individual color scaling. The contour lines are $[3\sigma_{map}, 5\sigma_{map}, 10\sigma_{map}, 20\sigma_{map}, \dots]$. Beams are drawn in the lower right corners. All lines have been smoothed and circularized using CASA’s IMSMOOTH function so that the beams are identical across each disk. Subplots outlined in black show the lines included in the rotational diagrams of Section 4.3. Subplots outlined in gray show the lines that were excluded from the diagrams due to low signal-to-noise in the outer disk regions.




FIG. 8.— **Radial Profiles and Spectra for Lines used in Rotational Diagrams.** The top and bottom rows show radial profiles and spectra, respectively, for the H₂CO lines used in the rotational diagrams (Section 4.3). Each column corresponds to a different disk, and each line style and color depict a different H₂CO line, as listed in the legends in the top row. The shading represents the 1σ uncertainties. The average half-power beam widths are represented by the horizontal bars in the lower right corners of the top row.




FIG. 9.— **Disk-Averaged Rotational Diagrams.** The purple points represent the $\overline{N_u/g_u}$ values extracted from the observations, while the black lines indicate the best linear fits taking into account calculated line opacities. Disk areas extend out to where at least one H₂CO line has azimuthally-averaged intensities above 1 σ . Errorbars are 1 σ and include 10% absolute flux uncertainties, added in quadrature to the velocity-integrated flux uncertainties.




FIG. 10.— **Radial H₂CO Excitation Temperatures and Column Densities.** The top and bottom rows show H₂CO excitation temperatures and column densities, respectively, as a function of radial distance from the host star. The shaded regions give the 1 σ errors in the fits and include 10% absolute flux uncertainties, added in quadrature to the flux uncertainties within each radial ring around the host star. The horizontal gray lines show the disk-averaged excitation temperatures and column densities from Table 5. All profiles are estimated out to where at least three H₂CO lines have azimuthally-averaged intensities above 1 σ . The subplots' x-axes are extended to show the disk area over which the disk-averaged temperatures and column densities are measured, out to where at least one H₂CO line has azimuthally-averaged intensities above 1 σ . The average half-power beam widths are represented by the horizontal bars in the lower right corners of the bottom panels.




FIG. 11.— **Radial H₂CO Column Density Profiles for the Disk Sample.** The panels show azimuthally-averaged H₂CO column densities for the disks with detected H₂CO 3₀₃-2₀₂ and 4₀₄-3₀₃ lines, as a function of radial distance from the host star. Colors and line styles indicate different assumed excitation temperatures (listed in the bottom panel). Fitted column densities from Figure 10 are overplotted in solid purple. Errors, which include 10% absolute flux uncertainties, are shown as shaded regions for the column densities at 30K and for the fitted column densities. The average half-power beam widths are represented by the horizontal bars in the lower right corners.


the neutral-neutral reaction (e.g. Fockenberg & Preses 2002; Atkinson et al. 2006):



Under laboratory conditions from ~ 290 - 930 K, the H₂CO yield from this reaction is roughly constant (e.g., Fockenberg & Preses 2002). Astrochemical disk modeling has shown that at low temperatures (< 100 K), this pathway is the dominant source of gas-phase H₂CO formation (van der Marel et al. 2014; Loomis et al. 2015), and that gas-phase production of H₂CO in general is most efficient in the warm, dense inner regions of the

disk (Loomis et al. 2015).

On grain surfaces, H₂CO can form through the sequential hydrogenation of CO ice, which also produces CH₃OH if the hydrogenation continues (e.g. Hiraoka et al. 1994; Hiraoka et al. 2002; Watanabe & Kouchi 2002; Hidaka et al. 2004; Watanabe et al. 2004; Fuchs et al. 2009):



We expect CO hydrogenation to produce H₂CO efficiently beyond the CO snowline. However, since H₂CO grain-surface production is possible as long as some CO is




FIG. 12.— **Radial H₂CO Column Density Profiles Grouped by Stellar Characteristics.** The panels plot statistical summaries of the azimuthally-averaged H₂CO column densities as a function of radial distance from the host star, assuming an excitation temperature of 30K (from Figure 11). The statistics are computed at intervals of 25 AU, and only for disks that are within the bounds of the Keplerian masks at that radius. The dots indicate the median values for each group and are connected by solid lines to guide the eye. The shaded regions span the 16th-84th percentile ranges. Colors group the column densities by stellar age on the left and stellar type on the right (from Table 1).

on the grains long enough to react with atomic hydrogen, H₂CO grain-surface production likely also occurs somewhat interior to the CO snowline (Öberg et al. 2017).

H₂CO is less volatile than CO; the desorption energies of H₂CO off of different icy surfaces range from 3300-3700K (Noble et al. 2012), while the desorption energies of ¹³CO range from 900-1600K (Fayolle et al. 2016). Therefore H₂CO freezes out at temperatures higher than ~80K, which is interior to the CO snowline at smaller radial distances in the disks than we are sensitive to. However, frozen-out H₂CO beyond the H₂CO snowline can still be released into the gas via non-thermal desorption (e.g. Walsh et al. 2014; Loomis et al. 2015; Öberg et al. 2017; Fraud et al. 2019).

5.2. H₂CO and the CO Snowline

To constrain H₂CO formation pathways in our survey, we compare the observed H₂CO emission profiles with estimated locations of CO snowlines at the disk midplane. Studies have inferred CO snowlines for disks using CO isotopologues and chemical tracers of CO freeze-out, and found freeze-out temperatures between ~17-26~K (e.g. Mathews et al. 2013; Qi et al. 2013, 2015; Öberg et al. 2015a; Schwarz et al. 2016; Pinte et al. 2018). For DM Tau, GM Aur, HD 163296, and LkCa 15, we use the CO snowline locations fitted using N₂H⁺ as a tracer by Qi et al. (2015) and Qi et al. (2019). For the nine other disks with detected H₂CO, we estimate midplane snowlines at locations corresponding to a similar temperature range of 17-25K, extracted from the disk toy models of Bergner et al. (2019) and Loomis et al. (subm.). Note that these model-based estimates are very uncertain, and in at least two cases (AS 209 and V4046 Sgr) they overestimate the CO snowline location compared to the location based on N₂H⁺ (Qi et al. 2019). Three disks DO Tau, HD 143006, and J1609-1908 have CO snowline ranges that are too broad to be informative, and we exclude them from further analysis.

Figure 5 compares the H₂CO profiles and the inferred

TABLE 6
CO Midplane Snowline Locations.

Disk	CO Midpl. Snowline Range [AU]	Reference
AS 209	50-111	Bergner et al. (2019)
CI Tau	64-166	Bergner et al. (2019)
DM Tau	45-85	Qi et al. (2019)
DO Tau	76-310	Bergner et al. (2019)
GM Aur	40-58	Qi et al. (2019)
HD 143006	59-251	Bergner et al. (2019)
HD 163296	69-83	Qi et al. (2015)
IM Lup	38-69	Bergner et al. (2019)
J1604-2130	43-64	Bergner et al. (2019)
J1609-1908	36-73	Bergner et al. (2019)
J1612-1859	36-78	Bergner et al. (2019)
J1614-1906	31-71	Bergner et al. (2019)
LkCa 15	48-64	Qi et al. (2019)
MWC 480	90-150*	Loomis et al. (subm.)
V4046 Sgr	40-85	Bergner et al. (2019)

NOTE. — All locations have been scaled to new GAIA distances (Table 1). *: The CO snowline was estimated as 120 AU by Loomis et al. (subm.); here we broaden the snowline by ± 30 AU to account for the resolution of the data.

CO snowline locations. There is significant H₂CO emission both interior and exterior to the CO snowline of all disks. A majority (6/10) of the H₂CO radial profiles peak interior to the CO snowline, but have rings or plateaus in H₂CO at or beyond the CO snowline. Two of the remaining disks, IM Lup and J1604-2130, have H₂CO radial profiles that peak beyond the CO snowline. One of the last two disks, MWC 480, may show more substructure at higher resolution. The simplest explanation for these H₂CO morphologies is that H₂CO is produced through both gas-phase and grain-surface chemistry in disks. The substantial central H₂CO emission found interior to the CO snowline implies an inner component of H₂CO produced in the gas phase and/or due to sublimation of inward-moving grains. The rings and local

enhancements in H₂CO emission found at or beyond the CO snowline are best explained by grain-surface production, since the efficiency of H₂CO gas-phase formation is expected to decrease with disk radius.

It is possible that H₂CO ices inherited from the preceding protostellar stage are contributing to the observed H₂CO emission in disks. While we would expect inherited H₂CO ices to mainly contribute to emission in the inner disk, where H₂CO desorption would be efficient, we cannot exclude a smaller contribution in the outer disk, which may arise from the mixing of midplane ices into the warm upper layers of the disk. However, there should be no relationship between this emission and the CO snowline, which suggests that inheritance alone cannot explain the observed H₂CO emission morphologies.

Our results thus indicate that the two-component H₂CO model discussed by Loomis et al. (2015), Carney et al. (2017), and Öberg et al. (2017) for three specific disks, containing one warm inner component and one cold component exterior to the CO snowline, likely holds for disks generally. Furthermore, it is likely that H₂CO is commonly produced through both gas-phase and grain-surface chemistry in disks.

We expect that the H₂CO formed through CO hydrogenation beyond the CO snowline will coincide with the formation of complex oxygen-bearing organics, particularly methanol (CH₃OH), as these molecules can form along the same hydrogenation pathway as H₂CO (e.g., Hiraoka et al. 1994; Watanabe & Kouchi 2002). However, since there is also some contribution from gas-phase chemistry to the total H₂CO budget at these radii, it is difficult to assess how much of these COMs are present. Detailed modeling, as well as a few benchmarking disks with both H₂CO and COM detections, are needed to quantify the relationship between H₂CO and COM production in disks. In the meantime, we conclude that a majority of disks do have an active organic ice chemistry, and that COM production should be expected in the outer regions of most protoplanetary disks.

The exception to this conclusion may be disks around Herbig Ae stars, because their warmer temperatures allow for limited CO freeze-out. Our survey tentatively shows that T Tauri disks have more H₂CO than Herbig Ae disks, despite how the Herbig Ae disks present the highest C¹⁸O flux densities. This finding is consistent with the single-dish survey of H₂CO in disks conducted by Guilloteau et al. (2013), which showed that warmer disks tended to have less H₂CO emission. It is also consistent with the Submillimeter Array (SMA) survey conducted by Öberg et al. (2010), which noted that the lower detection rates towards Herbig Ae disks could be because their midplanes are too warm for a cold CO-ice based chemistry. Therefore our results could be explained by a small or non-existing production of H₂CO through CO ice hydrogenation in Herbig Ae disks. More spatially-resolved observations of H₂CO towards Herbig Ae disks are needed to further evaluate this tentative trend.

5.3. Relationship between H₂CO and Dust Substructure

Carney et al. (2017) and Öberg et al. (2017) note three ways that H₂CO could be affected by a hole or edge in the dust continuum: (1) through enhanced temperatures due

to greater irradiation in regions with fewer solids (Cleeves et al. 2011; Cleeves 2016); (2) through increased photodesorption of H₂CO off the grains from the increased irradiation (Öberg et al. 2015a); and/or (3) through increased CO photodissociation in the upper layers, since more CO in the upper layers could produce more hydrocarbon radicals and atomic oxygen in the gas, and then possibly allow for more efficient gas-phase production of H₂CO (Carney et al. 2017; Öberg et al. 2017). Five of the disks in our survey (DM Tau, GM Aur, J1604-2130, LkCa 15, and V4046 Sgr) have inner holes in the sub-millimeter/millimeter dust continuum that are detected at our spatial resolution. For these five disks, we see no universal trend in H₂CO emission morphologies in the inner disks. Four of the five disks show centrally-peaked H₂CO emission despite the central depletions in dust continuum. The final disk, J1604-2130, does show an inner hole in the H₂CO emission. Notably J1604-2130 also has a large inner hole in its C¹⁸O emission (see also Zhang et al. 2014; van der Marel et al. 2015). We also do not see a consistent trend in H₂CO emission morphologies at the outer dust continuum edges, but four of the disks show clear bumps in H₂CO emission right at those edges. It therefore appears that, while dust substructure does influence the distribution of H₂CO in disks, there is no single relationship between the two.

5.4. H₂CO Excitation Temperatures

The H₂CO excitation temperature should reflect the temperature of the vertical layer in the disk within which the gas resides, as long as the densities are high enough for LTE. T Tauri disk models predict that beyond ~50 AU (the scale we are most sensitive to), the midplane will be 10-30K, the molecular layer 20-50K, and the atmosphere 50-500K (e.g., Walsh et al. 2010). Herbig Ae disks will have warmer temperatures at the same radial distances, while low-mass M-star disks will have colder temperatures. In the four disks where we could measure H₂CO excitation temperatures, three are consistent with the warm molecular layer, and one (DM Tau) with the disk midplane, indicating that the vertical distribution of H₂CO may vary between disks.

Our radially-resolved rotational diagram analysis is largely consistent with a constant radial temperature per disk in the outer disks beyond 100-200 AU. This suggests that for three of the four disks, H₂CO is consistently emitting from the molecular layer at/above CO freeze-out temperatures. This is in accordance with model predictions, which find the most abundant H₂CO in the molecular layer above 20K (Walsh et al. 2010). However, since error bars are currently large (typically 10~K or more), we need more precise observations in order to better constrain radial H₂CO excitation temperatures.

6. SUMMARY

We have conducted an ALMA survey of H₂CO towards 15 protoplanetary disks. We have analyzed the H₂CO morphologies, excitation temperatures, and column densities, and their relationships to stellar and disk characteristics. We summarize our main findings below:

1. We report a total of 26 H₂CO detections towards 13/15 disks, consisting of one line towards 8 disks,

- two lines towards 1 disk, three lines towards 2 disks, four lines towards 1 disk, and six lines towards 1 disk. We report an additional 11 tentative H₂CO detections across 7/15 disks.
2. Over half of the disks have centrally-peaked H₂CO emission, while the rest have central dips or holes in H₂CO emission. About half of the disks show rings or a plateau in H₂CO emission in the outer disk.
 3. We measure disk-averaged and azimuthally-averaged H₂CO excitation temperatures for four disks with multiple H₂CO line detections. Three of the four disks have disk-averaged excitation temperatures of 21-37K, while the fourth disk, DM Tau, is colder at 11K. The azimuthally-averaged temperatures show an increase at the inner dust edge of J1604-2130, but otherwise close to constant radial temperatures for the disks.
 4. In addition to the disks we could directly model, we estimate radial H₂CO column densities for all disks assuming excitation temperatures of 20-50K. The column densities between and across the disks of our survey span three orders of magnitude, from $\sim 5 - 5000 \times 10^{11} \text{cm}^{-2}$. The highest H₂CO column densities are found near the inner dust continuum edges of the disks GM Aur and J1604-2130. The lowest H₂CO column densities are found across the Herbig Ae disks HD 163296 and MWC 480, as well as in the old disk V4046 Sgr.
 5. Based on simple estimates of CO midplane snowline locations, significant H₂CO emission is present both interior and exterior to the CO snowline in all disks, with ring-like structures in the emission occurring beyond the CO snowline in half of the disks. This suggests that H₂CO is commonly formed through both gas-phase and grain-surface pathways in protoplanetary disks, and that a majority of disks present an active organic ice chemistry that is likely also producing COMs. The low H₂CO column densities toward the two Herbig Ae disks in the sample may be due to a smaller degree of CO ice chemistry in these warmer disks.

This paper has benefited from discussions with Alyssa Goodman, Dimitar Sasselov, and David Wilner.

Jamila Pegues gratefully acknowledges the support of the National Science Foundation (NSF) Graduate Research Fellowship through Grant Numbers DGE1144152 and DGE1745303. Karin Öberg gratefully acknowledges

the support of the Simons Foundation through a Simons Collaboration on the Origins of Life (SCOL) PI Grant (Number 321183). Jane Huang gratefully acknowledges support from the NSF Graduate Research Fellowship under Grant No. DGE-1144152. Geoffrey Blake gratefully acknowledges support from the NSF (Grant AST-1514670) and NASA (Grant NNX16AB48G). Jes K. Jørgensen gratefully acknowledges support from the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme through ERC Consolidator Grant "S4F" (Grant Agreement No. 646908). Kamber Schwarz, a Sagan Fellow, gratefully acknowledges the support of NASA through Hubble Fellowship Program Grant HST-HF2-51419.001, awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS5-26555.

This paper makes use of the following ALMA data:

- ADS/JAO.ALMA#2011.0.00629.S
- ADS/JAO.ALMA#2012.1.00681.S
- ADS/JAO.ALMA#2013.1.00226.S
- ADS/JAO.ALMA#2015.1.00657.S
- ADS/JAO.ALMA#2015.1.00678.S
- ADS/JAO.ALMA#2015.1.00964.S
- ADS/JAO.ALMA#2016.1.00627.S

ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), MOST and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement.

All computer code used for this research was written in Python (version 2.7). All plots were generated using Python's Matplotlib package (Hunter 2007). This research also made use of Astropy (<http://www.astropy.org>), a community-developed core Python package for Astronomy, and the NumPy (Oliphant 2006-) and SciPy (Jones et al. 2001-) Python packages.

APPENDIX

Fig. Set A1. Channel maps of detected H₂CO 3₀₃-2₀₂ and 4₀₄-3₀₃ above 2σ.

Fig. Set A2. Channel maps of detected C¹⁸O above 2σ.

Fig. Set A3. Channel maps of all detected H₂CO lines (other than 3₀₃-2₀₂ and 4₀₄-3₀₃) above 2σ.

TABLE B1
Keplerian Mask Parameters.

Disk	Position Angle* [°]	Inclination Angle* [°]	H ₂ CO and Cont. Mask Radius** ["]	C ¹⁸ O Mask Radius** ["]	Systemic Velocity [km s ⁻¹]
AS 209	6.6	31.2	2.0	2.9	4.600
CI Tau	260.4	41.6	3.4	4.7	5.840
DM Tau	110.3	23.2	4.0	4.1	6.070
DO Tau	113.4	18.5	3.5	2.3	5.900
GM Aur	33.8	37.2	2.7	-	5.550
HD 143006	277.1	24.0	1.7	1.7	7.755
HD 163296	136.9	43.7	4.8	4.1	5.790
IM Lup	305.3	46.2	3.4	3.4	4.400
J1604-2130	194.6	6.2	2.3	2.0	4.600
J1609-1908	103.4	32.1	1.7	-	3.800
J1612-1859	151.9	30.9	0.8	-	4.700
J1614-1906	262.5	40.9	0.8	-	3.800
LkCa 15	33.1	37.8	4.0	4.1	6.300
MWC 480	122	37.4	3.6	5.0	5.100
V4046 Sgr	191.9	29.3	5.6	4.0	2.900

NOTE. — All Keplerian masks were generated assuming the combined thermal and turbulent line width is $\Delta v \sim v_0(r_0/100\text{AU})^q$ (Yen et al. 2016). For all disks, we fixed r_0 and q to be 100 AU and -0.3, respectively. For most disks, we fixed v_0 to be 0.3km s^{-1} . The exceptions were the broader H₂CO 4₁₄-3₁₃ and 4₁₃-3₁₂ lines, for which we fixed v_0 to be 0.6km s^{-1} , as well as the nearly face-on disk J1604-2130, for which we fixed v_0 to be 0.15km s^{-1} . *: The position and inclination angles were estimated using a grid-search algorithm and the fixed broadening parameters. We stress that these estimates are not exact measurements of the disk angles. They are parametric values used only to maximize the masked emission. **: The Keplerian mask radii were set where the azimuthally-averaged emission intensities first reached zero. For disks with multiple H₂CO lines observed, we used the same (largest) mask radius for all lines to maintain a consistent area of H₂CO emission.




FIG. B1.— H₂CO 3₀₃-2₀₂ and 4₀₄-3₀₃ Spectra. H₂CO 3₀₃-2₀₂ and 4₀₄-3₀₃ lines are depicted in light and dark purple, respectively, above and below the horizontal black line. DM Tau appears twice because both lines are observed towards DM Tau.

TABLE C1
Dust Continuum Flux Measurements.

Disk	λ [mm]	Size* [AU]	Sum** [mJy]	Peak Em.** [mJy beam ⁻¹]	rms (σ) [†] [mJy beam ⁻¹]	Beam Size (P.A.)
AS 209	1.0	66.5	297	45	0.18	0.31" x 0.21" (-70.80 deg)
CI Tau	1.3	206.7	161	54	0.16	0.63" x 0.44" (30.36 deg)
DM Tau	1.3	159.5	99	37	0.10	0.60" x 0.44" (35.26 deg)
	1.0	65.2	89	8.7	0.15	0.18" x 0.15" (14.33 deg)
	0.9	159.5	224	91	0.31	0.65" x 0.44" (-28.59 deg)
DO Tau	1.3	98.0	127	102	0.27	0.68" x 0.44" (29.50 deg)
GM Aur	1.0	79.5	255	26	0.13	0.29" x 0.19" (-1.65 deg)
HD 143006	1.3	148.5	53	21	0.12	0.69" x 0.47" (-75.15 deg)
HD 163296	1.0	121.2	956	219	0.87	0.44" x 0.33" (-84.74 deg)
IM Lup	1.0	189.6	237	78	0.35	0.40" x 0.33" (78.16 deg)
J1604-2130	1.3	178.8	69	11	0.14	0.63" x 0.43" (-77.99 deg)
J1609-1908	1.3	95.9	22	18	0.14	0.63" x 0.43" (-76.80 deg)
J1612-1859	1.3	96.6	3.6	3.0	0.13	0.63" x 0.43" (-76.23 deg)
J1614-1906	1.3	100.1	19	17	0.14	0.62" x 0.43" (-76.99 deg)
LkCa 15	1.0	94.8	281	23	0.11	0.32" x 0.26" (-17.58 deg)
MWC 480	1.0	209.3	482	347	0.27	1.05" x 0.84" (-5.72 deg)
	0.9	161.0	848	448	0.44	0.78" x 0.42" (-23.28 deg)
V4046 Sgr	1.0	80.3	542	75	0.18	0.50" x 0.39" (-70.09 deg)

NOTE. — *: The radial extent of the dust continuum is defined as where the continuum emission is 5% of its peak. **: The fluxes and peak fluxes were measured within the same Keplerian mask radii used for H₂CO (Table B1). Errors are small relative to the fluxes. †: The average rms across 1000 1"-by-1" random samples, extracted 3"-7" away from the disk center.

TABLE C2
C¹⁸O Emission Flux Measurements.

Disk	Flux* [mJy × km s ⁻¹]	Peak Flux* [mJy beam ⁻¹ × km s ⁻¹]	Int. Vel. Range [km s ⁻¹]	Chan. Width [km s ⁻¹]	Chan. rms** [mJy beam ⁻¹]	Beam Size (P.A.)
AS 209	429 ± 30	53 ± 6.4	0.9 - 8.1	0.4	8.0	0.53" x 0.50" (37.32°)
CI Tau	549 ± 17	44 ± 3.3	2.6 - 9.0	0.2	5.4	0.67" x 0.46" (31.32°)
DM Tau	998 ± 17	66 ± 3.2	3.6 - 8.6	0.2	5.6	0.63" x 0.47" (36.01°)
DO Tau	210 ± 12	82 ± 3.6	3.2 - 8.8	0.2	5.3	0.72" x 0.46" (30.42°)
HD 143006	135 ± 10	43 ± 3.6	5.4 - 10.2	0.2	5.5	0.70" x 0.48" (-79.98°)
HD 163296	5040 ± 31	355 ± 6.7	-1.4 - 13.0	0.2	4.9	0.61" x 0.52" (62.26°)
IM Lup	1203 ± 14	80 ± 4.1	0.2 - 8.6	0.2	4.9	0.58" x 0.43" (-64.43°)
J1604-2130	1267 ± 7.2	141 ± 1.7	3.7 - 5.5	0.2	5.4	0.65" x 0.45" (-80.59°)
LkCa 15	619 ± 18	53 ± 3.7	2.3 - 10.3	0.2	4.7	0.55" x 0.43" (11.85°)
MWC 480	1648 ± 20	290 ± 11	-1.0 - 11.2	0.2	4.6	0.64" x 0.41" (13.23°)
V4046 Sgr	1184 ± 18	385 ± 5.1	-4.3 - 10.1	0.2	4.5	0.73" x 0.49" (89.68°)

NOTE. — *: The velocity-integrated fluxes (Column 2) were measured within the bounds of the Keplerian masks (Table B1). The peak fluxes (Column 3) are the peaks of the velocity-integrated emission maps; note the difference in unit compared to the velocity-integrated fluxes. The uncertainty in each peak flux is the standard deviation of the peaks across 1000 random samples. Uncertainties do not include absolute flux uncertainties. **: The channel rms was estimated as the standard deviation of 1000 1"-by-1" random samples.




FIG. D1.— **H₂CO Critical Densities as a Function of Gas Temperature.** The critical density n_{crit} is approximately $n_{crit} \approx A_{ul}/(\sigma v)$ (e.g., Condon & Ransom 2016), where A_{ul} is the Einstein coefficient (Table 2), σ is the collisional cross section (assumed here to be $\sim 9.9 \times 10^{-15} \text{cm}^2$ for H₂CO), $v \approx \sqrt{3k_B T_{gas}/m}$ is the average gas velocity, k_B is the Boltzmann constant, T_{gas} is the gas temperature, and m is the mass of the molecule. The critical densities for all H₂CO lines in this survey (plotted above, grouped by upper energy E_u) exceed 10^6cm^{-3} only at temperatures below 40K. Based on the T Tauri protoplanetary disk modeling of Walsh et al. (2010), we expect typical disk densities to be below 10^6cm^{-3} only in the disk’s atmospheric layer, where gas temperatures would be $\gtrsim 50\text{K}$.

REFERENCES

- Aikawa, Y., Momose, M., Thi, W.-F., et al. 2003, PASJ, 55, 11
 Andrews, S. M., Huang, J., Pérez, L. M., et al. 2018, ApJ, 869, L41
 Atkinson, R., Baulch, D. L., Cox, R. A., et al. 2006, Atmospheric Chemistry and Physics, 6, 3625. <https://www.atmos-chem-phys.net/6/3625/2006/>
 Bergin, E., Calvet, N., Sitko, M. L., et al. 2004, ApJ, 614, L133
 Bergner, J. B., Guzmán, V. G., Öberg, K. I., Loomis, R. A., & Pegues, J. 2018, ApJ, 857, 69
 Bergner, J. B., Öberg, K. I., Bergin, E. A., et al. 2019, ApJ, 876, 25
 Blake, G. A., Sutton, E. C., Masson, C. R., & Phillips, T. G. 1987, ApJ, 315, 621
 Boogert, A. C. A., Pontoppidan, K. M., Knez, C., et al. 2008, ApJ, 678, 985
 Carney, M. T., Hogerheijde, M. R., Loomis, R. A., et al. 2017, A&A, 605, A21
 Chapillon, E., Dutrey, A., Guilloteau, S., et al. 2012, ApJ, 756, 58
 Cleeves, L. I. 2016, ApJ, 816, L21
 Cleeves, L. I., Bergin, E. A., Bethell, T. J., et al. 2011, ApJ, 743, L2
 Condon, J. J., & Ransom, S. M. 2016, Spectral Lines (The National Radio Astronomy Observatory)
 Cridland, A. J., Pudritz, R. E., & Alessi, M. 2016, MNRAS, 461, 3274
 Cridland, A. J., Pudritz, R. E., Birnstiel, T., Cleeves, L. I., & Bergin, E. A. 2017, arXiv e-prints, arXiv:1705.02381
 Drozdovskaya, M. N., van Dishoeck, E. F., Rubin, M., Jørgensen, J. K., & Altwegg, K. 2019, MNRAS, 490, 50
 Drozdovskaya, M. N., Walsh, C., Visser, R., Harsono, D., & van Dishoeck, E. F. 2014, MNRAS, 445, 913
 Dutrey, A., Guilloteau, S., & Guélin, M. 1997, A&A, 317, L55
 Endres, C. P., Schlemmer, S., Schilke, P., Stutzki, J., & Müller, H. S. P. 2016, Journal of Molecular Spectroscopy, 327, 95
 Espaillat, C., D’Alessio, P., Hernández, J., et al. 2010, ApJ, 717, 441
 Favre, C., Fedele, D., Semenov, D., et al. 2018, ApJ, 862, L2
 Fayolle, E. C., Balfe, J., Loomis, R., et al. 2016, ApJ, 816, L28
 Flaherty, K. M., Hughes, A. M., Rosenfeld, K. A., et al. 2015, ApJ, 813, 99
 Fockenberg, C., & Preses, J. M. 2002, The Journal of Physical Chemistry A, 106, 2924. <http://dx.doi.org/10.1021/jp0141880>
 Fuchs, G. W., Cuppen, H. M., Ioppolo, S., et al. 2009, A&A, 505, 629
 Fraud, G., Bertin, M., Romanzin, C., et al. 2019, ACS Earth and Space Chemistry
 Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, ArXiv e-prints, arXiv:1804.09365
 Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016a, A&A, 595, A1
 Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2016b, A&A, 595, A2
 Gibb, E. L., Whittet, D. C. B., Boogert, A. C. A., & Tielens, A. G. G. M. 2004, ApJS, 151, 35
 Goldsmith, P. F., & Langer, W. D. 1999, ApJ, 517, 209
 Guilloteau, S., Di Folco, E., Dutrey, A., et al. 2013, A&A, 549, A92
 Guzmán, V. V., Öberg, K. I., Carpenter, J., et al. 2018, ApJ, 864, 170
 Henning, T., & Semenov, D. 2013, Chemical Reviews, 113, 9016
 Herbst, E., & van Dishoeck, E. F. 2009, ARA&A, 47, 427
 Hidaka, H., Watanabe, N., Shiraki, T., Nagaoka, A., & Kouchi, A. 2004, ApJ, 614, 1124
 Hiraoka, K., Ohashi, N., Kihara, Y., et al. 1994, Chemical Physics Letters, 229, 408. <http://www.sciencedirect.com/science/article/pii/0009261494010668>
 Hiraoka, K., Sato, T., Sato, S., et al. 2002, ApJ, 577, 265
 Huang, J., Öberg, K. I., Qi, C., et al. 2017, ApJ, 835, 231
 Hunter, J. D. 2007, Computing In Science & Engineering, 9, 90
 Jensen, E. L. N., & Mathieu, R. D. 1997, AJ, 114, 301
 Jones, E., Oliphant, T., Peterson, P., et al. 2001–, SciPy: Open source scientific tools for Python, . [http://www.scipy.org/](http://www.scipy.org/Kastner, J. H., Qi, C., Dickson-Vandervelde, D. A., et al. 2018, ApJ, 863, 106)
 Lee, J.-E., Lee, S., Baek, G., et al. 2019, Nature Astronomy, 3, 314
 Loomis, R. A., Cleeves, L. I., Öberg, K. I., et al. 2018, ArXiv e-prints, arXiv:1805.01458
 Loomis, R. A., Cleeves, L. I., Öberg, K. I., Guzman, V. V., & Andrews, S. M. 2015, ApJ, 809, L25
 Marsh, K. A., & Mahoney, M. J. 1992, ApJ, 395, L115
 Mathews, G. S., Williams, J. P., Ménard, F., et al. 2012, ApJ, 745, 23
 Mathews, G. S., Klaassen, P. D., Juhász, A., et al. 2013, A&A, 557, A132
 Mordasini, C., Alibert, Y., Benz, W., & Naef, D. 2008, in Astronomical Society of the Pacific Conference Series, Vol. 398, Extreme Solar Systems, ed. D. Fischer, F. A. Rasio, S. E. Thorsett, & A. Wolszczan, 235

- Müller, H. S. P., Schlöder, F., Stutzki, J., & Winnewisser, G. 2005, *Journal of Molecular Structure*, 742, 215
- Mumma, M. J., & Charnley, S. B. 2011, *ARA&A*, 49, 471
- Noble, J. A., Theule, P., Mispelaer, F., et al. 2012, *A&A*, 543, A5
- Öberg, K. I., & Bergin, E. A. 2016, *ApJ*, 831, L19
- Öberg, K. I., Furuya, K., Loomis, R., et al. 2015a, *ApJ*, 810, 112
- Öberg, K. I., Guzmán, V. V., Furuya, K., et al. 2015b, *Nature*, 520, 198
- Öberg, K. I., Qi, C., Fogel, J. K. J., et al. 2010, *ApJ*, 720, 480
- . 2011, *ApJ*, 734, 98
- Öberg, K. I., Guzmán, V. V., Merchantz, C. J., et al. 2017, *ApJ*, 839, 43
- Oliphant, T. 2006–, *NumPy: A guide to NumPy, USA: Trelgol Publishing*, . <http://www.numpy.org/>
- Pinte, C., Ménard, F., Duchêne, G., et al. 2018, *A&A*, 609, A47
- Podio, L., Bacciotti, F., Fedele, D., et al. 2019, *A&A*, 623, L6
- Pontoppidan, K. M., van Dishoeck, E. F., & Dartois, E. 2004, *A&A*, 426, 925
- Qi, C., Öberg, K. I., Andrews, S. M., et al. 2015, *ApJ*, 813, 128
- Qi, C., Öberg, K. I., & Wilner, D. J. 2013, *ApJ*, 765, 34
- Qi, C., Öberg, K. I., Espaillat, C. C., et al. 2019, *ApJ*, 882, 160
- Quast, G. R., Torres, C. A. O., de La Reza, R., da Silva, L., & Mayor, M. 2000, in *IAU Symposium, Vol. 200, IAU Symposium*, 28
- Rosenfeld, K. A., Andrews, S. M., Hughes, A. M., Wilner, D. J., & Qi, C. 2013, *ApJ*, 774, 16
- Rosenfeld, K. A., Andrews, S. M., Wilner, D. J., & Stempels, H. C. 2012, *ApJ*, 759, 119
- Salinas, V. N., Hogerheijde, M. R., Mathews, G. S., et al. 2017, *A&A*, 606, A125
- Schwarz, K. R., Bergin, E. A., Cleeves, L. I., et al. 2016, *ApJ*, 823, 91
- Thi, W.-F., van Zadelhoff, G.-J., & van Dishoeck, E. F. 2004, *A&A*, 425, 955
- van der Marel, N., van Dishoeck, E. F., Bruderer, S., Pérez, L., & Isella, A. 2015, *A&A*, 579, A106
- van der Marel, N., van Dishoeck, E. F., Bruderer, S., & van Kempen, T. A. 2014, *A&A*, 563, A113
- van 't Hoff, M. L. R., Tobin, J. J., Trapman, L., et al. 2018, *ApJ*, 864, L23
- Walsh, C., Millar, T. J., & Nomura, H. 2010, *ApJ*, 722, 1607
- Walsh, C., Millar, T. J., Nomura, H., et al. 2014, *A&A*, 563, A33
- Walsh, C., Loomis, R. A., Öberg, K. I., et al. 2016, *ApJ*, 823, L10
- Watanabe, N., & Kouchi, A. 2002, *ApJ*, 571, L173
- Watanabe, N., Nagaoka, A., Shiraki, T., & Kouchi, A. 2004, *ApJ*, 616, 638
- Yen, H.-W., Koch, P. M., Liu, H. B., et al. 2016, *ApJ*, 832, 204
- Zhang, K., Isella, A., Carpenter, J. M., & Blake, G. A. 2014, *ApJ*, 791, 42