# SPECTRAL ASPECTS OF COMMUTING CONJUGACY CLASS GRAPH OF FINITE GROUPS

#### PARTHAJIT BHOWAL AND RAJAT KANTI NATH\*

ABSTRACT. The commuting conjugacy class graph of a non-abelian group G, denoted by  $\mathcal{CCC}(G)$ , is a simple undirected graph whose vertex set is the set of conjugacy classes of the non-central elements of G and two distinct vertices  $x^G$  and  $y^G$  are adjacent if there exists some elements  $x' \in x^G$  and  $y' \in y^G$  such that x'y' = y'x'. In this paper we compute various spectra and energies of commuting conjugacy class graph of the groups  $D_{2n}, Q_{4m}, U_{(n,m)}, V_{8n}$  and  $SD_{8n}$ . Our computation shows that  $\mathcal{CCC}(G)$  is super integral for these groups. We compare various energies and as a consequence it is observed that  $\mathcal{CCC}(G)$  satisfy E-LE Conjecture of Gutman et al. We also provide negative answer to a question posed by Dutta et al. comparing Laplacian and Signless Laplacian energy. Finally, we conclude this paper by characterizing the above mentioned groups G such that  $\mathcal{CCC}(G)$  is hyperenergetic, L-hyperenergetic or Q-hyperenergetic.

#### 1. Introduction

The commuting graph of a finite non-abelian group G with center Z(G) is a simple undirected graph whose vertex set is  $G \setminus Z(G)$  and two distinct vertices x and y are adjacent if xy = yx. This graph was first considered by Brauer and Fowler [3]. Later on, many mathematicians have extended this graph by considering nilpotent graphs, solvable graphs, commuting conjugacy class graphs etc. The commuting conjugacy class graph of a non-abelian group G, denoted by  $\mathcal{CCC}(G)$ , is a simple undirected graph whose vertex set is the set of conjugacy classes of the non-central elements of G and two distinct vertices  $x^G$  and  $y^G$  are adjacent if there exists some elements  $x' \in x^G$  and  $y' \in y^G$  such that x'y' = y'x'. The notion of commuting conjugacy class graph of groups was introduced by Herzog, Longobardi and Maj [14] in the year 2009. However, in their definition of  $\mathcal{CCC}(G)$ , the vertex set is considered to be the set of all non-identity conjugacy classes of G. In the year 2016, Mohammadian et al. [18] have classified all finite groups such that  $\mathcal{CCC}(G)$  is triangle-free. Recently, in [20], Salahshour and Ashrafi have obtain the structure of  $\mathcal{CCC}(G)$  considering G to be the following groups:

$$D_{2n} = \langle x, y : x^n = y^2 = 1, yxy = x^{-1} \rangle \text{ for } n \geq 3,$$

$$Q_{4m} = \langle x, y : x^{2m} = 1, x^m = y^2, y^{-1}xy = x^{-1} \rangle \text{ for } m \geq 2,$$

$$U_{(n,m)} = \langle x, y : x^{2n} = y^m = 1, x^{-1}yx = y^{-1} \rangle \text{ for } m \geq 2 \text{ and } n \geq 2,$$

$$V_{8n} = \langle x, y : x^{2n} = y^4 = 1, yx = x^{-1}y^{-1}, y^{-1}x = x^{-1}y \rangle \text{ for } n \geq 2,$$

$$SD_{8n} = \langle x, y : x^{4n} = y^2 = 1, yxy = x^{2n-1} \rangle \text{ for } n \geq 2 \text{ and}$$

$$G(p, m, n) = \langle x, y : x^{p^m} = y^{p^n} = [x, y]^p = 1, [x, [x, y]] = [y, [x, y]] = 1 \rangle,$$

where p is any prime,  $m \ge 1$  and  $n \ge 1$ .

In this paper we compute various spectra and energies of commuting conjugacy class graph of the first five groups listed above due to the similar nature of their commuting conjugacy class graphs. In a subsequent paper we shall consider commuting conjugacy class graph of G(p, m, n). Computation of various spectra is helpful to check whether  $\mathcal{CCC}(G)$  is super integral. Recall that a graph  $\mathcal{G}$ 

 $<sup>2010\</sup> Mathematics\ Subject\ Classification.\ 20D99,\ 05C50,\ 15A18,\ 05C25.$ 

Key words and phrases. commuting conjugacy class graph, spectrum, energy, finite group.

<sup>\*</sup>Corresponding author.

is called super integral if it is integral, L-integral and Q-integral. In the year 1974, Harary and Schwenk [13] introduced the concept of integral graphs. Several results on these graphs can be found in [1, 2, 4, 15, 17, 21]. It is observed that  $\mathcal{CCC}(G)$  is super integral for the groups mentioned above. In Section 4, using the energies computed in Section 3, we determine whether the inequalities in [5, Conjecture 1] and [5, Question 1] satisfy for  $\mathcal{CCC}(G)$ . In Section 5, we determine whether  $\mathcal{CCC}(G)$  is hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic or Q-borderenergetic. It is worth mentioning that various spectra and energies of commuting graphs of finite groups have been computed in [6, 7, 19, 8, 5].

## 2. Definitions and useful results

Let  $A(\mathcal{G})$  and  $D(\mathcal{G})$  denote the adjacency matrix and degree matrix of a graph  $\mathcal{G}$  respectively. Then the Laplacian matrix and Signless Laplacian matrix of  $\mathcal{G}$  are given by  $L(\mathcal{G}) = D(\mathcal{G}) - A(\mathcal{G})$  and  $Q(\mathcal{G}) = D(\mathcal{G}) + A(\mathcal{G})$  respectively. We write  $\operatorname{Spec}(\mathcal{G})$ , L-spec( $\mathcal{G}$ ) and Q-spec( $\mathcal{G}$ ) to denote the spectrum, Laplacian spectrum and Signless Laplacian spectrum of  $\mathcal{G}$ . Also,  $\operatorname{Spec}(\mathcal{G}) = \{\alpha_1^{a_1}, \alpha_2^{a_2}, \dots, \alpha_l^{a_l}\}$ , L-spec( $\mathcal{G}$ ) =  $\{\beta_1^{b_1}, \beta_2^{b_2}, \dots, \beta_m^{b_m}\}$  and Q-spec( $\mathcal{G}$ ) =  $\{\gamma_1^{c_1}, \gamma_2^{c_2}, \dots, \gamma_n^{c_n}\}$  where  $\alpha_1, \alpha_2, \dots, \alpha_n$  are the eigenvalues of  $A(\mathcal{G})$  with multiplicities  $a_1, a_2, \dots, a_l; \beta_1, \beta_2, \dots, \beta_m$  are the eigenvalues of  $L(\mathcal{G})$  with multiplicities  $b_1, b_2, \dots, b_m$  and  $\gamma_1, \gamma_2, \dots, \gamma_n$  are the eigenvalues of  $Q(\mathcal{G})$  with multiplicities  $a_1, a_2, \dots, a_l$  is called integral or L-integral or Q-integral according as  $\operatorname{Spec}(\mathcal{G})$  or L-spec( $\mathcal{G}$ ) or Q-spec( $\mathcal{G}$ ) contains only integers. Following theorem is helpful in computing various spectra.

**Theorem 2.1.** If  $\mathcal{G} = l_1 K_{m_1} \sqcup l_2 K_{m_2}$ , where  $l_i K_{m_i}$  denotes the disjoint union of  $l_i$  copies of the complete graph  $K_{m_i}$  on  $m_i$  vertices for i = 1, 2, then

$$\begin{aligned} \operatorname{Spec}(\mathcal{G}) &= \left\{ (-1)^{\sum_{i=1}^{2} l_i(m_i - 1)}, (m_1 - 1)^{l_1}, (m_2 - 1)^{l_2} \right\} \\ \operatorname{L-spec}(\mathcal{G}) &= \left\{ 0^{l_1 + l_2}, m_1^{l_1(m_1 - 1)}, m_2^{l_2(m_2 - 1)} \right\} \ and \\ \operatorname{Q-spec}(\mathcal{G}) &= \left\{ (2m_1 - 2)^{l_1}, (m_1 - 2)^{l_1(m_1 - 1)}, (2m_2 - 2)^{l_2}, (m_2 - 2)^{l_2(m_2 - 1)} \right\}. \end{aligned}$$

Depending on the various spectra of a graph, there are various energies called *energy*, Laplacian energy and Signless Laplacian energy denoted by  $E(\mathcal{G})$ ,  $LE(\mathcal{G})$  and  $LE^+(\mathcal{G})$  respectively. These energies are defined as follows:

(2.1) 
$$E(\mathcal{G}) = \sum_{\lambda \in \text{Spec}(\mathcal{G})} |\lambda|,$$

(2.2) 
$$LE(\mathcal{G}) = \sum_{\mu \in L\text{-spec}(\mathcal{G})} \left| \mu - \frac{2|e(\Gamma)|}{|V(\Gamma)|} \right|,$$

(2.3) 
$$LE^{+}(\mathcal{G}) = \sum_{\nu \in Q\text{-spec}(\mathcal{G})} \left| \nu - \frac{2|e(\Gamma)|}{|V(\mathcal{G})|} \right|,$$

where  $V(\mathcal{G})$  and  $e(\mathcal{G})$  denote the set of vertices and edges of  $\Gamma$ .

In 2008, Gutman et al. [12] posed the following conjecture comparing  $E(\mathcal{G})$  and  $LE(\mathcal{G})$ .

Conjecture 2.2. (E-LE Conjecture)  $E(\mathcal{G}) \leq LE(\mathcal{G})$  for any graph  $\mathcal{G}$ .

However, in the same year, Stevanović et al. [22] disproved the above conjecture. In 2009, Liu and Lin [16] also disproved Conjecture 2.2 by providing some counter examples. Following Gutman et al. [12], recently Dutta et al. have posed the following question in [5] comparing Laplacian and singless Laplacian energies of graphs.

Question 2.3. Is  $LE(\mathcal{G}) \leq LE^+(\mathcal{G})$  for all graphs  $\mathcal{G}$ ?

### 3. Various spectra and energies

In this section we compute various spectra and energies of commuting conjugacy class graphs of the groups mentioned in the introduction.

Theorem 3.1. If  $G = D_{2n}$  then

*Proof.* We shall prove the result by considering the following cases.

Case 1. n is odd.

By [20, Proposition 2.1] we have  $\mathcal{CCC}(G) = K_1 \sqcup K_{\frac{n-1}{2}}$ . Therefore, by Theorem 2.1, it follows that

$$\begin{split} \operatorname{Spec}(\mathcal{CCC}(G)) &= \left\{ (-1)^{\frac{n-3}{2}}, 0^1, \left(\frac{n-3}{2}\right)^1 \right\}, \quad \operatorname{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^2, \left(\frac{n-1}{2}\right)^{\frac{n-3}{2}} \right\} \\ \text{and } \operatorname{Q-spec}(\mathcal{CCC}(G)) &= \left\{ 0^1, (n-3)^1, \left(\frac{n-5}{2}\right)^{\frac{n-3}{2}} \right\}. \end{split}$$

Hence, by (2.1), we get

$$E(\mathcal{CCC}(G)) = \frac{n-3}{2} + \frac{n-3}{2} = n-3.$$

We have  $|V(\mathcal{CCC}(G))| = \frac{n+1}{2}$  and  $|e(\mathcal{CCC}(G))| = \frac{(n-1)(n-3)}{8}$ . Therefore,  $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(n-1)(n-3)}{2(n+1)}$ . Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(n-1)(n-3)}{2(n+1)} \right| = \frac{(n-1)(n-3)}{2(n+1)} \quad \text{and} \quad \left| \frac{n-1}{2} - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| \frac{n-1}{2} - \frac{(n-1)(n-3)}{2(n+1)} \right| = \frac{2(n-1)}{n+1}.$$

Now, by (2.2), we have

$$LE(\mathcal{CCC}(G)) = 2 \times \frac{(n-1)(n-3)}{2(n+1)} + \frac{n-3}{2} \times \frac{2(n-1)}{n+1} = \frac{2(n-1)(n-3)}{n+1}.$$

Again,

$$\left| n - 3 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| n - 3 - \frac{(n-1)(n-3)}{2(n+1)} \right| = \frac{(n-3)(n+3)}{2(n+1)} \quad \text{and} \quad \left| \frac{n-5}{2} - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| \frac{n-5}{2} - \frac{(n-1)(n-3)}{2(n+1)} \right| = \left| \frac{-4}{n+1} \right| = \frac{4}{n+1}.$$

By (2.3), we have

$$LE^{+}(\mathcal{CCC}(G)) = \frac{(n-1)(n-3)}{2(n+1)} + \frac{(n-3)(n+3)}{2(n+1)} + \frac{n-3}{2} \times \frac{4}{n+1} = \frac{(n-3)(n+3)}{n+1}.$$

Case 2. n is even.

Consider the following subcases.

Subcase 2.1  $\frac{n}{2}$  is even.

By [20, Proposition 2.1] we have  $\mathcal{CCC}(G) = 2K_1 \sqcup K_{\frac{n}{2}-1}$ . Therefore, by Theorem 2.1, it follows that

$$\begin{split} \operatorname{Spec}(\mathcal{CCC}(G)) &= \left\{ (-1)^{\frac{n}{2}-2}, 0^2, \left(\frac{n}{2}-2\right)^1 \right\}, \quad \operatorname{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^3, \left(\frac{n}{2}-1\right)^{\frac{n}{2}-2} \right\} \\ \text{and } \operatorname{Q-spec}(\mathcal{CCC}(G)) &= \left\{ 0^2, (n-4)^1, \left(\frac{n}{2}-3\right)^{\frac{n}{2}-2} \right\}. \end{split}$$

Hence, by (2.1), we get

$$E(\mathcal{CCC}(G)) = \frac{n}{2} - 2 + \frac{n}{2} - 2 = n - 4.$$

We have  $|V(\mathcal{CCC}(G))| = \frac{n}{2} + 1$  and  $|e(\mathcal{CCC}(G))| = \frac{(n-2)(n-4)}{8}$ . So,  $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(n-2)(n-4)}{2(n+2)}$ . Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(n-2)(n-4)}{2(n+2)} \right| = \frac{(n-2)(n-4)}{2(n+2)} \quad \text{and} \quad \left| \frac{n}{2} - 1 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| \frac{n}{2} - 1 - \frac{(n-2)(n-4)}{2(n+2)} \right| = \frac{3(n-2)}{n+2}.$$

Now, by (2.2), we have

$$LE(\mathcal{CCC}(G)) = 3 \times \frac{(n-2)(n-4)}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{3(n-2)}{n+2} = \frac{3(n-2)(n-4)}{n+2}.$$

Again,

$$\left| n - 4 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| n - 4 - \frac{(n-2)(n-4)}{2(n+2)} \right| = \frac{(n-4)(n+6)}{2(n+2)} \quad \text{and} \quad$$

$$\left| \frac{n}{2} - 3 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| \frac{n}{2} - 3 - \frac{(n-2)(n-4)}{2(n+2)} \right| = \left| \frac{n-10}{n+2} \right| = \begin{cases} \frac{-n+10}{n+2}, & \text{if } n = 4, 8 \\ \frac{n-10}{n+2}, & \text{if } n \geq 12. \end{cases}$$

By (2.3), we have

$$LE^{+}(\mathcal{CCC}(G)) = 2 \times \frac{(n-2)(n-4)}{2(n+2)} + \frac{(n-4)(n+6)}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{-n+10}{n+2} = \frac{(n-4)(n+6)}{n+2},$$

if n = 4, 8. If  $n \ge 12$  then

$$LE^{+}(\mathcal{CCC}(G)) = 2 \times \frac{(n-2)(n-4)}{2(n+2)} + \frac{(n-4)(n+6)}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{n-10}{n+2} = \frac{2(n-2)(n-4)}{n+2}.$$

Subcase 2.2  $\frac{n}{2}$  is odd.

By [20, Proposition 2.1] we have  $\mathcal{CCC}(G) = K_2 \sqcup K_{\frac{n}{2}-1}$ . Therefore, by Theorem 2.1, it follows that

$$\begin{split} \operatorname{Spec}(\mathcal{CCC}(G)) &= \left\{ (-1)^{\frac{n}{2}-1}, 1^1, \left(\frac{n}{2}-2\right)^1 \right\}, \quad \operatorname{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^2, 2^1, \left(\frac{n}{2}-1\right)^{\frac{n}{2}-2} \right\} \\ \text{and } \operatorname{Q-spec}(\mathcal{CCC}(G)) &= \left\{ 2^1, 0^1, (n-4)^1, \left(\frac{n}{2}-3\right)^{\frac{n}{2}-2} \right\}. \end{split}$$

Hence, by (2.1), we get

$$E(\mathcal{CCC}(G)) = \frac{n}{2} - 1 + 1 + \frac{n}{2} - 2 = n - 2.$$

We have  $|V(\mathcal{CCC}(G))| = \frac{n}{2} + 1$  and  $|e(\mathcal{CCC}(G))| = \frac{(n-2)(n-4)+8}{8}$ . Therefore,  $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(n-2)(n-4)+8}{2(n+2)}$ . Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(n-2)(n-4)+8}{2(n+2)} \right| = \frac{(n-2)(n-4)+8}{2(n+2)},$$

$$\left| 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2 - \frac{(n-2)(n-4)+8}{2(n+2)} \right| = \left| \frac{-n^2+10n-8}{2(n+2)} \right| = \begin{cases} 1, & \text{if } n=6\\ \frac{n^2-10n+8}{2(n+2)}, & \text{if } n \ge 10 \end{cases}$$
and 
$$\left| \frac{n}{2} - 1 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| \frac{n}{2} - 1 - \frac{(n-2)(n-4)+8}{2(n+2)} \right| = \frac{3n-10}{n+2}.$$

Now, by (2.2), we have

$$LE(\mathcal{CCC}(G)) = 2 \times \frac{(n-2)(n-4) + 8}{2(n+2)} + 1 + \left(\frac{n}{2} - 2\right) \times \frac{3n - 10}{n+2} = 4,$$

if n = 6. If n > 10 then

$$LE(\mathcal{CCC}(G)) = 2 \times \frac{(n-2)(n-4)+8}{2(n+2)} + \frac{n^2 - 10n + 8}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{3n - 10}{n+2}$$
$$= \frac{3n^2 - 22n - 40}{n+2} = \frac{(n-4)(3n-10)}{n+2}.$$

Again,

$$\left| n - 4 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| n - 4 - \frac{(n-2)(n-4) + 8}{2(n+2)} \right| = \frac{n^2 + 2n - 32}{2(n+2)} \quad \text{and}$$

$$\left| \frac{n}{2} - 3 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| \frac{n}{2} - 3 - \frac{(n-2)(n-4) + 8}{2(n+2)} \right| = \left| \frac{n-14}{n+2} \right| = \begin{cases} \frac{-n+14}{n+2}, & \text{if } n = 6, 10\\ \frac{n-14}{n+2}, & \text{if } n \geq 14. \end{cases}$$

By (2.3), we have

$$LE^{+}(\mathcal{CCC}(G)) = 1 + \frac{(n-2)(n-4) + 8}{2(n+2)} + \frac{n^2 + 2n - 32}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{-n + 14}{n+2} = 4,$$

if n = 6. If n = 10 then

$$LE^{+}(\mathcal{CCC}(G)) = \frac{n^2 - 10n + 8}{2(n+2)} + \frac{(n-2)(n-4) + 8}{2(n+2)} + \frac{n^2 + 2n - 32}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{-n + 14}{n+2} = \frac{22}{3}.$$

If  $n \geq 14$  then

$$LE^{+}(\mathcal{CCC}(G)) = \frac{n^{2} - 10n + 8}{2(n+2)} + \frac{(n-2)(n-4) + 8}{2(n+2)} + \frac{n^{2} + 2n - 32}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{n - 14}{n+2}$$
$$= \frac{2(n-2)(n-6)}{n+2}.$$

This completes the proof.

**Theorem 3.2.** If  $G = Q_{4m}$  then

$$\text{heorem 3.2. } If G = Q_{4m} \ then$$

$$\text{(a) } \operatorname{Spec}(\mathcal{CCC}(G)) = \begin{cases} \left\{ (-1)^{m-1}, 1^1, (m-2)^1 \right\}, & \text{if } m \text{ is odd} \\ \left\{ (-1)^{m-2}, 0^2, (m-2)^1 \right\}, & \text{if } m \text{ is even} \end{cases}$$

$$and \ E(\mathcal{CCC}(G)) = \begin{cases} 2m-2, & \text{if } m \text{ is odd} \\ 2m-4, & \text{if } m \text{ is even}. \end{cases}$$

$$\text{(b) } \operatorname{L-spec}(\mathcal{CCC}(G)) = \begin{cases} \left\{ 0^2, 2^1, (m-1)^{m-2} \right\}, & \text{if } m \text{ is odd} \\ \left\{ 0^3, (m-1)^{m-2} \right\}, & \text{if } m \text{ is even} \end{cases}$$

$$and \ LE(\mathcal{CCC}(G)) = \begin{cases} 4, & \text{if } m = 3 \\ \frac{2(m-2)(3m-5)}{m+1}, & \text{if } m \text{ is odd and } m \geq 5 \\ \frac{6(m-1)(m-2)}{m+1}, & \text{if } m \text{ is even.} \end{cases}$$

$$\text{(c) } \operatorname{Q-spec}(\mathcal{CCC}(G)) = \begin{cases} \left\{ 2^1, 0^1, (2m-4)^1, (m-3)^{m-2} \right\}, & \text{if } m \text{ is odd} \\ \left\{ 0^2, (2m-4)^1, (m-3)^{m-2} \right\}, & \text{if } m \text{ is even.} \end{cases}$$

$$and \ LE^+(\mathcal{CCC}(G)) = \begin{cases} 4, & \text{if } m = 3 \\ \frac{22}{3}, & \text{if } m = 5 \\ \frac{4(m-1)(m-3)}{m+1}, & \text{if } m \text{ is odd and } m \geq 7 \\ \frac{2(m-2)(m+3)}{m+1}, & \text{if } m \text{ is even and } m \geq 6. \end{cases}$$

*Proof.* We shall prove the result by considering the following cases. Case 1. m is odd.

By [20, Proposition 2.2] we have  $\mathcal{CCC}(G) = K_2 \sqcup K_{m-1}$ . Therefore, by Theorem 2.1, it follows that

$$\begin{aligned} &\operatorname{Spec}(\mathcal{CCC}(G)) = \left\{ (-1)^{m-1}, 1^1, (m-2)^1 \right\}, \quad \operatorname{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^2, 2^1, (m-1)^{m-2} \right\} \\ &\operatorname{and} \ \operatorname{Q-spec}(\mathcal{CCC}(G)) = \left\{ 2^1, 0^1, (2m-4)^1, (m-3)^{m-2} \right\}. \end{aligned}$$

Hence, by (2.1), we get

$$E(\mathcal{CCC}(G)) = m - 1 + 1 + m - 2 = 2m - 2.$$

We have  $|V(\mathcal{CCC}(G))| = m+1$  and  $|e(\mathcal{CCC}(G))| = \frac{(m-1)(m-2)+2}{2}$ . Therefore,  $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(m-1)(m-2)+2}{2}$ .

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(m-1)(m-2) + 2}{m+1} \right| = \frac{(m-1)(m-2) + 2}{m+1},$$
 
$$\left| 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2 - \frac{(m-1)(m-2) + 2}{m+1} \right| = \left| \frac{-m^2 + 5m - 2}{m+1} \right| = \begin{cases} 1, & \text{if } m = 3 \\ \frac{m^2 - 5m + 2}{m+1}, & \text{if } m \geq 5 \end{cases}$$
 and 
$$\left| m - 1 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| m - 1 - \frac{(m-1)(m-2) + 2}{m+1} \right| = \frac{3m - 5}{m+1}.$$

Now, by (2.2), we have

$$LE(\mathcal{CCC}(G)) = 2 \times \frac{(m-1)(m-2)+2}{m+1} + 1 + (m-2) \times \frac{3m-5}{m+1} = 4,$$

if m=3. If  $m\geq 5$  then

$$LE(\mathcal{CCC}(G)) = 2 \times \frac{(m-1)(m-2)+2}{m+1} + \frac{m^2-5m+2}{m+1} + (m-2) \times \frac{3m-5}{m+1} = \frac{2(m-2)(3m-5)}{m+1}.$$

Again,

$$\left| 2m - 4 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2m - 4 - \frac{(m-1)(m-2) + 2}{m+1} \right| = \frac{m^2 + m - 8}{m+1} \quad \text{and} \quad$$

$$\left| m - 3 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| m - 3 - \frac{(m-1)(m-2) + 2}{m+1} \right| = \left| \frac{m-7}{m+1} \right| = \begin{cases} \frac{-m+7}{m+1}, & \text{if } m = 3, 5 \\ \frac{m-7}{m+1}, & \text{if } m \geq 7. \end{cases}$$

By (2.3), we have

$$LE^{+}(\mathcal{CCC}(G)) = 1 + \frac{(m-1)(m-2)+2}{m+1} + \frac{m^2+m-8}{m+1} + (m-2) \times \frac{-m+7}{m+1} = 4,$$

if m=3. If m=5 then

$$LE^{+}(\mathcal{CCC}(G)) = \frac{m^{2} - 5m + 2}{m+1} + \frac{(m-1)(m-2) + 2}{m+1} + \frac{m^{2} + m - 8}{m+1} + (m-2) \times \frac{-m+7}{m+1} = \frac{22}{3}.$$

If m > 7 then

$$LE^{+}(\mathcal{CCC}(G)) = \frac{m^{2} - 5m + 2}{m+1} + \frac{(m-1)(m-2) + 2}{m+1} + \frac{m^{2} + m - 8}{m+1} + (m-2) \times \frac{m-7}{m+1}$$
$$= \frac{4(m-1)(m-3)}{m+1}.$$

Case 2. m is even.

By [20, Proposition 2.2] we have  $\mathcal{CCC}(G) = 2K_1 \sqcup K_{m-1}$ . Therefore, by Theorem 2.1, it follows that

$$\begin{split} &\operatorname{Spec}(\mathcal{CCC}(G)) = \left\{ (-1)^{m-2}, 0^2, (m-2)^1 \right\}, \quad \operatorname{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^3, (m-1)^{m-2} \right\} \\ &\operatorname{and} \ \operatorname{Q-spec}(\mathcal{CCC}(G)) = \left\{ 0^2, (2m-4)^1, (m-3)^{m-2} \right\}. \end{split}$$

Hence, by (2.1), we get

$$E(CCC(G)) = m - 2 + m - 2 = 2m - 4.$$

We have  $|V(\mathcal{CCC}(G))| = m+1$  and  $|e(\mathcal{CCC}(G))| = \frac{(m-1)(m-2)}{2}$ . Therefore,  $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(m-1)(m-2)}{m+1}$ . Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(m-1)(m-2)}{m+1} \right| = \frac{(m-1)(m-2)}{m+1} \quad \text{and} \quad \left| m - 1 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| m - 1 - \frac{(m-1)(m-2)}{m+1} \right| = \frac{3(m-1)}{m+1}.$$

Now, by (2.2), we have

$$LE(\mathcal{CCC}(G)) = 3 \times \frac{(m-1)(m-2)}{m+1} + (m-2) \times \frac{3(m-1)}{m+1} = \frac{6(m-1)(m-2)}{m+1}.$$

Again,

$$\left|2m-4-\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|}\right|=\left|2m-4-\frac{(m-1)(m-2)}{m+1}\right|=\frac{(m-2)(m+3)}{m+1}\quad\text{ and }\quad$$

$$\left| m - 3 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| m - 3 - \frac{(m-1)(m-2)}{m+1} \right| = \left| \frac{m-5}{m+1} \right| = \begin{cases} \frac{-m+5}{m+1}, & \text{if } m = 2, 4\\ \frac{m-5}{m+1}, & \text{if } m \geq 6. \end{cases}$$

By (2.3), we have

$$LE^{+}(\mathcal{CCC}(G)) = 2 \times \frac{(m-1)(m-2)}{m+1} + \frac{(m-2)(m+3)}{m+1} + (m-2) \times \frac{-m+5}{m+1}$$
$$= \frac{2(m-2)(m+3)}{m+1},$$

if m=2,4. If  $m\geq 6$  then

$$LE^{+}(\mathcal{CCC}(G)) = 2 \times \frac{(m-1)(m-2)}{m+1} + \frac{(m-2)(m+3)}{m+1} + (m-2) \times \frac{m-5}{m+1}$$
$$= \frac{4(m-1)(m-2)}{m+1}.$$

This completes the proof.

**Theorem 3.3.** If  $G = U_{(n,m)}$  then

$$\text{ heorem 3.3. } \textit{If } G = U_{(n,m)} \textit{ then }$$

$$\text{ (a) } \operatorname{Spec}(\mathcal{CCC}(G)) = \begin{cases} \left\{ (-1)^{\frac{n(m+1)-4}{2}}, \left(\frac{n(m-1)-2}{2}\right)^1, (n-1)^1 \right\}, & \textit{if } m \textit{ is odd and } n \geq 2 \\ \left\{ (-1)^{\frac{n(m+2)-6}{2}}, \left(\frac{n(m-2)-2}{2}\right)^1, (n-1)^2 \right\}, & \textit{if } m \textit{ is even and } n \geq 2 \end{cases}$$

$$and \textit{ E(CCC}(G)) = \begin{cases} n(m+1)-4, & \textit{if } m \textit{ is odd and } n \geq 2 \\ 4(n-1), & \textit{if } m = 2 \textit{ and } n \geq 2 \\ n(m+2)-6, & \textit{if } m \textit{ is even, } m \geq 4 \textit{ and } n \geq 2. \end{cases}$$

$$\left\{ \begin{cases} 0^2, \left(\frac{n(m-1)}{2}\right)^{\frac{n(m-1)-2}{2}}, n^{n-1} \right\}, & \textit{if } m \textit{ is odd and } n \geq 2 \\ 0^3, \left(\frac{n(m-2)}{2}\right)^{\frac{n(m-2)-2}{2}}, n^{2n-2} \right\}, & \textit{if } m \textit{ is even and } n \geq 2 \end{cases}$$

$$and \textit{ LE(CCC}(G)) = \begin{cases} 4(n-1), & \textit{if } m = 3 \textit{ and } n \geq 2 \\ \frac{2(2n-1)(n+3)}{m+1}, & \textit{if } m = 5 \textit{ and } n \geq 2 \\ 4(n-1), & \textit{if } m = 5 \textit{ and } n \geq 2 \end{cases}$$

$$and \textit{ ned } 2 \end{cases}$$

$$and \textit{ lending } 3 \end{cases}$$

$$and \textit{ lending } 2 \end{cases}$$

$$and \textit{ lending } 3 \end{cases}$$

$$and \textit{ lending } 2 \end{cases}$$

$$and \textit{ lending } 3 \end{cases}$$

$$and \textit{ lending } 2 \end{cases}$$

$$and \textit{ lending } 3 \end{cases}$$

$$and \textit{ lending } 2 \end{cases}$$

$$and \textit{ lending } 3 \end{cases}$$

$$and \textit{ lending } 3 \end{cases}$$

$$and \textit{ lending } 4 \end{cases}$$

$$and \textit{ lending } 2 \end{cases}$$

$$and \textit{ lending } 3 \end{cases}$$

$$and \textit{ lending } 4 \end{cases}$$

$$and \textit{ lending } 2 \end{cases}$$

$$and \textit{ lending } 3 \end{cases}$$

$$and \textit{ lending } 4 \end{cases}$$

$$and \textit{ lending } 2 \end{cases}$$

$$and \textit{ lending } 3 \end{cases}$$

$$and \textit{ lending } 4 \end{cases}$$

$$and \textit{ lending } 2 \end{cases}$$

$$and \textit{ lending } 4 \end{cases}$$

$$and \textit{ lending } 2 \end{cases}$$

$$and \textit{ lending } 3 \end{cases}$$

$$and \textit{ lending } 4 \end{cases}$$

$$and \textit{ lending } 2 \end{cases}$$

$$and \textit{ lending } 3 \end{cases}$$

$$and \textit{ lending } 4 \end{cases}$$

$$an$$

$$and \ LE^{+}(\mathcal{CCC}(G)) = \begin{cases} 4(n-1), & \text{if } m=3 \ and \ n \geq 2 \\ \frac{22}{3}, & \text{if } m=5 \ and \ n=2 \\ \frac{2(2n+3)(n-1)}{3}, & \text{if } m=5 \ and \ n \geq 3 \\ \frac{n^2(m-1)(m-3)}{m+1}, & \text{if } m \ is \ odd, \ m \geq 7 \ and \ n \geq 2 \\ 4(n-1), & \text{if } m=2 \ and \ n \geq 2 \\ 6(n-1), & \text{if } m=4 \ and \ n \geq 2 \\ 2(n+2)(n-1), & \text{if } m=6 \ and \ n \geq 2 \\ \frac{2n^2(m-2)(m-4)}{m+2}, & \text{if } m \ is \ even, \ m \geq 8 \ and \ n \geq 2. \end{cases}$$

*Proof.* We shall prove the result by considering the following cases. Case 1. m is odd.

By [20, Proposition 2.3] we have  $\mathcal{CCC}(G) = K_{\frac{n(m-1)}{2}} \sqcup K_n$ . Therefore, by Theorem 2.1, it follows that

$$\begin{split} \operatorname{Spec}(\mathcal{CCC}(G)) &= \left\{ (-1)^{\frac{n(m+1)-4}{2}}, \left(\frac{n(m-1)-2}{2}\right)^1, (n-1)^1 \right\}, \\ \operatorname{L-spec}(\mathcal{CCC}(G)) &= \left\{ 0^2, \left(\frac{n(m-1)}{2}\right)^{\frac{n(m-1)-2}{2}}, n^{n-1} \right\} \\ \text{and Q-spec}(\mathcal{CCC}(G)) &= \left\{ (n(m-1)-2)^1, \left(\frac{n(m-1)-4}{2}\right)^{\frac{n(m-1)-2}{2}}, \quad (2n-2)^1, (n-2)^{n-1} \right\}. \end{split}$$

Hence, by (2.1), we get

$$E(\mathcal{CCC}(G)) = \frac{n(m+1)-4}{2} + \frac{n(m-1)-2}{2} + n - 1 = n(m+1) - 4.$$

We have  $|V(\mathcal{CCC}(G))| = \frac{n(m+1)}{2}$  and  $|e(\mathcal{CCC}(G))| = \frac{n^2(m-1)^2 - 2n(m-2n+1)}{8}$ . Therefore,  $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)}$ . Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} \right| = \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)},$$

since  $n(m-1)^2 - 2(m-2n+1) = m^2n - 2m(n+1) + 5n - 2 > 0$ ;

$$\left| \frac{n(m-1)}{2} - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| \frac{n(m-1)}{2} - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} \right| = \frac{n(m-3) + m + 1}{m+1} \text{ and }$$

$$\left|n - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|}\right| = \left|n - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)}\right| = \left|\frac{-f_1(m,n)}{2(m+1)}\right|,$$

where  $f_1(m,n) = n(m^2+3) - (4mn+2m+2)$ . For m = 3 and  $n \ge 2$  we have  $f_1(3,n) = -8$ . For m = 5 and  $n \ge 2$  we have  $f_1(5,n) = 8n - 12 > 0$ . For  $m \ge 7$  and  $n \ge 2$  we have  $m^2 + 3 > m^2 > 4m + 2m + 2$ . Therefore,  $n(m^2 + 3) > 4mn + (2m + 2)n > 4mn + 2m + 2$  and so  $f_1(m,n) > 0$ . Hence,

$$\left|\frac{-f_1(m,n)}{2(m+1)}\right| = \begin{cases} 1, & \text{if } m = 3 \text{ and } n \ge 2\\ \frac{2n-3}{3}, & \text{if } m = 5 \text{ and } n \ge 2\\ \frac{n(m^2+3)-(4mn+2m+2)}{2(m+1)}, & \text{if } m \ge 7 \text{ and } n \ge 2. \end{cases}$$

Now, by (2.2), we have

$$LE(\mathcal{CCC}(G)) = 2 \times \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} + \frac{n(m-1) - 2}{2} \times \frac{n(m-3) + m + 1}{m+1} + (n-1) \times 1$$
$$= 4(n-1),$$

if m=3 and  $n\geq 2$ . If m=5 and  $n\geq 2$  then

$$LE(\mathcal{CCC}(G)) = 2 \times \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} + \frac{n(m-1) - 2}{2} \times \frac{n(m-3) + m + 1}{m+1} + (n-1) \times \frac{2n-3}{3}$$
$$= \frac{2(2n-1)(n+3)}{2}.$$

If  $m \geq 7$  and  $n \geq 2$  then

$$LE(\mathcal{CCC}(G)) = 2 \times \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} + \frac{n(m-1) - 2}{2} \times \frac{n(m-3) + m + 1}{m+1} + (n-1) \times \frac{n(m^2+3) - (4mn + 2m + 2)}{2(m+1)}$$
$$= \frac{m^2n^2 - 4mn^2 + m^2n + 3n^2 - 2mn - 2m + 5n - 2}{m+1}.$$

Again,

$$\begin{vmatrix} n(m-1) - 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \end{vmatrix} = \begin{vmatrix} n(m-1) - 2 - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} \end{vmatrix}$$
$$= \begin{vmatrix} \frac{n(m-1)(m+3) - 2(m+2n+1)}{2(m+1)} \end{vmatrix}$$
$$= \frac{n(m-1)(m+3) - 2(m+2n+1)}{2(m+1)},$$

since  $n(m-1)(m+3) - 2(m+2n+1) = n(m^2-4) - 2 + n(m-3) + m(n-2) > 0$ ;

$$\left| \frac{n(m-1)-4}{2} - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| \frac{n(m-1)-4}{2} - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} \right| = \left| \frac{f_2(m,n)}{2(m+1)} \right|,$$

where  $f_2(m,n) = n(m-6) - 2 + m(n-2)$ . Clearly, for  $m \ge 7$  and  $n \ge 2$  we have  $f_2(m,n) \ge 0$ . For m = 3 and  $n \ge 2$  we have  $f_2(3,n) = -8$ . Also for m = 5 and  $n \ge 2$  we have  $f_2(5,n) = 4n - 12$ . Therefore,  $f_2(5,2) = -4$  and  $f_2(5,n) \ge 0$  for  $n \ge 3$ . Hence,

$$\left| \frac{f_2(m,n)}{2(m+1)} \right| = \begin{cases} 1, & \text{if } m = 3 \text{ and } n \ge 2\\ \frac{1}{3}, & \text{if } m = 5 \text{ and } n = 2\\ \frac{n-3}{3}, & \text{if } m = 5 \text{ and } n \ge 3\\ \frac{n(m-3)-m-1}{m+1}, & \text{if } m \ge 7 \text{ and } n \ge 2. \end{cases}$$

$$\left| 2n - 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n - 2 - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} \right| = \left| -\frac{f_3(m,n)}{2(m+1)} \right|,$$

where  $f_3(m,n) = mn(m-6) + 2m + n + 2$ . Clearly,  $f_3(m,n) > 0$  if  $m \ge 7$  and  $n \ge 2$ . For m = 3 and  $n \ge 2$  we have  $f_3(3,n) = -8n + 8 < 0$ . For m = 5 and  $n \ge 2$  we have  $f_3(5,n) = -4n + 12$ .

Therefore,  $f_3(5,2) = 4$  and  $f_3(5,n) \le 0$  if  $n \ge 3$ . Hence,

$$\left| -\frac{f_3(m,n)}{2(m+1)} \right| = \begin{cases} n-1, & \text{if } m = 3 \text{ and } n \ge 2\\ \frac{1}{3}, & \text{if } m = 5 \text{ and } n = 2\\ \frac{n-3}{3}, & \text{if } m = 5 \text{ and } n \ge 3\\ \frac{mn(m-6)+2m+n+2}{2(m+1)}, & \text{if } m \ge 7 \text{ and } n \ge 2. \end{cases}$$

$$\left| n - 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| n - 2 - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} \right| = \left| -\frac{f_4(m,n)}{2(m+1)} \right|,$$

where  $f_4(m, n) = mn(m-2) + 2 - (m(n-2) + n(m-3))$ . For m = 3 and  $n \ge 2$  we have  $f_4(3, n) = 8$ . Also, for  $m \ge 5$  and  $n \ge 2$  we have

$$mn(m-2) - 2mn + 2 = mn(m-4) + 2 > -2m - 3n.$$

Therefore,

$$mn(m-2) + 2 > 2mn - 2m - 3n = m(n-2) + n(m-3)$$

and so  $f_4(m,n) > 0$  for  $m \ge 5$  and  $n \ge 2$ . Hence,

$$\left| -\frac{f_4(m,n)}{2(m+1)} \right| = \frac{f_4(m,n)}{2(m+1)} = \frac{mn(m-2) + 2 - m(n-2) - n(m-3)}{2(m+1)}.$$

By (2.3), we have

$$LE^{+}(\mathcal{CCC}(G)) = \frac{n(m-1)(m+3) - 2(m+2n+1)}{2(m+1)} + \frac{n(m-1) - 2}{2} \times 1 + (n-1)$$
$$+ (n-1) \times \frac{mn(m-2) + 2 - m(n-2) - n(m-3)}{2(m+1)}$$
$$= 4(n-1),$$

if m=3 and  $n\geq 2$ . If m=5 and n=2 then

$$LE^{+}(\mathcal{CCC}(G)) = \frac{n(m-1)(m+3) - 2(m+2n+1)}{2(m+1)} + \frac{n(m-1) - 2}{2} \times \frac{1}{3} + \frac{1}{3} + (n-1) \times \frac{mn(m-2) + 2 - m(n-2) - n(m-3)}{2(m+1)}$$
$$= \frac{22}{3}.$$

If m = 5 and  $n \ge 3$  then

$$LE^{+}(\mathcal{CCC}(G)) = \frac{n(m-1)(m+3) - 2(m+2n+1)}{2(m+1)} + \frac{n(m-1) - 2}{2} \times \frac{n-3}{3} + \frac{n-3}{3} +$$

If  $m \geq 7$  and  $n \geq 2$  then

$$LE^{+}(\mathcal{CCC}(G)) = \frac{n(m-1)(m+3) - 2(m+2n+1)}{2(m+1)} + \frac{n(m-1) - 2}{2} \times \frac{n(m-3) - m - 1}{m+1} + \frac{mn(m-6) + 2m + n + 2}{2(m+1)} + (n-1) \times \frac{mn(m-2) + 2 - m(n-2) - n(m-3)}{2(m+1)} = \frac{n^{2}(m-1)(m-3)}{m+1}.$$

Case 2. m is even.

By [20, Proposition 2.3] we have  $\mathcal{CCC}(G) = K_{\frac{n(m-2)}{2}} \sqcup 2K_n$ . Therefore, by Theorem 2.1, it follows that

$$\begin{split} \operatorname{Spec}(\mathcal{CCC}(G)) &= \left\{ (-1)^{\frac{n(m+2)-6}{2}}, \left(\frac{n(m-2)-2}{2}\right)^1, (n-1)^2 \right\}, \\ \operatorname{L-spec}(\mathcal{CCC}(G)) &= \left\{ 0^3, \left(\frac{n(m-2)}{2}\right)^{\frac{n(m-2)-2}{2}}, n^{2n-2} \right\} \\ \text{and Q-spec}(\mathcal{CCC}(G)) &= \left\{ (n(m-2)-2)^1, \left(\frac{n(m-2)-4}{2}\right)^{\frac{n(m-2)-2}{2}}, \quad (2n-2)^2, (n-2)^{2n-2} \right\}. \end{split}$$

We have

$$\left| \frac{n(m-2)-2}{2} \right| = \begin{cases} 1, & \text{if } m=2\\ \frac{n(m-2)-2}{2}, & \text{if } m \ge 4. \end{cases}$$

Therefore, by (2.1), we have

$$E(\mathcal{CCC}(G)) = \frac{n(m+2) - 6}{2} + 1 + 2(n-1) = 4(n-1),$$

if m=2. If  $m\geq 4$  then

$$E(\mathcal{CCC}(G)) = \frac{n(m+2)-6}{2} + \frac{n(m-2)-2}{2} + 2(n-1) = n(m+2)-6.$$

We have  $|V(\mathcal{CCC}(G))| = \frac{n(m+2)}{2}$  and  $|e(\mathcal{CCC}(G))| = \frac{n^2(m-2)^2 - 2n(m-4n+2)}{8}$ . Therefore,  $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)}$ . Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| = \left| \frac{-f_5(m,n)}{2(m+2)} \right|,$$

where  $f_5(m,n) = m(n(m-4)-2) + 12n - 4$ . Note that for  $m \ge 6$  we have  $f_5(m,n) > 0$  since n(m-4) > 2 and 12n - 4 > 0. For m = 2 and  $n \ge 2$  we have  $f_5(2,n) = 8n - 8 > 0$ . For m = 4 and  $n \ge 2$  we have  $f_5(4,n) = 12n - 12 > 0$ . Therefore, for all  $m \ge 2$  and  $n \ge 2$ , we have

$$\left| \frac{-f_5(m,n)}{2(m+2)} \right| = \left| \frac{f_5(m,n)}{2(m+2)} \right| = \frac{m(n(m-4)-2)+12n-4}{2(m+2)}.$$

$$\left| \frac{n(m-2)}{2} - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| \frac{n(m-2)}{2} - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| = \left| \frac{f_6(m,n)}{m+2} \right|,$$

where  $f_6(m,n) = 2n(m-4) + m + 2$ . Clearly,  $f_6(m,n) > 0$  if  $m \ge 4$  and  $n \ge 2$ . For m = 2 and  $n \ge 2$  we have  $f_6(2,n) = -4n + 4 < 0$ . Therefore,

$$\left| \frac{f_6(m,n)}{m+2} \right| = \begin{cases} n-1, & \text{if } m=2 \text{ and } n \ge 2\\ \frac{2n(m-4)+m+2}{m+2}, & \text{if } m \ge 4 \text{ and } n \ge 2. \end{cases}$$
$$\left| n - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| n - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| = \left| \frac{-f_7(m,n)}{2(m+2)} \right|,$$

where  $f_7(m, n) = mn(m-6) - 2m + 8n - 4$ . For m = 2 and  $n \ge 2$  we have  $f_7(2, n) = -8$ . For m = 4 and  $n \ge 2$  we have  $f_7(4, n) = -12$ . For m = 6 and  $n \ge 2$  we have  $f_7(6, n) = 8n - 16 \ge 0$ . Also, for  $m \ge 8$  and  $n \ge 2$  we have  $m^2 \ge 8m$  which gives  $m(m-6) \ge 2m$  and so  $mn(m-6) \ge 2mn > 2m$ . Therefore, mn(m-6) - 2m > 0 and so  $f_7(m, n) > 0$  since 8n - 4 > 0. Hence,

$$\left| \frac{-f_7(m,n)}{2(m+2)} \right| = \begin{cases} 1, & \text{if } m = 2, 4 \text{ and } n \ge 2\\ \frac{mn(m-6)-2m+8n-4}{2(m+2)}, & \text{if } m \ge 6 \text{ and } n \ge 2. \end{cases}$$

Now, by (2.2), we have

$$LE(\mathcal{CCC}(G)) = 3 \times \frac{m(n(m-4)-2)+12n-4}{2(m+2)} + \frac{n(m-2)-2}{2} \times (n-1) + (2n-2) \times 1$$
  
= 4(n-1),

if m=2 and  $n\geq 2$ . If m=4 and  $n\geq 2$  then

$$LE(\mathcal{CCC}(G)) = 3 \times \frac{m(n(m-4)-2)+12n-4}{2(m+2)} + \frac{n(m-2)-2}{2} \times \frac{2n(m-4)+m+2}{m+2} + (2n-2) \times 1$$

$$=6(n-1).$$

If  $m \ge 6$  and  $n \ge 2$  then

$$LE(\mathcal{CCC}(G)) = 3 \times \frac{m(n(m-4)-2)+12n-4}{2(m+2)} + \frac{n(m-2)-2}{2} \times \frac{2n(m-4)+m+2}{m+2} + (2n-2) \times \frac{mn(m-6)-2m+8n-4}{2(m+2)}$$
$$= \frac{2m^2n^2 - 12mn^2 + m^2n + 16n^2 - 4mn - 2m + 12n - 4}{m+2}.$$

Again,

$$\left| n(m-2) - 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| n(m-2) - 2 - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| = \left| \frac{f_8(m,n)}{2(m+2)} \right|,$$

where  $f_8(m,n) = n(m^2 - 20) + 2m(n-1) + 2mn - 4$ . For m = 2 and  $n \ge 2$  we have  $f_8(2,n) = -8n - 8 < 0$ . For m = 4 and  $n \ge 2$  we have  $f_8(4,n) = 12n - 12 > 0$ . For  $m \ge 6$  and  $n \ge 2$  we have  $f_8(m,n) > 0$ . Therefore,

$$\left| \frac{f_8(m,n)}{2(m+2)} \right| = \begin{cases} n+1, & \text{if } m=2 \text{ and } n \ge 2\\ n-1, & \text{if } m=4 \text{ and } n \ge 2\\ \frac{n(m^2-20)+2m(n-1)+2mn-4}{2(m+2)}, & \text{if } m \ge 6 \text{ and } n \ge 2. \end{cases}$$

$$\left| \frac{n(m-2)-4}{2} - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| \frac{n(m-2)-4}{2} - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| = \left| \frac{f_9(m,n)}{m+2} \right|,$$

where  $f_9(m, n) = n(m - 8) + m(n - 1) - 2$ . For m = 2 and  $n \ge 2$  we have  $f_9(2, n) = -4n - 4 < 0$ . For m = 4 and  $n \ge 2$  we have  $f_9(4, n) = -6$ . For m = 6 and  $n \ge 2$  we have  $f_9(6, n) = 4n - 8 \ge 0$ . Further, if For  $m \ge 8$  and  $n \ge 2$  then  $f_9(m, n) > 0$  since  $n(m - 8) \ge 0$  and m(n - 1) - 2 > 0. Hence,

$$\left| \frac{f_9(m,n)}{m+2} \right| = \begin{cases} n+1, & \text{if } m=2 \text{ and } n \ge 2\\ 1, & \text{if } m=4 \text{ and } n \ge 2\\ \frac{n(m-8)+m(n-1)-2}{m+2}, & \text{if } m \ge 6 \text{ and } n \ge 2. \end{cases}$$

$$\left| 2n-2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n-2 - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| = \left| \frac{-f_{10}(m,n)}{2(m+2)} \right|,$$

where  $f_{10}(m,n) = n(m^2 - 8m + 4) + 2m + 4$ . Clearly,  $f_{10}(m,n) > 0$  for  $m \ge 8$  and  $n \ge 2$ . For m = 2 and  $n \ge 2$  we have  $f_{10}(2,n) = -8n + 8 < 0$ . For m = 4 and  $n \ge 2$  we have  $f_{10}(4,n) = -12n + 12 < 0$ . For m = 6 and  $n \ge 2$  we have  $f_{10}(6,n) = -8n + 16 \le 0$ . Hence,

$$\left| \frac{f_{10}(m,n)}{m+2} \right| = \begin{cases} n-1, & \text{if } m=2 \text{ and } n \ge 2\\ n-1, & \text{if } m=4 \text{ and } n \ge 2\\ \frac{n-2}{2}, & \text{if } m=6 \text{ and } n \ge 2\\ \frac{n(m^2-8m+4)+2m+4}{2(m+2)}, & \text{if } m \ge 8 \text{ and } n \ge 2. \end{cases}$$

$$\left| n - 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| n - 2 - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| = \left| \frac{-f_{11}(m,n)}{2(m+2)} \right|,$$

where  $f_{11}(m,n) = n(m-2)(m-4) + 2m + 4$ . Note that for  $m \ge 4$  and  $n \ge 2$  we have  $f_{11}(m,n) > 0$ . For m = 2 and  $n \ge 2$  we have  $f_{11}(m,n) = 8$ . Therefore,

$$\left| \frac{-f_{11}(m,n)}{2(m+2)} \right| = \frac{f_{11}(m,n)}{2(m+2)} = \frac{n(m-2)(m-4) + 2m + 4}{2(m+2)}.$$

By (2.3), we have

$$LE^{+}(\mathcal{CCC}(G)) = n + 1 + \frac{n(m-2) - 2}{2} \times (n+1) + 2 \times (n-1) + (2n-2) \times \frac{n(m-2)(m-4) + 2m + 4}{2(m+2)}$$

if m = 2 and n > 2. If m = 4 and n > 2 then

$$LE^{+}(\mathcal{CCC}(G)) = n - 1 + \frac{n(m-2) - 2}{2} \times 1 + 2 \times (n-1) + (2n-2) \times \frac{n(m-2)(m-4) + 2m + 4}{2(m+2)}$$
$$= 6(n-1).$$

If m = 6 and  $n \ge 2$  then

$$LE^{+}(\mathcal{CCC}(G)) = \frac{n(m^{2} - 20) + 2m(n - 1) + 2mn - 4}{2(m + 2)} + \frac{n(m - 2) - 2}{2} \times \frac{n(m - 8) + m(n - 1) - 2}{m + 2} + 2 \times \frac{n - 2}{2} + (2n - 2) \times \frac{n(m - 2)(m - 4) + 2m + 4}{2(m + 2)}$$
$$= 2(n + 2)(n - 1).$$

If  $m \geq 8$  and  $n \geq 2$  then

$$LE^{+}(\mathcal{CCC}(G)) = \frac{n(m^{2} - 20) + 2m(n - 1) + 2mn - 4}{2(m + 2)} + \frac{n(m - 2) - 2}{2} \times \frac{n(m - 8) + m(n - 1) - 2}{m + 2}$$
$$+ 2 \times \frac{n(m^{2} - 8m + 4) + 2m + 4}{2(m + 2)} + (2n - 2) \times \frac{n(m - 2)(m - 4) + 2m + 4}{2(m + 2)}$$
$$= \frac{2n^{2}(m - 2)(m - 4)}{m + 2}.$$

This completes the proof.

**Theorem 3.4.** If  $G = V_{8n}$  then

$$\text{(a) } \operatorname{Spec}(\mathcal{CCC}(G)) = \begin{cases} \left\{ (-1)^{2n-2}, 0^2, (2n-2)^1 \right\}, & \text{if $n$ is odd} \\ \left\{ (-1)^{2n-1}, 1^2, (2n-3)^1 \right\}, & \text{if $n$ is even} \end{cases}$$
 
$$\operatorname{and } E(\mathcal{CCC}(G)) = \begin{cases} 4n-4, & \text{if $n$ is odd} \\ 4n-2, & \text{if $n$ is even}. \end{cases}$$
 
$$\text{(b) } \operatorname{L-spec}(\mathcal{CCC}(G)) = \begin{cases} \left\{ 0^3, (2n-1)^{2n-2} \right\}, & \text{if $n$ is odd} \\ \left\{ 0^3, 2^2, (2n-2)^{2n-3} \right\}, & \text{if $n$ is even} \end{cases}$$
 
$$\operatorname{and } \operatorname{LE}(\mathcal{CCC}(G)) = \begin{cases} \frac{6(2n-1)(2n-2)}{2n+1}, & \text{if $n$ is odd} \\ 6, & \text{if $n=2$} \\ \frac{2(2n-3)(5n-7)}{n+1}, & \text{if $n$ is even and $n \geq 4$.} \end{cases}$$

$$\begin{aligned} \text{(c)} \ \ &\mathrm{Q\text{-}spec}(\mathcal{CCC}(G)) = \begin{cases} \left\{0^2, (4n-4)^1, (2n-3)^{2n-2}\right\}, & \text{if $n$ is odd} \\ \left\{2^2, 0^2, (4n-6)^1, (2n-4)^{2n-3}\right\}, & \text{if $n$ is even} \end{cases} \\ & \text{and } LE^+(\mathcal{CCC}(G)) = \begin{cases} \frac{4(2n-1)(2n-2)}{2n+1}, & \text{if $n$ is odd} \\ 6, & \text{if $n=2$} \\ \frac{16(n-1)(n-2)}{n+1}, & \text{if $n$ is even and $n \geq 4$.} \end{cases}$$

*Proof.* We shall prove the result by considering the following cases. Case 1. n is odd.

By [20, Proposition 2.4] we have  $\mathcal{CCC}(G) = 2K_1 \sqcup K_{2n-1}$ . Therefore, by Theorem 2.1, it follows that

$$\begin{aligned} &\operatorname{Spec}(\mathcal{CCC}(G)) = \left\{ (-1)^{2n-2}, 0^2, (2n-2)^1 \right\}, \quad \operatorname{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^3, (2n-1)^{2n-2} \right\} \\ &\operatorname{and} \ \operatorname{Q-spec}(\mathcal{CCC}(G)) = \left\{ 0^2, (4n-4)^1, (2n-3)^{2n-2} \right\}. \end{aligned}$$

Hence, by (2.1), we get

$$E(\mathcal{CCC}(G)) = 2n - 2 + 2n - 2 = 4n - 4.$$

We have  $|V(\mathcal{CCC}(G))| = 2n + 1$  and  $|e(\mathcal{CCC}(G))| = \frac{(2n-1)(2n-2)}{2}$ . Therefore,  $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(2n-1)(2n-2)}{2n+1}$ . Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{(2n-1)(2n-2)}{2n+1} \quad \text{and}$$

$$\left| 2n - 1 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n - 1 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{3(2n-1)}{2n+1}.$$

Now, by (2.2), we have

$$LE(\mathcal{CCC}(G)) = 3 \times \frac{(2n-1)(2n-2)}{2n+1} + (2n-2) \times \frac{3(2n-1)}{2n+1} = \frac{6(2n-1)(2n-2)}{2n+1}.$$

Again,

$$\left| 4n - 4 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 4n - 4 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{(2n-2)(2n+3)}{2n+1} \quad \text{and} \quad \left| 2n - 3 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n - 3 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{2n-5}{2n+1}.$$

By (2.3), we have

$$LE^{+}(\mathcal{CCC}(G)) = 2 \times \frac{(2n-1)(2n-2)}{2n+1} + \frac{(2n-2)(2n+3)}{2n+1} + (2n-2) \times \frac{2n-5}{2n+1}$$
$$= \frac{4(2n-1)(2n-2)}{2n+1}.$$

Case 2. n is even.

By [20, Proposition 2.4] we have  $\mathcal{CCC}(G) = 2K_2 \sqcup K_{2n-2}$ . Therefore, by Theorem 2.1, it follows that

$$\begin{split} \operatorname{Spec}(\mathcal{CCC}(G)) &= \left\{ (-1)^{2n-1}, 1^2, (2n-3)^1 \right\}, \quad \operatorname{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^3, 2^2, (2n-2)^{2n-3} \right\} \\ \text{and } \operatorname{Q-spec}(\mathcal{CCC}(G)) &= \left\{ 2^2, 0^2, (4n-6)^1, (2n-4)^{2n-3} \right\}. \end{split}$$

Hence, by (2.1), we get

$$E(\mathcal{CCC}(G)) = 2n - 1 + 2 + 2n - 3 = 4n - 2.$$

We have  $|V(\mathcal{CCC}(G))| = 2n + 2$  and  $|e(\mathcal{CCC}(G))| = \frac{(2n-2)(2n-3)+4}{2}$ . Therefore,  $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(n-1)(2n-3)+2}{n+1}$ . Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(n-1)(2n-3) + 2}{n+1} \right| = \frac{(n-1)(2n-3) + 2}{n+1}$$

$$\left| 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2 - \frac{(n-1)(2n-3)+2}{n+1} \right| = \left| \frac{-(2n-1)(n-3)}{n+1} \right| = \begin{cases} 1, & \text{if } n=2 \\ \frac{(2n-1)(n-3)}{n+1}, & \text{if } n \geq 4 \end{cases}$$

and

$$\left| 2n - 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n - 2 - \frac{(n-1)(2n-3) + 2}{n+1} \right| = \frac{5n-7}{n+1}.$$

Now, by (2.2), we have

$$LE(\mathcal{CCC}(G)) = 3 \times \frac{(n-1)(2n-3)+2}{n+1} + 2 \times 1 + (2n-3) \times \frac{5n-7}{n+1} = 6,$$

if n=2. If  $n\geq 4$  then

$$LE(\mathcal{CCC}(G)) = 3 \times \frac{(n-1)(2n-3)+2}{n+1} + 2 \times \frac{(2n-1)(n-3)}{n+1} + (2n-3) \times \frac{5n-7}{n+1}$$
$$= \frac{2(10n^2 - 29n + 21)}{n+1} = \frac{2(2n-3)(5n-7)}{n+1}.$$

Again,

$$\left| 4n - 6 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 4n - 6 - \frac{(n-1)(2n-3) + 2}{n+1} \right| = \frac{2n^2 + 3n - 11}{n+1} \quad \text{and} \quad$$

$$\left| 2n - 4 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n - 4 - \frac{(n-1)(2n-3) + 2}{n+1} \right| = \left| \frac{3n-9}{n+1} \right| = \begin{cases} 1, & \text{if } n = 2\\ \frac{3n-9}{n+1}, & \text{if } n \geq 4. \end{cases}$$

By (2.3), we have

$$LE^{+}(\mathcal{CCC}(G)) = 2 \times 1 + 2 \times \frac{(n-1)(2n-3)+2}{n+1} + \frac{2n^2+3n-11}{n+1} + (2n-3) \times 1 = 6,$$

if n=2. If  $n\geq 4$  then

 $LE^+(\mathcal{CCC}(G))$ 

$$= 2 \times \frac{(2n-1)(n-3)}{n+1} + 2 \times \frac{(n-1)(2n-3)+2}{n+1} + \frac{2n^2+3n-11}{n+1} + (2n-3) \times \frac{3n-9}{n+1}$$

$$= \frac{16(n-1)(n-2)}{n+1}.$$

This completes the proof.

Theorem 3.5. If  $G = SD_{8n}$  then

(a) 
$$\operatorname{Spec}(\mathcal{CCC}(G)) = \begin{cases} \left\{ (-1)^{2n}, 3^1, (2n-3)^1 \right\}, & \text{if } n \text{ is odd} \\ \left\{ (-1)^{2n-2}, 0^2, (2n-2)^1 \right\}, & \text{if } n \text{ is even} \end{cases}$$

$$and \ E(\mathcal{CCC}(G)) = \begin{cases} 4n, & \text{if } n \text{ is odd} \\ 4n-4, & \text{if } n \text{ is even.} \end{cases}$$
(b)  $\operatorname{L-spec}(\mathcal{CCC}(G)) = \begin{cases} \left\{ 0^2, 4^3, (2n-2)^{2n-3} \right\}, & \text{if } n \text{ is odd} \\ \left\{ 0^3, (2n-1)^{2n-2} \right\}, & \text{if } n \text{ is even.} \end{cases}$ 

$$and \ LE(\mathcal{CCC}(G)) = \begin{cases} 12, & if \ n = 3 \\ \frac{2(2n-3)(5n-11)}{n+1}, & if \ n \ is \ odd \ and \ n \geq 5 \\ \frac{6(2n-1)(2n-2)}{2n+1}, & if \ n \ is \ even. \end{cases}$$

$$(c) \ Q\text{-spec}(\mathcal{CCC}(G)) = \begin{cases} 6^1, 2^3, (4n-6)^1, (2n-4)^{2n-3} \\ 0^2, (4n-4)^1, (2n-3)^{2n-2} \\ 12, & if \ n \ is \ even \end{cases}$$

$$and \ LE^+(\mathcal{CCC}(G)) = \begin{cases} 12, & if \ n = 3 \\ 22, & if \ n = 5 \\ \frac{16(n-1)(n-3)}{n+1}, & if \ n \ is \ odd \ and \ n \geq 7 \\ \frac{28}{5}, & if \ n = 2 \\ \frac{4(2n-1)(2n-2)}{2n+1}, & if \ n \ is \ even \ and \ n \geq 4. \end{cases}$$

*Proof.* We shall prove the result by considering the following cases. Case 1. n is odd.

By [20, Proposition 2.5] we have  $\mathcal{CCC}(G) = K_4 \sqcup K_{2n-2}$ . Therefore, by Theorem 2.1, it follows that

$$\begin{aligned} &\operatorname{Spec}(\mathcal{CCC}(G)) = \left\{ (-1)^{2n}, 3^1, (2n-3)^1 \right\}, \quad \operatorname{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^2, 4^3, (2n-2)^{2n-3} \right\} \\ &\operatorname{and} \ \operatorname{Q-spec}(\mathcal{CCC}(G)) = \left\{ 6^1, 2^3, (4n-6)^1, (2n-4)^{2n-3} \right\}. \end{aligned}$$

Hence, by (2.1), we get

$$E(\mathcal{CCC}(G)) = 2n + 3 + 2n - 3 = 4n.$$

We have  $|V(\mathcal{CCC}(G))| = 2n + 2$  and  $|e(\mathcal{CCC}(G))| = \frac{(2n-2)(2n-3)+12}{2}$ . Therefore,  $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(n-1)(2n-3)+6}{n+1}$ . Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(n-1)(2n-3)+6}{n+1} \right| = \frac{(n-1)(2n-3)+6}{n+1},$$

$$\left| 4 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 4 - \frac{(n-1)(2n-3)+6}{n+1} \right| = \left| \frac{-2n^2+9n-5}{n+1} \right| = \begin{cases} 1, & \text{if } n=3\\ \frac{2n^2-9n+5}{n+1}, & \text{if } n \geq 5 \end{cases}$$
and
$$\left| 2n-2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n-2 - \frac{(n-1)(2n-3)+6}{n+1} \right| = \frac{5n-11}{n+1}.$$

Now, by (2.2), we have

$$LE(\mathcal{CCC}(G)) = 2 \times \frac{(n-1)(2n-3)+6}{n+1} + 3 \times 1 + (2n-3) \times \frac{5n-11}{n+1} = 12,$$

if n = 3. If  $n \ge 5$  then

$$LE(\mathcal{CCC}(G)) = 2 \times \frac{(n-1)(2n-3)+6}{n+1} + 3 \times \frac{2n^2 - 9n + 5}{n+1} + (2n-3) \times \frac{5n-11}{n+1}$$
$$= \frac{2(10n^2 - 37n + 33)}{n+1} = \frac{2(2n-3)(5n-11)}{n+1}.$$

Again,

$$\begin{vmatrix} 6 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \end{vmatrix} = \begin{vmatrix} 6 - \frac{(n-1)(2n-3) + 6}{n+1} \end{vmatrix} = \begin{vmatrix} -2n^2 + 11n - 3 \\ n+1 \end{vmatrix} = \begin{cases} \frac{-2n^2 + 11n - 3}{n+1}, & \text{if } n = 3, 5 \\ \frac{2n^2 - 11n + 3}{n+1}, & \text{if } n \ge 7, \end{cases}$$
$$\begin{vmatrix} 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \end{vmatrix} = \begin{vmatrix} 2 - \frac{(n-1)(2n-3) + 6}{n+1} \end{vmatrix} = \frac{2n^2 - 7n + 7}{n+1},$$
$$\begin{vmatrix} 4n - 6 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \end{vmatrix} = \begin{vmatrix} 4n - 6 - \frac{(n-1)(2n-3) + 6}{n+1} \end{vmatrix} = \frac{2n^2 + 3n - 15}{n+1} \quad \text{and}$$

$$\left|2n-4-\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|}\right| = \left|2n-4-\frac{(n-1)(2n-3)+6}{n+1}\right| = \left|\frac{3n-13}{n+1}\right| = \begin{cases} 1, & \text{if } n=3\\ \frac{3n-13}{n+1}, & \text{if } n\geq 5. \end{cases}$$

By (2.3), we have

$$LE^{+}(\mathcal{CCC}(G)) = \frac{-2n^2 + 11n - 3}{n+1} + 3 \times \frac{2n^2 - 7n + 7}{n+1} + \frac{2n^2 + 3n - 15}{n+1} + (2n-3) \times 1 = 12,$$

if n = 3. If n = 5 then

$$LE^{+}(\mathcal{CCC}(G)) = \frac{-2n^2 + 11n - 3}{n+1} + 3 \times \frac{2n^2 - 7n + 7}{n+1} + \frac{2n^2 + 3n - 15}{n+1} + (2n-3) \times \frac{3n - 13}{n+1}$$
$$= 22.$$

If  $n \geq 7$  then

$$LE^{+}(\mathcal{CCC}(G)) = \frac{2n^{2} - 11n + 3}{n+1} + 3 \times \frac{2n^{2} - 7n + 7}{n+1} + \frac{2n^{2} + 3n - 15}{n+1} + (2n-3) \times \frac{3n-13}{n+1}$$
$$= \frac{16(n-1)(n-3)}{n+1}.$$

Case 2. n is even.

By [20, Proposition 2.5] we have  $\mathcal{CCC}(G) = 2K_1 \sqcup K_{2n-1}$ . Therefore, by Theorem 2.1, it follows that

$$\begin{aligned} &\operatorname{Spec}(\mathcal{CCC}(G)) = \left\{ (-1)^{2n-2}, 0^2, (2n-2)^1 \right\}, \quad \operatorname{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^3, (2n-1)^{2n-2} \right\} \\ &\operatorname{and} \ \operatorname{Q-spec}(\mathcal{CCC}(G)) = \left\{ 0^2, (4n-4)^1, (2n-3)^{2n-2} \right\}. \end{aligned}$$

Hence, by (2.1), we get

$$E(\mathcal{CCC}(G)) = 2n - 2 + 2n - 2 = 4n - 4.$$

We have  $V(\mathcal{CCC}(G)) = 2n + 1$  and  $e(\mathcal{CCC}(G)) = \frac{(2n-1)(2n-2)}{2}$ . So,  $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(2n-1)(2n-2)}{2n+1}$ . Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{(2n-1)(2n-2)}{2n+1} \quad \text{and} \quad \left| 2n - 1 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n - 1 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{3(2n-1)}{2n+1}.$$

Now, by (2.2), we have

$$LE(\mathcal{CCC}(G)) = 3 \times \frac{(2n-1)(2n-2)}{2n+1} + (2n-2) \times \frac{3(2n-1)}{2n+1} = \frac{6(2n-1)(2n-2)}{2n+1}$$

Again,

$$\left| 4n - 4 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 4n - 4 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{(2n-2)(2n+3)}{2n+1} \quad \text{and}$$

$$\left| 2n - 3 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n - 3 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \left| \frac{2n-5}{2n+1} \right| = \begin{cases} \frac{1}{5}, & \text{if } n = 2\\ \frac{2n-5}{2n+1}, & \text{if } n \geq 4 \end{cases}$$

By (2.3), we have

$$LE^{+}(\mathcal{CCC}(G)) = 2 \times \frac{(2n-1)(2n-2)}{2n+1} + \frac{(2n-2)(2n+3)}{2n+1} + (2n-2) \times \frac{1}{5} = \frac{28}{5},$$

if n = 2. If n > 4 then

$$LE^{+}(\mathcal{CCC}(G)) = 2 \times \frac{(2n-1)(2n-2)}{2n+1} + \frac{(2n-2)(2n+3)}{2n+1} + (2n-2) \times \frac{2n-5}{2n+1}$$
$$= \frac{4(2n-1)(2n-2)}{2n+1}.$$

This completes the proof.

We conclude this section with the following corollary.

Corollary 3.6. If G is isomorphic to  $D_{2n}$ ,  $Q_{4m}$ ,  $U_{(n,m)}$ ,  $V_{8n}$  or  $SD_{8n}$  then CCC(G) is super integral.

#### 4. Comparing various energies

In this section we compare various energies of  $\mathcal{CCC}(G)$  obtained in Section 3 and derive the following relations.

Theorem 4.1. Let  $G = D_{2n}$ .

- (a) If n = 3, 4, 6 then  $E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G))$ .
- (b) If n = 5 then  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G))$ .
- (c) If n = 10 then  $LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .
- (d) If  $n \ge 7$  but  $n \ne 10$  then  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

*Proof.* We shall prove the result by considering the following cases.

Case 1. n is odd.

If n = 3 then, by Theorem 3.1, we have

$$E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 0.$$

If n = 5 then, by Theorem 3.1, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = n - 3 - \frac{(n-3)(n+3)}{n+1} = -\frac{4}{5} < 0$$

and  $LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = \frac{8}{3}$ . Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G))$ . If  $n \ge 7$  then, by Theorem 3.1, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = n - 3 - \frac{(n-3)(n+3)}{n+1} = -\frac{2(n-3)}{n+1} < 0$$

and

$$LE^{+}(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{(n-3)(n+3)}{n+1} - \frac{2(n-1)(n-3)}{n+1} = -\frac{(n-3)(n-5)}{n+1} < 0.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

Case 2. n is even.

Consider the following subcases.

Subcase 2.1  $\frac{n}{2}$  is even.

If n = 4 then, by Theorem 3.1, we have

$$E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 0.$$

If n = 8 then, by Theorem 3.1, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = n - 4 - \frac{(n-4)(n+6)}{n+2} = -\frac{8}{5} < 0$$

and

$$LE^+(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{(n-4)(n+6)}{n+2} - \frac{3(n-2)(n-4)}{n+2} = -\frac{8}{5} < 0.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

If n > 12 then, by Theorem 3.1, we have

$$E(\mathcal{CCC}(G)) - LE^+(\mathcal{CCC}(G)) = n - 4 - \frac{2(n-2)(n-4)}{n+2} = -\frac{(n-4)(n-6)}{n+2} < 0$$

and

$$LE^{+}(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{2(n-2)(n-4)}{n+2} - \frac{3(n-2)(n-4)}{n+2} = -\frac{(n-2)(n-4)}{n+2} < 0.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

Subcase 2.2  $\frac{n}{2}$  is odd.

If n = 6 then, by Theorem 3.1, we have

$$E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 4.$$

If n = 10 then, by Theorem 3.1, we have

$$LE^{+}(\mathcal{CCC}(G)) - E(\mathcal{CCC}(G)) = \frac{22}{3} - (n-2) = -\frac{2}{3} < 0$$

and

$$E(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = n - 2 - \frac{(n-4)(3n-10)}{n+2} = -2 < 0.$$

Therefore,  $LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

If n > 14 then, by Theorem 3.1, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = n - 2 - \frac{2(n-2)(n-6)}{n+2} = -\frac{(n-2)(n-10)}{n+2} < 0$$

and

$$LE^{+}(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{2(n-2)(n-6)}{n+2} - \frac{(n-4)(3n-10)}{n+2}$$
$$= -\frac{n^2 - 6n + 16}{n+2} = -\frac{n(n-14) + 8n + 10}{n+2} < 0.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ . Hence, the result follows.

**Theorem 4.2.** Let  $G = Q_{4m}$ .

- (a) If m = 2, 3 then  $E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G))$ .
- (b) If m = 5 then  $LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .
- (c) If m = 7 then  $LE^+(\mathcal{CCC}(G)) = E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .
- (d) If m = 4,6 or m > 8 then  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

*Proof.* We shall prove the result by considering the following cases.

Case 1. m is odd.

If m=3 then, by Theorem 3.2, we have

$$E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 4.$$

If m = 5 then, by Theorem 3.2, we have

$$LE^{+}(\mathcal{CCC}(G)) - E(\mathcal{CCC}(G)) = \frac{22}{3} - (2m - 2) = -\frac{2}{3} < 0$$

and

$$E(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = 2m - 2 - \frac{2(m-2)(3m-5)}{m+1} = -2 < 0.$$

Therefore,  $LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

If m = 7 then, by Theorem 3.2, we have

$$LE^+(\mathcal{CCC}(G)) = E(\mathcal{CCC}(G)) = 12$$

and

$$LE^{+}(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{4(m-1)(m-3)}{m+1} - \frac{2(m-2)(3m-5)}{m+1} = -\frac{2(m+4)(m-1)}{m+1} < 0.$$

Therefore,  $LE^+(\mathcal{CCC}(G)) = E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

If  $m \geq 9$  then, by Theorem 3.2, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = 2m - 2 - \frac{4(m-1)(m-3)}{m+1} = -\frac{2(m-1)(m-7)}{m+1} < 0$$

and

$$LE^{+}(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{4(m-1)(m-3)}{m+1} - \frac{2(m-2)(3m-5)}{m+1} = -\frac{2(m+4)(m-1)}{m+1} < 0.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

Case 2. m is even.

If m=2 then, by Theorem 3.2, we have

$$E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 0.$$

If m = 4 then, by Theorem 3.2, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = 2m - 4 - \frac{2(m-2)(m+3)}{m+1} = -\frac{8}{5} < 0$$

and

$$LE^{+}(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{2(m-2)(m+3)}{m+1} - \frac{6(m-1)(m-2)}{m+1} = -\frac{8}{5} < 0.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

If  $m \geq 6$  then, by Theorem 3.2, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = 2m - 4 - \frac{4(m-1)(m-2)}{m+1} = -\frac{2(m-2)(m-3)}{m+1} < 0$$

and

$$LE^{+}(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{4(m-1)(m-2)}{m+1} - \frac{6(m-1)(m-2)}{m+1} = -\frac{2(m-1)(m-2)}{m+1}.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ . Hence, the result follows.

**Theorem 4.3.** Let  $G = U_{(n,m)}$ .

- (a) If m = 2, 3, 4 and  $n \ge 2$  then  $LE^+(\mathcal{CCC}(G)) = E(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G))$ .
- (b) If m = 5 and n = 2, 3; or m = 6 and n = 2 then

$$LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

(c) If m = 5 and  $n \ge 4$ ;  $m \ge 6$  and  $n \ge 3$ ; or  $m \ge 8$  and  $n \ge 2$  then

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

(d) If 
$$m = 7$$
 and  $n = 2$  then  $E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

*Proof.* We shall prove the result by considering the following cases.

Case 1. If m is odd and  $n \geq 2$ .

If m = 3 and  $n \ge 2$  then, by Theorem 3.3, we have

$$LE^+(\mathcal{CCC}(G)) = E(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 4(n-1).$$

If m = 5 and n = 2 then, by Theorem 3.3, we have

$$LE^{+}(\mathcal{CCC}(G)) - E(\mathcal{CCC}(G)) = \frac{2n^{2} + 10n - 6}{3} - (n(m+1) - 4) = -\frac{2}{3} < 0$$

and

$$E(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = n(m+1) - 4 - \frac{2(2n-1)(n+3)}{3} = -2 < 0.$$

Therefore,  $LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

If m = 5 and n = 3 then, by Theorem 3.3, we have

$$LE^{+}(\mathcal{CCC}(G)) - E(\mathcal{CCC}(G)) = \frac{2(2n+3)(n-1)}{3} - (n(m+1)-4) = -2 < 0$$

and

$$E(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = n(m+1) - 4 - \frac{2(2n-1)(n+3)}{3} = -4 < 0.$$

Therefore,  $LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

If m = 5 and  $n \ge 4$  then, by Theorem 3.3, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = n(m+1) - 4 - \frac{2(2n+3)(n-1)}{3}$$
$$= \frac{-2(2n^{2} - 8n + 3)}{3} = \frac{-2(2n(n-4) + 3)}{3} < 0$$

and

$$LE^{+}(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{2(2n+3)(n-1)}{3} - \frac{2(2n-1)(n+3)}{3} = \frac{-8n}{3} < 0.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

If  $m \geq 7$  and  $n \geq 2$  then, by Theorem 3.3, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = n(m+1) - 4 - \frac{n^{2}(m-1)(m-3)}{m+1} = -\frac{f_{1}(m,n)}{m+1},$$

where  $f_1(m,n) = mn(m-4)(n-3) + 2mn(m-7) + 3n(n-1) + 4(m+1)$ . For  $m \ge 7$  and n = 2 we have  $f_1(m,n) = 2(m-1)(m-7) \ge 0$ . Hence,  $f_1(7,2) = 0$  and  $f_1(m,2) > 0$  if  $m \ge 9$ . Thus,  $E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G))$  and  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G))$  according as if m = 7, n = 2 and  $m \ge 9$ , n = 2. For  $m \ge 7$  and  $n \ge 3$  we have  $f_1(m,n) > 0$  and so  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G))$ .

If  $m \geq 7$  and  $n \geq 2$  then, by Theorem 3.3, we also have

$$\begin{split} LE^+(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) \\ &= \frac{n^2(m-1)(m-3)}{m+1} - \frac{m^2n^2 - 4mn^2 + m^2n + 3n^2 - 2mn - 2m + 5n - 2}{m+1} \\ &= -\frac{m^2n - 2mn - 2m + 5n - 2}{m+1} = -\frac{(mn-2)(m-2) + 5(n-2) + 4}{m+1} < 0. \end{split}$$

Therefore,  $LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ . Thus, if m=7 and n=2 then

$$E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$$

and if  $m \geq 7$  and  $n \geq 3$  or  $m \geq 9$  and n = 2 then

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

Case 2. m is even and  $n \geq 2$ .

If m=2 and  $n\geq 2$  then, by Theorem 3.3, we have

$$LE^+(\mathcal{CCC}(G)) = E(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 4(n-1).$$

If m = 4 and  $n \ge 2$  then, by Theorem 3.3, we have

$$LE^+(\mathcal{CCC}(G)) = E(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 6(n-1).$$

If m = 6 and n = 2 then, by Theorem 3.3, we have

$$LE^{+}(\mathcal{CCC}(G)) - E(\mathcal{CCC}(G)) = 2(n+2)(n-1) - (n(m+2)-6) = -4 < 0$$

and

$$E(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G))$$

$$= n(m+2) - 6 - \frac{2m^2n^2 - 12mn^2 + m^2n + 16n^2 - 4mn - 2m + 12n - 4}{m+2} = -2 < 0.$$

Therefore,  $LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

If m = 6 and  $n \ge 3$  then by Theorem 3.3

$$E(\mathcal{CCC}(G)) - LE^+(\mathcal{CCC}(G)) = n(m+2) - 6 - 2(n+2)(n-1) = 2n(3-n) - 2 < 0$$

and

$$\begin{split} LE^+(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) \\ &= 2(n+2)(n-1) - \frac{2m^2n^2 - 12mn^2 + m^2n + 16n^2 - 4mn - 2m + 12n - 4}{m+2} \\ &= -(n+2) < 0. \end{split}$$

Therefore  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

If  $m \geq 8$  and  $n \geq 2$  then, by Theorem 3.3, we have

$$\begin{split} E(\mathcal{CCC}(G)) - LE^+(\mathcal{CCC}(G)) &= n(m+2) - 6 - \frac{2n^2(m-2)(m-4)}{m+2} \\ &= -\frac{2m^2n^2 - 12mn^2 - m^2n + 16n^2 - 4mn + 6m - 4n + 12}{m+2} \\ &= -\frac{f_2(m,n)}{m+2}, \end{split}$$

where  $f_2(m,n) = mn(2n-1)(m-8) + 2m(2n(n-3)+3) + 4n(4n-1) + 12$ . For n = 2 and  $m \ge 8$  we have  $f_2(m,n) = (6m-2)(m-8) + 52 > 0$ . For  $n \ge 3$  and  $m \ge 8$  we have  $f_2(m,n) > 0$ . Therefore, if  $m \ge 8$  and  $n \ge 2$  then  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G))$ .

If  $m \geq 8$  and  $n \geq 2$  then, by Theorem 3.3, we also have

$$\begin{split} LE^+(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) \\ &= \frac{2n^2(m-2)(m-4)}{m+2} - \frac{2m^2n^2 - 12mn^2 + m^2n + 16n^2 - 4mn - 2m + 12n - 4}{m+2} \\ &= -\frac{m^2n - 4mn - 2m + 12n - 4}{m+2} \\ &= -\frac{mn(m-8) + 2m(2n-1) + 4(3n-1)}{m+2} < 0. \end{split}$$

Therefore,  $LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ . Thus, if  $m \geq 8$  and  $n \geq 2$  then

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

Hence, the result follows.

**Theorem 4.4.** If  $G = V_{8n}$  then  $E(\mathcal{CCC}(G)) \leq LE^+(\mathcal{CCC}(G)) \leq LE(\mathcal{CCC}(G))$ . The equality holds if and only if n = 2.

*Proof.* We shall prove the result by considering the following cases.

Case 1. n is odd.

By Theorem 3.4, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = 4n - 4 - \frac{4(2n-1)(2n-2)}{2n+1} = -\frac{4(n-1)(2n-3)}{2n+1}$$

and

$$LE^+(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{4(2n-1)(2n-2)}{2n+1} - \frac{6(2n-1)(2n-2)}{2n+1} = -\frac{2(2n-1)(2n-2)}{2n+1} < 0.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

Case 2. n is even.

If n = 2 then, by Theorem 3.4, we have

$$E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 6.$$

If  $n \geq 4$  then, by Theorem 3.4, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = 4n - 2 - \frac{16(n-1)(n-2)}{n+1}$$
$$= -\frac{2(6n^{2} + 25n - 17)}{n+1} = -\frac{2(6n(n-4) + 49n - 7)}{n+1} < 0$$

and

$$LE^{+}(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{16(n-1)(n-2)}{n+1} - \frac{2(2n-3)(5n-7)}{n+1}$$
$$= -\frac{2(2n^2 - 5n + 5)}{n+1} = -\frac{2(2n(n-4) + 3n + 5)}{n+1} < 0.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ . Hence, the result follows.

**Theorem 4.5.** If  $G = SD_{8n}$  then  $E(\mathcal{CCC}(G)) \leq LE^+(\mathcal{CCC}(G)) \leq LE(\mathcal{CCC}(G))$ . The equality holds if and only if n = 3.

*Proof.* We shall prove the result by considering the following cases.

Case 1. n is odd.

If n = 3 then, by Theorem 3.5, we have

$$E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 12.$$

If n = 5 then, by Theorem 3.5, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = 4n - 22 = -2 < 0$$

and

$$LE^{+}(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = 22 - \frac{2(2n-3)(5n-11)}{n+1} = -\frac{32}{3} < 0.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

If  $n \geq 7$  then, by Theorem 3.5, we have

$$\begin{split} E(\mathcal{CCC}(G)) - LE^+(\mathcal{CCC}(G)) &= 4n - \frac{16(n-1)(n-3)}{n+1} \\ &= -\frac{4(3n^2 - 17n + 12)}{n+1} = -\frac{4(3n(n-7) + 4n + 12)}{n+1} < 0 \end{split}$$

and

$$LE^{+}(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{16(n-1)(n-3)}{n+1} - \frac{2(2n-3)(5n-11)}{n+1}$$
$$= -\frac{2(2n^2 - 5n + 9)}{n+1} = -\frac{2(2n(n-7) + 9n + 9)}{n+1} < 0.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

Case 2. n is even.

If n = 2 then, by Theorem 3.5, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = 4n - 4 - \frac{28}{5} = -\frac{8}{5} < 0$$

and

$$LE^{+}(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{28}{5} - \frac{6(2n-1)(2n-2)}{2n+1} = -\frac{8}{5} < 0.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .

If  $n \geq 4$  then, by Theorem 3.5, we have

$$E(\mathcal{CCC}(G)) - LE^{+}(\mathcal{CCC}(G)) = 4n - 4 - \frac{4(2n-1)(2n-2)}{2n+1} = -\frac{4(n-1)(2n-3)}{2n+1} < 0$$

and

$$LE^{+}(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{4(2n-1)(2n-2)}{2n+1} - \frac{6(2n-1)(2n-2)}{2n+1} = -\frac{2(2n-1)(2n-2)}{2n+1} < 0.$$

Therefore,  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ . Hence, the result follows.

Note that Theorems 4.1–4.5 can be summarized in the following way.

**Theorem 4.6.** Let G be a finite non-abelian group. Then we have the following.

- (a) If G is isomorphic to  $D_6, D_8, D_{12}, Q_8, Q_{12}, U_{(n,2)}, U_{(n,3)}, U_{(n,4)} (n \ge 2), V_{16} \text{ or } SD_{24} \text{ then}$  $E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)).$
- (b) If G is isomorphic to  $D_{20}, Q_{20}, U_{(2,5)}, U_{(3,5)}$  or  $U_{(2,6)}$  then

$$LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

(c) If G is isomorphic to  $D_{14}, D_{16}, D_{18}, D_{2n} (n \ge 11), Q_{16}, Q_{24}, Q_{4m} (m \ge 8), U_{(n,5)}, (n \ge 4), U_{(n,m)} (m \ge 6 \text{ and } n \ge 3), U_{(n,m)} (m \ge 8 \text{ and } n \ge 2), V_{8n} (n \ge 3), SD_{16} \text{ or } SD_{8n} (n \ge 4)$ then

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

- (d) If G is isomorphic to  $Q_{28}$  or  $U_{(2,7)}$  then  $E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$ .
- (e) If G is isomorphic to  $D_{10}$  then  $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G))$ .

We conclude this section with the following remark regarding Conjecture 2.2 and Question 2.3.

**Remark 4.7.** By Theorem 4.6, it follows that  $E(\mathcal{CCC}(G)) \leq LE(\mathcal{CCC}(G))$  and  $LE^+(\mathcal{CCC}(G)) \leq LE(\mathcal{CCC}(G))$  for commuting conjugacy class graph of the groups  $D_{2n}, Q_{4m}, U_{(n,m)}, V_{8n}$  and  $SD_{8n}$ . Therefore, Conjecture 2.2 holds for commuting conjugacy class graph of these groups whereas the inequality in Question 2.3 does not. However,  $LE(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G))$  if  $G = D_6, D_8, D_{10}, D_{12}, Q_8, Q_{12}, V_{16}, SD_{24}$  and  $U_{(n,m)}$  where m = 2, 3, 4 and  $n \geq 2$ .

5. Hyperenergetic and borderenergetic graphs

It is well-known that

(5.1) 
$$E(K_n) = LE(K_n) = LE^+(K_n) = 2(n-1).$$

A graph  $\mathcal{G}$  with n vertices is called hyperenergetic, L-hyperenergetic or Q-hyperenergetic according as  $E(K_n) < E(\mathcal{G})$ ,  $LE(K_n) < LE(\mathcal{G})$  or  $LE^+(K_n) < LE^+(\mathcal{G})$ . Also,  $\mathcal{G}$  is called borderenergetic, L-borderenergetic and Q-borderenergetic if  $E(K_n) = E(\mathcal{G})$ ,  $LE(K_n) = LE(\mathcal{G})$  and  $LE^+(K_n) = LE^+(\mathcal{G})$  respectively. These graphs are considered in [24, 11, 10, 23, 9]. In this section we consider commuting conjugacy class graph  $\mathcal{CCC}(G)$  for the groups considered in Section 3 and determine whether they are hyperenergetic, L-hyperenergetic or Q-hyperenergetic. We shall also determine whether they are borderenergetic, L-borderenergetic or Q-borderenergetic.

Theorem 5.1. Let  $G = D_{2n}$ .

- (a) If n is odd or n = 4,6 then CCC(G) is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.
- (b) If n = 8, 10, 12, 14 then CCC(G) is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.
- (c) If n is even and  $n \ge 16$  then CCC(G) is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

*Proof.* We shall prove the result by considering the following cases. Case 1. n is odd.

By [20, Proposition 2.1] we have  $\mathcal{CCC}(G) = K_1 \sqcup K_{\frac{n-1}{2}}$ . Therefore,  $|V(\mathcal{CCC}(G))| = \frac{n+1}{2}$ . Using (5.1), we get

(5.2) 
$$E(K_{|V(\mathcal{CCC}(G))|}) = LE^+(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = n - 1$$

If n = 3 then, by Theorem 4.1 and Theorem 3.1, we get

$$(5.3) E(\mathcal{CCC}(G)) = LE^{+}(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 0 < 2 = E(K_{|V(\mathcal{CCC}(G))|}).$$

If n = 5 then, by Theorem 4.1 and Theorem 3.1, we get

$$(5.4) E(\mathcal{CCC}(G)) < LE^{+}(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = \frac{8}{3} < 4 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore  $\mathcal{CCC}(G)$  is neither hyperenergetic nor L-hyperenergetic nor Q-hyperenergetic for n = 5. If  $n \geq 7$  then, by Theorem 4.1 and Theorem 3.1, we get

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)) = \frac{(n-3)(n+3)}{n+1}.$$

Again,

$$\frac{(n-3)(n+3)}{n+1} - (n-1) = -\frac{8}{n+1} < 0$$

Therefore,

$$(5.5) E(\mathcal{CCC}(G)) < LE^{+}(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)) < n - 1 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Hence, in view of (5.2)–(5.5), it follows that  $\mathcal{CCC}(G)$  is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

Case 2. n is even.

By [20, Proposition 2.1] we have  $\mathcal{CCC}(G) = 2K_1 \sqcup K_{\frac{n}{2}-1}$ . Therefore,  $|V(\mathcal{CCC}(G))| = \frac{n}{2} + 1$ . Using (5.1), we get

$$(5.6) E(K_{|V(\mathcal{CCC}(G))|}) = LE^+(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = n.$$

Subcase 2.1.  $\frac{n}{2}$  is even.

If n = 4 then, by Theorem 4.1 and Theorem 3.1, we get

$$(5.7) E(\mathcal{CCC}(G)) = LE^{+}(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 0 < 4 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, by (5.6) and by (5.7), it follows that  $\mathcal{CCC}(G)$  is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If n = 8 then, by Theorem 4.1 and Theorem 3.1, we get

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = 6 < 8 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Also,

$$LE(\mathcal{CCC}(G)) = 9 > 8 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

So,  $\mathcal{CCC}(G)$  is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If  $n \geq 12$  then, by Theorem 3.1, we get

$$LE(\mathcal{CCC}(G)) = \frac{3(n-2)(n-4)}{n+2}.$$

We have

$$n - \frac{3(n-2)(n-4)}{n+2} = -\frac{2(n(n-12) + 2n + 12)}{n+2} < 0.$$

So,  $LE(K_{|V(\mathcal{CCC}(G))|}) < LE(\mathcal{CCC}(G))$  and so  $\mathcal{CCC}(G)$  is L-hyperenergetic but not L-borderenergetic. By Theorem 4.1 and Theorem 3.1, we also get

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = \frac{2(n-2)(n-4)}{n+2}.$$

We have

(5.8) 
$$\frac{2(n-2)(n-4)}{n+2} - n = \frac{n^2 - 14n + 16}{n+2} = \frac{n(n-16) + 2n + 16}{n+2} := f_1(n)$$

Therefore, for n = 12, we have  $f_1(n) < 0$  and so

$$E(\mathcal{CCC}(G)) < LE^{+}(\mathcal{CCC}(G)) = \frac{2(n-2)(n-4)}{n+2} < n = LE^{+}(K_{|V(\mathcal{CCC}(G))|}).$$

Thus, if n = 12 then  $\mathcal{CCC}(G)$  is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If  $n \geq 16$  then, by (5.8), we have  $f_1(n) > 0$  and so  $LE^+(\mathcal{CCC}(G)) > n = LE^+(K_{|V(\mathcal{CCC}(G))|})$ . Therefore,  $\mathcal{CCC}(G)$  is Q-hyperenergetic but not Q-borderenergetic. Also,  $E(\mathcal{CCC}(G)) = n-4 < n = E(K_{|V(\mathcal{CCC}(G))|})$  and so  $\mathcal{CCC}(G)$  is neither hyperenergetic nor borderenergetic. Thus, if  $n \geq 16$  then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

Subcase 2.2.  $\frac{n}{2}$  is odd.

If n = 6 then, by Theorem 4.1 and Theorem 3.1, we get

$$E(\mathcal{CCC}(G)) = LE^{+}(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 4 < 6 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore,  $\mathcal{CCC}(G)$  is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If n = 10 then, by Theorem 4.1 and Theorem 3.1, we get

$$LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)) = 10 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

So,  $\mathcal{CCC}(G)$  is L-border erenergetic but neither hyperenergetic, border energetic, L-border energetic, Q-hyperenergetic nor Q-border energetic.

If  $n \ge 14$  then, by Theorem 3.1, we get

$$LE(\mathcal{CCC}(G)) = \frac{(n-4)(3n-10)}{n+2}.$$

We have

$$n - \frac{(n-4)(3n-10)}{n+2} = -\frac{2n(n-14) + 4n + 40}{n+2} < 0.$$

So,  $LE(K_{|V(\mathcal{CCC}(G))|}) < LE(\mathcal{CCC}(G))$  and so  $\mathcal{CCC}(G)$  is L-hyperenergetic but not L-border By Theorem 4.1 and Theorem 3.1, we also get

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = \frac{2(n-2)(n-6)}{n+2}.$$

We have

(5.9) 
$$\frac{2(n-2)(n-6)}{n+2} - n = \frac{n^2 - 18n + 24}{n+2} = \frac{n(n-18) + 24}{n+2} := f_2(n).$$

Therefore, for n = 14, we have  $f_2(n) < 0$  and so

$$E(\mathcal{CCC}(G)) < LE^{+}(\mathcal{CCC}(G)) = \frac{2(n-2)(n-4)}{n+2} < n = LE^{+}(K_{|V(\mathcal{CCC}(G))|}).$$

Thus, if n=14 then  $\mathcal{CCC}(G)$  is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. If  $n \geq 18$  then, by (5.9), we have  $f_2(n) > 0$  and so  $LE^+(\mathcal{CCC}(G)) > n = LE^+(K_{|V(\mathcal{CCC}(G))|})$ . Therefore,  $\mathcal{CCC}(G)$  is Q-hyperenergetic but not Q-borderenergetic. Also,  $E(\mathcal{CCC}(G)) = n-2 < n = E(K_{|V(\mathcal{CCC}(G))|})$  and so  $\mathcal{CCC}(G)$  is neither hyperenergetic nor borderenergetic. Thus, if  $n \geq 18$  then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

Theorem 5.2. Let  $G = Q_{4m}$ .

- (a) If m = 2, 3, 4 then CCC(G) is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-hyperenergetic, Q-hyperenergetic nor Q-borderenergetic.
- (b) If m = 5 then CCC(G) is L-borderenergetic but neither hyperenergetic, borderenergetic, L-hyperenergetic, Q-hyperenergetic nor Q-borderenergetic.

- (c) If m=6,7 then  $\mathcal{CCC}(G)$  is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.
- (d) If  $m \geq 8$  then CCC(G) is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

*Proof.* We shall prove the result by considering the following cases.

Case 1. m is odd.

By [20, Proposition 2.2] we have  $\mathcal{CCC}(G) = K_2 \sqcup K_{m-1}$ . Therefore,  $|V(\mathcal{CCC}(G))| = m+1$ . Using (5.1), we get

$$(5.10) E(K_{|V(\mathcal{CCC}(G))|}) = LE^+(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = 2m.$$

If m = 3 then, by Theorem 4.2 and Theorem 3.2, we get

$$(5.11) E(\mathcal{CCC}(G)) = LE^{+}(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 4 < 6 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, by (5.10) and (5.11), CCC(G) is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If m = 5 then, by Theorem 4.2 and Theorem 3.2, we get

$$LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)) = 10 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

So,  $\mathcal{CCC}(G)$  is L-border energetic but neither hyperenergetic, border energetic, L-hyperenergetic, Q-hyperenergetic nor Q-border energetic.

If m = 7 then, by Theorem 4.2 and Theorem 3.2, we get

$$LE^+(\mathcal{CCC}(G)) = E(\mathcal{CCC}(G)) = 12 < 14 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Also,

$$LE(\mathcal{CCC}(G)) = 20 > 14 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

So,  $\mathcal{CCC}(G)$  is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If  $m \geq 9$  then, by Theorem 4.2 and Theorem 3.2, we get

$$\frac{4(m-1)(m-3)}{m+1} = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

We have

$$2m - \frac{4(m-1)(m-3)}{m+1} = -\frac{2(m(m-9)+6)}{m+1} < 0$$

and so  $LE^+(K_{|V(\mathcal{CCC}(G))|})=2m<\frac{4(m-1)(m-3)}{m+1}=LE^+(\mathcal{CCC}(G))< LE(\mathcal{CCC}(G)).$  Hence,  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither L-border energetic nor Q-border energetic. Also,

$$E(\mathcal{CCC}(G)) = 2m - 2 < 2m = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore,  $\mathcal{CCC}(G)$  is neither hyperenergetic nor borderenergetic. Thus, if  $m \geq 9$  then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

Case 2. m is even.

By [20, Proposition 2.2] we have  $\mathcal{CCC}(G) = 2K_1 \sqcup K_{m-1}$ . Therefore,  $|V(\mathcal{CCC}(G))| = m+1$ . Using (5.1), we get

$$(5.12) E(K_{|V(\mathcal{CCC}(G))|}) = LE^+(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = 2m.$$

If m=2 then, by Theorem 4.2 and Theorem 3.2, we get

$$(5.13) E(\mathcal{CCC}(G)) = LE^{+}(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 0 < 4 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, by (5.12) and (5.13),  $\mathcal{CCC}(G)$  is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If m = 4 then, by Theorem 4.2 and Theorem 3.2, we get

$$(5.14) E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)) = \frac{36}{5} < 8 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, by (5.12) and (5.14), CCC(G) is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If  $m \ge 6$  then, by Theorem 3.2, we get

$$LE(\mathcal{CCC}(G)) = \frac{6(m-1)(m-2)}{m+1}.$$

We have

$$2m - \frac{6(m-1)(m-2)}{m+1} = -\frac{4(m^2 - 5m + 3)}{m+1} = -\frac{4(m(m-6) + m + 3)}{m+1} < 0$$

and so  $LE(K_{|V(\mathcal{CCC}(G))|}) = 2m < \frac{6(m-1)(m-2)}{m+1} = LE(\mathcal{CCC}(G))$ . Hence,  $\mathcal{CCC}(G)$  is L-hyperenergetic but not L-border energetic.

By Theorem 4.2 and Theorem 3.2, we also get

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = \frac{4(m-1)(m-2)}{m+1}.$$

We have

(5.15) 
$$\frac{4(m-1)(m-2)}{m+1} - 2m = \frac{2(m^2 - 7m + 4)}{m+1} = \frac{2(m(m-8) + m + 4)}{m+1} = f(m).$$

Therefore, for m = 6, we have f(m) < 0 and so

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = \frac{4(m-1)(m-2)}{m+1} < 2m = LE^+(K_{|V(\mathcal{CCC}(G))|}).$$

Thus, if m=6 then  $\mathcal{CCC}(G)$  is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If  $n \geq 8$  then, by (5.15), we have f(m) > 0 and so  $LE^+(\mathcal{CCC}(G)) > 2m = LE^+(K_{|V(\mathcal{CCC}(G))|})$ . Therefore,  $\mathcal{CCC}(G)$  is Q-hyperenergetic but not Q-borderenergetic. Also,  $E(\mathcal{CCC}(G)) = 2m - 4 < 2m = E(K_{|V(\mathcal{CCC}(G))|})$  and so  $\mathcal{CCC}(G)$  is neither hyperenergetic nor borderenergetic. Thus, if  $n \geq 8$  then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

**Theorem 5.3.** Let  $G = U_{(n,m)}$ .

- (a) If m = 2, 3, 4 and  $n \ge 2$  or m = 6 and n = 2 then CCC(G) is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.
- (b) If m = 5 and n = 2 then CCC(G) is L-borderenergetic but neither hyperenergetic, borderenergetic, L-hyperenergetic, Q-hyperenergetic nor Q-borderenergetic.
- (c) If m = 5 and n = 3, m = 6 and n = 3 or m = 7 and n = 2 then  $\mathcal{CCC}(G)$  is L-hyper-energetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.
- (d) If m = 5, 6 and  $n \ge 4$ ; m = 7 and  $n \ge 3$  or  $m \ge 8$  and  $n \ge 2$  then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-border-energetic nor Q-borderenergetic.

*Proof.* We shall prove the result by considering the following cases.

Case 1. m is odd and  $n \geq 2$ .

By [20, Proposition 2.3] we have  $\mathcal{CCC}(G) = K_{\frac{n(m-1)}{2}} \sqcup K_n$ . Therefore,  $|V(\mathcal{CCC}(G))| = \frac{n(m+1)}{2}$ . Using (5.1), we get

(5.16) 
$$E(K_{|V(\mathcal{CCC}(G))|}) = LE^{+}(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = mn + n - 2.$$

By Theorem 3.3 we get

$$E(\mathcal{CCC}(G)) = mn + n - 4 < mn + n - 2.$$

Therefore,  $\mathcal{CCC}(G)$  is neither hyperenergetic nor borderenergetic.

If m=3 and  $n\geq 2$  then, by Theorem 3.3, we get

$$LE^{+}(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 4n - 4 < 4n - 2 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

So,  $\mathcal{CCC}(G)$  is neither L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if m=3 and  $n\geq 2$  then  $\mathcal{CCC}(G)$  is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If m = 5 and n = 2 then, by Theorem 4.3 and Theorem 3.3, we get

$$LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)) = 10 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore,  $\mathcal{CCC}(G)$  is L-borderenergetic but neither L-hyperenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if m=5 and n=2 then  $\mathcal{CCC}(G)$  is L-borderenergetic but neither hyperenergetic, borderenergetic, L-hyperenergetic, Q-hyperenergetic nor Q-borderenergetic.

If m = 5 and n = 3 then, by Theorem 3.3, we get

$$LE(\mathcal{CCC}(G)) = 20 > 16 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore,  $\mathcal{CCC}(G)$  is L-hyperenergetic but not L-borderenergetic. Also,

$$LE^+(\mathcal{CCC}(G)) = 12 < 16 = LE^+(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore,  $\mathcal{CCC}(G)$  is neither Q-hyperenergetic nor Q-border energetic. Thus, if m=5 and n=3 then  $\mathcal{CCC}(G)$  is L-hyperenergetic but neither hyperenergetic, border energetic, L-border energetic, Q-hyperenergetic nor Q-border energetic.

If m=5 and  $n\geq 4$  then, by Theorem 4.3 and Theorem 3.3, we get

$$\frac{2(2n+3)(n-1)}{3} = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

We have

$$6n - 2 - \frac{2(2n+3)(n-1)}{3} = -\frac{4(n^2 - n - 1)}{3} = -\frac{4(n(n-4) + 3n - 1)}{3} < 0$$

and so  $LE^+(K_{|V(\mathcal{CCC}(G))|})=6n-2<\frac{2(2n+3)(n-1)}{3}=LE^+(\mathcal{CCC}(G))< LE(\mathcal{CCC}(G))$ . Therefore,  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither L-borderenergetic nor Q-borderenergetic. Thus, if m=5 and  $n\geq 4$  then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

If  $m \geq 7$  and  $n \geq 2$  then, by Theorem 3.3, we get

$$LE(\mathcal{CCC}(G)) = \frac{m^2n^2 - 4mn^2 + m^2n + 3n^2 - 2mn - 2m + 5n - 2}{m+1}.$$

We have

$$\begin{split} mn + n - 2 - LE(\mathcal{CCC}(G)) &= -\frac{m^2n^2 - 4mn^2 - 4mn + 3n^2 + 4n}{m+1} \\ &= -\frac{mn^2(m-7) + 2mn(n-2) + mn^2 + 3n^2 + 4n}{m+1} < 0 \end{split}$$

and so  $LE(K_{|V(\mathcal{CCC}(G))|}) = mn + n - 2 < LE(\mathcal{CCC}(G))$ . Therefore,  $\mathcal{CCC}(G)$  is L-hyperenergetic but not L-borderenergetic. By Theorem 4.3 and Theorem 3.3, we also get

$$\frac{n^2(m-1)(m-3)}{m+1} = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

Let 
$$f_1(m,n) = \frac{n^2(m-1)(m-3)}{m+1} - (mn+n-2)$$
. Then

$$f_1(m,n) = \frac{2 + 2m - 2mn - m^2n - n + 3n^2 - 4mn^2 + m^2n^2}{m+1}$$
$$= \frac{mn^2(m-11) + mn^2 + m^2n(n-2) + 2mn(n-2) + 2n(3n-1) + 2(m+1)}{2(m+1)}.$$

For m=7 and n=2 we have  $f_1(m,n)=-2<0$  and so

$$LE^{+}(\mathcal{CCC}(G)) = \frac{n^{2}(m-1)(m-3)}{m+1} < mn+n-2 = LE^{+}(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, CCC(G) is neither Q-hyperenergetic nor Q-borderenergetic. Thus, if m = 7 and n = 2 then CCC(G) is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If m=7 and  $n\geq 3$  then  $f_1(m,n)=\frac{n(3n-8)+16}{8}>0$ . Therefore,

$$LE^{+}(K_{|V(\mathcal{CCC}(G))|}) = mn + n - 2 < \frac{n^{2}(m-1)(m-3)}{m+1} = LE^{+}(\mathcal{CCC}(G))$$

and so  $\mathcal{CCC}(G)$  is Q-hyperenergetic but not Q-border energetic. Thus, if m=7 and  $n\geq 3$  then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, border energetic, L-border energetic nor Q-border energetic.

Now, for m=9 and n=2 we have  $f_1(m,n)=\frac{6}{5}>0$ . For m=9 and  $n\geq 3$  we have  $f_1(m,n)=\frac{2n(12n-25)+10}{5}>0$ . For  $m\geq 11$  and  $n\geq 2$  we have  $f_1(m,n)>0$ . Therefore, for  $m\geq 9$  and  $n\geq 2$  we have

$$LE^{+}(K_{|V(\mathcal{CCC}(G))|}) = mn + n - 2 < \frac{n^{2}(m-1)(m-3)}{m+1} = LE^{+}(\mathcal{CCC}(G))$$

and so  $\mathcal{CCC}(G)$  is Q-hyperenergetic but not Q-border energetic. Thus, if  $m \geq 9$  and  $n \geq 2$  then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, border energetic, L-border energetic nor Q-border energetic.

Case 2. m is even and  $n \geq 2$ .

By [20, Proposition 2.3] we have  $\mathcal{CCC}(G) = K_{\frac{n(m-2)}{2}} \sqcup 2K_n$ . Therefore,  $|V(\mathcal{CCC}(G))| = \frac{n(m+2)}{2}$ . Using (5.1), we get

(5.17) 
$$E(K_{|V(\mathcal{CCC}(G))|}) = LE^{+}(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = mn + 2n - 2.$$

By Theorem 3.3 we get

$$E(CCC(G)) = 4n - 4 < 4n - 2 = E(K_{|V(CCC(G))|}),$$

if m=2. If  $m\geq 4$  then

$$E(\mathcal{CCC}(G)) = mn + 2n - 6 < mn + 2n - 2 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore,  $\mathcal{CCC}(G)$  is neither hyperenergetic nor borderenergetic.

If m=2 and  $n\geq 2$  then, by Theorem 3.3, we get

$$LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 4n - 4 < 4n - 2.$$

So,  $\mathcal{CCC}(G)$  is neither L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if m=2 and  $n\leq 2$  then  $\mathcal{CCC}(G)$  is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If m = 4 and  $n \ge 2$  then, by Theorem 3.3, we get

$$LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 6n - 6 < 6n - 2.$$

So,  $\mathcal{CCC}(G)$  is neither L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if m=4 and  $n\leq 2$  then  $\mathcal{CCC}(G)$  is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If m = 6 and n = 2 then, Theorem 4.3 and Theorem 3.3, we get

$$LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)) = 12 < 14 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

So,  $\mathcal{CCC}(G)$  is neither L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if m=6 and n=2 then  $\mathcal{CCC}(G)$  is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. If m=6 and  $n\geq 3$  then, by Theorem 3.3, we get

$$LE(\mathcal{CCC}(G)) = 2n^2 + 3n - 2.$$

We have

$$8n - 2 - (2n^2 + 3n - 2) = -n(2n - 5) < 0.$$

Therefore,  $LE(K_{|V(\mathcal{CCC}(G))|}) = 8n - 2 < 2n^2 + 3n - 2 < LE(\mathcal{CCC}(G))$  and so  $\mathcal{CCC}(G)$  is Lhyperenergetic but not L-borderenergetic. By Theorem 3.3, we also get

$$LE^{+}(\mathcal{CCC}(G)) = 2(n+2)(n-1).$$

Let g(n) = 2(n+2)(n-1) - (8n-2). Then g(n) = 2(n(n-4) + n - 1). Therefore, if n = 3 then g(n) = -2 < 0 and so

$$LE^{+}(\mathcal{CCC}(G)) = 2(n+2)(n-1) < 8n-2 = LE^{+}(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore,  $\mathcal{CCC}(G)$  is neither Q-hyperenergetic nor Q-borderenergetic. Thus, if m=6 and n=3then  $\mathcal{CCC}(G)$  is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If  $n \geq 4$  then g(n) > 0 and so

$$LE^{+}(K_{|V(\mathcal{CCC}(G))|}) = 8n - 2 < 2(n+2)(n-1) = LE^{+}(\mathcal{CCC}(G)).$$

Therefore,  $\mathcal{CCC}(G)$  is Q-hyperenergetic but not Q-borderenergetic. Thus, if m=6 and n>4then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

If m > 8 and n > 2 then, by Theorem 4.3 and Theorem 3.3, we get

$$\frac{2n^2(m-2)(m-4)}{m+2} = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

We have

$$mn + 2n - 2 - \frac{2n^2(m-2)(m-4)}{m+2} = -\frac{4 + 2m - m^2n - 4n - 4mn + 16n^2 - 12mn^2 + 2m^2n^2}{m+2}$$
$$= -f_2(m,n),$$

where  $f_2(m,n) = \frac{mn^2(m-12) + m^2n(n-2) + mn(m-6) + 2n(m-2) + 16n^2 + 2m + 4}{m+2}$ . For m=8 and n=2 we have  $f_2(m,n) = \frac{6}{5} > 0$ . For m=8 and  $n \geq 3$  we have  $f_2(m,n) = \frac{6}{5} > 0$ .  $\frac{2}{5}(12n^2-25n+5)=\frac{2}{5}(12n(n-3)+11n+5)>0$ . For m=10 and  $n\geq 2$  we have  $f_2(m,n)=1$  $2(4n^2-6n+1)=2(4n(n-2)+2n+1)>0$ . For  $m \ge 12$  and  $n \ge 2$  we have  $f_2(m,n)>0$ . Therefore,

$$LE^{+}(K_{|V(\mathcal{CCC}(G))|}) = mn + 2n - 2 < \frac{2n^{2}(m-2)(m-4)}{m+2} = LE^{+}(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$$

and so  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither L-borderenergetic nor Qborderenergetic. Thus, if m > 8 and n > 2 then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

Theorem 5.4. Let  $G = V_{8n}$ .

- (a) If n=2 then  $\mathcal{CCC}(G)$  is neither hyperenergetic, border-energetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.
- (b) If n = 3, 4 then  $\mathcal{CCC}(G)$  is L-hyperenergetic but neither hyperenergetic, borderenergetic, Lborderenergetic, Q-hyperenergetic nor Q-borderenergetic.
- (c) If n > 5 then CCC(G) is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic. borderenergetic, L-borderenergetic nor Q-borderenergetic.

*Proof.* We shall prove the result by considering the following cases.

Case 1. n is odd.

By [20, Proposition 2.4] we have  $\mathcal{CCC}(G) = 2K_1 \sqcup K_{2n-1}$ . Therefore,  $|V(\mathcal{CCC}(G))| = 2n + 1$ . Using (5.1), we get

$$(5.18) E(K_{|V(\mathcal{CCC}(G))|}) = LE^+(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = 4n.$$

By Theorem 3.4 we get

$$LE(\mathcal{CCC}(G)) = \frac{6(2n-1)(2n-2)}{2n+1}.$$

We have

$$4n - \frac{6(2n-1)(2n-2)}{2n+1} = -\frac{4(4n^2 - 10n + 3)}{2n+1} = -\frac{4(4n(n-3) + 2n + 3)}{2n+1} < 0$$

and so  $LE(K_{|V(\mathcal{CCC}(G))|}) = 4n < \frac{6(2n-1)(2n-2)}{2n+1} = LE(\mathcal{CCC}(G))$ . Hence,  $\mathcal{CCC}(G)$  is L-hyperenergetic but not L-border energetic.

By Theorem 4.4 and Theorem 3.4, we also get

$$E(\mathcal{CCC}(G)) < LE^{+}(\mathcal{CCC}(G)) = \frac{4(2n-1)(2n-2)}{2n+1}.$$

We have

(5.19) 
$$\frac{4(2n-1)(2n-2)}{2n+1} - 4n = \frac{4(2n^2 - 7n + 2)}{2n+1} = \frac{4(2n(n-5) + 3n + 2)}{2n+1} := g_1(n).$$

Therefore, for n = 3, we have  $g_1(n) < 0$  and so

$$E(\mathcal{CCC}(G)) < LE^{+}(\mathcal{CCC}(G)) = \frac{4(2n-1)(2n-2)}{2n+1} < 4n = LE^{+}(K_{|V(\mathcal{CCC}(G))|}).$$

Thus, if n=3 then  $\mathcal{CCC}(G)$  is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. If  $n \geq 5$  then, by (5.19), we have  $g_1(n) > 0$  and so  $LE^+(\mathcal{CCC}(G)) > 4n = LE^+(K_{|V(\mathcal{CCC}(G))|})$ . Therefore,  $\mathcal{CCC}(G)$  is Q-hyperenergetic but not Q-borderenergetic. Also,  $E(\mathcal{CCC}(G)) = 4n - 4 < 4n = E(K_{|V(\mathcal{CCC}(G))|})$  and so  $\mathcal{CCC}(G)$  is neither hyperenergetic nor borderenergetic. Thus, if  $n \geq 5$  then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

Case 2. n is even.

By [20, Proposition 2.4] we have  $\mathcal{CCC}(G) = 2K_2 \sqcup K_{2n-2}$ . Therefore  $|V(\mathcal{CCC}(G))| = 2n+2$ . Using (5.1), we get

(5.20) 
$$E(K_{|V(\mathcal{CCC}(G))|}) = LE^{+}(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = 4n + 2.$$

If n = 2 then, by Theorem 4.4 and Theorem 3.4, we get

$$(5.21) E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 6 < 10 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, by (5.20) and (5.21), we have  $\mathcal{CCC}(G)$  is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If  $n \geq 4$  then, Theorem 3.4, we get

$$E(\mathcal{CCC}(G)) = 4n - 2 < 4n + 2 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore,  $\mathcal{CCC}(G)$  is neither hyperenergetic nor borderenergetic.

By Theorem 4.4 and Theorem 3.4, we also get

$$\frac{16(n-1)(n-2)}{n+1} = LE^{+}(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

We have

$$(5.22) \frac{16(n-1)(n-2)}{n+1} - (4n+2) = \frac{6(2n^2 - 9n + 5)}{n+1} = \frac{6(2n(n-6) + 3n + 5)}{n+1} := g_2(n).$$

Therefore, for n=4 we have  $g_2(n)<0$  and so  $LE^+(\mathcal{CCC}(G))=\frac{16(n-1)(n-2)}{n+1}<4n+2=LE^+(K_{|V(\mathcal{CCC}(G))|})$ . Therefore,  $\mathcal{CCC}(G)$  is neither Q-hyperenergetic nor Q-borderenergetic. Also,

$$LE(\mathcal{CCC}(G)) = \frac{130}{5} = 26 > 18 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore,  $\mathcal{CCC}(G)$  is L-hyperenergetic but not L-borderenergetic. Thus, if n=4 then  $\mathcal{CCC}(G)$  is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If  $n \ge 6$  then, by (5.22), we have  $g_2(n) > 0$  and so

$$LE^{+}(K_{|V(\mathcal{CCC}(G))|}) = 4n + 2 < \frac{16(n-1)(n-2)}{n+1} = LE^{+}(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

Therefore,  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither L-borderenergetic nor Q-borderenergetic. Thus, if  $n \geq 6$  then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

Theorem 5.5. Let  $G = SD_{8n}$ .

- (a) If n = 2, 3 then CCC(G) is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-hyperenergetic, Q-hyperenergetic nor Q-borderenergetic.
- (b) If n = 5 then CCC(G) is L-hyperenergetic and Q-borderenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-hyperenergetic.
- (c) If n = 4 or  $n \ge 6$  then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

*Proof.* We shall prove the result by considering the following cases.

Case 1. n is odd.

By [20, Proposition 2.5] we have  $\mathcal{CCC}(G) = K_4 \sqcup K_{2n-2}$ . Therefore,  $|V(\mathcal{CCC}(G))| = 2n + 2$ . Using (5.1), we get

(5.23) 
$$E(K_{|V(\mathcal{CCC}(G))|}) = LE^{+}(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = 4n + 2.$$

By Theorem 3.5 we get

$$E(\mathcal{CCC}(G)) = 4n < 4n + 2.$$

Therefore,  $\mathcal{CCC}(G)$  is neither hyperenergetic nor borderenergetic.

If n = 3 then, by Theorem 3.5, we get

$$LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 12 < 14 = LE(K_{|V(\mathcal{CCC}(G))|})$$

So,  $\mathcal{CCC}(G)$  is neither L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if n=3 then  $\mathcal{CCC}(G)$  is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If n = 5 then, by Theorem 4.5 and Theorem 3.5, we get

$$LE^+(K_{|V(\mathcal{CCC}(G))|}) = 22 = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

Therefore,  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-borderenergetic but neither L-borderenergetic nor Q-hyperenergetic. Thus, if n=5 then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-borderenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-hyperenergetic.

If  $n \geq 7$  then, by Theorem 4.5 and Theorem 3.5, we get

$$\frac{16(n-1)(n-3)}{n+1} = LE^{+}(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

We have

$$4n+2-\frac{16(n-1)(n-3)}{n+1}=-\frac{2(6n^2-35n+23)}{n+1}=-\frac{2(6n(n-7)+7n+23)}{n+1}<0.$$

So,  $LE^+(K_{|V(\mathcal{CCC}(G))|}) = 4n + 2 < \frac{16(n-1)(n-3)}{n+1} = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$  and so  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither L-border energetic nor Q-border energetic. Thus, if  $n \geq 7$  then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

Case 2. n is even.

By [20, Proposition 2.5] we have  $\mathcal{CCC}(G) = 2K_1 \sqcup K_{2n-1}$ . Therefore,  $|V(\mathcal{CCC}(G))| = 2n + 1$ . Using (5.1), we get

$$(5.24) E(K_{|V(\mathcal{CCC}(G))|}) = LE^+(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = 4n.$$

By Theorem 3.5 we get

$$E(\mathcal{CCC}(G)) = 4n - 4 < 4n.$$

Therefore,  $\mathcal{CCC}(G)$  is neither hyperenergetic nor borderenergetic.

If n = 2 then, by Theorem 4.5 and Theorem 3.5, we get

$$LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)) = \frac{36}{5} < 8 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

So, CCC(G) is neither L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if n = 2 then CCC(G) is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If  $n \ge 4$  then, by Theorem 4.5 and Theorem 3.5, we get

$$\frac{4(2n-1)(2n-2)}{2n+1} = LE^{+}(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

We have

$$4n - \frac{4(2n-1)(2n-2)}{2n+1} = -\frac{4(2n^2 - 7n + 2)}{2n+1} = -\frac{4(2n(n-4) + n + 2)}{2n+1} < 0.$$

Therefore,  $LE^+(K_{|V(\mathcal{CCC}(G))|})=4n<\frac{4(2n-1)(2n-2)}{2n+1}=LE^+(\mathcal{CCC}(G))< LE(\mathcal{CCC}(G))$  and so  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither L-borderenergetic nor Q-borderenergetic. Thus, if  $n\geq 4$  then  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

We conclude this paper with the following characterization of commuting conjugacy class graph.

## **Theorem 5.6.** Let G be a finite non-abelian group. Then

- (a)  $\mathcal{CCC}(G)$  is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic if G is isomorphic to  $D_8$ ,  $D_{12}$ ,  $D_{2n}(n$  is odd),  $Q_8$ ,  $Q_{12}$ ,  $Q_{16}$ ,  $U_{(2,6)}$ ,  $U_{(n,2)}$ ,  $U_{(n,3)}$ ,  $U_{(n,4)}$   $(n \geq 2)$ ,  $V_{16}$ ,  $SD_{16}$  or  $SD_{24}$ .
- (b) CCC(G) is L-borderenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic if G is isomorphic to  $Q_{20}$  or  $U_{(2.5)}$ .
- (c)  $\mathcal{CCC}(G)$  is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic if G is isomorphic to  $D_{16}, D_{20}, D_{24}, D_{28}, Q_{24}, Q_{28}, U_{(3,5)}, U_{(3,6)}, U_{(2,7)}, V_{24}$  or  $V_{32}$ .
- (d) CCC(G) is L-hyperenergetic and Q-borderenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-hyperenergetic if G is isomorphic to  $SD_{40}$ .
- (e)  $\mathcal{CCC}(G)$  is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic if G is isomorphic to  $D_{2n}(n \text{ is even}, n \geq 16), Q_{4m}(m \geq 8), U_{(n,5)}(n \geq 4), U_{(n,6)}(n \geq 4), U_{(n,7)}(n \geq 3), U_{(n,m)}(n \geq 2 \text{ and } m \geq 8), V_{8n}(n \geq 5), SD_{32} \text{ or } SD_{8n}(n \geq 6).$

## **Theorem 5.7.** Let G be a finite non-abelian group. Then

- (a) If G is isomorphic to  $D_{2n}$ ,  $Q_{4m}$ ,  $U_{(n,m)}$ ,  $V_{8n}$  or  $SD_{8n}$  then CCC(G) is neither hyperenergetic nor borderenergetic.
- (b) If G is isomorphic to  $D_{2n}(n \text{ is even}, n \geq 8), Q_{4m}(m \geq 6), U_{(n,5)}(n \geq 3), U_{(n,6)}(n \geq 3), U_{(n,m)}(n \geq 2 \text{ and } m \geq 7), V_{8n}(n \geq 3) \text{ or } SD_{8n}(n \geq 4) \text{ then } CCC(G) \text{ is L-hyperenergetic}$
- (c) If G is isomorphic to  $Q_{20}$  or  $U_{(2.5)}$  then CCC(G) is L-borderenergetic.

- (d) If G is isomorphic to  $D_{2n}(n \text{ is even}, n \geq 16), Q_{4m}(m \geq 8), U_{(n,5)}(n \geq 4), U_{(n,6)}(n \geq 4), U_{(n,7)}(n \geq 3), U_{(n,m)}(n \geq 2 \text{ and } m \geq 8), V_{8n}(n \geq 5), SD_{32} \text{ or } SD_{8n}(n \geq 6) \text{ then } CCC(G) \text{ is } Q-hyperenergetic.}$
- (e) If G is isomorphic to  $SD_{40}$  then CCC(G) is Q-borderenergetic.

**Acknowledgements** The first author is thankful to Council of Scientific and Industrial Research for the fellowship (File No. 09/796(0094)/2019-EMR-I).

#### References

- [1] N. M. M. Abreu, C. T. M. Vinagre, A. S. Bonifácioa and I. Gutman, The Laplacian energy of some Laplacian integral graph, *MATCH Commun. Math. Comput. Chem.*, **60**, 447–460 (2008)
- [2] K. Balińska, D. Cvetković, Z. Radosavljević, S. Simić and D. Stevanović, A survey on integral graphs, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat., 13, 42–65 (2003).
- [3] R. Brauer and K. A. Fowler, On groups of even order, Ann. of Math. 62(2) 565-583 (1955).
- [4] D. Cvetković, P. Rowlinson, S. Simić, Signless Laplacian of finite graphs, Linear Algebra Appl., 423, 155–171 (2007).
- [5] P. Dutta, B. Bagchi and R. K. Nath, Various energies of commuting graphs of finite nonabelian groups, Khayyam J. Math., 6(1), 27-45 (2020).
- [6] J. Dutta and R. K. Nath, Spectrum of commuting graphs of some classes of finite groups, Matematika, 33(1), 87–95 (2017).
- [7] J. Dutta and R. K. Nath, Finite groups whose commuting graphs are integral, Matematicki Vesnik, 69(3), 226–230 (2017).
- [8] J. Dutta and R. K. Nath, Laplacian and signless Laplacian spectrum of commuting graphs of finite groups, Khayyam J. Math., 4(1), 77–87 (2018).
- [9] W. N. T. Fasfous, R. K. Nath and R. Sharafdini, Various spectra and energies of commuting graphs of finite rings, *Hacettepe Journal of Mathematics and Statistics*, accepted for publication.
- [10] S.C. Gong, X. Li, G.H. Xu, I. Gutman and B. Furtula, Borderenergetic graphs, MATCH Commun. Math. Comput. Chem. 74, 321-332, 2015.
- [11] I. Gutman, Hyperenergetic molecular graphs, J. Serb. Chem. Soc. 64, 199-205, 1999.
- [12] I. Gutman, N. M. M. Abreu, , C. T. M. Vinagre, A. S. Bonifácioa and S. Radenković, Relation between energy and Laplacian energy. MATCH Communications in Mathematical and in Computer Chemistry, 59, 343–354, (2008).
- [13] F. Harary and A. J. Schwenk, Which graphs have integral spectra?, *Graphs and Combin.*, Lect. Notes Math., Vol 406, Springer-Verlag, Berlin, 45–51 (1974).
- [14] M. Herzog, M. Longobardi and M. Maj, On a commuting graph on conjugacy classes of groups. Communications in Algebra, 37(10), 3369–3387, (2009).
- [15] S. Kirkland, Constructably Laplacian integral graphs, Linear Algebra Appl., 423, 3-21 (2007).
- [16] J. Liu, and B. Liu, On the relation between energy and Laplacian energy. MATCH Communications in Mathematical and in Computer Chemistry, 61, 403–406 (2009).
- [17] R. Merris, Degree maximal graphs are Laplacian integral, Linear Algebra Appl., 199, 381–389 (1994).
- [18] A. Mohammadian, A. Erfanian, D. G. M. Farrokhi and B. Wilkens, Triangle-free commuting conjugacy class graphs. *Journal of Group Theory*, 19, 1049–1061, (2016).
- [19] R. K. Nath, Various spectra of commuting graphs n-centralizer finite groups, International Journal of Engineering Science and Technology, accepted for publication.
- [20] M. A. Salahshour and A. R. Ashrafi, Commuting conjugacy class graph of finite CA-groups, Khayyam J. Math., 6(1), 108–118.
- [21] S. K. Simić and Z. Stanić, Q-integral graphs with edge-degrees at most five, Discrete Math., 308, 4625–4634 (2008).
- [22] D. Stevanović, I. Stanković, and M. Milošević, More on the relation between energy and Laplacian energy. MATCH Communications in Mathematical and in Computer Chemistry, 61, 395–401 (2008).
- [23] F. Tura, L-borderenergetic graphs, MATCH Commun. Math. Comput. Chem. 77, 37-44, 2017.
- [24] H.B. Walikar, H.S. Ramane and P.R. Hampiholi, On the energy of a graph, Graph Connections, Allied Publishers, New Delhi, 1999, pp. 120-123, Eds. R. Balakrishnan, H. M. Mulder, A. Vijayakumar.

DEPARTMENT OF MATHEMATICAL SCIENCES, TEZPUR UNIVERSITY, NAPAAM-784028, SONITPUR, ASSAM, INDIA. E-mail address: bhowal.parthajit8@gmail.com, rajatkantinath@yahoo.com (correcponding author)