

Dimension Independent Generalization Error by Stochastic Gradient Descent

Xi Chen* Qiang Liu[†] Xin T. Tong[‡]

January 5, 2021

Abstract

One classical canon of statistics is that large models are prone to overfitting, and model selection procedures are necessary for high dimensional data. However, many overparameterized models, such as neural networks, perform very well in practice, although they are often trained with simple online methods and regularization. The empirical success of overparameterized models, which is often known as benign overfitting, motivates us to have a new look at the statistical generalization theory for online optimization. In particular, we present a general theory on the generalization error of stochastic gradient descent (SGD) solutions for both convex and locally convex loss functions. We further discuss data and model conditions that lead to a “low effective dimension”. Under these conditions, we show that the generalization error either does not depend on the ambient dimension p or depends on p via a poly-logarithmic factor. We also demonstrate that in several widely used statistical models, the “low effective dimension” arises naturally in overparameterized settings. The studied statistical applications include both convex models such as linear regression and logistic regression and non-convex models such as M -estimator and two-layer neural networks.

1 Introduction

The study of overfitting phenomenon has been an important topic in statistics and machine learning. From classical statistical learning theory, we

*Stern School of Business, New York University, Email: xichen@nyu.edu

[†]Department of Mathematics, National University of Singapore, Email: matliuq@nus.edu.sg

[‡]Department of Mathematics, National University of Singapore, Email: mat-txin@nus.edu.sg

understand that when the number of model parameters is large compared to the amount of data, the generalization error can be excessively large even if the training error is small. This phenomenon is usually known as overfitting. For this reason, dimension reduction or feature selection mechanisms such as principle component analysis (PCA) and shrinkage methods are often required in the training phase to reduce model dimension and avoid overfitting.

In recent years, deep neural networks have achieved great successes in practical applications. Researchers have found out that overparameterized neural networks usually achieve superior performance (Golowich et al., 2018; Li and Liang, 2018; Neyshabur et al., 2019; Allen-Zhu et al., 2019; Arora et al., 2019). Moreover, these models are often trained with simple regularization and do not need dimension reduction procedures. This phenomenon is sometimes referred to as *benign overfitting* (Bartlett et al., 2019). To understand it, we need a new statistical framework to study generalization errors.

Although there is much practical evidence on the benefit of overparameterization, the existing theoretical study mainly focuses on linear models (see, e.g., Bartlett et al., 2019; Nakkiran, 2019; Ali et al., 2019) or neural networks with certain special data structures (see, e.g., Li and Liang, 2018). The main purpose of our paper is to systematically investigate the generalization error for a risk minimization problem when the number of parameters p is much larger than the sample size N . In particular, we establish a generalization error bound for stochastic gradient descent (SGD) solutions for both convex (e.g., linear regression and logistic regression) and non-convex problems (some M -estimators and neural networks). We focus our study on the SGD algorithm because it has been widely used in large-scale data learning due to its computational and memory efficiency.

Let us briefly introduce our setup of the overfitting problem and the SGD algorithm. We consider the following population risk minimization problem under a loss function F , which can be either convex or non-convex:

$$w^* = \operatorname{argmin}_w F(w), \quad F(w) := \mathbb{E}_\zeta f(w, \zeta). \quad (1)$$

In (1), $w \in \mathbb{R}^p$ is a p -dimensional parameter vector, ζ denotes a random sample from a certain probability distribution, and $f(\cdot, \zeta)$ is the loss function on each individual data ζ . The global minimizer w^* is often the true model parameter in statistical estimation problems. In practice, the distribution of ζ is usually unknown, and one only has the access to N *i.i.d.* samples ζ_1, \dots, ζ_N from the population. Instead of minimizing the population risk

$F(w)$ in (1), it is more practical to minimize the empirical loss function

$$\hat{F}(w) = \frac{1}{N} \sum_{i=1}^N f(w, \zeta_i). \quad (2)$$

Often, instead of directly minimizing the empirical loss, an extra regularization term is sometimes added to the empirical loss to avoid overfitting. In this paper, we consider the most commonly used ridge or Tikhonov regularization. The corresponding regularized empirical loss function takes the following form,

$$\hat{F}_\lambda(w) := \hat{F}(w) + \frac{\lambda}{2} \|w\|^2 = \frac{1}{N} \sum_{i=1}^N f_\lambda(w, \zeta_i), \quad f_\lambda(w, \zeta) := f(w, \zeta) + \frac{\lambda}{2} \|w\|^2. \quad (3)$$

The weight of regularization is controlled by λ , which is a tuning parameter. When $\lambda = 0$, this corresponds to the ridgeless regression. We will study this setup in this paper, which corresponds to “implicit regularization”. One popular way to optimize \hat{F}_λ in machine learning is via SGD. In particular, for a generic initialization parameter w_0 , SGD is an iterative algorithm, where the $(n + 1)$ -th iterate w_{n+1} is updated according to the following equation,

$$w_{n+1} := w_n - \eta \nabla f_\lambda(w_n, \zeta_n) = w_n - \eta (\nabla f(w_n, \zeta_n) + \lambda w_n). \quad (4)$$

By running through N samples, SGD outputs the N -th iterate w_N as the final estimator of w^* . Notably, SGD iterates are affected by the stochasticity of the data samples ζ . To reduce such noise and improve accuracy, the averaged SGD (ASGD) method uses the average iterate

$$\bar{w}_N = \frac{1}{N} \sum_{i=1}^N w_i$$

as the final estimator of w^* . In SGD iterations (4), the hyper-parameter η is known as the stepsize. In our paper, we consider using a constant stepsize, which is a popular choice in practice (Bach and Moulines, 2011). The value of η will be discussed later in our theoretical results. Moreover, in (4), the gradient is taken with respect to the parameter vector w . For notational simplicity, we will use “ ∇ ” as a short notation for “ ∇_w ” throughout the paper.

When the sample size N is much larger than the dimensionality p , it is expected that w_N would be close to w^* under certain conditions. However,

in an overparameterized setting where N is less than p , the solution w_N can be far away from w^* . In this case, estimating the underlying parameter accurately usually requires strong assumptions. However, for many machine learning tasks, it is of more interest in achieving small generalization error, which is defined as follows,

$$G(w_N) = F(w_N) - F(w^*). \quad (5)$$

The main purpose of the paper is to provide an upper bound of the generalization error in (5) in overparameterized settings. We will characterize the scenarios where such a generalization error bound is independent of p or only involves in poly-logarithmic factors of p .

1.1 Main results and paper organization

The main message of this paper is as follows. For a large class of statistical learning problems where the *effective dimension* is low (see the rigorous definition in Section 3), the stochastic gradient descent (SGD) algorithm with proper ridge regularization will not overfit even if the ambient model dimension is much larger than the sample size. In particular, we will show that the generalization error has at most poly-logarithmic dependence on the ambient model dimension p .

In Section 2, we present a framework for generalization error analysis. This framework is designed to handle high dimensional regression problems. We will separately discuss two scenarios. In the first scenario, the true parameter is weakly sparse with a dimension independent l_2 norm. In this case, we show ridgeless SGD has been sufficient to obtain dimension independent generalization error (Theorem 2.3). In the second scenario, the true parameter is dimension independent only under some problem-specific norms. In this more challenging case, we show SGD can achieve dimension independent generalization error by proper amount of ridge penalty (Theorem 2.4). The upper bound of the generalization error is provided. Using linear regression as an illustrative example, we show that each term in the generalization bound has a strong statistical interpretation (see Section 2.5). The upper bound also leads to practical guidelines on the rates of problem-related parameters, which are given by Corollary 2.6.

While Theorem 2.4 provides an error bound on generalization, for this error bound to be almost dimension-independent, we require the *effective dimension* to be small. Section 3 first formally defines the general notion of *low effective dimension*, which can essentially be described by 1) the loss

function has a fast decaying Hessian spectrum, and 2) the true parameter is either sparse or uniformly bounded along Hessian’s eigen-directions.

In Section 4, we carefully investigate generalization errors in various linear models. We consider the cases of finite projections of infinite-dimensional models and linear regression with redundant features. In these scenarios, we quantify when the overparameterization does not hurt the generalization performance.

Our generalization result can also be applied to a wide range of nonlinear models. In Section 5, we study both convex nonlinear models such as logistic regression and non-convex models such as M -estimator with the Tukey’s biweight loss function (Tukey, 1960) and two-layer neural networks. We show that the low effective dimension naturally occurs in these applications.

1.2 Related works

High dimensional statistical learning with low effective dimension is a common idea in principal component analysis (PCA) and functional data analysis applications (Jolliffe, 2002; Ramsay and Silverman, 2005). To implement PCA, we first find a k^* -dimensional subspace of which the truncated variability contains a significant proportion (e.g., 95%) of the full model variability. Then we project the problem onto this subspace and conduct the data analysis on it. In general, the truncated dimension k^* needs to be less than the sample size N , and the statistical problem needs to be well-conditioned in the k^* -dimensional subspace. While we also require the existence of a low effective dimension, the problem setting considered in this paper is fundamentally different from the classical PCA problem. In our setting, the SGD algorithm directly runs on the original problem space of dimension p , and there is no projection step. Moreover, we do not need the problem to be well-conditioned in the ambient space. We would argue that our setting is more useful for practical applications since it is difficult to find a proper truncation dimension k^* and PCA is intractable with rank-deficient data.

In recent years, the generalization error bounds for linear models in overparameterized settings have been carefully investigated in the literature, see, e.g., Bartlett et al. (2019); Nakkiran (2019); Ali et al. (2019); Hastie et al. (2019). Our result is different from these existing results in the following perspectives:

- 1) Nonlinearity: Most of these works focus on linear models. For example, Bartlett et al. (2019) defined two notions of *effective ranks* of the data covariance matrix in linear models and expressed the generalization error

bound in terms of effective ranks. A more detailed comparison between our low effective rank conditions and the ones in [Bartlett et al. \(2019\)](#) will be provided in Remark 3.5. In addition, [Hastie et al. \(2019\)](#) developed the generalization error bound for the composition of an activation function and a linear model. Our work also requires conditions similar to small effective ranks, but they are simpler in formulation, as discussed in Section 3. Moreover, our results can be applied to general nonlinear models.

- 2) Anisotropic spectrum and regularization: [Nakkiran \(2019\)](#) and [Hastie et al. \(2019\)](#) focus their studies on cases with isotropic or well-conditioned regressor covariance matrices. Our study focuses on cases with anisotropic regressor covariance, of which the minimum eigenvalue decays to zero. Moreover, since online learning has the implicit regularization effect and the ridge penalty is applied in certain scenarios, we do not have the “double descent” phenomenon as in other literature ([Belkin et al., 2018, 2019](#)). A recent paper by [Nakkiran et al. \(2020\)](#) also showed that for certain linear regression models with isotropic data distribution, the ridge penalty regularized regression (in the offline setting) can avoid the “double descent” phenomenon.
- 3) Online optimization: The aforementioned works mainly focus on offline optimization. For example, [Bartlett et al. \(2019\)](#), [Nakkiran \(2019\)](#), and [Hastie et al. \(2019\)](#) show that the generalization error of the offline linear regression solution is closely related to the spectrum of the design matrix. In particular, when the minimum eigenvalue is close to zero, the offline learning results become unstable because of singular matrix-inversions. Since the design matrix relies on data realization, such instability can only be studied through random matrix theories (RMT). While these studies of offline linear regression are interesting and technically deep, their dependence on RMT makes the extensions to nonlinear models difficult. In comparison, online learning methods process one data point at a time and do not involve the inversion of design matrices, which facilitates the study of general nonlinear models. Moreover, in terms of practical applications, online optimization methods such as SGD are appealing due to their low per-iteration complexity compared to offline optimization. Therefore, this paper focuses on the generalization error for online learning in overparameterized settings.

For online optimization, the stochastic gradient descent (SGD), which dates back to [Robbins and Monro \(1951\)](#), is perhaps the most widely used

method in practice. The convergence rates of the SGD for convex loss functions have been well studied in the literature (see, e.g., [Zhang, 2004](#), [Nesterov and Vial, 2008](#), [Bach and Moulines, 2011](#), [Ghadimi and Lan, 2012](#), [Roux et al., 2012](#), [Bach and Moulines, 2013](#)). For the constant stepsize SGD, [Bach and Moulines \(2011\)](#) provided the generalization error bound in Theorem 1 of their paper. However, their bound has an explicit linear dependence on p , which is not applicable to the overparameterized setting. Our paper provides a more refined generalization error bound, which incorporates the Hessian spectrum to capture the “low dimension effect” in the overparameterized setting. When ridge regularization is presented, existing results show that the generalization error is dimension independent ([Shalev-Shwartz and Ben-David, 2014](#); [Zhang, 2005](#); [Carratino et al., 2018](#)). But, in general, these results need the norm of population gradient ∇F to be dimension independent (See Section 14.5.3 in [Shalev-Shwartz and Ben-David \(2014\)](#)). Our new results do not have this restriction but only require stochastic gradient variance to be independent of the dimension. This is a much less stringent restriction because the variance can be reduced by various techniques (see, e.g., [Johnson and Zhang \(2013\)](#)).

Overparameterized neural network (NN) is a very active research direction. There are several existing works explaining why overfitting does not happen in large NN ([Golowich et al., 2018](#); [Li and Liang, 2018](#); [Neyshabur et al., 2019](#); [Allen-Zhu et al., 2019](#); [Arora et al., 2019](#); [E et al., 2019a,b](#)). Interestingly, the conditions they impose are largely similar to the ones we will use. Namely, they require the high dimensional input data and the Frobenius norm of true weight matrices to be bounded by constants. For this to be true, only a small portion of the data or model components can be significantly active, which satisfies the concept of low effective dimension.

On the other hand, our study of two-layer NN in Section 5.3 is different from existing results in the following perspectives. One popular way to analyze generalization error is by analyzing the Rademacher complexity ([Golowich et al., 2018](#); [Neyshabur et al., 2019](#); [E et al., 2019a](#)), which can be used to establish an upper bound of the difference between training error and generalization error (see, e.g., Theorem 3.3 of [Mohri et al. \(2018\)](#)). However, these results are usually derived in an offline optimization setting, while our paper focuses on the result from online SGD. [Li and Liang \(2018\)](#) and [Allen-Zhu et al. \(2019\)](#) both studied NN generalization error with SGD iterations. But they mainly focused on classification scenarios where the loss function is bounded. Moreover, [Li and Liang \(2018\)](#) required the loss function to be of a logistic form, and [Allen-Zhu et al. \(2019\)](#) studied the running average generalization error. Our results can be applied to regres-

sion NN with unbounded loss functions. Arora et al. (2019) and E et al. (2019b) studied NN generalization bound with deterministic gradient descent. Moreover, their studies assume a certain data angle or Gram matrix to have a strictly positive minimal eigenvalue. Please see Section 5.3 for more detailed comparisons with these works.

2 A Generalization Bound

In this section, we present a general result on the generalization bound for the SGD solution from (4).

2.1 Preliminaries: high dimensional norms and non-convex energy landscape

The main issue this paper tries to understand is the high dimensional generalization error when using SGD as the training method. In some simple scenarios, the true model parameters are “sparse”, so $\|w^*\|$ does not grow with the dimension. This allows us to measure the distance between SGD iterate w_n and w^* directly. In other scenarios, w^* can be very dense, and $\|w^*\|$ may grow linearly or even faster with the dimension. To resolve this issue, we define the following norms:

Definition 2.1. *Given a matrix $A \in \mathbb{R}^{p \times p}$ and λ , we decompose $\mathbb{R}^p = S_\lambda \oplus S_\perp$, where S_λ consists of eigenvectors of A with eigenvalues above $\lambda > 0$ and S_\perp is the orthogonal complement of S_λ . Given any vector v , denote its decomposition as $v = v_\lambda + v_\perp$, where $v_\lambda \in S_\lambda$ and $v_\perp \in S_\perp$. Then define*

$$\|v\|_A^2 = v^T A v, \quad \|v\|_{A,\lambda}^2 := \lambda \|v_\lambda\|^2 + v_\perp^T A v_\perp. \quad (6)$$

We introduce the norm $\|v\|_A^2$, whose value can be independent of the ambient dimension p . The second norm $\|v\|_{A,\lambda}^2$ is a truncated version of the first norm, it essentially truncates all eigenvalues of A above λ to λ . It is easy to see that $\|v\|_{A,\lambda}^2 \leq \|v\|_A^2$. We introduce the second norm because $\|v\|_{A,\lambda}^2$ converges to zero when the regularization parameter λ does, while $\|v\|_A^2$ is independent of λ .

It is well known that SGD works well for convex problems. This is also true for our theoretical analysis, which works best in convex settings. On the other hand, many practical problems are non-convex, which makes the statistical learning problem technically more challenging. First, the function F can have multiple local minima, and each local minimum has an attraction

basin, which is a “valley” in the graph of F . Within each valley, we assume that F is locally convex. Machine learning and theoretical deep learning literature often study the scenarios that local minima lie in large and shallow valleys, which lead to more stable generalization performance. Suppose \mathcal{D} is the attraction basin of the optimal solution w^* in (1), initializing SGD in \mathcal{D} will generate iterates converging to w^* with high probability. So a natural question is the gap between the estimator learned from SGD and w^* . On the other hand, if the SGD iterates take place outside \mathcal{D} , the output can be irrelevant to the properties of w^* . Therefore, we need to introduce a stopping time τ , which describes the first time SGD exits \mathcal{D} :

$$\tau = \min\{n : w_n \notin \mathcal{D}\},$$

where w_n is the n -th SGD iterate defined in (4). Our generalization analysis will assume the SGD iterates stay within \mathcal{D} . We will also provide bounds of probability that SGD leaves \mathcal{D} .

While it is reasonable to assume that w^* as a local minimum is in the valley \mathcal{D} , verifying this assumption can be difficult. For example, because we do not have access to the population loss function F or the Hessian directly, to check the convexity of F , we need to investigate \hat{F} instead. There will be a certain inaccuracy due to the randomness in \hat{F} . As another example, while we know the Hessian of F is positive semidefinite at w^* because w^* is a local minimum, F does not have to be convex in the neighborhood w^* . For both these examples, it is more accurate to say F is “approximately convex” in \mathcal{D} . Our analysis can also extend to such very challenging cases by introducing proper regularizations. On the other hand, if F is very non-convex in \mathcal{D} , then one should not expect that SGD will produce good learning results. Therefore, to achieve a reasonable generalization error, we need F to be very close to convex. In many applications, this can be done by choosing either a very large sample size N or a very small neighborhood around w^* .

Based on our discussion, we have the following assumption on the population risk function,

Assumption 2.2. *The optimal solution w^* of the population risk F has a neighborhood \mathcal{D} , such that for some positive semidefinite (PSD) matrix A and $\delta \in [0, 1/2)$,*

$$-\delta A \preceq \nabla^2 F(w) \preceq A, \quad \forall w \in \mathcal{D}. \quad (7)$$

In Assumption 2.2 and in the sequel, for two symmetric matrices C, D , $C \preceq D$ indicates that $D - C$ is positive semidefinite (PSD). The upper

bound on the Hessian matrix $\nabla^2 F(w) \preceq A$ is widely assumed in the statistical literature. The parameter δ above describes the level of non-convexity. In particular, $\delta = 0$ indicates that F is convex within \mathcal{D} . However, our condition in (7) is more general since δ can be strictly positive, which allows F to be non-convex. On the other hand, although our generalization error bound holds for $\delta \in [0, 1/2)$, for this upper bound to be smaller than a certain threshold, δ needs to be small. Please see Corollary 2.6 for the exact dependence of δ in the upper bound.

2.2 Generalization bound with sparse true parameter

We will discuss two possible scenarios with high dimensional machine learning. In the first case, the true parameter w^* is sparse or weak so that its l_2 norm does not depend on the dimension. In this case, we have the following results:

Theorem 2.3. *Under Assumption 2.2, suppose $w_0 \in \mathcal{D}$ and there is a constants r and c_r such that*

$$\mathbb{E}\|\nabla f(w, \zeta) - \nabla F(w)\|^2 \leq r^2 + c_r |(w - w^*)^T \nabla F(w)|, \quad \forall w \in \mathcal{D}. \quad (8)$$

Then if the ridgeless SGD stepsize η and the regularization parameter λ satisfy

$$\eta \leq \min \left\{ \frac{1}{4(1 + c_r)(\|A\| + \lambda)}, 1 \right\}, \quad 2\delta\|A\| \leq \lambda \leq 1,$$

we have the generalization error for the averaged SGD,

$$\mathbb{E}[1_{\tau \geq N-1} G(\bar{w}_N)] \leq \frac{2\mathbb{E}\|w_0 - w^*\|^2}{N\eta} + 2\eta r^2 + 8\lambda\|w^*\|^2, \quad (9)$$

where the generalization error G is defined in (5) and $\bar{w}_N = \frac{1}{N} \sum_{i=0}^{N-1} w_i$ is the averaged SGD iterate.

We note that when F is locally convex in \mathcal{D} (i.e., $\delta = 0$ in (7)), this result incorporates the implicit regularization case by setting $\lambda = 0$. The generalization error bound in (9) contains three terms. The first term decays with the sample size N . The second term is controlled by the stepsize η . The last term $8\lambda\|w^*\|^2$ is a bias caused by the regularization. Since F_λ is different from F , the minimizer of F_λ is also different from w^* . In other words, if F is convex in \mathcal{D} (i.e., $\delta = 0$), it is actually better to do ridgeless regression by setting $\lambda = 0$. Note that in this case of implicit regularization, the generalization error bound in (9) is dimensional independent when $\|w^*\|^2$ is dimension independent with the initialization $w_0 = \mathbf{0}$.

2.3 Generalization bound with non-sparse true parameter

In some scenarios, $\|w^*\|$ may grow with the dimension. While the results in Theorem 2.3 still hold, the estimate (9) is no longer dimension independent. In this case, the learning cannot rely only on implicit regularization. We will need to introduce regularization and high dimensional norms as in Definition 2.1 for these problems:

Theorem 2.4. *Under Assumption 2.2, suppose $w_0 \in \mathcal{D}$ and there is a constants r and c_r such that*

$$\mathbb{E}\|\nabla f(w, \zeta) - \nabla F(w)\|^2 \leq r^2 + c_r r^2 \min\{G(w), \|w\|^2\}, \quad \forall w \in \mathcal{D}. \quad (10)$$

Then if the SGD hyper-parameters, the stepsize η and the regularization parameter λ , satisfy

$$\eta \leq \min \left\{ 1, \frac{\lambda}{12\|A\|^2 + 6\lambda^2 + 6c_r r^2}, \frac{1}{12\|A\|}, \frac{\lambda}{6c_r \|A\| r^2} \right\}, \quad 4\delta\|A\| \leq \lambda \leq 1,$$

we have

$$\begin{aligned} \mathbb{E}[G(w_N)1_{\tau \geq N}] &\leq 4\|w^*\|_{A, \lambda}^2 + \frac{C_1}{\lambda}(\eta + \delta) \\ &\quad + \exp(-\frac{1}{4}\lambda N \eta)\mathbb{E}[G(w_0) + 4N\|A\|\|w_0\|^2], \end{aligned} \quad (11)$$

with $C_1 = 60\|A\| (r^2 + \|w^\|_A^2) + 10\|w^*\|_A^2$.*

While it is not completely new that SGD on convex ridge regression has dimension independent generalization error (Shalev-Shwartz and Ben-David, 2014; Zhang, 2005; Carratino et al., 2018), in general, these results need the norm of gradient ∇F to be dimension independent (See Section 14.5.3 in Shalev-Shwartz and Ben-David (2014)). Theorem 2.4 does not have this restriction. In fact, the population gradient ∇F can be unbounded in many applications (e.g., linear regression). In contrast, our assumption for the stochastic gradient is imposed on its variance, see (10). This is a much relaxed assumption because the variance can be reduced by various techniques (e.g., Johnson and Zhang (2013)) or simply by increasing the mini-batch size for stochastic gradient computation.

2.4 SGD configuration with given generalization error target

In Section 3, we will explicitly define the low effective dimension so that C_1 and the upper bound in (11) are independent of dimension p , or depend on

p only via a polynomial logarithmic factor. This differentiates our result from the estimates in existing literature on SGD, e.g., [Bach and Moulines \(2011\)](#). In particular, most existing results do not take the Hessian spectrum into consideration, so they will have linear or stronger dependence on the dimension p ; while our results do not necessarily have such dependence.

In the upper bound (11), each term carries a strong statistical interpretation, which will be illustrated using linear models in the next subsection (see Section 2.5). In particular, the term $\|w\|_{A,\lambda}^2$ in (11) can be interpreted as the bias caused by minimizing F_λ instead of F , it decays with the regularization parameter λ shrinking to zero. The term $\frac{C_1\eta}{\lambda}$ is the variance induced by the SGD algorithm, which increases as λ decreases. This reveals that under our current problem setting, λ controls a bias-variance tradeoff. Ideally, we can choose small λ and stepsize η to make both the bias and variance small. However, this comes with a price. As the convergence rate scales with $\lambda\eta$, so using small λ and η need to be compensated with a large sample size N (i.e., the number of iterations in SGD).

We can quantify these tradeoffs by considering a practical scenario where a generalization error is pre-fixed to be ϵ , then our results provide guidelines on how to tune the parameters:

Corollary 2.5 (Corollary of Theorem 2.3). *Suppose there is an universal constant C_0 such that*

$$\|w^*\|^2, \|w_0 - w^*\|^2 \leq C_0.$$

Given any $\epsilon > 0$, if the regularization parameter $\lambda(\epsilon)$, the stepsize $\eta(\epsilon)$, the non-convexity parameter $\delta(\epsilon)$, and the sample size $N(\epsilon)$ satisfy

$$\begin{aligned} \lambda(\epsilon) &\leq \min \left\{ \frac{\epsilon}{8C_0}, 1 \right\}, & \delta(\epsilon) &\leq \min \left\{ \frac{\epsilon}{16C_0\|A\|}, \frac{1}{2\|A\|} \right\}, \\ \eta(\epsilon) &< \min \left\{ \frac{1}{2(1+c_r)(\|A\| + \lambda(\epsilon))}, \frac{\epsilon}{2r^2}, 1 \right\}, & N(\epsilon) &> \frac{2C_0}{\epsilon\eta(\epsilon)}, \end{aligned} \quad (12)$$

then $\mathbb{E}[G(\bar{w}_N)1_{\tau \geq N}] \leq 3\epsilon$.

Corollary 2.6 (Corollary of Theorem 2.4). *Given any $\epsilon > 0$, if the regularization parameter $\lambda(\epsilon)$, the stepsize $\eta(\epsilon)$, the non-convexity parameter $\delta(\epsilon)$, and the sample size $N(\epsilon)$ satisfy*

$$\begin{aligned} 4\|w^*\|_{A,\lambda(\epsilon)}^2 &< \epsilon, & \delta(\epsilon) &\leq \frac{\lambda(\epsilon)\epsilon}{C_1}, & \eta(\epsilon) &< \frac{\lambda(\epsilon)\epsilon}{C_1}, \\ N(\epsilon) &> \max \left\{ \frac{-4 \log \{\epsilon/2\mathbb{E}[G(w_0)]\}}{\lambda(\epsilon)\eta(\epsilon)}, \frac{-8 \log \{\epsilon\lambda(\epsilon)\eta(\epsilon)/(64\|A\|\mathbb{E}[\|w_0\|^2])\}}{\lambda(\epsilon)\eta(\epsilon)} \right\}, \end{aligned} \quad (13)$$

and the conditions of Theorem 2.4 hold, then $\mathbb{E}[G(w_N)1_{\tau \geq N}] \leq 4\epsilon$.

It is noteworthy that (11) only discusses the scenario where SGD iterates stay in the domain \mathcal{D} . This is necessary since all our conditions are imposed only within \mathcal{D} . Once an SGD iterate leaves \mathcal{D} , there is no particular reason it can get back to \mathcal{D} . Another possible improvement is to find an upper bound for the conditional generalization error $\mathbb{E}[G(w_N)|\tau \geq N]$. But this is not feasible when \mathcal{D} is a general region. For example, if \mathcal{D} is the intersection between any set and $\{w : G(w) \geq G(w_0) - 1\}$, the conditional expectation will be larger than $G(w_0) - 1$, while $\{\tau \geq N\}$ may have a nonzero occurrence probability. In Section 6, we will briefly discuss how to remedy such a restriction in practice through data splitting.

In practice, SGD is often implemented with mini-batch data to reduce the noise within stochastic gradient. With regularization, the mini-batch SGD with batch-size J can be formally written as

$$w_{n+1} := w_n - \frac{1}{J}\eta \sum_{k=Jn+1}^{J(n+1)} (\nabla f(w_n, \zeta_k) + \lambda w_n),$$

where *i.i.d.* data $\{\zeta_{Jn+1}, \dots, \zeta_{J(n+1)}\}$ forms the n -th batch of data. Our results can be extended to mini-batch SGD as well. To see this, we simply let $z_n = \{\zeta_{Jn+1}, \dots, \zeta_{J(n+1)}\}$ and

$$\tilde{f}(w, z_n) = \frac{1}{J} \sum_{k=Jn+1}^{J(n+1)} f(w, \zeta_k).$$

Note that $\mathbb{E}\tilde{f}(w, z_n) = \mathbb{E}f(w, \zeta_i) = F(w)$. It is also straight forward to see that applying our SGD formulation (4) on \tilde{f} with z_n leads to the mini-batch SGD. Applying our results, e.g. Corollaries 2.5 and 2.6, to mini-batch SGD requires simple modifications for only two parameters. First, the variance of stochastic gradient $\nabla \tilde{f}(w, z)$ is only $\frac{1}{J}$ of the variance of $\nabla f(w, \zeta)$, so the parameter r^2 in mini-batch SGD should be $\frac{1}{J}$ of r^2 in the standard SGD. Second, because each iteration of the mini-batch SGD requires J data samples, so the overall sample size should be NJ .

2.5 Statistical interpretation in linear models and bias-variance tradeoff

To facilitate better understanding our results, we will use linear models to illustrate the statistical interpretation of each term in the generalization

upper bound in (11). In linear regression, each *i.i.d.* observation contains a pair of dependent and response variables, $\zeta_i = (x_i, y_i) \in \mathbb{R}^p \times \mathbb{R}$, where y_i is generated by the following model

$$y_i = x_i^T w^* + \xi_i. \quad (14)$$

In (14), w^* is the true regression coefficient, and the noise term ξ_i is independent of x_i with zero mean and a finite variance σ^2 . For the ease of illustration, we assume $x_i \sim \mathcal{N}(0, \Sigma)$. The generalization error of this linear model takes the following form,

$$G(w) = \mathbb{E} \left[\frac{1}{2} (y_i - w^T x_i)^2 - \frac{1}{2} (y_i - (w^*)^T x_i)^2 \right] = \frac{1}{2} (w - w^*)^T \Sigma (w - w^*). \quad (15)$$

From (15), we can see that $G(w)$ has a strong dependence on the structure of Σ . Let us denote the eigenvalues of Σ by $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p$, and the eigenvector corresponding to λ_i by v_i . Then the parameter error, $v_i^T (w - w^*)$, contributes to $G(w)$ via the factor λ_i .

It is well known that SGD can be interpreted as a stochastic approximation of the gradient descent (Robbins and Monro, 1951), namely we can rewrite (4) as

$$w_{n+1} = w_n - \eta \nabla F_\lambda(w_n) + \eta \xi_n, \quad (16)$$

where $\xi_n = -\nabla f(w_n, \zeta_n) + \nabla F(w_n)$ is the noise in stochastic gradient. For the quadratic loss with the ridge penalty F_λ , the SGD iterates in (16) take the following form,

$$w_{n+1} = w_n - \eta \Sigma (w_n - w^*) - \eta \lambda w_n + \eta \xi_n = (I - (\Sigma + \lambda I) \eta) (w_n - w_\lambda^*) + w_\lambda^* + \eta \xi_n, \quad (17)$$

where $w_\lambda^* := (\Sigma + \lambda I)^{-1} \Sigma w^*$ is the minimizer of $F_\lambda(w) = \mathbb{E} \frac{1}{2} (y - w^T x)^2 + \frac{\lambda}{2} \|w\|^2$.

It is easy to see that w_n then follows a vector autoregressive (VAR) model (see, e.g., Chapter 8 of Tsay, 2010). For the ease of discussion, we simply treat ξ_n as $\mathcal{N}(0, \frac{r^2}{p} I)$, so $\mathbb{E} \|\xi_n\|^2 = r^2$ as we assumed in Theorem 2.4. Admittedly, the independent Gaussian assumption here contradicts our anisotropic assumption. We are assuming it here to gain an intuitive understanding. Then the stationary distribution of w_n in (17) is a Gaussian $\mathcal{N}(\mu, V)$. Take expectation and covariance on both sides of (17), and the μ and V take the following form (see Chapter 8.2.2 of Tsay, 2010),

$$\mu = (I - (\Sigma + \lambda I) \eta) (\mu - w_\lambda^*) + w_\lambda^* \Rightarrow \mu = w_\lambda^*,$$

$$V = (I - (\Sigma + \lambda I)\eta)V(I - (\Sigma + \lambda I)\eta) + \frac{\eta^2 r^2}{p}I.$$

When the stepsize $\eta \leq \|\Sigma + \lambda I\|^{-1}$, $(\Sigma + \lambda I)^2\eta \preceq \Sigma + \lambda I$, so

$$V = \frac{r^2\eta}{p}(2(\Sigma + \lambda I) - (\Sigma + \lambda I)^2\eta)^{-1} \preceq \frac{r^2\eta}{p}(\Sigma + \lambda I)^{-1}.$$

These results give us the limiting average generalization error

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbb{E}G(w_n) &= \frac{1}{2}(w_\lambda^* - w^*)^T \Sigma (w_\lambda^* - w^*) + \frac{1}{2}\text{tr}(V\Sigma) \\ &\leq G(w_\lambda^*) + \frac{r^2\eta}{2p}\text{tr}((\Sigma + \lambda I)^{-1}\Sigma). \end{aligned} \quad (18)$$

The first term $G(w_\lambda^*)$ is the bias caused by using regularization. Indeed, the optimizer of F_λ is w_λ^* rather than w^* . Recall that (λ_i, v_i) are the eigenvalues and eigenvectors of Σ . We define $a_i = \langle v_i, w^* \rangle$ and further express $G(w_\lambda^*)$ in (18) as follows,

$$\begin{aligned} G(w_\lambda^*) &= \frac{1}{2}((\Sigma + \lambda I)^{-1}\Sigma w^* - w^*)^T \Sigma ((\Sigma + \lambda I)^{-1}\Sigma w^* - w^*) \\ &= \frac{1}{2}\lambda^2 (w^*)^T (\Sigma + \lambda I)^{-1} \Sigma (\Sigma + \lambda I)^{-1} w^* = \frac{1}{2} \sum_{i=1}^p \frac{\lambda^2 \lambda_i a_i^2}{(\lambda + \lambda_i)^2}. \end{aligned}$$

Note that when $\lambda_i \geq 0$, $\frac{\lambda^2 \lambda_i}{(\lambda_i + \lambda)^2} \leq \frac{\lambda^2 \lambda_i}{\lambda^2} = \lambda_i$, and by Young's inequality $\frac{\lambda^2 \lambda_i}{(\lambda_i + \lambda)^2} \leq \frac{\lambda^2 \lambda_i}{4\lambda_i \lambda} \leq \lambda$. Therefore, we have the following upper bound of $G(w_\lambda^*)$

$$G(w_\lambda^*) \leq \frac{1}{2} \sum_{i=1}^p (\lambda \wedge \lambda_i) a_i^2 = \frac{1}{2} \|w^*\|_{\Sigma, \lambda}^2. \quad (19)$$

The upper bound in (19) is essentially the first term in (11) by noticing that $\Sigma = A = \nabla^2 F(w)$ in linear regression, which gives an upper bound for the bias.

For the second variance term in (18),

$$\text{var}(\lambda) := \frac{r^2\eta}{2p} \text{tr}((\Sigma + \lambda I)^{-1}\Sigma) = \frac{r^2\eta}{2p} \sum_{i=1}^p \frac{\lambda_i}{\lambda_i + \lambda} \leq \frac{r^2\eta}{2p} \sum_{i=1}^p \frac{\lambda_i}{\lambda} = \frac{\eta r^2 \lambda_1}{2\lambda}. \quad (20)$$

This upper bound is essentially the second term in the generalization upper bound in (11), as it depends linearly on $\eta, \lambda^{-1}, \lambda_1 r^2$.

The first two terms in (11) are based on the limiting average generalization error. With finite SGD iterations, the iterate w_n may not reach the limiting distribution. On the other hand, for VAR models, it is well known that the speed of convergence for w_n is exponential, and the convergence rate is closely related to the minimum eigenvalue $\lambda_{\min}((\Sigma + \lambda I)\eta) = \lambda\eta$ (see, e.g., Tsay, 2010 Chapter 8.2.2). The finite iterate error leads to the third term of $\exp(-\frac{1}{4}\lambda\eta N)$ in generalization error bound in (11).

In the special case that $\lambda = 0$, (18) reduces to

$$\lim_{n \rightarrow \infty} \mathbb{E}G(w_n) \leq G(w^*) + \frac{r^2\eta}{2p} \text{tr}(I) = r^2\eta.$$

Finally, we consider the scenario where Σ is indefinite with $\delta = -\lambda_{\min}(\Sigma) > 0$. While the population loss F is non-convex, when adopting $\lambda > 2\delta$, we have that $\Sigma + \lambda I$ is positive definite and F_λ is convex. Then the generalization upper bounds need to be updated by replacing λ with $\lambda - \delta$, which leads to a perturbation on the order of δ . In particular, note that by Young's inequality, the derivative of the bias term with respect to λ is bounded by

$$|\partial_\lambda G(w_\lambda^*)| = \sum_{i=1}^p \frac{\lambda \lambda_i^2 a_i^2}{(\lambda + \lambda_i)^3} \leq \sum_{i=1}^p \frac{\lambda_i a_i^2}{4(\lambda + \lambda_i)} \leq \frac{1}{4\lambda} \sum_{i=1}^p \lambda_i a_i^2 = \frac{\|w^*\|_\Sigma^2}{4\lambda}.$$

The derivative of the variance term with respect to λ in (20) is bounded by Young's inequality,

$$|\partial_\lambda \text{var}(\lambda)| = \frac{r^2\eta}{2p} \sum_{i=1}^p \frac{\lambda_i}{(\lambda_i + \lambda)^2} \leq \frac{r^2\eta}{2p} \sum_{i=1}^p \frac{\lambda_1}{\lambda^2} \leq \frac{r^2\eta\lambda_1}{2\lambda^2}.$$

Therefore, replacing λ with $\lambda - \delta$ to handle non-convexity, we need to add the following term in the generalization error bound,

$$\delta |\partial_\lambda F(w_\lambda^*)| + \delta |\partial_\lambda \text{var}(\lambda)| \leq \delta \left(\frac{\|w^*\|_\Sigma^2}{4\lambda} + \frac{r^2\lambda_1\eta}{2\lambda^2} \right).$$

This term can be further upper bounded by the last term of (11).

3 Low Effective Dimension

Given the generalization error bound in Theorem 2.4, we introduce the concept of “low effective dimension” and show that the generalization error

bound in (11) can be independent (or dependent poly-logarithmically) of the ambient dimension p in an overparameterized regime. We will use the O and Ω notations to hide constants independent of p and use the \tilde{O} and $\tilde{\Omega}$ notations to hide constants depend poly-logarithmically on p . In particular, we introduce the following standard asymptotic notations: $A_\epsilon = O(f(\epsilon))$, $B_\epsilon = \tilde{O}(f(\epsilon))$, $C_\epsilon = \Omega(f(\epsilon))$, $D_\epsilon = \tilde{\Omega}(f(\epsilon))$. These notations mean that there exist some universal constants c and $C > 0$ such that,

$$A_\epsilon \leq C f(\epsilon), \quad B_\epsilon \leq C(\log p)^c f(\epsilon), \quad C_\epsilon \geq C f(\epsilon), \quad D_\epsilon \geq C(\log p)^c f(\epsilon).$$

3.1 Initialization and stochastic gradient variance

We investigate the terms that appear in the generalization bound (11): whether they can be independent of p ; and how they affect the necessary sample size $N(\epsilon)$ in (13).

First, we notice that the terms related to initialization w_0 , i.e., $\mathbb{E}\|w_0\|^2$ and $\mathbb{E}G(w_0)$, appear in the sample size $N(\epsilon)$ in (13). If the region $\mathcal{D} = \mathbb{R}^p$, we can often choose appropriate w_0 so that $\mathbb{E}\|w_0\|^2$ and $\mathbb{E}G(w_0)$ are independent of p . For example, for linear regression loss function in (15), we can pick $w_0 = 0$, then $\mathbb{E}G(w_0) = \frac{1}{2}\|w^*\|_A^2$ with $A = \Sigma$, which will be bounded by an $O(1)$ constant as shown below. For a restrictive region \mathcal{D} , although $\mathbb{E}\|w_0\|^2$ and $\mathbb{E}G(w_0)$ may scale as a polynomial function of p , $N(\epsilon)$ only depends logarithmically on these two terms. Therefore, the dimension dependence of $N(\epsilon)$ is only logarithmic.

Second, we consider the stochastic gradient variance r^2 , which contributes to the term C_1 in (11). In a typical setting, it scales roughly as the squared population gradient, i.e.,

$$\begin{aligned} \mathbb{E}\|\nabla f(w, \zeta) - \nabla F(w)\|^2 &\approx O(\mathbb{E}\|\nabla F(w)\|^2) \\ &= O(\mathbb{E}\|\nabla F(w) - \nabla(F(w^*))\|^2) \\ &= O(\mathbb{E}\|\nabla^2 F(w)(w - w^*)\|^2) \end{aligned}$$

Assume that $w \sim \mathcal{N}(0, I_p)$ and $\nabla^2 F \preceq A$

$$= O(\|A\|(\|w^*\|_A^2 + \text{tr}(A))).$$

We will see such an approximation holds for many applications of interest. Moreover, we have $\|A\| \leq \text{tr}(A)$, which can often be p -independent as discussed below. The scale of $\|w^*\|_A$ will also be discussed next.

From the discussion above, we only need to focus on two terms in (11), $\|w^*\|_A$ and $\|w^*\|_{A,\lambda}$. For the generalization error to be small and independent of p , we need to show $\|w^*\|_A$ is dimension independent and $\|w^*\|_{A,\lambda}$ decreases as λ decreases.

3.2 Low effective dimension settings

In this section, we formally define two settings of low effective dimension as Assumptions 3.1 and 3.3. In Sections 4 and 5, we will show that these assumptions easily hold for a wide range of convex and non-convex statistical models.

3.2.1 Sparse true parameter

The first setting is characterized in the following assumption.

Assumption 3.1. *The followings are true*

- 1) $\|A\|$ with A defined in Assumption 2.2 is bounded by an $O(1)$ constant.
- 2) $\|w^*\|$ is bounded by an $O(1)$ constant.
- 3) r^2, c_r defined in Theorem 2.4 are bounded by $O(1)$ constants.
- 4) The initial values $\mathbb{E}\|w_0\|^2$ and $\mathbb{E}G(w_0)$ grow polynomially with p .

Assumption 3.1 can be interpreted as a weak sparsity condition for w^* , since there can be only a few significant components in w^* . Sparsity assumption is a very common condition in the statistical literature. However, our assumption only assumes that the ℓ_2 -norm of w^* , instead of the ℓ_0 -norm, is bounded. As compared to the ℓ_0 -norm, the ℓ_2 -norm is rotation-free. In addition, we do not need to apply any projection or shrinkage procedures on the SGD iterates.

Under Assumption 3.1, Corollary 2.6 can be simplified as the following generalization error bound, which shows that the necessary sample size depends on p in a poly-logarithmic factor.

Proposition 3.2. *Under the conditions in Corollary 2.5 and Assumption 3.1, given any $\epsilon > 0$, when*

$$\lambda(\epsilon) = O(\epsilon), \quad \delta(\epsilon) = O(\epsilon), \quad \eta(\epsilon) = O(\epsilon), \quad N(\epsilon) = \Omega\left(\frac{1}{\epsilon^2}\right), \quad (21)$$

we have $\mathbb{E}[G(\bar{w}_N)1_{\tau \geq N-1}] \leq 3\epsilon$. Moreover, for any $p_0 \geq 0$, there are $A = O(1/p_0)$, if $\mathcal{D} = \{w : \|w - w^*\|^2 \leq A\}$, we have

$$\mathbf{P}(\tau \leq N) \leq p_0.$$

Alternatively, if $\delta = 0$, for any $\alpha > 0$, we take

$$\lambda(\epsilon) = 0, \quad \eta(\epsilon) = O(\epsilon^{1+\alpha}), \quad N(\epsilon) = \Omega\left(\frac{1}{\epsilon^{2+\alpha}}\right),$$

we have $\mathbb{E}[G(\bar{w}_N)1_{\tau \geq N-1}] \leq 3\epsilon$. Meanwhile, for any $A > \mathbb{E}\|w_0 - w^*\|^2$, if $\mathcal{D} = \{w : \|w - w^*\|^2 \leq A\}$, we have

$$\mathbf{P}(\tau \leq N) \leq \frac{\mathbb{E}\|w_0 - w^*\|^2 + O(\epsilon^\alpha)}{A}.$$

Proposition 3.2 consists of two parts. The first part shows that the generalization error is of order $O(1/\sqrt{N})$ if the non-convexity is of the same order. The second part demonstrates how to bound the probability of SGD escaping the convexity region \mathcal{D} if it is a ball centered at w^* . For all $\delta \geq 0$, A needs to be of order $1/\delta$ so that the chance of escaping is less than δ . If the problem is convex in \mathcal{D} , \mathcal{D} just need to include w_0 to ensure the chance of no-escaping is nonzero. In both cases, the escape is harder when the radius \sqrt{A} is larger. This also explains why machine learning literature is in favor of local-minima in large valleys.

3.2.2 Non-sparse true parameter

The second setting is technically more interesting, which assumes the data has a low effective dimension in the following sense. When we say a component or a linear combination of components of w is effective, it means that the loss function F has a significant dependence on it. This can be analyzed through the eigen-decomposition of $\nabla^2 F(w)$ or its upper bound A in (7). Let (λ_i, v_i) be the eigenvalue-eigenvectors of A , where λ_i are arranged in decreasing order. Then a small λ_i indicates that F has a weak dependence along the direction of v_i . For the model to have a low effective dimension, there will be only constantly many λ_i being significant, while the remaining eigenvalues in sum have a negligible contribution to the overall loss function. We formally formulate this setting into the following assumption.

Assumption 3.3. *The followings are true*

- 1) $\text{tr}(A)$ with A defined in Assumption 2.2 is bounded by an $\tilde{O}(1)$ constant.
- 2) the true parameter w^* is bounded in each of A 's eigen-direction, in the sense that

$$\|w^*\|_{A,S} := \max_i \{|\langle v_i, w^* \rangle|, i = 1, \dots, p\} = \tilde{O}(1). \quad (22)$$

- 3) r^2, c_r defined in Theorem 2.4 are bounded by $\tilde{O}(1)$ constants.
- 4) The initial values $\mathbb{E}\|w_0\|^2$ and $\mathbb{E}G(w_0)$ grow polynomially with p .

By Cauchy Schwartz inequality, we have $\|w^*\|_{A,S} \leq \|w^*\|$. So Assumption 3.3 condition 2) is weaker than Assumption 3.1 condition 2). In particular, it can include important cases where we only have upper and lower bounds on each of w^* 's components, and A is known to be a diagonal matrix. These cases are not covered by Assumption 3.1. On the other hand, the spectrum profile of A will be required to choose the regularization parameter as shown in the following proposition.

Proposition 3.4. *By the following inequalities,*

$$\|w^*\|_A^2 \leq \text{tr}(A)\|w^*\|_{A,S}^2, \quad \|w^*\|_{A,\lambda}^2 \leq \|w^*\|_{A,S}^2 \sum_{i=1}^p \lambda \wedge \lambda_i.$$

Assumption 3.3 implies that $\|w^*\|_A = \tilde{O}(1)$ and $\|w^*\|_{A,\lambda}^2 = \tilde{O}(\sum_{i=1}^p \lambda \wedge \lambda_i)$. Moreover, under the conditions in Corollary 2.6 and Assumption 3.3, given any $\epsilon > 0$, if the eigenvalues of A follows,

1) *Exponential decay:* $\lambda_i = e^{-ci}$ for some constant $c > 0$, and setting

$$\lambda = \tilde{O}\left(\frac{\epsilon}{|\log \epsilon|}\right), \quad \delta(\epsilon) = \tilde{O}\left(\frac{\epsilon^3}{|\log \epsilon|^2}\right), \quad \eta(\epsilon) = \tilde{O}\left(\frac{\epsilon^2}{|\log \epsilon|}\right), \quad N(\epsilon) = \tilde{\Omega}\left(\frac{|\log \epsilon|^3}{\epsilon^3}\right),$$

we have $\mathbb{E}[G(w_N)\mathbf{1}_{\tau \geq N}] \leq 4\epsilon$.

2) *Polynomial decay:* $\lambda_i = i^{-c}$ for some constant $c > 0$, and setting

$$\lambda(\epsilon) = \tilde{O}\left(\epsilon^{\frac{c+1}{c}}\right), \quad \delta(\epsilon) = \tilde{O}\left(\epsilon^{\frac{3c+2}{c}}\right), \quad \eta(\epsilon) = \tilde{O}\left(\epsilon^{\frac{2c+1}{c}}\right), \quad N(\epsilon) = \tilde{\Omega}\left(\frac{|\log(\epsilon)|}{\epsilon^{\frac{3c+2}{c}}}\right),$$

we have $\mathbb{E}[G(w_N)\mathbf{1}_{\tau \geq N}] \leq 4\epsilon$.

In both cases, we have

$$\mathbb{E}[G(w_{N \wedge \tau})] \leq \mathbb{E}[G(w_0)] + \tilde{O}(\epsilon |\log \epsilon|).$$

So if $\mathcal{D} = \{w : G(w) \leq (1+a)\mathbb{E}[G(w_0)]\}$, then

$$\mathbf{P}(\tau < N) \leq \frac{1}{1+a} + \tilde{O}(\epsilon |\log \epsilon|).$$

We remark that the parameter of the spectrum decay (e.g., the constant c in polynomial decay spectrum) is often assumed to be known for many functional data analysis problems (Hall and Horowitz, 2007; Cai and Hall, 2008). From Proposition 3.4, for both exponential decay and polynomial

decay of the Hessian spectrum, the sample size N only depends on p in a poly-logarithmic factor.

Similar to Proposition 3.2, the second part of this result demonstrates how to bound the probability of SGD escaping the convexity region \mathcal{D} if it is the sub-level set with w^* inside. The parameter a controls the size of \mathcal{D} . A larger a produces a larger \mathcal{D} and hence a smaller escape probability.

Remark 3.5. *It is interesting to compare our low effective dimension settings with the conditions used in Bartlett et al. (2019). For their main result, Theorem 4 in Bartlett et al. (2019), to yield dimension-independent generalization error bound, three conditions (formulated in our notation) need to hold: 1) $\|w^*\|^2$ is bounded by a constant; 2) $\text{tr}(A)$ is bounded by a constant; 3) the spectrum of A decays not so fast so that, for some k , $\sum_{i \geq k} \lambda_i \geq bN\lambda_k$ with some constant b . In comparison, our Assumption 3.1 only requires conditions 1) and 2), but not the technical condition 3). Moreover, our result can also work under Assumption 3.3 where only $\|w^*\|_{A,S}^2$, instead of $\|w^*\|^2$, needs to be bounded.*

4 Overparameterization in Linear Regression

In general, overparameterization may lead to overfitting, but this sometimes can be avoided. Our main result, Theorem 2.4, provides a general tool to understand why overfitting sometimes happens and sometimes does not. In this section, we will demonstrate how to apply our results on linear regression models in various high dimensional settings. This section is technically straightforward and is mainly used for pedagogical purpose. The discussions on more technically challenging cases for nonlinear and non-convex models are provided in the next section.

4.1 Linear regression

First of all, we will find out the problem related parameters in Theorem 2.4 when applying to linear regression models. As in Section 2.5, we consider *i.i.d.* data points form $\zeta_i = (x_i, y_i) \in \mathbb{R}^p \times \mathbb{R}$, where the response is generated by

$$y_i = x_i^T w^* + \xi_i. \quad (23)$$

In (23), $w^* \in \mathbb{R}^p$ is the true model-parameter to be estimated. $\xi_i \in \mathbb{R}$ are observation noise terms in the observation process, and we assume they are *i.i.d.* with zero mean and variance σ^2 . For simplicity, we assume that the

data x_i are *i.i.d.* Gaussian distributed, i.e., $x_i \sim \mathcal{N}(0, \Sigma)$. As a remark, our proof also allows the non-Gaussian distribution with finite fourth moments.

The regression loss of parameter w on data ζ_i is

$$f(w, \zeta_i) = \frac{1}{2}(x_i^T w - y_i)^2. \quad (24)$$

Plugging (23) into (24) and taking expectation, and we find the population loss function

$$F(w) = \frac{1}{2}(w - w^*)^T \Sigma (w - w^*) + \frac{1}{2}\sigma^2. \quad (25)$$

Now we show the problem related parameters in Theorem 2.4 can be set as below:

Proposition 4.1. *For linear regression, Assumption 2.2 holds with $A = \Sigma$, $\delta = 0$, $\mathcal{D} = \mathbb{R}^p$. When $w_0 = 0$, $\mathbb{E}G(w_0) = \frac{1}{2}\|w^*\|_\Sigma^2$, the stochastic gradient variance bounds in (8) and (10) hold with*

$$r^2 = 2\sigma^2 \text{tr}(\Sigma) + 12\text{tr}(\Sigma)\|w^*\|_\Sigma^2, \quad c_r = \frac{6}{\sigma^2} \max\{\|\Sigma\|, 1\}.$$

As a consequence, by Proposition 3.2, Assumption 3.1 holds if $\|w^*\|^2$ and $\text{tr}(\Sigma)$ are $O(1)$ constants, and the necessary sample size $N(\epsilon)$ in Corollary 2.6 is independent of p . Similarly, by Proposition 3.4, Assumption 3.3 holds if $\|w^*\|_{A,S}^2$ and $\text{tr}(\Sigma)$ are $\tilde{O}(1)$ constants, and the sample size $N(\epsilon)$ depends on p only via a polynomial logarithmic factor.

4.2 High dimensional data with principle components

The low effective dimension settings in Section 3 naturally rise in many high dimensional problems. For example, in image processing or functional data analysis (see e.g. Ramsay and Silverman, 2005; Hall and Horowitz, 2007; Cai and Hall, 2008; Fan et al., 2015; Xue and Yao, 2018), the data are in general assumed to take place in a Hilbert space $(H, \langle \cdot, \cdot \rangle)$ with potentially infinitely many orthonormal basis functions $\{e^j, j = 1, 2, \dots\}$. Each data can be written as

$$x_i = \sum_{j=1}^{\infty} a_i^j e^j. \quad (26)$$

Suppose a_i^j are independent Gaussian random variables with mean zero and variance σ_j^2 . Note that $\mathbb{E}\langle x_i, x_i \rangle = \sum_{j=1}^{\infty} \sigma_j^2$. Therefore, for each data $x_i \in H$,

we assume that $\sum_{j=1}^{\infty} \sigma_j^2 < \infty$ so that the norm of the data is bounded, which implicitly requires σ_j decaying to zero (Hall and Horowitz, 2007). Given the form of x_i in (26), the linear regression model takes the following form,

$$y_i = \langle x_i, w^* \rangle + \xi_i, \quad (27)$$

where $w^* = \sum_{j=1}^{\infty} w^{*,j} e^j$. If we assume $w^* \in H$, then $\langle w^*, w^* \rangle = \sum_{j=1}^{\infty} (w^{*,j})^2 < \infty$.

When training this “infinite dimensional” linear regression model in (27), we would need a finite projection $\mathcal{P}_p : H \mapsto \mathbb{R}^p$. When the basis functions are available, one natural choice of the projection is

$$\mathcal{P}_p x_i = \mathcal{P}_p \left(\sum_{j=1}^{\infty} a_i^j e^j \right) := [a_i^1, \dots, a_i^p]^T.$$

Then the p -dimensional linear regression model is formulated as

$$y_i = (\mathcal{P}_p x_i)^T w_p^* + \xi_i^p. \quad (28)$$

It is worthwhile noticing that the true infinite dimensional model in (27) is compatible with the finite dimensional model in (28), in the sense that

$$w_p^* = \mathcal{P}_p w^* = [w^{*,1}, \dots, w^{*,p}]^T, \quad \xi_i^p = \xi_i + \sum_{j=p+1}^{\infty} w^{*,j} a_i^j.$$

Since a_i^j are independent Gaussian random variables, we have $\xi_i^p \sim \mathcal{N}(0, \sigma_{\xi,p}^2)$ with

$$\sigma_{\xi,p}^2 := \sigma^2 + \sum_{j=p+1}^{\infty} \sigma_j^2 (w^{*,j})^2 \leq \sigma^2 + \|w^*\|_{\Sigma}^2, \quad \|w^*\|_{\Sigma}^2 := \sum_{j=1}^{\infty} \sigma_j^2 (w^{*,j})^2.$$

In the finite dimensional model (28), the data $\mathcal{P}_p x_i$ has the population covariance matrix $\Sigma_p = \text{diag}(\sigma_1^2, \dots, \sigma_p^2)$, whose trace is bounded by

$$\text{tr}(\Sigma_p) = \sum_{j=1}^p \sigma_j^2 \leq \sum_{j=1}^{\infty} \sigma_j^2.$$

Therefore, by Proposition 4.1, the problem related parameters in Theorem

2.4 are

$$\begin{aligned}
A_p &= \Sigma_p, \quad \text{with} \quad \text{tr}(A_p) \leq \sum_{j=1}^{\infty} \sigma_j^2, \quad \|A_p\| = \sigma_1^2, \\
r_p^2 &= 2\sigma_{\xi,p}^2 \text{tr}(\Sigma_p) + 12\text{tr}(\Sigma_p) \|w_p^*\|_{\Sigma_p}^2 \leq 2(\sigma^2 + 7\|w^*\|_{\Sigma}^2) \sum_{j=1}^{\infty} \sigma_j^2, \quad (29) \\
c_{r,p}^2 &= \frac{6}{\sigma_{\xi,p}^2} \max\{1, \sigma_1^2\} \leq \frac{6}{\sigma^2} \max\{1, \sigma_1^2\}.
\end{aligned}$$

Moreover, if we use $w_0 = 0$, then $\mathbb{E}\|w_0\|^2 = 0$ and

$$\mathbb{E}G(w_0) = \frac{1}{2} \|w_p^*\|_{\Sigma_p}^2 \leq \|w^*\|_{\Sigma}^2.$$

Note that

$$\|w^*\|_{\Sigma}^2 = \sum_{j=1}^{\infty} \sigma_j^2 (w^{*,j})^2 \leq \|w^*\|_{\infty}^2 \sum_{j=1}^{\infty} \sigma_j^2, \quad \|w^*\|_{\infty} := \max_{1 \leq j} |w^{*,j}|.$$

So as long as $\|w^*\|_{\infty}$ is finite, the upper bounds above are independent of dimension p .

When the true loading parameter w^* is an element of H , $\|w_p^*\|^2 \leq \langle w^*, w^* \rangle < \infty$. Then we can check that all items of Assumption 3.1 hold. So by Proposition 3.2, we know that the generalization error is dimension-independent. Moreover, this does not require any information of the spectrum decay profile.

More generally, we only need that each component of the true loading parameter w^* is bounded, and w^* does not need to be an element of H itself. In particular, we note that

$$\|w_p^*\|_{\Sigma_p, S} = \max_{1 \leq j \leq p} |w^{*,j}| \leq \|w^*\|_{\infty}.$$

Therefore, if $\|w^*\|_{\infty}$ is finite, Assumption 3.3 holds (but in general Assumption 3.1 does not). Then by Proposition 3.4, the generalization error can be dimension independent when we know the spectrum decay profile.

As a simple demonstration, we run some simulations of SGD on linear regression model (28) and present them in Figure 1. We run SGD on (28) with the sample size $N = 500$ and the regularization parameter $\lambda = 0.01$. The covariance spectrum of predictors is set to be $\sigma_j^2 = j^{-2}$ so that $\text{tr}(A_p)$ in (29) is a constant, and the true parameter is set to be $w^{*,j} = j^{-1}$ for




Figure 1: Generalization error bar plot with high dimensional linear regression for different settings of w_p^* and Σ_p . The x -axis is the dimension and y -axis is the generalization error.

$1 \leq j \leq p$ so that $\|w^*\|_\Sigma$ is bounded. The problem dimension ranges from $p = 100$ to $p = 1000$, which can be larger than the sample size. We use the final SGD output w_{500} as the estimator and compute the generalization error as in (15). We repeat this experiment 1000 times and compute the mean and standard deviation. We plot the error bar plot in the upper left panel of Figure 1. As one can see, the generalization error does not increase as the dimension increases, even when $p \gg N$. As a comparison experiment, we run simulations with the same settings except for $\sigma_j^2 \equiv 1$. We plot the generalization error in the upper right panel of Figure 1, which clearly shows the overfitting phenomenon, even when the dimension is in a lower range.

The similar story repeats when the true parameter $w^{*,j} \equiv 1$ (i.e., the case when $\|w^*\|_\infty$ is bounded), where the plots are given by the lower panels

in Figure 1. As one can see, the generalization error with the decaying spectrum still remains stable against the increase of dimension, and it does not change much from the previous setting where the components of w^* are decaying. Meanwhile, overfitting with constant spectrum (i.e., $\sigma_j^2 \equiv 1$) becomes stronger. This simple illustrative example justifies that the low effective dimension helps to address the overfitting issue, even when the dimension p is much larger than the sample size N .

4.3 Overfitting with redundant features

Another interesting setting of overparameterization is to consider adding redundant predictors to an existing model. In this scenario, the true model is low dimensional with the true parameter $w^* \in \mathbb{R}^d$. Suppose we do not know the true model and collect additional features $z \in \mathbb{R}^{p-d}$, so that the overparameterized linear model is written as

$$y_i = x_i^T w^* + z_i^T u^* + \xi_i, \quad (30)$$

where $w^* \in \mathbb{R}^d$, $u^* = 0$, and $[x_i; z_i]$ is jointly Gaussian with mean zero and covariance

$$\Sigma_p = \begin{bmatrix} \Sigma_x & B \\ B^T & \Sigma_z \end{bmatrix}.$$

We assume that $\|\Sigma_z\| \leq \|\Sigma_x\|$ for the ease of discussion. Then, Σ_p being PSD implies that $\|B\| \leq \|\Sigma_x\|$. Since we do not impose any restriction on B other than Σ_p being PSD, our setting allows the possibility that some components of z_i to be highly correlated or even identical with the ones of x_i . This, in general, leads to highly singular design matrices and unstable offline learning results.

We apply Proposition 4.1 and find $A_p = \Sigma_p$. By triangular inequality, for any vectors x and z ,

$$\|\Sigma_p[x; z]\| = \|\Sigma_x x + Bz; B^T x + \Sigma_z z\| \leq 2\|\Sigma_x\| \| [x; z] \| \Rightarrow \|\Sigma_p\| \leq 2\|\Sigma_x\|.$$

For simplicity, we initialize with $[w_0; u_0] = 0$, so

$$\mathbb{E}G(w_0, u_0) = \frac{1}{2}\|w_0 - w^*\|_{\Sigma_x}^2 + \frac{1}{2}\|u_0\|_{\Sigma_z}^2 = \frac{1}{2}\|w^*\|_{\Sigma_x}^2.$$

Moreover, we have

$$c_{r,p} = \frac{6}{\sigma^2} \max\{\|\Sigma_p\|, 1\} \leq \frac{6}{\sigma^2} \max\{2\|\Sigma_x\|, 1\},$$

and

$$\|[w^*, u^*]\|_{\Sigma_p, S} = \|w^*\|_\infty, \quad \|[w^*, u^*]\|^2 = \|w^*\|^2.$$

These upper bounds are all independent of p , or the choice of Σ_z and B .

Meanwhile,

$$r_p^2 = 2(\sigma^2 + 6\|w^*\|_{\Sigma_x}^2)(\text{tr}(\Sigma_x) + \text{tr}(\Sigma_z)), \quad \text{tr}(A_p) = \text{tr}(\Sigma_x) + \text{tr}(\Sigma_z). \quad (31)$$

Given these simple calculation, we find that the only problem related parameters that depend on z are r_p^2 and $\text{tr}(A_p)$ in (31) through $\text{tr}(\Sigma_z)$. Therefore, our theory indicates that there is a simple dichotomy on whether the model (30) will overfit.

If $\text{tr}(\Sigma_z)$ is bounded by a constant independent of p , Proposition 3.2 applies, which indicates that the generalization error is also independent of the ambient dimension p . This can happen if we select data features in z as PCA components. For example, suppose that the redundant data is in the form of $\sum_{j=1}^{\infty} a_j^j e^j$ as in the setting of (26), and we collect the $p-d$ dimensional principle components as $z_i = [a_i^1, \dots, a_i^{p-d}]^T$. Then $\text{tr}(\Sigma_z) = \sum_{j=1}^{p-d} \sigma_j^2 < \sum_{j=1}^{\infty} \sigma_j^2$, which is independent of p .

If $\text{tr}(\Sigma_z)$ grows with p , model (30) may overfit. For simplicity, we consider a special case where $\Sigma_z = I_{p-d}, B = 0$. In other words, the redundant features are independent with each other and the features of x . Then our derivation shows that $r_p^2 = O(p)$. This indicates that the learning results may overfit.

To demonstrate this dichotomy, we simulate the SGD learning results and present their generalization error in Figure 2. In particular, we set $d = 5$ with true parameter $w^* = [1, 1, 1, 1, 1]$, $\Sigma_x = I_5, \sigma^2 = 1$. We let $B = 0$ and choose first that Σ_z to be diagonal with decaying entries $\frac{1}{j^2}$. We run SGD with 500 iterations and compute the generalization error of the final iterate. We repeat this 1000 times and plot the error bar plot in the left panel of Figure 2. As we can see, the generalization error is stable against the increase of the dimension p . In comparison, if we use $\Sigma_z = I_{p-d}$, the learning results overfit, as we can see from the right panel of Figure 2.

5 Overparameterization for Nonlinear and Non-convex Models

In this section, we apply our main theorem and corollaries in Section 3 to several important nonlinear and non-convex statistical problems, such as logistic regression, M-estimator with Tukey's biweight loss function, and two-layer neural networks.




Figure 2: Generalization error bar plot with high dimensional redundant features for two different cases of Σ_z . The x -axis is the dimension p and y -axis is the generalization error.

5.1 Logistic regression

We consider the logistic regression for binary classification with N *i.i.d.* data $\zeta_i = (x_i, y_i)$. The binary response y_i takes values within $\{-1, 1\}$ with probability

$$\mathbf{P}(y_i = y|x_i) = \frac{1}{1 + \exp(-yx_i^T w^*)}, \quad y = \pm 1,$$

where $w^* \in \mathbb{R}^p$ is the true parameter to be estimated. We assume the predictors x_i are *i.i.d.* with $\mathbb{E}x_i x_i^T = \Sigma$. For each data, we adopt the negative log-likelihood as the loss function

$$f(w, \zeta_i) := \log(1 + \exp(-y_i x_i^T w)),$$

and the corresponding population loss is given by

$$F(w) = \mathbb{E}f(w, \zeta) = \mathbb{E} \log(1 + \exp(-yx^T w)).$$

The problem related parameters in Theorem 2.4 can be set by the following proposition.

Proposition 5.1. *For logistic regression, Assumption 2.2 holds with $A = \Sigma, \delta = 0, \mathcal{D} = \mathbb{R}^p$. When $w_0 = 0, \mathbb{E}G(w_0) = \log 2 = O(1)$, the stochastic gradient variance bound in (10) holds with*

$$r^2 = \text{tr}(\Sigma), \quad c_r = 0.$$

As a consequence, by Proposition 3.2, Assumption 3.1 holds if $\|w^*\|^2$ and $\text{tr}(\Sigma)$ are $O(1)$ constants, and the sample size $N(\epsilon)$ in Corollary 2.6 is independent of p . Similarly, by Proposition 3.4, Assumption 3.3 holds if $\|w^*\|_{\Sigma,S}^2$ and $\text{tr}(\Sigma)$ are $\tilde{O}(1)$ constants, and the sample size $N(\epsilon)$ depends on p only via a poly-logarithmic factor.

5.2 M -estimator with Tukey's biweight loss function

In this non-convex example, we assume that the data $\zeta_i = (x_i, y_i)$ are generated from a linear model

$$y_i = x_i^T w^* + \xi_i. \quad (32)$$

We assume $x_i \sim \mathcal{N}(0, \Sigma)$, and ξ_i are i.i.d. mean-zero noises with finite fourth moment. We adopt the *non-convex* Tukey's biweight loss function as follows for the purpose of robust estimation

$$\rho(u) = \begin{cases} \frac{c^2}{6} [1 - (1 - (u/c)^2)^3] & \text{if } |u| \leq c; \\ \frac{c^2}{6} & \text{if } |u| > c. \end{cases}$$

Then the individual data loss function and the population loss are given by,

$$f(w, \zeta) = \rho(x^T w - y) = \rho(x^T (w - w^*) - \xi), \quad F(w) = \mathbb{E} \rho(x^T (w - w^*) - \xi).$$

Proposition 5.2. *For the M -estimator with Tukey's biweight loss in (32), the model true parameter w^* is a local minimum if and only if*

$$c_0 = \mathbb{E}[(1 - (\xi/c)^2)(1 - 5(\xi/c)^2)1_{|\xi| \leq c}] > 0.$$

In that case, Assumption 2.2 holds with any $\delta \geq 0$, $A = \Sigma$ and

$$\mathcal{D} = \{w : \|w - w^*\|_{\Sigma} \leq \frac{c_0 + \delta}{16}\}.$$

Moreover, the stochastic gradient variance bound in (10) holds with

$$r^2 = \text{tr}(\Sigma), \quad c_r = 0.$$

Since $G(w_0) \leq \max_u \rho(u) = \frac{c^2}{6}$, by Proposition 3.2, Assumption 3.1 holds if $\|w^*\|^2$, $\|w_0\|^2$ and $\text{tr}(\Sigma)$ are $O(1)$ constants, and the sample size $N(\epsilon)$ in Corollary 2.6 is independent of p . Similarly, by Proposition 3.4, Assumption 3.3 holds if $\|w_0\|_{\infty}$, $\|w^*\|_{\Sigma,S}^2$ and $\text{tr}(\Sigma)$ are $\tilde{O}(1)$ constants, and the sample size $N(\epsilon)$ depends on p only via a polynomial logarithmic factor.

5.3 Two-layer neural network

In this example, we consider applying our result to two-layer neural networks (NN). We assume that every data point $\zeta = (x, y)$ consists of a p -dimensional predictor $x \sim \mathcal{N}(0, \Sigma)$ and a univariate response $y \in \mathbb{R}$. We assume that the response is generated by

$$y = g(w, x) + \xi, \quad \mathbb{E}\xi = 0, \quad \mathbb{E}\xi^2 = \sigma_0^2.$$

The function g takes the form of a two-layer NN:

$$g(w, x) = c^T \psi(bx + a) = \sum_{i=1}^k c_i \psi(b_i^T x + a_i). \quad (33)$$

In (33), a and c are k -dimensional vectors with a_i and c_i being their components. The notation b is a p by k matrix, and b_i denotes the i -th column of b with $i = 1, \dots, k$. We impose no restriction on k and it can depend on p in general. We denote all the parameters by $w = [a; b_1, \dots, b_k; c] \in \mathbb{R}^{(p+2)k}$. In (33), ψ denotes the activation function. Popular choices of ψ include the rectified linear unit (ReLU), sigmoid function, and the hyperbolic tangent. Here, we do not require ψ to take a specific form but only satisfy certain regularity assumptions for some constant $C > 0$,

$$\psi(0) = 0, \quad |\dot{\psi}(x)| \leq C, \quad |\ddot{\psi}(x)| \leq C. \quad (34)$$

It is easy to verify that hyperbolic tangent satisfies these requirements, and the sigmoid also satisfies these if we shift its center to zero. The condition $\psi(0) = 0$ is mainly for the ease of technical derivations. Although ReLU does not have continuous derivatives, one can find a smooth approximation to meet these requirements.

Since we consider the regression problem, the squared loss function is given by

$$f(w, \zeta) = (y - g(w, x))^2 = (g(w^*, x) + \xi - g(w, x))^2.$$

We also introduce the following $(p+2)k$ by $(p+2)k$ block-diagonal matrix

$$\Sigma^* = \text{diag}\{I_k, \Sigma, \Sigma, \dots, \Sigma, I_k\}.$$

This matrix introduces a high dimensional norm

$$\|w\|_{\Sigma^*}^2 = w^T \Sigma^* w = \|a\|^2 + \sum_{i=1}^k \|b_i\|_{\Sigma}^2 + \|c\|^2.$$

Recall that b_i is of dimension p , its contribution to $\|w\|_{\Sigma^*}^2$ is $\|b_i\|_{\Sigma}^2$. By Proposition 3.4, $\|b_i\|_{\Sigma}^2 \leq \text{tr}(\Sigma)\|b_i\|_{\Sigma,S}^2$, which can be independent of p under suitable conditions.

We are ready to show that the two-layer NN will not overfit in some overparameterized settings.

Proposition 5.3. *Assume the activation function satisfies the condition in (34). With the two-layer NN defined in (33), Assumption 2.2 holds for any $\delta \in (0, 1/4]$ with*

$$A = C_0(w^*)\Sigma^*, \quad \mathcal{D} = \{w : \|w - w^*\|_{\Sigma^*} \leq \delta C_1(w^*)\|w^*\|_{\Sigma^*}\}.$$

For any $w_0 \in \mathcal{D}$, $G(w_0) \leq C_2(w^*)\|w^*\|_{\Sigma^*}^4$, the stochastic gradient variance bound in (10) holds with

$$r^2 = C_3(w^*), c_r = 0.$$

The exact values of the problem related parameters are given by

$$\begin{aligned} C_0(w^*) &= 7C^2\|w^*\|_{\Sigma^*}^2, & C_1(w^*) &= \frac{2}{9\sqrt{2}(2\|w^*\|_{\Sigma^*} + 1)}, & C_2(w^*) &= C^2\|w^*\|_{\Sigma^*}^4, \\ C_3(w^*) &= 8\sqrt{3}(1 + \text{tr}(\Sigma))C^2\|w^*\|_{\Sigma^*}^2(C^2\|w^*\|_{\Sigma^*}^4 + \sigma_0^2). \end{aligned}$$

As a consequence, by Proposition 3.2, Assumption 3.1 holds if

$$\max \left\{ \|a^*\|^2 + \sum_{i=1}^k \|b_i^*\|^2 + \|c^*\|^2, \text{tr}(\Sigma), \|w_0\|_{\infty} \right\} = O(1), \quad (35)$$

and the sample size $N(\epsilon)$ is independent of p . Similarly, by Proposition 3.4, Assumption 3.3 holds if

$$\max \{k, \text{tr}(\Sigma), \|w_0\|_{\infty}, |a_i|, |c_i|, |v_j^T b_i|, i = 1, \dots, k, j = 1, \dots, p\} = \tilde{O}(1), \quad (36)$$

where v_j are the eigenvectors of Σ . Then the sample size $N(\epsilon)$ depends on p only via a polynomial logarithmic factor.

It is worthwhile mentioning that a similar version of Condition (35) can also be found in Neyshabur et al. (2019). In particular, Neyshabur et al. (2019) assumed $\|c^*\|^2, \sum_{i=1}^k \|b_i^*\|^2$ to be $O(1)$ while the parameter a is set to be 0. There is no variance assumption of x_i in Neyshabur et al. (2019), but it is assumed that $\mathbb{E}\|x_i\|^2$ is $O(1)$, which is equivalent to requesting $\text{tr}(\Sigma) = O(1)$ in our setting.

When the width of the hidden layer k is a fixed constant, the second condition (36) is in general less restrictive than the first one (35), since it allows $\|b_i^*\|$ to grow with p . When k grows with p , only the first condition is applicable, and it requires that $\|a^*\|^2 + \sum_{i=1}^k \|b_i^*\|^2 + \|c^*\|^2$ is bounded by $O(1)$. In other words, we need either $k = O(1)$ or the true parameters to be bounded by $O(1)$ to prevent overfitting. This can also be understood intuitively. Note that the output of the two-layer NN in (33) is a sum of k objects. Therefore, if k grows with p , the output of (33) will diverge, which contradicts the common assumption that g is bounded (see, e.g., Li and Liang, 2018; Neyshabur et al., 2019; Allen-Zhu et al., 2019). Our result is consistent with the results in Allen-Zhu et al. (2019), in the sense that Allen-Zhu et al. (2019) also showed that the sample size needs to grow with k . It is also possible to rescale g by multiplying (33) with a factor $\frac{1}{k}$ or $\frac{1}{\sqrt{k}}$, as done by Arora et al. (2019), so that the generalization error is independent of the parameter k .

6 Conclusions and Future Works

One classical canon of statistics is that high dimensional models are prone to overfitting when the data sample size is not sufficiently large. However, many existing models, such as neural networks (NN), exhibit stable generalization error despite being overparameterized. This paper developed an analysis framework of the generalization error for high dimensional regularized online learning. The error bound can be interpreted as a bias-variance tradeoff through a simplified stochastic approximation. This result indicates that overparameterization does not lead to overfitting if the model has a low effective dimension. We demonstrated how to apply this framework on various models such as linear regression, logistic regression, M -estimator with Tukey’s biweight loss, and two-layer NN.

There are a few future directions. First, the generalization bound in Theorem 2.4 only applies when the SGD iterates stay in a local region \mathcal{D} near the true parameter w^* . Although it is a common assumption when analyzing non-convex models, this assumption might be difficult to check in practice. To address this challenge, a simple data splitting can serve as a remedy. In practice, we may split the data into two parts and run SGD through the first half. If $w_{N/2}$ gives us some rough ideas on how to construct \mathcal{D} , we can use $w_{N/2}$ as an initialization and run SGD on the second half of the data. In the process, we can check whether the iterates stay in \mathcal{D} , and then Theorem 2.4 provides us an upper bound for the generalization error of

w_N . Of course, it would be an interesting future work on how to construct \mathcal{D} based on the SGD solution from the first half of the data.

Second, our framework indicates that the ambient model dimension itself may not be a good indicator of model complexity, especially in overparameterized settings. The quantity that characterizes the data variability may lead to new information criterion for model selection in the overparameterized setting. Such results may extend the classical criteria such as the AIC and BIC.

Last but not least, regularization and overparameterization are good tools to handle misspecified models. [Hastie et al. \(2019\)](#) has discussed this issue for linear regression problems. How to extend our results to nonlinear misspecified models will be very interesting.

Acknowledgement

Xi Chen would like to thank the support from NSF via the grant IIS-1845444. Qiang Liu would like to thank the support from Singapore MOE via the grant R-146-000-258-114. Xin T. Tong would like to thank the support from Singapore MOE via the grant R-146-000-292-114. The authors also thank the suggestions made by the editor and anonymous reviewers.

A Proof of the Main Results in Section 2.2

A.1 Preliminaries

Lemma A.1. *For any vector $v \in \mathbb{R}^p$ and PSD matrix $A \in \mathbb{R}^{p \times p}$, the following results hold*

1) *For any $-\delta A \preceq B \preceq A$, let $B = V\Lambda V^T$ be the eigenvalue decomposition of B , and denote $|\Lambda|$ as taking absolute value on each element of the diagonal matrix Λ . Denote $|B| = V|\Lambda|V^T$. Then for any vectors v and w , $a > 0$*

$$2\langle v, Bw \rangle \leq a\langle v, |B|v \rangle + \frac{1}{a}\langle w, |B|w \rangle \leq a(1 + \delta)\|v\|_A^2 + \frac{1 + \delta}{a}\|w\|_A^2.$$

2) *For any $-\delta A \preceq B \preceq A$, and any vectors u and v , $a > 0$*

$$2|\langle u, Bv \rangle| \leq a\langle u, Bu \rangle + 2a\delta\|u\|_A^2 + \frac{1 + 2\delta}{a}\|v\|_A^2.$$

Proof. Claim 1). Let (l_i, u_i) be the eigenvalue-eigenvectors of B . Assume also that

$$v = \sum_{i=1}^p a_i u_i, \quad w = \sum_{i=1}^p b_i u_i.$$

Then by Young's inequality

$$2\langle v, Bw \rangle = 2 \sum_{i=1}^p l_i a_i b_i \leq a \sum_{i=1}^p |l_i| |a_i|^2 + \frac{1}{a} \sum_{i=1}^p |l_i| |b_i|^2 = a\langle v, |B|v \rangle + \frac{1}{a}\langle w, |B|w \rangle.$$

Next, we denote the positive part of Λ as Λ_+ and the negative part as Λ_- , so that

$$\Lambda = \Lambda_+ + \Lambda_-, \quad |\Lambda| = \Lambda_+ - \Lambda_-, \quad \Lambda_- \preceq 0 \preceq \Lambda_+.$$

Then by checking eigen-space with nonnegative eigenvalues, $B \preceq A$ indicates that $V\Lambda_+V^T \preceq A$. Likewise, we have $-\delta A \preceq V\Lambda_-V^T$. In combination, we have

$$|B| = V\Lambda_+V^T - V\Lambda_-V^T \preceq (1 + \delta)A.$$

Therefore

$$a\langle v, |B|v \rangle + \frac{1}{a}\langle w, |B|w \rangle \leq (1 + \delta)a\|v\|_A^2 + \frac{1 + \delta}{a}\|w\|_A^2.$$

For claim 2), denote $B_\delta = B + \delta A \succeq 0$. Then

$$\begin{aligned}
2|\langle u, Bv \rangle| &\leq 2|\langle u, B_\delta v \rangle| + 2\delta|\langle u, Av \rangle| \\
&\leq a\|u\|_{B_\delta}^2 + \frac{1}{a}\|v\|_{B_\delta}^2 + a\delta\|u\|_A^2 + \frac{\delta}{a}\|v\|_A^2 \\
&= a\langle u, Bu \rangle + a\delta\|u\|_A^2 + \frac{1}{a}\langle v, Bv \rangle + \frac{\delta}{a}\|v\|_A^2 + a\delta\|u\|_A^2 + \frac{\delta}{a}\|v\|_A^2 \\
&\leq a\langle u, Bu \rangle + 2a\delta\|u\|_A^2 + \frac{1+2\delta}{a}\|v\|_A^2.
\end{aligned}$$

□

A.2 Proof of the main results

Proof of Theorem 2.3. We rewrite SGD update as

$$w_{n+1} = w_n - \eta \nabla f_\lambda(w_n, \zeta_n) = w_n - \eta \nabla F_\lambda(w_n) + \eta \xi_n, \quad (37)$$

where

$$\xi_n = \nabla F_\lambda(w_n) - \nabla f_\lambda(w_n, \zeta_n) = \nabla F(w_n) - \nabla f(w_n, \zeta_n).$$

Let \mathcal{F}_n be the σ -algebra generated by $\{w_{i+1}, \zeta_i, i = 1, \dots, n-1\}$. We use $\mathbb{E}_n(\cdot)$ to denote the conditional expectation $\mathbb{E}(\cdot | \mathcal{F}_n)$. Then ξ_n is a martingale sequence since $\mathbb{E}_n \xi_n \equiv 0$.

From (37), we find

$$\|w_{n+1} - w^*\|^2 = \|w_n - w^*\|^2 - 2\eta \langle w_n - w^*, \nabla F_\lambda(w_n) - \xi_n \rangle + \eta^2 \|\nabla F_\lambda(w_n) - \xi_n\|^2. \quad (38)$$

We first try to find a bound of $\langle -(w_n - w^*), \nabla F_\lambda(w_n) \rangle$. We define

$$B_n := \int_0^1 \nabla^2 F_\lambda(sw_n + (1-s)w^*) ds = \lambda I + \int_0^1 \nabla^2 F(sw_n + (1-s)w^*) ds,$$

and apply fundamental theorem of calculus on ∇F_λ . Note that $\nabla F(w^*) = 0$, we obtain

$$\begin{aligned}
\nabla F_\lambda(w_n) &= \nabla F_\lambda(w^*) + \int_0^1 \nabla^2 F_\lambda(sw_n + (1-s)w^*)(w_n - w^*) ds \\
&= \lambda w^* + B_n(w_n - w^*).
\end{aligned} \quad (39)$$

Note that $\frac{1}{2}\lambda I \preceq -\delta A + \lambda I \preceq B_n \preceq A + \lambda I$ and B_n is symmetric, we have

$$\begin{aligned} \langle -(w_n - w^*), \nabla F_\lambda(w_n) \rangle &= -\|w_n - w^*\|_{B_n}^2 - \lambda \langle w^*, w_n - w^* \rangle \\ &\leq -\frac{1}{2}\|w_n - w^*\|_{B_n}^2 - \frac{\lambda}{4}\|w_n - w^*\|^2 - \lambda \langle w^*, w_n - w^* \rangle \\ &\leq -\frac{1}{2}\|w_n - w^*\|_{B_n}^2 + \lambda \|w^*\|^2. \end{aligned} \quad (40)$$

Furthermore, note that by $B_n \preceq A + \lambda I$, we have

$$\|\nabla F_\lambda(w_n)\|^2 = \|\lambda w^* + B_n(w_n - w^*)\|^2 \leq 2\lambda^2\|w^*\|^2 + 2(\|A\| + \lambda)\|w_n - w^*\|_{B_n}^2. \quad (41)$$

Similarly, we find

$$\nabla F(w_n) = \int_0^1 \nabla^2 F(sw_n + (1-s)w^*) ds (w_n - w^*) = (B_n - \lambda I)(w_n - w^*),$$

thus

$$\|\nabla F(w_n)\|^2 \leq \|A\| \|w_n - w^*\|_{B_n}^2. \quad (42)$$

Recall that w_n is \mathcal{F}_n -measurable and $\mathbb{E}_n \xi_n = 0$. Also

$$\mathbb{E}\|\xi_n\|^2 \leq r^2 + c_r |(w_n - w^*)^T \nabla F(w_n)| \leq r^2 + c_r \|w_n - w^*\|_{B_n}^2.$$

So plugging (40) and (41) into (38), using (8) with (42), and by Cauchy Schwarz inequality, we then have

$$\begin{aligned} \mathbb{E}_n \|w_{n+1} - w^*\|^2 &= \|w_n - w^*\|^2 - 2\eta \mathbb{E}_n \langle w_n - w^*, \nabla F_\lambda(w_n) - \xi_n \rangle + \eta^2 \mathbb{E}_n \|\nabla F_\lambda(w_n) - \xi_n\|^2 \\ &= \mathbb{E}_n [\|w_n - w^*\|^2 - 2\eta \langle w_n - w^*, \nabla F_\lambda(w_n) \rangle + \eta^2 (\|\nabla F_\lambda(w_n)\|^2 + \|\xi_n\|^2)] \\ &\leq \|w_n - w^*\|^2 - \eta \|w_n - w^*\|_{B_n}^2 + 2\lambda\eta(1 + \lambda\eta)\|w^*\|^2 \\ &\quad + \eta^2 2(1 + c_r)(\|A\| + \lambda)\|w_n - w^*\|_{B_n}^2 + \eta^2 r^2. \end{aligned}$$

Under the condition

$$\eta \leq \min \left\{ \frac{1}{4(1 + c_r)(\|A\| + \lambda)}, 1 \right\}, \quad \lambda \leq 1,$$

and since $0 \leq 1_{\tau \geq n+1} \leq 1_{\tau \geq n}$, we have

$$\begin{aligned} \mathbb{E}[1_{\tau \geq n+1} \|w_{n+1} - w^*\|^2] &\leq \mathbb{E}[1_{\tau \geq n} \|w_{n+1} - w^*\|^2] \\ &= \mathbb{E}[1_{\tau \geq n} \mathbb{E}_n \|w_{n+1} - w^*\|^2] \\ &\leq \mathbb{E}[1_{\tau \geq n} (\|w_n - w^*\|^2 - \frac{1}{2}\eta \|w_n - w^*\|_{B_n}^2)] + 4\lambda\eta \|w^*\|^2 + \eta^2 r^2. \end{aligned}$$

Summing this inequality over all $n = 0, \dots, (N \wedge \tau) - 1$, we find that

$$\mathbb{E}[\|w_{\tau \wedge N} - w^*\|^2] \leq \mathbb{E}\|w_0 - w^*\|^2 + N(4\lambda\eta\|w^*\|^2 + \eta^2 r^2). \quad (43)$$

Summing the same inequality over all $n = 0, \dots, N - 1$, we find that

$$\mathbb{E} \left[\mathbf{1}_{\tau \geq N-1} \left(\frac{1}{2} \eta \sum_{n=0}^{N-1} \|w_n - w^*\|_{B_n}^2 \right) \right] \leq \mathbb{E}\|w_0 - w^*\|^2 + N(4\lambda\eta\|w^*\|^2 + \eta^2 r^2). \quad (44)$$

To continue, recall $G(w_n) = F(w_n) - F(w^*)$. Apply fundamental theorem of calculus to $F(w)$, we obtain

$$\begin{aligned} G(w_n) &= F(w_n) - F(w^*) \\ &= \left[\int_0^1 \nabla F(sw_n + (1-s)w^*) ds \right]^T (w_n - w^*) \\ &= \left[\int_0^1 \left(\nabla F(w^*) + \int_0^s \nabla^2 F(tw_n + (1-t)w^*) (w_n - w^*) dt \right) ds \right]^T (w_n - w^*) \\ &= (w_n - w^*)^T \left[\int_0^1 (1-s) \nabla^2 F(sw_n + (1-s)w^*) ds \right] (w_n - w^*) \\ &= (w_n - w^*)^T A_n (w_n - w^*), \end{aligned} \quad (45)$$

with

$$A_n = \int_0^1 (1-s) \nabla^2 F(sw_n + (1-s)w^*) ds.$$

Under Assumption 3.1, we observe that

$$\begin{aligned} A_n + \frac{1}{2} \lambda I &= \int_0^1 (1-s) \nabla^2 F_\lambda(sw_n + (1-s)w^*) ds \\ &\preceq \int_0^1 \nabla^2 F_\lambda(sw_n + (1-s)w^*) ds = B_n. \end{aligned}$$

Namely, we have $H(w_n) = G(w_n) + \frac{\lambda}{2} \|w_n - w^*\|^2 \leq \|w_n - w^*\|_{B_n}^2$. Together with (44), we obtain

$$\mathbb{E} \left[\mathbf{1}_{\tau \geq N-1} \left(\frac{1}{2} \eta \sum_{n=0}^{N-1} H(w_n) \right) \right] \leq \mathbb{E}\|w_0 - w^*\|^2 + N(4\lambda\eta\|w^*\|^2 + \eta^2 r^2).$$

Then because H is convex within \mathcal{D} , we have $H(\bar{w}_N) \leq \frac{1}{N} \sum_{n=0}^{N-1} H(w_n)$ and

$$\mathbb{E}[1_{\tau \geq N-1}(\eta N H(\bar{w}_N))] \leq 2\mathbb{E}\|w_0 - w^*\|^2 + 2N(4\lambda\eta\|w^*\|^2 + \eta^2 r^2).$$

This leads to our claim

$$\mathbb{E}[1_{\tau \geq N-1} G(\bar{w}_N)] \leq \mathbb{E}[1_{\tau \geq N-1} H(\bar{w}_N)] \leq \frac{2\mathbb{E}\|w_0 - w^*\|^2}{N\eta} + 8\lambda\|w^*\|^2 + 2\eta r^2.$$

□

Proof of Theorem 2.4. Step 1: we build a bound for $\|w_n\|^2$. We rewrite SGD update as

$$w_{n+1} = w_n - \eta \nabla f_\lambda(w_n, \zeta_n) = w_n - \eta \nabla F_\lambda(w_n) + \eta \xi_n, \quad (46)$$

where

$$\xi_n = \nabla F_\lambda(w_n) - \nabla f_\lambda(w_n, \zeta_n) = \nabla F(w_n) - \nabla f(w_n, \zeta_n).$$

Let \mathcal{F}_n be the σ -algebra generated by $\{w_{i+1}, \zeta_i, i = 1, \dots, n-1\}$. We use $\mathbb{E}_n(\cdot)$ to denote the conditional expectation $\mathbb{E}(\cdot | \mathcal{F}_n)$. Then ξ_n is a martingale sequence since $\mathbb{E}_n \xi_n \equiv 0$.

From (46), we find

$$\|w_{n+1}\|^2 = \|w_n\|^2 - 2\eta \langle w_n, \nabla F_\lambda(w_n) - \xi_n \rangle + \eta^2 \|\nabla F_\lambda(w_n) - \xi_n\|^2. \quad (47)$$

To continue, we try to find a bound of $-2\eta \langle w_n, \nabla F_\lambda(w_n) \rangle$. We define

$$B_n := \int_0^1 \nabla^2 F(sw_n + (1-s)w^*) ds,$$

and apply fundamental theorem of calculus on ∇F . Note that $\nabla F(w^*) = 0$, we obtain

$$\nabla F(w_n) = \nabla F(w^*) + \int_0^1 \nabla^2 F(sw_n + (1-s)w^*)(w_n - w^*) ds = B_n(w_n - w^*). \quad (48)$$

Note that $-\delta A \preceq B_n \preceq A$. We have

$$\begin{aligned} \langle -w_n, \nabla F(w_n) \rangle &= -\langle w_n, B_n(w_n - w^*) \rangle \\ &= -\langle w_n, (B_n + \delta A)w_n \rangle + \langle w_n, (B_n + \delta A)w^* \rangle + \delta(\|w_n\|_A^2 - \langle w_n, Aw^* \rangle) \\ &\leq \frac{1}{4}\|w^*\|_{B_n + \delta A}^2 + \delta(2\|w_n\|_A^2 + \frac{1}{4}\|w^*\|_A^2) \\ &\leq 2\delta\|A\|\|w_n\|^2 + \frac{1}{2}\|w^*\|_A^2 \text{ since } \delta \leq \frac{1}{2} \text{ and } \|w_n\|_A^2 \leq \|A\|\|w_n\|^2. \end{aligned}$$

Recall that $\delta\|A\| \leq \frac{\lambda}{4}$, we find

$$\begin{aligned}
-2\eta\langle w_n, \nabla F_\lambda(w_n) \rangle &= -2\eta\langle w_n, \nabla F(w_n) + \lambda w_n \rangle \\
&= -2\lambda\eta\|w_n\|^2 + 2\eta\langle -w_n, \nabla F(w_n) \rangle \\
&\leq -\lambda\eta\|w_n\|^2 + \eta\|w^*\|_A^2.
\end{aligned} \tag{49}$$

If $B_n = Q\Lambda Q^T$ is the eigendecomposition of B_n , let $|B_n| = Q|\Lambda|Q^T$, where $|\Lambda|$ takes absolute value on each element of the diagonal matrix Λ . From the proof of Lemma A.1 claim 1), we know $|B_n| \preceq (1 + \delta)A \preceq (1 + \delta)\|A\|I$. Thus by $B_n \preceq A$, we have

$$\begin{aligned}
\|\nabla F(w_n)\|^2 &= \|B_n(w^* - w_n)\|^2 \leq 2\|B_n w^*\|^2 + 2\|B_n w_n\|^2 \\
&\leq 2(w^*)^T B_n^{1/2} |B_n| B_n^{1/2} w^* + 2\|A\|^2 \|w_n\|^2 \\
&\leq 2(1 + \delta)\|A\| \|B_n^{1/2} w^*\|^2 + 2\|A\|^2 \|w_n\|^2 \\
&\leq 4\|A\| \|w^*\|_A^2 + 2\|A\|^2 \|w_n\|^2.
\end{aligned} \tag{50}$$

Recall that w_n is \mathcal{F}_n -measurable and $\mathbb{E}_n \xi_n = 0$, we have $\mathbb{E}_n \langle w_n, \xi_n \rangle = 0$. So plugging (49) and (50) into (47), and by Cauchy Schwarz inequality, we then have

$$\begin{aligned}
&\mathbb{E}_n \|w_{n+1}\|^2 \\
&= \mathbb{E}_n [\|w_n\|^2 - 2\eta\langle w_n, \nabla F_\lambda(w_n) \rangle + \eta^2 \|\nabla F(w_n) + \lambda w_n - \xi_n\|^2] \\
&\leq \mathbb{E}_n [\|w_n\|^2 - 2\eta\langle w_n, \nabla F_\lambda(w_n) \rangle + 3\eta^2 (\|\nabla F(w_n)\|^2 + \|\xi_n\|^2 + \lambda^2 \|w_n\|^2)] \\
&\leq \|w_n\|^2 - \lambda\eta\|w_n\|^2 + 6\eta^2 \|A\|^2 \|w_n\|^2 + 3\eta^2 \lambda^2 \|w_n\|^2 \\
&\quad + (\eta + 12\|A\|\eta^2) \|w^*\|_A^2 + 3\eta^2 r^2 (1 + c_r \|w_n\|^2).
\end{aligned}$$

Under the condition

$$\eta \leq \frac{\lambda}{12\|A\|^2 + 6\lambda^2 + 6c_r r^2},$$

we have that if $\tau \geq n$

$$\mathbb{E}_n \|w_{n+1}\|^2 \leq (1 - \frac{1}{2}\lambda\eta) \|w_n\|^2 + \eta M_w,$$

which can also leads to

$$\begin{aligned}
\mathbb{E}_n \|w_{n+1}\|^2 \mathbf{1}_{\tau \geq n+1} &\leq \mathbb{E}_n \|w_{n+1}\|^2 \mathbf{1}_{\tau \geq n} \leq (1 - \frac{1}{2}\lambda\eta) \|w_n\|^2 \mathbf{1}_{\tau \geq n} + \eta M_w, \\
\mathbb{E}_n \|w_{n+1}\|^2 \mathbf{1}_{\tau \geq n+1} - \|w_n\|^2 \mathbf{1}_{\tau \geq n} &\leq \eta M_w,
\end{aligned}$$

where

$$M_w := (1 + 12\|A\|\eta) \|w^*\|_A^2 + 3\eta r^2 = \|w^*\|_A^2 + \eta (12\|A\|\|w^*\|_A^2 + 3r^2).$$

Then iterating the inequalities above gives us

$$\mathbb{E}[\|w_n\|^2 \mathbf{1}_{\tau \geq n}] \leq \left(1 - \frac{\lambda\eta}{2}\right)^n \mathbb{E}\|w_0\|^2 + \frac{2}{\lambda} M_w, \quad (51)$$

$$\mathbb{E}[\|w_{n \wedge \tau}\|^2] \leq \mathbb{E}\|w_0\|^2 + \eta n M_w. \quad (52)$$

Step 2: we derive how does the generalization error evolve. According to Taylor's expansion and (46), we know that there exists a v_n , such that

$$\begin{aligned} F(w_{n+1}) &= F(w_n) - \eta \|\nabla F(w_n)\|^2 - \eta \lambda \nabla F(w_n)^T w_n + \eta \xi_n^T \nabla F(w_n) \\ &\quad + \frac{\eta^2}{2} (\nabla F(w_n) + \lambda w_n - \xi_n)^T \nabla^2 F(v_n) (\nabla F(w_n) + \lambda w_n - \xi_n) \\ &\leq F(w_n) - \eta \|\nabla F(w_n)\|^2 - \eta \lambda \nabla F(w_n)^T w_n + \eta \xi_n^T \nabla F(w_n) \\ &\quad + \frac{\eta^2}{2} (\nabla F(w_n) + \lambda w_n - \xi_n)^T A (\nabla F(w_n) + \lambda w_n - \xi_n) \\ &\leq F(w_n) - \eta \|\nabla F(w_n)\|^2 - \eta \lambda \nabla F(w_n)^T (w_n - w^*) + \eta \lambda \|\nabla F(w_n)\|^T w^* \\ &\quad + \eta \xi_n^T \nabla F(w_n) + \frac{3\eta^2}{2} \|A\| (\|\nabla F(w_n)\|^2 + \lambda^2 \|w_n\|^2 + \|\xi_n\|^2). \end{aligned} \quad (53)$$

Step 3: we bound each term in (53) through interpolation. We observe that

$$|\lambda \nabla F(w_n)^T w^*| \leq |\lambda \nabla F(w_n)^T w_\perp^*| + |\lambda \nabla F(w_n)^T w_\lambda^*|, \quad (54)$$

where $w^* = w_\perp^* + w_\lambda^*$ is the decomposition introduced in Definition 2.1. Next

$$|\lambda \nabla F(w_n)^T w_\lambda^*| \leq \frac{1}{2} \|\nabla F(w_n)\|^2 + \frac{1}{2} \lambda^2 \|w_\lambda^*\|^2. \quad (55)$$

Recall $\nabla F(w_n) = B_n(w_n - w_*)$ in (48), and by Lemma A.1 claim 2), we further have

$$\begin{aligned} |\lambda \nabla F(w_n)^T w_\perp^*| &\leq \frac{1}{2} \lambda (w_n - w_*)^T B_n (w_n - w_*) \\ &\quad + \delta \lambda \|w_n - w_*\|_A^2 + \frac{1 + 2\delta}{2} \lambda w_\perp^{*T} A w_\perp^*. \end{aligned} \quad (56)$$

Plugging (55), (56) into (54), applying the result to (53) gives us

$$\begin{aligned}
F(w_{n+1}) &\leq F(w_n) - \frac{1}{2}\eta\|\nabla F(w_n)\|^2 - \frac{1}{2}\lambda\eta\nabla F(w_n)^T(w_n - w_*) + \frac{1+2\delta}{2}\lambda\eta\|w^*\|_{A,\lambda}^2 \\
&\quad + \delta\lambda\eta\|w_n - w^*\|_A^2 + \eta\xi_n^T\nabla F(w_n) + \frac{3\eta^2}{2}\|A\|(\|\nabla F(w_n)\|^2 + \lambda^2\|w_n\|^2 + \|\xi_n\|^2) \\
&\quad (\text{Recall that } \eta \leq 1, \delta < \frac{1}{2} \text{ and } \eta/2 - 3\eta^2\|A\|/2 \geq 0) \\
&\leq F(w_n) - \frac{1}{2}\lambda\eta\nabla F(w_n)^T(w_n - w_*) + \lambda\eta\|w^*\|_{A,\lambda}^2 \tag{57} \\
&\quad + \delta\lambda\eta\|w_n - w^*\|_A^2 + \eta\xi_n^T\nabla F(w_n) + \frac{3\eta^2}{2}\|A\|(\lambda^2\|w_n\|^2 + \|\xi_n\|^2).
\end{aligned}$$

To continue, recall $G(w_n) = F(w_n) - F(w^*)$. Apply fundamental theorem of calculus to $F(w)$, we obtain

$$\begin{aligned}
G(w_n) &= F(w_n) - F(w^*) \\
&= \left[\int_0^1 \nabla F(sw_n + (1-s)w^*) ds \right]^T (w_n - w^*) \\
&= \left[\int_0^1 \left(\nabla F(w^*) + \int_0^s \nabla^2 F(tw_n + (1-t)w^*)(w_n - w^*) dt \right) ds \right]^T (w_n - w^*) \\
&= (w_n - w^*)^T \left[\int_0^1 (1-s)\nabla^2 F(sw_n + (1-s)w^*) ds \right] (w_n - w^*) \\
&= (w_n - w^*)^T A_n (w_n - w^*), \tag{58}
\end{aligned}$$

with

$$A_n = \int_0^1 (1-s)\nabla^2 F(sw_n + (1-s)w^*) ds.$$

Under Assumption 3.1, namely $0 \preceq \nabla^2 F(w_n) + \delta A \preceq A + \delta A$, we observe that

$$\begin{aligned}
\frac{1}{2}\delta A + A_n &= \frac{1}{2}\delta A + \int_0^1 (1-s)\nabla^2 F(sw_n + (1-s)w^*) ds \\
&= \int_0^1 (1-s)(\nabla^2 F(sw_n + (1-s)w^*) + \delta A) ds \\
&\preceq \int_0^1 (\nabla^2 F(sw_n + (1-s)w^*) + \delta A) ds \\
&= B_n + \delta A \preceq (1 + \delta)A.
\end{aligned}$$

Namely, we have

$$A_n \preceq B_n + \frac{1}{2}\delta A \preceq (1 + \frac{\delta}{2})A.$$

Thus

$$G(w_n) - \frac{1}{2}\delta\|w_n - w^*\|_A^2 \leq (w_n - w^*)^T B_n (w_n - w^*) = \nabla F(w_n)^T (w_n - w_*).$$

Plug this into (57), together with $\|w_n - w^*\|_A^2 \leq 2\|w_n\|_A^2 + 2\|w^*\|_A^2 \leq 2\|A\|\|w_n\|^2 + 2\|w^*\|_A^2$, we have

$$\begin{aligned} G(w_{n+1}) &\leq G(w_n) - \frac{1}{2}\eta\lambda G(w_n) + \eta\xi_n^T \nabla F(w_n) \\ &\quad + \frac{5}{4}\delta\lambda\eta\|w_n - w^*\|_A^2 + \lambda\eta\|w^*\|_{A,\lambda}^2 + \frac{3\eta^2}{2}\|A\|(\lambda^2\|w_n\|^2 + \|\xi_n\|^2). \\ &\leq G(w_n) - \frac{1}{2}\eta\lambda G(w_n) + \eta\xi_n^T \nabla F(w_n) \\ &\quad + \lambda\eta(\|w^*\|_{A,\lambda}^2 + \frac{5}{2}\delta\|w^*\|_A^2) + \left(\frac{3\eta^2\lambda^2}{2} + \frac{5}{2}\lambda\eta\delta\right)\|A\|\|w_n\|^2 + \frac{3\eta^2}{2}\|A\|\|\xi_n\|^2. \end{aligned}$$

Step 4: summarizing arguments. We will first establish a rough estimate, which is useful to the escape probability. Since $\mathbb{E}_n \xi_n^T \nabla F(w_n) \equiv 0$, we have

$$\begin{aligned} \mathbb{E}_{n \wedge \tau}[G(w_{n \wedge \tau+1})] &\leq (1 - \frac{1}{2}\eta\lambda)G(w_{n \wedge \tau}) + \left(\frac{3\eta^2\lambda^2}{2} + \frac{5}{2}\lambda\eta\delta\right)\|A\|\|w_{n \wedge \tau}\|^2 \\ &\quad + \lambda\eta\left(\|w^*\|_{A,\lambda}^2 + \frac{5}{2}\delta\|w^*\|_A^2\right) + \frac{3\eta^2\|A\|}{2}r^2(1 + c_r G(w_{n \wedge \tau})) \end{aligned}$$

$$\begin{aligned} \text{Because } \eta &\leq \frac{\lambda}{6c_r\|A\|r^2} \\ &\leq G(w_{n \wedge \tau}) + \left(\frac{3\eta^2\lambda^2}{2} + \frac{5}{2}\lambda\eta\delta\right)\|A\|\|w_{n \wedge \tau}\|^2 \\ &\quad + \lambda\eta\left(\|w^*\|_{A,\lambda}^2 + \frac{5}{2}\delta\|w^*\|_A^2\right) + \frac{3\eta^2\|A\|}{2}r^2. \end{aligned}$$

Recall that $\mathbb{E}[\|w_{n \wedge \tau}\|^2] \leq \mathbb{E}\|w_0\|^2 + \eta n M_w$ with

$$M_w = \|w^*\|_A^2 + \eta(12\|A\|\|w^*\|_A^2 + 3r^2).$$

Iterating above result gives us

$$\begin{aligned} \mathbb{E}[G(w_{n \wedge \tau})] &\leq \mathbb{E}[G(w_0)] + \left(\frac{3\eta^2\lambda^2}{2} + \frac{5}{2}\lambda\eta\delta\right)\|A\|(n\mathbb{E}\|w_0\|^2 + n^2\eta M_w) \quad (59) \\ &\quad + \lambda n \eta \left(\|w^*\|_{A,\lambda}^2 + \frac{5}{2}\delta\|w^*\|_A^2\right) + \frac{3n\eta^2\|A\|}{2}r^2. \end{aligned}$$

We can further improve this bound by using $0 \leq 1_{\tau \geq n+1} \leq 1_{\tau \geq n} \leq 1$ and taking conditional expectation for both sides. Since $\mathbb{E}_n \xi_n^T \nabla F(w_n) \equiv 0$, we have

$$\begin{aligned}
\mathbb{E}[G(w_{n+1})1_{\tau \geq n+1}] &\leq \mathbb{E}[G(w_{n+1})1_{\tau \geq n}] = \mathbb{E}[1_{\tau \geq n} \mathbb{E}_n[G(w_{n+1})]] \\
&\leq (1 - \frac{1}{2}\eta\lambda)\mathbb{E}[G(w_n)1_{\tau \geq n}] + \left(\frac{3\eta^2\lambda^2}{2} + \frac{5}{2}\lambda\eta\delta\right) \|A\| \mathbb{E}[\|w_n\|^2 1_{\tau \geq n}] \\
&\quad + \lambda\eta \left(\|w^*\|_{A,\lambda}^2 + \frac{5}{2}\delta\|w^*\|_A^2\right) + \frac{3\eta^2\|A\|}{2} r^2 (1 + c_r \mathbb{E}[G(w_n)1_{\tau \geq n}])
\end{aligned}$$

Because $\eta \leq \frac{\lambda}{6c_r\|A\|r^2}$

$$\begin{aligned}
&\leq (1 - \frac{1}{4}\eta\lambda)\mathbb{E}[G(w_n)1_{\tau \geq n}] + \left(\frac{3\eta^2\lambda^2}{2} + \frac{5}{2}\lambda\eta\delta\right) \|A\| \mathbb{E}[\|w_n\|^2 1_{\tau \geq n}] \\
&\quad + \lambda\eta \left(\|w^*\|_{A,\lambda}^2 + \frac{5}{2}\delta\|w^*\|_A^2\right) + \frac{3\eta^2\|A\|}{2} r^2. \tag{60}
\end{aligned}$$

Since $\eta\lambda \leq 1$, we have $0 \leq 1 - \frac{1}{4}\eta\lambda \leq \exp(-\frac{1}{4}\eta\lambda)$, then iterating above result gives us

$$\begin{aligned}
\mathbb{E}[G(w_n)1_{\tau \geq n}] &\leq \exp(-\frac{1}{4}\lambda n\eta)\mathbb{E}[G(w_0)] + 4\|w^*\|_{A,\lambda}^2 + 10\delta\|w^*\|_A^2 + \frac{6\eta\|A\|}{\lambda} r^2 \\
&\quad + \left(\frac{3\eta^2\lambda^2}{2} + \frac{5}{2}\lambda\eta\delta\right) \|A\| \sum_{i=0}^n (1 - \frac{1}{4}\eta\lambda)^{n-i} \mathbb{E}[\|w_i\|^2 1_{\tau \geq i}].
\end{aligned}$$

Applying (51), together with $\lambda \leq 1, \eta \leq 1, \delta \leq \frac{1}{2}, 12\eta\|A\| \leq 1$ and $1 - \frac{1}{4}\eta\lambda \leq$

$\exp(-\frac{1}{4}\lambda\eta)$, we obtain

$$\begin{aligned}
& \mathbb{E}[G(w_n)1_{\tau \geq n}] \\
& \leq \exp(-\frac{1}{4}\lambda n\eta)\mathbb{E}[G(w_0)] + 4\|w^*\|_{A,\lambda}^2 + 10\delta\|w^*\|_A^2 + \frac{6\eta\|A\|}{\lambda}r^2 \\
& \quad + \left(\frac{3\eta^2\lambda^2}{2} + \frac{5}{2}\lambda\eta\delta\right)\|A\|\sum_{i=0}^n \left((1 - \frac{1}{4}\lambda\eta)^n \mathbb{E}[\|w_0\|^2] + (1 - \frac{1}{4}\lambda\eta)^{n-i} \frac{2}{\lambda}M_w \right) \\
& \leq \exp(-\frac{1}{4}\lambda n\eta)\mathbb{E}[G(w_0) + 4n\|A\|\|w_0\|^2] + 4\|w^*\|_{A,\lambda}^2 + 10\delta\|w^*\|_A^2 + \frac{6\eta\|A\|}{\lambda}r^2 \\
& \quad + \frac{(12\lambda\eta + 20\delta)\|A\|}{\lambda} (\|w^*\|_A^2 + \eta(12\|A\|\|w^*\|_A^2 + 3r^2)) \\
& \leq \exp(-\frac{1}{4}\lambda n\eta)\mathbb{E}[G(w_0) + 4n\|A\|\|w_0\|^2] + 4\|w^*\|_{A,\lambda}^2 + 10\delta\|w^*\|_A^2 + \frac{6\eta\|A\|}{\lambda}r^2 \\
& \quad + \frac{(12\lambda\eta + 20\delta)\|A\|}{\lambda} (2\|w^*\|_A^2 + 3r^2\eta) \\
& \leq 4\|w^*\|_{A,\lambda}^2 + \frac{C_1}{\lambda}(\eta + \delta) + \exp(-\frac{1}{4}\lambda n\eta)\mathbb{E}[G(w_0) + 4n\|A\|\|w_0\|^2],
\end{aligned}$$

with $C_1 = 60\|A\|(r^2 + \|w^*\|_A^2) + 10\|w^*\|_A^2$.

□

Proof of Corollary 2.5. By Theorem 2.3, $\mathbb{E}[G(\bar{w}_N)1_{\tau \geq N}] \leq 3\epsilon$ holds if we choose λ, η, N, δ such that the following results hold

$$\frac{2\mathbb{E}\|w_0 - w^*\|^2}{N\eta} \leq \epsilon, \quad 8\lambda\|w^*\|^2 \leq \epsilon, \quad 2\eta r^2 \leq \epsilon,$$

and the following conditions are satisfied

$$\eta \leq \left\{ \frac{1}{2(1+c_r)(\|A\|+\lambda)}, 1 \right\}, \quad 2\delta\|A\| \leq \lambda \leq 1.$$

Solving $8\lambda\|w^*\|^2 \leq 8\lambda C_0 \leq \epsilon$ gives us $\lambda(\epsilon) \leq \frac{\epsilon}{8C_0}$. The condition on $\delta(\epsilon)$ is obtained from $\lambda \geq 2\delta\|A\|$. The condition of $\eta(\epsilon)$ ensures that $2\eta r^2 \leq \epsilon$ and $\eta \leq \frac{1}{2(1+c_r)(\|A\|+\lambda)}$. With chosen $\eta(\epsilon)$, the scale of $N(\epsilon)$ is obtained by solving $\frac{2\mathbb{E}\|w_0 - w^*\|^2}{N\eta} \leq \frac{2C_0}{N\eta} \leq \epsilon$. □

Proof of Corollary 2.6. By Theorem 2.4, $\mathbb{E}[G(w_N)1_{\tau \geq N}] \leq 4\epsilon$ holds if we choose λ, η, N, δ such that the following results hold

$$4\|w^*\|_{A,\lambda}^2 \leq \epsilon, \quad \frac{C_1\eta}{\lambda} \leq \epsilon, \quad \frac{C_1\delta}{\lambda} \leq \epsilon, \quad \exp(-\frac{1}{4}\lambda N\eta)\mathbb{E}[G(w_0) + 4N\|A\|\|w_0\|^2] \leq \epsilon.$$

We first choose $\lambda(\epsilon)$ such that $4\|w^*\|_{A,\lambda(\epsilon)}^2 < \epsilon$. The conditions on $\eta(\epsilon), \delta(\epsilon)$ ensure that $\frac{C_1\eta}{\lambda} \leq \epsilon$ and $\frac{C_1\delta}{\lambda} \leq \epsilon$. With chosen $\lambda(\epsilon), \eta(\epsilon)$, the scale of $N(\epsilon)$ is obtained by solving $\exp(-\frac{1}{4}\lambda N\eta)\mathbb{E}[G(w_0)] \leq \frac{\epsilon}{2}$ and $\exp(-\frac{1}{4}\lambda N\eta)4N\|A\|\mathbb{E}[\|w_0\|^2] \leq \frac{\epsilon}{2}$ by using

$$\exp(-\frac{1}{4}\lambda N\eta)N = \frac{4}{\lambda\eta} \exp(-\frac{1}{4}\lambda N\eta)\frac{1}{4}N\lambda\eta \leq \frac{8}{\lambda\eta} \exp(-\frac{1}{8}\lambda N\eta),$$

which is derived from $\exp(-x)x \leq 2\exp(-\frac{1}{2}x)$, since by Taylor expansion $x \leq 2\exp(\frac{x}{2})$. \square

B Proof for Results in Low Effective Dimension in Section 3.2

Proof of Proposition 3.2. Under Assumption 3.1, applying Corollary 2.5 results in

$$\lambda(\epsilon) = O(\epsilon), \quad \delta(\epsilon) = O(\epsilon), \quad \eta(\epsilon) = O(\epsilon), \quad N(\epsilon) = \Omega\left(\frac{1}{\epsilon^2}\right),$$

for guaranteeing $\mathbb{E}[G(\bar{w}_N)1_{\tau \geq N}] \leq 3\epsilon$. In this case, by (43), according to Chebyshev's inequality and recall that

$$\mathbb{E}\|w_{N \wedge \tau} - w^*\|^2 \leq \mathbb{E}\|w_0 - w^*\|^2 + N(4\lambda\eta\|w^*\|^2 + \eta^2 r^2) = O(1),$$

we have

$$\mathbf{P}(\tau \leq N) = \mathbf{P}(\{w_{N \wedge \tau} \notin \mathcal{D}\}) = \mathbf{P}(\{\|w_{N \wedge \tau} - w^*\|^2 > A\}) \leq \frac{\mathbb{E}\|w_{N \wedge \tau} - w^*\|^2}{A} \leq \delta.$$

If $\delta = 0$, according to Theorem 2.3,

$$\mathbb{E}[1_{\tau \geq N-1}G(\bar{w}_N)] \leq \frac{2\mathbb{E}\|w_0 - w^*\|^2}{N\eta} + 8\lambda\|w^*\|^2 + 2\eta r^2,$$

taking

$$\lambda(\epsilon) = 0, \quad \eta(\epsilon) = O(\epsilon^{1+\alpha}), \quad N(\epsilon) = \Omega\left(\frac{1}{\epsilon^{2+\alpha}}\right),$$

we obtain $\mathbb{E}[G(\bar{w}_N)1_{\tau \geq N}] \leq 3\epsilon$. In this case, according to Chebyshev's inequality and recall that

$$\mathbb{E}\|w_{N \wedge \tau} - w^*\|^2 \leq \mathbb{E}\|w_0 - w^*\|^2 + N(4\lambda\eta\|w^*\|^2 + \eta^2 r^2) = \mathbb{E}\|w_0 - w^*\|^2 + O(\epsilon^\alpha),$$

we have

$$\mathbf{P}(\tau \leq N) = \mathbf{P}(\{w_{N \wedge \tau} \notin \mathcal{D}\}) = \mathbf{P}(\{\|w_{N \wedge \tau} - w^*\|^2 > A\}) \leq \frac{\mathbb{E}\|w_0 - w^*\|^2 + O(\epsilon^\alpha)}{A} < 1,$$

if $A > \mathbb{E}\|w_0 - w^*\|^2$. \square

Proof of Proposition 3.4. Recall that (λ_i, v_i) , for $i = 1, \dots, p$, are the eigenvalue-eigenvectors of A with λ_i decreasingly sorted. Therefore, we have

$$\begin{aligned} \|w^*\|_A^2 &= w^{*T} A w^* = \sum_{i=1}^p \lambda_i \langle w^*, v_i \rangle^2 \leq \|w^*\|_{A,S}^2 \text{tr}(A), \\ \|w^*\|_{A,\lambda}^2 &= \sum_{i=1}^p \lambda_i \wedge \lambda \langle w^*, v_i \rangle^2 \leq \|w^*\|_{A,S}^2 \sum_{i=1}^p \lambda_i \wedge \lambda. \end{aligned}$$

For an exponential spectrum, given any k and p ,

$$\sum_{i=k+1}^p \lambda_i = \sum_{i=k+1}^p e^{-ci} = \frac{e^{-(k+1)c}(1 - e^{-(p-k)c})}{1 - e^{-c}} \leq \frac{1}{e^{kc}(e^c - 1)}.$$

Thus, to make $\sum_{i=k+1}^p \lambda_i \leq \frac{\epsilon}{8} \|w^*\|_{A,S}^2$, it is sufficient for us to take $k \geq \frac{1}{c} \log \left\{ \frac{8 \|w^*\|_{A,S}^2}{\epsilon(e^c - 1)} \right\}$. And to make $k\lambda = \frac{\epsilon}{8 \|w^*\|_{A,S}^2}$, we take $\lambda = \frac{\epsilon}{8k \|w^*\|_{A,S}^2} = \tilde{O}\left(\frac{\epsilon}{|\log \epsilon|}\right)$. By these choices, we have

$$\|w^*\|_{A,\lambda}^2 \leq \sum_{i=1}^p \lambda \wedge \lambda_i \|w^*\|_{A,S}^2 \leq \frac{\epsilon}{4}.$$

Next, we find that $\|A\| \leq \text{tr}(A) = \tilde{O}(1)$, so $C_1 = \tilde{O}(1), C_2 = \tilde{O}(1)$. We implement Corollary 2.6 and find

$$\delta(\epsilon) = \tilde{O}\left(\frac{\epsilon^3}{|\log \epsilon|^2}\right), \quad \eta(\epsilon) = \tilde{O}\left(\frac{\epsilon^2}{|\log \epsilon|}\right), \quad N(\epsilon) = \tilde{\Omega}\left(\frac{|\log \epsilon|^3}{\epsilon^3}\right).$$

Recall that

$$\begin{aligned} \mathbb{E}[G(w_{n \wedge \tau})] &\leq \mathbb{E}[G(w_0)] + \left(\frac{3\eta^2 \lambda^2}{2} + \frac{5}{2} \lambda \eta \delta\right) n^2 \eta C_1 \\ &\quad + \lambda n \eta \left(\|w^*\|_{A,\lambda}^2 + \frac{5}{2} \delta \|w^*\|_A^2\right) + \frac{3n\eta^2 \|A\|}{2} r^2, \end{aligned}$$

together with

$$\left(\frac{3\eta^2\lambda^2}{2} + \frac{5}{2}\lambda\eta\delta\right) N^2\eta = \tilde{O}(\epsilon^2|\log \epsilon|), \quad \lambda N\eta \left(\|w^*\|_{A,\lambda}^2 + \frac{5}{2}\delta\|w^*\|_A^2\right) = \tilde{O}(\epsilon|\log \epsilon|),$$

$$N\eta^2\frac{3\|A\|r^2}{2} = \tilde{O}(\epsilon|\log \epsilon|),$$

we have

$$\mathbb{E}[G(w_{N\wedge\tau})] \leq \mathbb{E}[G(w_0)] + \tilde{O}(\epsilon|\log \epsilon|).$$

For a polynomial spectrum, the derivation is similar. Given any k and p

$$\sum_{i=k+1}^p \lambda_i = \sum_{i=k+1}^p i^{-(1+c)} \leq \sum_{i=k+1}^p \int_{i-1}^i \frac{1}{x^{1+c}} dx = \sum_{i=k+1}^p \frac{-1}{c} x^{-c} \Big|_{i-1}^i = \frac{1}{c}(k^{-c} - p^{-c}) \leq \frac{1}{ck^c}.$$

Thus to make $\|w^*\|_{A,S}^2 \sum_{i=k+1}^p \lambda_i \leq \frac{1}{8}\epsilon$, we take $k \geq (\frac{8\|w^*\|_{A,S}^2}{c\epsilon})^{1/c}$. Next, we take $\lambda(\epsilon) = \frac{\epsilon}{8\|w^*\|_{A,S}^2} = \tilde{O}(\epsilon^{\frac{c+1}{c}})$. This leads to $\|w^*\|_{A,\lambda}^2 \leq \epsilon/4$. Again we find that $C_1 = \tilde{O}(1), C_2 = \tilde{O}(1)$. The order of $\delta(\epsilon), \eta(\epsilon)$ and $N(\epsilon)$ can be derived by Corollary 2.6, that is,

$$\delta(\epsilon) = \tilde{O}\left(\epsilon^{\frac{3c+2}{c}}\right), \quad \eta(\epsilon) = \tilde{O}\left(\epsilon^{\frac{2c+1}{c}}\right), \quad N(\epsilon) = \tilde{\Omega}\left(\frac{|\log(\epsilon)|}{\epsilon^{\frac{3c+2}{c}}}\right).$$

Recall that

$$\mathbb{E}[G(w_{n\wedge\tau})] \leq \mathbb{E}[G(w_0)] + \left(\frac{3\eta^2\lambda^2}{2} + \frac{5}{2}\lambda\eta\delta\right) n^2\eta C_1$$

$$+ \lambda n\eta \left(\|w^*\|_{A,\lambda}^2 + \frac{5}{2}\delta\|w^*\|_A^2\right) + \frac{3n\eta^2\|A\|r^2}{2},$$

together with

$$\left(\frac{3\eta^2\lambda^2}{2} + \frac{5}{2}\lambda\eta\delta\right) N^2\eta = \tilde{O}(\epsilon^{\frac{2c+1}{c}}(\log \epsilon)^2), \quad \lambda N\eta = \tilde{O}(|\log \epsilon|),$$

$$\|w^*\|_{A,\lambda}^2 + \frac{5}{2}\delta\|w^*\|_A^2 = O(\epsilon), \quad N\eta^2\frac{3\|A\|r^2}{2} = \tilde{O}(\epsilon|\log \epsilon|),$$

we have

$$\mathbb{E}[G(w_{N\wedge\tau})] \leq \mathbb{E}[G(w_0)] + \tilde{O}(\epsilon|\log \epsilon|).$$

Given $\mathbb{E}[G(w_{N \wedge \tau})] \leq \mathbb{E}[G(w_0)] + \tilde{O}(\epsilon |\log \epsilon|)$ for both cases, according to Chebyshev's inequality, we have

$$\begin{aligned} \mathbf{P}(\tau \leq N) &= \mathbf{P}(\{w_{N \wedge \tau} \notin \mathcal{D}\}) = \mathbf{P}(\{G(w_{N \wedge \tau}) > (1+a)\mathbb{E}[G(w_0)]\}) \\ &\leq \frac{\mathbb{E}[G(w_{N \wedge \tau})]}{(1+a)\mathbb{E}[G(w_0)]} \leq \frac{\mathbb{E}[G(w_0)] + \tilde{O}(\epsilon |\log \epsilon|)}{(1+a)\mathbb{E}[G(w_0)]} \leq \frac{1}{1+a} + \tilde{O}(\epsilon |\log \epsilon|). \end{aligned}$$

□

C Proofs of Results for Overparameterization in Statistical Models

C.1 Linear regression

Proof of Proposition 4.1. It is straightforward to find the gradient and Hessian of F as:

$$\nabla F(w) = \Sigma(w - w^*), \quad \nabla^2 F(w) = \Sigma. \quad (61)$$

This leads to $A = \Sigma$, $\delta = 0$, $\mathcal{D} = \mathbb{R}^p$.

Next, note that

$$\nabla f(w, \zeta) = (x^T w - y)x = (x^T(w - w^*) - \xi)x.$$

By Cauchy Schwarz inequality, we have

$$\begin{aligned} \mathbb{E}\|\nabla f(w, \zeta) - \nabla F(w)\|^2 &\leq \mathbb{E}\|\nabla f(w, \zeta)\|^2 \\ &\leq 2\mathbb{E}\|xx^T(w - w^*)\|^2 + 2\mathbb{E}\|x\xi\|^2 \\ &= 2(w - w^*)^T \mathbb{E}[xx^T xx^T](w - w^*) + 2\sigma^2 \text{tr}(\Sigma). \end{aligned} \quad (62)$$

Next we compute $\mathbb{E}[xx^T xx^T]$. Since $x \sim \mathcal{N}(0, \Sigma)$, it can be decomposed as $x = \Sigma^{1/2}z$ with $z \sim \mathcal{N}(0, I_p)$. Let the eigen-decomposition of Σ be $V^T \Lambda V$ and denote $\Sigma^{1/2} = V^T \Lambda^{1/2} V$. We notice that $z' = Vz \sim \mathcal{N}(0, I_p)$, then the (i, j) -th element of $Vz z^T V^T \Lambda Vz z^T V^T$ is $\sum_{k=1}^p \lambda_k z'_i z'_j (z'_k)^2$, and taking expectation results in

$$\mathbb{E}[Vz z^T V^T \Lambda Vz z^T V^T] = \text{diag} \left[2\lambda_1 + \sum_{j=1}^p \lambda_j, \dots, 2\lambda_j + \sum_{j=1}^p \lambda_j, \dots, 2\lambda_p + \sum_{j=1}^p \lambda_j \right].$$

Thus we have

$$\begin{aligned}
\mathbb{E}[xx^T xx^T] &= V^T \Lambda^{1/2} \mathbb{E}[V z z^T V^T \Lambda V z z^T V^T] \Lambda^{1/2} V \\
&= V^T \Lambda^{1/2} \text{diag} \left[2\lambda_1 + \sum_{j=1}^p \lambda_j, \dots, 2\lambda_j + \sum_{j=1}^p \lambda_j, \dots, 2\lambda_p + \sum_{j=1}^p \lambda_j \right] \Lambda^{1/2} V \\
&\leq 3\text{tr}(\Sigma)\Sigma.
\end{aligned}$$

Plugging this upper bound in (62) gives us

$$\mathbb{E}\|\nabla f(w, \zeta) - \nabla F(w)\|^2 \leq 6\text{tr}(\Sigma)\|w - w^*\|_\Sigma^2 + 2\sigma^2\text{tr}(\Sigma).$$

Finally, we note that $\|w - w^*\|_\Sigma^2 = 2G(w)$, and by Young's inequality

$$\|w - w^*\|_\Sigma^2 \leq 2\|w\|_\Sigma^2 + 2\|w^*\|_\Sigma^2 \leq 2\|\Sigma\|\|w\|^2 + 2\|w^*\|_\Sigma^2,$$

$$\|w - w^*\|_\Sigma^2 = (w - w^*)^T \Sigma (w - w^*) = \frac{1}{2}(w - w^*)^T \nabla F(w).$$

Therefore, we conclude that

$$\mathbb{E}\|\nabla f(w, \zeta) - \nabla F(w)\|^2 \leq 2\sigma^2\text{tr}(\Sigma) + 12\text{tr}(\Sigma)\|w^*\|_\Sigma^2 + 12\text{tr}(\Sigma) \min\{G(w), \|\Sigma\|\|w\|^2\}.$$

□

Remark C.1. *In the proof above, we used the Gaussian distribution assumption only to obtain the first, second and fourth moments of x . This proof can be extended to scenarios where x has a non-Gaussian distribution, as long as an upper bound of $\mathbb{E}[xx^T xx^T]$ is available. Similar extensions can be made for other proofs below as well.*

C.2 Logistic regression

Proof for Proposition 5.1. By Fubini's theorem,

$$\nabla F(w) = \mathbb{E}\nabla f(w, \zeta) = \mathbb{E}\frac{-yx}{1 + \exp(yx^T w)},$$

and

$$\nabla^2 F(w) = \mathbb{E}\nabla \frac{-yx}{1 + \exp(yx^T w)} = \mathbb{E}\frac{y^2 \exp(yx^T w) xx^T}{(1 + \exp(yx^T w))^2}.$$

Because $0 < \frac{y^2 \exp(yx^T w)}{(1 + \exp(yx^T w))^2} < 1$ and $0 \preceq xx^T$, we find $0 \preceq \nabla^2 F(w) \preceq \Sigma$.

Next, we observe

$$\nabla f(w, \zeta) = \frac{-yx}{1 + \exp(yx^T w)}.$$

Then, because $y = \pm 1$, we obtain

$$\mathbb{E}[\|\nabla f(w, \zeta)\|^2] = \mathbb{E}\left[\left(\frac{-y}{1 + \exp(yx^T w)}\right)^2 \|x\|^2\right] \leq \mathbb{E}[\|x\|^2] = \text{tr}(\Sigma).$$

□

C.3 M -estimator with Tukey's biweight loss

Proof for Proposition 5.2. First of all, let $v = w - w^*$, $u = x^T v - \xi$. We find that

$$\nabla f(w, \zeta) = (1 - (u/c)^2)^2 ux 1_{|u| \leq c}.$$

Then, by Fubini theorem, we have

$$\begin{aligned} \nabla F(w) &= \nabla_v F(w) = \mathbb{E} \nabla[\rho(x^T v - \xi)] = \mathbb{E}[(1 - (u/c)^2)^2 ux 1_{\{|u| \leq c\}}], \\ \nabla^2 F(w) &= \mathbb{E}[xx^T (1 - (u/c)^2)(1 - 5(u/c)^2) 1_{|u| \leq c}]. \end{aligned}$$

For the first two claims, note that

$$\begin{aligned} \mathbb{E}\|\nabla f(w, \zeta)\|^2 &= \mathbb{E}[(1 - (u/c)^2)^4 (u/c)^2 1_{|u| \leq c} \|x\|^2] \leq \mathbb{E}\|x\|^2 = \text{tr}(\Sigma), \\ \nabla^2 F(w) &= \mathbb{E}[xx^T (1 - (u/c)^2)(1 - 5(u/c)^2) 1_{|u| \leq c}] \preceq \mathbb{E}xx^T = \Sigma. \end{aligned}$$

At $w = w^*$,

$$\nabla^2 F(w^*) = \mathbb{E}[xx^T (1 - (\xi/c)^2)(1 - 5(\xi/c)^2) 1_{|\xi| \leq c}] = c_0 \Sigma.$$

We consider the directional derivative along the v direction

$$\begin{aligned} \langle v, \nabla^3 F(w) \rangle &:= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} (\nabla^2 F(w + \epsilon v) - \nabla^2 F(w)) \\ &= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} (\mathbb{E}[xx^T (1 - (u/c + \epsilon x^T v)^2)(1 - 5(u/c + \epsilon x^T v)^2) 1_{|u| \leq c}] \\ &\quad - \mathbb{E}[xx^T (1 - (u/c)^2)(1 - 5(u/c)^2) 1_{|u| \leq c}]) \\ &= \mathbb{E}[4xx^T x^T v (3u/c - 5(u/c)^3) 1_{|u| \leq c}]. \end{aligned}$$

We find

$$\begin{aligned} \pm \langle v, \nabla^3 F(w) \rangle &= \pm \mathbb{E}[4xx^T x^T v (3u/c - 5(u/c)^3) 1_{|u| \leq c}] \\ &\preceq \mathbb{E}[4xx^T |x^T v| |5(u/c)^3 - 3u/c| 1_{|u| \leq c}] \preceq \mathbb{E}[8xx^T |x^T v|]. \end{aligned}$$

For any test vector ψ ,

$$\begin{aligned} |\psi^T \langle v, \nabla^3 F(w) \rangle \psi| &\leq 8\mathbb{E}[(x^T \psi)^2 |x^T v|] \leq 8\sqrt{\mathbb{E}[(x^T \psi)^4] \mathbb{E}[|x^T v|^2]} \\ &= 8\sqrt{3(\psi^T \Sigma \psi)^2 (v^T \Sigma v)} \leq 16\|v\|_{\Sigma} \psi^T \Sigma \psi. \end{aligned}$$

Therefore,

$$-16\|v\|_{\Sigma} \Sigma \preceq \langle v, \nabla^3 F(w) \rangle \preceq 16\|v\|_{\Sigma} \Sigma.$$

Furthermore, since $w = w^* + v$, from

$$\nabla^2 F(v + w^*) = \nabla^2 F(w^*) + \int_0^1 \langle v, \nabla^3 F(w^* + sv) \rangle ds,$$

we find

$$\nabla^2 F(w) \succeq -\delta \Sigma,$$

if $16\|v\|_{\Sigma} \leq c_0 + \delta$. □

C.4 Two-layer neural network

First of all, we provide a simple upper bound when computing the fourth order moments of Gaussian random variables.

Lemma C.2. *If $x \in \mathbb{R}^p$ is Gaussian with mean being zero, for any PSD $A \in \mathbb{R}^{p \times p}$ and $a > 0$*

$$\mathbb{E}(x^T A x + a)^2 \leq 3(\mathbb{E}(x^T A x + a))^2.$$

Proof. Let Σ be the covariance matrix of x . Since replacing x with $\Sigma^{-1/2}x$, the statement of the Lemma remains the same, therefore we can assume $x \sim \mathcal{N}(0, I_p)$. Let $A = V^T \Lambda V$ be the eigenvalue decomposition of A , and the eigenvalues of A be $\lambda_1, \dots, \lambda_p$. Let $z = Vx \sim \mathcal{N}(0, I_p)$. Note that

$$\mathbb{E}(x^T A x + a)^2 = \mathbb{E}(\|z\|_{\Lambda}^4 + 2a\|z\|_{\Lambda}^2 + a^2),$$

and further,

$$\mathbb{E}\|z\|_{\Lambda}^4 = \sum_{i,j} \lambda_i \lambda_j \mathbb{E}(z_i^2 z_j^2) \leq 3 \sum_{i,j} \lambda_i \lambda_j \mathbb{E}z_i^2 \mathbb{E}z_j^2 = 3(\mathbb{E}\|z\|_{\Lambda}^2)^2.$$

As a consequence, we obtain

$$\mathbb{E}(x^T A x + a)^2 \leq 3(\mathbb{E}\|z\|_{\Lambda}^2 + a)^2 = 3(\mathbb{E}(x^T A x + a))^2. \quad \square$$

Lemma C.3. Assume that $\psi(0) = 0$ and $|\dot{\psi}|, |\ddot{\psi}| \leq C$. Denote

$$\Sigma^* = \text{diag}\{I_k, \Sigma, \dots, \Sigma, I_k\} \in \mathbb{R}^{(p+2)k \times (p+2)k},$$

and $\Delta w = w - w^*$. Then the followings hold

$$1) \mathbb{E}\|\nabla f(w)\|^2 \leq 8\sqrt{3}(1+\text{tr}(\Sigma))(6C^2\|\Delta w\|_{\Sigma^*}^2(\|w^*\|_{\Sigma^*}^2 + \|w\|_{\Sigma^*}^2) + \sigma_0^2)C^2\|w\|_{\Sigma^*}^2.$$

$$2) \mathbb{E}\nabla g(w, x)\nabla g(w, x)^T \preceq 6C^2\|w\|_{\Sigma^*}^2\Sigma^*.$$

$$3) -M_w \preceq \mathbb{E}(g(w, x) - g(w^*, x) - \xi)\nabla^2 g \preceq M_w, \text{ where}$$

$$M_w := 6\sqrt{2}C^2(\|c\|_\infty + 1)\|\Delta w\|_{\Sigma^*}(\|w^*\|_{\Sigma^*} + \|w\|_{\Sigma^*})\Sigma^*,$$

$$\text{with } \|c\|_\infty := \max_i \{c_i\}.$$

$$4) G(w) \leq 6C^2\|\Delta w\|_{\Sigma^*}^2(\|w^*\|_{\Sigma^*}^2 + \|w\|_{\Sigma^*}^2).$$

Proof. For simplicity of discussion, we denote $z_i = b_i^T x + a_i$ and $z = bx + a$.

Proof for Claim 1) We note that $\nabla f(w) = 2(g(w, x) - g(w^*, x) - \xi)\nabla g(w, x)$, thus

$$\begin{aligned} \mathbb{E}\|\nabla f(w)\|^2 &= 4\mathbb{E}[(g(w, x) - g(w^*, x))^2\|\nabla g(w, x)\|^2] + 4\sigma_0^2\mathbb{E}\|\nabla g(w, x)\|^2 \\ &\leq 4\sqrt{\mathbb{E}(g(w, x) - g(w^*, x))^4}\sqrt{\mathbb{E}\|\nabla g(w, x)\|^4} + 4\sigma_0^2\sqrt{\mathbb{E}\|\nabla g(w, x)\|^4}. \end{aligned} \quad (63)$$

Note that

$$\nabla g = [c \circ \dot{\psi}(z); c_1 \dot{\psi}(z_1)x; \dots; c_k \dot{\psi}(z_k)x; \psi(z)]^T \in \mathbb{R}^{2k+kp},$$

as a consequence, we have

$$\begin{aligned} \mathbb{E}\|\nabla g(w, x)\|^4 &= \mathbb{E}\left(\|c \circ \dot{\psi}(z)\|^2 + \sum_{i=1}^k \|c_i \dot{\psi}(z_i)x\|^2 + \sum_{i=1}^k \|\psi(z_i)\|^2\right)^2 \\ &\leq \mathbb{E}\left(C^2\|c\|^2 + \sum_{i=1}^k (c_i)^2 C^2\|x\|^2 + 2C^2 \sum_{i=1}^k (b_i^T x)^2 + 2C^2\|a\|^2\right)^2 \end{aligned}$$

Since x is mean zero Gaussian, by Lemma C.2

$$\begin{aligned} &\leq 3\left(C^2\|c\|^2 + \sum_{i=1}^k (c_i)^2 C^2\mathbb{E}\|x\|^2 + 2C^2\mathbb{E}\sum_{i=1}^k (b_i^T x)^2 + 2C^2\|a\|^2\right)^2 \\ &\leq 3C^4\left(\|c\|^2 + \|c\|^2\text{tr}(\Sigma) + 2\sum_{i=1}^k \|b_i\|_\Sigma^2 + 2\|a\|^2\right)^2 \\ &\leq 12C^4(1 + \text{tr}(\Sigma))^2\|w\|_{\Sigma^*}^4. \end{aligned} \quad (64)$$

Next, we let $w^s = sw + (1 - s)w^*$ and $C_w^2 = \|w\|_{\Sigma^*}^2 + \|w^*\|_{\Sigma^*}^2$. By the convexity of $\|\cdot\|_{\Sigma^*}^2$, we get

$$\|w^s\|_{\Sigma^*}^4 \leq \max\{\|w\|_{\Sigma^*}^4, \|w^*\|_{\Sigma^*}^4\} \leq \|w\|_{\Sigma^*}^4 + \|w^*\|_{\Sigma^*}^4 \leq C_w^4.$$

Then, we have

$$\begin{aligned} |g(w, x) - g(w^*, x)|^2 &= \left(\int_0^1 \Delta w^T \nabla g(w^s, x) ds \right)^2 \\ &\leq \int_0^1 \left(\Delta a^T c^s \circ \psi(z^s) + \sum_{i=1}^k c_i^s \psi(z_i^s) \Delta b_i^T x + \Delta c^T \psi(z^s) \right)^2 ds \\ &\leq \int_0^1 \left(C \|\Delta a\| \|c^s\| + C \sum_{i=1}^k |c_i^s| |\Delta b_i^T x| + \|\Delta c\| \|\psi(z^s)\| \right)^2 ds \\ &\leq \int_0^1 \left(C_w^2 \|\Delta a\|^2 + C_w^2 \sum_{i=1}^k |\Delta b_i^T x|^2 + \|\Delta c\|^2 \|\psi(z^s)\|^2 / C^2 \right) ds \\ &\quad \cdot \int_0^1 \left(\frac{C^2 \|c^s\|^2}{C_w^2} + \frac{C^2}{C_w^2} \sum_{i=1}^k |c_i^s|^2 + C^2 \right) ds \\ &\leq \int_0^1 \left(C_w^2 \|\Delta a\|^2 + C_w^2 \sum_{i=1}^k |\Delta b_i^T x|^2 + \|\Delta c\|^2 \|\psi(z^s)\|^2 / C^2 \right) ds \\ &\quad \cdot \int_0^1 (2C^2 \|c^s\|^2 / C_w^2 + C^2) ds \\ &\leq 3C^2 \int_0^1 \left(C_w^2 \|\Delta a\|^2 + C_w^2 \sum_{i=1}^k |\Delta b_i^T x|^2 + \|\Delta c\|^2 \|\psi(z^s)\|^2 / C^2 \right) ds. \quad (65) \end{aligned}$$

By Lemma C.2 and $\mathbb{E}|\Delta b_i^T x|^2 = \Delta b_i^T \Sigma \Delta b_i$, we have

$$\begin{aligned}
& \mathbb{E} \left(C_w^2 \|\Delta a\|^2 + C_w^2 \sum_{i=1}^k |\Delta b_i^T x|^2 + \|\Delta c\|^2 \|\psi(z^s)\|^2 / C^2 \right)^2 \\
& \leq \mathbb{E} \left(C_w^2 \|\Delta a\|^2 + C_w^2 \sum_{i=1}^k |\Delta b_i^T x|^2 + 2\|\Delta c\|^2 (\|a^s\|^2 + \sum_{i=1}^k |(b_i^s)^T x|^2) \right)^2 \\
& \text{Note that } \|a^s\|^2 + \sum_{i=1}^k |(b_i^s)^T x|^2 \leq \max\{\|w\|_{\Sigma^*}^2, \|w^*\|_{\Sigma^*}^2\} \leq C_w^2 \\
& \leq \left(C_w^2 \|\Delta a\|^2 + C_w^2 \sum_{i=1}^k \|\Delta b_i\|_{\Sigma}^2 + 2C_w^2 \|\Delta c\|^2 \right)^2 \\
& \leq 4(\|w\|_{\Sigma^*}^2 + \|w^*\|_{\Sigma^*}^2)^2 \|\Delta w\|_{\Sigma^*}^4.
\end{aligned}$$

Replace these bounds into the square of (65), we find

$$\mathbb{E}|g(w, x) - g(w^*, x)|^4 \leq 36C^4 \|\Delta w\|_{\Sigma^*}^4 (\|w^*\|_{\Sigma^*}^2 + \|w\|_{\Sigma^*}^2)^2. \quad (66)$$

Furthermore, we combine this with (64) into (63), we find that

$$\mathbb{E}\|\nabla f(w)\|^2 \leq 8\sqrt{3}(1 + \text{tr}(\Sigma))(6C^2 \|\Delta w\|_{\Sigma^*}^2 (\|w^*\|_{\Sigma^*}^2 + \|w\|_{\Sigma^*}^2) + \sigma_0^2) C^2 \|w\|_{\Sigma^*}^2.$$

Proof for Claim 2): Recall that

$$\nabla g = [c \circ \dot{\psi}(z); c_1 \dot{\psi}(z_1)x; \dots; c_k \dot{\psi}(z_k)x; \psi(z)] \in \mathbb{R}^{2k+kp}.$$

With $u \in \mathbb{R}^k$, $v_1 \in \mathbb{R}^p, \dots, v_k \in \mathbb{R}^p$, $w \in \mathbb{R}^k$, we define

$$W = [u; v_1; \dots; v_k; w] \in \mathbb{R}^{2k+kp},$$

and show that $W^T \mathbb{E} \nabla g \nabla g^T W \preceq 6C^2 \|w\|_{\Sigma^*}^2 W^T \Sigma^* W$. Note that

$$\begin{aligned}
& W^T \mathbb{E} \nabla g \nabla g^T W \tag{67} \\
& = \mathbb{E} u^T [c \circ \dot{\psi}(z) (c \circ \dot{\psi}(z))^T] u + v_i^T \mathbb{E} [c_i^2 \dot{\psi}(z_i) \dot{\psi}(z_i) x x^T] v_i + w^T \mathbb{E} [\psi(z) (\psi(z))^T] w \\
& \quad + 2u^T \mathbb{E} c \circ \dot{\psi}(z) (\psi(z))^T w + 2 \sum_{i=1}^k u^T \mathbb{E} c \circ \dot{\psi}(z) c_i \dot{\psi}(z_i) x^T v_i \\
& \quad + 2 \sum_{i < j} v_i^T \mathbb{E} c_i \dot{\psi}(z_i) c_j \dot{\psi}(z_j) x x^T v_j + 2 \sum_{i=1}^k v_i^T \mathbb{E} c_i \dot{\psi}(z_i) x (\psi(z))^T w.
\end{aligned}$$

For the diagonal terms, note that

$$\begin{aligned}\mathbb{E}[c \circ \dot{\psi}(z)(c \circ \dot{\psi}(z))^T] &\preceq \mathbb{E}[\|c \circ \dot{\psi}(z)\|^2 I_k] \preceq C^2 \|c\|^2 I_k, \\ \mathbb{E}[c_i^2 \dot{\psi}(z_i) \dot{\psi}(z_i) x x^T] &\preceq C^2 c_i^2 \mathbb{E}[x x^T] = C^2 c_i^2 \Sigma, \\ \mathbb{E}[\psi(z)(\psi(z))^T] &\preceq \mathbb{E}[\|\psi(z)\|^2 I_k] \preceq 2C^2 \left(\|a\|^2 + \sum_{i=1}^k \|b_i\|_\Sigma^2 \right) I_k.\end{aligned}$$

For the cross terms, note that by Cauchy Schwarz inequality

$$\begin{aligned}u^T c \circ \dot{\psi}(z) c_i \dot{\psi}(z_i) x x^T v_i &\leq |u^T c \circ \dot{\psi}(z)| |c_i \dot{\psi}(z_i) x x^T v_i| \\ &= (u^T c \circ \dot{\psi}(z) (c \circ \dot{\psi}(z))^T u)^{1/2} (v_i^T (c_i \dot{\psi}(z_i))^2 x x^T v_i)^{1/2} \\ &\leq \frac{c_i^2}{2\|c\|^2} (u^T c \circ \dot{\psi}(z) (c \circ \dot{\psi}(z))^T u) + \frac{\|c\|^2}{2} (v_i^T (\dot{\psi}(z_i))^2 x x^T v_i),\end{aligned}$$

and similarly,

$$\begin{aligned}u^T c \circ \dot{\psi}(z) (\psi(z))^T w &\leq \frac{1}{2} u^T c \circ \dot{\psi}(z) (c \circ \dot{\psi}(z))^T u + \frac{1}{2} w^T \psi(z) (\psi(z))^T w, \\ v_i^T c_i \dot{\psi}(z_i) c_j \dot{\psi}(z_j) x x^T v_j &\leq \frac{c_j^2}{2} v_i^T (\dot{\psi}(z_i))^2 x x^T v_i + \frac{c_i^2}{2} v_j^T (\dot{\psi}(z_j))^2 x x^T v_j, \\ v_i^T c_i \dot{\psi}(z_i) x (\psi(z))^T w &\leq \frac{\|c\|^2}{2} v_i^T (\dot{\psi}(z_i))^2 x x^T v_i + \frac{c_i^2}{2\|c\|^2} w^T (\psi(z)) (\psi(z))^T w.\end{aligned}$$

Plugging the results above into (67) gives us

$$\begin{aligned}\mathbb{E} \nabla g(w, x) \nabla g(w, x)^T &\preceq C^2 \begin{bmatrix} 2\|c\|^2 I_k & \mathbf{0}_{k \times p} & \mathbf{0}_{k \times p} & \mathbf{0}_{k \times p} & \mathbf{0}_{k \times k} \\ \mathbf{0}_{p \times k} & 3\|c\|^2 \Sigma & \mathbf{0}_{p \times p} & \mathbf{0}_{p \times p} & \mathbf{0}_{p \times k} \\ \mathbf{0}_{p \times k} & \mathbf{0}_{p \times p} & \ddots & \mathbf{0}_{p \times p} & \mathbf{0}_{p \times k} \\ \mathbf{0}_{p \times k} & \mathbf{0}_{p \times p} & \mathbf{0}_{p \times p} & 3\|c\|^2 \Sigma & \mathbf{0}_{p \times k} \\ \mathbf{0}_{k \times k} & \mathbf{0}_{k \times p} & \mathbf{0}_{k \times p} & \mathbf{0}_{k \times p} & 6 \left(\|a\|^2 + \sum_{i=1}^k \|b_i\|_\Sigma^2 \right) I_k \end{bmatrix} \\ &\preceq 6C^2 \|w\|_{\Sigma^*}^2 \Sigma^*.\end{aligned}$$

Proof for Claim 3): First of all, we find that

$$\nabla^2 g = \begin{bmatrix} D_{c \circ \ddot{\psi}(z)} & c_1 \ddot{\psi}(z_1) e_1 x^T & c_2 \ddot{\psi}(z_2) e_2 x^T & \cdots & c_k \dot{\psi}(z_k) x e_k^T & D_{\dot{\psi}(z)} \\ c_1 \ddot{\psi}(z_1) x e_1^T & c_1 \ddot{\psi}(z_1) x x^T & \mathbf{0}_{p \times p} & \cdots & \mathbf{0}_{p \times p} & \dot{\psi}(z_1) x e_1^T \\ c_2 \ddot{\psi}(z_2) x e_2^T & \mathbf{0}_{p \times p} & c_2 \ddot{\psi}(z_2) x x^T & \cdots & \mathbf{0}_{p \times p} & \dot{\psi}(z_2) x e_2^T \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ c_k \ddot{\psi}(z_k) x e_k^T & \mathbf{0}_{p \times p} & \cdots & \mathbf{0}_{p \times p} & c_k \ddot{\psi}(z_k) x x^T & \dot{\psi}(z_k) x e_k^T \\ D_{\dot{\psi}(z)} & \dot{\psi}(z_1) e_1 x^T & \dot{\psi}(z_2) e_2 x^T & \cdots & \dot{\psi}(z_k) e_k x^T & \mathbf{0}_{k \times k} \end{bmatrix}.$$

In above, we use D_v to denote the diagonal matrix with diagonal entries being components of v . We will first show that $\nabla^2 g \preceq Q_x \preceq (2\|c\|_\infty + 2)C\Sigma_x^*$, where

$$\begin{aligned} Q_x &:= C \text{diag}\{(2\|c\|_\infty + 1)I_k, (2\|c\|_\infty + 1)xx^T, \dots, (2\|c\|_\infty + 1)xx^T, 2I_k\}, \\ \Sigma_x^* &:= \text{diag}\{I_k, xx^T, \dots, xx^T, I_k\}. \end{aligned}$$

Recall $W = [u; v_1; \dots; v_k; w] \in \mathbb{R}^{2k+kp}$. Note that

$$\begin{aligned} W^T \nabla^2 g W &= u^T D_{\text{co}\ddot{\psi}(z)} u + \sum_{i=1}^k c_i \ddot{\psi}(z_i) (v_i^T x)^2 + 2w^T D_{\dot{\psi}(z)} u \\ &\quad + 2 \sum_{i=1}^k c_i \ddot{\psi}(z_i) (v_i^T x) (u^T e_i) + 2 \sum_{i=1}^k \dot{\psi}(z_i) (v_i^T x) (w^T e_i). \end{aligned} \quad (68)$$

And further

$$\begin{aligned} u^T D_{\text{co}\ddot{\psi}(z)} u &\leq \|D_{\text{co}\ddot{\psi}(z)}\| \|u\|^2 \leq C \|c\|_\infty \|u\|^2, \\ c_i \ddot{\psi}(z_i) (v_i^T x)^2 &\leq C \|c\|_\infty (v_i^T x)^2, \\ 2w^T D_{\dot{\psi}(z)} u &\leq C \|w\|^2 + C \|u\|^2, \\ 2c_i \ddot{\psi}(z_i) (v_i^T x) (u^T e_i) &\leq \|c\|_\infty C ((v_i^T x)^2 + (u^T e_i)^2), \\ 2\dot{\psi}(z_i) (v_i^T x) (w^T e_i) &\leq C ((v_i^T x)^2 + (w^T e_i)^2). \end{aligned}$$

Replace these upper bounds to terms in (68), we find

$$W^T \nabla^2 g W \leq W^T Q_x W,$$

because $\sum_i (u^T e_i)^2 = \|u\|^2$. Since this holds for all W , we have $\nabla^2 g \preceq Q_x$. Finally, we note that

$$\begin{aligned} |W^T \mathbb{E}(g(w, x) - g(w^*, x) - \xi) \nabla^2 g W| &= |W^T \mathbb{E}[(g(w, x) - g(w^*, x)) \nabla^2 g] W| \\ &\leq \sqrt{\mathbb{E}(g(w, x) - g(w^*, x))^2} \sqrt{\mathbb{E}(W^T \nabla^2 g W)^2}. \end{aligned}$$

Recall (66), we have

$$\sqrt{\mathbb{E}(g(w, x) - g(w^*, x))^2} \leq (\mathbb{E}(g(w, x) - g(w^*, x))^4)^{1/4} \leq \sqrt{6} C \|\Delta w\|_{\Sigma^*} (\|w^*\|_{\Sigma^*} + \|w\|_{\Sigma^*}).$$

Then, by Lemma C.2, we have

$$\mathbb{E}(W^T \nabla^2 g W)^2 \leq 4(\|c\|_\infty + 1)^2 C^2 \mathbb{E}(W^T \Sigma_x^* W)^2 \leq 12C^2 (\|c\|_\infty + 1)^2 (W^T \Sigma^* W)^2.$$

In combination, we find

$$|W^T \mathbb{E}(g(w, x) - g(w^*, x) - \xi) \nabla^2 g W| \leq 6\sqrt{2}C^2(\|c\|_\infty + 1) \|\Delta w\|_{\Sigma^*} (\|w^*\|_{\Sigma^*} + \|w\|_{\Sigma^*}) W^T \Sigma^* W.$$

This verifies our claim 3).

Proof for Claim 4): Simply note that by (66), we get

$$G(w) = \mathbb{E}|g(w, x) - g(w^*, x)|^2 \leq 6C^2 \|\Delta w\|_{\Sigma^*}^2 (\|w^*\|_{\Sigma^*}^2 + \|w\|_{\Sigma^*}^2).$$

□

Proof for Proposition 5.3. First, we find that, when $w \in \mathcal{D}$,

$$\|c\|_\infty^2 \leq \|w\|_{\Sigma^*}^2 \leq (1 + \frac{1}{4})^2 \|w^*\|_{\Sigma^*}^2 \leq 2\|w^*\|_{\Sigma^*}^2. \quad (69)$$

Note that

$$\begin{aligned} \nabla^2 F &= \mathbb{E} \nabla g(w, x) \nabla g(w, x)^T + \mathbb{E}(g(w, x) - g(w^*, x) - \xi) \nabla^2 g(w, x) \\ &= \mathbb{E} \nabla g(w, x) \nabla g(w, x)^T + \mathbb{E}(g(w, x) - g(w^*, x)) \nabla^2 g(w, x). \end{aligned}$$

By Lemma C.3 claim 2) and claim 3) and (69), we have

$$\begin{aligned} \mathbb{E} \nabla g(w, x) \nabla g(w, x)^T &\preceq 6C^2 \|w^*\|_{\Sigma^*}^2 \Sigma^*, \\ \mathbb{E}(g(w, x) - g(w^*, x)) \nabla^2 g(w, x) &\leq 6\sqrt{2}C^2 (\|c\|_\infty + 1) \delta C_1(w^*) \|w^*\|_{\Sigma^*} (\|w^*\|_{\Sigma^*} + \|w\|_{\Sigma^*}) \Sigma^* \\ &\leq 18\sqrt{2}C^2 (2\|w^*\|_{\Sigma^*} + 1) \delta C_1(w^*) \|w^*\|_{\Sigma^*}^2 \Sigma^* \\ &\leq 4\delta C^2 \|w^*\|_{\Sigma^*}^2 \Sigma^*. \end{aligned}$$

So $\nabla^2 F \preceq C_0(w^*) \Sigma^*$. Also note that $\mathbb{E} \nabla g(w, x) \nabla g(w, x)^T \succeq 0$, we have

$$\nabla^2 F \succeq \mathbb{E}(g(w, x) - g(w^*, x)) \nabla^2 g(w, x) \succeq -4\delta C^2 \|w^*\|_{\Sigma^*}^2 \Sigma^* \succeq -\delta A.$$

Then, by Lemma C.3 claim 1) and (69), we find that

$$\begin{aligned} \mathbb{E} \|\nabla f(w) - \nabla F(w)\|^2 &\leq \mathbb{E} \|\nabla f(w)\|^2 \\ &\leq 8\sqrt{3}(1 + \text{tr}(\Sigma))(6C^2 \|\Delta w\|_{\Sigma^*}^2 (\|w^*\|_{\Sigma^*}^2 + \|w\|_{\Sigma^*}^2) + \sigma_0^2) C^2 \|w\|_{\Sigma^*}^2 \\ &\leq 8\sqrt{3}(1 + \text{tr}(\Sigma))(18C^2 \|\Delta w\|_{\Sigma^*}^2 \|w^*\|_{\Sigma^*}^2 + \sigma_0^2) C^2 \|w^*\|_{\Sigma^*}^2 \\ &\leq 8\sqrt{3}(1 + \text{tr}(\Sigma))(18\delta^2 (C_1(w^*))^2 C^2 \|w^*\|_{\Sigma^*}^2 \|w^*\|_{\Sigma^*}^2 + \sigma_0^2) C^2 \|w^*\|_{\Sigma^*}^2 \\ &\leq 8\sqrt{3}(1 + \text{tr}(\Sigma))(C^2 \|w^*\|_{\Sigma^*}^4 + \sigma_0^2) C^2 \|w^*\|_{\Sigma^*}^2. \end{aligned}$$

Finally, by claim 4) of Lemma C.3, when $w_0 \in \mathcal{D}$, we have

$$G(w_0) \leq 18C^2 (C_1(w^*))^2 \delta^2 \|w^*\|_{\Sigma^*}^4 \leq C^2 \|w^*\|_{\Sigma^*}^4.$$

□

References

- Alnur Ali, Zico Kolter, and Ryan Tibshirani. A continuous-time view of early stopping for least squares regression. In *Proceedings of the International Conference on Artificial Intelligence and Statistics*, 2019.
- Zeyuan Allen-Zhu, Yuanzhi Li, and Yingyu Liang. Learning and generalization in overparameterized neural networks, going beyond two layers. In *Proceedings of the Advance of Neural Information Processing Systems*, 2019.
- Sanjeev Arora, Simon S. Du, Wei Hu, Zhiyuan Li, and Ruosong Wang. Fine-grained analysis of optimization and generalization for overparameterized two-layer neural networks. In *Proceedings of the International Conference on Machine Learning*, 2019.
- Francis Bach and Eric Moulines. Non-asymptotic analysis of stochastic approximation algorithms for machine learning. In *Proceedings of the Advances in Neural Information Processing Systems*, 2011.
- Francis Bach and Eric Moulines. Non-strongly-convex smooth stochastic approximation with convergence rate $o(1/n)$. In *Proceedings of the Advances in Neural Information Processing Systems*, 2013.
- Peter L. Bartlett, Philip M. Long, Gabor Lugosi, and Alexander Tsigler. Benign overfitting in linear regression. *arXiv preprint arXiv:1906.11300*, 2019.
- Mikhail Belkin, Daniel Hsu, and Partha Mitra. Overfitting or perfect fitting? Risk bounds for classification and regression rules that interpolate. In *Proceedings of the Advance of Neural Information Processing Systems*, 2018.
- Mikhail Belkin, Daniel Hsu, Siyuan Ma, and Soumik Mandal. Reconciling modern machine-learning practice and the classical bias–variance trade-off. *Proceedings of the National Academy of Sciences of the United States of America*, 116(32):15849–15854, 2019.
- T. Cai and P. Hall. Prediction in function linear regression. *Ann. Statist.*, 34:2159–2179, 2008.
- Luigi Carratino, Alessandro Rudi, and Lorenzo Rosasco. Learning with sgd and random features. In *Advances in Neural Information Processing Systems*, pages 10192–10203, 2018.

- W. E, C. Ma, and Q. Wang. A priori estimates of the population risk for residual networks. *arXiv:1903.02154*, 2019a.
- W. E, C. Ma, and L. Wu. A comparative analysis of the optimization and generalization property of two-layer neural network and random feature models under gradient descent dynamics. *arXiv:1904.04326*, 2019b.
- Y. Fan, M. James, and P. Radchenko. Functional additive regression. *Ann. Statist.*, 43:2296–2325, 2015.
- Saeed Ghadimi and Guanghui Lan. Optimal stochastic approximation algorithms for strongly convex stochastic composite optimization i: A generic algorithmic framework. *SIAM J. Optim.*, 22(4):1469–1492, 2012.
- Noah Golowich, Alexander Rakhlin, and Ohad Shamir. Size-independent sample complexity of neural networks. In *Proceedings of the Annual Conference on Learning Theory*, 2018.
- P. Hall and J.L. Horowitz. Methodology and convergence rates for functional linear regression. *Ann. Statist.*, 35:70–91, 2007.
- T. Hastie, A. Montanari, S. Rosset, and R. Tibshirani. Surprises high-dimensional ridgeless least squares interpolation. *arXiv:1903.08560*, 2019.
- Rie Johnson and Tong Zhang. Accelerating stochastic gradient descent using predictive variance reduction. *Advances in neural information processing systems*, 26:315–323, 2013.
- I.T. Jolliffe. *Principal component analysis, 2nd ed.* Springer, 2002.
- Yuanzhi Li and Yingyu Liang. Learning overparameterized neural networks via stochastic gradient descent on structured data. In *Proceedings of the Advance of Neural Information Processing Systems*, 2018.
- M. Mohri, A. Rostamizadeh, and A. Talwakar. *Foundations of machine learning.* The MIT press, 2018.
- P. Nakkiran. More data can hurt for linear regression: sample-wise double descent. *arxiv:1912.07242*, 2019.
- Preetum Nakkiran, Prayaag Venkat, Sham Kakade, and Tengyu Ma. Optimal regularization can mitigate double descent. *arXiv preprint arXiv:2003.01897*, 2020.

- Yu Nesterov and J-Ph Vial. Confidence level solutions for stochastic programming. *Automatica*, 44(6):1559–1568, 2008.
- Behnam Neyshabur, Zhiyuan Li, Srinadh Bhojanapalli, Yann LeCun, and Nathan Srebro. Towards understanding the role of over-parametrization in generalization of neural networks. In *Proceedings of the International Conference on Learning Representations*, 2019.
- J. O. Ramsay and B. W. Silverman. *Functional data analysis (2nd ed.)*. Springer, 2005.
- Herbert Robbins and Sutton Monro. A stochastic approximation method. *Ann. Math. Statist.*, 22(3):400–407, 1951.
- Nicolas Le Roux, Mark Schmidt, and Francis Bach. A stochastic gradient method with an exponential convergence rate for strongly-convex optimization with finite training sets. In *Proceedings of the Advances in Neural Information Processing Systems*, 2012.
- Shai Shalev-Shwartz and Shai Ben-David. *Understanding machine learning: From theory to algorithms*. Cambridge university press, 2014.
- R. Tsay. *Analysis of financial time series, 3rd edition*. Wiley, 2010.
- J. W. Tukey. A survey of sampling from contaminated distributions. *Contributions to Probability and Statistics*, 2:448–485, 1960.
- K. Xue and F. Yao. Hypothesis testing in large-scale functional linear regression. Accepted by *Statistica Sinica*, SS-2018-0456, 2018.
- Tong Zhang. Solving large scale linear prediction problems using stochastic gradient descent algorithms. In *Proceedings of the International Conference on Machine Learning*, 2004.
- Tong Zhang. Learning bounds for kernel regression using effective data dimensionality. *Neural Computation*, 17(9):2077–2098, 2005.