

L^p -trace-free generalized Korn inequalities for incompatible tensor fields in three space dimensions

Peter Lewintan^{1,*} and Patrizio Neff¹

October 14, 2021

*In memoriam of Sergio Dain [1970-2016],
who gave the first proof of the trace-free Korn's inequality
on bounded Lipschitz domains.*

Abstract

For $1 < p < \infty$ we prove an L^p -version of the generalized trace-free Korn inequality for incompatible tensor fields P in $W_0^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$. More precisely, let $\Omega \subset \mathbb{R}^3$ be a bounded Lipschitz domain. Then there exists a constant $c > 0$ such that

$$\|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c (\|\text{dev sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})})$$

holds for all tensor fields $P \in W_0^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$, i.e., for all $P \in W^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$ with vanishing tangential trace $P \times \nu = 0$ on $\partial\Omega$ where ν denotes the outward unit normal vector field to $\partial\Omega$ and $\text{dev } P := P - \frac{1}{3} \text{tr}(P) \cdot \mathbb{1}$ denotes the deviatoric (trace-free) part of P . We also show the norm equivalence

$$\|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{Curl } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c (\|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})})$$

for tensor fields $P \in W^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$. These estimates also hold true for tensor fields with vanishing tangential trace only on a relatively open (non-empty) subset $\Gamma \subseteq \partial\Omega$ of the boundary.

AMS 2010 subject classification: Primary: 35A23; Secondary: 35B45, 35Q74, 46E35.

Keywords: $W^{1,p}(\text{Curl})$ -Korn's inequality, Poincaré's inequality, Lions lemma, Nečas estimate, incompatibility, Curl-spaces, Maxwell problems, gradient plasticity, dislocation density, relaxed micromorphic model, Cosserat elasticity, Kröner's incompatibility tensor, Saint-Venant compatibility, trace-free Korn's inequality, conformal mappings, conformal Killing vector field, Nye's formula.

1 Introduction

Korn-type inequalities are crucial for a priori estimates in linear elasticity and fluid mechanics. They allow to bound the L^p -norm of the gradient Du in terms of the symmetric gradient, i.e. Korn's first inequality states

$$\exists c > 0 \forall u \in W_0^{1,p}(\Omega, \mathbb{R}^n) : \quad \|Du\|_{L^p(\Omega, \mathbb{R}^{n \times n})} \leq c \|\text{sym } Du\|_{L^p(\Omega, \mathbb{R}^{n \times n})}. \quad (1.1)$$

Generalizations to many different settings have been obtained in the literature, including the geometrically nonlinear counterpart [24, 39, 23], mixed growth conditions [15], incompatible fields (also with dislocations)

¹Chair for Nonlinear Analysis and Modeling, Faculty of Mathematics, University of Duisburg-Essen, Thea-Leymann-Str. 9, 45127 Essen, Germany

*Corresponding author, email: peter.lewintan@uni-due.de

[48, 58, 57, 55, 56, 6, 43, 42, 41, 40], as well as the case of non-constant coefficients [50, 37, 59, 62] and on Riemannian manifolds [9]. In this paper we focus on their improvement towards the trace-free case:

$$\exists c > 0 \forall u \in W_0^{1,p}(\Omega, \mathbb{R}^n) : \quad \|Du\|_{L^p(\Omega, \mathbb{R}^{n \times n})} \leq c \|\text{dev}_n \text{sym} Du\|_{L^p(\Omega, \mathbb{R}^{n \times n})}, \quad (1.2)$$

where $\text{dev}_n X := X - \frac{1}{n} \text{tr}(X) \cdot \mathbb{1}$ denotes the deviatoric (trace-free) part of the square matrix X . Note in passing that (1.2) implies (1.1).

There exist many different proofs and generalizations of the trace-free classical Korn's inequality in the literature, see [63, Theorem 2] but also [64, 17, 27, 65, 33, 6] as well as [67] for trace-free Korn's inequalities in pseudo-Euclidean space and [17, 32] for trace-free Korn inequalities on manifolds, [8, 25] for trace-free Korn inequalities in Orlicz spaces and [45, 18] for weighted trace-free Korn inequalities in Hölder and John domains. Such coercive inequalities found application in micro-polar Cosserat-type models [34, 49, 33, 27] and general relativity [17]. On the other hand, corresponding trace-free coercive inequalities for incompatible tensor fields are useful in infinitesimal gradient plasticity as well as in linear relaxed micromorphic elasticity, see [51, 31] but also [6, sec. 7] and the references contained therein.

Notably, in case $n = 2$, the condition $\text{dev}_2 \text{sym} Du \equiv 0$ becomes the system of Cauchy-Riemann equations, so that the corresponding kernel is infinite-dimensional and an adequate quantitative version of the trace-free classical Korn's inequality does not hold true. Nevertheless, in [27] it is proved that

$$\|Du\|_{L^p(\Omega, \mathbb{R}^{2 \times 2})} \leq c \|\text{dev}_2 \text{sym} Du\|_{L^p(\Omega, \mathbb{R}^{2 \times 2})} \quad (1.3)$$

holds for each $u \in W_0^{1,p}(\Omega, \mathbb{R}^2)$,¹ but, again, this result ceases to be valid if the Dirichlet conditions are prescribed only on a part of the boundary, cf. the counterexample in [6, sec. 6.6].

Korn-type inequalities fail for the limiting cases $p = 1$ and $p = \infty$. Indeed, from the counterexamples traced back in [16, 38, 61, 47] it follows that $\int_{\Omega} |\text{sym} Du| dx$ does not dominate each quantity $\int_{\Omega} |\partial_i u_j| dx$ for any vector field $u \in W_0^{1,1}(\Omega, \mathbb{R}^n)$. Hence, also trace-free versions fail for $p = 1$ and $p = \infty$. On the other hand, Poincaré-type inequalities estimating certain integral norms of the deformation u in terms of the total variation of the symmetric strain tensor $\text{sym} Du$ are still valid. In particular, for Poincaré-type inequalities for functions of bounded deformation involving the deviatoric part of the symmetric gradient we refer to [26].

The classical Korn's inequalities need compatibility, i.e. a gradient Du ; giving up the compatibility necessitates controlling the distance of P to a gradient by adding the incompatibility measure (the dislocation density tensor) $\text{Curl} P$. We showed in [43] the following quantitative version of Korn's inequality for incompatible tensor fields $P \in W^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$:

$$\inf_{\tilde{A} \in \mathfrak{so}(3)} \|P - \tilde{A}\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \left(\|\text{sym} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{Curl} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \right). \quad (1.4)$$

Note that the constant skew-symmetric matrix fields (restricted to Ω) represent the elements from the kernel of the right-hand side of (1.4). For compatible $P = Du$ recover from (1.4) the quantitative version of the classical Korn's inequality, namely for $u \in W^{1,p}(\Omega, \mathbb{R}^3)$:

$$\inf_{\tilde{A} \in \mathfrak{so}(3)} \|Du - \tilde{A}\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \|\text{sym} Du\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \quad (1.5)$$

and for skew-symmetric matrix fields $P = A \in \mathfrak{so}(3)$ the corresponding Poincaré inequality for squared skew-symmetric matrix fields $A \in W^{1,p}(\Omega, \mathfrak{so}(3))$ (and thus for vectors in \mathbb{R}^3):

$$\inf_{\tilde{A} \in \mathfrak{so}(3)} \|A - \tilde{A}\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \|\text{Curl} A\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq \tilde{c} \|DA\|_{L^p(\Omega, \mathbb{R}^{3 \times 3^2})}, \quad (1.6)$$

¹A simple proof using partial integration is given in the Appendix for the case $p = 2$ and all dimensions.

where in the last step we have used that Curl consists of linear combinations from D. Interestingly, for skew-symmetric A also the converse is true, more precisely, the entries of DA are linear combinations of the entries from $\text{Curl } A$, cf. e.g. [43, Cor. 2.3]:

$$DA = L(\text{Curl } A) \quad \text{for skew-symmetric } A, \quad (1.7)$$

where $L(\cdot)$ denotes a corresponding linear operator with constant coefficients, not necessarily the same in any two places in the present paper. In fact, the mentioned results also hold in higher dimensions $n > 3$, see [42] and the discussion contained therein. In our proof of (1.4) we were highly inspired by a proof of (1.5) advocated by P. G. Ciarlet and his collaborators [11, 12, 13, 14, 29, 19, 10], which uses the Lions lemma resp. Nečas estimate, the compact embedding $W^{1,p} \subset\subset L^p$ and the representation of the second distributional derivatives of the displacement u by a linear combination of the first derivatives of the symmetrized gradient Du :

$$D^2u = L(D \text{sym } Du). \quad (1.8)$$

It is worth mentioning that the role of the latter ingredient (1.8) was taken over by (1.7) in our proof of (1.4) in [43] resp. [42]. In $n = 3$ dimensions the relation (1.7) is an easy consequence of the so called *Nye's formula* [60, eq. (7)]:

$$\text{Curl } A = \text{tr}(D \text{axl } A) \cdot \mathbf{1} - (D \text{axl } A)^T, \quad (1.9a)$$

resp.

$$D \text{axl } A = \frac{1}{2} \text{tr}(\text{Curl } A) \cdot \mathbf{1} - (\text{Curl } A)^T, \quad (1.9b)$$

where we identify the vectorspace of skew-symmetric matrices $\mathfrak{so}(3)$ and \mathbb{R}^3 via $\text{axl} : \mathfrak{so}(3) \rightarrow \mathbb{R}^3$ which is defined by the cross product:

$$A b =: \text{axl}(A) \times b \quad \forall b \in \mathbb{R}^3, \quad (1.10)$$

and associates with a skew-symmetric matrix $A \in \mathfrak{so}(3)$ the vector $\text{axl } A := (-A_{23}, A_{13}, -A_{12})^T$. The relation (1.9a) admits moreover a counterpart on the group of orthogonal matrices $O(3)$ and even in higher spatial dimensions, see [54]. In fact, Nye's formula is (formally) a consequence of the following algebraic identity:

$$(\text{Anti } a) \times b = b \otimes a - \langle b, a \rangle \cdot \mathbf{1} \quad \forall a, b \in \mathbb{R}^3, \quad (1.11)$$

where the vector product of a matrix and a vector is to be seen row-wise and $\text{Anti} : \mathbb{R}^3 \rightarrow \mathfrak{so}(3)$ is the inverse of axl . Despite the absence of the simple algebraic relations in the higher dimensional case a corresponding relation to (1.7) also holds true in $n > 3$, see e.g. [42].

Moreover, the kernel in quantitative versions of Korn's inequalities is killed by corresponding boundary conditions, namely by a vanishing trace condition $u|_{\partial\Omega} = 0$ in the case of (1.5) and (1.6) and by a vanishing tangential trace condition $P \times \nu|_{\partial\Omega} = 0$ in the general case (1.4), cf. [43, 42].

The objective of the present paper is to improve on inequality (1.4) by showing that it already suffices to consider the deviatoric (trace-free) parts on the right-hand side, hence, further contributing to the problems proposed in [58]. More precisely, the main results are

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^3$ be a bounded Lipschitz domain and $1 < p < \infty$. There exists a constant $c = c(p, \Omega) > 0$ such that for all $P \in L^p(\Omega, \mathbb{R}^{3 \times 3})$ we have*

$$\inf_{T \in K_{dS, dC}} \|P - T\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \left(\|\text{dev sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} \right), \quad (1.12)$$

where $\text{dev } X := X - \frac{1}{3} \text{tr}(X) \cdot \mathbf{1}$ denotes the deviatoric part of a square tensor $X \in \mathbb{R}^{3 \times 3}$ and $K_{dS, dC}$ represent the kernel of the right-hand side and is given by

$$K_{dS, dC} = \{T : \Omega \rightarrow \mathbb{R}^{3 \times 3} \mid T(x) = \text{Anti}(\tilde{A}x + \beta x + b) + (\langle \text{axl } \tilde{A}, x \rangle + \gamma) \cdot \mathbf{1}, \\ \tilde{A} \in \mathfrak{so}(3), b \in \mathbb{R}^3, \beta, \gamma \in \mathbb{R}\}. \quad (1.13)$$

By killing the kernel with tangential trace conditions (note that $\text{dev}(P \times \nu) = 0$ iff $P \times \nu = 0$) we arrive at the following Korn's first type inequality

Theorem 1.2. *Let $\Omega \subset \mathbb{R}^3$ be a bounded Lipschitz domain and $1 < p < \infty$. There exists a constant $c = c(p, \Omega) > 0$ such that we have*

$$\|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c (\|\text{dev sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}) \quad (1.14)$$

for all

$$P \in W_0^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3}) := \{P \in L^p(\Omega, \mathbb{R}^{3 \times 3}) \mid \text{Curl } P \in L^p(\Omega, \mathbb{R}^{3 \times 3}), P \times \nu \equiv 0 \text{ on } \partial\Omega\}.$$

The appearance of the term $\text{dev Curl } P$ on the right hand side of (1.14) would suggest to consider p -integrable tensor fields P with ‘only’ p -integrable $\text{dev Curl } P$. However, this would not lead to a new Banach space, since we show that for all $m \in \mathbb{Z}$ it holds that

$$\text{Curl } P \in W^{m,p}(\Omega, \mathbb{R}^{3 \times 3}) \Leftrightarrow \text{dev Curl } P \in W^{m,p}(\Omega, \mathbb{R}^{3 \times 3}). \quad (1.15)$$

The estimate (1.14) generalizes the corresponding result in [6] from the L^2 -setting to the L^p -setting, whereas the trace-free second type inequality (1.12) is completely new. Generalizations to different right hand sides and higher dimensions have been obtained in the recent papers [40, 41]. Note however that the estimates (1.12) and (1.14) are restricted to the case of three dimensions since the deviatoric operator acts on square matrices and only in the three-dimensional setting the matrix Curl returns a square matrix.

Again, for compatible $P = Du$ we get back a tangential trace-free classical Korn inequality for the displacement gradient, namely

$$\|Du\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \|\text{dev sym } Du\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \quad \text{with } Du \times \nu = 0 \text{ on } \partial\Omega \quad (1.16)$$

as well as

$$\inf_{T \in K_{dS,C}} \|Du - T\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \|\text{dev sym } Du\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \quad (1.17)$$

respectively

$$\|u - \Pi u\|_{W^{1,p}(\Omega, \mathbb{R}^3)} \leq c \|\text{dev sym } Du\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}, \quad (1.18)$$

where Π denotes an arbitrary projection operator from $W^{1,p}(\Omega, \mathbb{R}^3)$ onto the space of *conformal Killing vectors*, here the finite dimensional kernel of $\text{dev sym } D$, which is given by quadratic polynomials of the form

$$\varphi_c(x) = \langle a, x \rangle x - \frac{1}{2} a \|x\|^2 + \text{Anti}(b)x + \beta x + c, \quad \text{with } a := axl \tilde{A}, b, c \in \mathbb{R}^3 \text{ and } \beta \in \mathbb{R},$$

namely the *infinitesimal conformal mappings*, cf. [63, 49, 33, 17, 64, 65], see Figure 1 for an illustration in 2D.

A first proof of (1.18), even in all dimensions $n \geq 3$, was given by Reshetnyak [63] over domains which are star-like with respect to a ball. Over bounded Lipschitz domains the trace-free Korn's second inequality in all dimensions $n \geq 3$, namely

$$\exists c > 0 \forall u \in W^{1,p}(\Omega, \mathbb{R}^n) : \|u\|_{W^{1,p}(\Omega, \mathbb{R}^n)} \leq c (\|u\|_{L^p(\Omega, \mathbb{R}^n)} + \|\text{dev}_n \text{sym } Du\|_{L^p(\Omega, \mathbb{R}^{n \times n})}), \quad (1.19)$$

was justified by Dain [17] in the case $p = 2$ and by Schirra [65] for all $p > 1$. Their proofs use again the Lions lemma and the ‘higher order’ analogues of the differential relation (1.8):

$$D\Delta u = L(D^2 \text{dev}_n \text{sym } Du). \quad (1.20)$$

However, the differential operators $\text{sym } D$ and $\text{dev}_n \text{sym } D$ are particular cases of the so-called *coercive elliptic operators* whose study began with Aronszajn [5].




Figure 1: In the planar case, the condition $\text{dev}_2 \text{sym } Du = 0$ coincides with the Cauchy-Riemann equations for the function u (see Appendix). Therefore, infinitesimal conformal mappings in 2D are holomorphic functions which preserve angles exactly. This ceases to be the case for 3D infinitesimal conformal mappings defined by $\text{dev}_3 \text{sym } Du = 0$.

Let us go back to

$$\|P\|_{L^2(\Omega, \mathbb{R}^{3 \times 3})} \leq c (\|\text{dev } \text{sym } P\|_{L^2(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev } \text{Curl } P\|_{L^2(\Omega, \mathbb{R}^{3 \times 3})}) \quad (1.21)$$

whose first proof for $P \in W_0^{1,2}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$ was given in [6] via the trace-free classical Korn's inequality, a Maxwell estimate and a Helmholtz decomposition and is not directly amenable to the L^p -case. Here, we catch up with the latter.

In the following section we start by summarizing the notations and collect some preliminary results from algebraic calculations which are needed in the subsequent vector calculus to establish relations of the type:

$$D^3(A + \zeta \cdot \mathbf{1}) = L(D^2 \text{dev } \text{Curl}(A + \zeta \cdot \mathbf{1})) \quad (1.22)$$

for skew-symmetric tensor fields A and scalar functions ζ , where L denotes a corresponding constant coefficients linear operator. Based on this “higher order” analogue of the differential relation (1.7) we prove our main results in the last section using a similar argumentation as in [17, 65] which argue by the Lions lemma resp. Nečas estimate and the compact embedding $W^{1,p}(\Omega) \subset\subset L^p(\Omega)$.

2 Notations and preliminaries

Let $n \geq 2$. We consider for vectors $a, b \in \mathbb{R}^n$ the scalar product $\langle a, b \rangle := \sum_{i=1}^n a_i b_i \in \mathbb{R}$, the (squared) norm $\|a\|^2 := \langle a, a \rangle$ and the dyadic product $a \otimes b := (a_i b_j)_{i,j=1,\dots,n} \in \mathbb{R}^{n \times n}$. Similarly, we define the scalar product for matrices $P, Q \in \mathbb{R}^{n \times n}$ by $\langle P, Q \rangle := \sum_{i,j=1}^n P_{ij} Q_{ij} \in \mathbb{R}$ and the (squared) Frobenius-norm by $\|P\|^2 := \langle P, P \rangle$. We highlight by \cdot the scalar multiplication of a scalar with a matrix, whereas matrix multiplication is denoted only by juxtaposition.

Moreover, $P^T := (P_{ji})_{i,j=1,\dots,n}$ denotes the transposition of the matrix $P = (P_{ij})_{i,j=1,\dots,n}$. The latter decomposes orthogonally into the symmetric part $\text{sym } P := \frac{1}{2}(P + P^T)$ and the skew-symmetric part $\text{skew } P := \frac{1}{2}(P - P^T)$. We will denote by $\mathfrak{so}(n) := \{A \in \mathbb{R}^{n \times n} \mid A^T = -A\}$ the Lie-Algebra of skew-symmetric matrices.

For the identity matrix we will write $\mathbf{1}$, so that the trace of a squared matrix P is given by $\text{tr } P := \langle P, \mathbf{1} \rangle$. The deviatoric (trace-free) part of P is given by $\text{dev}_n P := P - \frac{1}{n} \text{tr}(P) \cdot \mathbf{1}$ and in three dimensions its index will be suppressed, i.e. we write dev instead of dev_3 .

We will denote by $\mathcal{D}'(\Omega)$ the space of distributions on a bounded Lipschitz domain $\Omega \subset \mathbb{R}^n$ and by $W^{-k,p}(\Omega)$ the dual space of $W_0^{k,p'}(\Omega)$, where $p' = \frac{p}{p-1}$ is the Hölder dual exponent to p .

Throughout the paper we use c as a generic positive constant, which is not necessarily the same in any two places, and we use $L(\cdot)$ as a generic linear operator with constant coefficients, which also may differ in any two places within the paper.

In 3-dimensions we make use of the vector product $\times : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$. Since the vector product $a \times \cdot$ with a fixed vector $a \in \mathbb{R}^3$ is linear in the second component, there exists a unique matrix $\text{Anti}(a)$ such that

$$a \times b =: \text{Anti}(a)b \quad \forall b \in \mathbb{R}^3, \quad (2.1)$$

and direct calculations show that for $a = (a_1, a_2, a_3)^T$ the matrix $\text{Anti}(a)$ has the form

$$\text{Anti}(a) = \begin{pmatrix} 0 & -a_3 & a_2 \\ a_3 & 0 & -a_1 \\ -a_2 & a_1 & 0 \end{pmatrix}. \quad (2.2)$$

The inverse of $\text{Anti} : \mathbb{R}^3 \rightarrow \mathfrak{so}(3)$ is denoted by $\text{axl} : \mathfrak{so}(3) \rightarrow \mathbb{R}^3$ and fulfills $\text{axl}(A) \times b = Ab$ for all skew-symmetric (3×3) -matrices A and vectors $b \in \mathbb{R}^3$. The matrix representation of the cross product allows for a generalization towards a cross product of a matrix $P \in \mathbb{R}^{3 \times 3}$ and a vector $b \in \mathbb{R}^3$ via

$$P \times b := P \text{Anti}(b), \quad (2.3)$$

so, especially, for $P = \mathbb{1}$ it holds

$$\mathbb{1} \times b = \mathbb{1} \text{Anti}(b) = \text{Anti}(b) \quad \forall b \in \mathbb{R}^3. \quad (2.4)$$

We repeat the following crucial algebraic identity:

$$(\text{Anti } a) \times b = b \otimes a - \langle b, a \rangle \cdot \mathbb{1} \quad \forall a, b \in \mathbb{R}^3. \quad (2.5)$$

Observation 2.1. For $P \in \mathbb{R}^{3 \times 3}$ and $b \in \mathbb{R}^3$ we have

$$\text{dev}(P \times b) = 0 \Leftrightarrow P \times b = 0. \quad (2.6)$$

Proof. We decompose P into its symmetric and skew-symmetric part, i.e.,

$$P = S + A = S + \text{Anti}(a), \quad \text{for some } S \in \text{Sym}(3), A \in \mathfrak{so}(3) \text{ and with } a = \text{axl}(A).$$

For a symmetric matrix S it holds $\text{tr}(S \times b) = 0$ for any $b \in \mathbb{R}^3$, since²

$$\text{tr}(S \times b) = \langle S \times b, \mathbb{1} \rangle_{\mathbb{R}^{3 \times 3}} = \langle S \text{Anti}(b), \mathbb{1} \rangle_{\mathbb{R}^{3 \times 3}} = -\langle S, \text{Anti}(b) \rangle_{\mathbb{R}^{3 \times 3}} \stackrel{S \in \text{Sym}(3)}{=} 0. \quad (2.7)$$

Thus, using the decomposition $P = S + \text{Anti}(a)$, we have:

$$\begin{aligned} \text{dev}(P \times b) &= P \times b - \frac{1}{3} \text{tr}(P \times b) \cdot \mathbb{1} \stackrel{(2.7)}{=} P \times b - \frac{1}{3} \text{tr}((\text{Anti } a) \times b) \cdot \mathbb{1} \\ &\stackrel{(2.5)}{=} P \times b - \frac{1}{3} \text{tr}(b \otimes a - \langle b, a \rangle \cdot \mathbb{1}) \cdot \mathbb{1} = P \times b + \frac{2}{3} \langle a, b \rangle \cdot \mathbb{1}. \end{aligned} \quad (2.8)$$

Moreover, for any matrix $P \in \mathbb{R}^{3 \times 3}$ we note that

$$(P \times b) b = (P \text{Anti}(b)) b = P(\text{Anti}(b) b) = P(b \times b) = 0. \quad (2.9)$$

²Cf. the Appendix for component-wise calculations.

Thus, we obtain

$$\langle b, \operatorname{dev}(P \times b) b \rangle \stackrel{(2.8)}{=} \langle b, (P \times b + \frac{2}{3} \langle a, b \rangle \cdot \mathbf{1}) b \rangle \stackrel{(2.9)}{=} \frac{2}{3} \langle a, b \rangle \|b\|^2, \quad (2.10)$$

and the conclusion follows from the identity

$$\|b\|^2 \cdot P \times b \stackrel{(2.8)}{=} \|b\|^2 \cdot \operatorname{dev}(P \times b) - \frac{2}{3} \|b\|^2 \langle a, b \rangle \cdot \mathbf{1} \stackrel{(2.10)}{=} \|b\|^2 \cdot \operatorname{dev}(P \times b) - \langle b, \operatorname{dev}(P \times b) b \rangle \cdot \mathbf{1}. \quad (2.11)$$

An application of the Cauchy-Bunyakovsky-Schwarz inequality on the right hand side of (2.11) shows that

$$\|\operatorname{dev}(P \times b)\| \leq \|P \times b\| \leq (1 + \sqrt{3}) \|\operatorname{dev}(P \times b)\|. \quad (2.12)$$

■

Observation 2.2. *Let $a \in \mathbb{R}^3$ and $\alpha \in \mathbb{R}$, then*

$$(\operatorname{Anti}(a) + \alpha \cdot \mathbf{1}) \times b = 0 \quad \text{for } b \in \mathbb{R}^3 \setminus \{0\} \quad \Rightarrow \quad a = 0 \text{ and } \alpha = 0.$$

Proof. By (2.5) and (2.4) we have:

$$0 = (\operatorname{Anti}(a) + \alpha \cdot \mathbf{1}) \times b = b \otimes a - \langle b, a \rangle \cdot \mathbf{1} + \alpha \cdot \operatorname{Anti}(b). \quad (2.13)$$

Taking the trace on both sides we obtain

$$0 = \operatorname{tr}(b \otimes a - \langle b, a \rangle \cdot \mathbf{1} + \alpha \cdot \operatorname{Anti}(b)) = \langle a, b \rangle - 3 \langle a, b \rangle = -2 \langle a, b \rangle.$$

Thus, reinserting $\langle b, a \rangle = 0$ in (2.13) and applying sym on both sides, this implies $\operatorname{sym}(b \otimes a) = 0$. Since

$$\|\operatorname{sym}(a \otimes b)\|^2 = \frac{1}{2} \|a\|^2 \|b\|^2 + \frac{1}{2} \langle a, b \rangle^2 \quad (2.14)$$

and $b \neq 0$ we must have $a = 0$. Hence, by (2.13) also $\alpha = 0$. ■

Formally the gradient and the curl of a vector field $a : \Omega \rightarrow \mathbb{R}^3$ can be seen as

$$\operatorname{Daxl} a = a \otimes \nabla \quad \text{and} \quad \operatorname{curl} a = a \times (-\nabla).$$

The latter also generalizes to (3×3) -matrix fields $P : \Omega \rightarrow \mathbb{R}^{3 \times 3}$ row-wise:³

$$\operatorname{Curl} P = P \times (-\nabla) = \begin{pmatrix} (P^T e_1)^T \\ (P^T e_2)^T \\ (P^T e_3)^T \end{pmatrix} \times (-\nabla) = \begin{pmatrix} (\operatorname{curl}(P^T e_1))^T \\ (\operatorname{curl}(P^T e_2))^T \\ (\operatorname{curl}(P^T e_3))^T \end{pmatrix} \in \mathbb{R}^{3 \times 3}. \quad (2.15)$$

Replacing b by ∇ in (2.5) we obtain Nye's formulas

$$\operatorname{Curl} A = \operatorname{tr}(\operatorname{Daxl} A) \cdot \mathbf{1} - (\operatorname{Daxl} A)^T, \quad (2.16a)$$

and

$$\operatorname{Daxl} A = \frac{1}{2} \operatorname{tr}(\operatorname{Curl} A) \cdot \mathbf{1} - (\operatorname{Curl} A)^T \quad (2.16b)$$

for all skew-symmetric (3×3) -matrix fields A .

³In the literature, the matrix Curl operator is sometimes defined as our transposed $(\operatorname{Curl} P)^T$, cf. Ciarlet [12, Problem 6.18-4].

Remark 2.3. Formal calculations (e.g. replacing b by ∇) have to be performed very carefully. Indeed, they are allowed in algebraic identities but fail, in general, for implications, e.g. for $A \in \mathfrak{so}(3)$ and $b \in \mathbb{R}^3$ we have $A \times b = 0$ if and only if $\operatorname{dev}(A \times b) = 0$, since the following expression holds true, cf. Observation 2.1 and (2.11):

$$\|b\|^2 \cdot A \times b = \|b\|^2 \cdot \operatorname{dev}(A \times b) - \langle b, \operatorname{dev}(A \times b) \rangle \cdot \mathbb{1}. \quad (2.17)$$

However, $\operatorname{dev}(\operatorname{Curl} A) = \operatorname{dev}(A \times (-\nabla)) = 0$ does not imply already that $\operatorname{Curl} A = A \times (-\nabla) = 0$, due to the counterexample $A = \operatorname{Anti}(x)$, since by Nye's formula (2.16) we have $\operatorname{Curl}(\operatorname{Anti}(x)) = 2 \cdot \mathbb{1}$. Of course, we can interpret (2.17) also in the sense of vector calculus, which gives then an expression for $\Delta \operatorname{Curl} A$ in terms of the second distributional derivatives of $\operatorname{dev}(\operatorname{Curl} A)$, but, the latter would have no meaning for the relation of $\operatorname{Curl} A$ and $\operatorname{dev} \operatorname{Curl} A$.

Lemma 2.4. *Let $A \in \mathcal{D}'(\Omega, \mathfrak{so}(3))$ and $\zeta \in \mathcal{D}'(\Omega, \mathbb{R})$. Then*

- (a) *the entries of $D^2(A + \zeta \cdot \mathbb{1})$ are linear combinations of the entries of $D \operatorname{Curl}(A + \zeta \cdot \mathbb{1})$.*
- (b) *the entries of $D^2 A$ are linear combinations of the entries of $D \operatorname{dev} \operatorname{Curl} A$.*
- (c) *the entries of $D^3(A + \zeta \cdot \mathbb{1})$ are linear combinations of the entries of $D^2 \operatorname{dev} \operatorname{Curl}(A + \zeta \cdot \mathbb{1})$.*

Proof. Observe that applying (2.4) to the vector field $\nabla \zeta$ we obtain:

$$\operatorname{Curl}(\zeta \cdot \mathbb{1}) \stackrel{(2.15)}{=} \mathbb{1} \times (-\nabla \zeta) \stackrel{(2.4)}{=} -\operatorname{Anti}(\nabla \zeta). \quad (2.18)$$

Let us first start by proving part (b). From Nye's formula (2.16a) we obtain

$$\operatorname{dev} \operatorname{Curl} A = \frac{1}{3} \operatorname{tr}(\operatorname{Daxl} A) \cdot \mathbb{1} - (\operatorname{Daxl} A)^T \quad (2.19)$$

so that taking the Curl of the transpositions on both sides gives

$$\operatorname{Curl}([\operatorname{dev} \operatorname{Curl} A]^T) \stackrel{\operatorname{Curl} \circ D \equiv 0}{\stackrel{(2.19)}{=}} \frac{1}{3} \operatorname{Curl}(\operatorname{tr}(\operatorname{Daxl} A) \cdot \mathbb{1}) \stackrel{(2.18)}{=} -\frac{1}{3} \operatorname{Anti}(\nabla \operatorname{tr}(\operatorname{Daxl} A)). \quad (2.20)$$

In other words, we have that $\operatorname{Curl}([\operatorname{dev} \operatorname{Curl} A]^T) \in \mathfrak{so}(3)$, and applying axl on both sides of (2.20) we obtain

$$\nabla \operatorname{tr}(\operatorname{Daxl} A) = -3 \operatorname{axl}(\operatorname{Curl}([\operatorname{dev} \operatorname{Curl} A]^T)) = L_0(D \operatorname{dev} \operatorname{Curl} A). \quad (2.21)$$

Taking the ∂_j -derivative of (2.19) for $j = 1, 2, 3$ we conclude

$$\partial_j (\operatorname{Daxl} A)^T \stackrel{(2.19)}{=} \frac{1}{3} \partial_j \operatorname{tr}(\operatorname{Daxl} A) - \partial_j \operatorname{dev} \operatorname{Curl} A \stackrel{(2.21)}{=} \tilde{L}_0(D \operatorname{dev} \operatorname{Curl} A), \quad (2.22)$$

which establishes part (b), namely $D^2 A = L_2(D(\operatorname{dev} \operatorname{Curl} A))$ for skew-symmetric tensor fields A .

The proof of part (a) is divided into the following two key observations:

- (a.i) $D^2 \zeta = \tilde{L}_1(D \operatorname{Curl}(A + \zeta \cdot \mathbb{1}))$,
- (a.ii) $D^2 A = \tilde{L}_2(D \operatorname{Curl}(A + \zeta \cdot \mathbb{1}))$.

To show that each entry of the Hessian matrix $D^2 \zeta$ is a linear combination of the entries of $D \operatorname{Curl}(A + \zeta \cdot \mathbb{1})$ we make use of the second-order differential operator **inc** given for $B \in \mathcal{D}'(\Omega, \mathbb{R}^{3 \times 3})$ via⁴

$$\mathbf{inc} B := \operatorname{Curl}([\operatorname{Curl} B]^T) \quad (2.23)$$

⁴See Kröner [35, §8] for a component-wise expression of the incompatibility operator **inc**.

so that

$$\begin{aligned} \mathbf{inc}(\zeta \cdot \mathbb{1}) &= \text{Curl}([\text{Curl}(\zeta \cdot \mathbb{1})]^T) \stackrel{(2.18)}{=} \text{Curl}(-[\text{Anti}(\nabla\zeta)]^T) = \text{Curl}(\text{Anti}(\nabla\zeta)) \\ &\stackrel{(2.16a)}{=} \text{tr}(\text{D}\nabla\zeta) \cdot \mathbb{1} - (\text{D}\nabla\zeta)^T = \Delta\zeta \cdot \mathbb{1} - \text{D}^2\zeta \in \text{Sym}(3) \end{aligned} \quad (2.24)$$

is symmetric. On the other hand, for a skew-symmetric matrix field $A \in \mathcal{D}'(\Omega, \mathfrak{so}(3))$ we have that

$$\begin{aligned} \mathbf{inc} A &= \text{Curl}([\text{Curl} A]^T) \stackrel{(2.16)}{=} \text{Curl}(\text{tr}(\text{Daxl} A) \cdot \mathbb{1} - \text{Daxl} A) \\ &\stackrel{\text{Curl} \circ \text{D} \equiv 0}{=} \text{Curl}(\text{tr}(\text{Daxl} A) \cdot \mathbb{1}) \stackrel{(2.18)}{=} -\text{Anti}(\nabla \text{tr}(\text{Daxl} A)) \in \mathfrak{so}(3) \end{aligned} \quad (2.25)$$

is skew-symmetric. Hence,

$$\text{sym}(\mathbf{inc}(A + \zeta \cdot \mathbb{1})) = \Delta\zeta \cdot \mathbb{1} - \text{D}^2\zeta \quad \text{and} \quad \text{tr}(\mathbf{inc}(A + \zeta \cdot \mathbb{1})) = 2\Delta\zeta. \quad (2.26)$$

In other words, the entries of the Hessian matrix of ζ are linear combinations of entries from $\mathbf{inc}(A + \zeta \cdot \mathbb{1})$:

$$\begin{aligned} \text{D}^2\zeta &= \Delta\zeta \cdot \mathbb{1} - \text{sym}(\mathbf{inc}(A + \zeta \cdot \mathbb{1})) = \frac{1}{2} \text{tr}(\mathbf{inc}(A + \zeta \cdot \mathbb{1})) \cdot \mathbb{1} - \text{sym}(\mathbf{inc}(A + \zeta \cdot \mathbb{1})) \\ &= \tilde{L}_1(\text{DCurl}(A + \zeta \cdot \mathbb{1})), \end{aligned} \quad (2.27)$$

where we have used that the entries of $\mathbf{inc} B$ are, of course, linear combinations of entries of $\text{DCurl} B$.

To establish (a.ii) from (a.i), recall that for a skew-symmetric matrix field A the entries of $\text{D}A$ are linear combinations of the entries from $\text{Curl} A$:

$$\text{D}A \stackrel{(1.7)}{=} L(\text{Curl} A) = L(\text{Curl}(A + \zeta \cdot \mathbb{1})) - L(\text{Curl}(\zeta \cdot \mathbb{1})) \stackrel{(2.18)}{=} L(\text{Curl}(A + \zeta \cdot \mathbb{1})) + L(\text{Anti}(\nabla\zeta)). \quad (2.28)$$

We conclude by taking the ∂_j -derivative of (2.28) for $j = 1, 2, 3$, namely

$$\partial_j \text{D}A = L(\partial_j \text{Curl}(A + \zeta \cdot \mathbb{1})) + L(\partial_j \text{Anti}(\nabla\zeta)) \stackrel{(a.i)}{=} \tilde{L}_3(\text{DCurl}(A + \zeta \cdot \mathbb{1})).$$

Finally, we establish part (c) arguing in a similar way by showing the following linear combinations:

$$(c.i) \quad \text{D}^2\zeta = \tilde{L}_4(\text{Ddev Curl}(A + \zeta \cdot \mathbb{1})), \quad (c.ii) \quad \text{D}^3A = \tilde{L}_7(\text{D}^2 \text{dev Curl}(A + \zeta \cdot \mathbb{1})).$$

Regarding (2.18) and (2.16) we have

$$\begin{aligned} \text{dev Curl}(A + \zeta \cdot \mathbb{1}) &\stackrel{(2.18)}{=} \text{dev}[\text{Curl} A - \text{Anti}(\nabla\zeta)] = \text{dev Curl} A - \text{Anti}(\nabla\zeta) \\ &\stackrel{(2.16)}{=} \frac{1}{3} \text{tr}(\text{Daxl} A) \cdot \mathbb{1} - (\text{Daxl} A)^T - \text{Anti}(\nabla\zeta). \end{aligned} \quad (2.29)$$

Transposing and taking the Curl on both sides yields

$$\text{Curl}([\text{dev Curl}(A + \zeta \cdot \mathbb{1})]^T) \stackrel{(2.18), (2.16)}{\stackrel{\text{Curl} \circ \text{D} \equiv 0}{=}} -\frac{1}{3} \underbrace{\text{Anti}(\nabla \text{tr}(\text{Daxl} A))}_{\in \mathfrak{so}(3)} + \underbrace{\Delta\zeta \cdot \mathbb{1} - \text{D}^2\zeta}_{\in \text{Sym}(3)} \quad (2.30)$$

and we obtain, similar to the decomposition in (2.27):

$$\begin{aligned} \text{D}^2\zeta &= \frac{1}{2} \text{tr}(\text{Curl}([\text{dev Curl}(A + \zeta \cdot \mathbb{1})]^T)) \cdot \mathbb{1} - \text{sym}(\text{Curl}([\text{dev Curl}(A + \zeta \cdot \mathbb{1})]^T)) \\ &= \tilde{L}_4(\text{Ddev Curl}(A + \zeta \cdot \mathbb{1})). \end{aligned} \quad (2.31)$$

On the other hand, taking \mathbf{inc} of the transpositions on both sides of (2.29) gives

$$\mathbf{inc}([\text{dev Curl}(A + \zeta \cdot \mathbb{1})]^T) \stackrel{(2.24)}{=} \frac{1}{3} \Delta \text{tr}(\text{Daxl } A) \cdot \mathbb{1} - \frac{1}{3} \text{D}^2 \text{tr}(\text{Daxl } A) - \text{Anti}(\nabla \Delta \zeta), \quad (2.32)$$

yielding the relation

$$\begin{aligned} \text{D}^2 \text{tr}(\text{Daxl } A) &= \frac{3}{2} \text{tr}(\mathbf{inc}([\text{dev Curl}(A + \zeta \cdot \mathbb{1})]^T)) \cdot \mathbb{1} \\ &\quad - \text{sym}(\mathbf{inc}([\text{dev Curl}(A + \zeta \cdot \mathbb{1})]^T)) \\ &= \tilde{L}_5(\text{D}^2 \text{dev Curl}(A + \zeta \cdot \mathbb{1})). \end{aligned} \quad (2.33)$$

Considering the second distributional derivatives in (2.29) we conclude

$$\begin{aligned} \text{D}^3 \text{axl } A &= \frac{1}{3} \text{D}^2 \text{tr}(\text{Daxl } A) \cdot \mathbb{1} - \text{D}^2([\text{dev Curl}(A + \zeta \cdot \mathbb{1})]^T) + \text{D}^2 \text{Anti}(\nabla \zeta) \\ &\stackrel{(2.31)}{=} \tilde{L}_6(\text{D}^2 \text{dev Curl}(A + \zeta \cdot \mathbb{1})). \end{aligned} \quad \blacksquare$$

Remark 2.5. In the above proof we have used that the second-order differential operator \mathbf{inc} does not change the symmetry property after application on square matrix fields, cf. the Appendix. Further properties are collected e.g. in [52, Appendix], [1, Sec. 2] and [12, Sec. 6.18].

The incompatibility operator \mathbf{inc} arises in dislocation models, e.g., in the modeling of elastic materials with dislocations or in the modeling of dislocated crystals, since the strain cannot be a symmetric gradient of a vector field as soon as dislocations are present and the notion of incompatibility is at the basis of a new paradigm to describe the inelastic effects, cf. [20, 4, 3, 46], cf. the Appendix for further comments. Moreover, the equation $\mathbf{inc} \text{sym } e \equiv 0$ is equivalent to the *Saint-Venant compatibility condition*⁵ defining the relation between the symmetric strain $\text{sym } e$ and the displacement vector field u :

$$\mathbf{inc} \text{sym } e \equiv 0 \quad \Leftrightarrow \quad \text{sym } e = \text{sym } \text{D}u \quad (2.34)$$

over simply connected domains, cf. [1, 46]. In the appendix we show that the operators \mathbf{inc} and sym can be interchanged, so that

$$\mathbf{inc} \text{sym } e = \text{sym } \mathbf{inc} e = \text{sym } \text{Curl}([\text{Curl } e]^T). \quad (2.35)$$

Investigations over multiply connected domains can be found e.g. in [66, 30].

Returning to our proof, a crucial ingredient in our following argumentation is

Theorem 2.6 (Lions lemma and Nečas estimate). *Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain. Let $m \in \mathbb{Z}$ and $p \in (1, \infty)$. Then $f \in \mathcal{D}'(\Omega, \mathbb{R}^d)$ and $\text{D}f \in W^{m-1, p}(\Omega, \mathbb{R}^{d \times n})$ imply $f \in W^{m, p}(\Omega, \mathbb{R}^d)$. Moreover,*

$$\|f\|_{W^{m, p}(\Omega, \mathbb{R}^d)} \leq c \left(\|f\|_{W^{m-1, p}(\Omega, \mathbb{R}^d)} + \|\text{D}f\|_{W^{m-1, p}(\Omega, \mathbb{R}^{d \times n})} \right), \quad (2.36)$$

with a constant $c = c(m, p, n, d, \Omega) > 0$.

⁵Those compatibility conditions are contained in the third appendix §32 p. 597 et seq. of the third edition of the lecture notes *Résistance des corps solides* given by Navier and extended with several notes and appendices by Barré de Saint-Venant and published as *Résumé des Leçons données à l'École des Ponts et Chaussées sur l'Application de la Mécanique*, vol. I, Paris, 1864. Their coordinate-free version can be found in Lagally's monograph on vector calculus from 1928 [36, Ziff. 191] where it reads:

$$\nabla \times (\text{sym } \text{D}u) \times \nabla \equiv 0$$

and formally follows from the definitions of those operators, see [36, Ziff. 191], since

$$\nabla \times (\text{sym } \text{D}u) \times \nabla = \frac{1}{2} \nabla \times (\nabla \otimes u + u \otimes \nabla) \times \nabla = \frac{1}{2} [(\nabla \times \nabla) \otimes u \times \nabla + \nabla \times u \otimes (\nabla \times \nabla)] \equiv 0.$$

For the proof we refer to [2, Proposition 2.10 and Theorem 2.3], [7]. However, since we are dealing with higher order derivatives we also need a “higher order” version of the Lions lemma resp. Nečas estimate.

Corollary 2.7. *Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain, $m \in \mathbb{Z}$ and $p \in (1, \infty)$. Denote by $D^k f$ the collection of all distributional derivatives of order k . Then $f \in \mathcal{D}'(\Omega, \mathbb{R}^d)$ and $D^k f \in W^{m-k, p}(\Omega, \mathbb{R}^{d \times n^k})$ imply $f \in W^{m, p}(\Omega, \mathbb{R}^d)$. Moreover,*

$$\|f\|_{W^{m, p}(\Omega, \mathbb{R}^d)} \leq c \left(\|f\|_{W^{m-1, p}(\Omega, \mathbb{R}^d)} + \|D^k f\|_{W^{m-k, p}(\Omega, \mathbb{R}^{d \times n^k})} \right), \quad (2.37)$$

with a constant $c = c(m, p, n, d, \Omega) > 0$.

Proof. The assertion $f \in W^{m, p}(\Omega, \mathbb{R}^d)$ and the estimate (2.37) follow by inductive application of Theorem 2.6 to $D^l f$ with $l = k - 1, k - 2, \dots, 0$. Indeed, starting by applying Theorem 2.6 to $D^{k-1} f$ gives $D^{k-1} f \in W^{m-k+1, p}(\Omega, \mathbb{R}^{d \times n^{k-1}})$ as well as

$$\begin{aligned} \|D^{k-1} f\|_{W^{m-k+1, p}(\Omega, \mathbb{R}^{d \times n^{k-1}})} &\leq c \left(\|D^{k-1} f\|_{W^{m-k, p}(\Omega, \mathbb{R}^{d \times n^{k-1}})} + \|D^k f\|_{W^{m-k, p}(\Omega, \mathbb{R}^{d \times n^k})} \right) \\ &\leq c \left(\|f\|_{W^{m-1, p}(\Omega, \mathbb{R}^d)} + \|D^k f\|_{W^{m-k, p}(\Omega, \mathbb{R}^{d \times n^k})} \right). \end{aligned} \quad (2.38)$$

Now, we can apply Theorem 2.6 to $D^{k-2} f$ to deduce $D^{k-2} f \in W^{m-k+2, p}(\Omega, \mathbb{R}^{d \times n^{k-2}})$ and moreover

$$\begin{aligned} \|D^{k-2} f\|_{W^{m-k+2, p}(\Omega, \mathbb{R}^{d \times n^{k-2}})} &\leq c \left(\|D^{k-2} f\|_{W^{m-k+1, p}(\Omega, \mathbb{R}^{d \times n^{k-1}})} + \|D^{k-1} f\|_{W^{m-k+1, p}(\Omega, \mathbb{R}^{d \times n^{k-1}})} \right) \\ &\leq c \left(\|f\|_{W^{m-1, p}(\Omega, \mathbb{R}^d)} + \|D^{k-1} f\|_{W^{m-k+1, p}(\Omega, \mathbb{R}^{d \times n^{k-1}})} \right) \\ &\stackrel{(2.38)}{\leq} c \left(\|f\|_{W^{m-1, p}(\Omega, \mathbb{R}^d)} + \|D^k f\|_{W^{m-k, p}(\Omega, \mathbb{R}^{d \times n^k})} \right). \end{aligned} \quad (2.39)$$

Consequently, for all $l = k - 1, k - 2, \dots, 0$ we deduce $D^l f \in W^{m-l, p}(\Omega, \mathbb{R}^{d \times n^l})$ as well as

$$\|D^l f\|_{W^{m-l, p}(\Omega, \mathbb{R}^{d \times n^l})} \leq c \left(\|f\|_{W^{m-1, p}(\Omega, \mathbb{R}^d)} + \|D^k f\|_{W^{m-k, p}(\Omega, \mathbb{R}^{d \times n^k})} \right). \quad (2.40)$$

■

Remark 2.8. The need to consider higher order derivatives is indicated by the appearance of linear terms in the kernel of Korn’s quantitative versions, similar to the situation at the classical trace-free Korn inequalities [17, 65]. In our case we have:

Lemma 2.9. *Let $A \in L^p(\Omega, \mathfrak{so}(3))$ and $\zeta \in L^p(\Omega, \mathbb{R})$. Then we have in the distributional sense*

- (a) $\text{Curl}(A + \zeta \cdot \mathbf{1}) \equiv 0$ if and only if $A + \zeta \cdot \mathbf{1} = \text{Anti}(\tilde{A}x + b) + (\langle \text{axl } \tilde{A}, x \rangle + \beta) \cdot \mathbf{1}$ a.e. on Ω ,
- (b) $\text{dev Curl } A \equiv 0$ if and only if $A = \text{Anti}(\beta x + b)$ a.e. on Ω ,
- (c) $\text{dev Curl}(A + \zeta \cdot \mathbf{1}) \equiv 0$ if and only if $A + \zeta \cdot \mathbf{1} = \text{Anti}(\tilde{A}x + \beta x + b) + (\langle \text{axl } \tilde{A}, x \rangle + \gamma) \cdot \mathbf{1}$ a.e. on Ω ,

with constant $\tilde{A} \in \mathfrak{so}(3)$, $b \in \mathbb{R}^3$, $\beta, \gamma \in \mathbb{R}$.

Proof. Although the deductions have already been partially indicated in the literature, cf. e.g. [53, sec. 3.4] and [63, 17, 6, 64], we include it here for the sake of completeness. The “if”-parts are seen by direct calculations, cf. the relations (2.16) and (2.18):

- (a) $\text{Curl}(\text{Anti}(\tilde{A}x + b) + (\langle \text{axl } \tilde{A}, x \rangle + \beta) \cdot \mathbf{1}) = \tilde{A} - \text{Anti}(\text{axl } \tilde{A}) \equiv 0$,
- (b) $\text{dev Curl}(\text{Anti}(\beta x + b)) = \text{dev}(\text{tr}(\beta \cdot \mathbf{1}) \cdot \mathbf{1} - \beta \cdot \mathbf{1}) = \text{dev}(2\beta \cdot \mathbf{1}) \equiv 0$,

$$(c) \quad \text{dev Curl}(\text{Anti}(\tilde{A}x + \beta x + b) + (\langle \text{axl } \tilde{A}, x \rangle + \gamma) \cdot \mathbf{1}) \\ = \text{dev}(\tilde{A} + 2\beta \cdot \mathbf{1} - \text{Anti}(\text{axl } \tilde{A})) \equiv 0.$$

Now, we focus on the “only if”-directions, starting with

$$\text{Curl}(A + \zeta \cdot \mathbf{1}) \equiv 0 \stackrel{(2.18)}{\iff} \text{Anti}(\nabla \zeta) = \text{Curl } A \stackrel{(2.16)}{=} \text{tr}(\text{Daxl } A) \cdot \mathbf{1} - (\text{Daxl } A)^T.$$

Taking the trace on both sides we obtain $\text{tr}(\text{Daxl } A) = 0$ and consequently

$$\text{Anti}(\nabla \zeta) = -(\text{Daxl } A)^T, \quad (2.41)$$

hence $\text{sym}(\text{Daxl } A) = 0$. By the classical Korn’s inequality (1.5) it follows that there exists a constant skew-symmetric matrix $\tilde{A} \in \mathfrak{so}(3)$ so that $\text{Daxl } A \equiv \tilde{A}$, which implies $A = \text{Anti}(\tilde{A}x + b)$ with $b \in \mathbb{R}^3$. Furthermore, by (2.41) we obtain

$$\text{Anti}(\nabla \zeta) = \tilde{A} \quad \Rightarrow \quad \zeta = \langle \text{axl } \tilde{A}, x \rangle + \beta \quad \text{with } \beta \in \mathbb{R},$$

which establishes (a).

For part (b) we start with the relation $\text{dev Curl } A \equiv 0$ in (2.20) and have

$$\text{Anti}(\nabla \text{tr}(\text{Daxl } A)) \equiv 0 \quad \Rightarrow \quad \nabla \text{tr}(\text{Daxl } A) \equiv 0, \quad (2.42)$$

so that

$$\frac{1}{3} \text{tr}(\text{Daxl } A) = \beta \quad (2.43)$$

for some $\beta \in \mathbb{R}$. Reinserting in the deviatoric counterpart of Nye’s formula (2.19) gives

$$0 = \beta \cdot \mathbf{1} - (\text{Daxl } A)^T \quad \text{resp.} \quad \text{Daxl } A = \beta \cdot \mathbf{1} \quad \Rightarrow \quad \text{axl } A = \beta x + b \quad (2.44)$$

for some $b \in \mathbb{R}^3$ and thus $A = \text{Anti}(\beta x + b)$.

Finally, for part (c), let now $\text{dev Curl}(A + \zeta \cdot \mathbf{1}) \equiv 0$. Then considering the skew-symmetric parts of (2.30) we obtain

$$\text{Anti}(\nabla \text{tr}(\text{Daxl } A)) \equiv 0 \quad \Rightarrow \quad \nabla \text{tr}(\text{Daxl } A) \equiv 0.$$

Hence, again

$$\frac{1}{3} \text{tr}(\text{Daxl } A) = \beta \quad (2.45)$$

for some $\beta \in \mathbb{R}$, so that considering the symmetric parts of (2.29) we get

$$0 = \frac{1}{3} \text{tr}(\text{Daxl } A) \cdot \mathbf{1} - \text{sym}(\text{Daxl } A) \stackrel{(2.45)}{=} \beta \cdot \mathbf{1} - \text{sym}(\text{Daxl } A). \quad (2.46)$$

In other words, we have

$$\text{sym}(\text{D}(\text{axl } A - \beta x)) \equiv 0$$

and by (1.5), it follows that $\text{D}(\text{axl } A - \beta x)$ must be a constant skew-symmetric matrix. Thus

$$\text{axl } A = \tilde{A}x + \beta x + b \quad (2.47)$$

for some $\tilde{A} \in \mathfrak{so}(3)$, $b \in \mathbb{R}^3$ and $\beta \in \mathbb{R}$. Furthermore, by (2.29) we have

$$\text{Anti}(\nabla \zeta) \stackrel{(2.29)}{=} \text{skew}(\text{Daxl } A) \stackrel{(2.47)}{=} \tilde{A}$$

so that ζ is of the form

$$\zeta = \langle \text{axl } \tilde{A}, x \rangle + \gamma \quad (2.48)$$

for some $\gamma \in \mathbb{R}$, and we arrive at (c):

$$A + \zeta \cdot \mathbf{1} \stackrel{(2.47)}{=} \text{Anti}(\tilde{A}x + \beta x + b) + (\langle \text{axl } \tilde{A}, x \rangle + \gamma) \cdot \mathbf{1}. \quad \blacksquare$$

We are now prepared to proceed as in the proof of the generalized Korn inequality for incompatible tensor fields .

3 Main results

We will make use of the Banach space

$$W^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3}) := \{P \in L^p(\Omega, \mathbb{R}^{3 \times 3}) \mid \text{Curl } P \in L^p(\Omega, \mathbb{R}^{3 \times 3})\} \quad (3.1a)$$

equipped with the norm

$$\|P\|_{W^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})} := \left(\|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}^p + \|\text{Curl } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}^p \right)^{\frac{1}{p}}, \quad (3.1b)$$

as well as its subspace

$$W_0^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3}) := \{P \in W^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3}) \mid P \times \nu = 0 \text{ on } \partial\Omega\},$$

where ν denotes the outward unit normal vector field to $\partial\Omega$, and the tangential trace $P \times \nu$ is understood in the sense of $W^{-\frac{1}{p}, p}(\partial\Omega, \mathbb{R}^{3 \times 3})$ which is justified by partial integration, so that its trace is defined by

$$\forall Q \in W^{1-\frac{1}{p}, p'}(\partial\Omega, \mathbb{R}^{3 \times 3}) : \quad \langle P \times (-\nu), Q \rangle_{\partial\Omega} = \int_{\Omega} \langle \text{Curl } P, \tilde{Q} \rangle - \langle P, \text{Curl } \tilde{Q} \rangle dx, \quad (3.2)$$

where $\tilde{Q} \in W^{1, p'}(\Omega, \mathbb{R}^{3 \times 3})$ denotes any extension of Q in Ω . Here, $\langle \cdot, \cdot \rangle_{\partial\Omega}$ indicates the duality pairing between $W^{-\frac{1}{p}, p}(\partial\Omega, \mathbb{R}^{3 \times 3})$ and $W^{1-\frac{1}{p}, p'}(\partial\Omega, \mathbb{R}^{3 \times 3})$.

However, the appearance of the operator dev Curl on the right hand side of our designated results in this paper would suggest to work in

$$W^{1,p}(\text{dev Curl}; \Omega, \mathbb{R}^{3 \times 3}) := \{P \in L^p(\Omega, \mathbb{R}^{3 \times 3}) \mid \text{dev Curl } P \in L^p(\Omega, \mathbb{R}^{3 \times 3})\} \quad (3.3)$$

but this is, surprisingly at first glance, not a new space:

Lemma 3.1. $W^{1,p}(\text{dev Curl}; \Omega, \mathbb{R}^{3 \times 3}) = W^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$.

It is sufficient to show that the p -integrability of $\text{dev Curl } P$ already implies the p -integrability of $\text{Curl } P$, and follows from the general case:

Lemma 3.2. *Let $P \in \mathcal{D}'(\Omega, \mathbb{R}^{3 \times 3})$. Then we have for all $m \in \mathbb{Z}$ that*

$$\text{Curl } P \in W^{m,p}(\Omega, \mathbb{R}^{3 \times 3}) \Leftrightarrow \text{dev Curl } P \in W^{m,p}(\Omega, \mathbb{R}^{3 \times 3}). \quad (3.4)$$

Proof. We again consider the decomposition of P into its symmetric and skew-symmetric part, i.e.

$$P = S + A = S + \text{Anti}(a) \quad \text{for some } S \in \text{Sym}(3), A \in \mathfrak{so}(3) \text{ and with } a = \text{axl}(A).$$

Then by Nye's formula (2.16a) we have

$$\text{Curl } P = \text{Curl}(S + \text{Anti}(a)) \stackrel{(2.16)}{=} \text{Curl } S + \text{div } a \cdot \mathbf{1} - (\text{D}a)^T \quad (3.5)$$

and in view of $\text{tr}(\text{Curl } S) = 0$ we obtain

$$\text{dev Curl } P = \text{Curl } S - (\text{D}a)^T + \frac{1}{3} \text{div } a \cdot \mathbf{1} \quad (3.6)$$

so that taking the Curl of the transpositions on both sides gives

$$\text{Curl}([\text{dev Curl } P]^T) \stackrel{\text{Curl} \circ \text{D} \equiv 0}{\stackrel{(2.18)}{=}} \underbrace{\text{inc } S}_{\in \text{Sym}(3)} - \frac{1}{3} \underbrace{\text{Anti}(\nabla \text{div } a)}_{\in \mathfrak{so}(3)}, \quad (3.7)$$

which gives

$$\text{skew Curl}([\text{dev Curl } P]^T) = -\frac{1}{3} \text{Anti}(\nabla \text{div } a). \quad (3.8)$$

Thus, $\text{dev Curl } P \in W^{m,p}(\Omega, \mathbb{R}^{3 \times 3})$ implies $\text{Curl}([\text{dev Curl } P]^T) \in W^{m-1,p}(\Omega, \mathbb{R}^{3 \times 3})$ as well as

$$\begin{aligned} \text{skew Curl}([\text{dev Curl } P]^T) &= \frac{1}{2} (\text{Curl}([\text{dev Curl } P]^T) - [\text{Curl}([\text{dev Curl } P]^T)]^T) \\ &\in W^{m-1,p}(\Omega, \mathbb{R}^{3 \times 3}), \end{aligned} \quad (3.9)$$

so that we obtain

$$\nabla \text{div } a \stackrel{(3.8)}{=} -3 \text{axl skew Curl}([\text{dev Curl } P]^T) \in W^{m-1,p}(\Omega, \mathbb{R}^3). \quad (3.10)$$

Since $a = \text{axl skew } P \in \mathcal{D}'(\Omega, \mathbb{R}^3)$, we apply Theorem 2.6 to $\text{div } a \in \mathcal{D}'(\Omega, \mathbb{R})$ to conclude from (3.10) that $\text{div } a \in W^{m,p}(\Omega, \mathbb{R})$. The statement of the Lemma then follows from the decompositions (3.5) and (3.6) which give the expression

$$\text{Curl } P = \text{dev Curl } P + \frac{2}{3} \text{div } a \cdot \mathbf{1} \in W^{m,p}(\Omega, \mathbb{R}^{3 \times 3}). \quad (3.11) \quad \blacksquare$$

Corollary 3.3. *The classical Hilbert space $H(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$ coincides with the Hilbert space $H(\text{dev Curl}; \Omega, \mathbb{R}^{3 \times 3}) := \{P \in L^2(\Omega, \mathbb{R}^{3 \times 3}) \mid \text{dev Curl } P \in L^2(\Omega, \mathbb{R}^{3 \times 3})\}$.*

Remark 3.4 (Equivalence of norms). In view of (3.10) an application of the Lions lemma to $\text{div } a$, with $a = \text{axl skew } P$, gives us $\text{div } a \in W^{m,p}(\Omega, \mathbb{R})$. Moreover, by the Nečas estimate we have

$$\begin{aligned} \|\text{div } a\|_{W^{m,p}(\Omega, \mathbb{R})} &\leq c_1 (\|\text{div } a\|_{W^{m-1,p}(\Omega, \mathbb{R})} + \|\nabla \text{div } a\|_{W^{m-1,p}(\Omega, \mathbb{R}^3)}) \\ &\stackrel{(3.10)}{\leq} c_2 (\|\text{div axl skew } P\|_{W^{m-1,p}(\Omega, \mathbb{R})} + \|\text{Curl}([\text{dev Curl } P]^T)\|_{W^{m-1,p}(\Omega, \mathbb{R}^{3 \times 3})}) \\ &\leq c_3 (\|P\|_{W^{m,p}(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{W^{m,p}(\Omega, \mathbb{R}^{3 \times 3})}), \end{aligned} \quad (3.12)$$

provided that $P \in W^{m,p}(\Omega, \mathbb{R}^{3 \times 3})$. Together with (3.11) we conclude:

$$\|\text{Curl } P\|_{W^{m,p}(\Omega, \mathbb{R}^{3 \times 3})} \leq c_4 (\|P\|_{W^{m,p}(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{W^{m,p}(\Omega, \mathbb{R}^{3 \times 3})}) \quad (3.13)$$

as well as

$$\|P\|_{W^{m,p}(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{Curl } P\|_{W^{m,p}(\Omega, \mathbb{R}^{3 \times 3})} \leq c_5 (\|P\|_{W^{m,p}(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{W^{m,p}(\Omega, \mathbb{R}^{3 \times 3})}) \quad (3.14)$$

and especially for $m = 0$:

$$\|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{Curl } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c_5 (\|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}) \quad (3.15)$$

for all $P \in W^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$.⁶

⁶This result also follows from the open mapping theorem (also known as Banach-Schauder theorem [12, Thm 5.6-1]) in functional analysis. More precisely, the latter provides the following sufficient condition for two norms to be equivalent in an infinite-dimensional space, see [12, Thm 5.6-4]:

Corollary 3.5. *Let $\|\cdot\|$ and $\|\cdot\|'$ be two norms on the same vector space X , with the following properties: both spaces $(X, \|\cdot\|)$ and $(X, \|\cdot\|')$ are complete, and there exists a constant C such that*

$$\|x\|' \leq C \|x\| \quad \text{for all } x \in X.$$

Then the two norms $\|\cdot\|$ and $\|\cdot\|'$ are equivalent.

Remark 3.6. The last identity in (3.11), which could also be formally obtained from (2.8) with $b = -\nabla$, together with the expression (3.10) gives for general matrix field $P \in \mathcal{D}'(\Omega, \mathbb{R}^{3 \times 3})$:

$$\text{DCurl } P = L(\text{D dev Curl } P). \quad (3.16)$$

Thus, recalling (1.7), we arrive directly at the case (b) of Lemma 2.4.

Corollary 3.7. *Notably, the trace condition in $W_0^{1,p}(\text{dev Curl}; \Omega, \mathbb{R}^{3 \times 3})$ would read $\text{dev}(P \times \nu) = 0$ on $\partial\Omega$, to be understood by partial integration via*

$$\begin{aligned} \forall Q \in W^{1-\frac{1}{p'}, p'}(\partial\Omega, \mathbb{R}^{3 \times 3}) : \quad & \langle \text{dev}(P \times (-\nu)), Q \rangle_{\partial\Omega} = \int_{\Omega} \langle \text{dev Curl } P, \tilde{Q} \rangle - \langle P, \text{Curl dev } \tilde{Q} \rangle \, dx \quad (3.17) \\ & = \int_{\Omega} \langle \text{Curl } P, \text{dev } \tilde{Q} \rangle - \langle P, \text{Curl dev } \tilde{Q} \rangle \, dx \\ & \stackrel{(3.2)}{=} \langle P \times (-\nu), \text{dev } Q \rangle_{\partial\Omega}, \end{aligned}$$

where $\tilde{Q} \in W^{1, p'}(\Omega, \mathbb{R}^{3 \times 3})$ denotes any extension of Q in Ω . However, it follows from Observation 2.1 that the boundary conditions $P \times \nu = 0$ and $\text{dev}(P \times \nu) = 0$ on $\partial\Omega$ are the same.

Lemma 3.8. *Let $\Omega \subset \mathbb{R}^3$ be a bounded Lipschitz domain, $1 < p < \infty$ and $P \in \mathcal{D}'(\Omega, \mathbb{R}^{3 \times 3})$. Then either of the conditions*

- (a) $\text{dev sym } P \in L^p(\Omega, \mathbb{R}^{3 \times 3})$ and $\text{Curl } P \in W^{-1, p}(\Omega, \mathbb{R}^{3 \times 3})$,
- (b) $\text{sym } P \in L^p(\Omega, \mathbb{R}^{3 \times 3})$ and $\text{dev Curl } P \in W^{-1, p}(\Omega, \mathbb{R}^{3 \times 3})$,
- (c) $\text{dev sym } P \in L^p(\Omega, \mathbb{R}^{3 \times 3})$ and $\text{dev Curl } P \in W^{-1, p}(\Omega, \mathbb{R}^{3 \times 3})$,

implies $P \in L^p(\Omega, \mathbb{R}^{3 \times 3})$. Moreover, we have the corresponding estimates

$$\begin{aligned} \|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \left(\|\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1}\|_{W^{-1, p}(\Omega, \mathbb{R}^{3 \times 3})} \right. \\ \left. + \|\text{dev sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{Curl } P\|_{W^{-1, p}(\Omega, \mathbb{R}^{3 \times 3})} \right), \quad (3.18a) \end{aligned}$$

$$\begin{aligned} \|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \left(\|\text{skew } P\|_{W^{-1, p}(\Omega, \mathbb{R}^{3 \times 3})} \right. \\ \left. + \|\text{sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{W^{-1, p}(\Omega, \mathbb{R}^{3 \times 3})} \right), \quad (3.18b) \end{aligned}$$

$$\begin{aligned} \|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \left(\|\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1}\|_{W^{-1, p}(\Omega, \mathbb{R}^{3 \times 3})} \right. \\ \left. + \|\text{dev sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{W^{-1, p}(\Omega, \mathbb{R}^{3 \times 3})} \right), \quad (3.18c) \end{aligned}$$

each with a constant $c = c(p, \Omega) > 0$.

Proof. We start by proving part (b). For that purpose we will follow the proof of [43, Lemma 3.1]. Thus, for part (b) it remains to deduce that $\text{skew } P \in L^p(\Omega, \mathbb{R}^{3 \times 3})$. We have

$$\begin{aligned} \|\text{D}^2 \text{skew } P\|_{W^{-2, p}(\Omega, \mathbb{R}^{3 \times 3 \times 3})} & \stackrel{\text{Lem. 2.4(b)}}{\leq} c \|\text{D dev Curl skew } P\|_{W^{-2, p}(\Omega, \mathbb{R}^{3 \times 3 \times 2})} \\ & \leq c \|\text{dev Curl}(P - \text{sym } P)\|_{W^{-1, p}(\Omega, \mathbb{R}^{3 \times 3})} \\ & \leq c (\|\text{dev Curl } P\|_{W^{-1, p}(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{Curl sym } P\|_{W^{-1, p}(\Omega, \mathbb{R}^{3 \times 3})}) \\ & \leq c (\|\text{sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{W^{-1, p}(\Omega, \mathbb{R}^{3 \times 3})}). \quad (3.19) \end{aligned}$$

Hence, the assumptions of part (b) yield $D^2 \text{skew } P \in W^{-2,p}(\Omega, \mathbb{R}^{3 \times 3^3})$, so that, by Corollary 2.7, we obtain $\text{skew } P \in L^p(\Omega, \mathbb{R}^{3 \times 3})$ and moreover the estimate

$$\begin{aligned} \|\text{skew } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} &\leq c(\|\text{skew } P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} + \|D^2 \text{skew } P\|_{W^{-2,p}(\Omega, \mathbb{R}^{3 \times 3^3})}) \\ &\stackrel{(3.19)}{\leq} c\left(\|\text{skew } P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})}\right). \end{aligned} \quad (3.20)$$

Then by adding $\|\text{sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}$ on both sides we obtain (3.18b).

Clearly, the conclusion of (a) as well as the estimate (3.18a) follow from (c) and (3.18c), respectively. To establish (c), we make use of the orthogonal decomposition $P = \text{dev sym } P + (\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1})$. Then, to obtain $\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1} \in L^p(\Omega, \mathbb{R}^{3 \times 3})$ for (c), we consider

$$\begin{aligned} &\|D^2 \text{dev Curl}(\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1})\|_{W^{-3,p}(\Omega, \mathbb{R}^{3 \times 3^3})} \\ &\leq c \|\text{dev Curl}(P - \text{dev sym } P)\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} \\ &\leq c(\|\text{dev Curl } P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{Curl dev sym } P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})}) \\ &\leq c(\|\text{dev Curl } P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}). \end{aligned} \quad (3.21)$$

Therefore, $D^2 \text{dev Curl}(\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1}) \in W^{-3,p}(\Omega, \mathbb{R}^{3 \times 3^3})$ follows from the assumptions of (c) and Lemma 2.4 (c) implies

$$D^3(\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1}) \in W^{-3,p}(\Omega, \mathbb{R}^{3 \times 3^4}). \quad (3.22)$$

Applying Corollary 2.7 again, this time to $\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1}$, we arrive at $\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1} \in L^p(\Omega, \mathbb{R}^{3 \times 3})$ and, moreover,

$$\begin{aligned} \|\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1}\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} &\leq c\left(\|\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1}\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} \right. \\ &\quad \left. + \|D^3(\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1})\|_{W^{-3,p}(\Omega, \mathbb{R}^{3 \times 3^4})}\right) \\ &\stackrel{\text{Lem. 2.4 (c)}}{\leq} c\left(\|\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1}\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} \right. \\ &\quad \left. + \|D^2 \text{dev Curl}(\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1})\|_{W^{-3,p}(\Omega, \mathbb{R}^{3 \times 3^3})}\right) \\ &\stackrel{(3.21)}{\leq} c\left(\|\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1}\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} \right. \\ &\quad \left. + \|\text{dev sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})}\right). \quad \blacksquare \end{aligned} \quad (3.23)$$

Remark 3.9. Of course, part (a) can also be proven independently of part (c). Indeed, using Lemma 2.4 (a) we obtain

$$\begin{aligned} \|D^2(\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1})\|_{W^{-2,p}(\Omega, \mathbb{R}^{3 \times 3^3})} &\stackrel{\text{Lem. 2.4 (a)}}{\leq} c \|D \text{Curl}(\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1})\|_{W^{-2,p}(\Omega, \mathbb{R}^{3 \times 3^2})} \\ &\leq c \|\text{Curl}(P - \text{dev sym } P)\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} \\ &\leq c(\|\text{Curl } P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}) \end{aligned} \quad (3.24)$$

and the conclusion follows from an application of Corollary 2.7 to $\text{skew } P + \frac{1}{3} \text{tr } P \cdot \mathbb{1}$.

The rigidity results now follow by elimination of the corresponding first term on the right-hand side.

Theorem 3.10. *Let $\Omega \subset \mathbb{R}^3$ be a bounded Lipschitz domain and $1 < p < \infty$. There exists a constant $c = c(p, \Omega) > 0$ such that for all $P \in L^p(\Omega, \mathbb{R}^{3 \times 3})$ we have*

$$\inf_{T \in K_{dS,C}} \|P - T\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \left(\|\operatorname{dev} \operatorname{sym} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\operatorname{Curl} P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} \right), \quad (3.25a)$$

$$\inf_{T \in K_{S,dC}} \|P - T\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \left(\|\operatorname{sym} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\operatorname{dev} \operatorname{Curl} P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} \right), \quad (3.25b)$$

$$\inf_{T \in K_{dS,dC}} \|P - T\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \left(\|\operatorname{dev} \operatorname{sym} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\operatorname{dev} \operatorname{Curl} P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} \right), \quad (3.25c)$$

where the kernels are given, respectively, by

$$K_{dS,C} = \{T : \Omega \rightarrow \mathbb{R}^{3 \times 3} \mid T(x) = \operatorname{Anti}(\tilde{A}x + b) + (\langle \operatorname{axl} \tilde{A}, x \rangle + \beta) \cdot \mathbb{1}, \tilde{A} \in \mathfrak{so}(3), b \in \mathbb{R}^3, \beta \in \mathbb{R}\}, \quad (3.26a)$$

$$K_{S,dC} = \{T : \Omega \rightarrow \mathbb{R}^{3 \times 3} \mid T(x) = \operatorname{Anti}(\beta x + b), b \in \mathbb{R}^3, \beta \in \mathbb{R}\}, \quad (3.26b)$$

$$K_{dS,dC} = \{T : \Omega \rightarrow \mathbb{R}^{3 \times 3} \mid T(x) = \operatorname{Anti}(\tilde{A}x + \beta x + b) + (\langle \operatorname{axl} \tilde{A}, x \rangle + \gamma) \cdot \mathbb{1}, \\ \tilde{A} \in \mathfrak{so}(3), b \in \mathbb{R}^3, \beta, \gamma \in \mathbb{R}\}. \quad (3.26c)$$

Proof. We proceed as in the proof of Korn's inequalities (1.4) resp. (1.5), see [43, Theorem 3.3] resp. [12, Theorem 6.15-3], and start by characterizing the kernel of the right-hand side,

$$K_{dS,C} := \{P \in L^p(\Omega, \mathbb{R}^{3 \times 3}) \mid \operatorname{dev} \operatorname{sym} P = 0 \text{ a.e. and} \\ \operatorname{Curl} P = 0 \text{ in the distributional sense}\},$$

so that $P \in K_{dS,C}$ if and only if $P = \operatorname{skew} P + \frac{1}{3} \operatorname{tr} P \cdot \mathbb{1}$ and $\operatorname{Curl}(\operatorname{skew} P + \frac{1}{3} \operatorname{tr} P \cdot \mathbb{1}) \equiv 0$. Hence, (3.26a) follows by virtue of Lemma 2.9 (a).

Let us denote by e_1, \dots, e_M a basis of $K_{dS,C}$, where $M := \dim K_{dS,C} = 7$, and by ℓ_1, \dots, ℓ_M the corresponding continuous linear forms on $K_{dS,C}$ given by

$$\ell_\alpha(e_j) := \delta_{\alpha j}. \quad (3.27)$$

By the Hahn-Banach theorem in a normed vector space (see e.g. [12, Theorem 5.9-1]), we extend ℓ_α to continuous linear forms - again denoted by ℓ_α - on the Banach space $L^p(\Omega, \mathbb{R}^{3 \times 3})$, $1 \leq \alpha \leq M$. Notably,

$$T \in K_{dS,C} \text{ is equal to } 0 \quad \Leftrightarrow \quad \ell_\alpha(T) = 0 \quad \forall \alpha \in \{1, \dots, M\}.$$

Following the proof of [43, Theorem 3.4] we eliminate the first term on the right-hand side of (3.18a) by exploiting the compactness $L^p(\Omega, \mathbb{R}^{3 \times 3}) \subset\subset W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})$ and arrive at

$$\|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \left(\|\operatorname{dev} \operatorname{sym} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\operatorname{Curl} P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} + \sum_{\alpha=1}^M |\ell_\alpha(P)| \right). \quad (3.28)$$

Indeed, if (3.28) were false, there would exist a sequence $P_k \in L^p(\Omega, \mathbb{R}^{3 \times 3})$ such that

$$\|P_k\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} = 1 \quad \text{and} \quad \left(\|\operatorname{dev} \operatorname{sym} P_k\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\operatorname{Curl} P_k\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} + \sum_{\alpha=1}^M |\ell_\alpha(P_k)| \right) < \frac{1}{k}.$$

Thus, for a subsequence $P_k \rightharpoonup P^*$ in $L^p(\Omega, \mathbb{R}^{3 \times 3})$ with $\operatorname{dev} \operatorname{sym} P^* = 0$ a.e., $\operatorname{sym} \operatorname{Curl} P^* = 0$ in the distributional sense and $\ell_\alpha(P_k) = 0$ for all $\alpha = 1, \dots, M$, so that $P^* = 0$ a.e.. By the compact embedding $L^p(\Omega, \mathbb{R}^{3 \times 3}) \subset\subset W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})$ there exists a subsequence P_k , so that $\operatorname{skew} P_k + \frac{1}{3} \operatorname{tr} P_k \cdot \mathbb{1} \rightarrow 0$ in $W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})$. This is a contradiction to (3.18a).

Considering now the projection $\pi_\alpha : L^p(\Omega, \mathbb{R}^{3 \times 3}) \rightarrow K_{dS,C}$ given by

$$\pi_\alpha(P) := \sum_{j=1}^M \ell_j(P) e_j \quad (3.29)$$

we obtain $\ell_\alpha(P - \pi_\alpha(P)) \stackrel{(3.27)}{=} 0$ for all $1 \leq \alpha \leq M$, so that (3.25a) follows after applying (3.28) to $P - \pi_\alpha(P)$.

Furthermore, we obtain the characterizations (3.26b) and (3.26c) by Lemma 2.9 (b) and (c), respectively, since

$$K_{S,dC} := \{P \in L^p(\Omega, \mathbb{R}^{3 \times 3}) \mid \text{sym } P = 0 \text{ a.e. and } \text{dev Curl } P = 0 \text{ in the distributional sense}\} \quad (3.30)$$

$$\stackrel{\text{Lemma 2.9 (b)}}{=} \{T : \Omega \rightarrow \mathbb{R}^{3 \times 3} \mid T(x) = \text{Anti}(\beta x + b), \ b \in \mathbb{R}^3, \beta \in \mathbb{R}\}$$

and

$$K_{dS,dC} := \{P \in L^p(\Omega, \mathbb{R}^{3 \times 3}) \mid \text{dev sym } P = 0 \text{ a.e. and } \text{dev Curl } P = 0 \text{ in the distributional sense}\} \quad (3.31)$$

$$\stackrel{\text{Lemma 2.9 (c)}}{=} \{T : \Omega \rightarrow \mathbb{R}^{3 \times 3} \mid T(x) = \text{Anti}(\tilde{A}x + \beta x + b) + (\langle \text{axl } \tilde{A}, x \rangle + \gamma) \cdot \mathbf{1},$$

$$\tilde{A} \in \mathfrak{so}(3), b \in \mathbb{R}^3, \beta, \gamma \in \mathbb{R}\}$$

with $\dim K_{S,dC} = 4$ and $\dim K_{dS,dC} = 8$. Hence, we can argue as above to deduce (3.25b) and (3.25c) from (3.18b) and (3.18c), respectively, since we end up with

$$\|P - \pi_b(P)\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c (\|\text{sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})}) \quad (3.32)$$

and

$$\|P - \pi_c(P)\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c (\|\text{dev sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})}) \quad (3.33)$$

respectively, with projections $\pi_b : L^p(\Omega, \mathbb{R}^{3 \times 3}) \rightarrow K_{S,dC}$ and $\pi_c : L^p(\Omega, \mathbb{R}^{3 \times 3}) \rightarrow K_{dS,dC}$. \blacksquare

Finally, the kernel is killed by the tangential trace condition $P \times \nu \equiv 0$ ($\Leftrightarrow \text{dev}(P \times \nu) = 0$, cf. Obs. 2.1):

Theorem 3.11. *Let $\Omega \subset \mathbb{R}^3$ be a bounded Lipschitz domain and $1 < p < \infty$. There exists a constant $c = c(p, \Omega) > 0$ such that for all $P \in W_0^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$ we have*

$$\|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c (\|\text{dev sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}). \quad (3.34)$$

Proof. We argue as in the proof of [43, Theorem 3.5] and consider a sequence $\{P_k\}_{k \in \mathbb{N}} \subset W_0^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$ which converges weakly in $L^p(\Omega, \mathbb{R}^{3 \times 3})$ to P^* so that $\text{dev sym } P^* = 0$ a.e. and $\text{dev Curl } P^* = 0$ in the distributional sense, i.e. $P^* \in K_{dS,dC}$, where

$$K_{dS,dC} \stackrel{(3.26c)}{=} \{T : \Omega \rightarrow \mathbb{R}^{3 \times 3} \mid T(x) = \text{Anti}(\tilde{A}x + \beta x + b) + (\langle \text{axl } \tilde{A}, x \rangle + \gamma) \cdot \mathbf{1},$$

$$\tilde{A} \in \mathfrak{so}(3), b \in \mathbb{R}^3, \beta, \gamma \in \mathbb{R}\}.$$

By (3.17) it further follows that $\langle \text{dev}(P^* \times (-\nu)), Q \rangle_{\partial\Omega} = 0$ for all $Q \in W^{1-\frac{1}{p'}, p'}(\partial\Omega, \mathbb{R}^{3 \times 3})$. However, since $P^* \in K_{dS,dC}$ also has an explicit representation, the boundary condition $\text{dev}(P^* \times \nu) = 0$ is also valid in the classical sense. Furthermore, we deduce by Observation 2.1 that $P^* \times \nu = 0$ on $\partial\Omega$, so that $P^* \in W_0^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$. Again, using the explicit representation of $P^* = \text{Anti}(\tilde{A}x + \beta x + b) + (\langle \text{axl } \tilde{A}, x \rangle + \gamma) \cdot \mathbf{1}$, we conclude with Observation 2.2 that, in fact, $P^* \equiv 0$:

$$[\text{Anti}(\tilde{A}x + \beta x + b) + (\langle \text{axl } \tilde{A}, x \rangle + \gamma) \cdot \mathbf{1}] \times \nu = 0$$

$$\stackrel{\text{Obs. 2.2}}{\Rightarrow} \tilde{A}x + \beta x + b = 0 \quad \text{and} \quad \langle \text{axl } \tilde{A}, x \rangle + \gamma = 0 \quad \text{for all } x \in \partial\Omega$$

$$\Rightarrow \gamma = 0, \tilde{A} = 0 \quad \Rightarrow \quad b = 0, \beta = 0. \quad \blacksquare$$

Remark 3.12. Similarly, the following estimates can also be deduced, even independently of (3.34), for $P \in W_0^{1,p}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$:

$$\|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c (\|\text{dev sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{Curl } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}), \quad (3.35)$$

$$\|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c (\|\text{sym } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{dev Curl } P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}). \quad (3.36)$$

Since by [6, Theorem 3.1 (ii)] it holds

$$\|\operatorname{Curl} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \|\operatorname{dev} \operatorname{Curl} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \quad \text{for } P \in W_0^{1,p}(\operatorname{Curl}; \Omega, \mathbb{R}^{3 \times 3}), \quad (3.37)$$

we can recover (3.34) from (3.35) and (3.37).

However, without boundary conditions the Nečas estimate provides for $P \in W^{1,p}(\operatorname{Curl}; \Omega, \mathbb{R}^{3 \times 3})$:

$$\begin{aligned} \|\operatorname{Curl} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} &\stackrel{(2.36)}{\leq} c (\|\operatorname{Curl} P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} + \|\operatorname{DCurl} P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3^2})}) \\ &\stackrel{(3.16)}{\leq} c (\|\operatorname{Curl} P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} + \|\operatorname{Ddev} \operatorname{Curl} P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3^2})}) \\ &\leq c (\|\operatorname{Curl} P\|_{W^{-1,p}(\Omega, \mathbb{R}^{3 \times 3})} + \|\operatorname{dev} \operatorname{Curl} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}). \end{aligned} \quad (3.38)$$

Remark 3.13. Among the inequalities (3.34), (3.35) and (3.36) we expect (3.35) also to hold true in higher space dimensions $n > 3$, see the discussion in our Introduction.

Remark 3.14. Regarding (3.15) and (3.34) or (3.37) and (3.34) we obtain the norm equivalence

$$\|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\operatorname{Curl} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c (\|\operatorname{dev} \operatorname{sym} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\operatorname{dev} \operatorname{Curl} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})})$$

for tensor fields $P \in W_0^{1,p}(\operatorname{Curl}; \Omega, \mathbb{R}^{3 \times 3})$.

For $P = \operatorname{D}u$ in (3.34) we recover the following tangential trace-free Korn inequality:

Corollary 3.15. *Let $\Omega \subset \mathbb{R}^3$ be a bounded Lipschitz domain and $1 < p < \infty$. There exists a constant $c = c(p, \Omega) > 0$ such that for all $u \in W^{1,p}(\Omega, \mathbb{R}^3)$ with $\operatorname{D}u \times \nu = 0$ on $\partial\Omega$ we have*

$$\|\operatorname{D}u\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c \|\operatorname{dev} \operatorname{sym} \operatorname{D}u\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}. \quad (3.39)$$

For skew-symmetric $P = \operatorname{Anti}(a)$ we recover from (3.34) a Poincaré inequality involving only the deviatoric (trace-free) part of the gradient:

Corollary 3.16. *Let $\Omega \subset \mathbb{R}^3$ be a bounded Lipschitz domain and $1 < p < \infty$. There exists a constant $c = c(p, \Omega) > 0$ such that for all $a \in W_0^{1,p}(\Omega, \mathbb{R}^3)$ we have*

$$\|a\|_{L^p(\Omega, \mathbb{R}^3)} \leq c \|\operatorname{dev} \operatorname{D}a\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}. \quad (3.40)$$

Proof. This follows from Theorem 3.11 by setting $P = \operatorname{Anti}(a)$ and the following observations:

$\operatorname{Anti}(a) \times \nu = 0 \Leftrightarrow a = 0$ on $\partial\Omega$, see Observation 2.2, $\operatorname{Curl}(\operatorname{Anti}(a)) = L(\operatorname{D}a)$, see (2.16a) and the form of $\operatorname{Anti}(a)$, see (2.2). \blacksquare

Remark 3.17. The previous results also hold true for functions with vanishing tangential trace only on a relatively open (non-empty) subset $\Gamma \subseteq \partial\Omega$ of the boundary. So, e.g., we have

$$\|P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} \leq c (\|\operatorname{dev} \operatorname{sym} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})} + \|\operatorname{dev} \operatorname{Curl} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}) \quad (3.41)$$

for all $P \in W_{\Gamma,0}^{1,p}(\operatorname{Curl}; \Omega, \mathbb{R}^{3 \times 3})$, which is the completion of $C_{\Gamma,0}^\infty(\Omega, \mathbb{R}^{3 \times 3})$ with respect to the $W^{1,p}(\operatorname{Curl}; \Omega, \mathbb{R}^{3 \times 3})$ -norm.

Remark 3.18. In [28] the authors proved that in $n = 2$ dimensions, for $p = 2$ a Korn inequality for incompatible fields also holds true when $\operatorname{Curl} P$ is only in L^1 (actually when it is a measure with bounded total variation) under the normalization condition $\int_\Omega \operatorname{skew} P \, dx = 0$. In terms of scaling, it is interesting to involve in (3.34) the Sobolev exponent. So, we will show in a forthcoming paper that for $1 < p < 3$ the following estimate holds true on an arbitrary open set $\Omega \subseteq \mathbb{R}^3$:

$$\|P\|_{L^{p^*}(\Omega, \mathbb{R}^{3 \times 3})} \leq c (\|\operatorname{dev} \operatorname{sym} P\|_{L^{p^*}(\Omega, \mathbb{R}^{3 \times 3})} + \|\operatorname{dev} \operatorname{Curl} P\|_{L^p(\Omega, \mathbb{R}^{3 \times 3})}) \quad (3.42)$$

for all $P \in C_c^\infty(\Omega, \mathbb{R}^{3 \times 3})$, where $p^* = \frac{3p}{3-p}$. However, we do not know if such a result still holds in the borderline case $p = 1$.

Acknowledgment

The authors are grateful for inspiring discussions with Stefan Müller (Hausdorff Center for Mathematics, Bonn, Germany) and thank also the anonymous referee for his valuable comments and suggestions. This work was initiated in the framework of the Priority Programme SPP 2256 'Variational Methods for Predicting Complex Phenomena in Engineering Structures and Materials' funded by the Deutsche Forschungsgemeinschaft (DFG, German research foundation), Project-ID 422730790. The second author was supported within the project 'A variational scale-dependent transition scheme - from Cauchy elasticity to the relaxed micromorphic continuum' (Project-ID 440935806). Moreover, both authors were supported in the Project-ID 415894848 by the Deutsche Forschungsgemeinschaft.

References

- [1] C. Amrouche, P.G. Ciarlet, L. Gratie, and S. Kesavan. "On Saint Venant's compatibility conditions and Poincaré's lemma". *Comptes Rendus. Mathématique. Académie des Sciences, Paris* 342.11 (2006). Pp. 887–891.
- [2] C. Amrouche and V. Girault. "Decomposition of vector spaces and application to the Stokes problem in arbitrary dimension". *Czechoslovak Mathematical Journal* 44(119).1 (1994). Pp. 109–140.
- [3] S. Amstutz and N. Van Goethem. "Analysis of the incompatibility operator and application in intrinsic elasticity with dislocations". *SIAM Journal on Mathematical Analysis* 48.1 (2016). Pp. 320–348.
- [4] S. Amstutz and N. Van Goethem. "The incompatibility operator: From Riemann's intrinsic view of geometry to a new model of elasto-plasticity". *CIM Series in Mathematical Sciences* (2019). Ed. by J.F. Rodrigues and M. Hintermüller. Pp. 33–70.
- [5] N. Aronszajn. "On coercive integro-differential quadratic forms". *Conference on Partial Differential Equations, Univ. Kansas, Summer 1954*. 1955, pp. 94–106.
- [6] S. Bauer, P. Neff, D. Pauly, and G. Starke. "Dev-Div- and DevSym-DevCurl-inequalities for incompatible square tensor fields with mixed boundary conditions". *ESAIM: Control, Optimisation and Calculus of Variations* 22.1 (2016). Pp. 112–133.
- [7] W. Borchers and H. Sohr. "On the equations $\operatorname{rot} \mathbf{v} = \mathbf{g}$ and $\operatorname{div} \mathbf{u} = f$ with zero boundary conditions". *Hokkaido Mathematical Journal* 19.1 (1990). Pp. 67–87.
- [8] D. Breit, A. Cianchi, and L. Diening. "Trace-free Korn inequalities in Orlicz spaces". *SIAM Journal on Mathematical Analysis* 49.4 (2017). Pp. 2496–2526.
- [9] W. Chen and J. Jost. "A Riemannian version of Korn's inequality". *Calculus of Variations and Partial Differential Equations* 14.4 (2002). Pp. 517–530.
- [10] P.G. Ciarlet. "An introduction to differential geometry with applications to elasticity". *Journal of Elasticity* 78-79.1-3 (2005). Pp. 3–201.
- [11] P.G. Ciarlet. "On Korn's inequality". *Chinese Annals of Mathematics. Series B* 31.5 (2010). Pp. 607–618.
- [12] P.G. Ciarlet. *Linear and Nonlinear Functional Analysis with Applications*. Society for Industrial and Applied Mathematics, Philadelphia, PA, 2013.
- [13] P.G. Ciarlet and P. Ciarlet Jr. "Another approach to linearized elasticity and a new proof of Korn's inequality". *Mathematical Models and Methods in Applied Sciences* 15.2 (2005). Pp. 259–271.
- [14] P.G. Ciarlet, M. Malin, and C. Mardare. "On a vector version of a fundamental lemma of J. L. Lions". *Chinese Annals of Mathematics. Series B* 39.1 (2018). Pp. 33–46.
- [15] S. Conti, G. Dolzmann, and S. Müller. "Korn's second inequality and geometric rigidity with mixed growth conditions". *Calculus of Variations and Partial Differential Equations* 50.1-2 (2014). Pp. 437–454.
- [16] S. Conti, D. Faraco, and F. Maggi. "A new approach to counterexamples to L^1 estimates: Korn's inequality, geometric rigidity, and regularity for gradients of separately convex functions". *Archive for Rational Mechanics and Analysis* 175.2 (2005). Pp. 287–300.
- [17] S. Dain. "Generalized Korn's inequality and conformal Killing vectors". *Calculus of Variations and Partial Differential Equations* 25.4 (2006). Pp. 535–540.
- [18] Z. Ding and B. Li. "A conformal Korn inequality on Hölder domains". *Journal of Mathematical Analysis and Applications* 481.1 (2020). P. 14, Id/No 123440.
- [19] G. Duvaut and J.-L. Lions. *Les Inéquations en Mécanique et en Physique*. Dunod, Paris, 1972.

- [20] F. Ebobisse and P. Neff. “A fourth order gauge-invariant gradient plasticity model for polycrystals based on Kröner’s incompatibility tensor”. *Mathematics and Mechanics of Solids* 25.2 (2020). Pp. 129–159.
- [21] F. Ebobisse, P. Neff, and E. C. Aifantis. “Existence result for a dislocation based model of single crystal gradient plasticity with isotropic or linear kinematic hardening”. *The Quarterly Journal of Mechanics and Applied Mathematics* 71.1 (2018). Pp. 99–124.
- [22] F. Ebobisse, P. Neff, and S. Forest. “Well-posedness for the microcurl model in both single and polycrystal gradient plasticity”. *International Journal of Plasticity* 107 (2018). Pp. 1–26.
- [23] D. Faraco and X. Zhong. “Geometric rigidity of conformal matrices”. *Annali della Scuola Normale Superiore di Pisa. Classe di Scienze. Serie V* 4.4 (2005). Pp. 557–585.
- [24] G. Friesecke, R. D. James, and S. Müller. “A theorem on geometric rigidity and the derivation of nonlinear plate theory from three-dimensional elasticity”. *Communications on Pure and Applied Mathematics* 55.11 (2002). Pp. 1461–1506.
- [25] M. Fuchs. “Generalizations of Korn’s inequality based on gradient estimates in Orlicz spaces and applications to variational problems in 2D involving the trace free part of the symmetric gradient”. *Journal of Mathematical Sciences (New York)* 167.3 (2010). Pp. 418–434.
- [26] M. Fuchs and S. Repin. “Some Poincaré-type inequalities for functions of bounded deformation involving the deviatoric part of the symmetric gradient.” *Journal of Mathematical Sciences (New York)* 178.3 (2011). Pp. 367–372.
- [27] M. Fuchs and O. Schirra. “An application of a new coercive inequality to variational problems studied in general relativity and in Cosserat elasticity giving the smoothness of minimizers”. *Archiv der Mathematik* 93.6 (2009). Pp. 587–596.
- [28] A. Garroni, G. Leoni, and M. Ponsiglione. “Gradient theory for plasticity via homogenization of discrete dislocations”. *Journal of the European Mathematical Society* 12.5 (2010). Pp. 1231–1266.
- [29] G. Geymonat and P. Suquet. “Functional spaces for Norton-Hoff materials”. *Mathematical Methods in the Applied Sciences* 8.2 (1986). Pp. 206–222.
- [30] G. Geymonat and F. Krasucki. “Some remarks on the compatibility conditions in elasticity”. *Rendiconti Accademia Nazionale delle Scienze detta dei XL* 29 (2005). Pp. 175–181.
- [31] I.-D. Ghiba, P. Neff, A. Madeo, and I. Münch. “A variant of the linear isotropic indeterminate couple-stress model with symmetric local force-stress, symmetric nonlocal force-stress, symmetric couple-stresses and orthogonal boundary conditions”. *Mathematics and Mechanics of Solids* 22.6 (2017). Pp. 1221–1266.
- [32] M. Holst, J. Kommemi, and G. Nagy. *Rough solutions of the Einstein constraint equations with nonconstant mean curvature*. 2007. arXiv: [0708.3410](https://arxiv.org/abs/0708.3410) [gr-qc].
- [33] J. Jeong and P. Neff. “Existence, uniqueness and stability in linear Cosserat elasticity for weakest curvature conditions”. *Mathematics and Mechanics of Solids* 15.1 (2010). Pp. 78–95.
- [34] J. Jeong, H. Ramézani, I. Münch, and P. Neff. “A numerical study for linear isotropic Cosserat elasticity with conformally invariant curvature”. *Zeitschrift für Angewandte Mathematik und Mechanik* 89.7 (2009). Pp. 552–569.
- [35] E. Kröner. “Die Spannungsfunktionen der dreidimensionalen isotropen Elastizitätstheorie”. German. *Zeitschrift für Physik* 139 (1954). Pp. 175–188, English version: *The stress functions of three-dimensional, isotropic, elasticity theory*, translated by D. H. Delphenich, [available online](#).
- [36] M. Lagally. *Vorlesungen über Vektor-Rechnung*. German. Ed. by E. Hilb. Mathematik und ihre Anwendungen in Monographien und Lehrbüchern. Akademische Verlagsgesellschaft m.B.H., Leipzig, 1928.
- [37] J. Lankeit, P. Neff, and D. Pauly. “Uniqueness of integrable solutions to $\nabla\zeta = G\zeta$, $\zeta|_{\Gamma} = 0$ for integrable tensor coefficients G and applications to elasticity”. *Zeitschrift für angewandte Mathematik und Physik* 64.6 (2013). Pp. 1679–1688.
- [38] K. de Leeuw and H. Mirkil. “A priori estimates for differential operators in L_{∞} norm”. *Illinois Journal of Mathematics* 8 (1964). Pp. 112–124.
- [39] M. Lewicka and S. Müller. “On the optimal constants in Korn’s and geometric rigidity estimates, in bounded and unbounded domains, under Neumann boundary conditions”. *Indiana University Mathematics Journal* 65.2 (2016). Pp. 377–397.
- [40] P. Lewintan, S. Müller, and P. Neff. “Korn inequalities for incompatible tensor fields in three space dimensions with conformally invariant dislocation energy”. *Calculus of Variations and Partial Differential Equations* 60.150 (2021).
- [41] P. Lewintan and P. Neff. “ L^p -trace-free version of the generalized Korn inequality for incompatible tensor fields in arbitrary dimensions”. *Zeitschrift für angewandte Mathematik und Physik* 72.127 (2021).
- [42] P. Lewintan and P. Neff. “ L^p -versions of generalized Korn inequalities for incompatible tensor fields in arbitrary dimensions with p -integrable exterior derivative”. *Comptes Rendus. Mathématique* 359.6 (2021). Pp. 749–755.
- [43] P. Lewintan and P. Neff. “Nečas-Lions lemma revisited: An L^p -version of the generalized Korn inequality for incompatible tensor fields”. *Mathematical Methods in the Applied Sciences* 44 (2021). Pp. 11392–11403.
- [44] S. Li. “On variational symmetry of defect potentials and multiscale configurational force”. *Philosophical Magazine* 88.7 (2008). Pp. 1059–1084.

- [45] F. López-García. “Weighted generalized Korn inequalities on John domains”. *Mathematical Methods in the Applied Sciences* 41.17 (2018). Pp. 8003–8018. ISSN: 0170-4214; 1099-1476/e.
- [46] G. B. Maggiani, R. Scala, and N. Van Goethem. “A compatible-incompatible decomposition of symmetric tensors in L^p with application to elasticity”. *Mathematical Methods in the Applied Sciences* 38.18 (2015). Pp. 5217–5230.
- [47] B. S. Mityagin. “On second mixed derivative”. Russian. *Doklady Akademii Nauk SSSR* 123 (1958). Pp. 606–609.
- [48] S. Müller, L. Scardia, and C. I. Zeppieri. “Geometric rigidity for incompatible fields, and an application to strain-gradient plasticity”. *Indiana University Mathematics Journal* 63.5 (2014). Pp. 1365–1396.
- [49] P. Neff, J. Jeong, and H. Ramezani. “Subgrid interaction and micro-randomness - novel invariance requirements in infinitesimal gradient elasticity.” *Int. J. Solids Struct.* 46.25-26 (2009). Pp. 4261–4276.
- [50] P. Neff. “On Korn’s first inequality with non-constant coefficients”. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics* 132.01 (2002). Pp. 221–243.
- [51] P. Neff, I.-D. Ghiba, M. Lazar, and A. Madeo. “The relaxed linear micromorphic continuum: well-posedness of the static problem and relations to the gauge theory of dislocations”. *The Quarterly Journal of Mechanics and Applied Mathematics* 68.1 (2015). Pp. 53–84.
- [52] P. Neff, I.-D. Ghiba, A. Madeo, L. Placidi, and G. Rosi. “A unifying perspective: the relaxed linear micromorphic continuum”. *Continuum Mechanics and Thermodynamics* 26.5 (2014). Pp. 639–681.
- [53] P. Neff and J. Jeong. “A new paradigm: the linear isotropic Cosserat model with conformally invariant curvature energy”. *Zeitschrift für Angewandte Mathematik und Mechanik* 89.2 (2009). Pp. 107–122.
- [54] P. Neff and I. Münch. “Curl bounds Grad on $SO(3)$ ”. *ESAIM: Control, Optimisation and Calculus of Variations* 14.1 (2008). Pp. 148–159.
- [55] P. Neff, D. Pauly, and K.-J. Witsch. “A canonical extension of Korn’s first inequality to $H(\text{Curl})$ motivated by gradient plasticity with plastic spin”. *Comptes Rendus Mathématique. Académie des Sciences. Paris* 349.23 (2011). Pp. 1251–1254.
- [56] P. Neff, D. Pauly, and K.-J. Witsch. “Maxwell meets Korn: A new coercive inequality for tensor fields in $\mathbb{R}^{n \times n}$ with square-integrable exterior derivative”. *Mathematical Methods in the Applied Sciences* 35.1 (2012). Pp. 65–71.
- [57] P. Neff, D. Pauly, and K.-J. Witsch. “On a canonical extension of Korn’s first and Poincaré’s inequalities to $H(\text{Curl})$ ”. *Journal of Mathematical Sciences* 185.5 (2012). Pp. 721–727.
- [58] P. Neff, D. Pauly, and K.-J. Witsch. “Poincaré meets Korn via Maxwell: Extending Korn’s first inequality to incompatible tensor fields”. *Journal of Differential Equations* 258.4 (2015). Pp. 1267–1302.
- [59] P. Neff and W. Pompe. “Counterexamples in the theory of coerciveness for linear elliptic systems related to generalizations of Korn’s second inequality”. *Zeitschrift für Angewandte Mathematik und Mechanik* 94.9 (2014). Pp. 784–790.
- [60] J. F. Nye. “Some geometrical relations in dislocated crystals”. *Acta Metallurgica* 1 (1953). Pp. 153–162.
- [61] D. Ornstein. “A non-equality for differential operators in the L_1 norm”. *Archive for Rational Mechanics and Analysis* 11 (1962). Pp. 40–49.
- [62] W. Pompe. “Korn’s first inequality with variable coefficients and its generalization”. *Commentationes Mathematicae Universitatis Carolinae* 44.1 (2003). Pp. 57–70.
- [63] Y. G. Reshetnyak. “Estimates for certain differential operators with finite-dimensional kernel”. *Siberian Mathematical Journal* 11.2 (1970). Pp. 315–326.
- [64] Y. G. Reshetnyak. *Stability Theorems in Geometry and Analysis*. Springer Science+Business Media Dordrecht, 1994.
- [65] O. D. Schirra. “New Korn-type inequalities and regularity of solutions to linear elliptic systems and anisotropic variational problems involving the trace-free part of the symmetric gradient”. *Calculus of Variations and Partial Differential Equations* 43.1 (2012). Pp. 147–172.
- [66] T. W. Ting. “St. Venant’s compatibility conditions and basic problems in elasticity”. *Rocky Mountain Journal of Mathematics* 7 (1977). Pp. 47–52.
- [67] W. Wang. “Korn’s inequality and Donati’s theorem for the conformal Killing operator on pseudo-Euclidean space”. *Journal of Mathematical Analysis and Applications* 345.2 (2008). Pp. 777–782.

A Appendix

A.1 On the trace-free Korn’s first inequality in L^2

Using partial integration (see also [58, Appendix A.1]) we catch up with a simple proof of

Lemma A.1. Let $n \geq 2$, $\Omega(\text{open}) \subset \mathbb{R}^n$, $u \in W_0^{1,2}(\Omega, \mathbb{R}^n)$. Then

$$\int_{\Omega} \|Du\|^2 dx \leq 2 \int_{\Omega} \|\text{dev}_n \text{sym } Du\|^2 dx. \quad (\text{A.1})$$

Proof. For $u \in C_c^\infty(\Omega, \mathbb{R}^n)$ we have

$$\begin{aligned} 2 \int_{\Omega} \|\text{sym } Du\|^2 dx &= \int_{\Omega} \|Du\|^2 + \sum_{i,j=1}^n (\partial_i u_j)(\partial_j u_i) dx \stackrel{\text{part.}=\text{int.}}{=} \int_{\Omega} \|Du\|^2 + \sum_{i,j=1}^n (\partial_j u_j)(\partial_i u_i) dx \\ &= \int_{\Omega} \|Du\|^2 + (\text{div } u)^2 dx, \end{aligned} \quad (\text{A.2})$$

from where the "baby" Korn inequality $\int_{\Omega} \|Du\|^2 dx \leq 2 \int_{\Omega} \|\text{sym } Du\|^2 dx$ for $u \in W_0^{1,2}(\Omega, \mathbb{R}^n)$ follows. Its improvement is obtained in regard with the decomposition

$$\|\text{dev}_n \text{sym } Du\|^2 = \|\text{sym } Du - \underbrace{\frac{1}{n} \text{tr}(\text{sym } Du)}_{=\text{div } u} \cdot \mathbf{1}\|^2 = \|\text{sym } Du\|^2 - \frac{1}{n} (\text{div } u)^2, \quad (\text{A.3})$$

since we obtain

$$\begin{aligned} 2 \int_{\Omega} \|\text{dev}_n \text{sym } Du\|^2 dx &\stackrel{(\text{A.3})}{=} 2 \int_{\Omega} \|\text{sym } Du\|^2 dx - \frac{2}{n} \int_{\Omega} (\text{div } u)^2 dx \stackrel{(\text{A.2})}{=} \int_{\Omega} \|Du\|^2 dx + \frac{n-2}{n} \int_{\Omega} (\text{div } u)^2 dx \\ &\stackrel{n \geq 2}{\geq} \int_{\Omega} \|Du\|^2 dx. \quad \blacksquare \end{aligned}$$

Remark A.2. The trace-free Korn's first inequality (A.1) is also valid in L^p , $p > 1$, see [27, Prop. 1] for the $n = 2$ case and [65, Thm. 2.3] for all $n \geq 2$ where again the justification was based on the Lions lemma.

A.2 Infinitesimal planar conformal mappings

Infinitesimal conformal mappings are defined by $\text{dev}_n \text{sym } Du \equiv 0$ and in $n > 2$ they have the representation

$$\langle a, x \rangle x - \frac{1}{2} a \|x\|^2 + Ax + \beta x + c, \quad \text{with } A \in \mathfrak{so}(n), a, c \in \mathbb{R}^n \text{ and } \beta \in \mathbb{R},$$

cf. [63, 49, 33, 17, 64, 65].

In the planar case, the situation is quite different. Indeed, the condition $\text{dev}_2 \text{sym } Du \equiv 0$ reads

$$\begin{aligned} &\begin{pmatrix} u_{1,x} & \frac{1}{2}(u_{1,y} + u_{2,x}) \\ \frac{1}{2}(u_{1,y} + u_{2,x}) & u_{2,y} \end{pmatrix} - \frac{1}{2}(u_{1,x} + u_{2,y}) \cdot \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = 0 \\ \Leftrightarrow &\begin{pmatrix} \frac{1}{2}(u_{1,x} - u_{2,y}) & \frac{1}{2}(u_{1,y} + u_{2,x}) \\ \frac{1}{2}(u_{1,y} + u_{2,x}) & \frac{1}{2}(u_{2,y} - u_{1,x}) \end{pmatrix} = 0 \Leftrightarrow \begin{cases} u_{1,x} = u_{2,y} \\ u_{1,y} = -u_{2,x} \end{cases} \end{aligned}$$

and corresponds to the validity of the Cauchy-Riemann-equations. Thus, in the planar case, infinitesimal conformal mappings are conformal mappings.

A.3 Kröner's relation in infinitesimal elasto-plasticity

At the macroscopic scale, in infinitesimal elasto-plastic theory, see e.g. [amstutz:hal-01789190, 22, 21, 20, 44, 3, 46], the incompatibility of the elastic strain is related to the Curl of the *contortion tensor* $\kappa := \alpha^T - \frac{1}{2} \text{tr}(\alpha) \cdot \mathbf{1}$, where $\alpha := \text{Curl } P$ is the dislocation density tensor, by Kröner's relation [35]:

$$\text{inc}(\text{sym } e) = -\text{Curl } \kappa, \quad (\text{A.4})$$

where the additive decomposition of the displacement gradient into non-symmetric elastic and plastic distortions is assumed:

$$Du = e + P. \quad (\text{A.5})$$

Indeed, (A.4) follows from Nye's formula (1.9) and the identities

$$\text{tr } \text{Curl } \text{sym } e = 0 \quad \text{as well as} \quad \alpha := \text{Curl } P \stackrel{(\text{A.5})}{=} -\text{Curl } e,$$

since we have

$$\begin{aligned} \text{Daxl skew } e &\stackrel{(1.9)_2}{=} \frac{1}{2} \text{tr}(\text{Curl skew } e) \cdot \mathbf{1} - (\text{Curl skew } e)^T \quad \text{tr } \text{Curl } \text{sym } e = 0 \quad \frac{1}{2} \text{tr}(\text{Curl skew } e + \text{Curl } \text{sym } e) \cdot \mathbf{1} - (\text{Curl skew } e)^T \\ &= \frac{1}{2} \text{tr}(\text{Curl } e) \cdot \mathbf{1} - (\text{Curl } e)^T + (\text{Curl } \text{sym } e)^T \stackrel{\alpha = -\text{Curl } e}{=} -\frac{1}{2} \text{tr}(\alpha) \cdot \mathbf{1} + \alpha^T + (\text{Curl } \text{sym } e)^T \\ &= \kappa + (\text{Curl } \text{sym } e)^T. \end{aligned} \quad (\text{A.6})$$

Thus, applying Curl on both sides of (A.6) establishes (A.4), since $\text{Curl} \circ \text{D} \equiv 0$:

$$0 = \text{Curl Daxl skew } e \stackrel{(A.6)}{=} \text{Curl } \kappa + \text{Curl}([\text{Curl sym } e]^T) = \text{Curl } \kappa + \mathbf{inc}(\text{sym } e). \quad (\text{A.7})$$

From the decomposition $\text{sym } \text{D}u = \text{sym } e + \text{sym } P$ it follows moreover $\mathbf{inc}(\text{sym } e) = -\mathbf{inc}(\text{sym } P)$, see also last calculation in footnote 5.

In finite strain elasticity [10], the Riemann-Christoffel tensor \mathcal{R} expresses the compatibility of strain tensors in the sense of

$$C \in C^2(\Omega, \text{Sym}^+(3)) : \mathcal{R}(C) = 0 \Leftrightarrow C = (D\varphi)^T D\varphi \text{ in simply connected domains.} \quad (\text{A.8})$$

Writing $C = (\mathbb{1} + P)^T(\mathbb{1} + P) = \mathbb{1} + 2 \text{sym } P + P^T P$ for $P \in C^2(\Omega, \mathbb{R}^{3 \times 3})$, the incompatibility operator is the linearization of the Riemann-Christoffel tensor at the identity, since

$$\mathcal{R}(\mathbb{1} + 2 \text{sym } P + P^T P) = \mathcal{R}(\mathbb{1}) + 2 \text{D}\mathcal{R}(\mathbb{1}) \text{sym } P + \text{h.o.t.} = 0 + 2 \mathbf{inc}(\text{sym } P) + \text{h.o.t.} \quad (\text{A.9})$$

see also [20] and the references contained therein.

A.4 Further identities

Symmetric tensors play an important role in the above considerations. We mention here the full expression of $S \times b$ for $S \in \text{Sym}(3)$ and $b \in \mathbb{R}^3$:

$$S \times b = \begin{pmatrix} S_{12} b_3 - S_{13} b_2 & S_{13} b_1 - S_{11} b_3 & S_{11} b_2 - S_{12} b_1 \\ S_{22} b_3 - S_{23} b_2 & S_{23} b_1 - S_{12} b_3 & S_{12} b_2 - S_{22} b_1 \\ S_{23} b_3 - S_{33} b_2 & S_{33} b_1 - S_{13} b_3 & S_{13} b_2 - S_{23} b_1 \end{pmatrix} \quad (\text{A.10})$$

which is an example of a trace-free matrix with non-zero entries on the diagonal:

$$\text{tr}(S \times b) = S_{12} b_3 - S_{13} b_2 + S_{23} b_1 - S_{12} b_3 + S_{13} b_2 - S_{23} b_1 = 0.$$

Moreover, we outline some basic identities which played useful roles in our considerations:

<p>1. from linear algebra:</p> <p>(a) $P \times b$ row-wise,</p> <p>(b) $\mathbb{1} \times b = \text{anti}(b) \in \mathfrak{so}(3)$,</p> <p>(c) $(\text{anti } a) \times b = b \otimes a - \langle b, a \rangle \mathbb{1}$,</p> <p>(d) $\text{tr}(S \times b) = 0$,</p> <p>(e) $(\mathbb{1} \times b)^T \times b = \ b\ ^2 \cdot \mathbb{1} - b \otimes b \in \text{Sym}(3)$,</p> <p>(f) $((\text{anti } a) \times b)^T \times b = -\langle b, a \rangle \text{anti}(b) \in \mathfrak{so}(3)$,</p> <p>(g) $(S \times b)^T \times b \in \text{Sym}(3)$,</p> <p>(h) $\text{dev}(P \times b) = P \times b + \frac{2}{3} \langle \text{axl skew } P, b \rangle \cdot \mathbb{1}$,</p> <p>(i) $\text{sym}[(P \times b)^T \times b] = ((\text{sym } P) \times b)^T \times b$,</p> <p>(j) $\text{skew}[(P \times b)^T \times b] = ((\text{skew } P) \times b)^T \times b$,</p> <p>(k) $a \otimes b = 0 \Leftrightarrow \begin{aligned} &\text{sym}(a \otimes b) = 0 \\ &\text{dev}(a \otimes b) = 0 \\ &\text{dev sym}(a \otimes b) = 0, \end{aligned}$</p> <p>(l) $\text{dev}(P \times b) = 0 \Leftrightarrow P \times b = 0$,</p> <p>for $a, b \in \mathbb{R}^3$, $S \in \text{Sym}(3)$ and $P \in \mathbb{R}^{3 \times 3}$,</p>	<p>2. and their formal equivalents from calculus:</p> <p>(a) $\text{Curl } P = P \times (-\nabla)$,</p> <p>(b) $\text{Curl}(\zeta \cdot \mathbb{1}) = -\text{anti}(\nabla \zeta) \in \mathfrak{so}(3)$,</p> <p>(c) $\text{Curl } A = \text{tr}(\text{Daxl } A) \mathbb{1} - (\text{Daxl } A)^T$,</p> <p>(d) $\text{tr}(\text{Curl } S) = 0$,</p> <p>(e) $\mathbf{inc}(\zeta \cdot \mathbb{1}) = \Delta \zeta \cdot \mathbb{1} - \text{D}^2 \zeta \in \text{Sym}(3)$,</p> <p>(f) $\mathbf{inc } A = -\text{anti}(\nabla \text{tr}(\text{Daxl } A)) \in \mathfrak{so}(3)$,</p> <p>(g) $\mathbf{inc } S \in \text{Sym}(3)$,</p> <p>(h) $\text{dev } \text{Curl } P = \text{Curl } P - \frac{2}{3} \text{div axl skew } P \cdot \mathbb{1}$,</p> <p>(i) $\text{sym } \mathbf{inc } P = \mathbf{inc } \text{sym } P$,</p> <p>(j) $\text{skew } \mathbf{inc } P = \mathbf{inc } \text{skew } P$,</p> <p>for $\zeta \in \mathcal{D}'(\Omega, \mathbb{R})$, $A \in \mathcal{D}'(\Omega, \mathfrak{so}(3))$, $S \in \mathcal{D}'(\Omega, \text{Sym}(3))$ and $P \in \mathcal{D}'(\Omega, \mathbb{R}^{3 \times 3})$.</p>
---	---

We catch up with the verification of the identities not contained in our considerations explicitly:

- $(\mathbb{1} \times b)^T \times b \stackrel{1.(a)}{=} (\text{anti}(b))^T \times b = -(\text{anti } b) \times b \stackrel{1.(b)}{=} -b \otimes b + \langle b, b \rangle \mathbb{1} \Rightarrow 1.(d)$,
- we have the decompositions:

$$(P \times b)^T \times b = (\text{sym } P \times b + \text{skew } P \times b)^T \times b = \underbrace{((\text{sym } P) \times b)^T \times b}_{\in \text{Sym}(3)} + \underbrace{((\text{skew } P) \times b)^T \times b}_{\in \mathfrak{so}(3)}$$

but also

$$\mathbf{inc } P = \mathbf{inc}(\text{sym } P + \text{skew } P) = \underbrace{\mathbf{inc } \text{sym } P}_{\in \text{Sym}(3)} + \underbrace{\mathbf{inc } \text{skew } P}_{\in \mathfrak{so}(3)}$$

where we have used (e) and (f), so that (h) and (i) follow,

- the equivalence $a \otimes b = 0 \Leftrightarrow \text{dev sym}(a \otimes b) = 0$ follows from the expression:

$$\frac{\|b\|^4}{2} \|a \otimes b\|^2 = \|b\|^4 \|\text{dev sym}(a \otimes b)\|^2 + \frac{1}{2} \left(\frac{n}{n-1} \right)^2 \langle b, \text{dev sym}(a \otimes b) \rangle^2.$$