



# Measurement of the branching fraction of the $B^0 \rightarrow D_s^+ \pi^-$ decay

LHCb collaboration<sup>†</sup>

## Abstract

A branching fraction measurement of the  $B^0 \rightarrow D_s^+ \pi^-$  decay is presented using proton-proton collision data collected with the LHCb experiment, corresponding to an integrated luminosity of  $5.0 \text{ fb}^{-1}$ . The branching fraction is found to be  $\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-) = (19.4 \pm 1.8 \pm 1.3 \pm 1.2) \times 10^{-6}$ , where the first uncertainty is statistical, the second systematic and the third is due to the uncertainty on the  $B^0 \rightarrow D^- \pi^+$ ,  $D_s^+ \rightarrow K^+ K^- \pi^+$  and  $D^- \rightarrow K^+ \pi^- \pi^-$  branching fractions. This is the most precise single measurement of this quantity to date. As this decay proceeds through a single amplitude involving a  $b \rightarrow u$  charged-current transition, the result provides information on non-factorisable strong interaction effects and the magnitude of the Cabibbo-Kobayashi-Maskawa matrix element  $V_{ub}$ . Additionally, the collision energy dependence of the hadronisation-fraction ratio  $f_s/f_d$  is measured through  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$  and  $B^0 \rightarrow D^- \pi^+$  decays.

Published in Eur. Phys. J. C81 (2021) 314

© 2021 CERN for the benefit of the LHCb collaboration. CC BY 4.0 licence.

<sup>†</sup>Authors are listed at the end of this paper.



# 1 Introduction

To test the Cabibbo-Kobayashi-Maskawa (CKM) sector of the Standard Model (SM), it is crucial to perform accurate measurements of the quark-mixing matrix elements. Any discrepancy among the numerous measurements of CKM matrix elements could reveal effects from new particles or forces beyond the SM. The knowledge of the magnitude of the matrix element  $V_{ub}$  governing the strength of  $b \rightarrow u$  transitions is key in the consistency checks of the SM and its naturally motivated extensions [1, 2].

The hadronic  $B^0 \rightarrow D_s^+ \pi^-$  decay<sup>1</sup> proceeds in the SM through the  $b \rightarrow u$  transition as shown in Fig. 1. Its branching fraction is proportional to  $|V_{ub}|^2$ ,

$$\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-) = \Phi |V_{ub}|^2 |V_{cs}|^2 |F(B^0 \rightarrow \pi^-)|^2 f_{D_s^+}^2 |a_{\text{NF}}|^2, \quad (1)$$

where  $\Phi$  is a phase-space factor,  $F(B^0 \rightarrow \pi^-)$  is a form factor,  $f_{D_s^+}$  is the  $D_s^+$  decay constant,  $V_{cs}$  is the CKM matrix element representing  $c \rightarrow s$  transitions, and  $|a_{\text{NF}}|$  encapsulates non-factorisable effects. The form factor and the decay constant can be obtained from light-cone sum rules [3, 4] and lattice QCD calculations [5, 6], and since  $|V_{cs}|$  is known to be close to unity, the  $B^0 \rightarrow D_s^+ \pi^-$  branching fraction can be used to probe the product  $|V_{ub}| |a_{\text{NF}}|$ . The assumption of factorisation is expected to hold, *i.e.*  $|a_{\text{NF}}|$  is close to unity, for  $B$  meson decays into a heavy and a light meson, where the  $W$  emission of the decay corresponds to the light meson and the spectator quark forms part of the heavy meson. This is not the case for the  $B^0 \rightarrow D_s^+ \pi^-$  decay, as shown in Fig 1, and consequently  $|a_{\text{NF}}|$  may be significantly different from unity [7].

The measurement of the  $B^0 \rightarrow D_s^+ \pi^-$  branching fraction can also be used to estimate the ratio of the amplitudes of the Cabibbo-suppressed  $B^0 \rightarrow D^+ \pi^-$  and the Cabibbo-favoured  $B^0 \rightarrow D^- \pi^+$  decays,

$$r_{D\pi} = \left| \frac{A(B^0 \rightarrow D^+ \pi^-)}{A(B^0 \rightarrow D^- \pi^+)} \right|, \quad (2)$$

which is necessary for the measurement of charge-parity ( $CP$ ) asymmetries in  $B^0 \rightarrow D^\mp \pi^\pm$




Figure 1: Tree diagram of the  $B^0 \rightarrow D_s^+ \pi^-$  decay, in which a  $B^0$  meson decays through the weak interaction to a  $D_s^+$  meson and a charged pion. This diagram represents the only (leading order) process contributing to this decay. Strong interaction between the  $D_s^+$  meson and the pion lead to a non-factorisable contribution to the decay amplitude.

<sup>1</sup>Inclusion of charge-conjugate modes is implied unless explicitly stated.

decays [8–13]. Assuming SU(3) flavour symmetry, Eq. (2) can be written as [14, 15]

$$r_{D\pi} = \tan \theta_c \frac{f_{D^+}}{f_{D_s^+}} \sqrt{\frac{\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-)}{\mathcal{B}(B^0 \rightarrow D^- \pi^+)}} \quad (3)$$

where  $\theta_c$  is the Cabibbo angle and  $f_{D^+}$  is the decay constant of the  $D^+$  meson. SU(3) symmetry breaking is caused by different non-factorisable effects in  $B^0 \rightarrow D_s^+ \pi^-$  and  $B^0 \rightarrow D^- \pi^+$  decays.

This article presents measurements of  $\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-)$  and  $r_{D\pi}$  using proton-proton ( $pp$ ) collision data collected with the LHCb detector at centre-of-mass energies of 7, 8 and 13 TeV corresponding to an integrated luminosity of  $5 \text{ fb}^{-1}$ . The data samples recorded in the years 2011 and 2012 (2015 and 2016) at 7 and 8 (13) TeV will be referred to as Run 1 (Run 2). The  $B^0 \rightarrow D_s^+ \pi^-$  branching ratio is measured relative to the  $B^0 \rightarrow D^- \pi^+$  normalisation channel, which is well measured and experimentally similar to the  $B^0 \rightarrow D_s^+ \pi^-$  decay. The  $B^0 \rightarrow D_s^+ \pi^-$  ( $B^0 \rightarrow D^- \pi^+$ ) candidates are reconstructed via the  $D_s^+ \rightarrow K^+ K^- \pi^+$  ( $D^- \rightarrow K^+ \pi^- \pi^-$ ) decay. The branching fraction of the  $B^0 \rightarrow D_s^+ \pi^-$  decay is determined by

$$\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-) = \mathcal{B}(B^0 \rightarrow D^- \pi^+) \frac{N_{B^0 \rightarrow D_s^+ \pi^-} \epsilon_{B^0 \rightarrow D^- \pi^+}}{N_{B^0 \rightarrow D^- \pi^+} \epsilon_{B^0 \rightarrow D_s^+ \pi^-}} \frac{\mathcal{B}(D^- \rightarrow K^+ \pi^- \pi^-)}{\mathcal{B}(D_s^+ \rightarrow K^+ K^- \pi^+)}, \quad (4)$$

where  $N_X$  denotes the selected candidate yield and  $\epsilon_X$  the related efficiency for the decay mode X. In this measurement, extended maximum-likelihood fits to unbinned invariant mass distributions are performed in order to obtain the yields, while the efficiencies are obtained from simulated events and using calibration data samples.

The relative production of  $B_s^0$  and  $B^0$  mesons, described by the ratio  $f_s/f_d$  where  $f_s$  and  $f_d$  are the  $B_s^0$  and  $B^0$  hadronisation fractions, is shown to slightly depend on the  $pp$  collision energy [16]. The efficiency-corrected yield ratio  $\mathcal{R}$ ,

$$\mathcal{R} \equiv \frac{N_{\bar{B}_s^0 \rightarrow D_s^+ \pi^-} \epsilon_{B^0 \rightarrow D^- \pi^+}}{N_{B^0 \rightarrow D^- \pi^+} \epsilon_{\bar{B}_s^0 \rightarrow D_s^+ \pi^-}} \propto \frac{f_s}{f_d}, \quad (5)$$

is proportional to the relative production ratio and its dependence on the centre-of-mass energy is also reported here. This is measured using  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$  and  $B^0 \rightarrow D^- \pi^+$  decays. Accurate knowledge of  $f_s/f_d$  is a crucial input for every  $B_s^0$  branching fraction measurement, *e.g.*  $\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-)$ , since it dominates in most cases the systematic uncertainty [17]. Following the method described in Ref. [18], the value of  $f_s/f_d$  can be calculated as

$$\frac{f_s}{f_d} = 0.982 \frac{\tau_{B_d}}{\tau_{B_s}} \frac{\mathcal{R}}{\mathcal{N}_a \mathcal{N}_F \mathcal{N}_E} \frac{\mathcal{B}(D^- \rightarrow K^+ \pi^- \pi^-)}{\mathcal{B}(D_s^+ \rightarrow K^+ K^- \pi^+)}, \quad (6)$$

where  $\mathcal{R}$  is defined in Eq. (5), the numerical factor takes phase-space effects into account,  $\mathcal{N}_a$  describes non-factorisable SU(3) breaking effects,  $\mathcal{N}_F$  is the ratio of the form factors,  $\mathcal{N}_E$  takes into account the contribution of the  $W$ -exchange diagram in the  $B^0 \rightarrow D^- \pi^+$  decay, and  $\tau_{B_d}$  ( $\tau_{B_s}$ ) is the  $B^0$  ( $B_s^0$ ) lifetime.

## 2 Detector and simulation

The LHCb detector [19, 20] is a single-arm forward spectrometer covering the pseudorapidity range  $2 < \eta < 5$ , designed for the study of particles containing  $b$  or

$c$  quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the  $pp$  interaction region [21], a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes [22, 23] placed downstream of the magnet. The tracking system provides a measurement of the momentum,  $p$ , of charged particles with a relative uncertainty that varies from about 0.5% below 20 GeV/ $c$  to 1.0% at 200 GeV/ $c$ . The minimum distance of a track to a primary vertex (PV), the impact parameter (IP), is measured with a resolution of  $(15 + 29/p_T)$   $\mu\text{m}$ , where  $p_T$  is the component of the momentum transverse to the beam, in GeV/ $c$ . Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov (RICH) detectors [24]. Hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers [25].

The online event selection is performed by a trigger [26], which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction.

Simulation is required to calculate geometrical, reconstruction and selection efficiencies, and to determine shapes of invariant mass distributions. In the simulation,  $pp$  collisions are generated using PYTHIA [27] with a specific LHCb configuration [28]. Decays of unstable particles are described by EVTGEN [29], in which final-state radiation is generated using PHOTOS [30]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [31] as described in Ref. [32].

### 3 Selection

The  $B^0 \rightarrow D_s^+ \pi^-$  ( $B^0 \rightarrow D^- \pi^+$ ) decays are reconstructed by forming a  $D_s^+ \rightarrow K^+ K^- \pi^+$  ( $D^- \rightarrow K^+ \pi^- \pi^-$ ) candidate and combining it with an additional pion of opposite charge, referred to as the companion. The same reconstruction and selection procedure is applied to the  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$  decay. For the  $B^0 \rightarrow D_s^+ \pi^-$  decay, the invariant mass of the  $K^+ K^-$  pair is required to be within 20 MeV/ $c^2$  of the  $\phi(1020)$  mass to select only the  $D_s^+ \rightarrow \phi(1020) \pi^+$  decays, which significantly improves the signal-to-background ratio compared to other decays with a  $K^+ K^- \pi^+$  combination in the final state. Selecting  $D_s^+ \rightarrow \phi(1020) \pi^+$  decays has an efficiency of about 40%.

At the hardware trigger stage, events are required to have a muon with high  $p_T$  or a hadron, photon or electron with high transverse energy in the calorimeters. For hadrons, the transverse-energy threshold varied between 3 and 4 GeV between 2011 and 2016. The software trigger requires a two-, three- or four-track secondary vertex with significant displacement from any primary  $pp$  interaction vertex (PV). At least one charged particle must have transverse momentum  $p_T > 1.6$  GeV/ $c$  and be inconsistent with originating from a PV. A multivariate algorithm [33] is used for the identification of secondary vertices consistent with the decay of a  $b$  hadron.

After the trigger selection, a preselection is applied to the reconstructed candidates to ensure good quality for the vertex of the  $b$ -hadron and  $c$ -hadron candidates comprising of tracks with large total and transverse momentum. Combinatorial background is suppressed using a gradient boosted decision tree (BDTG) algorithm [34, 35], trained

on Run 1  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$  data. A set of 15 variables is used to train the BDTG classifier, the ones with highest importance in the training being the transverse momentum of the companion pion, the radial flight distance of the  $\bar{B}_s^0$  and of the  $D_s^+$  candidates, the minimum transverse momentum of the  $D_s^+$  decay products and the minimum  $\chi_{\text{IP}}^2$  of the companion and the  $\bar{B}_s^0$  candidates, where  $\chi_{\text{IP}}^2$  is defined as the difference in the vertex-fit  $\chi^2$  of a given PV reconstructed with and without the particle under consideration. The correlation among the input variables has been studied and was found to be small. The BDTG classifier used in this measurement is described in Ref. [36].

To improve the  $B^0$  and  $\bar{B}_s^0$  invariant mass resolutions, the  $D_s^+$  and  $D^-$  invariant masses are constrained to their known values [37]. All  $D_s^+ \pi^-$  ( $D^- \pi^+$ ) candidates are required to have their invariant masses,  $m(D_s^+ \pi^-)$  ( $m(D^- \pi^+)$ ), within the range 5150–5800 (5000–5800) MeV/ $c^2$  and the  $K^+ K^- \pi^+$  ( $K^+ \pi^- \pi^-$ ) invariant mass within 1930–2065 (1830–1920) MeV/ $c^2$ . The range of the  $K^+ K^- \pi^+$  invariant mass includes a large upper sideband to model properly the combinatorial background shape, as described in Sec. 4.

To reduce the background due to misidentified final-state particles, particle identification (PID) information from the RICH detectors is used. The companion pion is required to pass a strict PID requirement to reduce the number of  $\bar{B}_s^0 \rightarrow D_s^+ K^-$  ( $B^0 \rightarrow D^- K^+$ ) decays where the kaon companion is misidentified as a pion. For  $D_s^+ \rightarrow \phi(1020) \pi^+$  candidates, loose PID requirements are applied to both kaons and the pion, which imply a signal efficiency of about 96%. In the case of the pion, the PID requirement is used primarily to remove protons originating from the  $\Lambda_c^+ \rightarrow \phi p$  decay. Further PID requirements are applied to veto  $\Lambda_b^0 \rightarrow \Lambda_c^+ (\rightarrow p K^- \pi^+) \pi^-$  and  $\bar{B}^0 \rightarrow D^+ (\rightarrow K^- \pi^+ \pi^+) \pi^-$  and  $\bar{A}_b^0 \rightarrow \bar{\Lambda}_c^- (\rightarrow \bar{p} K^+ \pi^-) \pi^+$  and  $B_s^0 \rightarrow D_s^- (\rightarrow K^- K^+ \pi^-) \pi^+$  events, which are misidentified as the final-state particles of  $D_s^+ (\rightarrow K^+ K^- \pi^+) \pi^-$  and  $D^- (\rightarrow K^+ \pi^- \pi^-) \pi^+$  decays, respectively. These vetoes are applied if candidates are consistent with the above mentioned decays when a mass hypothesis is changed. The PID requirements result in 75% efficiency for  $B^0 \rightarrow D_s^+ \pi^-$  signal decays, which is dominated by the strict PID requirement on the companion pion, while the retention is about 9% for the  $\bar{B}_s^0 \rightarrow D_s^+ K^-$  misidentified background contribution.

The event selection efficiencies are calculated from simulation with the exception of the efficiency of the PID requirements which is determined using calibration data samples.

## 4 Signal and background parametrisation

After the full event selection, unbinned maximum-likelihood fits are performed to obtain the yields of the signal  $B^0 \rightarrow D_s^+ \pi^-$  and the normalisation  $B^0 \rightarrow D^- \pi^+$  candidates. A two-dimensional fit to the  $D_s^+ \pi^-$  and the  $K^+ K^- \pi^+$  invariant mass distributions is performed to determine the  $B^0 \rightarrow D_s^+ \pi^-$  signal yield, while the yield of the normalisation channel is obtained from a fit to the  $D^- \pi^+$  invariant mass distribution. Due to the  $D_s^+$  mass constraint, the correlation between  $m(D_s^+ \pi^-)$  and  $m(K^+ K^- \pi^+)$  is found to be small, thus the two variables are factorised in the fit model [38]. The two-dimensional fit is performed in order to constrain the combinatorial background (see further in this Section for details).

The  $B^0 \rightarrow D_s^+ \pi^-$  decay is Cabibbo-suppressed and is therefore considerably less abundant than the Cabibbo-favoured  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$  decay, which produces the same final-

state particles. The  $m(D_s^+\pi^-)$  and  $m(D^-\pi^+)$  shapes for  $\bar{B}_s^0 \rightarrow D_s^+\pi^-$  and  $B^0 \rightarrow D^-\pi^+$  candidates, respectively, are described by the sum of a double-sided Hypatia function [39] and a Johnson  $S_U$  function [40]. The left tail of the  $\bar{B}_s^0 \rightarrow D_s^+\pi^-$  invariant mass distribution overlaps with the  $B^0 \rightarrow D_s^+\pi^-$  signal peak and therefore special attention is given to the description of the lower mass range of the  $\bar{B}_s^0 \rightarrow D_s^+\pi^-$  peak, shaped by the combination of detector resolution and radiative effects. The  $B^0 \rightarrow D_s^+\pi^-$  signal is described with the same model as the  $\bar{B}_s^0 \rightarrow D_s^+\pi^-$  decay, shifted by the known  $B^0-B_s^0$  mass difference [37]. The left tail of this distribution is described by two parameters,  $a_1$  and  $n_1$ , which are found to be correlated and therefore the parameter  $n_1$  is fixed to the value obtained from simulation, whereas  $a_1$  is obtained from simulated  $\bar{B}_s^0 \rightarrow D_s^+\pi^-$  and  $B^0 \rightarrow D^-\pi^+$  events, as well as from  $B^0 \rightarrow D^-\pi^+$  data. In the invariant mass fit to  $B^0 \rightarrow D^-\pi^+$  candidates the common mean of the double-sided Hypatia and the Johnson  $S_U$  functions, the widths and the left-tail parameter  $a_1$  are left free in the fit, while this parameter is constrained in the  $D_s^+\pi^-$  invariant mass distribution, as the background does not allow to determine the shape of the radiative tail reliably. All other parameters are fixed from simulation. In the  $K^+K^-\pi^+$  invariant mass fit a sum of two Crystal Ball functions with a common mean is used. The common mean and a scale factor for the widths are left free, while the other shape parameters are fixed from simulation.

The combinatorial background in  $B^0 \rightarrow D_s^+\pi^-$  candidates is split in two components, referred to as random- $D_s^+$  and true- $D_s^+$ . The random- $D_s^+$  combinatorial background consists of random combinations of tracks that do not peak in the  $K^+K^-\pi^+$  invariant mass, while the true- $D_s^+$  combinatorial background consists of events with a true  $D_s^+$  meson, combined with a random companion track. The upper mass range of the  $K^+K^-\pi^+$  candidate sample is used to account accurately for the random- $D_s^+$  component, modelled with a single exponential distribution, while the true- $D_s^+$  background is described by the signal shape. In the  $D_s^+\pi^-$  invariant mass fit, the random- $D_s^+$  background is described by an exponential distribution and the true- $D_s^+$  background is described by the sum of an exponential and a constant function. The exponential parameters are left free in both invariant mass fits.

The combinatorial background in the  $m(D^-\pi^+)$  fit of the normalisation channel is described by the sum of an exponential and a constant function, with the relative weight of the two functions and exponential parameter left free.

Decays where one or more final-state particles are not reconstructed are referred to as partially reconstructed backgrounds. In the  $D_s^+\pi^-$  and  $D^-\pi^+$  invariant mass fits these background contributions are described by an upward-open parabola or a parabola exhibiting a maximum, whose ranges are defined by the kinematic endpoints of the decay, which are convolved with Gaussian resolution functions, and which are known to describe decays involving a missing neutral pion or a missing photon, as defined in Ref. [41]. In the fit to the  $K^+K^-\pi^+$  invariant mass, the partially reconstructed background contributions are described by the signal mass shape.

The  $m(D_s^+\pi^-)$  fit requires two partially reconstructed background components from  $\bar{B}_s^0 \rightarrow D_s^{*+}(\rightarrow D_s^+\gamma/\pi^0)\pi^-$  and  $\bar{B}_s^0 \rightarrow D_s^+\rho^-(\rightarrow \pi^-\pi^0)$  decays. The fit model describing the  $D^-\pi^+$  invariant mass accounts analogously for two partially reconstructed background contributions:  $B^0 \rightarrow D^{*-}(\rightarrow D^-\pi^0)\pi^+$  and  $B^0 \rightarrow D^-\rho^+(\rightarrow \pi^+\pi^0)$ . In the case of the  $\bar{B}_s^0 \rightarrow D_s^{*+}\pi^-$  background the previously mentioned upward-open parabola together with a parabola exhibiting a maximum is used to parameterise the components with  $D_s^{*+} \rightarrow D_s^+\gamma$  and  $D_s^{*+} \rightarrow D_s^+\pi^0$  decays, respectively. The  $\bar{B}_s^0 \rightarrow D_s^+\rho^-$  background is described by the

upward-open parabola, to take into account the missing neutral pion. The  $B^0 \rightarrow D^{*-}\pi^+$  decay uses an upward-open parabola function and exhibits a double-peaked shape. Most parameters are obtained from simulated events and fixed, aside from the relevant invariant mass shifts and widths. For the  $B^0 \rightarrow D^-\rho^+$  background a single upward-open parabola function is taken, with a floating width and a floating mass shift parameter that is shared with the  $B^0 \rightarrow D^{*-}\pi^+$  contribution. The widths of the partially reconstructed background contributions in the  $m(D_s^+\pi^-)$  fits are fixed to the values obtained from  $B^0 \rightarrow D^-\pi^+$  candidates in data, corrected for differences between the  $m(D_s^+\pi^-)$  and  $m(D^-\pi^+)$  distributions, as obtained from simulation.

The  $B^0 \rightarrow D^-\pi^+$  candidate sample is contaminated by the  $B_s^0 \rightarrow D_s^-\pi^+$ ,  $\bar{A}_b^0 \rightarrow \bar{A}_c^-\pi^+$  and  $B^0 \rightarrow D^-K^+$  decays, resulting from the misidentification of one or two of the final-state particles. Analogously, the  $\bar{B}_s^0 \rightarrow D_s^+K^-$ ,  $\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-$  and  $\bar{B}^0 \rightarrow D^+\pi^-$  decays are misidentified background contributions of the  $B^0 \rightarrow D_s^+\pi^-$  candidate sample. Their shapes are determined from simulation using a non-parametric kernel estimation method [42]. The yields of the misidentified background contributions are estimated by using known branching fractions [37] and efficiencies that are determined from simulated background decays. Each yield of a misidentified background in the fit model is constrained to be close to its estimated value and is allowed to vary within the corresponding uncertainty.

## 5 Signal yields

The  $m(D^-\pi^+)$  data distributions, with overlaid fit projections for the total, the  $B^0 \rightarrow D^-\pi^+$  signal and the background components, are shown in Fig. 2. The resulting signal yields are  $(4.971 \pm 0.013) \times 10^5$  and  $(6.294 \pm 0.016) \times 10^5$  for Run 1 and Run 2 samples, respectively. The fit results are also used to constrain the left tail of the signal shape and the widths of the partially reconstructed backgrounds to the invariant mass distribution of  $B^0 \rightarrow D_s^+\pi^-$  candidates.

The two-dimensional fit to  $B^0 \rightarrow D_s^+\pi^-$  candidates is performed in the  $D_s^+\pi^-$  and  $K^+K^-\pi^+$  invariant mass distributions. The  $B^0 \rightarrow D_s^+\pi^-$  branching fraction is determined using the yields of the signal and normalisation modes, their selection efficiencies and the known  $B^0 \rightarrow D^-\pi^+$ ,  $D^- \rightarrow K^+\pi^-\pi^-$  and  $D_s^+ \rightarrow K^+K^-\pi^+$  branching fractions [37]. The two-dimensional fit is performed simultaneously for Run 1 and Run 2 data samples in which the  $\mathcal{B}(B^0 \rightarrow D_s^+\pi^-)$  and left-tail parameter are shared. The fit results in  $B^0 \rightarrow D_s^+\pi^-$  signal yields of  $(8.9 \pm 0.8) \times 10^2$  and  $(1.12 \pm 0.11) \times 10^3$  and  $\bar{B}_s^0 \rightarrow D_s^+\pi^-$  yields of  $(3.370 \pm 0.023) \times 10^4$  and  $(4.647 \pm 0.027) \times 10^4$  for Run 1 and Run 2 samples, respectively. Figure 3 shows the  $D_s^+\pi^-$  invariant mass distributions together with the fit projections and background contributions overlaid. Additionally, the invariant mass fits to  $B^0 \rightarrow D^-\pi^+$  and  $B^0 \rightarrow D_s^+\pi^-$  candidates are performed simultaneously to 2011, 2012 and Run 2 data in order to study the collision energy dependence of  $f_s/f_d$ , as is described in Sec. 7.

## 6 Systematic uncertainties

Systematic uncertainties on the  $\mathcal{B}(B^0 \rightarrow D_s^+\pi^-)$  measurement arise from choices in the fit model and the determination of trigger, BDT and PID efficiencies. Many possible






Figure 2: The invariant mass distributions of normalisation  $B^0 \rightarrow D^- \pi^+$  candidates, for (left) Run 1 and (right) Run 2 data samples. Overlaid are the fit projections along with the signal and background contributions.




Figure 3: The (top)  $D_s^+ \pi^-$  and (bottom)  $K^+ K^- \pi^+$  invariant mass distributions of signal  $B^0 \rightarrow D_s^+ \pi^-$  candidates, for (left) Run 1 and (right) Run 2 data samples. Overlaid are the fit projections along with the signal and background contributions.

sources of systematic uncertainty cancel in the ratio of either the yields or the efficiencies of  $B^0 \rightarrow D_s^+ \pi^-$  and  $B^0 \rightarrow D^- \pi^+$  events. A summary of all the systematic uncertainties is shown in Table 1. The precision of the measurement relies mostly on the accurate modelling of the signal shape and of the partially reconstructed backgrounds.

The most critical aspect of the signal shape is the description of the left tail of the  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$  signal, affecting the composition of signal and background around the  $B^0$  mass. The shape of the left tail was determined from  $B^0 \rightarrow D^- \pi^+$  candidates, taking into account differences between the final states, as obtained from simulation, and was Gaussian constrained in the fit. A systematic uncertainty is assigned for the assumption of the signal shape. This is done by repeating the signal fit with a different parametrisation, *i.e.* the sum of a double-sided Hypatia function and a Gaussian function, which leads to a systematic uncertainty of 5.1%. This parametrisation was found to be the only alternative parametrisation that satisfactorily described simulated signal candidates. Furthermore, a systematic uncertainty is assigned by fixing the mean of the  $B^0 \rightarrow D_s^+ \pi^-$  signal shape to the result of the  $B^0 \rightarrow D^- \pi^+$  fit, rather than shifting by the known  $B^0 - B_s^0$  mass difference. Moreover, the width of the  $B^0 \rightarrow D_s^+ \pi^-$  signal shape is scaled by the ratio of the known  $B^0$  and  $B_s^0$  masses. The widths of the partially reconstructed backgrounds is varied by  $\pm 1 \text{ MeV}/c^2$ , in order to cover the differences between data and simulation as well as the differences between the  $D_s^+ \pi^-$  and  $D^- \pi^+$  invariant mass distributions. The resulting difference between the signal yields is assigned as a systematic uncertainty.

The simulated samples are corrected for an imperfect modelling of the response of the particle identification algorithms as a function of the kinematical properties of the particle, using samples of  $D^{*+}$  calibration data. A systematic uncertainty associated with the PID efficiency evaluation is assigned by varying the corrections within their uncertainties. Proton misidentification is the most difficult to control accurately from data calibration samples, as relatively little calibration data is available in the kinematic region that overlaps with the  $B$  decay products. In addition, the Cherenkov angles of photons emitted by protons and kaons are more similar than those of kaons and pions. Thus, a systematic uncertainty is estimated from the difference between the nominal signal yields and a fit where the misidentified background  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$  decay yield is left free to vary.

The systematic uncertainty assigned to the hardware trigger efficiency takes into account a difference in detection efficiency between kaons and pions. This mostly cancels in the ratio of  $B^0 \rightarrow D^- \pi^+$  and  $B^0 \rightarrow D_s^+ \pi^-$  efficiencies, but the difference of one final-state particle is sensitive to this detection asymmetry. Moreover, an uncertainty related to the reconstruction efficiency of charged particles is taken into account, which mainly arises from the uncertainty on the LHCb material and the different interaction cross-section of pions and kaons with the material [43]. Additionally, a systematic uncertainty is determined on the BDT efficiency due to the difference between simulation and data. This is determined by weighting all the BDT input variables in the simulated signal sample to the signal distributions in data, which are obtained using signal weights for each candidate using the *sPlot* technique [44].

The systematic uncertainties on the collision energy dependence of the efficiency-corrected  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$  and  $B^0 \rightarrow D^- \pi^+$  yield ratios are shown in Table 2. The sources of these systematic uncertainties are the same as for the  $B^0 \rightarrow D_s^+ \pi^-$  branching fraction. Exceptions are the uncertainties on the  $B^0 \rightarrow D_s^+ \pi^-$  signal and the partially reconstructed backgrounds, which are found to be negligible, and the uncertainty on the charged-particle reconstruction efficiency, which cancels out in the double ratio of efficiencies.

Table 1: Relative systematic uncertainty  $\sigma$  on the  $B^0 \rightarrow D_s^+ \pi^-$  branching fraction measurement.

Source	$\sigma(\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-))$ [%]
Fit model	
Signal shape parametrisation	5.1
$B^0 \rightarrow D_s^+ \pi^-$ signal width	1.5
$B^0 \rightarrow D_s^+ \pi^-$ mean	0.2
Partially reconstructed backgrounds	4.2
Misidentified backgrounds	0.6
Efficiencies	
Hardware trigger efficiency	0.3
Reconstruction efficiency	0.5
BDT efficiency	0.7
PID efficiency	1.1
Total	6.9

Table 2: Relative systematic uncertainty  $\sigma$  on the ratio of the efficiency-corrected  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$  and  $B^0 \rightarrow D^- \pi^+$  yield ratios. The ratios  $\mathcal{R}_{13\text{ TeV}}/\mathcal{R}_{7\text{ TeV}}$  and  $\mathcal{R}_{13\text{ TeV}}/\mathcal{R}_{8\text{ TeV}}$  are reported together as the difference of the systematic uncertainty for 7 and 8 TeV is negligible.

Source	$\sigma\left(\frac{\mathcal{R}_{13\text{ TeV}}}{\mathcal{R}_{7,8\text{ TeV}}}\right)$ [%]	$\sigma\left(\frac{\mathcal{R}_{8\text{ TeV}}}{\mathcal{R}_{7\text{ TeV}}}\right)$ [%]
Fit model		
Signal shape parametrisation	0.2	–
Misidentified backgrounds	0.2	–
Efficiencies		
Hardware trigger efficiency	0.4	0.4
BDT efficiency	1.1	1.3
PID efficiency	1.4	1.4
Total	1.9	2.0

## 7 Results

Table 3 gathers all measurements and inputs to determine the branching fraction according to Eq. (4). The branching fraction ratio of  $B^0 \rightarrow D_s^+ \pi^-$  and  $B^0 \rightarrow D^- \pi^+$  decays is found to be

$$\frac{\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-)}{\mathcal{B}(B^0 \rightarrow D^- \pi^+)} = (7.7 \pm 0.7 \pm 0.5 \pm 0.3) \times 10^{-3},$$

where the first uncertainty is statistical, the second systematic and the third stems from knowledge of the  $D^- \rightarrow K^+ \pi^- \pi^-$  and  $D_s^- \rightarrow K^- K^+ \pi^-$  branching fractions.

Using the known value of  $\mathcal{B}(B^0 \rightarrow D^- \pi^+)$  [37], the  $B^0 \rightarrow D_s^+ \pi^-$  branching fraction is found to be

$$\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-) = (19.4 \pm 1.8 \pm 1.3 \pm 1.2) \times 10^{-6},$$

where the first uncertainty is statistical, the second systematic and the third refers to the uncertainty due to the branching fractions listed in Table 3. This result represents the most precise single measurement of  $\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-)$  to date.

Table 3: Results of  $B^0 \rightarrow D_s^+ \pi^-$  and  $B^0 \rightarrow D^- \pi^+$  signal efficiencies and yields, as well as the branching fractions used as input for this measurement [37].

	Run 1	Run 2
$\epsilon_{B^0 \rightarrow D_s^+ \pi^-}$ (%)	$0.1412 \pm 0.0010$	$0.1922 \pm 0.0012$
$\epsilon_{B^0 \rightarrow D^- \pi^+}$ (%)	$0.3485 \pm 0.0016$	$0.4536 \pm 0.0016$
$N_{B^0 \rightarrow D^- \pi^+}$	$(4.971 \pm 0.013) \times 10^5$	$(6.294 \pm 0.016) \times 10^5$
$N_{B^0 \rightarrow D_s^+ \pi^-}$	$(8.9 \pm 0.8) \times 10^2$	$(1.12 \pm 0.11) \times 10^3$
$\mathcal{B}(B^0 \rightarrow D^- \pi^+)$	$(2.52 \pm 0.13) \times 10^{-3}$	
$\mathcal{B}(D^- \rightarrow K^+ \pi^- \pi^-)$	$(9.38 \pm 0.16) \times 10^{-2}$	
$\mathcal{B}(D_s^+ \rightarrow K^+ K^- \pi^+)$	$(5.39 \pm 0.15) \times 10^{-2}$	

The  $B^0 \rightarrow D_s^+ \pi^-$  branching fraction depends on both  $|a_{\text{NF}}|$  and  $|V_{ub}|$ . Using the measurement of  $\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-)$ , the product

$$|V_{ub}| |a_{\text{NF}}| = (3.14 \pm 0.20 \pm 0.25) \times 10^{-3}$$

is obtained, where the first uncertainty is from the  $B^0 \rightarrow D_s^+ \pi^-$  branching fraction measurement and the second from the CKM and QCD parameters. The form factor  $F(B^0 \rightarrow \pi^-)|_{q^2=m_{D_s^+}^2} = 0.327 \pm 0.025$  is obtained using light-cone sum rules [3, 4] and lattice QCD calculations are used for the decay constant  $f_{D_s^+} = 0.2499 \pm 0.0005$  GeV [5, 6]. A phase-space factor  $\Phi = 296.2 \pm 0.8$  GeV $^{-2}$  is used in order to relate the branching fraction to  $|V_{ub}| |a_{\text{NF}}|$ . Additionally, the CKM matrix element  $|V_{cs}|$  is well measured and used as an input [37]. The determination of  $|V_{ub}| |a_{\text{NF}}|$  can be compared to the known inclusive and exclusive determinations of  $|V_{ub}|$  to provide a constraint on the  $|a_{\text{NF}}|$  parameter as displayed in Fig. 4.

The branching fraction ratio of  $B^0 \rightarrow D_s^+ \pi^-$  and  $B^0 \rightarrow D^- \pi^+$  decays can be used to determine the parameter  $r_{D\pi}$ , as shown in Eq. (3). Inserting the measured branching fraction ratio  $\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-)/\mathcal{B}(B^0 \rightarrow D^- \pi^+)$ , the tangent of  $\theta_c$  [37] and the fraction between the decay constants  $f_{D_s^+}$  and  $f_{D^+}$  [5, 6] into Eq. (3) gives

$$r_{D\pi} = 0.0163 \pm 0.0007 \pm 0.0007 \pm 0.0033,$$

where the first uncertainty is statistical, the second systematic and the third arises from possible non-factorisable SU(3)-breaking effects, estimated to be 20% according to Ref. [12]. SU(3)-breaking effects of about 20% are consistent with the measured  $|a_{\text{NF}}|$  in this analysis, see Fig. 4.

Finally, the potential dependence of the hadronisation fraction  $f_s/f_d$  on collision energy is probed using the  $B^0 \rightarrow D^- \pi^+$  and  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$  signal yields obtained in the invariant mass fits, using Eq. (5). To determine these, the fit to Run 1 data is split based on collision energy into 2011 (7 TeV) and 2012 (8 TeV), sharing the shape parameters. The measured double ratios for the different collision energies are

$$\mathcal{R}_{13 \text{ TeV}}/\mathcal{R}_{7 \text{ TeV}} = 1.020 \pm 0.013 \pm 0.021,$$

$$\mathcal{R}_{13 \text{ TeV}}/\mathcal{R}_{8 \text{ TeV}} = 1.035 \pm 0.011 \pm 0.021,$$

$$\mathcal{R}_{8 \text{ TeV}}/\mathcal{R}_{7 \text{ TeV}} = 0.986 \pm 0.013 \pm 0.021,$$




Figure 4: Result of the determination of  $|V_{ub}||a_{\text{NF}}|$ . The blue line represents the result of this measurement, the vertical bands are the known exclusive and inclusive measurements of  $|V_{ub}|$ , which are  $(3.70 \pm 0.16) \times 10^{-3}$  and  $(4.49 \pm 0.28) \times 10^{-3}$ , respectively [37]. The horizontal dashed line at  $|a_{\text{NF}}| = 1.0$  represents exact factorisation. The error bands represent an uncertainty of one standard deviation.

where the first uncertainty is statistical and the second systematic. The average transverse momentum of the  $B$  meson after full event selection is found to be 10.4, 10.6 and 10.9 GeV/ $c$  for  $pp$  collision centre-of-mass energies of 7 TeV, 8 TeV and 13 TeV, respectively. The separate values of  $\mathcal{R}$  at the three collision energies are

$$\begin{aligned}\mathcal{R}_{7\text{ TeV}} &= 0.1631 \pm 0.0018 \pm 0.0025 \pm 0.0014, \\ \mathcal{R}_{8\text{ TeV}} &= 0.1609 \pm 0.0013 \pm 0.0024 \pm 0.0014, \\ \mathcal{R}_{13\text{ TeV}} &= 0.1665 \pm 0.0011 \pm 0.0023 \pm 0.0012,\end{aligned}$$

where the first uncertainty is statistical and the following are the uncorrelated and correlated systematic uncertainties, respectively. The value of  $\mathcal{R}$  at 7 TeV shows good agreement with the previous hadronic  $f_s/f_d$  measurement at 7 TeV, which was performed using  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$ ,  $B^0 \rightarrow D^- \pi^+$  and  $B^0 \rightarrow D^- K^+$  decays [45]. A visualisation of the dependence of  $\mathcal{R}$  on the centre-of-mass energy is given in Fig. 5. The resulting centre-of-mass energy dependence is obtained from a linear fit using the statistical and uncorrelated systematic uncertainties and is found to be  $\mathcal{R} = 0.156(6) + 0.0008(6)\sqrt{s}$ , where  $\sqrt{s}$  is in TeV. The observed trend is in agreement with the LHCb measurement of the  $f_s/f_u$  dependence upon the  $pp$  collision energy [16]. The values for  $\mathcal{R}$  will be used in a future work and can be used to obtain  $f_s/f_d$  by correcting  $\mathcal{R}$  for the relative  $D$  branching fractions, the ratio of  $B$  lifetimes, the form factor ratio, the contribution from non-factorisable SU(3)-breaking effects and the contribution from the exchange diagram, as given by Eq. (6).




Figure 5: Visualisation of the  $pp$  collision energy dependence of the efficiency-corrected yield ratio of  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$  and  $B^0 \rightarrow D^- \pi^+$  decays, which scales with  $f_s/f_d$ . The inner error bars indicate the statistical uncertainty only, whereas the outer indicate the uncorrelated, including statistical, uncertainties. The correlated systematic uncertainty is not shown. The red dotted line represents a linear fit through the three values of  $\mathcal{R}$  with uncorrelated, including statistical, uncertainties.

## 8 Summary

A branching fraction measurement of the  $B^0 \rightarrow D_s^+ \pi^-$  decay is performed using  $pp$  collision data taken between 2011 and 2016, leading to

$$\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-) = (19.4 \pm 1.8 \pm 1.3 \pm 1.2) \times 10^{-6},$$

where the first uncertainty is statistical, the second systematic and the third is due to the branching fractions used as normalisation inputs. This is the most precise single measurement of  $\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-)$  to date, and is in agreement with the current world average [37]. Using this branching fraction, the product of  $|V_{ub}|$  and the non-factorisation constant  $|a_{\text{NF}}|$  is determined to be

$$|V_{ub}| |a_{\text{NF}}| = (3.14 \pm 0.20 \pm 0.25) \times 10^{-3}.$$

Comparison with independently measured values of  $V_{ub}$  [37] indicate that  $|a_{\text{NF}}|$  may deviate from unity by around 20%, indicating significant non-factorisable corrections.

The measurement of the ratio of the  $B^0 \rightarrow D_s^+ \pi^-$  and  $B^0 \rightarrow D^- \pi^+$  branching fractions is used to determine the  $r_{D\pi}$  parameter,

$$r_{D\pi} = 0.0163 \pm 0.0007 \pm 0.0007 \pm 0.0033,$$

where the first uncertainty is statistical, the second systematic and the third arises from possible non-factorisable SU(3)-breaking effects, estimated to be 20% [12]. Knowledge of this parameter is essential to interpret the  $CP$  asymmetries in  $B^0 \rightarrow D^\mp \pi^\pm$  decays.

Finally, the efficiency-corrected yield ratio of  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$  and  $B^0 \rightarrow D^- \pi^+$  decays,  $\mathcal{R}$ , is used to probe the collision energy dependence of the hadronisation fraction  $f_s/f_d$ .

## Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MSHE (Russia); MICINN (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); A\*MIDEX, ANR, Labex P2IO and OCEVU, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, Thousand Talents Program, and Sci. & Tech. Program of Guangzhou (China); RFBR, RSF and Yandex LLC (Russia); GVA, XuntaGal and GENCAT (Spain); the Royal Society and the Leverhulme Trust (United Kingdom).

## References

- [1] CKMfitter group, J. Charles *et al.*, *Current status of the standard model CKM fit and constraints on  $\Delta F = 2$  new physics*, Phys. Rev. **D91** (2015) 073007, [arXiv:1501.05013](https://arxiv.org/abs/1501.05013), updated results and plots available at <http://ckmfitter.in2p3.fr/>.
- [2] UTfit collaboration, M. Bona *et al.*, *The unitarity triangle fit in the standard model and hadronic parameters from lattice QCD: A reappraisal after the measurements of  $\Delta m_s$  and  $BR(B \rightarrow \tau\nu_\tau)$* , JHEP **10** (2006) 081, [arXiv:hep-ph/0606167](https://arxiv.org/abs/hep-ph/0606167), updated results and plots available at <http://www.utfit.org/>.
- [3] P. Ball and R. Zwicky, *New results on  $B \rightarrow \pi, K, \eta$  decay form factors from light-cone sum rules*, Phys. Rev. **D71** (2005) 014015, [arXiv:hep-ph/0406232](https://arxiv.org/abs/hep-ph/0406232).
- [4] P. Ball,  *$|V_{ub}|$  from UTangles and  $B^0 \rightarrow \pi^- \ell^+ \nu$* , Phys. Lett. **B644** (2007) 38, [arXiv:hep-ph/0611108](https://arxiv.org/abs/hep-ph/0611108).
- [5] A. Bazavov *et al.*,  *$B$ - and  $D$ -meson leptonic decay constants from four-flavor lattice QCD*, Phys. Rev. **D98** (2018) 074512, [arXiv:1712.09262](https://arxiv.org/abs/1712.09262).
- [6] N. Carrasco *et al.*, *Leptonic decay constants  $f_K, f_D$ , and  $f_{D_s}$  with  $N_f = 2 + 1 + 1$  twisted-mass lattice QCD*, Phys. Rev. **D91** (2015) 054507, [arXiv:1411.7908](https://arxiv.org/abs/1411.7908).
- [7] M. Beneke, G. Buchalla, M. Neubert, and C. T. Sachrajda, *QCD factorization for exclusive, nonleptonic  $B$  meson decays: General arguments and the case of heavy light final states*, Nucl. Phys. **B591** (2000) 313, [arXiv:hep-ph/0006124](https://arxiv.org/abs/hep-ph/0006124).

- [8] BaBar collaboration, B. Aubert *et al.*, *Measurement of time-dependent CP-violating asymmetries and constraints on  $\sin(2\beta + \gamma)$  with partial reconstruction of  $B \rightarrow D^{*\mp}\pi^\pm$  decays*, Phys. Rev. **D71** (2005) 112003, [arXiv:hep-ex/0504035](#).
- [9] BaBar collaboration, B. Aubert *et al.*, *Measurement of time-dependent CP asymmetries in  $B^0 \rightarrow D^{(*)\pm}\pi^\mp$  and  $B^0 \rightarrow D^\pm\rho^\mp$  decays*, Phys. Rev. **D73** (2006) 111101, [arXiv:hep-ex/0602049](#).
- [10] Belle collaboration, F. J. Ronga *et al.*, *Measurements of CP violation in  $B^0 \rightarrow D^{*-}\pi^+$  and  $B^0 \rightarrow D^-\pi^+$  decays*, Phys. Rev. **D73** (2006) 092003, [arXiv:hep-ex/0604013](#).
- [11] Belle collaboration, S. Bahinipati *et al.*, *Measurements of time-dependent CP asymmetries in  $B \rightarrow D^{*\mp}\pi^\pm$  decays using a partial reconstruction technique*, Phys. Rev. **D84** (2011) 021101, [arXiv:1102.0888](#).
- [12] K. De Bruyn *et al.*, *Exploring  $B_s \rightarrow D_s^{(*)\pm}K^\mp$  decays in the presence of a sizable width difference  $\Delta\Gamma_s$* , Nucl. Phys. **B868** (2013) 351, [arXiv:1208.6463](#).
- [13] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP violation in  $B^0 \rightarrow D^\pm\pi^\mp$  decays*, JHEP **06** (2018) 084, [arXiv:1805.03448](#).
- [14] BaBar collaboration, B. Aubert *et al.*, *Measurement of the branching fractions of the rare decays  $B^0 \rightarrow D_s^{(*)+}\pi^-$ ,  $B^0 \rightarrow D_s^{(*)+}\rho^-$ , and  $B^0 \rightarrow D_s^{(*)-}K^{(*)+}$* , Phys. Rev. **D78** (2008) 032005, [arXiv:0803.4296](#).
- [15] Belle collaboration, A. Das *et al.*, *Measurements of branching fractions for  $B^0 \rightarrow D_s^+\pi^-$  and  $\bar{B}^0 \rightarrow D_s^+K^-$* , Phys. Rev. **D82** (2010) 051103, [arXiv:1007.4619](#).
- [16] LHCb collaboration, R. Aaij *et al.*, *Measurement of  $f_s/f_u$  variation with proton-proton collision energy and B-meson kinematics*, Phys. Rev. Lett. **124** (2020) 122002, [arXiv:1910.09934](#).
- [17] LHCb collaboration, R. Aaij *et al.*, *Measurement of the  $B_s^0 \rightarrow \mu^+\mu^-$  branching fraction and effective lifetime and search for  $B^0 \rightarrow \mu^+\mu^-$  decays*, Phys. Rev. Lett. **118** (2017) 191801, [arXiv:1703.05747](#).
- [18] R. Fleischer, N. Serra, and N. Tuning, *Tests of factorization and  $SU(3)$  relations in B decays into heavy-light final states*, Phys. Rev. **D83** (2011) 014017, [arXiv:1012.2784](#).
- [19] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [20] LHCb collaboration, R. Aaij *et al.*, *LHCb detector performance*, Int. J. Mod. Phys. **A30** (2015) 1530022, [arXiv:1412.6352](#).
- [21] R. Aaij *et al.*, *Performance of the LHCb Vertex Locator*, JINST **9** (2014) P09007, [arXiv:1405.7808](#).
- [22] R. Arink *et al.*, *Performance of the LHCb Outer Tracker*, JINST **9** (2014) P01002, [arXiv:1311.3893](#).



- [23] P. d'Argent *et al.*, *Improved performance of the LHCb Outer Tracker in LHC Run 2*, JINST **12** (2017) P11016, [arXiv:1708.00819](#).
- [24] M. Adinolfi *et al.*, *Performance of the LHCb RICH detector at the LHC*, Eur. Phys. J. **C73** (2013) 2431, [arXiv:1211.6759](#).
- [25] A. A. Alves Jr. *et al.*, *Performance of the LHCb muon system*, JINST **8** (2013) P02022, [arXiv:1211.1346](#).
- [26] R. Aaij *et al.*, *The LHCb trigger and its performance in 2011*, JINST **8** (2013) P04022, [arXiv:1211.3055](#).
- [27] T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, Comput. Phys. Commun. **178** (2008) 852, [arXiv:0710.3820](#).
- [28] I. Belyaev *et al.*, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, J. Phys. Conf. Ser. **331** (2011) 032047.
- [29] D. J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [30] P. Golonka and Z. Was, *PHOTOS Monte Carlo: A precision tool for QED corrections in Z and W decays*, Eur. Phys. J. **C45** (2006) 97, [arXiv:hep-ph/0506026](#).
- [31] Geant4 collaboration, J. Allison *et al.*, *Geant4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; Geant4 collaboration, S. Agostinelli *et al.*, *Geant4: A simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [32] M. Clemencic *et al.*, *The LHCb simulation application, Gauss: Design, evolution and experience*, J. Phys. Conf. Ser. **331** (2011) 032023.
- [33] V. V. Gligorov and M. Williams, *Efficient, reliable and fast high-level triggering using a bonsai boosted decision tree*, JINST **8** (2013) P02013, [arXiv:1210.6861](#).
- [34] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, *Classification and regression trees*, Wadsworth international group, Belmont, California, USA, 1984.
- [35] B. P. Roe *et al.*, *Boosted decision trees, an alternative to artificial neural networks*, Nucl. Instrum. Meth. **A543** (2005) 577, [arXiv:physics/0408124](#).
- [36] U. P. Eitschberger, *Flavour-tagged measurement of CP observables in  $B_s^0 \rightarrow D_s^\mp K^\pm$  decays with the LHCb experiment*, PhD thesis, Tech. U., Dortmund (main), 2018, doi: 10.17877/DE290R-18881.
- [37] Particle Data Group, P. A. Zyla *et al.*, *Review of particle physics*, Prog. Theor. Exp. Phys. **2020** (2020) 083C01.
- [38] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP asymmetry in  $B_s^0 \rightarrow D_s^\mp K^\pm$  decays*, JHEP **03** (2018) 059, [arXiv:1712.07428](#).
- [39] D. Martínez Santos and F. Dupertuis, *Mass distributions marginalized over per-event errors*, Nucl. Instrum. Meth. **A764** (2014) 150, [arXiv:1312.5000](#).

- [40] N. L. Johnson, *Systems of frequency curves generated by methods of translation*, *Biometrika* **36** (1949) 149.
- [41] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP observables in  $B^\pm \rightarrow D^{(*)}K^\pm$  and  $B^\pm \rightarrow D^{(*)}\pi^\pm$  decays*, *Phys. Lett.* **B777** (2018) 16, [arXiv:1708.06370](#).
- [42] K. S. Cranmer, *Kernel estimation in high-energy physics*, *Comput. Phys. Commun.* **136** (2001) 198, [arXiv:hep-ex/0011057](#).
- [43] LHCb collaboration, R. Aaij *et al.*, *Measurement of the branching fraction and CP asymmetry in  $B^+ \rightarrow J/\psi \rho^+$  decays*, *Eur. Phys. J.* **C79** (2019) 537, [arXiv:1812.07041](#).
- [44] M. Pivk and F. R. Le Diberder, *sPlot: A statistical tool to unfold data distributions*, *Nucl. Instrum. Meth.* **A555** (2005) 356, [arXiv:physics/0402083](#).
- [45] LHCb collaboration, R. Aaij *et al.*, *Measurement of the fragmentation fraction ratio  $f_s/f_d$  and its dependence on B meson kinematics*, *JHEP* **04** (2013) 001, [arXiv:1301.5286](#).

## LHCb collaboration

R. Aaij<sup>32</sup>, C. Abellán Beteta<sup>50</sup>, T. Ackernley<sup>60</sup>, B. Adeva<sup>46</sup>, M. Adinolfi<sup>54</sup>, H. Afsharnia<sup>9</sup>, C.A. Aidala<sup>85</sup>, S. Aiola<sup>25</sup>, Z. Ajaltouni<sup>9</sup>, S. Akar<sup>65</sup>, J. Albrecht<sup>15</sup>, F. Alessio<sup>48</sup>, M. Alexander<sup>59</sup>, A. Alfonso Alberio<sup>45</sup>, Z. Aliouche<sup>62</sup>, G. Alkhazov<sup>38</sup>, P. Alvarez Cartelle<sup>55</sup>, S. Amato<sup>2</sup>, Y. Amhis<sup>11</sup>, L. An<sup>48</sup>, L. Anderlini<sup>22</sup>, A. Andreianov<sup>38</sup>, M. Andreotti<sup>21</sup>, F. Archilli<sup>17</sup>, A. Artamonov<sup>44</sup>, M. Artuso<sup>68</sup>, K. Arzymatov<sup>42</sup>, E. Aslanides<sup>10</sup>, M. Atzeni<sup>50</sup>, B. Audurier<sup>12</sup>, S. Bachmann<sup>17</sup>, M. Bachmayer<sup>49</sup>, J.J. Back<sup>56</sup>, S. Baker<sup>61</sup>, P. Baladron Rodriguez<sup>46</sup>, V. Balagura<sup>12</sup>, W. Baldini<sup>21</sup>, J. Baptista Leite<sup>1</sup>, R.J. Barlow<sup>62</sup>, S. Barsuk<sup>11</sup>, W. Barter<sup>61</sup>, M. Bartolini<sup>24,g</sup>, F. Baryshnikov<sup>82</sup>, J.M. Basels<sup>14</sup>, G. Bassi<sup>29</sup>, B. Batsukh<sup>68</sup>, A. Battig<sup>15</sup>, A. Bay<sup>49</sup>, M. Becker<sup>15</sup>, F. Bedeschi<sup>29</sup>, I. Bediaga<sup>1</sup>, A. Beiter<sup>68</sup>, V. Belavin<sup>42</sup>, S. Belin<sup>27</sup>, V. Bellee<sup>49</sup>, K. Belous<sup>44</sup>, I. Belov<sup>40</sup>, I. Belyaev<sup>41</sup>, G. Bencivenni<sup>23</sup>, E. Ben-Haim<sup>13</sup>, A. Berezhnoy<sup>40</sup>, R. Bernet<sup>50</sup>, D. Berninghoff<sup>17</sup>, H.C. Bernstein<sup>68</sup>, C. Bertella<sup>48</sup>, A. Bertolin<sup>28</sup>, C. Betancourt<sup>50</sup>, F. Betti<sup>20,c</sup>, I.a. Bezshyiko<sup>50</sup>, S. Bhasin<sup>54</sup>, J. Bhom<sup>35</sup>, L. Bian<sup>73</sup>, M.S. Bieker<sup>15</sup>, S. Bifani<sup>53</sup>, P. Billoir<sup>13</sup>, M. Birch<sup>61</sup>, F.C.R. Bishop<sup>55</sup>, A. Bizzeti<sup>22,j</sup>, M. Bjørn<sup>63</sup>, M.P. Blago<sup>48</sup>, T. Blake<sup>56</sup>, F. Blanc<sup>49</sup>, S. Blusk<sup>68</sup>, D. Bobulska<sup>59</sup>, J.A. Boelhauve<sup>15</sup>, O. Boente Garcia<sup>46</sup>, T. Boettcher<sup>64</sup>, A. Boldyrev<sup>81</sup>, A. Bondar<sup>43</sup>, N. Bondar<sup>38</sup>, S. Borghi<sup>62</sup>, M. Borisyak<sup>42</sup>, M. Borsato<sup>17</sup>, J.T. Borsuk<sup>35</sup>, S.A. Bouchiba<sup>49</sup>, T.J.V. Bowcock<sup>60</sup>, A. Boyer<sup>48</sup>, C. Bozzi<sup>21</sup>, M.J. Bradley<sup>61</sup>, S. Braun<sup>66</sup>, A. Brea Rodriguez<sup>46</sup>, M. Brodski<sup>48</sup>, J. Brodzicka<sup>35</sup>, A. Brossa Gonzalo<sup>56</sup>, D. Brundu<sup>27</sup>, A. Buonauro<sup>50</sup>, C. Burr<sup>48</sup>, A. Bursche<sup>27</sup>, A. Butkevich<sup>39</sup>, J.S. Butter<sup>32</sup>, J. Buytaert<sup>48</sup>, W. Byczynski<sup>48</sup>, S. Cadeddu<sup>27</sup>, H. Cai<sup>73</sup>, R. Calabrese<sup>21,e</sup>, L. Calefice<sup>15,13</sup>, L. Calero Diaz<sup>23</sup>, S. Cali<sup>23</sup>, R. Calladine<sup>53</sup>, M. Calvi<sup>26,i</sup>, M. Calvo Gomez<sup>84</sup>, P. Camargo Magalhaes<sup>54</sup>, A. Camboni<sup>45,84</sup>, P. Campana<sup>23</sup>, A.F. Campoverde Quezada<sup>6</sup>, S. Capelli<sup>26,i</sup>, L. Capriotti<sup>20,c</sup>, A. Carbone<sup>20,c</sup>, G. Carboni<sup>31</sup>, R. Cardinale<sup>24,g</sup>, A. Cardini<sup>27</sup>, I. Carli<sup>4</sup>, P. Carniti<sup>26,i</sup>, K. Carvalho Akiba<sup>32</sup>, A. Casais Vidal<sup>46</sup>, G. Casse<sup>60</sup>, M. Cattaneo<sup>48</sup>, G. Cavallero<sup>48</sup>, S. Celani<sup>49</sup>, J. Cerasoli<sup>10</sup>, A.J. Chadwick<sup>60</sup>, M.G. Chapman<sup>54</sup>, M. Charles<sup>13</sup>, Ph. Charpentier<sup>48</sup>, G. Chatzikonstantinidis<sup>53</sup>, C.A. Chavez Barajas<sup>60</sup>, M. Chefdeville<sup>8</sup>, C. Chen<sup>3</sup>, S. Chen<sup>27</sup>, A. Chernov<sup>35</sup>, S.-G. Chitic<sup>48</sup>, V. Chobanova<sup>46</sup>, S. Cholak<sup>49</sup>, M. Chruszcz<sup>35</sup>, A. Chubykin<sup>38</sup>, V. Chulikov<sup>38</sup>, P. Ciambone<sup>23</sup>, M.F. Cicala<sup>56</sup>, X. Cid Vidal<sup>46</sup>, G. Ciezarek<sup>48</sup>, P.E.L. Clarke<sup>58</sup>, M. Clemencic<sup>48</sup>, H.V. Cliff<sup>55</sup>, J. Closier<sup>48</sup>, J.L. Cobbedick<sup>62</sup>, V. Coco<sup>48</sup>, J.A.B. Coelho<sup>11</sup>, J. Cogan<sup>10</sup>, E. Cogneras<sup>9</sup>, L. Cojocariu<sup>37</sup>, P. Collins<sup>48</sup>, T. Colombo<sup>48</sup>, L. Congedo<sup>19,b</sup>, A. Contu<sup>27</sup>, N. Cooke<sup>53</sup>, G. Coombs<sup>59</sup>, G. Corti<sup>48</sup>, C.M. Costa Sobral<sup>56</sup>, B. Couturier<sup>48</sup>, D.C. Craik<sup>64</sup>, J. Crkovská<sup>67</sup>, M. Cruz Torres<sup>1</sup>, R. Currie<sup>58</sup>, C.L. Da Silva<sup>67</sup>, E. Dall'Occo<sup>15</sup>, J. Dalseno<sup>46</sup>, C. D'Ambrosio<sup>48</sup>, A. Danilina<sup>41</sup>, P. d'Argent<sup>48</sup>, A. Davis<sup>62</sup>, O. De Aguiar Francisco<sup>62</sup>, K. De Bruyn<sup>78</sup>, S. De Capua<sup>62</sup>, M. De Cian<sup>49</sup>, J.M. De Miranda<sup>1</sup>, L. De Paula<sup>2</sup>, M. De Serio<sup>19,b</sup>, D. De Simone<sup>50</sup>, P. De Simone<sup>23</sup>, J.A. de Vries<sup>79</sup>, C.T. Dean<sup>67</sup>, D. Decamp<sup>8</sup>, L. Del Buono<sup>13</sup>, B. Delaney<sup>55</sup>, H.-P. Dembinski<sup>15</sup>, A. Dendek<sup>34</sup>, V. Denysenko<sup>50</sup>, D. Derkach<sup>81</sup>, O. Deschamps<sup>9</sup>, F. Desse<sup>11</sup>, F. Dettori<sup>27,d</sup>, B. Dey<sup>73</sup>, P. Di Nezza<sup>23</sup>, S. Didenko<sup>82</sup>, L. Dieste Maronas<sup>46</sup>, H. Dijkstra<sup>48</sup>, V. Dobishuk<sup>52</sup>, A.M. Donohoe<sup>18</sup>, F. Dordei<sup>27</sup>, A.C. dos Reis<sup>1</sup>, L. Douglas<sup>59</sup>, A. Dovbnya<sup>51</sup>, A.G. Downes<sup>8</sup>, K. Dreimanis<sup>60</sup>, M.W. Dudek<sup>35</sup>, L. Dufour<sup>48</sup>, V. Duk<sup>77</sup>, P. Durante<sup>48</sup>, J.M. Durham<sup>67</sup>, D. Dutta<sup>62</sup>, M. Dziewiecki<sup>17</sup>, A. Dziurda<sup>35</sup>, A. Dzyuba<sup>38</sup>, S. Easo<sup>57</sup>, U. Egede<sup>69</sup>, V. Egorychev<sup>41</sup>, S. Eidelman<sup>43,u</sup>, S. Eisenhardt<sup>58</sup>, S. Ek-In<sup>49</sup>, L. Eklund<sup>59,v</sup>, S. Ely<sup>68</sup>, A. Ene<sup>37</sup>, E. Epple<sup>67</sup>, S. Escher<sup>14</sup>, J. Eschle<sup>50</sup>, S. Esen<sup>32</sup>, T. Evans<sup>48</sup>, A. Falabella<sup>20</sup>, J. Fan<sup>3</sup>, Y. Fan<sup>6</sup>, B. Fang<sup>73</sup>, N. Farley<sup>53</sup>, S. Farry<sup>60</sup>, D. Fazzini<sup>26,i</sup>, P. Fedin<sup>41</sup>, M. Féo<sup>48</sup>, P. Fernandez Declara<sup>48</sup>, A. Fernandez Prieto<sup>46</sup>, J.M. Fernandez-tenllado Arribas<sup>45</sup>, F. Ferrari<sup>20,c</sup>, L. Ferreira Lopes<sup>49</sup>, F. Ferreira Rodrigues<sup>2</sup>, S. Ferreres Sole<sup>32</sup>, M. Ferrillo<sup>50</sup>, M. Ferro-Luzzi<sup>48</sup>, S. Filippov<sup>39</sup>, R.A. Fini<sup>19</sup>, M. Fiorini<sup>21,e</sup>, M. Firlej<sup>34</sup>, K.M. Fischer<sup>63</sup>, C. Fitzpatrick<sup>62</sup>, T. Fiutowski<sup>34</sup>, F. Fleuret<sup>12</sup>, M. Fontana<sup>13</sup>, F. Fontanelli<sup>24,g</sup>, R. Forty<sup>48</sup>, V. Franco Lima<sup>60</sup>, M. Franco Sevilla<sup>66</sup>, M. Frank<sup>48</sup>, E. Franzoso<sup>21</sup>, G. Frau<sup>17</sup>, C. Frei<sup>48</sup>, D.A. Friday<sup>59</sup>, J. Fu<sup>25</sup>, Q. Fuehring<sup>15</sup>,

W. Funk<sup>48</sup>, E. Gabriel<sup>32</sup>, T. Gaintseva<sup>42</sup>, A. Gallas Torreira<sup>46</sup>, D. Galli<sup>20,c</sup>, S. Gambetta<sup>58,48</sup>,  
 Y. Gan<sup>3</sup>, M. Gandelman<sup>2</sup>, P. Gandini<sup>25</sup>, Y. Gao<sup>5</sup>, M. Garau<sup>27</sup>, L.M. Garcia Martin<sup>56</sup>,  
 P. Garcia Moreno<sup>45</sup>, J. García Pardiñas<sup>26</sup>, B. Garcia Plana<sup>46</sup>, F.A. Garcia Rosales<sup>12</sup>,  
 L. Garrido<sup>45</sup>, C. Gaspar<sup>48</sup>, R.E. Geertsema<sup>32</sup>, D. Gerick<sup>17</sup>, L.L. Gerken<sup>15</sup>, E. Gersabeck<sup>62</sup>,  
 M. Gersabeck<sup>62</sup>, T. Gershon<sup>56</sup>, D. Gerstel<sup>10</sup>, Ph. Ghez<sup>8</sup>, V. Gibson<sup>55</sup>, M. Giovannetti<sup>23,o</sup>,  
 A. Gioventù<sup>46</sup>, P. Gironella Gironell<sup>45</sup>, L. Giubega<sup>37</sup>, C. Giugliano<sup>21,e,48</sup>, K. Gizdov<sup>58</sup>,  
 E.L. Gkougkousis<sup>48</sup>, V.V. Gligorov<sup>13</sup>, C. Göbel<sup>70</sup>, E. Golobardes<sup>84</sup>, D. Golubkov<sup>41</sup>,  
 A. Golutvin<sup>61,82</sup>, A. Gomes<sup>1,a</sup>, S. Gomez Fernandez<sup>45</sup>, F. Goncalves Abrantes<sup>70</sup>, M. Goncerz<sup>35</sup>,  
 G. Gong<sup>3</sup>, P. Gorbounov<sup>41</sup>, I.V. Gorelov<sup>40</sup>, C. Gotti<sup>26,i</sup>, E. Govorkova<sup>48</sup>, J.P. Grabowski<sup>17</sup>,  
 R. Graciani Diaz<sup>45</sup>, T. Grammatico<sup>13</sup>, L.A. Granado Cardoso<sup>48</sup>, E. Graugés<sup>45</sup>, E. Graverini<sup>49</sup>,  
 G. Graziani<sup>22</sup>, A. Grecu<sup>37</sup>, L.M. Greeven<sup>32</sup>, P. Griffith<sup>21,e</sup>, L. Grillo<sup>62</sup>, S. Gromov<sup>82</sup>,  
 B.R. Gruberg Cazon<sup>63</sup>, C. Gu<sup>3</sup>, M. Guarise<sup>21</sup>, P. A. Günther<sup>17</sup>, E. Gushchin<sup>39</sup>, A. Guth<sup>14</sup>,  
 Y. Guz<sup>44,48</sup>, T. Gys<sup>48</sup>, T. Hadavizadeh<sup>69</sup>, G. Haefeli<sup>49</sup>, C. Haen<sup>48</sup>, J. Haimberger<sup>48</sup>,  
 T. Halewood-leagas<sup>60</sup>, P.M. Hamilton<sup>66</sup>, Q. Han<sup>7</sup>, X. Han<sup>17</sup>, T.H. Hancock<sup>63</sup>,  
 S. Hansmann-Menzemer<sup>17</sup>, N. Harnew<sup>63</sup>, T. Harrison<sup>60</sup>, C. Hasse<sup>48</sup>, M. Hatch<sup>48</sup>, J. He<sup>6</sup>,  
 M. Hecker<sup>61</sup>, K. Heijhoff<sup>32</sup>, K. Heinicke<sup>15</sup>, A.M. Hennequin<sup>48</sup>, K. Hennessy<sup>60</sup>, L. Henry<sup>25,47</sup>,  
 J. Heuel<sup>14</sup>, A. Hicheur<sup>2</sup>, D. Hill<sup>49</sup>, M. Hilton<sup>62</sup>, S.E. Hollitt<sup>15</sup>, J. Hu<sup>72</sup>, J. Hu<sup>17</sup>, W. Hu<sup>7</sup>,  
 W. Huang<sup>6</sup>, X. Huang<sup>73</sup>, W. Hulsbergen<sup>32</sup>, R.J. Hunter<sup>56</sup>, M. Hushchyn<sup>81</sup>, D. Hutchcroft<sup>60</sup>,  
 D. Hynds<sup>32</sup>, P. Ibis<sup>15</sup>, M. Idzik<sup>34</sup>, D. Ilin<sup>38</sup>, P. Ilten<sup>65</sup>, A. Inglese<sup>38</sup>, A. Ishteev<sup>82</sup>, K. Ivshin<sup>38</sup>,  
 R. Jacobsson<sup>48</sup>, S. Jakobsen<sup>48</sup>, E. Jans<sup>32</sup>, B.K. Jashal<sup>47</sup>, A. Jawahery<sup>66</sup>, V. Jevtic<sup>15</sup>,  
 M. Jezabek<sup>35</sup>, F. Jiang<sup>3</sup>, M. John<sup>63</sup>, D. Johnson<sup>48</sup>, C.R. Jones<sup>55</sup>, T.P. Jones<sup>56</sup>, B. Jost<sup>48</sup>,  
 N. Jurik<sup>48</sup>, S. Kandybei<sup>51</sup>, Y. Kang<sup>3</sup>, M. Karacson<sup>48</sup>, N. Kazeev<sup>81</sup>, F. Keizer<sup>55,48</sup>, M. Kenzie<sup>56</sup>,  
 T. Ketel<sup>33</sup>, B. Khanji<sup>15</sup>, A. Kharisova<sup>83</sup>, S. Kholodenko<sup>44</sup>, K.E. Kim<sup>68</sup>, T. Kirn<sup>14</sup>,  
 V.S. Kirsebom<sup>49</sup>, O. Kitouni<sup>64</sup>, S. Klaver<sup>32</sup>, K. Klimaszewski<sup>36</sup>, S. Koliiev<sup>52</sup>, A. Kondybayeva<sup>82</sup>,  
 A. Konoplyannikov<sup>41</sup>, P. Kopciewicz<sup>34</sup>, R. Kopecna<sup>17</sup>, P. Koppenburg<sup>32</sup>, M. Korolev<sup>40</sup>,  
 I. Kostiuik<sup>32,52</sup>, O. Kot<sup>52</sup>, S. Kotriakhova<sup>38,30</sup>, P. Kravchenko<sup>38</sup>, L. Kravchuk<sup>39</sup>,  
 R.D. Krawczyk<sup>48</sup>, M. Kreps<sup>56</sup>, F. Kress<sup>61</sup>, S. Kretzschmar<sup>14</sup>, P. Krokovny<sup>43,u</sup>, W. Krupa<sup>34</sup>,  
 W. Krzemien<sup>36</sup>, W. Kucewicz<sup>35,s</sup>, M. Kucharczyk<sup>35</sup>, V. Kudryavtsev<sup>43,u</sup>, H.S. Kuindersma<sup>32</sup>,  
 G.J. Kunde<sup>67</sup>, T. Kvaratskheliya<sup>41</sup>, D. Lacarrere<sup>48</sup>, G. Lafferty<sup>62</sup>, A. Lai<sup>27</sup>, A. Lampis<sup>27</sup>,  
 D. Lancierini<sup>50</sup>, J.J. Lane<sup>62</sup>, R. Lane<sup>54</sup>, G. Lanfranchi<sup>23</sup>, C. Langenbruch<sup>14</sup>, J. Langer<sup>15</sup>,  
 O. Lantwin<sup>50,82</sup>, T. Latham<sup>56</sup>, F. Lazzari<sup>29,p</sup>, R. Le Gac<sup>10</sup>, S.H. Lee<sup>85</sup>, R. Lefèvre<sup>9</sup>, A. Leflat<sup>40</sup>,  
 S. Legotin<sup>82</sup>, O. Leroy<sup>10</sup>, T. Lesiak<sup>35</sup>, B. Leverington<sup>17</sup>, H. Li<sup>72</sup>, L. Li<sup>63</sup>, P. Li<sup>17</sup>, Y. Li<sup>4</sup>, Y. Li<sup>4</sup>,  
 Z. Li<sup>68</sup>, X. Liang<sup>68</sup>, T. Lin<sup>61</sup>, R. Lindner<sup>48</sup>, V. Lisovskyi<sup>15</sup>, R. Litvinov<sup>27</sup>, G. Liu<sup>72</sup>, H. Liu<sup>6</sup>,  
 S. Liu<sup>4</sup>, X. Liu<sup>3</sup>, A. Loi<sup>27</sup>, J. Lomba Castro<sup>46</sup>, I. Longstaff<sup>59</sup>, J.H. Lopes<sup>2</sup>, G. Loustau<sup>50</sup>,  
 G.H. Lovell<sup>55</sup>, Y. Lu<sup>4</sup>, D. Lucchesi<sup>28,k</sup>, S. Luchuk<sup>39</sup>, M. Lucio Martinez<sup>32</sup>, V. Lukashenko<sup>32</sup>,  
 Y. Luo<sup>3</sup>, A. Lupato<sup>62</sup>, E. Luppi<sup>21,e</sup>, O. Lupton<sup>56</sup>, A. Lusiani<sup>29,l</sup>, X. Lyu<sup>6</sup>, L. Ma<sup>4</sup>,  
 S. Maccolini<sup>20,c</sup>, F. Machefert<sup>11</sup>, F. Maciuc<sup>37</sup>, V. Macko<sup>49</sup>, P. Mackowiak<sup>15</sup>,  
 S. Maddrell-Mander<sup>54</sup>, O. Madejczyk<sup>34</sup>, L.R. Madhan Mohan<sup>54</sup>, O. Maev<sup>38</sup>, A. Maevskiy<sup>81</sup>,  
 D. Maisuzenko<sup>38</sup>, M.W. Majewski<sup>34</sup>, S. Malde<sup>63</sup>, B. Malecki<sup>48</sup>, A. Malinin<sup>80</sup>, T. Maltsev<sup>43,u</sup>,  
 H. Malygina<sup>17</sup>, G. Manca<sup>27,d</sup>, G. Mancinelli<sup>10</sup>, R. Manera Escalero<sup>45</sup>, D. Manuzzi<sup>20,c</sup>,  
 D. Marangotto<sup>25,h</sup>, J. Maratas<sup>9,r</sup>, J.F. Marchand<sup>8</sup>, U. Marconi<sup>20</sup>, S. Mariani<sup>22,f,48</sup>,  
 C. Marin Benito<sup>11</sup>, M. Marinangeli<sup>49</sup>, P. Marino<sup>49,l</sup>, J. Marks<sup>17</sup>, P.J. Marshall<sup>60</sup>,  
 G. Martellotti<sup>30</sup>, L. Martinazzoli<sup>48,i</sup>, M. Martinelli<sup>26,i</sup>, D. Martinez Santos<sup>46</sup>,  
 F. Martinez Vidal<sup>47</sup>, A. Massafferri<sup>1</sup>, M. Materok<sup>14</sup>, R. Matev<sup>48</sup>, A. Mathad<sup>50</sup>, Z. Mathe<sup>48</sup>,  
 V. Matiunin<sup>41</sup>, C. Matteuzzi<sup>26</sup>, K.R. Mattioli<sup>85</sup>, A. Mauri<sup>32</sup>, E. Maurice<sup>12</sup>, J. Mauricio<sup>45</sup>,  
 M. Mazurek<sup>36</sup>, M. McCann<sup>61</sup>, L. Mcconnell<sup>18</sup>, T.H. Mcgrath<sup>62</sup>, A. McNab<sup>62</sup>, R. McNulty<sup>18</sup>,  
 J.V. Mead<sup>60</sup>, B. Meadows<sup>65</sup>, C. Meaux<sup>10</sup>, G. Meier<sup>15</sup>, N. Meinert<sup>76</sup>, D. Melnychuk<sup>36</sup>,  
 S. Meloni<sup>26,i</sup>, M. Merk<sup>32,79</sup>, A. Merli<sup>25</sup>, L. Meyer Garcia<sup>2</sup>, M. Mikhasenko<sup>48</sup>, D.A. Milanese<sup>74</sup>,  
 E. Millard<sup>56</sup>, M. Milovanovic<sup>48</sup>, M.-N. Minard<sup>8</sup>, L. Minzoni<sup>21,e</sup>, S.E. Mitchell<sup>58</sup>, B. Mitreska<sup>62</sup>,  
 D.S. Mitzel<sup>48</sup>, A. Mödden<sup>15</sup>, R.A. Mohammed<sup>63</sup>, R.D. Moise<sup>61</sup>, T. Mombächer<sup>15</sup>,

I.A. Monroy<sup>74</sup>, S. Monteil<sup>9</sup>, M. Morandin<sup>28</sup>, G. Morello<sup>23</sup>, M.J. Morello<sup>29,l</sup>, J. Moron<sup>34</sup>,  
 A.B. Morris<sup>75</sup>, A.G. Morris<sup>56</sup>, R. Mountain<sup>68</sup>, H. Mu<sup>3</sup>, F. Muheim<sup>58</sup>, M. Mukherjee<sup>7</sup>,  
 M. Mulder<sup>48</sup>, D. Müller<sup>48</sup>, K. Müller<sup>50</sup>, C.H. Murphy<sup>63</sup>, D. Murray<sup>62</sup>, P. Muzzetto<sup>27,48</sup>,  
 P. Naik<sup>54</sup>, T. Nakada<sup>49</sup>, R. Nandakumar<sup>57</sup>, T. Nanut<sup>49</sup>, I. Nasteva<sup>2</sup>, M. Needham<sup>58</sup>, I. Neri<sup>21,e</sup>,  
 N. Neri<sup>25,h</sup>, S. Neubert<sup>75</sup>, N. Neufeld<sup>48</sup>, R. Newcombe<sup>61</sup>, T.D. Nguyen<sup>49</sup>, C. Nguyen-Mau<sup>49,w</sup>,  
 E.M. Niel<sup>11</sup>, S. Nieswand<sup>14</sup>, N. Nikitin<sup>40</sup>, N.S. Nolte<sup>48</sup>, C. Nunez<sup>85</sup>, A. Oblakowska-Mucha<sup>34</sup>,  
 V. Obraztsov<sup>44</sup>, D.P. O'Hanlon<sup>54</sup>, R. Oldeman<sup>27,d</sup>, M.E. Olivares<sup>68</sup>, C.J.G. Onderwater<sup>78</sup>,  
 A. Ossowska<sup>35</sup>, J.M. Otorola Goicochea<sup>2</sup>, T. Ovsianikova<sup>41</sup>, P. Owen<sup>50</sup>, A. Oyanguren<sup>47</sup>,  
 B. Pagare<sup>56</sup>, P.R. Pais<sup>48</sup>, T. Pajero<sup>29,l,48</sup>, A. Palano<sup>19</sup>, M. Palutan<sup>23</sup>, Y. Pan<sup>62</sup>, G. Panshin<sup>83</sup>,  
 A. Papanestis<sup>57</sup>, M. Pappagallo<sup>19,b</sup>, L.L. Pappalardo<sup>21,e</sup>, C. Pappenheimer<sup>65</sup>, W. Parker<sup>66</sup>,  
 C. Parkes<sup>62</sup>, C.J. Parkinson<sup>46</sup>, B. Passalacqua<sup>21</sup>, G. Passaleva<sup>22</sup>, A. Pastore<sup>19</sup>, M. Patel<sup>61</sup>,  
 C. Patrignani<sup>20,c</sup>, C.J. Pawley<sup>79</sup>, A. Pearce<sup>48</sup>, A. Pellegrino<sup>32</sup>, M. Pepe Altarelli<sup>48</sup>,  
 S. Perazzini<sup>20</sup>, D. Pereima<sup>41</sup>, P. Perret<sup>9</sup>, K. Petridis<sup>54</sup>, A. Petrolini<sup>24,g</sup>, A. Petrov<sup>80</sup>,  
 S. Petrucci<sup>58</sup>, M. Petruzzo<sup>25</sup>, A. Philippov<sup>42</sup>, L. Pica<sup>29</sup>, M. Piccini<sup>77</sup>, B. Pietrzyk<sup>8</sup>,  
 G. Pietrzyk<sup>49</sup>, M. Pili<sup>63</sup>, D. Pinci<sup>30</sup>, F. Pisani<sup>48</sup>, A. Piucci<sup>17</sup>, Resmi P.K<sup>10</sup>, V. Placinta<sup>37</sup>,  
 J. Plews<sup>53</sup>, M. Plo Casasus<sup>46</sup>, F. Polci<sup>13</sup>, M. Poli Lener<sup>23</sup>, M. Poliakov<sup>68</sup>, A. Poluektov<sup>10</sup>,  
 N. Polukhina<sup>82,t</sup>, I. Polyakov<sup>68</sup>, E. Polycarpo<sup>2</sup>, G.J. Pomery<sup>54</sup>, S. Ponce<sup>48</sup>, D. Popov<sup>6,48</sup>,  
 S. Popov<sup>42</sup>, S. Poslavskii<sup>44</sup>, K. Prasanth<sup>35</sup>, L. Promberger<sup>48</sup>, C. Prouve<sup>46</sup>, V. Pugatch<sup>52</sup>,  
 H. Pullen<sup>63</sup>, G. Punzi<sup>29,m</sup>, W. Qian<sup>6</sup>, J. Qin<sup>6</sup>, R. Quagliani<sup>13</sup>, B. Quintana<sup>8</sup>, N.V. Raab<sup>18</sup>,  
 R.I. Rabadan Trejo<sup>10</sup>, B. Rachwal<sup>34</sup>, J.H. Rademacker<sup>54</sup>, M. Rama<sup>29</sup>, M. Ramos Pernas<sup>56</sup>,  
 M.S. Rangel<sup>2</sup>, F. Ratnikov<sup>42,81</sup>, G. Raven<sup>33</sup>, M. Reboud<sup>8</sup>, F. Redi<sup>49</sup>, F. Reiss<sup>13</sup>,  
 C. Remon Alepuz<sup>47</sup>, Z. Ren<sup>3</sup>, V. Renaudin<sup>63</sup>, R. Ribatti<sup>29</sup>, S. Ricciardi<sup>57</sup>, D.S. Richards<sup>57</sup>,  
 K. Rinnert<sup>60</sup>, P. Robbe<sup>11</sup>, A. Robert<sup>13</sup>, G. Robertson<sup>58</sup>, A.B. Rodrigues<sup>49</sup>, E. Rodrigues<sup>60</sup>,  
 J.A. Rodriguez Lopez<sup>74</sup>, A. Rollings<sup>63</sup>, P. Roloff<sup>48</sup>, V. Romanovskiy<sup>44</sup>, M. Romero Lamas<sup>46</sup>,  
 A. Romero Vidal<sup>46</sup>, J.D. Roth<sup>85</sup>, M. Rotondo<sup>23</sup>, M.S. Rudolph<sup>68</sup>, T. Ruf<sup>48</sup>, J. Ruiz Vidal<sup>47</sup>,  
 A. Ryzhikov<sup>81</sup>, J. Ryzka<sup>34</sup>, J.J. Saborido Silva<sup>46</sup>, N. Sagidova<sup>38</sup>, N. Sahoo<sup>56</sup>, B. Saitta<sup>27,d</sup>,  
 D. Sanchez Gonzalo<sup>45</sup>, C. Sanchez Gras<sup>32</sup>, R. Santacesaria<sup>30</sup>, C. Santamarina Rios<sup>46</sup>,  
 M. Santimaria<sup>23</sup>, E. Santovetti<sup>31,o</sup>, D. Saranin<sup>82</sup>, G. Sarpis<sup>59</sup>, M. Sarpis<sup>75</sup>, A. Sarti<sup>30</sup>,  
 C. Satriano<sup>30,n</sup>, A. Satta<sup>31</sup>, M. Saur<sup>15</sup>, D. Savrina<sup>41,40</sup>, H. Sazak<sup>9</sup>, L.G. Scantlebury Smead<sup>63</sup>,  
 S. Schael<sup>14</sup>, M. Schellenberg<sup>15</sup>, M. Schiller<sup>59</sup>, H. Schindler<sup>48</sup>, M. Schmelling<sup>16</sup>, B. Schmidt<sup>48</sup>,  
 O. Schneider<sup>49</sup>, A. Schopper<sup>48</sup>, M. Schubiger<sup>32</sup>, S. Schulte<sup>49</sup>, M.H. Schune<sup>11</sup>, R. Schwemmer<sup>48</sup>,  
 B. Sciascia<sup>23</sup>, A. Sciubba<sup>23</sup>, S. Sellam<sup>46</sup>, A. Semennikov<sup>41</sup>, M. Senghi Soares<sup>33</sup>, A. Sergi<sup>53,48</sup>,  
 N. Serra<sup>50</sup>, L. Sestini<sup>28</sup>, A. Seuthe<sup>15</sup>, P. Seyfert<sup>48</sup>, D.M. Shangase<sup>85</sup>, M. Shapkin<sup>44</sup>,  
 I. Shchemerov<sup>82</sup>, L. Shchutska<sup>49</sup>, T. Shears<sup>60</sup>, L. Shekhtman<sup>43,u</sup>, Z. Shen<sup>5</sup>, V. Shevchenko<sup>80</sup>,  
 E.B. Shields<sup>26,i</sup>, E. Shmanin<sup>82</sup>, J.D. Shupperd<sup>68</sup>, B.G. Siddi<sup>21</sup>, R. Silva Coutinho<sup>50</sup>, G. Simi<sup>28</sup>,  
 S. Simone<sup>19,b</sup>, I. Skiba<sup>21,e</sup>, N. Skidmore<sup>62</sup>, T. Skwarnicki<sup>68</sup>, M.W. Slater<sup>53</sup>, J.C. Smallwood<sup>63</sup>,  
 J.G. Smeaton<sup>55</sup>, A. Smetkina<sup>41</sup>, E. Smith<sup>14</sup>, M. Smith<sup>61</sup>, A. Snoch<sup>32</sup>, M. Soares<sup>20</sup>,  
 L. Soares Lavra<sup>9</sup>, M.D. Sokoloff<sup>65</sup>, F.J.P. Soler<sup>59</sup>, A. Solovov<sup>38</sup>, I. Solovyevev<sup>38</sup>,  
 F.L. Souza De Almeida<sup>2</sup>, B. Souza De Paula<sup>2</sup>, B. Spaan<sup>15</sup>, E. Spadaro Norella<sup>25,h</sup>, P. Spradlin<sup>59</sup>,  
 F. Stagni<sup>48</sup>, M. Stahl<sup>65</sup>, S. Stahl<sup>48</sup>, P. Stefko<sup>49</sup>, O. Steinkamp<sup>50,82</sup>, S. Stemmler<sup>17</sup>,  
 O. Stenyakin<sup>44</sup>, H. Stevens<sup>15</sup>, S. Stone<sup>68</sup>, M.E. Stramaglia<sup>49</sup>, M. Straticiu<sup>37</sup>, D. Strelakina<sup>82</sup>,  
 S. Strovkov<sup>83</sup>, F. Suljik<sup>63</sup>, J. Sun<sup>27</sup>, L. Sun<sup>73</sup>, Y. Sun<sup>66</sup>, P. Svihra<sup>62</sup>, P.N. Swallow<sup>53</sup>,  
 K. Swientek<sup>34</sup>, A. Szabelski<sup>36</sup>, T. Szumlak<sup>34</sup>, M. Szymanski<sup>48</sup>, S. Taneja<sup>62</sup>, F. Teubert<sup>48</sup>,  
 E. Thomas<sup>48</sup>, K.A. Thomson<sup>60</sup>, M.J. Tilley<sup>61</sup>, V. Tisserand<sup>9</sup>, S. T'Jampens<sup>8</sup>, M. Tobin<sup>4</sup>,  
 S. Tolk<sup>48</sup>, L. Tomassetti<sup>21,e</sup>, D. Torres Machado<sup>1</sup>, D.Y. Tou<sup>13</sup>, M. Traill<sup>59</sup>, M.T. Tran<sup>49</sup>,  
 E. Trifonova<sup>82</sup>, C. Trippi<sup>49</sup>, G. Tuci<sup>29,m</sup>, A. Tully<sup>49</sup>, N. Tuning<sup>32</sup>, A. Ukleja<sup>36</sup>, D.J. Unverzagt<sup>17</sup>,  
 A. Usachov<sup>32</sup>, A. Ustyuzhanin<sup>42,81</sup>, U. Uwer<sup>17</sup>, A. Vagner<sup>83</sup>, V. Vagnoni<sup>20</sup>, A. Valassi<sup>48</sup>,  
 G. Valenti<sup>20</sup>, N. Valls Canudas<sup>45</sup>, M. van Beuzekom<sup>32</sup>, E. van Herwijnen<sup>82</sup>, C.B. Van Hulse<sup>18</sup>,  
 M. van Veghel<sup>78</sup>, R. Vazquez Gomez<sup>46</sup>, P. Vazquez Regueiro<sup>46</sup>, C. Vázquez Sierra<sup>48</sup>, S. Vecchi<sup>21</sup>,  
 J.J. Velthuis<sup>54</sup>, M. Veltri<sup>22,q</sup>, A. Venkateswaran<sup>68</sup>, M. Veronesi<sup>32</sup>, M. Vesterinen<sup>56</sup>, D. Vieira<sup>65</sup>,

M. Vieites Diaz<sup>49</sup>, H. Viemann<sup>76</sup>, X. Vilasis-Cardona<sup>84</sup>, E. Vilella Figueras<sup>60</sup>, P. Vincent<sup>13</sup>, G. Vitali<sup>29</sup>, A. Vollhardt<sup>50</sup>, D. Vom Bruch<sup>13</sup>, A. Vorobyev<sup>38</sup>, V. Vorobyev<sup>43,u</sup>, N. Voropaev<sup>38</sup>, R. Waldi<sup>76</sup>, J. Walsh<sup>29</sup>, C. Wang<sup>17</sup>, J. Wang<sup>73</sup>, J. Wang<sup>3</sup>, J. Wang<sup>5</sup>, J. Wang<sup>4</sup>, M. Wang<sup>3</sup>, R. Wang<sup>54</sup>, Y. Wang<sup>7</sup>, Z. Wang<sup>50</sup>, H.M. Wark<sup>60</sup>, N.K. Watson<sup>53</sup>, S.G. Weber<sup>13</sup>, D. Websdale<sup>61</sup>, C. Weisser<sup>64</sup>, B.D.C. Westhenry<sup>54</sup>, D.J. White<sup>62</sup>, M. Whitehead<sup>54</sup>, D. Wiedner<sup>15</sup>, G. Wilkinson<sup>63</sup>, M. Wilkinson<sup>68</sup>, I. Williams<sup>55</sup>, M. Williams<sup>64,69</sup>, M.R.J. Williams<sup>58</sup>, F.F. Wilson<sup>57</sup>, W. Wislicki<sup>36</sup>, M. Witek<sup>35</sup>, L. Witola<sup>17</sup>, G. Wormser<sup>11</sup>, S.A. Wotton<sup>55</sup>, H. Wu<sup>68</sup>, K. Wyllie<sup>48</sup>, Z. Xiang<sup>6</sup>, D. Xiao<sup>7</sup>, Y. Xie<sup>7</sup>, A. Xu<sup>5</sup>, J. Xu<sup>6</sup>, L. Xu<sup>3</sup>, M. Xu<sup>7</sup>, Q. Xu<sup>6</sup>, Z. Xu<sup>6</sup>, Z. Xu<sup>5</sup>, D. Yang<sup>3</sup>, Y. Yang<sup>6</sup>, Z. Yang<sup>3</sup>, Z. Yang<sup>66</sup>, Y. Yao<sup>68</sup>, L.E. Yeomans<sup>60</sup>, H. Yin<sup>7</sup>, J. Yu<sup>71</sup>, X. Yuan<sup>68</sup>, O. Yushchenko<sup>44</sup>, K.A. Zarebski<sup>53</sup>, M. Zavertyaev<sup>16,t</sup>, M. Zdybal<sup>35</sup>, O. Zenaiev<sup>48</sup>, M. Zeng<sup>3</sup>, D. Zhang<sup>7</sup>, L. Zhang<sup>3</sup>, S. Zhang<sup>5</sup>, Y. Zhang<sup>5</sup>, Y. Zhang<sup>63</sup>, A. Zhelezov<sup>17</sup>, Y. Zheng<sup>6</sup>, X. Zhou<sup>6</sup>, Y. Zhou<sup>6</sup>, X. Zhu<sup>3</sup>, V. Zhukov<sup>14,40</sup>, J.B. Zonneveld<sup>58</sup>, S. Zucchelli<sup>20,c</sup>, D. Zuliani<sup>28</sup>, G. Zunica<sup>62</sup>.

<sup>1</sup>Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

<sup>2</sup>Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

<sup>3</sup>Center for High Energy Physics, Tsinghua University, Beijing, China

<sup>4</sup>Institute Of High Energy Physics (IHEP), Beijing, China

<sup>5</sup>School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

<sup>6</sup>University of Chinese Academy of Sciences, Beijing, China

<sup>7</sup>Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China

<sup>8</sup>Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

<sup>9</sup>Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France

<sup>10</sup>Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

<sup>11</sup>Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France

<sup>12</sup>Laboratoire Leprince-ringuet (lir), Palaiseau, France

<sup>13</sup>LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France

<sup>14</sup>I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

<sup>15</sup>Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

<sup>16</sup>Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

<sup>17</sup>Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

<sup>18</sup>School of Physics, University College Dublin, Dublin, Ireland

<sup>19</sup>INFN Sezione di Bari, Bari, Italy

<sup>20</sup>INFN Sezione di Bologna, Bologna, Italy

<sup>21</sup>INFN Sezione di Ferrara, Ferrara, Italy

<sup>22</sup>INFN Sezione di Firenze, Firenze, Italy

<sup>23</sup>INFN Laboratori Nazionali di Frascati, Frascati, Italy

<sup>24</sup>INFN Sezione di Genova, Genova, Italy

<sup>25</sup>INFN Sezione di Milano, Milano, Italy

<sup>26</sup>INFN Sezione di Milano-Bicocca, Milano, Italy

<sup>27</sup>INFN Sezione di Cagliari, Monserrato, Italy

<sup>28</sup>Università degli Studi di Padova, Università e INFN, Padova, Padova, Italy

<sup>29</sup>INFN Sezione di Pisa, Pisa, Italy

<sup>30</sup>INFN Sezione di Roma La Sapienza, Roma, Italy

<sup>31</sup>INFN Sezione di Roma Tor Vergata, Roma, Italy

<sup>32</sup>Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands

<sup>33</sup>Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands

<sup>34</sup>AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland

<sup>35</sup>Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland

<sup>36</sup>National Center for Nuclear Research (NCBJ), Warsaw, Poland

<sup>37</sup>Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania

<sup>38</sup>Petersburg Nuclear Physics Institute NRC Kurchatov Institute (PNPI NRC KI), Gatchina, Russia

- <sup>39</sup>*Institute for Nuclear Research of the Russian Academy of Sciences (INR RAS), Moscow, Russia*
- <sup>40</sup>*Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- <sup>41</sup>*Institute of Theoretical and Experimental Physics NRC Kurchatov Institute (ITEP NRC KI), Moscow, Russia*
- <sup>42</sup>*Yandex School of Data Analysis, Moscow, Russia*
- <sup>43</sup>*Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia*
- <sup>44</sup>*Institute for High Energy Physics NRC Kurchatov Institute (IHEP NRC KI), Protvino, Russia, Protvino, Russia*
- <sup>45</sup>*ICCUB, Universitat de Barcelona, Barcelona, Spain*
- <sup>46</sup>*Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- <sup>47</sup>*Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain*
- <sup>48</sup>*European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- <sup>49</sup>*Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- <sup>50</sup>*Physik-Institut, Universität Zürich, Zürich, Switzerland*
- <sup>51</sup>*NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- <sup>52</sup>*Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- <sup>53</sup>*University of Birmingham, Birmingham, United Kingdom*
- <sup>54</sup>*H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- <sup>55</sup>*Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- <sup>56</sup>*Department of Physics, University of Warwick, Coventry, United Kingdom*
- <sup>57</sup>*STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- <sup>58</sup>*School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- <sup>59</sup>*School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- <sup>60</sup>*Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- <sup>61</sup>*Imperial College London, London, United Kingdom*
- <sup>62</sup>*Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- <sup>63</sup>*Department of Physics, University of Oxford, Oxford, United Kingdom*
- <sup>64</sup>*Massachusetts Institute of Technology, Cambridge, MA, United States*
- <sup>65</sup>*University of Cincinnati, Cincinnati, OH, United States*
- <sup>66</sup>*University of Maryland, College Park, MD, United States*
- <sup>67</sup>*Los Alamos National Laboratory (LANL), Los Alamos, United States*
- <sup>68</sup>*Syracuse University, Syracuse, NY, United States*
- <sup>69</sup>*School of Physics and Astronomy, Monash University, Melbourne, Australia, associated to <sup>56</sup>*
- <sup>70</sup>*Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to <sup>2</sup>*
- <sup>71</sup>*Physics and Micro Electronic College, Hunan University, Changsha City, China, associated to <sup>7</sup>*
- <sup>72</sup>*Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter, South China Normal University, Guangzhou, China, associated to <sup>3</sup>*
- <sup>73</sup>*School of Physics and Technology, Wuhan University, Wuhan, China, associated to <sup>3</sup>*
- <sup>74</sup>*Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to <sup>13</sup>*
- <sup>75</sup>*Universität Bonn - Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany, associated to <sup>17</sup>*
- <sup>76</sup>*Institut für Physik, Universität Rostock, Rostock, Germany, associated to <sup>17</sup>*
- <sup>77</sup>*INFN Sezione di Perugia, Perugia, Italy, associated to <sup>21</sup>*
- <sup>78</sup>*Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to <sup>32</sup>*
- <sup>79</sup>*Universiteit Maastricht, Maastricht, Netherlands, associated to <sup>32</sup>*
- <sup>80</sup>*National Research Centre Kurchatov Institute, Moscow, Russia, associated to <sup>41</sup>*
- <sup>81</sup>*National Research University Higher School of Economics, Moscow, Russia, associated to <sup>42</sup>*
- <sup>82</sup>*National University of Science and Technology "MISIS", Moscow, Russia, associated to <sup>41</sup>*
- <sup>83</sup>*National Research Tomsk Polytechnic University, Tomsk, Russia, associated to <sup>41</sup>*
- <sup>84</sup>*DS4DS, La Salle, Universitat Ramon Llull, Barcelona, Spain, associated to <sup>45</sup>*
- <sup>85</sup>*University of Michigan, Ann Arbor, United States, associated to <sup>68</sup>*

<sup>a</sup>*Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil*

<sup>b</sup>*Università di Bari, Bari, Italy*

<sup>c</sup>*Università di Bologna, Bologna, Italy*

<sup>d</sup>*Università di Cagliari, Cagliari, Italy*

<sup>e</sup>*Università di Ferrara, Ferrara, Italy*

- <sup>f</sup> *Università di Firenze, Firenze, Italy*  
<sup>g</sup> *Università di Genova, Genova, Italy*  
<sup>h</sup> *Università degli Studi di Milano, Milano, Italy*  
<sup>i</sup> *Università di Milano Bicocca, Milano, Italy*  
<sup>j</sup> *Università di Modena e Reggio Emilia, Modena, Italy*  
<sup>k</sup> *Università di Padova, Padova, Italy*  
<sup>l</sup> *Scuola Normale Superiore, Pisa, Italy*  
<sup>m</sup> *Università di Pisa, Pisa, Italy*  
<sup>n</sup> *Università della Basilicata, Potenza, Italy*  
<sup>o</sup> *Università di Roma Tor Vergata, Roma, Italy*  
<sup>p</sup> *Università di Siena, Siena, Italy*  
<sup>q</sup> *Università di Urbino, Urbino, Italy*  
<sup>r</sup> *MSU - Iligan Institute of Technology (MSU-IIT), Iligan, Philippines*  
<sup>s</sup> *AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*  
<sup>t</sup> *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*  
<sup>u</sup> *Novosibirsk State University, Novosibirsk, Russia*  
<sup>v</sup> *Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden*  
<sup>w</sup> *Hanoi University of Science, Hanoi, Vietnam*