EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH (CERN)

CERN-EP-2021-130 LHCb-PAPER-2021-015 June 21, 2022

Measurement of the nuclear modification factor and prompt charged particle production in pPb and pp collisions at $\sqrt{s_{NN}} = 5 \text{ TeV}$

LHC_b collaboration^{[†](#page-0-0)}

Abstract

The production of prompt charged particles in proton-lead collisions and in protonproton collisions at the nucleon-nucleon centre-of-mass energy $\sqrt{s_{NN}} = 5$ TeV is studied at LHCb as a function of pseudorapidity (η) and transverse momentum (p_T) with respect to the proton beam direction. The nuclear modification factor for charged particles is determined as a function of η between $-4.8 < \eta < -2.5$ (backward region) and $2.0 < \eta < 4.8$ (forward region), and p_T between $0.2 < p_T < 8.0$ GeV/c. The results show a suppression of charged particle production in proton-lead collisions relative to proton-proton collisions in the forward region and an enhancement in the backward region for p_T larger than 1.5 GeV/c. This measurement constrains nuclear PDFs and saturation models at previously unexplored values of the parton momentum fraction down to 10⁻⁶.

Published in Phys.Rev.Lett. 128 (2022), 142004

© 2022 CERN for the benefit of the LHCb collaboration. [CC BY 4.0 licence.](https://creativecommons.org/licenses/by/4.0/)

[†]Authors are listed at the end of this paper.

Charged particle production in hadronic collisions is a fundamental observable for studying the properties of the strong interaction governed by quantum chromodynamics (QCD). In high-energy collisions at the Large Hadron Collider (LHC), charged particles can be produced in soft and hard interactions which correspond to small and large momentum exchanges between the interacting partons of the hadrons, respectively. While hard interactions can be described by perturbative QCD (pQCD), the soft regime is less well understood and predictions currently rely on phenomenological considerations [\[1,](#page-9-0) [2\]](#page-9-1). Even at LHC energies, charged particles from soft interactions dominate over those from hard interactions. For this reason, experimental input is crucial to improve models and generators for hadron collider and cosmic ray physics [\[3](#page-9-2)[–5\]](#page-9-3).

The study of the hard regime, which corresponds to charged particles of high transverse momentum (p_T) with respect to the axis of the colliding hadrons, provides valuable information on the physics of heavy-ion collisions [\[6\]](#page-9-4). Modifications of the charged particle production rate in proton-lead (pPb) collisions relative to proton-proton (pp) collisions can be modelled assuming a variety of cold nuclear matter (CNM) effects [\[7,](#page-9-5) [8\]](#page-10-0). Recent indications of collective fluid-like phenomena in small systems suggest the presence of dynamics not generally classified as CNM effects but as signatures of a quark gluon plasma [\[9\]](#page-10-1).

For charged particles, these modifications are generally associated with initial-state effects, parameterised in nuclear parton distribution functions (nPDFs) [\[10](#page-10-2)[–12\]](#page-10-3). Other nuclear effects are related to initial- or final-state multiple scatterings of incoming and outgoing partons [\[13,](#page-10-4) [14\]](#page-10-5), and could manifest in a Cronin enhancement [\[15\]](#page-10-6). Another approach considers models based on parton saturation, an effect arising at low values of the parton momentum fraction, x, and heavy nuclei [\[16\]](#page-10-7). In this regime, the QCD dynamics can be described by the colour glass condensate (CGC) effective field theory [\[17\]](#page-10-8). Pion production at central rapidity [\[18\]](#page-10-9) is well described by modified nPDFs, energy loss and CGC calculations [\[10,](#page-10-2) [11,](#page-10-10) [19,](#page-10-11) [20\]](#page-10-12). Low values of x, where saturation effects are most likely to occur, can be probed with high-energy collisions at the most forward rapidities.

Previous studies at the LHC [\[21](#page-10-13)[–23\]](#page-11-0) have measured prompt charged particle production in pPb collisions at the centre-of-mass energy in the nucleon-nucleon system $\sqrt{s_{\text{NN}}} = 5 \text{ TeV}$ in the central pseudorapidity region. At the Relativistic Heavy Ion Collider (RHIC), measurements with deuteron-gold and proton-gold (pAu) collisions at more forward rapidities but lower energy ($\sqrt{s_{\text{NN}}}$ = 200 GeV) have been performed [\[24](#page-11-1)[–26\]](#page-11-2). The LHCb experiment can uniquely probe the lowest x ranges currently accessible, given its forward rapidity coverage and higher collision energy.

This Letter presents the measurement of the prompt charged particle spectra in pPb and pp collisions at $\sqrt{s_{\text{NN}}}$ = 5 TeV in the 0.2 < p_T < 8.0 GeV/c range, thus covering the soft and hard production regimes. The pPb measurement covers the *backward* pseudorapidity range of $-5.2 < \eta < -2.5$, where the lead beam enters the LHCb spectrometer at the interaction point, and the *forward* pseudorapidity range of $1.6 < \eta < 4.3$, where the proton beam enters the LHCb spectrometer at the interaction point. The pp measurement spans over $2.0 < \eta < 4.8$, and complements the recent measurement of prompt charged particle production at $\sqrt{s_{\text{NN}}}$ = 13 TeV [\[27\]](#page-11-3). Throughout the text η is expressed in the nucleonnucleon centre-of-mass system, and is related with the pseudorapidity in the laboratory frame η_{lab} by $\eta = \eta_{\text{lab}} - 0.465$ for pPb and $\eta = \eta_{\text{lab}}$ for pp collisions.

The double-differential production cross-section for prompt charged particles is mea-

sured as

$$
\frac{\mathrm{d}^2 \sigma^{\mathrm{ch}}(\eta, p_{\mathrm{T}})}{\mathrm{d}p_{\mathrm{T}} \mathrm{d}\eta} \equiv \frac{1}{\mathcal{L}} \frac{N^{\mathrm{ch}}(\eta, p_{\mathrm{T}})}{\Delta p_{\mathrm{T}} \Delta \eta}.
$$
\n(1)

Here, N^{ch} is the number of prompt charged particles produced in a given interval of η and $p_{\rm T}$, $\Delta \eta$ and $\Delta p_{\rm T}$, respectively, and $\mathcal L$ is the integrated luminosity of the corresponding data sample. In this study a prompt charged particle is any charged hadron or lepton with a mean lifetime above 0.3×10^{-10} s produced directly in the collision or from decays of shorter-lifetime particles [\[28\]](#page-11-4). The nuclear modification factor, R_{pPb} , is defined as

$$
R_{p\text{Pb}}(\eta, p_{\text{T}}) \equiv \frac{1}{A} \frac{\mathrm{d}^2 \sigma_{p\text{Pb}}^{\text{ch}}(\eta, p_{\text{T}})/\mathrm{d}p_{\text{T}} \mathrm{d}\eta}{\mathrm{d}^2 \sigma_{pp}^{\text{ch}}(\eta, p_{\text{T}})/\mathrm{d}p_{\text{T}} \mathrm{d}\eta},\tag{2}
$$

where $A = 208$ is the number of nucleons in the lead ion and $d^2 \sigma_{pPb,pp}^{ch}(\eta, p_T) / dp_T d\eta$ is the double-differential cross-section in pPb and pp collisions, respectively.

The LHCb detector is a single-arm forward spectrometer described in Refs. [\[29,](#page-11-5) [30\]](#page-11-6). The detector elements that are particularly relevant to this analysis are a silicon-strip vertex detector (VELO) surrounding the interaction region that allows the determination of the position of the collision point, known as the primary vertex (PV) , a tracking system that provides a measurement of the momentum, p , of charged particles and two ring-imaging Cherenkov detectors that are able to discriminate between different species of charged particles.

The corresponding integrated luminosity for the forward (backward) pPb data sample is $42.7 \pm 1.0 \,\mathrm{\mu b}^{-1}$ $(38.7 \pm 1.0 \,\mathrm{\mu b}^{-1})$ [\[31\]](#page-11-7), where the uncertainties are uncorrelated between the two data samples. Events are required to pass a minimum-bias trigger which requires at least one reconstructed track in the VELO detector. Additionally, only events with one reconstructed PV within three standard deviations from the mean PV position of the full sample are considered.

The pp data correspond to an integrated luminosity of 3.49 ± 0.07 nb⁻¹. An unbiased trigger for pp events is used to select every leading bunch crossing that occurred during the data taking period, thus avoiding potential contamination between neighbouring bunches.

Simulation is used to model the reconstruction efficiency, the effects of the selection requirements, and the contribution from background tracks. In the simulation, pPb collisions are generated using EPOS-LHC [\[32\]](#page-11-8), while pp collisions are generated using Pythia [\[33\]](#page-11-9) with a specific LHCb configuration [\[34\]](#page-11-10). Particle decays are described by EVTGEN [\[35\]](#page-11-11), while the interaction of particles with the detector, and its response in simulation, are implemented using the GEANT4 toolkit [\[36,](#page-11-12)37] as described in Ref. [\[38\]](#page-11-14).

Prompt charged particle candidates are defined as tracks with hits in the VELO and the tracking stations after the LHCb detector dipole. This last condition requires the measured particle to have $p > 2 \text{GeV}/c$. Background contributions due to fake tracks and secondary particles are considered in this study. Fake tracks are reconstruction artefacts that do not correspond to actual charged particles, which are particularly relevant for large detector occupancies and in the high- p_T region. Secondary particles are tracks produced by charged particles that do not meet the prompt particle definition and originate from interactions of particles with the detector material or from decays of prompt particles.

A selection is applied to reduce these background contributions. Fake tracks are suppressed with a tight requirement on the output of a neural-net based algorithm (ghost probability) [\[39\]](#page-12-0). To suppress further the fake track background, when two or more candidates from the same event share a segment reconstructed in the VELO, only the candidate with the best track fit-quality is retained. Secondary particles are reduced by requiring small impact parameters with respect to the mean of the PV distribution in the full sample. This criterion is particularly effective at removing hadrons produced in decays of prompt K^0_S and Λ particles and in interactions of hadrons with the detector material without inducing a bias by requiring a PV.

The prompt charged particle yield, N^{ch} , for the pPb and pp samples is obtained from the number of candidates, which is corrected with the reconstruction efficiency, the selection efficiency and the signal purity. The reconstruction efficiency accounts for detector inefficiencies or acceptance effects. The track-finding efficiency from simulation is corrected with a tag-and-probe method applied to data and simulation in two-dimensional intervals of η_{ab} and p_T using $J/\psi \to \mu^+\mu^-$ decays in the range $5 < p < 200$ GeV/c [\[40\]](#page-12-1). Since the reconstruction efficiency depends on the detector occupancy, the simulated samples are weighted to reproduce the occupancy distributions from different LHCb subdetectors in data. Additionally, the reconstruction efficiency depends on the particle type. The relative abundance of particles determined from simulation are validated with data from the ALICE [\[41–](#page-12-2)[44\]](#page-12-3) and LHCb [\[45\]](#page-12-4) experiments. The LHCb Pythia tune for pp collisions does not reproduce the kaon and prompt hyperons relative abundance at high p_T in data. Therefore, a dedicated simulated sample generated with EPOS-LHC [\[32\]](#page-11-8) is used to parameterise the particle composition in pp collisions. The relative abundances produced with the EPOS-LHC generator are in agreement with the data within 30% in pp and pPb collisions. As a cross-check, the relative abundances from EPOS-LHC in the LHCb acceptance are found to be compatible with those produced with Pythia using the rope hadronisation model [\[46,](#page-12-5) [47\]](#page-12-6). The use of the uncorrected LHCb Pythia tune would imply an overestimation of the reconstruction efficiency up to a 7% at high p_T .

The selection efficiency accounts for the fraction of prompt charged particles removed from the candidate sample by the selection. The efficiency is estimated using simulation and a dedicated calibration procedure using a tag-and-probe technique applied to $\phi(1020) \rightarrow K^+K^-$ decays in data and simulation.

The signal purity is determined in simulation and corrected with background-enriched samples of data. Two independent samples dominated by fake tracks are constructed: using tracks with high ghost probability, and tracks which share their reconstructed VELO segment with a better fit-quality track. For secondary particles, the contributions from hadrons and electrons are studied separately. An enriched sample of hadrons from promptparticle decays, such as Λ baryons and K^0_S mesons, and hadrons produced in material interactions is obtained using tracks with a large estimated effective impact parameter with respect to the mean of the PV distribution. The abundance of electrons from γ conversions, which is considerable at low p_T , is validated using particle identification detectors.

Additionally, bin migration effects due to the resolution of the detector are found to have a negligible contribution to the measured yields. A minor correction is made since the mass of the particle is ignored in the expression $\eta = \eta_{\rm lab} - 0.465$ which is used to translate the pseudorapidity in the laboratory system to the centre-of-mass system of the nucleon-nucleon collision in pPb collisions.

Several sources of systematic uncertainty are considered. For candidates in the range $5 < p < 200$ GeV/c the track-finding efficiency carries an uncertainty due to the limited size of the calibration samples and the difference between hadron and muon

Table 1: Relative uncertainties for pPb and pp charged particle cross-sections. The interval indicates the minimum and the maximum value among the (η, p_T) bins. The systematic uncertainty due to luminosity is fully correlated across the bins. The other sources of systematic uncertainties are fully uncorrelated between different bins.

Uncertainty source	$pPb[\%]$ (forward)	$pPb[\%]$ (backward)	pp [%]
Track-finding efficiency	$1.5 - 5.0$	$1.5 - 5.0$	$1.6 - 5.3$
Detector occupancy	$0.0 - 2.8$	$0.6 - 2.9$	$0.1 - 1.6$
Particle composition	$0.4 - 4.1$	$0.4 - 4.6$	$0.3 - 2.4$
Selection efficiency	$0.7 - 2.2$	$0.7 - 3.0$	$1.0 - 1.7$
Purity	$0.1 - 1.8$	$0.1 -11.7$	$0.1 - 5.8$
Truth-matching	$0.0 - 0.1$	$0.0 - 0.1$	$0.1 - 0.2$
Luminosity	2.3	2.5	2.0
Statistical uncertainty	$0.0 - 0.6$	$0.0 - 1.0$	$0.0 - 1.1$
Total (in $d^2\sigma/d\eta dp_T$)	$3.0 - 6.7$	$3.3 - 14.5$	$2.8 - 8.7$
Total (in R_{pPb})	$4.2 - 9.2$	$4.4 - 16.9$	

material interactions. For candidates outside this range, a conservative 5% uncertainty is assigned based on the efficiency variation in adjacent intervals. An uncertainty is assigned accounting for the detector occupancy description, which is estimated considering alternative weights. The uncertainty due to imperfect knowledge of the relative particle composition is determined from a 30% variation in the relative abundances of particles obtained from simulation. The uncertainty on the selection efficiency originates primarily from the limited size of the calibration sample. For the purity, the systematic uncertainty is estimated from the background abundance in the background-enriched samples and the data-simulation discrepancy in the background fraction from the independent samples. This uncertainty has a large η and p_T dependence: while negligible in regions with a small background level, it is the dominant contribution for intervals with large background contributions. These intervals correspond to high p_T for fake tracks in pPb collisions in the backward region. The uncertainties are given in Table [1,](#page-5-0) where all uncertainties are treated as uncorrelated.

The measured prompt charged particle cross-sections for pp and pPb are presented in Fig. [1.](#page-6-0) The total uncertainty is the sum in quadrature of statistical, systematic, and luminosity uncertainties. On average 0.1035 ± 0.0029 charged particles (with $0.961 < p_T < 1.249 \,\text{GeV}/c$ and $3.0 < \eta < 3.5$) are produced in pp collisions, when scaled by the total inelastic nucleon-nucleon cross-section of 67.6 ± 0.6 mb at $\sqrt{s_{NN}} = 5$ TeV [\[48\]](#page-12-7). This is two orders of magnitude smaller than for pPb collisions, assuming the same total inelastic nucleon-nucleon cross-section. The cross-section for pp collisions at $\sqrt{s_{\rm NN}} = 5$ TeV is compared with the result at $\sqrt{s_{\text{NN}}}$ = 13 TeV [\[27\]](#page-11-3). Both results are consistent, showing an increase in the cross-section at 13 TeV of a factor 1 to 3, depending on p_T .

The result for R_{pPb} in different (η, p_T) intervals is presented in Fig. [2,](#page-7-0) where the uncertainties arise from statistical, systematic and luminosity sources. In the forward region, the measurement indicates a suppression of charged particle production in pPb collisions relative to that in pp collisions, which increases towards forward pseudorapidities. In the low p_T regime, R_{pPb} reaches values of about 0.3 in the most forward pseudorapidities.

Figure 1: Differential cross-section of prompt charged particle production as a function of p_T in different η intervals in (left) pp, (middle) forward pPb and (right) backward pPb collisions. The cross-section values for pp are scaled by the lead mass number $(A = 208)$ for comparison with the pPb cross-sections.

In the backward region, a significant enhancement of R_{pPb} is observed for $p_T > 1.5 \text{ GeV}/c$. This can be interpreted as Cronin enhancement [\[15\]](#page-10-6). The enhancement reaches a maximum at different p_T values depending on η , followed by a decreasing trend towards unity. This decrease is more pronounced in the most backward pseudorapidities. The maximum value of R_{pPb} is found to be ~ 1.3 and depends slightly on η .

The R_{pPb} measurement is compared in Fig. [2](#page-7-0) with predictions from phenomenological models covering the $p_T \gtrsim 1.5 \,\text{GeV}/c$ region. The prediction in Ref. [\[49\]](#page-12-8) (shaded green) is based on the nPDF set EPPS16 [\[10\]](#page-10-2) for the lead nucleus and the PDF set CT14 [\[50\]](#page-12-9) for the proton. The calculation also employs the parton-to-hadron fragmentation functions set DSS [\[51\]](#page-12-10). The prediction reproduces the data in the forward region although with large uncertainties. However, it fails to reproduce the R_{pPb} enhancement in the backward region for $p_T > 2 \text{ GeV}/c$.

The second prediction (violet) is based on the CGC effective field theory [\[20\]](#page-10-12). The model is only applicable to the saturation region at low- x and thus to forward rapidities. The predicted gradual decrease of R_{pPb} with η is observed in the data, although the prediction overestimates R_{pPb} in the lower p_T intervals. The prediction does not include an uncertainty estimation.

The third prediction (shaded orange) is a pQCD calculation within the high-twist factorisation formalism in the backward region [\[13,](#page-10-4)[52\]](#page-12-11). The calculation shows an enhancement due to incoherent multiple scattering inside the nucleus before and after the hard scattering, and reproduces the enhancement seen in pAu collisions in the backward region by the PHENIX experiment at $\sqrt{s_{NN}} = 200 \,\text{GeV}$ [\[25\]](#page-11-15). The prediction shows a p_{T} trend similar to data for $p_T > 3$ GeV/c in the most backward η interval, although it does not

Figure 2: Nuclear modification factor as a function of p_T in different η intervals for the (top) forward and (bottom) backward regions, compared with the predictions from Refs. [\[13,](#page-10-4) [20,](#page-10-12) [49,](#page-12-8) [52\]](#page-12-11). Vertical error bars correspond to statistical uncertainties, open boxes to uncorrelated systematic uncertainty and the filled box at $R_{pPb} = 1$ to the correlated uncertainty from the luminosity.

reproduce the data for the other intervals in the backward configuration.

Understanding the evolution of R_{pPb} with x and the momentum transfer Q^2 , is a critical point for the study of CNM effects. However, x and Q^2 are partonic quantities and cannot be directly measured. Instead, experimental proxies for x and $Q²$ [\[53\]](#page-12-12), defined as

$$
Q_{\text{exp}}^2 \equiv m^2 + p_{\text{T}}^2 \quad \text{and} \quad x_{\text{exp}} \equiv \frac{Q_{\text{exp}}}{\sqrt{s_{\text{NN}}}} e^{-\eta},\tag{3}
$$

are considered to compare the R_{pPb} results among different LHC experiments. Here, m is the mass of the produced particle and is taken as $m = 256 \text{ MeV}/c^2$, the average charged particle mass in pPb collisions determined with EPOS-LHC. The variable x_{exp} is approximately x for a two body scattering, and Q_{exp} is the transverse mass of the produced particle.

Figure [3](#page-8-0) shows the R_{pPb} evolution with x_{exp} for four Q_{exp}^2 intervals from this study and the results from the ALICE [\[21\]](#page-10-13) and CMS [\[22\]](#page-10-14) collaborations. Since the p_T binning is different among the three experiments, the Q_{exp}^2 ranges are selected to contain at least one p_T interval from each experiment. A consistent trend between this measurement in the forward region, the measurements in the central region from ALICE and CMS and the result in the backward region is observed for the four $Q_{\rm exp}^2$ intervals. The evolution of R_{pPb} with x_{\exp} is Q_{\exp}^2 -dependent.

In summary, the differential production cross-sections have been measured in p_T and η in summary, the untertainty production cross-sections have been measured in p_T and η intervals for prompt charged particles produced in pp and pPb collisions at $\sqrt{s_{NN}} = 5$ TeV. The measurement corresponds to $p > 2$ GeV/c and $0.2 < p_T < 8.0$ GeV/c prompt charged particles with $2.0 < \eta < 4.8$ in pp and $-5.2 < \eta < -2.5$ and $1.5 < \eta < 4.3$ in pPb collisions.

Figure 3: Evolution of the nuclear modification factor with x_{exp} from this study, ALICE [\[21\]](#page-10-13), and CMS [\[22\]](#page-10-14), for different Q_{exp}^2 ranges. Each plot includes all the R_{pPb} (η, p_T) intervals with a p_T centre within the Q_{exp}^2 range specified in the plot. Horizontal error bars account for the minimum and maximum x_{\exp} value for a given (η, p_T) interval. Vertical error bars correspond to statistical, systematic and luminosity (normalisation) uncertainties for LHCb (ALICE, CMS), added in quadrature.

This is the first determination of such cross-sections in pPb collisions in the forward and backward regions at the LHC, and the first measurement in pp collisions at $\sqrt{s_{NN}} = 5$ TeV. The total uncertainty is around 3% for most kinematic intervals both in pp and pPb collisions. As a result, the data place stringent constraints on non-perturbative QCD models in high-energy nuclear collisions.

The nuclear modification factor R_{pPb} is also determined and is one of the most precise to date. The total uncertainty, including the normalisation contribution, is below 5% for most of the (η, p_T) intervals. In the forward region, a suppression of the charged particle production is observed, especially for low p_T and the most forward η . In the backward region, the production of charged particles with $p_T > 1.5$ GeV/c is significantly enhanced. The R_{pPb} shape exhibits a clear pseudorapidity dependence. These data cannot be simultaneously described across the entire measured η range by nPDFs alone. Contrary to what is observed at central rapidity [\[18\]](#page-10-9), the forward data are inconsistent with CGC calculations at the lowest p_T . Multiple scattering calculations, which successfully reproduce PHENIX results [\[25\]](#page-11-15), fail to describe the backward region. These measurements provide strong constraints on nuclear PDFs at the lowest accessible x ranges, and show that additional, previously unconsidered mechanisms are required to provide a consistent description of particle production in nuclear collisions at the LHC.

Acknowledgements

We would like to thank to Hannu Paukkunen, Ilka Helenius, Heikki Mäntysaari and Zhong-Bo Kang, for providing LHCb specific theoretical predictions. We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MSHE (Russia); MICINN (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); A*MIDEX, ANR, Labex P2IO and OCEVU, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, Thousand Talents Program, and Sci. & Tech. Program of Guangzhou (China); RFBR, RSF and Yandex LLC (Russia); GVA, XuntaGal and GENCAT (Spain); the Royal Society and the Leverhulme Trust (United Kingdom).

References

- [1] A. B. Kaidalov and K. A. Ter-Martirosyan, Multihadron production at high energies in the model of quark gluon strings, Sov. J. Nucl. Phys. 40 (1984) 135.
- [2] A. Capella, U. Sukhatme, C.-I. Tan, and J. Tran Thanh Van, *Dual parton model*, [Physics Reports](https://doi.org/https://doi.org/10.1016/0370-1573(94)90064-7) 236 (1994) 225.
- [3] A. Buckley *et al., General-purpose event generators for LHC physics*, [Phys. Rept.](https://doi.org/10.1016/j.physrep.2011.03.005) 504 [\(2011\) 145,](https://doi.org/10.1016/j.physrep.2011.03.005) [arXiv:1101.2599](http://arxiv.org/abs/1101.2599).
- [4] D. d'Enterria et al., Constraints from the first LHC data on hadronic event generators for ultra-high energy cosmic-ray physics, [Astropart. Phys.](https://doi.org/10.1016/j.astropartphys.2011.05.002) 35 (2011) 98, [arXiv:1101.5596](http://arxiv.org/abs/1101.5596).
- [5] J. Albrecht et al., The Muon Puzzle in cosmic-ray induced air showers and its connection to the Large Hadron Collider, [Astrophys. Space Sci.](https://doi.org/10.1007/s10509-022-04054-5) 367 (2022) 27, [arXiv:2105.06148](http://arxiv.org/abs/2105.06148).
- [6] C. A. Salgado et al., Proton-Nucleus collisions at the LHC: scientific opportunities and requirements, J. Phys. **G39** [\(2012\) 015010,](https://doi.org/10.1088/0954-3899/39/1/015010) $arXiv:1105.3919$.
- [7] J. L. Albacete et al., Predictions for $p+Pb$ collisions at $\sqrt{s_{NN}} = 5$ TeV, [Int. J. Mod.](https://doi.org/10.1142/S0218301313300075) Phys. E22 [\(2013\) 1330007,](https://doi.org/10.1142/S0218301313300075) [arXiv:1301.3395](http://arxiv.org/abs/1301.3395).
- [8] J. L. Albacete *et al.*, Predictions for cold nuclear matter effects in $p+Pb$ collisions at $\sqrt{s_{NN}} = 8.16 \text{ TeV}$, [Nucl. Phys.](https://doi.org/10.1016/j.nuclphysa.2017.11.015) **A972** (2018) 18, [arXiv:1707.09973](http://arxiv.org/abs/1707.09973).
- [9] J. L. Nagle and W. A. Zajc, Small system collectivity in relativistic hadronic and nuclear collisions, [Ann. Rev. Nucl. Part. Sci.](https://doi.org/10.1146/annurev-nucl-101916-123209) 68 (2018) 211, $arXiv:1801.03477$.
- [10] K. J. Eskola, P. Paakkinen, H. Paukkunen, and C. A. Salgado, EPPS16: Nuclear par-ton distributions with LHC data, [Eur. Phys. J.](https://doi.org/10.1140/epjc/s10052-017-4725-9) $C77$ (2017) 163, $arXiv:1612.05741$.
- [11] K. Kovarik et al., nCTEQ15 - Global analysis of nuclear parton distributions with uncertainties in the CTEQ framework, Phys. Rev. **D93** [\(2016\) 085037,](https://doi.org/10.1103/PhysRevD.93.085037) [arXiv:1509.00792](http://arxiv.org/abs/1509.00792).
- [12] D. de Florian, R. Sassot, P. Zurita, and M. Stratmann, Global analysis of nuclear parton distributions, Phys. Rev. **D85** [\(2012\) 074028,](https://doi.org/10.1103/PhysRevD.85.074028) $arXiv:1112.6324$.
- [13] Z.-B. Kang, I. Vitev, and H. Xing, Multiple scattering effects on inclusive particle production in the large-x regime, Phys. Rev. $DS8$ [\(2013\) 054010,](https://doi.org/10.1103/PhysRevD.88.054010) $arXiv:1307.3557$.
- [14] A. Accardi, Cronin effect in proton nucleus collisions: A survey of theoretical models, [arXiv:hep-ph/0212148](http://arxiv.org/abs/hep-ph/0212148).
- [15] J. W. Cronin et al., Production of hadrons with large transverse momentum at 200, 300, and 400 GeV, Phys. Rev. D11 [\(1975\) 3105.](https://doi.org/10.1103/PhysRevD.11.3105)
- [16] H. Kowalski, T. Lappi, and R. Venugopalan, Nuclear enhancement of universal dynam-ics of high parton densities, [Phys. Rev. Lett.](https://doi.org/10.1103/PhysRevLett.100.022303) 100 (2008) 022303, [arXiv:0705.3047](http://arxiv.org/abs/0705.3047).
- [17] L. D. McLerran and R. Venugopalan, Gluon distribution functions for very large nuclei at small transverse momentum, Phys. Rev. D49 [\(1994\) 3352,](https://doi.org/10.1103/PhysRevD.49.3352) [arXiv:hep-ph/9311205](http://arxiv.org/abs/hep-ph/9311205).
- [18] ALICE collaboration, S. Acharya et al., Nuclear modification factor of light neutralmeson spectra up to high transverse momentum in p–Pb collisions at $\sqrt{s_{NN}}$ =8.16 TeV, Phys. Lett. B827 [\(2022\) 136943,](https://doi.org/10.1016/j.physletb.2022.136943) [arXiv:2104.03116](http://arxiv.org/abs/2104.03116).
- [19] F. Arleo, F. Cougoulic, and S. Peigné, Fully coherent energy loss effects on light hadron production in pA collisions, JHEP 09 [\(2020\) 190,](https://doi.org/10.1007/JHEP09(2020)190) arXiv: 2003.06337.
- [20] T. Lappi and H. Mäntysaari, *Single inclusive particle production at high energy* from HERA data to proton-nucleus collisions, Phys. Rev. D88 [\(2013\) 114020,](https://doi.org/10.1103/PhysRevD.88.114020) [arXiv:1309.6963](http://arxiv.org/abs/1309.6963).
- [21] ALICE collaboration, S. Acharya et al., Transverse momentum spectra and nuclear modification factors of charged particles in pp, p-Pb and Pb-Pb collisions at the LHC, JHEP 11 [\(2018\) 013,](https://doi.org/10.1007/JHEP11(2018)013) [arXiv:1802.09145](http://arxiv.org/abs/1802.09145).
- [22] CMS collaboration, V. Khachatryan et al., Charged-particle nuclear modification factors in PbPb and pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, JHEP 04 [\(2017\) 039,](https://doi.org/10.1007/JHEP04(2017)039) [arXiv:1611.01664](http://arxiv.org/abs/1611.01664).
- [23] ATLAS collaboration, G. Aad et al., Transverse momentum, rapidity, and central-ATLAS COMMODIATION, C. Add *et al.*, Transverse momentum, raptatly, and central-
ity dependence of inclusive charged-particle production in $\sqrt{s_{NN}} = 5.02$ TeV p + Pb collisions measured by the ATLAS experiment, [Phys. Lett.](https://doi.org/10.1016/j.physletb.2016.10.053) **B763** (2016) 313, [arXiv:1605.06436](http://arxiv.org/abs/1605.06436).
- [24] BRAHMS collaboration, I. Arsene et al., On the evolution of the nuclear modification Factors with rapidity and centrality in $d + Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV, [Phys.](https://doi.org/10.1103/PhysRevLett.93.242303) Rev. Lett. 93 [\(2004\) 242303,](https://doi.org/10.1103/PhysRevLett.93.242303) [arXiv:nucl-ex/0403005](http://arxiv.org/abs/nucl-ex/0403005).
- [25] PHENIX collaboration, C. Aidala et al., Nuclear-modification factor of charged hadrons at forward and backward rapidity in p+Al and p+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, Phys. Rev. $C101$ [\(2020\) 034910,](https://doi.org/10.1103/PhysRevC.101.034910) [arXiv:1906.09928](http://arxiv.org/abs/1906.09928).
- [26] PHOBOS collaboration, B. B. Back et al., Pseudorapidity dependence of charged hadron transverse momentum spectra in $d + Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV, [Phys.](https://doi.org/10.1103/PhysRevC.70.061901) Rev. C70 [\(2004\) 061901,](https://doi.org/10.1103/PhysRevC.70.061901) [arXiv:nucl-ex/0406017](http://arxiv.org/abs/nucl-ex/0406017).
- [27] LHCb collaboration, R. Aaij et al., Measurement of prompt charged-particle production EITCD conaboration, K. Aaif et al., Measurement of prompt charged-particle pr
in pp collisions at $\sqrt{s} = 13$ TeV, JHEP 01 [\(2022\) 166,](https://doi.org/10.1007/JHEP01(2022)166) [arXiv:2107.10090](http://arxiv.org/abs/2107.10090).
- [28] ALICE collaboration, S. Acharya et al., The ALICE definition of primary particles, [ALICE-PUBLIC-2017-005,](https://cds.cern.ch/record/2270008) CERN, 2017.
- [29] LHCb collaboration, A. A. Alves Jr. et al., The LHCb detector at the LHC, [JINST](https://doi.org/10.1088/1748-0221/3/08/S08005) 3 [\(2008\) S08005.](https://doi.org/10.1088/1748-0221/3/08/S08005)
- [30] LHCb collaboration, R. Aaij et al., *LHCb detector performance*, [Int. J. Mod. Phys.](https://doi.org/10.1142/S0217751X15300227) A30 [\(2015\) 1530022,](https://doi.org/10.1142/S0217751X15300227) [arXiv:1412.6352](http://arxiv.org/abs/1412.6352).
- [31] LHCb collaboration, R. Aaij et al., Precision luminosity measurements at LHCb, JINST 9 [\(2014\) P12005,](https://doi.org/10.1088/1748-0221/9/12/P12005) [arXiv:1410.0149](http://arxiv.org/abs/1410.0149).
- [32] T. Pierog et al., EPOS LHC: Test of collective hadronization with data measured at the CERN Large Hadron Collider, Phys. Rev. $C92$ [\(2015\) 034906,](https://doi.org/10.1103/PhysRevC.92.034906) [arXiv:1306.0121](http://arxiv.org/abs/1306.0121).
- [33] T. Sjöstrand, S. Mrenna, and P. Skands, A brief introduction to PYTHIA 8.1, [Comput.](https://doi.org/10.1016/j.cpc.2008.01.036) [Phys. Commun.](https://doi.org/10.1016/j.cpc.2008.01.036) 178 (2008) 852, [arXiv:0710.3820](http://arxiv.org/abs/0710.3820).
- [34] I. Belyaev et al., Handling of the generation of primary events in Gauss, the LHCb simulation framework, [J. Phys. Conf. Ser.](https://doi.org/10.1088/1742-6596/331/3/032047) 331 (2011) 032047.
- [35] D. J. Lange, The EvtGen particle decay simulation package, [Nucl. Instrum. Meth.](https://doi.org/10.1016/S0168-9002(01)00089-4) A462 [\(2001\) 152.](https://doi.org/10.1016/S0168-9002(01)00089-4)
- [36] Geant4 collaboration, S. Agostinelli et al., Geant4: A simulation toolkit, [Nucl. Instrum.](https://doi.org/10.1016/S0168-9002(03)01368-8) Meth. A506 [\(2003\) 250.](https://doi.org/10.1016/S0168-9002(03)01368-8)
- [37] Geant4 collaboration, J. Allison *et al., Geant4 developments and applications*, [IEEE](https://doi.org/10.1109/TNS.2006.869826) [Trans. Nucl. Sci.](https://doi.org/10.1109/TNS.2006.869826) 53 (2006) 270.
- [38] M. Clemencic et al., The LHCb simulation application, Gauss: Design, evolution and experience, [J. Phys. Conf. Ser.](https://doi.org/10.1088/1742-6596/331/3/032023) 331 (2011) 032023.
- [39] M. De Cian, S. Farry, P. Seyfert, and S. Stahl, Fast neural-net based fake track rejection in the LHCb reconstruction, [LHCb-PUB-2017-011,](http://cdsweb.cern.ch/search?p=LHCb-PUB-2017-011&f=reportnumber&action_search=Search&c=LHCb+Notes) 2017.
- [40] LHCb collaboration, R. Aaij et al., Measurement of the track reconstruction efficiency at LHCb, JINST 10 [\(2015\) P02007,](https://doi.org/10.1088/1748-0221/10/02/P02007) [arXiv:1408.1251](http://arxiv.org/abs/1408.1251).
- [41] ALICE collaboration, J. Adam et al., Multiplicity dependence of charged pion, kaon, and (anti)proton production at large transverse momentum in p-Pb collisions at \overline{C} [00 \overline{T}]. Let \overline{P} \overline{Z} (0016) 700 \overline{X} 1.4004.00550 $\sqrt{s_{NN}}$ = 5.02 TeV, [Phys. Lett.](https://doi.org/10.1016/j.physletb.2016.07.050) **B760** (2016) 720, [arXiv:1601.03658](http://arxiv.org/abs/1601.03658).
- [42] ALICE collaboration, J. Adam et al., Enhanced production of multi-strange hadrons in high-multiplicity proton-proton collisions, [Nature Phys.](https://doi.org/10.1038/nphys4111) 13 (2017) 535, [arXiv:1606.07424](http://arxiv.org/abs/1606.07424).
- [43] ALICE collaboration, J. Adam et al., Multi-strange baryon production in p-Pb EXECT COMMODIATION, 3. Adam et al., Matti-strange our gon production in p-
collisions at $\sqrt{s_{NN}}$ = 5.02 TeV, [Phys. Lett.](https://doi.org/10.1016/j.physletb.2016.05.027) **B758** (2016) 389, [arXiv:1512.07227](http://arxiv.org/abs/1512.07227).
- [44] ALICE collaboration, B. B. Abelev et al., Multiplicity Dependence of pion, kaon, proton and lambda production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, [Phys. Lett.](https://doi.org/10.1016/j.physletb.2013.11.020) B728 [\(2014\) 25,](https://doi.org/10.1016/j.physletb.2013.11.020) [arXiv:1307.6796](http://arxiv.org/abs/1307.6796).
- [45] LHCb collaboration, R. Aaij et al., Measurement of prompt hadron production EFFCD conaboration, K. Aaij et al., Measurement of prompt haaron production
ratios in pp collisions at $\sqrt{s} = 0.9$ and 7 TeV, [Eur. Phys. J.](https://doi.org/10.1140/epjc/s10052-012-2168-x) C72 (2012) 2168, [arXiv:1206.5160](http://arxiv.org/abs/1206.5160).
- [46] T. Sjöstrand et al., An introduction to PYTHIA 8.2, [Comput. Phys. Commun.](https://doi.org/10.1016/j.cpc.2015.01.024) 191 [\(2015\) 159,](https://doi.org/10.1016/j.cpc.2015.01.024) [arXiv:1410.3012](http://arxiv.org/abs/1410.3012).
- [47] C. Bierlich, Microscopic collectivity: The ridge and strangeness enhancement from string–string interactions, [Nucl. Phys.](https://doi.org/10.1016/j.nuclphysa.2018.07.015) A982 (2019) 499, [arXiv:1807.05271](http://arxiv.org/abs/1807.05271).
- [48] C. Loizides, J. Kamin, and D. d'Enterria, Improved Monte Carlo Glauber predictions at present and future nuclear colliders, Phys. Rev. **C97** [\(2018\) 054910,](https://doi.org/10.1103/PhysRevC.97.054910) [arXiv:1710.07098](http://arxiv.org/abs/1710.07098), [Erratum: Phys. Rev. C99, 019901 (2019)].
- [49] I. Helenius, K. J. Eskola, and H. Paukkunen, Probing the small-x nuclear gluon distributions with isolated photons at forward rapidities in $p+Pb$ collisions at the LHC, JHEP 09 [\(2014\) 138,](https://doi.org/10.1007/JHEP09(2014)138) [arXiv:1406.1689](http://arxiv.org/abs/1406.1689).
- [50] S. Dulat et al., New parton distribution functions from a global analysis of quantum chromodynamics, Phys. Rev. D93 [\(2016\) 033006,](https://doi.org/10.1103/PhysRevD.93.033006) [arXiv:1506.07443](http://arxiv.org/abs/1506.07443).
- [51] D. de Florian, R. Sassot, and M. Stratmann, Global analysis of fragmentation functions for pions and kaons and their uncertainties, Phys. Rev. D75 [\(2007\) 114010,](https://doi.org/10.1103/PhysRevD.75.114010) [arXiv:hep-ph/0703242](http://arxiv.org/abs/hep-ph/0703242).
- [52] Z.-B. Kang et al., Multiple scattering effects on heavy meson production in $p+A$ collisions at backward rapidity, [Phys. Lett.](https://doi.org/10.1016/j.physletb.2014.11.024) $B740$ (2015) 23, $arXiv:1409.2494$.
- [53] N. Armesto, *Nuclear shadowing*, J. Phys. **G32** [\(2006\) R367,](https://doi.org/10.1088/0954-3899/32/11/R01) [arXiv:hep-ph/0604108](http://arxiv.org/abs/hep-ph/0604108).

LHCb collaboration

R. Aaij 32 , C. Abellán Beteta 50 , T. Ackernley 60 , B. Adeva 46 , M. Adinolfi 54 , H. Afsharnia 9 , C.A. Aidala⁸⁶, S. Aiola²⁵, Z. Ajaltouni⁹, S. Akar⁶⁵, J. Albrecht¹⁵, F. Alessio⁴⁸, M. Alexander⁵⁹, A. Alfonso Albero⁴⁵, Z. Aliouche⁶², G. Alkhazov³⁸, P. Alvarez Cartelle⁵⁵, S. Amato², J.L. Amey⁵⁴, Y. Amhis¹¹, L. An⁴⁸, L. Anderlini²², A. Andreianov³⁸, M. Andreotti²¹, F. Archilli¹⁷, A. Artamonov⁴⁴, M. Artuso⁶⁸, K. Arzymatov⁴², E. Aslanides¹⁰, M. Atzeni⁵⁰, B. Audurier¹², S. Bachmann¹⁷, M. Bachmayer⁴⁹, J.J. Back⁵⁶, P. Baladron Rodriguez⁴⁶, V. Balagura¹², W. Baldini²¹, J. Baptista Leite¹, R.J. Barlow⁶², S. Barsuk¹¹, W. Barter⁶¹, M. Bartolini^{24,h}, F. Baryshnikov⁸³, J.M. Basels¹⁴, G. Bassi²⁹, B. Batsukh⁶⁸, A. Battig¹⁵, A. Bay⁴⁹, M. Becker¹⁵, F. Bedeschi²⁹, I. Bediaga¹, A. Beiter⁶⁸, V. Belavin⁴², S. Belin²⁷, V. Bellee⁴⁹, K. Belous⁴⁴, I. Belov⁴⁰, I. Belyaev⁴¹, G. Bencivenni²³, E. Ben-Haim¹³, A. Berezhnoy⁴⁰, R. Bernet⁵⁰, D. Berninghoff¹⁷, H.C. Bernstein⁶⁸, C. Bertella⁴⁸, A. Bertolin²⁸, C. Betancourt⁵⁰, F. Betti⁴⁸, Ia. Bezshyiko⁵⁰, S. Bhasin⁵⁴, J. Bhom³⁵, L. Bian⁷³, M.S. Bieker¹⁵, S. Bifani⁵³, P. Billoir¹³, M. Birch⁶¹, F.C.R. Bishop⁵⁵, A. Bitadze⁶², A. Bizzeti^{22,k}, M. Bjørn⁶³, M.P. Blago⁴⁸, T. Blake⁵⁶, F. Blanc⁴⁹, S. Blusk⁶⁸, D. Bobulska⁵⁹, J.A. Boelhauve¹⁵, O. Boente Garcia⁴⁶, T. Boettcher⁶⁵, A. Boldyrev⁸², A. Bondar⁴³, N. Bondar^{38,48}, S. Borghi⁶², M. Borisyak⁴², M. Borsato¹⁷, J.T. Borsuk³⁵, S.A. Bouchiba⁴⁹, T.J.V. Bowcock⁶⁰, A. Boyer⁴⁸, C. Bozzi²¹, M.J. Bradley⁶¹, S. Braun⁶⁶, A. Brea Rodriguez⁴⁶, M. Brodski⁴⁸, J. Brodzicka³⁵, A. Brossa Gonzalo⁵⁶, D. Brundu²⁷, A. Buonaura⁵⁰, A.T. Burke⁶², C. Burr⁴⁸, A. Bursche⁷², A. Butkevich³⁹, J.S. Butter³², J. Buytaert⁴⁸, W. Byczynski⁴⁸, S. Cadeddu²⁷, H. Cai⁷³, R. Calabrese^{21,f}, L. Calefice^{15,13}, L. Calero Diaz²³, S. Cali²³, R. Calladine⁵³, M. Calvi^{26,j}, M. Calvo Gomez⁸⁵, P. Camargo Magalhaes⁵⁴, P. Campana²³, A.F. Campoverde Quezada⁶, S. Capelli^{26,j}, L. Capriotti^{20,d}, A. Carbone^{20,d}, G. Carboni³¹, R. Cardinale^{24,h}, A. Cardini²⁷, I. Carli⁴, P. Carniti^{26,j}, L. Carus¹⁴, K. Carvalho Akiba³², A. Casais Vidal⁴⁶, G. Casse⁶⁰, M. Cattaneo⁴⁸, G. Cavallero⁴⁸, S. Celani⁴⁹, J. Cerasoli¹⁰, A.J. Chadwick⁶⁰, M.G. Chapman⁵⁴, M. Charles¹³, Ph. Charpentier⁴⁸, G. Chatzikonstantinidis⁵³, C.A. Chavez Barajas⁶⁰, M. Chefdeville⁸, C. Chen³, S. Chen⁴, A. Chernov³⁵, V. Chobanova⁴⁶, S. Cholak⁴⁹, M. Chrzaszcz³⁵, A. Chubykin³⁸, V. Chulikov³⁸, P. Ciambrone²³, M.F. Cicala⁵⁶, X. Cid Vidal⁴⁶, G. Ciezarek⁴⁸, P.E.L. Clarke⁵⁸, M. Clemencic⁴⁸, H.V. Cliff⁵⁵, J. Closier⁴⁸, J.L. Cobbledick⁶², V. Coco⁴⁸, J.A.B. Coelho¹¹, J. Cogan¹⁰, E. Cogneras⁹, L. Cojocariu³⁷, P. Collins⁴⁸, T. Colombo⁴⁸, L. Congedo^{19,c}, A. Contu²⁷, N. Cooke⁵³, G. Coombs⁵⁹, I. Corredoira⁴⁶, G. Corti⁴⁸, C.M. Costa Sobral⁵⁶, B. Couturier⁴⁸, D.C. Craik⁶⁴, J. Crkovská⁶⁷, M. Cruz Torres¹, R. Currie⁵⁸, C.L. Da Silva⁶⁷, S. Dadabaev⁸³, L. Dai⁷¹, E. Dall'Occo¹⁵, J. Dalseno⁴⁶, C. D'Ambrosio⁴⁸, A. Danilina⁴¹, P. d'Argent⁴⁸, J.E. Davies⁶², A. Davis⁶², O. De Aguiar Francisco⁶², K. De Bruyn⁷⁹, S. De Capua⁶², M. De Cian⁴⁹, J.M. De Miranda¹, L. De Paula², M. De Serio^{19,c}, D. De Simone⁵⁰, P. De Simone²³, J.A. de Vries⁸⁰, C.T. Dean⁶⁷, D. Decamp⁸, L. Del Buono¹³, B. Delaney⁵⁵, H.-P. Dembinski¹⁵, A. Dendek³⁴, V. Denysenko⁵⁰, D. Derkach⁸², O. Deschamps⁹, F. Desse¹¹, F. Dettori^{27,e}, B. Dey⁷⁷, A. Di Cicco²³, P. Di Nezza²³, S. Didenko⁸³, L. Dieste Maronas⁴⁶, H. Dijkstra⁴⁸, V. Dobishuk⁵², A.M. Donohoe¹⁸, F. Dordei²⁷, A.C. dos Reis¹, L. Douglas⁵⁹, A. Dovbnya⁵¹, A.G. Downes⁸, K. Dreimanis⁶⁰, M.W. Dudek³⁵, L. Dufour⁴⁸, V. Duk⁷⁸, P. Durante⁴⁸, J.M. Durham⁶⁷, D. Dutta⁶², A. Dziurda³⁵, A. Dzyuba³⁸, S. Easo⁵⁷, U. Egede⁶⁹, V. Egorychev⁴¹, S. Eidelman^{43,*v*}, S. Eisenhardt⁵⁸, S. Ek-In⁴⁹, L. Eklund^{59,w}, S. Ely⁶⁸, A. Ene³⁷, E. Epple⁶⁷, S. Escher¹⁴, J. Eschle⁵⁰, S. Esen¹³, T. Evans⁴⁸, A. Falabella²⁰, J. Fan³, Y. Fan⁶, B. Fang⁷³, S. Farry⁶⁰, D. Fazzini^{26,j}, M. Féo⁴⁸, A. Fernandez Prieto⁴⁶, A.D. Fernez⁶⁶, F. Ferrari^{20,d}, L. Ferreira Lopes⁴⁹, F. Ferreira Rodrigues², S. Ferreres Sole³², M. Ferrillo⁵⁰, M. Ferro-Luzzi⁴⁸, S. Filippov³⁹, R.A. Fini¹⁹, M. Fiorini^{21,f}, M. Firlej³⁴, K.M. Fischer⁶³, D.S. Fitzgerald⁸⁶, C. Fitzpatrick⁶², T. Fiutowski³⁴, A. Fkiaras⁴⁸,

- F. Fleuret¹², M. Fontana¹³, F. Fontanelli^{24,h}, R. Forty⁴⁸, V. Franco Lima⁶⁰, M. Franco Sevilla⁶⁶,
- M. Frank⁴⁸, E. Franzoso²¹, G. Frau¹⁷, C. Frei⁴⁸, D.A. Friday⁵⁹, J. Fu²⁵, Q. Fuehring¹⁵,

W. Funk⁴⁸, E. Gabriel³², T. Gaintseva⁴², A. Gallas Torreira⁴⁶, D. Galli^{20,d}, S. Gambetta^{58,48}, Y. Gan³, M. Gandelman², P. Gandini²⁵, Y. Gao⁵, M. Garau²⁷, L.M. Garcia Martin⁵⁶, P. Garcia Moreno⁴⁵, J. García Pardiñas^{26,j}, B. Garcia Plana⁴⁶, F.A. Garcia Rosales¹², L. Garrido⁴⁵, C. Gaspar⁴⁸, R.E. Geertsema³², D. Gerick¹⁷, L.L. Gerken¹⁵, E. Gersabeck⁶², M. Gersabeck⁶², T. Gershon⁵⁶, D. Gerstel¹⁰, Ph. Ghez⁸, V. Gibson⁵⁵, H.K. Giemza³⁶, M. Giovannetti^{23,p}, A. Gioventù⁴⁶, P. Gironella Gironell⁴⁵, L. Giubega³⁷, C. Giugliano^{21,f,48}, K. Gizdov⁵⁸, E.L. Gkougkousis⁴⁸, V.V. Gligorov¹³, C. Göbel⁷⁰, E. Golobardes⁸⁵, D. Golubkov⁴¹, A. Golutvin^{61,83}, A. Gomes^{1,a}, S. Gomez Fernandez⁴⁵, F. Goncalves Abrantes⁶³, M. Goncerz³⁵, G. Gong³, P. Gorbounov⁴¹, I.V. Gorelov⁴⁰, C. Gotti²⁶, E. Govorkova⁴⁸, J.P. Grabowski¹⁷, T. Grammatico¹³, L.A. Granado Cardoso⁴⁸, E. Graugés⁴⁵, E. Graverini⁴⁹, G. Graziani²², A. Grecu³⁷, L.M. Greeven³², P. Griffith^{21,f}, L. Grillo⁶², S. Gromov⁸³, B.R. Gruberg Cazon⁶³, C. Gu³, M. Guarise²¹, P. A. Günther¹⁷, E. Gushchin³⁹, A. Guth¹⁴, Y. Guz⁴⁴, T. Gys⁴⁸, T. Hadavizadeh⁶⁹, G. Haefeli⁴⁹, C. Haen⁴⁸, J. Haimberger⁴⁸, T. Halewood-leagas⁶⁰, P.M. Hamilton⁶⁶, J.P. Hammerich⁶⁰, Q. Han⁷, X. Han¹⁷, T.H. Hancock⁶³, S. Hansmann-Menzemer¹⁷, N. Harnew⁶³, T. Harrison⁶⁰, C. Hasse⁴⁸, M. Hatch⁴⁸, J. He^{6,6}, M. Hecker⁶¹, K. Heijhoff³², K. Heinicke¹⁵, A.M. Hennequin⁴⁸, K. Hennessy⁶⁰, L. Henry⁴⁸, J. Heuel¹⁴, A. Hicheur², D. Hill⁴⁹, M. Hilton⁶², S.E. Hollitt¹⁵, J. Hu¹⁷, J. Hu⁷², W. Hu⁷, X. Hu³, W. Huang⁶, X. Huang⁷³, W. Hulsbergen³², R.J. Hunter⁵⁶, M. Hushchyn⁸², D. Hutchcroft⁶⁰, D. Hynds³², P. Ibis¹⁵, M. Idzik³⁴, D. Ilin³⁸, P. Ilten⁶⁵, A. Inglessi³⁸, A. Ishteev⁸³, K. Ivshin³⁸, R. Jacobsson⁴⁸, S. Jakobsen⁴⁸, E. Jans³², B.K. Jashal⁴⁷, A. Jawahery⁶⁶, V. Jevtic¹⁵, F. Jiang³, M. John⁶³, D. Johnson⁴⁸, C.R. Jones⁵⁵, T.P. Jones⁵⁶, B. Jost⁴⁸, N. Jurik⁴⁸, S. Kandybei⁵¹, Y. Kang³, M. Karacson⁴⁸, M. Karpov⁸², F. Keizer⁴⁸, M. Kenzie⁵⁶, T. Ketel³³, B. Khanji¹⁵, A. Kharisova⁸⁴, S. Kholodenko⁴⁴, T. Kirn¹⁴, V.S. Kirsebom⁴⁹, O. Kitouni⁶⁴, S. Klaver³², K. Klimaszewski³⁶, S. Koliiev⁵², A. Kondybayeva⁸³, A. Konoplyannikov⁴¹, P. Kopciewicz³⁴, R. Kopecna¹⁷, P. Koppenburg³², M. Korolev⁴⁰, I. Kostiuk^{32,52}, O. Kot⁵², S. Kotriakhova^{21,38}, P. Kravchenko³⁸, L. Kravchuk³⁹, R.D. Krawczyk⁴⁸, M. Kreps⁵⁶, F. Kress⁶¹, S. Kretzschmar¹⁴, P. Krokovny^{43,*v*}, W. Krupa³⁴, W. Krzemien³⁶, W. Kucewicz^{35,*t*}, M. Kucharczyk³⁵, V. Kudryavtsev^{43,v}, H.S. Kuindersma^{32,33}, G.J. Kunde⁶⁷, T. Kvaratskheliya⁴¹, D. Lacarrere⁴⁸, G. Lafferty⁶², A. Lai²⁷, A. Lampis²⁷, D. Lancierini⁵⁰, J.J. Lane⁶², R. Lane⁵⁴, G. Lanfranchi²³, C. Langenbruch¹⁴, J. Langer¹⁵, O. Lantwin⁵⁰, T. Latham⁵⁶, F. Lazzari^{29,q}, R. Le Gac¹⁰, S.H. Lee⁸⁶, R. Lefèvre⁹, A. Leflat⁴⁰, S. Legotin⁸³, O. Leroy¹⁰, T. Lesiak³⁵, B. Leverington¹⁷, H. Li⁷², L. Li⁶³, P. Li¹⁷, S. Li⁷, Y. Li⁴, Y. Li⁴, Z. Li⁶⁸, X. Liang⁶⁸, T. Lin⁶¹, R. Lindner⁴⁸, V. Lisovskyi¹⁵, R. Litvinov²⁷, G. Liu⁷², H. Liu⁶, S. Liu⁴, A. Loi²⁷, J. Lomba Castro⁴⁶, I. Longstaff⁵⁹, J.H. Lopes², S. Lopez Solino⁴⁶, G.H. Lovell⁵⁵, Y. Lu⁴, D. Lucchesi^{28,*l*}, S. Luchuk³⁹, M. Lucio Martinez³², V. Lukashenko^{32,52}, Y. Luo³, A. Lupato⁶², E. Luppi^{21,f}, O. Lupton⁵⁶, A. Lusiani^{29,*m*}, X. Lyu⁶, L. Ma⁴, R. Ma⁶, S. Maccolini^{20,d}, F. Machefert¹¹, F. Maciuc³⁷, V. Macko⁴⁹, P. Mackowiak¹⁵, S. Maddrell-Mander⁵⁴, O. Madejczyk³⁴, L.R. Madhan Mohan⁵⁴, O. Maev³⁸, A. Maevskiy⁸², D. Maisuzenko³⁸, M.W. Majewski³⁴, J.J. Malczewski³⁵, S. Malde⁶³, B. Malecki⁴⁸, A. Malinin⁸¹, T. Maltsev^{43,*v*}, H. Malygina¹⁷, G. Manca^{27,e}, G. Mancinelli¹⁰, D. Manuzzi^{20,d}, D. Marangotto^{25,i}, J. Maratas^{9,s}, J.F. Marchand⁸, U. Marconi²⁰, S. Mariani^{22,g}, C. Marin Benito⁴⁸, M. Marinangeli⁴⁹, J. Marks¹⁷, A.M. Marshall⁵⁴, P.J. Marshall⁶⁰, G. Martellotti³⁰, L. Martinazzoli^{48,j}, M. Martinelli^{26,j}, D. Martinez Santos⁴⁶, F. Martinez Vidal⁴⁷, A. Massafferri¹, M. Materok¹⁴, R. Matev⁴⁸, A. Mathad⁵⁰, Z. Mathe⁴⁸, V. Matiunin⁴¹, C. Matteuzzi²⁶, K.R. Mattioli⁸⁶, A. Mauri³², E. Maurice¹², J. Mauricio⁴⁵, M. Mazurek⁴⁸, M. McCann⁶¹, L. Mcconnell¹⁸, T.H. Mcgrath⁶², A. McNab⁶², R. McNulty¹⁸, J.V. Mead⁶⁰, B. Meadows⁶⁵, G. Meier¹⁵, N. Meinert⁷⁶, D. Melnychuk³⁶, S. Meloni^{26,j}, M. Merk^{32,80}, A. Merli²⁵, L. Meyer Garcia², M. Mikhasenko⁴⁸, D.A. Milanes⁷⁴, E. Millard⁵⁶, M. Milovanovic⁴⁸, M.-N. Minard⁸, A. Minotti²¹, L. Minzoni^{21,f}, S.E. Mitchell⁵⁸, B. Mitreska⁶², D.S. Mitzel⁴⁸, A. Mödden ¹⁵, R.A. Mohammed⁶³, R.D. Moise⁶¹, T. Mombächer⁴⁶, I.A. Monroy⁷⁴, S. Monteil⁹, M. Morandin²⁸, G. Morello²³, M.J. Morello^{29,*m*},

J. Moron³⁴, A.B. Morris⁷⁵, A.G. Morris⁵⁶, R. Mountain⁶⁸, H. Mu³, F. Muheim^{58,48}, M. Mulder⁴⁸, D. Müller⁴⁸, K. Müller⁵⁰, C.H. Murphy⁶³, D. Murray⁶², P. Muzzetto^{27,48}, P. Naik⁵⁴, T. Nakada⁴⁹, R. Nandakumar⁵⁷, T. Nanut⁴⁹, I. Nasteva², M. Needham⁵⁸, I. Neri²¹, N. Neri^{25,*i*}, S. Neubert⁷⁵, N. Neufeld⁴⁸, R. Newcombe⁶¹, T.D. Nguyen⁴⁹, C. Nguyen-Mau^{49,x}, E.M. Niel¹¹, S. Nieswand¹⁴, N. Nikitin⁴⁰, N.S. Nolte⁶⁴, C. Normand⁸, C. Nunez⁸⁶, A. Oblakowska-Mucha³⁴, V. Obraztsov⁴⁴, D.P. O'Hanlon⁵⁴, R. Oldeman^{27,e}, M.E. Olivares⁶⁸, C.J.G. Onderwater⁷⁹, R.H. O'neil⁵⁸, A. Ossowska³⁵, J.M. Otalora Goicochea², T. Ovsiannikova⁴¹, P. Owen⁵⁰, A. Oyanguren⁴⁷, B. Pagare⁵⁶, P.R. Pais⁴⁸, T. Pajero⁶³, A. Palano¹⁹, M. Palutan²³, Y. Pan⁶², G. Panshin⁸⁴, A. Papanestis⁵⁷, M. Pappagallo^{19,c}, L.L. Pappalardo^{21,f}, C. Pappenheimer⁶⁵, W. Parker⁶⁶, C. Parkes⁶², C.J. Parkinson⁴⁶, B. Passalacqua²¹, G. Passaleva²², A. Pastore¹⁹, M. Patel⁶¹, C. Patrignani^{20,d}, C.J. Pawley⁸⁰, A. Pearce⁴⁸, A. Pellegrino³², M. Pepe Altarelli⁴⁸, S. Perazzini²⁰, D. Pereima⁴¹, P. Perret⁹, M. Petric^{59,48}, K. Petridis⁵⁴, A. Petrolini^{24,h}, A. Petrov⁸¹, S. Petrucci⁵⁸, M. Petruzzo²⁵, T.T.H. Pham⁶⁸, A. Philippov⁴², L. Pica^{29,*m*}, M. Piccini⁷⁸, B. Pietrzyk⁸, G. Pietrzyk⁴⁹, M. Pili⁶³, D. Pinci³⁰, F. Pisani⁴⁸, Resmi P.K¹⁰, V. Placinta³⁷, J. Plews⁵³, M. Plo Casasus⁴⁶, F. Polci¹³, M. Poli Lener²³, M. Poliakova⁶⁸, A. Poluektov¹⁰, N. Polukhina^{83,*u*}, I. Polyakov⁶⁸, E. Polycarpo², G.J. Pomery⁵⁴, S. Ponce⁴⁸, D. Popov^{6,48}, S. Popov⁴², S. Poslavskii⁴⁴, K. Prasanth³⁵, L. Promberger⁴⁸, C. Prouve⁴⁶, V. Pugatch⁵², H. Pullen⁶³, G. Punzi^{29,n}, H. Qi³, W. Qian⁶, J. Qin⁶, N. Qin³, R. Quagliani¹³, B. Quintana⁸, N.V. Raab¹⁸, R.I. Rabadan Trejo¹⁰, B. Rachwal³⁴, J.H. Rademacker⁵⁴, M. Rama²⁹, M. Ramos Pernas⁵⁶, M.S. Rangel², F. Ratnikov^{42,82}, G. Raven³³, M. Reboud⁸, F. Redi⁴⁹, F. Reiss⁶², C. Remon Alepuz⁴⁷, Z. Ren³, V. Renaudin⁶³, R. Ribatti²⁹, S. Ricciardi⁵⁷, K. Rinnert⁶⁰, P. Robbe¹¹, G. Robertson⁵⁸, A.B. Rodrigues⁴⁹, E. Rodrigues⁶⁰, J.A. Rodriguez Lopez⁷⁴, E.R.R. Rodriguez Rodriguez⁴⁶, A. Rollings⁶³, P. Roloff⁴⁸, V. Romanovskiy⁴⁴, M. Romero Lamas⁴⁶, A. Romero Vidal⁴⁶, J.D. Roth⁸⁶, M. Rotondo²³, M.S. Rudolph⁶⁸, T. Ruf⁴⁸, J. Ruiz Vidal⁴⁷, A. Ryzhikov⁸², J. Ryzka³⁴, J.J. Saborido Silva⁴⁶, N. Sagidova³⁸, N. Sahoo⁵⁶, B. Saitta^{27,e}, M. Salomoni⁴⁸, C. Sanchez Gras³², R. Santacesaria³⁰, C. Santamarina Rios⁴⁶, M. Santimaria²³, E. Santovetti^{31,p}, D. Saranin⁸³, G. Sarpis¹⁴, M. Sarpis⁷⁵, A. Sarti³⁰, C. Satriano^{30,o}, A. Satta³¹, M. Saur¹⁵, D. Savrina^{41,40}, H. Sazak⁹, L.G. Scantlebury Smead⁶³, A. Scarabotto¹³, S. Schael¹⁴, M. Schiller⁵⁹, H. Schindler⁴⁸, M. Schmelling¹⁶, B. Schmidt⁴⁸, O. Schneider⁴⁹, A. Schopper⁴⁸, M. Schubiger³², S. Schulte⁴⁹, M.H. Schune¹¹, R. Schwemmer⁴⁸, B. Sciascia²³, S. Sellam⁴⁶, A. Semennikov⁴¹, M. Senghi Soares³³, A. Sergi^{24,h}, N. Serra⁵⁰, L. Sestini²⁸, A. Seuthe¹⁵, P. Seyfert⁴⁸, Y. Shang⁵, D.M. Shangase⁸⁶, M. Shapkin⁴⁴, I. Shchemerov⁸³, L. Shchutska⁴⁹, T. Shears⁶⁰, L. Shekhtman^{43,*v*}, Z. Shen⁵, V. Shevchenko⁸¹, E.B. Shields^{26,j}, E. Shmanin⁸³, J.D. Shupperd⁶⁸, B.G. Siddi²¹, R. Silva Coutinho⁵⁰, G. Simi²⁸, S. Simone^{19,c}, N. Skidmore⁶², T. Skwarnicki⁶⁸, M.W. Slater⁵³, I. Slazyk^{21,f}, J.C. Smallwood⁶³, J.G. Smeaton⁵⁵, A. Smetkina⁴¹, E. Smith⁵⁰, M. Smith⁶¹, A. Snoch³², M. Soares²⁰, L. Soares Lavra⁹, M.D. Sokoloff⁶⁵, F.J.P. Soler⁵⁹, A. Solovev³⁸, I. Solovyev³⁸, F.L. Souza De Almeida², B. Souza De Paula², B. Spaan¹⁵, E. Spadaro Norella^{25,*i*}, P. Spradlin⁵⁹, F. Stagni⁴⁸, M. Stahl⁶⁵, S. Stahl⁴⁸, P. Stefko⁴⁹, O. Steinkamp^{50,83}, O. Stenyakin⁴⁴, H. Stevens¹⁵, S. Stone⁶⁸, M.E. Stramaglia⁴⁹, M. Straticiuc³⁷, D. Strekalina⁸³, F. Suljik⁶³, J. Sun²⁷, L. Sun⁷³, Y. Sun⁶⁶, P. Svihra⁶², P.N. Swallow⁵³, K. Swientek³⁴, A. Szabelski³⁶, T. Szumlak³⁴, M. Szymanski⁴⁸, S. Taneja⁶², A.R. Tanner⁵⁴, A. Terentev⁸³, F. Teubert⁴⁸, E. Thomas⁴⁸, D.J.D. Thompson⁵³, K.A. Thomson⁶⁰, V. Tisserand⁹, S. T'Jampens⁸, M. Tobin⁴, L. Tomassetti^{21,f}, D. Torres Machado¹, D.Y. Tou¹³, M.T. Tran⁴⁹, E. Trifonova⁸³, C. Trippl⁴⁹, G. Tuci^{29,*n*}, A. Tully⁴⁹, N. Tuning^{32,48}, A. Ukleja³⁶, D.J. Unverzagt¹⁷, E. Ursov⁸³, A. Usachov³², A. Ustyuzhanin^{42,82}, U. Uwer¹⁷, A. Vagner⁸⁴, V. Vagnoni²⁰, A. Valassi⁴⁸, G. Valenti²⁰, N. Valls Canudas⁸⁵, M. van Beuzekom³², M. Van Dijk⁴⁹, E. van Herwijnen⁸³, C.B. Van Hulse¹⁸, M. van Veghel⁷⁹, R. Vazquez Gomez⁴⁵, P. Vazquez Regueiro⁴⁶, C. Vázquez Sierra⁴⁸, S. Vecchi²¹,

J.J. Velthuis⁵⁴, M. Veltri^{22, r}, A. Venkateswaran⁶⁸, M. Veronesi³², M. Vesterinen⁵⁶, D. Vieira⁶⁵,

M. Vieites Diaz⁴⁹, H. Viemann⁷⁶, X. Vilasis-Cardona⁸⁵, E. Vilella Figueras⁶⁰, A. Villa²⁰,

P. Vincent¹³, D. Vom Bruch¹⁰, A. Vorobyev³⁸, V. Vorobyev^{43,*v*}, N. Voropaev³⁸, K. Vos⁸⁰,

R. Waldi¹⁷, J. Walsh²⁹, C. Wang¹⁷, J. Wang⁵, J. Wang⁴, J. Wang³, J. Wang⁷³, M. Wang³,

R. Wang⁵⁴, Y. Wang⁷, Z. Wang⁵⁰, Z. Wang³, H.M. Wark⁶⁰, N.K. Watson⁵³, S.G. Weber¹³,

D. Websdale⁶¹, C. Weisser⁶⁴, B.D.C. Westhenry⁵⁴, D.J. White⁶², M. Whitehead⁵⁴,

D. Wiedner¹⁵, G. Wilkinson⁶³, M. Wilkinson⁶⁸, I. Williams⁵⁵, M. Williams⁶⁴, M.R.J. Williams⁵⁸,

F.F. Wilson⁵⁷, W. Wislicki³⁶, M. Witek³⁵, L. Witola¹⁷, G. Wormser¹¹, S.A. Wotton⁵⁵, H. Wu⁶⁸,

K. Wyllie⁴⁸, Z. Xiang⁶, D. Xiao⁷, Y. Xie⁷, A. Xu⁵, J. Xu⁶, L. Xu³, M. Xu⁷, Q. Xu⁶, Z. Xu⁵,

Z. Xu^6 , D. Yang³, S. Yang⁶, Y. Yang⁶, Z. Yang³, Z. Yang⁶⁶, Y. Yao⁶⁸, L.E. Yeomans⁶⁰, H. Yin⁷,

J. Yu⁷¹, X. Yuan⁶⁸, O. Yushchenko⁴⁴, E. Zaffaroni⁴⁹, M. Zavertyaev^{16,*u*}, M. Zdybal³⁵,

O. Zenaiev⁴⁸, M. Zeng³, D. Zhang⁷, L. Zhang³, S. Zhang⁷¹, S. Zhang⁵, Y. Zhang⁵, Y. Zhang⁶³,

A. Zharkova⁸³, A. Zhelezov¹⁷, Y. Zheng⁶, X. Zhou⁶, Y. Zhou⁶, X. Zhu³, Z. Zhu⁶, V. Zhukov^{14,40}, J.B. Zonneveld⁵⁸, Q. Zou⁴, S. Zucchelli^{20,d}, D. Zuliani²⁸, G. Zunica⁶².

Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

Center for High Energy Physics, Tsinghua University, Beijing, China

Institute Of High Energy Physics (IHEP), Beijing, China

School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

University of Chinese Academy of Sciences, Beijing, China

Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China

Univ. Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France

Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

¹³LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France

I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

School of Physics, University College Dublin, Dublin, Ireland

INFN Sezione di Bari, Bari, Italy

INFN Sezione di Bologna, Bologna, Italy

INFN Sezione di Ferrara, Ferrara, Italy

INFN Sezione di Firenze, Firenze, Italy

INFN Laboratori Nazionali di Frascati, Frascati, Italy

INFN Sezione di Genova, Genova, Italy

INFN Sezione di Milano, Milano, Italy

INFN Sezione di Milano-Bicocca, Milano, Italy

INFN Sezione di Cagliari, Monserrato, Italy

Universita degli Studi di Padova, Universita e INFN, Padova, Padova, Italy

INFN Sezione di Pisa, Pisa, Italy

INFN Sezione di Roma La Sapienza, Roma, Italy

INFN Sezione di Roma Tor Vergata, Roma, Italy

Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands

Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands

AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland

Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland National Center for Nuclear Research (NCBJ), Warsaw, Poland

Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania

Petersburg Nuclear Physics Institute NRC Kurchatov Institute (PNPI NRC KI), Gatchina, Russia ³⁹Institute for Nuclear Research of the Russian Academy of Sciences (INR RAS), Moscow, Russia

Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia

Institute of Theoretical and Experimental Physics NRC Kurchatov Institute (ITEP NRC KI), Moscow, Russia

Yandex School of Data Analysis, Moscow, Russia

Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia

Institute for High Energy Physics NRC Kurchatov Institute (IHEP NRC KI), Protvino, Russia, Protvino, Russia

ICCUB, Universitat de Barcelona, Barcelona, Spain

 46 Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela. Santiago de Compostela, Spain

Instituto de Fisica Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain European Organization for Nuclear Research (CERN), Geneva, Switzerland

 49 Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland Physik-Institut, Universität Zürich, Zürich, Switzerland

NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine

Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine

University of Birmingham, Birmingham, United Kingdom

H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom

Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom

Department of Physics, University of Warwick, Coventry, United Kingdom

STFC Rutherford Appleton Laboratory, Didcot, United Kingdom

School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom

School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom

Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom

Imperial College London, London, United Kingdom

Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom

Department of Physics, University of Oxford, Oxford, United Kingdom

Massachusetts Institute of Technology, Cambridge, MA, United States

University of Cincinnati, Cincinnati, OH, United States

University of Maryland, College Park, MD, United States

Los Alamos National Laboratory (LANL), Los Alamos, United States

Syracuse University, Syracuse, NY, United States

School of Physics and Astronomy, Monash University, Melbourne, Australia, associated to 56

 70 Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to 2

Physics and Micro Electronic College, Hunan University, Changsha City, China, associated to 7

Guangdong Provincial Key Laboratory of Nuclear Science, Guangdong-Hong Kong Joint Laboratory of Quantum Matter, Institute of Quantum Matter, South China Normal University, Guangzhou, China, associated to ³

 $^{73}School$ of Physics and Technology, Wuhan University, Wuhan, China, associated to 3

 74 Departamento de Fisica, Universidad Nacional de Colombia, Bogota, Colombia, associated to 13

 75 Universität Bonn - Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany, associated to 17

 76 Institut für Physik, Universität Rostock, Rostock, Germany, associated to 17

 $77Eotvos$ Lorand University, Budapest, Hungary, associated to 48

INFN Sezione di Perugia, Perugia, Italy, associated to ²¹

 79 Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to 32

Universiteit Maastricht, Maastricht, Netherlands, associated to 32

National Research Centre Kurchatov Institute, Moscow, Russia, associated to 41

National Research University Higher School of Economics, Moscow, Russia, associated to 42

National University of Science and Technology "MISIS", Moscow, Russia, associated to 41

National Research Tomsk Polytechnic University, Tomsk, Russia, associated to 41

 $85DS4DS$, La Salle, Universitat Ramon Llull, Barcelona, Spain, associated to 45

University of Michigan, Ann Arbor, United States, associated to 68

 a Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil

 b Hangzhou Institute for Advanced Study, UCAS, Hangzhou, China

 c Università di Bari, Bari, Italy

 d Università di Bologna, Bologna, Italy

 e Università di Cagliari, Cagliari, Italy

 f Università di Ferrara, Ferrara, Italy

 g Università di Firenze, Firenze, Italy

 h Università di Genova, Genova, Italy

 i Università degli Studi di Milano, Milano, Italy

 j Università di Milano Bicocca, Milano, Italy

 k Università di Modena e Reggio Emilia, Modena, Italy

 $^l Università di Padova, Padova, Italy$ </sup>

^mScuola Normale Superiore, Pisa, Italy

 n Università di Pisa, Pisa, Italy

 \degree Università della Basilicata, Potenza, Italy

 p Università di Roma Tor Vergata, Roma, Italy

^qUniversità di Siena, Siena, Italy

 r Università di Urbino, Urbino, Italy

 ^{s}MSU - Iligan Institute of Technology (MSU-IIT), Iligan, Philippines

^tAGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland

 U^u P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia

 v Novosibirsk State University, Novosibirsk, Russia

^wDepartment of Physics and Astronomy, Uppsala University, Uppsala, Sweden

^xHanoi University of Science, Hanoi, Vietnam