## Massive Star-Forming Galaxies Have Converted Most of Their Halo Gas into Stars

Ziwen Zhang<sup>1,2</sup>, Huiyuan Wang<sup>1,2</sup>, Wentao Luo<sup>1,2</sup>, Jun Zhang<sup>3</sup>, H. J. Mo<sup>4</sup>, YiPeng Jing<sup>3,1</sup>, Xiaohu Yang<sup>3</sup>, and Hao  $Li^{1,2}$ 

CAS Key Laboratory for Research in Galaxies and Cosmology, Department of Astronomy, University of Science and Technology of China, Hefei, Anhui 230026, China; Email: ziwen@mail.ustc.edu.cn, whywang@ustc.edu.cn

2

School of Astronomy and Space Science, University of Science and Technology of China, Hefei 230026, China 3

Department of Astronomy, and Tsung-Dao Lee Institute, Shanghai Jiao Tong University, Shanghai 200240, China

Department of Astronomy, University of Massachusetts, Amherst MA 01003-9305, USA

#### ABSTRACT

In the local Universe, the efficiency for converting baryonic gas into stars is very low. In dark matter halos where galaxies form and evolve, the average efficiency varies with galaxy stellar mass and has a maximum of about twenty percent for Milky-Way-like galaxies. The low efficiency at higher mass is believed to be produced by some quenching processes, such as the feedback from active galactic nuclei. We perform an analysis of weak lensing and satellite kinematics for SDSS central galaxies. Our results reveal that the efficiency is much higher, more than sixty percent, for a large population of massive star-forming galaxies around  $10^{11} M_{\odot}$ . This suggests that these galaxies acquired most of the gas in their halos and converted it into stars without being affected significantly by quenching processes. This population of galaxies is not reproduced in current galaxy formation models, indicating that our understanding of galaxy formation is incomplete. The implications of our results on circumgalactic media, star formation quenching and disc galaxy rotation curves are discussed. We also examine systematic uncertainties in halo-mass and stellar-mass measurements that might influence

Key words. gravitational lensing - galaxies: formation - galaxies: haloes - dark matter - large-scale structure of Universe

Technology of China, Hefei, Anhui 230026, China; Email
<sup>2</sup> School of Astronomy and Space Science, University of Sc.
<sup>3</sup> Department of Astronomy, and Tsung-Dao Lee Institute,
<sup>4</sup> Department of Astronomy, University of Massachusetts,
May 23, 2022 **ABST**In the local Universe, the efficiency for converting baryonic g form and evolve, the average efficiency varies with galaxy st Milky-Way-like galaxies. The low efficiency at higher mass is as the feedback from active galactic nuclei. We perform an central galaxies. Our results reveal that the efficiency is mutassive star-forming galaxies around 10<sup>11</sup>M<sub>☉</sub>. This suggest and converted it into stars without being affected significant reproduced in current galaxy formation models, indicating The implications of our results on circumgalactic media, st. discussed. We also examine systematic uncertainties in hald our results. **Key words**. gravitational lensing - galaxies: formation - gala - methods: statistical - galaxies: general **1. Introduction**In the standard A cold dark matter cosmogony, galaxies are believed to form and evolve within dark matter halos. The baryonic gas in halos cools radiatively and condenses and is then converted into stars(White & Rees 1978; Fall & Efstathiou 1980). The global efficiency for converting baryonic gas into stars is very low (Bregman 2007), about 10%. Within halos, the efficiency is usually defined as *M*<sub>\*</sub>/*M*<sub>h</sub>/*f*<sub>b</sub>, where *M*<sub>\*</sub>, *M*<sub>h</sub> and *f*<sub>b</sub> are stellar mass, halo mass and cosmic mean baryon fraction, respectively. So the efficiency reaches a dominant galaxies in the halos. The efficiency reaches a dominant galaxies in the halos. The efficiency reaches a dominant galaxies in the halos. The efficiency reaches and command the part of radiatively on a contract of the stellar mass of central galaxies, which are the dominant galaxies in the halos. The efficiency reaches and command the pare stellar mass of central galaxies, which are th ies and declines quickly towards the lower and higher masses (e.g. Yang et al. 2003, 2009; Leauthaud et al. 2012; Moster et al. 2013; Lu et al. 2014; Hudson et al. 2015; Mandelbaum et al. 2016; Wechsler & Tinker 2018; Behroozi et al. 2019).

The efficiency is expected to be the result of many physical processes in and around galaxies, and its massdependence may reflect the relative importance of individual processes. For example, the decline at the low-mass side may be produced by supernova feedback and stellar winds (e.g. Kauffmann & Charlot 1998; Cole et al. 2000). The gravitational potential wells associated with these low-mass

galaxies are expected to be shallow, so that these processes can effectively prevent star formation and the growth of galaxies. In contrast, at the high mass end, where effects of supernova feedback may not be important, the suppression of the star formation efficiency is usually believed to be caused by the energetic feedback from active galactic nuclei (AGNs, e.g. Silk & Rees 1998; Croton et al. 2006; Fabian 2012; Heckman & Best 2014), although other processes might also be at work, such as morphological quenching and virial shock heating (e.g. Dekel & Birnboim 2006; Martig et al. 2009).

The low efficiency described above is the average for galaxies of a given stellar mass. It is clearly interesting to check whether the efficiency varies with galaxy properties other than the stellar mass. There is a growing amount of evidence that red/quiescent/early type galaxies reside in more massive halos than blue/star-forming/late type galaxies of the same stellar mass (e.g. More et al. 2011; Rodríguez-Puebla et al. 2015; Mandelbaum et al. 2016; Behroozi et al. 2019; Lange et al. 2019b; Bilicki et al. 2021; Posti & Fall 2021; Xu & Jing 2022; Zhang et al. 2021), indicating that the efficiency for star-forming galaxies is higher than that for quiescent galaxies. Moreover, the efficiency also seems to be related to galaxy morphology (e.g. Mandelbaum et al. 2006; Xu & Jing 2022). For example, Posti et al. (2019) found the efficiency for disc galaxies increases monotonously with increasing stellar mass and deviates significantly from that for the total galaxy population at the massive end. More recently, Zhang et al. (2021) found that the host galaxies of optical AGNs have stellar to halo mass ratio (hereafter SHMR) similar to star-forming galaxies but different from quiescent galaxies. The mean halo mass of these AGNs is around  $10^{12} M_{\odot}$  (see also Mandelbaum et al. 2009), where the star formation efficiency peaks. This hints that AGNs tend to be triggered in galaxies with high star formation efficiency. Another interesting finding is that the peak efficiency declines, and the peak position shifts to lower mass as cosmic time progresses (e.g. Hudson et al. 2015). Although other works using very different approaches gave different results (e.g. Moster et al. 2013; Lu et al. 2015; Behroozi et al. 2019).

One important step to evaluate the efficiency is to measure the halo mass for a galaxy sample or for individual galaxies. In the literature, many methods have been developed to infer the halo masses from observational data, such as weak lensing, satellite kinematics, rotational velocity, galaxy clustering, galaxy abundance, X-ray emission and the Sunyaev-Zel'dovich effect (SZ effect). Galaxy-galaxy lensing and satellite kinematics are two powerful tools to measure halo mass, and have been investigated in great details (e.g. van den Bosch et al. 2004; Mandelbaum et al. 2006; Conroy et al. 2007; More et al. 2011; van Uitert et al. 2011; Leauthaud et al. 2012; Tinker et al. 2013; Wojtak & Mamon 2013; Velander et al. 2014; Hudson et al. 2015; Viola et al. 2015; Zu & Mandelbaum 2015; Mandelbaum et al. 2016; Shan et al. 2017; Luo et al. 2018; Lange et al. 2019a; Zhang et al. 2021). In this paper, we combine the data of both galaxy-galaxy lensing and satellite kinematics to measure halo masses of galaxy samples with different stellar mass and star formation rate, and then to evaluate the efficiency of converting baryonic gas into stars for those samples. We also check our results by using galaxy clustering.

The paper is organized as follows. Section 2.1 presents the sample selection and our method of using lensing and satellite kinematics to infer halo mass. In Section 3, we show our main results for different galaxies, test uncertainties, and make comparisons with other results. We discuss the implications of our results in Section 4. Finally, we summarize our results in Section 5. Throughout this paper, we assume the Planck Cosmology (Planck Collaboration et al. 2016):  $\Omega_{\rm m} = 0.307$ ,  $\Omega_{\rm b} = 0.048$ ,  $\Omega_{\Lambda} = 0.693$ , h = 0.678. The cosmic mean baryon fraction is  $f_{\rm b} = \Omega_{\rm b}/\Omega_{\rm m} = 0.157$ .

#### 2. Samples and Methods of Analysis

#### 2.1. Sample properties

Our galaxy sample is selected from the New York University Value Added Galaxy Catalog (NYU-VAGC) (Blanton et al. 2005b) of the Sloan Digital Sky Survey (SDSS) DR7 (Abazajian et al. 2009). Galaxies with *r*-band Petrosian magnitudes  $r \leq 17.72$ , with redshift completeness > 0.7, and with redshift range of  $0.01 \leq z \leq 0.2$ , are selected. We only focus on central galaxies that are the most massive galaxies within dark matter halos, and the halo-based group catalog of Yang et al. (2007) is adopted to identify centrals <sup>1</sup>. NYU-VAGC provides measurements of the sizes of a galaxy,  $R_{50}$  and  $R_{90}$ , which are the radii enclosing 50 and 90 percent of the Petrosian *r*-band flux of the galaxy, respectively <sup>2</sup>. The concentration of a galaxy, defined as  $C = R_{90}/R_{50}$ , is usually used to indicate the morphology of the galaxy (Shimasaku et al. 2001; Strateva et al. 2001). It also provides colors, e.g.  $(g - r)^{0.1}$ , and the Sérsic radial profile fits for galaxies.

We cross-match our sample with the MPA-JHU DR7 catalog<sup>3</sup> to obtain the measurements of stellar mass  $(M_*)$ (Kauffmann et al. 2003a) and star formation rate (SFR) (Brinchmann et al. 2004). The stellar mass is obtained by fitting the SDSS *ugriz* photometry to models of galaxy spectral energy distribution (SED), and the results are excellent agreement with those obtained by Moustakas et al. (2013) using photometry in 12 UV, optical and infrared bands. The SFR is derived using both the spectroscopic and photometric data of the SDSS. We define star-forming galaxies as the ones on the star-formation main sequence given by  $\log(SFR) = 0.73 \log M_* - 7.3$  (Bluck et al. 2016). The dispersion of the main sequence is about  $0.3 \, \text{dex}$  (e.g. Speagle et al. 2014; Kurczynski et al. 2016). We, therefore, identify galaxies above  $\log(SFR) = 0.73 \log M_* - 7.6$ as star-forming galaxies.

As shown below, we will use the weak lensing shear catalog measured from the Dark Energy Camera Legacy Survey (hereafter DECaLS Dey et al. 2019) to measure the halo masses of our galaxies. We thus only focus on galaxies within the DECaLS region. This selection excludes about thirty-two percent of galaxies. In this way we select two samples of central galaxies, one for star-forming galaxies, which consists of 129,278 galaxies with  $M_* \ge 10^{8.8} M_{\odot}$ , and the other for the total population, which consists of 304,162 galaxies in the same mass range and is used for comparison. We divide each of the two samples into six subsamples, equally spaced in the logarithm of stellar mass with a bin size of 0.5 dex. The second most massive subsample, which is of particular importance for our analysis, has a mean stellar mass of  $M_* \sim 10^{11} \mathrm{M}_{\odot}$  and contains a total of 106,125 galaxies, including 22,099 star-forming galaxies. See Table 1 for the stellar mass range, the mean stellar mass, and the galaxy number for each of the subsamples.

#### 2.2. Weak lensing measurements


The shear catalog <sup>4</sup> used here to measure galaxy-galaxy lensing signals is based on the Dark Energy Camera Legacy Survey (DECaLS) DR8 imaging data (Dey et al. 2019; Zou et al. 2019). The shape of each galaxy is measured using the FOURIER\_QUAD pipeline, which has been shown to yield accurate shear measurements even for extremely faint galaxy images (signal-to-noise ratio < 10) when applied to both simulations (Zhang et al. 2015) and real data (Zhang et al. (2019) for the CFHTLenS data, and Wang et al. (2021) for the DECaLS data). The whole shear catalog covers about 9,000 square degrees in g/r/z bands, containing shear estimators from 190/246/300 million galaxy images, respectively. Note that the images of the same galaxy in

<sup>&</sup>lt;sup>1</sup> The galaxy group catalog is publicly available at https://gax.sjtu.edu.cn/data/Group.html

 $<sup>^2</sup>$  The galaxy size data can be downloaded at http://sdss.physics.nyu.edu/vagc/

<sup>&</sup>lt;sup>3</sup> https://wwwmpa.mpa-garching.mpg.de/SDSS/DR7/

<sup>&</sup>lt;sup>4</sup> The DECaLS shear catalog is publicly available at https://gax.sjtu.edu.cn/data/DESI.html



**Fig. 1.** Excess surface density from DECaLS shear catalog and the corresponding best fitting results. The symbols in twelve panels show the excess surface density (ESD) profiles obtained by stacking the lensing signals for galaxies in each stellar mass bin, as indicated in each panel. Results are shown separately for star-forming galaxies (1st and 3rd columns) and the total population (2nd and 4th columns). The error bars correspond to the standard deviation of 150 bootstrap samples. We fit the ESD by using three components, the stellar mass term (dotted lines with stars), the one-halo term (dashed lines), and the two-halo term (dash-dotted lines). The sum of those components are shown by solid lines.




Fig. 2. The distribution of line-of-sight velocity difference  $(\Delta v)$  between central galaxies and their satellite candidates. The solid lines with error bars show the probability distribution functions (PDF) of  $\Delta v$ . The dashed lines show the Gaussian plus a constant fits to the data points. The error bars correspond to the standard deviation of 100 bootstrap samples. The upper-row shows the results for star-forming centrals, while the lower-row is for the total central populations. The results of star-forming/total samples in different stellar mass bins are shown in different columns, respectively. Note that the scales of the vertical axis are different for different panels.

Table 1. The properties of sta	r-forming and total galaxy	samples in the DECaLS region.
--------------------------------	----------------------------	-------------------------------

$\log M_* \text{ range}^{(a)}$	$100  { m M_{*}/M_{\odot}}^{(b)}$	$N_{\rm c,gal}^{\rm (c)}$	$N_{\rm s,gal}^{\rm (d)}$	$\log M_{\rm h}/M_{\odot}^{\rm (e)}$	Efficiency <sup>(f)</sup>	LcSK Mass <sup>(g)</sup>	LcSK Eff. <sup>(h)</sup>
		-	-	Star-forming			
[8.84, 9.34]	9.14	$7,\!940$	213	$10.99^{+0.43}_{-0.51}$	$0.089\substack{+0.197\\-0.056}$	$11.27^{+0.37}_{-0.32}$	$0.047^{+0.052}_{-0.027}$
[9.34,  9.84]	9.64	$16,\!903$	756	$11.44^{+0.22}_{-0.25}$	$0.102^{+0.127}_{-0.04}$	$11.13^{+0.11}_{-0.11}$	$0.207^{+0.061}_{-0.046}$
[9.84,  10.34]	10.13	$33,\!826$	$1,\!452$	$11.57^{+0.11}_{-0.14}$	$0.233_{-0.053}^{+0.087}$	$11.56^{+0.06}_{-0.07}$	$0.241_{-0.031}^{+0.042}$
[10.34, 10.84]	10.61	47,759	2,829	$11.92^{+0.06}_{-0.07}$	$0.308\substack{+0.05\\-0.04}$	$11.89^{+0.04}_{-0.04}$	$0.335_{-0.031}^{+0.034}$
[10.84, 11.34]	11.03	22,099	$1,\!272$	$12.03^{+0.08}_{-0.09}$	$0.629^{+0.142}_{-0.109}$	$12.01\substack{+0.05\\-0.04}$	$0.66^{+0.069}_{-0.067}$
[11.34, 11.82]	11.43	749	85	$12.45_{-0.35}^{+0.22}$	$0.61_{-0.238}^{+0.755}$	$12.47_{-0.25}^{+0.35}$	$0.58_{-0.32}^{+0.454}$
				Total			
[8.84, 9.34]	9.14	8,984	357	$11.15^{+0.42}_{-0.57}$	$0.063^{+0.169}_{-0.039}$	$11.02^{+0.15}_{-0.16}$	$0.084^{+0.036}_{-0.024}$
[9.34, 9.84]	9.64	19,777	$1,\!417$	$11.63^{+0.17}_{-0.23}$	$0.066^{+0.047}_{-0.021}$	$11.49^{+0.09}_{-0.08}$	$0.09^{+0.018}_{-0.017}$
[9.84, 10.34]	10.15	48,753	$3,\!975$	110+0.09	$0.141^{+0.036}$	$11.82^{+0.04}$	$0.135^{+0.016}_{-0.012}$
[10.34, 10.84]	10.64	$105,\!577$	$15,\!408$	$12.2^{+0.04}_{-0.04}$	$0.175^{+0.018}_{-0.016}$	$12.19^{+0.02}_{-0.02}$	$0.178^{+0.008}_{-0.009}$
[10.84, 11.34]	11.09	$106,\!125$	$27,\!873$	$12.59^{+0.03}$	+ 0.013	+ 0.02	$0.186^{+0.008}$
[11.34, 11.84]	11.47	$14,\!946$	$16,\!584$	$13.37^{+0.02}_{-0.02}$	$0.08^{+0.004}_{-0.004}$	$\frac{12.62^{+0.02}_{-0.02}}{13.32^{+0.03}_{-0.03}}$	$0.09^{+0.007}_{-0.007}$
(a) (T) (1)							

<sup>(a)</sup> The stellar mass range.

<sup>(b)</sup> The mean stellar mass.

<sup>(c)</sup> The number of central galaxies.

<sup>(d)</sup> The number of satellite candidates.

<sup>(e)</sup> The halo mass obtained from galaxy-galaxy lensing.

<sup>(f)</sup> The efficiency obtained from galaxy-galaxy lensing.

 $^{(g)}$  The halo mass  $(\log M_h/M_{\odot})$  obtained from lensing calibrated satellite kinematics method.

<sup>(h)</sup> The efficiency obtained from lensing calibrated satellite kinematics method.

different exposures are counted as different images in the FOURIER\_QUAD method.

Photometric redshifts of galaxies in the shear catalog are calculated using the random forest regression method (Breiman 2001), a machine learning algorithm based on decision trees. Eight parameters were used in training the algorithm, including the r-band magnitude, (q-r), (r-z), (z - W1), and (W1 - W2) colors, half-light radius, axial ratio, and shape probability. The photo-z error is estimated for each individual shear galaxy by perturbing the photometry of the galaxy. Specifically, the uncertainty is assumed to follow a Gaussian distribution with the standard deviation equal to the photometric error; a random value generated from the distribution is added to the observed flux in each band to obtain a 'perturbed' flux; the perturbation is repeated multiple times and the standard deviation of the photo-z estimates from the perturbations is used as the error of the photo-z (see Zhou et al. 2021, for more details).

Here we only use the r and z band data, because we find that the g-band images have some quality issues. The details of the image processing pipeline of the DECaLS data are given in another work (Zhang et al., in preparation). The overlapping region of DECaLS with SDSS DR7 is about 4744 square degrees. The estimator, Excess Surface Density (ESD),

$$\Delta\Sigma(\mathbf{R}) = \gamma_{\rm t}(\mathbf{R})\Sigma_{\rm crit},\tag{1}$$

is measured using the PDF-Symmetrization method (Zhang et al. 2017), which minimizes the statistical uncertainty. Note that due to the scatter as well as the uncertainty of the background galaxy redshifts, the PDF-Symmetrization method should be modified slightly. The details are given in a companion paper (Li et al., in preparation), which also includes a general discussion about different source of system-

Article number, page 4 of 15

atic errors in the measurement of the excess surface density within the framework of the PDF-Symmetrization method. In Figure 1, we show the ESD for both star-forming and total galaxy populations. The error bars are estimated by using 150 bootstrap samples (Barrow et al. 1984).

Following previous studies (e.g. Mandelbaum et al. 2008; Leauthaud et al. 2010; Luo et al. 2018), we model the ESD using three components:

$$\Delta \Sigma = \Delta \Sigma_{\text{stellar}} + \Delta \Sigma_{\text{NFW}} + \Delta \Sigma_{2h}.$$
 (2)

The first term is the contribution of the stellar mass of galaxies. We adopt the stellar mass directly from the observational data and modeled it as a point mass. The second term is the contribution of the dark matter halo, assumed to have a Navarro-Frenk-White(NFW, Navarro et al. 1997) density profile, described by two free parameters: the mass  $m_{\rm h}$  and the concentration c. Specifically,  $m_{\rm h}$  is the mass contained in a spherical region of radius  $r_{200m}$ , within which the mean mass density is equal to 200 times the mean matter density of the Universe. The distributions of the halo mass and concentration for a fixed galaxy stellar mass are usually quite broad. In our modeling, we use a single NFW profile with two free parameters,  $m_{\rm h}$  and concentration, to fit the data point, ignoring the dispersion in them. Since our analysis focuses only on the average information of halos, the bias produced by ignoring the dispersion is expected to be small, as shown in Mandelbaum et al. (2016) and to be discussed further in Section 3.3. Following Mandelbaum et al. (2016), we also ignore the effect of off-centering, which is negligible for the  $m_{\rm h}$  estimation of central galaxies according to our tests. The third term, referred to as the two-halo term, is calculated by projecting the halo-matter cross correlation function,  $\xi_{\rm hm}$ , along the line-of-sight. Here  $\xi_{hm} = b(m_h)\xi_{mm}$ , with  $\xi_{mm}$  being the linear matter-matter correlation function and  $b(m_h)$ the linear halo bias (Tinker et al. 2010), both generated using *COLOSSUS* (Diemer 2018). We sample the posterior distribution of the two parameters,  $m_h$  and c, using the Monte Carlo Markov Chain (MCMC) provided by a public open software, *emcee* (Foreman-Mackey et al. 2013). Both *COLOSSUS* and *emcee* are under the MIT License. The best-fitting profiles are presented in Figure 1. The quoted mass,  $m_h$ , is the median value of the posterior and its error bar indicates the 16 and 84 percentage range of the posterior.

We note that  $m_{\rm h}$  actually accounts for the contribution from cold dark matter, diffuse gas and satellites around centrals, but does not include the contribution from the central galaxies that is modeled by  $\Delta \Sigma_{\rm stellar}$ . The total mass of the halo, which is used to calculate the conversion efficiency in the following and should include all the components within the virial radius of the halo, is thus  $M_{\rm h} = m_{\rm h} + M_*$ . In general,  $M_* \ll m_{\rm h}$  and so  $M_{\rm h}$  is very close to  $m_{\rm h}$ . However, for halos with very high efficiency, central galaxies can also have a significant contribution to the total halo mass. In the rest of the paper, halo mass refers to the total mass of the halo,  $M_{\rm h}$ .

#### 2.3. Weak lensing calibrated satellite kinematics method

The kinematics of satellite galaxies provide an important probe of the gravitational potential wells of the dark matter halos (e.g. McKay et al. 2002; van den Bosch et al. 2004; More et al. 2011; Wojtak & Mamon 2013; Lange et al. 2019a,b; Abdullah et al. 2020; Seo et al. 2020). For a central galaxy of mass  $M_*$  in a given subsample, we identify its satellite candidates from a reference galaxy sample. For our analysis, we define the reference sample as a magnitudelimited sample following the selection criteria: r-band Petrosian apparent magnitude of r < 17.6, r-band Petrosian absolute magnitude in the range of (-24, -16), and redshift in range of 0.01 < z < 0.2 (Wang & Li 2019; Zhang et al. 2021). The candidates are defined as the ones that satisfy the following criteria:  $|\Delta v| \leq 3v_{\rm vir}, r_{\rm p} \leq r_{\rm vir}$  and  $M_{\rm s} < M_{*}$ . Here  $r_{\rm p}$  and  $\Delta v$  are the projected distance and the line-ofsight velocity difference between the central in question and it's satellite, respectively,  $r_{\rm vir}$  and  $v_{\rm vir}$  are, respectively, the virial radius and virial velocity calculated using the mean halo mass of the subsample derived from weak lensing, and  $M_{\rm s}$  is the stellar mass of satellite. The numbers of satellite candidates for different central galaxy samples are listed in Table 1.

Figure 2 shows the probability distribution functions (PDFs) of  $\Delta v$  for the selected satellites associated with different (total/star-forming) central galaxy subsamples. We use a Gaussian plus a constant,

$$\frac{A}{\sqrt{2\pi}\sigma_v} \mathrm{e}^{-\Delta v^2/2\sigma_v^2} + d,\tag{3}$$

to fit the PDFs. Here, the Gaussian component represents the true satellites, and the constant component is used to account for the interlopers that are not physically associated with the centrals (see also McKay et al. 2002; Brainerd & Specian 2003; van den Bosch et al. 2004; Conroy et al. 2007). A MCMC technique is used to perform the fitting. As can be seen from the figure, the  $\Delta v$  distributions are well fitted by the two-component model and the




Fig. 3. Halo mass-satellite velocity dispersion relation. The open (solid) diamonds show the velocity dispersion ( $\sigma_s$ ) of satellites versus the host halo masses for total (star-forming) central galaxies. The halo masses are measured from DECaLS lensing data. Different colors indicate different stellar mass bins of central galaxies. The error bars for halo mass indicate the 16 and 84 percentiles of the posterior distribution obtained from the MCMC fitting to the stacked lensing profiles. The error bars for  $\sigma_s$  represent the 16 and 84 percentiles of the posterior distribution obtained from the satellite-central velocity difference. The solid line represents the best-fitting to the data points for total galaxy sample.

contribution of the interlopers is negligible. Given that the uncertainty of a SDSS galaxy redshift is about  $35 \,\mathrm{km}\,\mathrm{s}^{-1}$ , the error of  $\Delta v$  is  $\sigma_e = \sqrt{2} \times 35 = 49.5 \,\mathrm{km}\,\mathrm{s}^{-1}$ . So the velocity dispersion,  $\sigma_{\rm s}$ , of satellites can be estimated from the best-fitting Gaussian after correcting for the redshift uncertainties,  $\sigma_{\rm s} = \sqrt{\sigma_v^2 - \sigma_e^2}$ . The estimate of  $\sigma_{\rm s}$  may be affected by the uncertainty of the halo masses that are used to determine  $r_{\rm vir}$  and  $v_{\rm vir}$ . As a check, for each sample we compare the  $\sigma_{\rm s}$  calculated using three different halo masses, corresponding to the 16, 50 and 84 percentiles of the posterior distribution obtained from the MCMC fitting to the stacked lensing profiles of the sample in question. We find that the three values of  $\sigma_{\rm s}$  so obtained are in general consistent with each other (see Appendix A for the detail). We thus conclude that the uncertainty in the halo mass has little impact on the estimate of  $\sigma_{\rm s}$ .

In Figure 3, we show  $\sigma_s$  versus  $M_h$ , the lensing mass measurement, for both total and star-forming samples. One can see a strong correlation between the two parameters. We fit the data points for the total sample with a power-law model, and constrained relation is

$$\log(M_{\rm h}/M_{\odot}) = (3.08 \pm 0.05) \log(\sigma_{\rm s}/\,{\rm km\,s^{-1}}) + (5.33 \pm 0.11). \tag{4}$$

Remarkably, the slope of the relation is in excellent agreement with what is expected from the virial scaling relation  $(M \propto \sigma^3)$  seen in numerical simulations (Evrard et al. 2008). A similar slope has been obtained in previous studies measuring halo mass using abundance matching, satellite kinematics, caustic technique, SZ effect, and virial theorem (e.g. Yang et al. 2007; More et al. 2011; Rines et al. 2013, 2016; Abdullah et al. 2020), although some other studies found different results (e.g. Viola et al. 2015). Our tests show that both weak lensing and satellite kinematics can provide robust and consistent measurements of the host halo masses of galaxies. The star-forming galaxies also closely follow the trend defined by the total sample, demonstrating that the lensing masses for star-forming galaxies are also robust.

Thus, we can derive the host halo masses for the starforming and total subsamples from their measured  $\sigma_{\rm s}$  (Figure 2) using the  $\sigma_{\rm s}$ - $M_{\rm h}$  relation. But,  $\sigma_{\rm s}$  is also used in fitting the relation, the uncertainties in  $\sigma_{\rm s}$  and in the relation are expected to be correlated. In order to avoid this correlation, we design a new way to calculate the  $\sigma_{\rm s}$ - $M_{\rm h}$ relation. Specifically, for each stellar mass bin, we only use data points ( $\sigma_{\rm s}$  versus  $M_{\rm h}$ ) in the rest of the stellar mass bins to fit the  $\sigma_{\rm s}$ - $M_{\rm h}$  relation, so that the derived relation is independent of the galaxy sample in question. In Appendix B, we show the best-fitting  $\sigma_{\rm s}$ - $M_{\rm h}$  relations for all six stellar mass bins.

We then combine the  $\sigma_s$  of galaxy sample with its corresponding  $\sigma_s$ - $M_h$  relation to derive the halo mass. The errors of the halo masses are obtained by considering both the uncertainties in  $\sigma_s$  and the  $M_h$ - $\sigma_s$  relation. Both  $\sigma_s$  and the  $M_h$ - $\sigma_s$  relation are derived by using MCMC fitting, which can provide the posterior distributions for them. For galaxies in a given stellar mass bin, we can randomly generate a  $\sigma_s$  value and a  $M_h$ - $\sigma_s$  relation respectively from their posterior distributions, and predict a halo mass by combining the two. In practice, we generate 30,000 predictions of the halo mass for each subsample, and use the 16 and 84 percentiles of the mass distribution to represent the uncertainties. The halo mass estimated in this way is referred to as the weak lensing calibrated satellite kinematics (LcSK) halo mass, and is denoted also by  $M_h$ .

Finally, we want to note that the uncertainties of the LcSK masses in different stellar mass bins are correlated. Therefore, if one wants to use these data points to constrain a galaxy formation model, one should consider the covariance among different stellar mass bins.

#### 2.4. Galaxy clustering analysis

Galaxy clustering can be used to measure the large scale environment and to infer the halo mass for a sample of selected galaxies. In this paper, we adopt the projected twopoint cross-correlation function (2PCCF) to quantify the clustering of galaxies. We first estimate the 2PCCF using

$$\xi(r_{\rm p}, r_{\pi}) = \frac{N_{\rm R}}{N_{\rm D}} \frac{GD(r_{\rm p}, r_{\pi})}{GR(r_{\rm p}, r_{\pi})} - 1,$$
(5)

where  $N_{\rm R}$  and  $N_{\rm D}$  are the galaxy numbers in the random and reference samples, respectively;  $r_{\rm p}$  and  $r_{\pi}$  are the separations perpendicular and parallel to the line of sight, respectively; GD is the number of cross pairs between the selected galaxy sample and the reference sample and GRis that between the selected galaxy sample and the random sample. To obtain the projected 2PCCF, we integrate  $\xi$  along the line of sight ,  $w_{\rm p}(r_{\rm p}) = \int_{-\Delta s}^{\Delta s} \xi(r_{\rm p}, r_{\pi}) dr_{\pi}$ , with  $\Delta s = 40 h^{-1}$ Mpc, sufficiently large so as to include almost all correlated pairs. The errors on the measurements of the 2PCCF are estimated by using 100 bootstrap samples.

Article number, page 6 of 15





Fig. 4. Stellar mass - halo mass relation (SHMR). The red symbols with error bars show the SHMR calculated by using the halo mass obtained from the fits to the stacked lensing mass profiles from the DECaLS shear catalog, while the black ones are based on the weak lensing calibrated satellite kinematics(LcSK) method. For clarity, we shift the black symbols towards the left slightly. The error bars reflect the 16 and 84 percentiles of the posterior distribution. The diamonds and squares show the results for star-forming and total populations, respectively. The shadow region represents the range covered by curves published in Yang et al. (2009); Moster et al. (2010); Leauthaud et al. (2012); Kravtsov et al. (2018); Behroozi et al. (2019).

The reference sample used here is the same as that described in Subsection 2.3. Based on this reference sample, we construct a random sample in the following way (see also Li et al. 2006, for more details). We generate ten duplicates for each galaxy in the reference sample and randomly place them in the SDSS sky coverage. All other properties, including stellar mass and redshift of the duplicate galaxies, are the same as those of the original galaxy. Thus, the random sample has the same survey geometry and the same distribution of galaxy properties as the reference sample.

The 2PCCFs of the star-forming galaxy sample cannot be directly compared with that of the total galaxy sample, because the two samples may have different redshift distributions. In order to make a fair comparison, it is necessary to construct a control sample that matches the starforming sample. For a star-forming galaxy in a given stellar mass bin, we select, from the total galaxy subsample in the same stellar mass bin, n galaxies whose redshift are within 0.005 from the star-forming galaxy in question. We choose n = [1, 1, 1, 2, 4, 19] for the six galaxy stellar mass bins (see Section 2.1). The number of control galaxies, n, is chosen according to the ratio in size between total and star-forming galaxy subsamples. This control sample is then used to estimate the 2PCCFs to compare with the corresponding starforming galaxies.



**Fig. 5.** Baryon conversion efficiency as a function of stellar mass. In the left panel, the symbols with error bars show the efficiency calculated by using the halo mass obtained from the fits to the stacked lensing mass profiles. The black ones show the results from the DECaLS shear catalog, while the green ones are obtained from the SDSS shear catalog (see Appendix C). The solid and open symbols show the results for star-forming and total populations, respectively. Error bars indicate the 16 and 84 percentiles of the posterior distribution obtained from the MCMC fitting. In the right panel, the symbols with error bars show the efficiency calculated by using the LcSK method. The error bars reflect the 16 and 84 percentiles of the posterior distribution obtained from the results for star-forming (total) galaxies in the hydro-simulation IllustrisTNG. The shadow region in two panels covers the range obtained before, as in Figure 4.

#### 3. Results

#### 3.1. The stellar mass - halo mass relation

Figure 4 and Table 1 show the halo mass  $(M_{\rm h})$  obtained from weak lensing (lensing mass) and from the weak lensing calibrated satellite kinematics (LcSK) method, separately for the total and star-forming central galaxies of different stellar masses. Both estimates give very similar results, as is expected from the tight correlation between the velocity dispersion and the lensing mass (Figure 3). For comparison, we also show the results in the literature obtained using various methods, including galaxy groups, abundance matching, conditional luminosity function, weak lensing, and empirical model (Yang et al. 2009; Moster et al. 2010; Leauthaud et al. 2012; Kravtsov et al. 2018; Behroozi et al. 2019). As one can see from the figure, the stellar mass-halo mass relation (SHMR) for our total galaxy sample follows closely the trend defined by previous results. In contrast, for a given stellar mass, the halo mass of starforming galaxies is lower than that of the total sample, and the difference becomes larger and more significant at the high-mass end. This is broadly consistent with previous weak-lensing and satellite kinematics studies that found star-forming/blue galaxies reside in less massive halos than quiescent/red galaxies of the same stellar mass (see e.g. Mandelbaum et al. 2006; More et al. 2011; Mandelbaum et al. 2016; Lange et al. 2019b; Zhang et al. 2021), and with results obtained from some empirical models constrained by observational data (e.g. Rodríguez-Puebla et al. 2015; Behroozi et al. 2019).

#### 3.2. The conversion efficiency

Assuming that the baryon fraction within halos is equal to the cosmic fraction  $(f_{\rm b})$ , the overall conversion efficiency can be represented by  $M_*/(f_b M_h)$ . Here  $M_h$  is the total mass of a dark matter halo,  $M_*$  is the stellar mass of the central galaxy within the halo. In the left panel of Figure 5, we show this conversion efficiency as a function of stellar mass for the total galaxy sample, with the halo mass measured directly from the weak lensing data. The efficiency for the total galaxy sample peaks roughly at  $M_* \sim 10^{10.6} M_{\odot}$ and decreases towards both lower and higher mass ends, in good agreement with previous results (e.g. Wechsler & Tinker 2018). The same panel also shows the results for star-forming central galaxies defined in Section 2.1. The efficiency of star-forming galaxies follows roughly the trend of the total population at the low-mass end, but deviates from it and continues the increasing trend toward the massive end. For star-forming galaxies of  $M_* \sim 10^{11} \mathrm{M}_{\odot}$ , the efficiency reaches a value of  $0.629^{+0.142}_{-0.109}$ , much higher than the maximum value of  $\sim 0.2$  for Milky-way-like galaxies with  $M_* \sim 10^{10.6} \mathrm{M}_{\odot}$ . At  $M_* > 10^{11} \mathrm{M}_{\odot}$ , the high efficiency seems to remain, although with a lager uncertainty. This high efficiency indicates that the most of the baryonic gas in the host halos of those massive star-forming galaxies has already assembled into galaxies and been converted into stars.

We can also use the halo mass obtained from the LcSK method to estimate the conversion efficiency (Section 2.3), and results are presented in the right panel of Figure 5 and




Fig. 6. The ratio of the 2PCCF between star-forming galaxies and the total control galaxies as a function of the projected distance  $(r_p)$  in different stellar mass bins. Error bars are the standard deviation of the 2PCCF ratio among 100 bootstrap samples. The horizontal dashed line in each panel indicates the theoretical ratio from a halo bias model (Tinker et al. 2010) using the halo mass measured from weak lensing as the input. For clarity, we only show the results for the four highest-mass bins, where the uncertainties in lensing mass measurements are relatively small.

in Table 1. The results are very similar to those obtained from the weak lensing data. For example, the efficiency for the second most massive star-forming sample is  $0.66^{+0.069}_{-0.067}$ . However, the LcSK method significantly reduces the uncertainty of the efficiency in most of the stellar mass bins.

As mentioned above, many previous studies found that star-forming or blue galaxies tend to reside in smaller halos than quenched or red galaxies of the same stellar mass(e.g. More et al. 2011; Hudson et al. 2015; Mandelbaum et al. 2016; Lange et al. 2019b; Behroozi et al. 2019; Posti et al. 2019; Bilicki et al. 2021). This is consistent with our results that the conversion efficiency is higher for star-forming galaxies than for quenched galaxies. However, in comparison with our results, the implied conversion efficiency is significantly lower and has larger uncertainties in many of the earlier studies. For example, Mandelbaum et al. (2006) and Mandelbaum et al. (2016) found a peak efficiency of  $0.35^{+0.92}_{-0.17}$  for late-type galaxies and  $0.351^{+0.165}_{-0.089}$  for blue galaxies, respectively. Dutton et al. (2011) found that the mean peak efficiency for their late-type galaxies is around 0.3. Rodríguez-Puebla et al. (2015) found that the  $M_*$ -to- $M_{\rm h}$  ratio of their blue centrals has a peak value of 0.051, corresponding to a conversion efficiency of 0.325 assuming  $f_{\rm b} = 0.157$ . These results appear to be in conflict with ours, and we will come back to this issue later. Recently, Posti et al. (2019) modeled the rotation curves of local disc galaxies and inferred the halo mass of individual galaxies. They found that the mean conversion efficiency for about 20 massive disc galaxies is about 0.5 but with large uncertainties. Our results are in broad agreement with theirs. However, our results are obtained from a large sample of 22,099

star-forming galaxies, indicating that the high conversion efficiency is a common property for massive star-forming galaxies.

#### 3.3. Tests of uncertainties

A number of factors may affect the estimates of the halo mass. As a test, we repeat our analysis using an independent weak-lensing shear catalog obtained by Luo et al. (2017) from the Sloan Digital Sky Survey (SDSS) DR7 (Abazajian et al. 2009) imaging data using a totally different method (see Appendix C). As shown in Figure 5, the results obtained from the SDSS data agree well with those from the DECaLS data. However, the SDSS results have much larger uncertainties because of the shallower imaging data used to measure the weak lensing shear.

Galaxy clustering provides another test because of its dependence on halo mass (Mo & White 1996; Tinker et al. 2010). Figure 6 shows the ratio of the 2PCCF, in the range of  $5 \,\mathrm{Mpc} < r_{\mathrm{p}} < 20 \,\mathrm{Mpc}$ , between star-forming galaxies and total population in four high-mass bins (see Section 2.4 for the method to estimate the 2PCCF). The horizontal dashed line in each panel shows the theoretical prediction using the linear halo bias model (Tinker et al. 2010) and the halo mass was derived from the DECaLS weak lensing measurements. As one can see, the model predictions agree very well with the observational results on large scales where the linear halo bias model is valid. However halo bias may also depends on halo assembly history, in addition to halo mass (e.g. Gao et al. 2005; Wang et al. 2007). If there is a correlation between the properties of the central galaxy in a halo and the assembly of the halo, then the halo bias model used here may not be an accurate description for star-forming and quenched centrals. Unfortunately, this potential correlation is not understood well enough to quantify the effect which may produces on our results.

As discussed above, the conversion efficiency obtained here for massive star-forming galaxies appears to be higher than that obtained in some previous investigations. One cause of the discrepancy may be that galaxy samples used in these investigations are different from ours. For example, Mandelbaum et al. (2016) and Bilicki et al. (2021) split galaxy samples according to galaxy color, instead of the SFR used here. As a test, we have repeated our analysis by separating red and blue galaxies according to equation (1)in van den Bosch et al. (2008), a way similar to that in Mandelbaum et al. (2016). The SHMRs obtained from the red and blue populations are shown in Figure 7, in comparison with the results taken from the table B1 of Mandelbaum et al. (2016)<sup>5</sup> and from Bilicki et al. (2021). As one can see, our results are in good agreement with theirs, which provides additional support to the reliability of our mass estimates. In particular, blue galaxies of  $M_* \sim 10^{11} M_{\odot}$  have an average halo mass of  $12.31^{+0.07}_{-0.07}$  from Mandelbaum et al. (2016) and  $12.39^{+0.15}_{-0.08}$  from Bilicki et al. (2021), corresponding to a conversion efficiency of  $0.338^{+0.062}_{-0.052}$  and  $0.262^{+0.053}_{-0.077}$ respectively, much lower than that for star-forming galaxies of the same stellar mass (Fig. 5 and Tab. 1). This suggests that the calculated efficiencies are sensitive to the sample selection, and that the discrepancy between our results

 $<sup>^{5}</sup>$  To be consistent, the stellar masses are adopted from the MPA/JHU catalog, and their halo masses are uncorrected; see the paper for the correction.




Fig. 7. Comparison of our SHMR with Mandelbaum et al. (2016) and Bilicki et al. (2021). The diamonds, the lines with error bars and the shaded regions are the SHMR from our results, Mandelbaum et al. (2016) and Bilicki et al. (2021), respectively. The solid symbols show the results calculated by using DECaLS shear catalog. The red and blue color represent the red and blue galaxy samples, respectively.

(samples selected by the star formation rate) and their results (samples selected by galaxy colors) is entirely due to the difference in the sample selection.

Other potential problems may also exist in the halo mass estimation. If a fraction of the selected galaxies are satellites, instead of centrals, a systematic bias can be introduced, as our modeling of the lensing measurements assumes that all galaxies are centrals. Contamination by satellites is expected to lead to an overestimation of the halo mass (Mandelbaum et al. 2006, 2016), and so the conversion efficiency may be underestimated in our results. To check the impact of this effect, we make an analysis by using an additional selection criterion to reduce the contamination of satellites in our central galaxy samples. Specifically, we require that a central galaxy be the most massive one among all its neighbors that have projected distances less than 1 Mpc, and line-of-sight velocity differences smaller than  $1000 \,\mathrm{km \, s^{-1}}$ , relative to the galaxy in question. This leads to a sample of 4,900 star-forming galaxies in the second most massive bin. The halo mass obtained from lensing for this sample is  $\log M_{\rm h}/{\rm M}_{\odot} = 12.06^{+0.16}_{-0.21}$ , very close to the value obtained above, demonstrating that the effect of satellite contamination is negligible in our results.

Another bias in the halo mass estimate may arise because halos of galaxies in a given sample can span a large range in mass. The stacked lensing signal around the sample galaxies is, therefore, an average of many different halo profiles, and using a single NFW profile to fit the average profile may introduce a bias. Detailed tests by Mandelbaum et al. (2016) using a mock catalog constructed from a semi-analytic galaxy formation model suggest that the bestfitting mass underestimates the mean halo mass by about 10%, quite independent of the stellar mass and galaxy color. At  $M_* \sim 10^{11} {\rm M}_{\odot}$ , the actual mean halo mass may be about




Fig. 8. Comparison of the MPA/JHU stellar mass with the stellar masses from NYU-VGAC (blue) and Salim et al. (2016) (red) for central star-forming galaxies. The solid lines show the median value of the residual, while the shaded regions show the 25 and 75 percentiles of the residual distribution.

1.14 times the best fitting halo mass. Taking this correction into account, the conversion efficiency for star-forming galaxies in this mass range changes to 0.629/1.14 = 0.552. One should keep in mind that the correction factor may depends on the used galaxy formation model. As we will see below, our results are not well reproduced by the current galaxy formation models.

Yet another important source of uncertainty comes from the estimate of the stellar mass. The statistical uncertainty of the stellar mass is usually 0.3 dex (e.g. Kauffmann et al. 2003b). Our galaxy sample is large enough, so that the statistical error in the mean stellar mass is small. However, the systematic uncertainties, such as those produced by the adopted initial mass function, the star formation history, the stellar library, and the dust attenuation, may not be negligible (see Moustakas et al. 2013, for a brief introduction). It is in general difficult to evaluate such systematic uncertainties within a specific model of the stellar population. One common practice is to make a consistency check by comparing the stellar masses of the same object measured with different techniques and/or from different data. The MPA masses used here have been used in many comparison studies in the literature. For example, Moustakas et al. (2013) showed that the MPA masses are in excellent agreement with theirs based on the fits to SEDs in 12 UV, optical and infrared bands. As a check, here we make a similar comparison for central star-forming galaxies, as they are the most relevant to our results. In Figure 8, we compare the MPA masses with those given in NYU-VAGC and those obtained by Salim et al. (2016). Different approaches and/or data were used to derive stellar masses in these three databases. As one can see, there are some systematic differences among the three masses. At  $M_* \sim 10^{11} M_{\odot}$ , the difference between the MPA mass and the other two is about 0.05 dex, and MPA mass appears to lie between the two masses. Thus, our results are robust as long as the systematic bias in the stellar mass is not much larger than the difference among the three mass estimates compared here.




Fig. 9. The probability distributions of galaxy concentration  $(C = R_{90}/R_{50})$  for total galaxy sample (green line) and massive star-forming galaxy sample (black line) of  $M_* \sim 10^{11} M_{\odot}$ .

#### 4. Discussion

One possibility for the large  $M_*$  to  $M_{\rm h}$  ratio reported here is that their host halos are splashback halos (e.g. Ludlow et al. 2009), which have ever entered the virial radii of more massive halos, and were severely stripped by the tidal field. However, splashback halos are usually much more strongly clustered than other halos of the same mass (Wang et al. 2009), because they are spatially close to massive halos. This is clearly inconsistent with the fact that the massive star-forming galaxies are less clustered than the total population of the same mass (Figure 6). Moreover, environmental effects may quench the star formation in splashback halos, and so their galaxies are not expected to be star forming. Thus, this possibility can be ruled out.

Another possibility is that some processes that are supposed to prevent the growth of massive galaxies did not operate on these galaxies in the past. Figure 9 shows the distributions of galaxy concentration for star-forming and total galaxies with  $M_* \sim 10^{11} \mathrm{M}_{\odot}$ . One can see that starforming galaxies have much smaller concentrations than the total galaxies of the same mass. AGN activities, which are thought to be capable of quenching star formation, are expected to be more prominent in more concentrated galaxies (e.g. Kauffmann et al. 2003a; Zhang et al. 2021). Previous theoretical and observational studies found that the efficiency of star-forming galaxies may be suppressed by several feedback processes, such as supernova and AGN feedback in the low and high stellar mass ends, respectively. Our results suggest that AGN feedback must be inefficient in suppressing cold gas acquisition and star formation in massive star-forming galaxies. AGN feedback may still be effective in quenching the star formation in other massive galaxies of  $M_* \sim 10^{11} M_{\odot}$ , and thus to produce a much lower conversion efficiency in them. As shown in Table 1, the number of the massive star-forming galaxies  $(M_* \sim 10^{11} M_{\odot})$  is much smaller than that of the total galaxy population of the same mass. Thus this absence of effective AGN feedback only applies to a relatively small fraction of the total galaxy population.

These massive star-forming galaxies have already converted more than sixty percent of their halo gas into stars. Based on CO and HI observations, Saintonge et al. (2016) found that these galaxies on average contain about  $\sim 10^{10} M_{\odot}$  in cold gas. Thus the total baryon mass in these galaxies is more than seventy percent of the total baryons in their halos. This leaves less than thirty percent of the baryons in the circumgalactic medium (CGM). Thus, observations of the CGM may provide an independent way to check our results. CGM can be probed in a number of ways, such as quasar absorption line systems (e.g. Tumlinson et al. 2017), extended X-ray emission (e.g. Anderson et al. 2013), and SZ effect (e.g. Lim et al. 2018). Many studies found evidence for the existence of CGM in galaxies with a wide range of stellar mass (e.g. Tumlinson et al. 2017). However, a detailed comparison with our results is not straightforward. As shown in Table 1, star-forming galaxies of  $10^{11} M_{\odot}$ make only about one-fifth of the total population of the same mass. It is thus inappropriate to compare our results directly with those that did not separate the star-forming galaxies from the total population. For low-ionization line systems observed in star forming galaxies, such as MgII, the absorbing gas is believed to be associated with outflow (e.g. Lan & Mo 2018). The total amount of gas involved may not be large, which is consistent with our results.

The mean density of the CGM around massive starforming galaxies is expected to be less than that around other galaxies. Therefore, the timescale for gas cooling, which is inversely proportional to the gas density, is much longer. The ability of massive star-forming galaxies to acquire additional gas to maintain a high star formation rate should be significantly suppressed. Our results suggest that the well-known flattening of the SFR- $M_*$  relation for starforming galaxies at the massive end (Whitaker et al. 2014; Saintonge et al. 2017) is caused by the decrease of gas supply. It is consistent with the analysis based on atomic and molecular gas within galaxies(Saintonge et al. 2016).

The high galaxy-to-halo mass ratio for massive starforming galaxies may have important implications for their rotation curves. To demonstrate this, we select galaxies with  $M_{*}\,\sim\,10^{11}{\rm M}_{\odot}$  and Sérsic index of  $n\,\sim\,1$  from the SDSS galaxy sample. These massive disc galaxies have a wide distribution of Sérsic  $r_0$  (which is equal to the disk scale length for n = 1), ranging from 2.5 to 6.3 Kpc, where both n and  $r_0$  are taken from the NYU-VAGC catalog (Blanton et al. 2005a). To derive the rotation curve, we assume an exponential mass profile with a typical disk scale length of 4.4 Kpc for a massive disc galaxy and a NFW profile for its host dark matter halo. In Figure 10, the blue line shows the rotation curve for a halo mass of  $M_{\rm h} = 10^{12} {\rm M}_\odot$ (with concentration c = 13), about the average value for star-forming galaxies with  $\dot{M}_* \sim 10^{11} M_{\odot}$ . For comparison, we also show the rotation curve for a disc galaxy residing in a halo of  $10^{12.6}$  M<sub> $\odot$ </sub> (c = 11), as expected from our to-tal galaxy sample with  $M_* \sim 10^{11}$  M<sub> $\odot$ </sub> (see Figure 4). For  $M_h = 10^{12.6}$  M<sub> $\odot$ </sub>, the predicted rotation curve is quite flat at large radius, as is observed for many disk galaxies. In contrast, for  $M_{\rm h} = 10^{12} {\rm M}_{\odot}$ , the rotation curve reaches a peak at  $\sim 10 \,\mathrm{Kpc}$  and gradually decreases at larger radius. This type of rotation curve is not common for the general disk population, but has been found for some local galaxies (e.g. Corbelli et al. 2010; Posti et al. 2019). Modeling their rotation curves show that these galaxies indeed have a high conversion efficiency (Posti et al. 2019). Our results




Fig. 10. The figure shows rotation curves for two disk galaxies with the stellar mass of  $10^{11} M_{\odot}$  but with the halo mass of  $10^{12} M_{\odot}$  (star-forming, blue) and  $10^{12.6} M_{\odot}$  (total, green), respectively. The two disc galaxies have the same stellar mass exponential profiles with disk scale length of 4.4 Kpc and Sérsic index of n = 1. Dark matter halos are assumed to be NFW profile with concentration-mass relation from Zhao et al. (2009), c = 13 for the blue line and c = 11 for the green line.

show that this type of rotation curve should be expected for massive star-forming disks.

Finally, we examine whether current galaxy formation models can reproduce our results. In the right panel of Figure 5, we show the results obtained from the Illustris-TNG simulation (Pillepich et al. 2018), which implements a series of baryonic physics, such as AGN feedback, to suppress star formation in massive galaxies (see Appendix D for a brief description of the simulation). As can be seen from the figure, the simulations result for the total sample has a peak around a stellar mass between  $10^{10.5}$  and  $10^{11} M_{\odot}$ , consistent with the observational results. However, the simulation fails to reproduce the high conversion efficiency for the observed massive star-forming galaxies. Indeed, massive starforming galaxies in the simulation follow closely with the total population over almost the entire stellar mass range. It is likely that the AGN feedback implemented in the simulation is too strong for these galaxies. We have also examined the Eagle simulation (Schaye et al. 2015), and found also that it cannot reproduce the high conversion efficiency observed for massive star-forming galaxies. Thus, our finding presents a challenging problem for current simulations in their modeling of feedback and star formation.

#### 5. Summary

Based on the shear catalog of DECaLS imaging data, we derive the halo mass of central galaxies selected from the SDSS. We develop a weak lensing calibrated satellite kinematics method and to improve the halo mass measurements. We then obtain the efficiency for converting baryons into stars within halos, defined as  $M_*/M_{\rm h}/f_{\rm b}$ , for both the total galaxy population and galaxies in the star-forming main sequence. Our main results are summarized as follows.

- The stellar mass-halo mass relation for the total galaxy population we obtained is in good agreement with previous studies. The conversion efficiency peaks around Milky-Way-like galaxies and declines towards both lower and higher stellar mass ends.
- The conversion efficiency of star-forming galaxies increases monotonically with stellar mass, and reaches a value of more than sixty percents at  $M_* \gtrsim 10^{11} M_{\odot}$ . Thus, these galaxies have converted most of their halo gas into stars.
- Our tests show that the measurements of the halo mass are consistent with the results obtained from the SDSS shear catalog and from galaxy clustering.
- Massive star-forming galaxies are expected to have rotation curves that are peaked at about two disk scalelengths and decline at larger distances, quite different from the flat rotation curves commonly observed for the general disk population.
  The high conversion efficiency observed for massive star-
- The high conversion efficiency observed for massive starforming galaxies is not reproduced by current cosmological gas simulations.
- We have tested a number of systematic effects that may affect our results and found that none of them can change our conclusions significantly.

Our finding has important implications for understanding galaxy formation and star formation quenching. The high conversion efficiency observed for massive star-forming galaxies indicates that AGN feedback may not have played an important role in affecting the conversion of gas into stars in these particular galaxies. The fact that current cosmological hydrodynamic simulations cannot reproduce such a high conversion efficiency indicates that our current understanding of feedback is still incomplete, at least for massive star-forming galaxies. Clearly, the observational results presented here will provide an important constraint on modeling feedback processes in galaxy formation, and we will come back to this in a future paper.

#### Acknowledgments

We thank the referee for useful comments. This work is supported by the National Key R&D Program of China (grant No. 2018YFA0404503; grant No. 2018YFA0404504), the National Natural Science Foundation of China (NSFC, Nos. 11733004, 12192224, 11890693, 11421303, 11890691, 11621303, 11833005 and 11890692), and the Fundamental Research Funds for the Central Universities. The authors gratefully acknowledge the support of Cyrus Chun Ying Tang Foundations. We acknowledge the science research grants from the China Manned Space Project with No. CMS-CSST-2021-A03. The work is also supported by the Supercomputer Center of University of Science and Technology of China.

The Legacy Surveys consist of three individual and complementary projects: the Dark Energy Camera Legacy Survey (DECaLS; Proposal ID #2014B-0404; PIs: David Schlegel and Arjun Dey), the Beijing-Arizona Sky Survey (BASS; NOAO Prop. ID #2015A-0801; PIs: Zhou Xu and Xiaohui Fan), and the Mayall z-band Legacy Survey (MzLS; Prop. ID #2016A-0453; PI: Arjun Dey). DECaLS, BASS and MzLS together include data obtained, respectively, at the Blanco telescope, Cerro Tololo Inter-American Observatory, NSF's NOIRLab; the Bok telescope, Steward Observatory, University of Arizona; and the Mayall telescope, Kitt Peak National Observatory, NOIRLab. The Legacy Surveys project is honored to be permitted to conduct astronomical research on Iolkam Du'ag (Kitt Peak), a mountain with particular significance to the Tohono O'odham Nation.

NOIRLab is operated by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the National Science Foundation.

This project used data obtained with the Dark Energy Camera (DECam), which was constructed by the Dark Energy Survey (DES) collaboration. Funding for the DES Projects has been provided by the U.S. Department of Energy, the U.S. National Science Foundation, the Ministry of Science and Education of Spain, the Science and Technology Facilities Council of the United Kingdom, the Higher Education Funding Council for England, the National Center for Supercomputing Applications at the University of Illinois at Urbana-Champaign, the Kavli Institute of Cosmological Physics at the University of Chicago, Center for Cosmology and Astro-Particle Physics at the Ohio State University, the Mitchell Institute for Fundamental Physics and Astronomy at Texas A&M University, Financiadora de Estudos e Projetos, Fundacao Carlos Chagas Filho de Amparo, Financiadora de Estudos e Projetos, Fundacao Carlos Chagas Filho de Amparo a Pesquisa do Estado do Rio de Janeiro, Conselho Nacional de Desenvolvimento Cientifico e Tecnologico and the Ministerio da Ciencia, Tecnologia e Inovacao, the Deutsche Forschungsgemeinschaft and the Collaborating Institutions in the Dark Energy Survey. The Collaborating Institutions are Argonne National Laboratory, the University of California at Santa Cruz, the University of Cambridge, Centro de Investigaciones Energeticas, Medioambientales y Tecnologicas-Madrid, the University of Chicago, University College London, the DES-Brazil Consortium, the University of Edinburgh, the Eidgenossische Technische Hochschule (ETH) Zurich, Fermi National Accelerator Laboratory, the University of Illinois at Urbana-Champaign, the Institut de Ciencies de l'Espai (IEEC/CSIC), the Institut de Fisica d'Altes Energies, Lawrence Berkeley National Laboratory, the Ludwig Maximilians Universitat Munchen and the associated Excellence Cluster Universe, the University of Michigan, NSF's NOIRLab, the University of Nottingham, the Ohio State University, the University of Pennsylvania, the University of Portsmouth, SLAC National Accelerator Laboratory, Stanford University, the University of Sussex, and Texas A&M University.

BASS is a key project of the Telescope Access Program (TAP), which has been funded by the National Astronomical Observatories of China, the Chinese Academy of Sciences (the Strategic Priority Research Program "The Emergence of Cosmological Structures" Grant # XDB09000000), and the Special Fund for Astronomy from the Ministry of Finance. The BASS is also supported by the External Cooperation Program of Chinese Academy of Sciences (Grant # 114A11KYSB20160057), and Chinese National Natural Science Foundation (Grant # 11433005).

The Legacy Survey team makes use of data products from the Near-Earth Object Wide-field Infrared Survey Explorer (NEOWISE), which is a project of the Jet Propulsion Laboratory/California Institute of Technology. NEOWISE is funded by the National Aeronautics and Space Administration.

The Legacy Surveys imaging of the DESI footprint is supported by the Director, Office of Science, Office of High Energy Physics of the U.S. Department of Energy under Contract No. DE-AC02-05CH1123, by the National Energy Research Scientific Computing Center, a DOE Office of Science User Facility under the same contract; and by the U.S. National Science Foundation, Division of Astronomical Sciences under Contract No. AST-0950945 to NOAO.

The Photometric Redshifts for the Legacy Surveys (PRLS) catalog used in this paper was produced thanks to funding from the U.S. Department of Energy Office of Science, Office of High Energy Physics via grant DE-SC0007914.

#### References

- Abazajian, K. N., Adelman-McCarthy, J. K., Agüeros, M. A., et al. 2009, ApJS, 182, 543
- Abdullah, M. H., Wilson, G., Klypin, A., et al. 2020, ApJS, 246, 2
- Anderson, M. E., Bregman, J. N., & Dai, X. 2013, ApJ, 762, 106
- Barrow, J. D., Bhavsar, S. P., & Sonoda, D. H. 1984, MNRAS, 210, 19P
- Behroozi, P., Wechsler, R. H., Hearin, A. P., & Conroy, C. 2019, MN-RAS, 488, 3143
- Bilicki, M., Dvornik, A., Hoekstra, H., et al. 2021, A&A, 653, A82
- Blanton, M. R., Eisenstein, D., Hogg, D. W., Schlegel, D. J., & Brinkmann, J. 2005a, ApJ, 629, 143
- Blanton, M. R., Schlegel, D. J., Strauss, M. A., et al. 2005b, AJ, 129, 2562
- Bluck, A. F. L., Mendel, J. T., Ellison, S. L., et al. 2016, MNRAS, 462, 2559
- Brainerd, T. G. & Specian, M. A. 2003, ApJ, 593, L7
- Bregman, J. N. 2007, ARA&A, 45, 221
- Breiman, L. 2001, Machine Learning, 45, 5
- Brinchmann, J., Charlot, S., White, S. D. M., et al. 2004, MNRAS, 351.1151
- Cole, S., Lacey, C. G., Baugh, C. M., & Frenk, C. S. 2000, MNRAS, 319, 168
- Conroy, C., Prada, F., Newman, J. A., et al. 2007, ApJ, 654, 153
- Corbelli, E., Lorenzoni, S., Walterbos, R., Braun, R., & Thilker, D. 2010, A&A, 511, A89
- Croton, D. J., Springel, V., White, S. D. M., et al. 2006, MNRAS, 365.11
- Dekel, A. & Birnboim, Y. 2006, MNRAS, 368, 2
- Dey, A., Schlegel, D. J., Lang, D., et al. 2019, AJ, 157, 168
- Diemer, B. 2018, ApJS, 239, 35
- Dutton, A. A., Conroy, C., van den Bosch, F. C., et al. 2011, MNRAS, 416, 322
- Evrard, A. E., Bialek, J., Busha, M., et al. 2008, ApJ, 672, 122 Fabian, A. C. 2012, ARA&A, 50, 455
- Fall, S. M. & Efstathiou, G. 1980, MNRAS, 193, 189
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, PASP, 125, 306
- Gao, L., Springel, V., & White, S. D. M. 2005, MNRAS, 363, L66
- Heckman, T. M. & Best, P. N. 2014, ARA&A, 52, 589
- Hudson, M. J., Gillis, B. R., Coupon, J., et al. 2015, MNRAS, 447, 298
  - Kauffmann, G. & Charlot, S. 1998, MNRAS, 294, 705
  - Kauffmann, G., Heckman, T. M., Tremonti, C., et al. 2003a, MNRAS, 346, 1055
  - Kauffmann, G., Heckman, T. M., White, S. D. M., et al. 2003b, MN-RAS, 341, 33
  - Kravtsov, A. V., Vikhlinin, A. A., & Meshcheryakov, A. V. 2018, Astronomy Letters, 44, 8


Kurczynski, P., Gawiser, E., Acquaviva, V., et al. 2016, ApJ, 820, L1 Lan, T.-W. & Mo, H. 2018, ApJ, 866, 36

- Lange, J. U., van den Bosch, F. C., Zentner, A. R., Wang, K., & Villarreal, A. S. 2019a, MNRAS, 482, 4824
- Lange, J. U., van den Bosch, F. C., Zentner, A. R., Wang, K., & Villarreal, A. S. 2019b, MNRAS, 487, 3112
- Leauthaud, A., Finoguenov, A., Kneib, J.-P., et al. 2010, ApJ, 709, 97
- Leauthaud, A., Tinker, J., Bundy, K., et al. 2012, ApJ, 744, 159
- Li, C., Kauffmann, G., Jing, Y. P., et al. 2006, MNRAS, 368, 21
- Lim, S. H., Mo, H. J., Li, R., et al. 2018, ApJ, 854, 181

- Lu, Z., Mo, H. J., Lu, Y., et al. 2014, MNRAS, 439, 1294 Lu, Z., Mo, H. J., Lu, Y., et al. 2015, MNRAS, 450, 1604 Ludlow, A. D., Navarro, J. F., Springel, V., et al. 2009, ApJ, 692, 931

Article number, page 12 of 15

- Luo, W., Yang, X., Lu, T., et al. 2018, ApJ, 862, 4
- Luo, W., Yang, X., Zhang, J., et al. 2017, ApJ, 836, 38
- Mandelbaum, R., Li, C., Kauffmann, G., & White, S. D. M. 2009, MNRAS, 393, 377
- Mandelbaum, R., Seljak, U., & Hirata, C. M. 2008, JCAP, 2008, 006 Mandelbaum, R., Seljak, U., Kauffmann, G., Hirata, C. M., & Brinkmann, J. 2006, MNRAS, 368, 715
- Mandelbaum, R., Wang, W., Zu, Y., et al. 2016, MNRAS, 457, 3200
- Martig, M., Bournaud, F., Teyssier, R., & Dekel, A. 2009, ApJ, 707, 250
- McKay, T. A., Sheldon, E. S., Johnston, D., et al. 2002, ApJ, 571, L85
- Mo, H. J. & White, S. D. M. 1996, MNRAS, 282, 347
- More, S., van den Bosch, F. C., Cacciato, M., et al. 2011, MNRAS, 410, 210
- Moster, B. P., Naab, T., & White, S. D. M. 2013, MNRAS, 428, 3121
- Moster, B. P., Somerville, R. S., Maulbetsch, C., et al. 2010, ApJ, 710, 903
- Moustakas, J., Coil, A. L., Aird, J., et al. 2013, ApJ, 767, 50
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, ApJ, 490, 493 Pillepich, A., Springel, V., Nelson, D., et al. 2018, MNRAS, 473, 4077
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016, A&A, 594, A13
- Posti, L. & Fall, S. M. 2021, A&A, 649, A119
- Posti, L., Fraternali, F., & Marasco, A. 2019, A&A, 626, A56
- Rines, K., Geller, M. J., Diaferio, A., & Kurtz, M. J. 2013, ApJ, 767, 15
- Rines, K. J., Geller, M. J., Diaferio, A., & Hwang, H. S. 2016, ApJ, 819, 63
- Rodríguez-Puebla, A., Avila-Reese, V., Yang, X., et al. 2015, ApJ, 799.130
- Saintonge, A., Catinella, B., Cortese, L., et al. 2016, MNRAS, 462, 1749
- Saintonge, A., Catinella, B., Tacconi, L. J., et al. 2017, ApJS, 233, 22
- Salim, S., Lee, J. C., Janowiecki, S., et al. 2016, ApJS, 227, 2 Schaye, J., Crain, R. A., Bower, R. G., et al. 2015, MNRAS, 446, 521 Seo, G., Sohn, J., & Lee, M. G. 2020, ApJ, 903, 130
- Shan, H., Kneib, J.-P., Li, R., et al. 2017, ApJ, 840, 104
- Shimasaku, K., Fukugita, M., Doi, M., et al. 2001, AJ, 122, 1238
- Silk, J. & Rees, M. J. 1998, A&A, 331, L1
- Speagle, J. S., Steinhardt, C. L., Capak, P. L., & Silverman, J. D. 2014, ApJS, 214, 15
- Springel, V. 2010, MNRAS, 401, 791
- Strateva, I., Ivezić, Ż., Knapp, G. R., et al. 2001, AJ, 122, 1861
- Tinker, J. L., Leauthaud, A., Bundy, K., et al. 2013, ApJ, 778, 93
- Tinker, J. L., Robertson, B. E., Kravtsov, A. V., et al. 2010, ApJ, 724.878
- Tumlinson, J., Peeples, M. S., & Werk, J. K. 2017, ARA&A, 55, 389 van den Bosch, F. C., Aquino, D., Yang, X., et al. 2008, MNRAS, 387, 79
- van den Bosch, F. C., Norberg, P., Mo, H. J., & Yang, X. 2004, MN-RAS, 352, 1302
- van Uitert, E., Hoekstra, H., Velander, M., et al. 2011, A&A, 534, A14
- Velander, M., van Uitert, E., Hoekstra, H., et al. 2014, MNRAS, 437, 2111
- Viola, M., Cacciato, M., Brouwer, M., et al. 2015, MNRAS, 452, 3529
- Wang, H., Mo, H. J., & Jing, Y. P. 2009, MNRAS, 396, 2249
- Wang, H., Zhang, J., Li, H., & Shen, Z. 2021, ApJ, 911, 10 Wang, H. Y., Mo, H. J., & Jing, Y. P. 2007, MNRAS, 375, 633
- Wang, L. & Li, C. 2019, MNRAS, 483, 1452
- Wechsler, R. H. & Tinker, J. L. 2018, ARA&A, 56, 435
- Whitaker, K. E., Franx, M., Leja, J., et al. 2014, ApJ, 795, 104 White, S. D. M. & Rees, M. J. 1978, MNRAS, 183, 341
- Wojtak, R. & Mamon, G. A. 2013, MNRAS, 428, 2407
- Xu, K. & Jing, Y. 2022, ApJ, 926, 130
- Yang, X., Mo, H. J., & van den Bosch, F. C. 2003, MNRAS, 339, 1057
- Yang, X., Mo, H. J., & van den Bosch, F. C. 2009, ApJ, 695, 900 Yang, X., Mo, H. J., van den Bosch, F. C., et al. 2007, ApJ, 671, 153 Zhang, J., Dong, F., Li, H., et al. 2019, The Astrophysical Journal, 875, 48
- Zhang, J., Luo, W., & Foucaud, S. 2015, JCAP, 2015, 024
- Zhang, J., Zhang, P., & Luo, W. 2017, ApJ, 834, 8
- Zhang, Z., Wang, H., Luo, W., et al. 2021, A&A, 650, A155 Zhao, D. H., Jing, Y. P., Mo, H. J., & Börner, G. 2009, ApJ, 707, 354
- Zhou, R., Newman, J. A., Mao, Y.-Y., et al. 2021, MNRAS, 501, 3309
- Zou, H., Gao, J., Zhou, X., & Kong, X. 2019, The Astrophysical Journal Supplement Series, 242, 8
- Zu, Y. & Mandelbaum, R. 2015, MNRAS, 454, 1161



Appendix A: Testing the impact of halo mass uncertainties on  $\sigma_s$ 

Fig. A.1. Comparison of the satellite velocity dispersion obtained using three different halo masses in the estimates of  $r_{\rm vir}$ and  $v_{\rm vir}$ . The left column shows the results for star-forming samples, and the right column for total samples. Different rows represent different stellar mass bins. In each panel, the three data points are for halo masses that correspond to 16%, 50% and 84% percentage points of the mass distribution, respectively, as labeled in the horizontal axis.

We use the halo mass to determine  $r_{\rm vir}$  and  $v_{\rm vir}$  that are used to identify satellite candidates around centrals. The satellite velocity dispersion can thus be affected by the uncertainties in the halo mass obtained from weak lensing. To test this, we adopt two additional halo masses to derive the velocity dispersion. These two halo masses correspond to the 16 and 84 percentiles of the posterior distribution ob-

Article number, page 14 of 15

tained from the MCMC fitting to the stacked lensing profiles, and may be considered as the lower and upper halo mass limits, respectively. Figure A.1 compares the  $\sigma_s$  obtained from the two halo mass limits and the medium halo mass adopted in the main text. The results show that, for most of the stellar mass bins where the sample sizes are sufficiently large, the value of  $\sigma_s$  is not affected significantly by the halo mass uncertainty. For some of the stellar mass bins where the sample sizes are small, such as the lowest stellar mass bin (see Table 1), the halo mass uncertainty is large and may have a significant impact on the estimate of  $\sigma_s$ .

# Appendix B: The $M_{\rm h}$ - $\sigma_{\rm s}$ relations for different stellar mass bins

To avoid the correlation between the uncertainties in  $\sigma_{\rm s}$  and in the  $M_{\rm h}$ - $\sigma_{\rm s}$  relation, we derive the  $M_{\rm h}$ - $\sigma_{\rm s}$  relation using a slightly different method. For a given galaxy sample in a stellar mass bin, we use the complementary sample consisting of all galaxies in other stellar mass bins to fit the  $M_{\rm h}$ - $\sigma_{\rm s}$  relation. The relations so obtained for different stellar mass bins are shown in Table B.1. As one can see, these relations are consistent with each other and Eq. 4 within error bars. It is clear that the  $\sigma_{\rm s}$  of a galaxy sample in a stellar mass bin is quite independent of the complementary sample used for the calibration. Thus, we can use the value of the  $\sigma_{\rm s}$  for the galaxy sample, in combination with the corresponding  $M_{\rm h}$ - $\sigma_{\rm s}$  relation, to derive the halo mass and conversion efficiency for the sample. The results are shown in Table 1.

 Table B.1. Halo mass-satellite velocity dispersion relations.

$\log M_*$ range	$\sigma_s M_h$ relation <sup>(b)</sup>
[8.84, 9.34]	$\log(M_{\rm h}/M_{\odot}) = (3.08 \pm 0.05) \log(\sigma_s/{\rm s}^{-1} {\rm km}) + (5.35 \pm 0.11).$
[9.34, 9.84]	$\log(M_{\rm h}/M_{\odot}) = (3.09 \pm 0.05) \log(\sigma_s/s^{-1} \text{ km}) + (5.32 \pm 0.12).$
[9.84, 10.34]	$\log(M_{\rm h}/M_{\odot}) = (3.07 \pm 0.05) \log(\sigma_s/s^{-1} \text{ km}) + (5.37 \pm 0.12).$
[10.34, 10.84]	$\log(M_{\rm h}/M_{\odot}) = (3.08 \pm 0.05) \log(\sigma_s/s^{-1} \text{ km}) + (5.34 \pm 0.13).$
[10.84, 11.34]	$\log(M_{\rm h}/M_{\odot}) = (3.05 \pm 0.05) \log(\sigma_s/s^{-1} \text{ km}) + (5.42 \pm 0.11).$
[11.34, 11.84]	$\log(M_{\rm h}/M_{\odot}) = (2.92 \pm 0.1) \log(\sigma_s/{\rm s}^{-1} {\rm km}) + (5.71 \pm 0.22).$
(n)	

<sup>(a)</sup> The stellar mass bins.

 $^{\rm (b)}$  The corresponding  $\sigma_s\text{-}M_{\rm h}$  relations.

### Appendix C: Measurements with SDSS shear catalog

As an independent test, we also use a different shear catalog (Luo et al. 2017) based on SDSS DR7 (Abazajian et al. 2009). The difference between the two samples is twofold. First, the shape measurement method here uses a traditional second-moment estimator to evaluate the ellipticity; secondly, the coverage of the SDSS DR7 is much larger than the DECaLS region overlapping with our lens samples. There are 190,730 galaxies with  $M_* \ge 10^{8.8} \mathrm{M}_{\odot}$  in the starforming sample, and 445,135 galaxies with  $M_* \geq 10^{8.8} M_{\odot}$ in the total sample. However, the deeper imaging, together with the PDF-symmetrization method, gives much smaller statistical errors than the SDSS DR7 catalog. We repeat the modeling as described above to extract the halo mass. The results are consistent with those obtained from the DECaLS shear data, but with larger error bars as shown in Figure 5. The ESDs and best-fitting results obtained from the SDSS shear catalog are presented in Figure C.1.





Fig. C.1. The same as in Figure 1 but for SDSS shear catalog.

### Appendix D: The IllustrisTNG Simulation used for comparison

IllustrisTNG The simulations are run with AREPO(Springel 2010), which implements a movingmesh technique. The simulations include subgrid physical models for gas cooling, star formation, metal enrichment, and stellar and AGN feedback. In this paper, we use the simulation TNG100-1, which has a box size of  $\sim 100$  Mpc. The dark matter particle mass is  $7.5\times10^6 M_{\odot}$  and the average gas cell mass is about  $1.39 \times 10^6 M_{\odot}$ . Galaxies with  $M_* \geq 10^{8.8} M_{\odot}$  are well resolved in the simulation. Central galaxies are defined as the most massive galaxies in their host halos. Galaxy stellar mass is the sum of all stellar particles within the gravitationally bound substructure, and the SFR is calculated from all gas cells in the same region. The halo mass, directly taken from the simulation, is the mass contained in spherical regions, within each of which the mean mass density is 200 times of the cosmic mean matter density.

In Figure D.1, we show the SFR-stellar mass relation for central galaxies in the simulation. The simulation can well reproduce the star-forming main sequence, and most star-forming galaxies in the stellar mass range plotted are well above the demarcation line used in our observational data. We thus decide to use the same demarcation criteria to identify star-forming galaxies. The efficiency as a func-



**Fig. D.1.** The SFR- $M_*$  diagram for model galaxies. The figure shows the result for central galaxies in IllustrisTNG. The solid line in the panel is the demarcation line used to identify star-forming galaxies, which is the same as the one used in the observation. Note that a large fraction of the galaxies have very low SFR. These galaxies fall outside the boundary of the plot.

tion of stellar mass for both star-forming galaxies and total galaxy populations is presented in Figure 5.