

ALP Searches at the LHC: FASER as a Light Shining through Walls Experiment

Felix Kling ^{1,*} and Pablo Quílez ^{1,†}

¹*Deutsches Elektronen-Synchrotron DESY, Germany*

We propose the use of FASER as a light-shining-through-walls experiment to search for axions and axion-like particles (ALPs). LHC collisions generate a high intensity and high energy photon flux in the forward direction which can oscillate into ALPs in the magnetic fields that are used to confine the beam. These ALPs then pass through about 100 m of rock before reaching the magnetic fields of FASER, where they can convert back into photons and be detected by an electromagnetic calorimeter. In the next years, FASER and its successor FASER2 at the Forward Physics Facility will be able to explore regions of the ALP parameter space inaccessible by other laboratory-based experiments.

I. INTRODUCTION

Axions are among the best motivated scenarios of new physics beyond the Standard Model (BSM). Most famously, the QCD axion arises as a low-energy consequence of the elegant Peccei-Quinn solution to the strong CP problem of the SM [1–6]. But the interest in axion-like particles goes well beyond the QCD axion. Indeed these pseudo-Nambu-Goldstone bosons arise in a plethora of BSM constructions such as theories with extra dimensions [7], Majoron models [8], theories of flavor [9, 10] and are ubiquitous within string theory [11]. They all share the basic properties that stem from its pseudo Nambu-Goldstone nature: axions are light and present feeble derivative or anomalous interactions suppressed by the axion decay constant f_a . Moreover, they are excellent dark matter candidates [12–14].

As a consequence, a strong experimental program has been developed and in the next decades we will explore large regions of the ALP parameter space. Most of the experiments only take advantage of the ALP coupling to photons. In this case the relevant Lagrangian reads

$$\mathcal{L} \supset -\frac{1}{2}m_a^2a^2 - \frac{1}{4}g_{a\gamma\gamma}aF_{\mu\nu}\tilde{F}^{\mu\nu} \quad (1)$$

where $F_{\mu\nu}$ is the electromagnetic field strength and $\tilde{F}^{\mu\nu}$ its dual. The resulting parameter space is therefore spanned by the ALP mass m_a and the ALP coupling $g_{a\gamma\gamma}$. ALP search experiments utilize a variety of experimental techniques that range from laboratory-based light-shining-through walls (LSW) experiments such as ALPS II [15], helioscopes looking for solar ALPs such as CAST [16, 17] and the future IAXO [18, 19], haloscopes looking for dark matter ALPs [20–25] but also colliders [26, 27], flavor probes [28–31], as well as beam dump [32] and neutrino [33, 34] experiments for larger masses. There are also a number of astrophysical constraints (typically through energy-loss arguments) and cosmological bounds which provide a nice complemen-

tarity with respect to proposed experiments. However, these considerations vary in their robustness. For example, the supernova bound from the SN1987 neutrino signal [35] has been recently put in question [36] although not everyone in the community agrees with this criticism and it has been argued that the SN1987 also constrains the ALP through the γ -ray signal and the diffuse cosmic γ -ray background [37]. Similarly some cosmological bounds can be significantly relaxed in non-standard cosmological scenarios [38, 39]. This is why there is a growing community interest in purely laboratory based ALP experiments, which has been advocated for (e.g. in Ref. [40]).

In this letter we contribute to this effort by noting that the LHC accelerator in combination with the recently installed FASER experiment [41–43] can be used as a LSW experiment without requiring any modification of the current experimental setup. The LSW detection principle is based on the coherent ALP-photon oscillation in the presence of an external magnetic field. A strong photon beam is directed at a thick wall with a photon detector placed behind it. In the absence of ALPs, no signal is expected in the detector. However, if ALPs exist the strong magnetic fields perpendicular to the beam located at both sides of the wall may induce the photons to convert to ALPs which can freely pass through the wall. These ALPs can then oscillate back to photons and be detected on the other side of the wall. Effectively, this double conversion mechanism allows a small fraction of the photon beam to pass through the wall where it can be detected. LSW experiments use this mechanism to search for ALPs.

In this study we explore the sensitivity of both FASER and future forward detectors proposed in the context of the Forward Physics Facility (FPF) [46, 47] for light ALP searches. The main advantage of this proposal with respect to existing LSW experiments lies on the high energies of LHC photons which allows us to explore higher ALP masses. With respect to other accelerator searches, this proposal enjoys a better sensitivity at lower masses since it relies on coherent conversion in an external macroscopic magnetic field instead of production/detection through scattering or perturbative decay. In the following, we will discuss the production of ALPs

*Electronic address: felix.kling@desy.de

†Electronic address: pablo.quilez@desy.de




FIG. 1: Considered magnetic fields at the HL-LHC (top) and LHC Run 3 (bottom) downstream from the ATLAS collisions point. We show dipoles in blue and quadrupoles in red, and indicate their corresponding length L , magnetic field B and field gradient B' . Here we have used the magnetic fields as presented in Refs. [44] and [45].

at the LHC in Section II, their detection in Section III, and the resulting sensitivity in Section IV.

II. ALP PRODUCTION AT THE LHC

Since more than half a century, hadron colliders have been the primary tools to probe the fundamental laws of nature at the highest energies. The currently most energetic realization of these hadron collider experiments is the LHC at CERN, which started to collect data in 2010 and is expected to run until 2040. With a collision energy of 14 TeV, it is able to produce heavy particles with masses at the weak scale and beyond, and large detectors like ATLAS and CMS have been build around the LHC collision point to detect their decay products. Their biggest achievement thus far is the discovery of the Higgs boson in 2012 [48, 49] and the main objectives for their remaining operation are to study the properties of the Higgs and to search for new heavy particles at the TeV scale. In addition, the LHC also produces an enormous number of energetic light particles, which are mainly produced in the forward direction. As first pointed out by the authors of Ref. [41], this provides an additional opportunity to search for light and weakly coupled particles. In the following, we will apply this idea in the context of ALPs.

As a byproduct of proton-proton collisions, the ATLAS experiment also produces an intense and strongly collimated beam of photons with energy $E_\gamma \sim \text{TeV}$ in the far forward direction. These photons mainly result from the decay of neutral pions, which were for example formed in the hadronization of the beam remnants. In this study, we use the double differential photon spectrum $d^2\sigma/dEd\theta$ (with respect to the photon energy and angle with respect to the beam axis) provided by the FORESEE [50] tool, which were obtained by the EPOS-LHC [51] event generator as implemented in the CRMC [52] package. The

photon distribution is shown in Fig. 2. Roughly speaking, we expect about 10^{14} photons per fb^{-1} of integrated luminosity to be produced within 1 mrad of the beam axis.

These photons travel down the beam pipe, where they pass through a variety of magnets before being absorbed in the LHC infrastructure. An overview of the magnetic fields that are relevant for our study is presented in Fig. 1. This includes the so called inner triplet, which is a set of quadrupole magnets to focus the proton beam before the collision, and the D1 dipole magnet, which separates the incoming and outgoing proton beams before they enter their separate beam pipes. Note that the exact dimensions and strength of these magnets will undergo some changes in the future. The LHC in its current configuration will operate until about 2025. Afterwards, it will be upgraded in preparation of the following high-luminosity LHC (HL-LHC) era. When modelling the magnetic fields, we assume that the dipole magnets have a constant field strength B , while quadrupole magnets are specified by a field gradient B' and their field strength is given by $B = B' \cdot r$ where r denotes the distance from the center [53]. We further model the beam pipe as a cylinder with diameter of 5.3 cm and assume that photons are quickly absorbed once they leave the vacuum.

When passing these strong magnetic fields, the photons could oscillate into ALPs. The corresponding conversion probability can be written as [54]

$$P_{\gamma \rightarrow a} \simeq \frac{1}{2} \times \frac{g_a^2}{4} \left| \int dz \vec{B}(z) e^{iqz} \right|^2, \quad (2)$$

where we integrate the magnetic field coherently along the trajectory of the photon. The additional factor 1/2 in front accounts for the fact that the photon beam is unpolarized and only the photons with a polarization parallel to the magnetic field can convert into ALPs since $F\bar{F} \sim \vec{E} \cdot \vec{B}$. Here $q = E_\gamma - (E_\gamma^2 - m_a^2)^{1/2} \simeq m_a^2/(2E_\gamma)$ is




FIG. 2: Differential production rate of photons at 14 TeV LHC predicted by EPOS-LHC in the (θ, p) plane, where θ and p are the proton's angle with respect to the beam axis and momentum, respectively. The angular coverage for FASER and FASER 2 are indicated by the vertical dotted lines.

the momentum transfer. In our case, the magnetic fields along a single photon's trajectory are piece-wise constant which allows us to write

$$P_{\gamma \rightarrow a} \simeq \frac{g_{a\gamma\gamma}^2}{8} \left| \sum_i \vec{B}_i L_i e^{iqz_i} \frac{\sin(qL_i/2)}{qL_i/2} \right|^2. \quad (3)$$

where \vec{B}_i is the field strength of the dipole and quadrupole magnets seen by the photon, L_i is the length and z_i is the central position of each magnet. Using that the typical length of the magnets is $L \sim 100$ m and the typical energy of the photons is $E_\gamma \sim 1$ TeV, we can distinguish two relevant kinematic regimes. For small masses $m_a \ll \mathcal{O}(100)$ eV and hence $qL \ll 1$, the conversion takes place coherently along the full length of the magnetic field. This can be understood from the fact that the phase velocity of both the ALP and the photon are approximately the same and thus the probability amplitude of the conversion at any point interferes constructively with that in the rest of the conversion region. As a consequence of the probability is enhanced, scaling with the square of the length $\propto L^2$ and becomes mass independent. However, for large masses $m_a \gg \mathcal{O}(100)$ eV and hence $qL \gg 1$ the ALP and photon states do not remain in phase along all the conversion length. Hence, the probability becomes suppressed $P_{\gamma \rightarrow a} \sim 1/m_a^4$ and other production mechanisms become more dominant.

Eventually, the photons produced at the LHC will collide with elements of the LHC infrastructure and be absorbed. For very forward going photons, this occurs in the TAN neutral particle absorber, which is located at about 140 m downstream from the ATLAS interaction point. This provides a further opportunity for ALP production via the Primakoff process, in which the photon

converts into an axion when interacting with nuclei in the target material. In the limit of small ALP masses m_a and small momentum transfer q , the corresponding cross section can be written as [55–57]

$$\sigma_{\text{Prim}} \simeq \frac{g_{a\gamma\gamma}^2}{2} \alpha Z^2 \int F^2(q^2) d\log q = 9.8 \times g_{a\gamma\gamma}^2. \quad (4)$$

Here we have assumed that the target material is made of iron with atomic number $Z = 26$ and integrated the form factor $F(q^2)$ as in used in Refs. [56, 57]. The photon to ALP conversion probability is given by

$$P_{\gamma \rightarrow a} = \frac{\sigma_{\text{Prim}}}{\sigma_{\text{conv}}} = 7.4 \cdot 10^{-4} \times (g_{a\gamma\gamma} \cdot \text{GeV})^2, \quad (5)$$

where $\sigma_{\text{conv}} \simeq 5$ barn is the conventional cross section for the photon interaction with the target material (dominated by pair-production). At the LHC, this production mechanism dominates for high ALP masses $m_a > 1$ keV.

After the TAN neutral particle absorber, essentially all photons originating from the primary collision have been absorbed. The produced ALPs will then continue to propagate along the beam collision axis. Roughly 270 m downstream from the ATLAS interaction point, the LHC magnets start to deflect the beam to bring it on its roughly circular orbit and LHC tunnels curve away from the beam collision axis.

III. AXION DETECTION

To detect the produced beam of ultrarelativistic ALPs an additional magnetic field is needed to convert them back into photons. As we will see, the recently installed FASER experiment is already placed directly in the beam of LHC ALPs and equipped with a strong magnetic field to perform that task.

About 480 m downstream from ATLAS, the beam collision axis intersects with the service tunnel TI12. This location provides a unique opportunity to access the beam of light and weakly interacting particles that are produced in the forward direction of the LHC [41]. The FASER experiment has recently been installed there to exploit this opportunity [41–43, 58–60]. The main goal of FASER search for light long-lived particles at the LHC [41, 61–64]. In addition, the experiment also contains the FASERν neutrino detector which will for the first time detect and study of TeV energy neutrinos produced at the LHC [65–67]. The FASER experiment will operate during LHC Run 3, from 2022 to 2025, with an expected integrated luminosity of 150 fb^{-1} .

The experiment consists of a decay volume and tracking spectrometer which are placed inside an otherwise empty magnetized area of length of 3.5 m, diameter of 20 cm and field strength of 0.6 T. Located behind the spectrometer is a preshower detector [68] for photon identification and an electromagnetic calorimeter to measure the photon energy. FASER therefore already provides an

ideal setup for ALP detection. The corresponding probability of ALP to photon conversion can be written as

$$P_{a \rightarrow \gamma} \simeq \frac{g_{a\gamma\gamma}^2}{4} \left| \int dz B(z) e^{iqz} \right|^2 = \frac{g_{a\gamma\gamma}^2}{4} B^2 L^2 \frac{\sin^2(qL/2)}{(qL/2)^2}. \quad (6)$$

Similarly to the production, the conversion probability is mass independent for small ALP masses and decreases for $m_a \gg 100$ eV. The expected signal then would be a single high energy photon in the preshower detector and electromagnetic calorimeter¹.

A continuation of FASER's physics program during the HL-LHC era has been proposed in the context of the FPF [46, 47]. The FPF would be a new cavern located about 620 m downstream from ATLAS where it will provide space for a variety of experiments and collect 3 ab^{-1} of data. In particular, this would include a larger version of the FASER detector, called FASER2. This detector is envisioned to be roughly 20 m long with an aperture of 2 m and a magnetic field of 1 T which would increase the ALP to photon conversion probability by roughly two orders of magnitude in comparison to FASER.

In addition to FASER2, the FPF contains three neutrino detectors: the emulsion based neutrino detector FASER ν 2, the electronic neutrino detector AdvSND and the noble gas neutrino detector FLArE. These could provide an additional opportunity for ALP detection, as these could also undergo the inverse Primakoff scattering process and convert into photons via collisions with the detector material. The signal would be a single highly energetic photon that emerges inside the detector. Perhaps the best suited detector for this purpose is FLArE as it provides timing capabilities, a high spatial resolution, particle identification capabilities as well as a good energy estimate. In particular the particle identification would allow to distinguish photons and electrons, and hence suppress a possible background from neutrino electron scattering that would also provide an isolated electromagnetic shower. The FLArE detector is envisioned to have a $1\text{m} \times 1\text{m} \times 7\text{m}$ target volume filled with liquid argon. The corresponding ALP conversion probability is given by

$$P_{a \rightarrow \gamma} = \sigma_{\text{Prim}} n_{Ar} L = 0.028 \times (g_{a\gamma\gamma} \cdot \text{GeV})^2, \quad (7)$$

where n is the number density of liquid argon nuclei and σ_{Prim} is the cross section for the Primakoff process defined in Eq. (4), which has now been evaluated for $Z = 18$.

IV. RESULTS AND DISCUSSION

We can now combine our previous results to estimate the sensitivity of the LSW setup at the LHC to probe ALPs². For this, we compute the expected number of events as

$$N = \mathcal{L} \int dE \int d\theta \frac{d^2\sigma}{dEd\theta} \cdot P_{\gamma \rightarrow a}(E) \cdot P_{a \rightarrow \gamma}(E), \quad (8)$$

where \mathcal{L} is the LHC's integrated luminosity, $d^2\sigma/(dEd\theta)$ is the double differential photon production cross section with respect to the photon energy and angle with respect to the beam axis, $P_{\gamma \rightarrow a}$ is the probability of photon to ALP conversion in the LHC infrastructure and $P_{a \rightarrow \gamma}$ is the probability of ALP to photon conversion in the detector. The angular integral is performed within the detector's angular acceptance. In order to obtain the sensitivity contours we require the number of events to be larger than 3 over the full run time of the experiment for a point in the parameter space to be excluded at 95% confidence level. Here we use that the converted photons can be identified by FASER's electromagnetic calorimeter and preshower detector and assume that backgrounds can be reduced to a negligible level by the experiment's veto system. A more detailed discussion of possible background sources for the mono-photon signature and how they can be rejected can be found in Ref. [71] in the context of sterile neutrino decays $N \rightarrow \nu\gamma$ arising in neutrino dipole portal models.

We present our obtained sensitivity on the ALP parameter space in Fig. 3 alongside existing constraints. In the left panel, we show the broad picture including all existing constraints on ALPs arising from helioscopes [72–75], haloscopes [76–100] and LSW experiments [101–106] (red); astrophysics³ [37, 109–143] (green); cosmology [38, 144] (blue); and accelerator experiments [32, 145–159] (gray). The projected sensitivity for searches at FASER and future experiments at the FPF is indicated by the blue solid lines. Even though the regions that can be explored by LSW searches at FASER and FASER2 are excluded by helioscopes, astrophysical or cosmological arguments, these searches will be able to probe regions of the parameter space that have not been explored by any purely laboratory based experiment. This is shown in the right panel, where we show the bounds from current ground-based experiments together with the projections of future experiments and proposals. Note that these experiments are based on different ALP detection strategies which do not rely on extra astrophysical or cosmological assumptions and thus can be considered specially

¹ An interesting characteristic of the photons produced via $a \rightarrow \gamma$ conversion in FASER is that they are linearly polarized in the direction of the magnetic field since $F\tilde{F} \sim \vec{E} \cdot \vec{B}$. However, this feature is not experimentally accessible at FASER and thus FASER cannot distinguish between a scalar and a pseudoscalar.

² A notebook containing this analysis has been added to the `FORESEE` package

³ The supernova bounds are taken from Refs. [37, 107, 108] and should be considered conservative regarding the upper end of the bound that corresponds to the trapping limit [37].




FIG. 3: Sensitivity projections of FASER, FASER2 and FLArE as a LSW experiment for ALP searches (shown as blue solid lines) as compared with all current ALP bounds (left) and only purely laboratory based bounds and projections (right). The theoretical predictions for different QCD axion models are shown in orange where $g_{a\gamma\gamma} = \alpha/(2\pi f_a)(E/N - 1.92)$ [69]. See text for details. We provide the data of the sensitivity projections of this work as arXiv ancillary files. Existing experimental bounds adapted from [AxionLimits](#) [70].

robust. Classical LSW experiments such as ALPS [101], PVLAS [104], CROWS [102] and OSQAR [103] as well as a search at NOMAD [145] utilize the coherent conversion of ALPs into photons in an external magnetic field and are shown in reddish tones. In contrast, many high energy physics experiment look for the ALP decay into photons and thus are sensitive to heavier ALPs with larger couplings, as shown in grey tones. This includes both searches for long-lived ALPs at beam dump experiments such as CHARM [151], E141 [152, 153], E137 [154, 155], NuCal [32, 156, 160] and NA64 [157] as well as resonance searches for promptly decaying ALPs at Belle 2 [158], PrimEx [159] and LEP [146–150].

As explained in the previous section, our LSW proposal at the LHC benefits from both the coherent conversion in the external magnetic fields and the Primakoff conversion via scattering with nuclei in the target material. The LSW@FASER line corresponds to the reach of the currently installed FASER detector, while LSW@FASER2 represents the reach that will be achieved by the future FASER2 detector at the FPF. Similarly to other LWS experiments it is easy to understand the shape of the bound. For small ALP masses, $m_a \ll \mathcal{O}(100)$ eV the factor $qL \ll 1$ and the conversion probability in Eq. (3) becomes independent on the ALP mass giving rise to a flat bound on $g_{a\gamma\gamma}$. For higher ALP masses, $qL \gg 1$, the coherence is lost and the probability is suppressed by $1/m_a^8$ (the suppression factor $1/q^2 \sim E_\gamma^2/m_a^4$ arises both at production and detection). The bound on $g_{a\gamma\gamma}$ thus grows as $\propto m_a^2$. For even larger masses, the Primakoff conversion in nuclei becomes the dominant ALP

production channel, therefore one of the suppression factors disappears and the bound on $g_{a\gamma\gamma}$ grows only as $\propto m_a$. Furthermore, we also show the projections for the neutrino detector FLArE that will be installed as part of the FPF and could detect the ALPs via Primakoff conversion in nuclei and becomes dominant for larger masses. At large masses $m_a \gtrsim 10$ keV, both the production and detection proceeds through the Primakoff process and the sensitivity in $g_{a\gamma\gamma}$ becomes independent of the ALP mass. At high masses and large couplings the ALP lifetime becomes so small that the majority of ALPs decay before reaching the detector, hence limiting the reach at FLArE. ALP decays within the detector volume also provide an additional search channel that has been investigated in Ref. [63, 64]. The corresponding reach of these long-lived particle searches is shown by the LLP@FASER and LLP@FASER2 lines.

For comparison, we also show the projected sensitivity of other proposed searches and experiments for ALPs as dashed lines in Fig. 3. Using a high intensity LASER beam and optical resonant cavities, the ALPS II experiment [106] plans to probe ALP masses smaller than $\sim 10^{-4}$ eV with exceptional sensitivity down to couplings of $\sim 2 \times 10^{-11} \text{ GeV}^{-1}$. A proposal to probe higher masses has been recently put forward in Ref. [161] for the International Linear Collider (ILC) [162] using the photon beam that would be required for positron production followed by 4km of 1T magnets (adding up conversion and re-conversion lengths). As for the configuration of the magnetic field, on top of the uniform case they also consider non-homogeneous (or *wiggled*) magnetic field pro-

files which improve the sensitivity for higher ALP masses $m_a \gtrsim 0.1\text{eV}$ (shown with a dotted line). In addition, the authors of Refs. [163, 164] have proposed a class of experiments dubbed PASSAT which use helioscope magnets as detectors for ALPs produced via Primakoff effect at the target of beam dump experiments. In Fig. 3, we show two realizations of this idea, one using the CAST magnets placed at the proposed beam dump facility (BDF) and one using the multi-purpose detector (MPD) of the proposed DUNE experiment. To probe ALPs with even higher masses, several searches for their interactions via inverse Primakoff scattering as well as their decays at neutrino experiments have been proposed. We show the corresponding sensitivity both for CONUS as an example of coherent elastic neutrino scattering experiments at nuclear reactors [165] as well as DUNE as example of high energy accelerator experiments [33]. Further searches for the decay of long-lived ALPs have also been proposed for other beam dump type experiments, see Ref. [32] or Ref. [166] for recent reviews.

Regarding the possible UV complete models that FASER as a LSW experiment will be able to explore, we find that the projections are not able to reach the KSVZ and DFSZ axion benchmarks and they barely touch the preferred QCD axion window [167–169], which is illustrated by the diagonal yellow shaded region. Therefore this proposal mainly probes other axion-like particles. Nonetheless, FASER could still probe QCD axions that solve the strong CP problem in the context of photophilic axion models where the axion coupling to photon is enhanced [170–172] or the Z_N axion model [173–175] where the axion mass is suppressed.

Another interesting feature of our proposal is the ability of measuring the ALP mass given a positive signal. Both helioscopes and LSW experiments, provide the best bounds in the maximum coherence limit $qL \ll 1$ and when that condition is fulfilled they are not sensitive to the ALP mass. However, for ALP masses such that $qL \sim 1$ the conversion probability depends on the energy of the photon (as one can see by substituting $q \simeq m_a^2/(2E_\gamma)$ in Eq. (6)). This oscillatory behaviour as a function of the photon energy depends on the axion mass and thus may allow to extract it [176, 177]. Our proposal is most competitive in the region where the ALP-photon conversion is suppressed due to the loss of coherence $qL \gtrsim \mathcal{O}(1)$ and therefore in the hypothetical

scenario of a positive detection we could benefit from the excellent energy resolution of the FASER calorimeter [43] of about 1% to extract the ALP mass.

It is also worth emphasizing that the coherent ALP-photon oscillation in a magnetic field could be relevant for other ALP searches where the ALP couples not only to photons but also to other particles such as electrons or heavy electroweak gauge bosons. For example, some searches could use coherent conversion for ALP production and ALP decay or scattering via the coupling to electrons for detection. Alternatively, ALPs could be produced through another mechanism (e.g. rare meson decay via the ALP coupling to electroweak gauge bosons or quarks) and be detected via coherent conversion [30, 178, 179].

Regarding the outlook and possible improvements of this idea, there exist several techniques that could enhance the reach of this proposal and have already been proposed in the past for helioscopes or LSW [180, 181]. One option is the use of a buffer gas in the conversion cavity that generates an effective photon mass that maintains the coherence of the oscillation for larger masses. Another option is the use of non-homogeneous magnetic field [161, 180] or the use of resonant cavities in the conversion chambers [54, 101, 182]. Although it does not seem trivial to implement the buffer gas and the resonant cavity improvements for such energetic photons, the use of non-homogeneous magnetic field profiles might be a viable option at the FPF.

Acknowledgments

We thank Axel Lindner for discussions on the strategies to improve the sensitivity of LSW experiments. We also thank Javier Redondo and Igor Irastorza for useful comments regarding the endpoint of the CAST bound. We thank Andrea Caputo for clarifications regarding the extent of the Supernova bound. F.K. and P.Q. acknowledge support by the Deutsche Forschungsgemeinschaft under Germany’s Excellence Strategy - EXC 2121 Quantum Universe - 390833306. This project has received funding/support from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 860881-HIDDeN.

-
- [1] R. D. Peccei and H. R. Quinn, “CP Conservation in the Presence of Instantons,” *Phys. Rev. Lett.* **38** (1977) 1440–1443. [,328(1977)]. 1
 - [2] J. E. Kim, “Weak Interaction Singlet and Strong CP Invariance,” *Phys. Rev. Lett.* **43** (1979) 103.
 - [3] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, “Can Confinement Ensure Natural CP Invariance of Strong Interactions?,” *Nucl. Phys.* **B166** (1980) 493–506.
 - [4] A. R. Zhitnitsky, “On Possible Suppression of the Axion Hadron Interactions. (In Russian),” *Sov. J. Nucl. Phys.* **31** (1980) 260. [Yad. Fiz.31,497(1980)].
 - [5] M. Dine, W. Fischler, and M. Srednicki, “A Simple Solution to the Strong CP Problem with a Harmless Axion,” *Phys. Lett.* **B104** (1981) 199–202.
 - [6] J. E. Kim, “A Composite Invisible Axion,” *Phys. Rev.*

- D31** (1985) 1733. 1
- [7] K. R. Dienes, E. Dudas, and T. Gherghetta, “Invisible axions and large radius compactifications,” *Phys. Rev. D* **62** (2000) 105023, [arXiv:hep-ph/9912455](#). 1
- [8] G. B. Gelmini and M. Roncadelli, “Left-Handed Neutrino Mass Scale and Spontaneously Broken Lepton Number,” *Phys. Lett.* **99B** (1981) 411–415. 1
- [9] A. Davidson and K. C. Wali, “Minimal flavour unification via multigenerational Peccei-Quinn symmetry,” *Phys. Rev. Lett.* **48** (1982) 11. 1
- [10] F. Wilczek, “Axions and Family Symmetry Breaking,” *Phys. Rev. Lett.* **49** (1982) 1549–1552. 1
- [11] M. Cicoli, “Axion-like Particles from String Compactifications,” in *Proceedings, 9th Patras Workshop on Axions, WIMPs and WISPs (AXION-WIMP 2013): Mainz, Germany, June 24–28, 2013*, pp. 235–242. 2013. [arXiv:1309.6988 \[hep-th\]](#). 1
- [12] L. F. Abbott and P. Sikivie, “A Cosmological Bound on the Invisible Axion,” *Phys. Lett.* **B120** (1983) 133–136. [,URL(1982)]. 1
- [13] M. Dine and W. Fischler, “The Not So Harmless Axion,” *Phys. Lett. B* **120** (1983) 137–141.
- [14] J. Preskill, M. B. Wise, and F. Wilczek, “Cosmology of the Invisible Axion,” *Phys. Lett. B* **120** (1983) 127–132. 1
- [15] **ALPS** Collaboration, A. Spector, “ALPSII Status Report,” in *14th Patras Workshop on Axions, WIMPs and WISPs*. 6, 2019. [arXiv:1906.09011 \[physics.ins-det\]](#). 1
- [16] **IAXO** Collaboration, I. Irastorza *et al.*, “The International Axion Observatory IAXO. Letter of Intent to the CERN SPS committee,”. 1
- [17] **CAST** Collaboration, V. Anastassopoulos *et al.*, “New CAST Limit on the Axion-Photon Interaction,” *Nature Phys.* **13** (2017) 584–590, [arXiv:1705.02290 \[hep-ex\]](#). 1
- [18] E. Armengaud *et al.*, “Conceptual Design of the International Axion Observatory (IAXO),” *JINST* **9** (2014) T05002, [arXiv:1401.3233 \[physics.ins-det\]](#). 1
- [19] M. Giannotti, J. Ruz, and J. K. Vogel, “IAXO, next-generation of helioscopes,” *PoS ICHEP2016* (2016) 195, [arXiv:1611.04652 \[physics.ins-det\]](#). 1
- [20] **MADMAX interest Group** Collaboration, B. Majorovits *et al.*, “MADMAX: A new road to axion dark matter detection,” in *15th International Conference on Topics in Astroparticle and Underground Physics (TAUP 2017) Sudbury, Ontario, Canada, July 24–28, 2017*. 2017. [arXiv:1712.01062 \[physics.ins-det\]](#). 1
- [21] D. Budker, P. W. Graham, M. Ledbetter, S. Rajendran, and A. Sushkov, “Proposal for a Cosmic Axion Spin Precession Experiment (CASPER),” *Phys. Rev. X* **4** no. 2, (2014) 021030, [arXiv:1306.6089 \[hep-ph\]](#).
- [22] G. Ruoso, A. Lombardi, A. Ortolan, R. Pengo, C. Braggio, G. Carugno, C. S. Gallo, and C. C. Speake, “The QUAX proposal: a search of galactic axion with magnetic materials,” *J. Phys. Conf. Ser.* **718** no. 4, (2016) 042051, [arXiv:1511.09461 \[hep-ph\]](#).
- [23] **FUNK Experiment** Collaboration, D. Veberič *et al.*, “Search for hidden-photon dark matter with the FUNK experiment,” *PoS ICRC2017* (2018) 880, [arXiv:1711.02958 \[hep-ex\]](#).
- [24] **nEDM** Collaboration, J. Zenner, “The nEDM experiment at the Paul Scherrer Institute, Switzerland,” *AIP Conf. Proc.* **1560** no. 1, (2013) 254–256.
- [25] C. B. Adams *et al.*, “Axion Dark Matter,” [arXiv:2203.14923 \[hep-ex\]](#). 1
- [26] M. Bauer, M. Neubert, and A. Thamm, “Collider Probes of Axion-Like Particles,” *JHEP* **12** (2017) 044, [arXiv:1708.00443 \[hep-ph\]](#). 1
- [27] G. Alonso-Álvarez, M. B. Gavela, and P. Quilez, “Axion couplings to electroweak gauge bosons,” *Eur. Phys. J. C* **79** no. 3, (2019) 223, [arXiv:1811.05466 \[hep-ph\]](#). 1
- [28] E. Izaguirre, T. Lin, and B. Shuve, “Searching for Axionlike Particles in Flavor-Changing Neutral Current Processes,” *Phys. Rev. Lett.* **118** no. 11, (2017) 111802, [arXiv:1611.09355 \[hep-ph\]](#). 1
- [29] X. Cid Vidal, A. Mariotti, D. Redigolo, F. Sala, and K. Tobioka, “New Axion Searches at Flavor Factories,” *JHEP* **01** (2019) 113, [arXiv:1810.09452 \[hep-ph\]](#). [Erratum: JHEP 06, 141 (2020)].
- [30] M. B. Gavela, R. Houtz, P. Quilez, R. Del Rey, and O. Sumensari, “Flavor constraints on electroweak ALP couplings,” *Eur. Phys. J. C* **79** no. 5, (2019) 369, [arXiv:1901.02031 \[hep-ph\]](#). 6
- [31] M. Bauer, M. Neubert, S. Renner, M. Schnubel, and A. Thamm, “Flavor probes of axion-like particles,” [arXiv:2110.10698 \[hep-ph\]](#). 1
- [32] B. Döbrich, J. Jaeckel, and T. Spadaro, “Light in the beam dump - ALP production from decay photons in proton beam-dumps,” *JHEP* **05** (2019) 213, [arXiv:1904.02091 \[hep-ph\]](#). [Erratum: JHEP 10, 046 (2020)]. 1, 4, 5, 6
- [33] V. Brdar, B. Dutta, W. Jang, D. Kim, I. M. Shoemaker, Z. Tabrizi, A. Thompson, and J. Yu, “Axionlike Particles at Future Neutrino Experiments: Closing the Cosmological Triangle,” *Phys. Rev. Lett.* **126** no. 20, (2021) 201801, [arXiv:2011.07054 \[hep-ph\]](#). 1, 6
- [34] P. Coloma, P. Hernández, and S. Urrea, “New bounds on axion-like particles from MicroBooNE,” [arXiv:2202.03447 \[hep-ph\]](#). 1
- [35] G. G. Raffelt, “Astrophysical methods to constrain axions and other novel particle phenomena,” *Phys. Rept.* **198** (1990) 1–113. 1
- [36] N. Bar, K. Blum, and G. D’Amico, “Is there a supernova bound on axions?,” *Phys. Rev. D* **101** no. 12, (2020) 123025, [arXiv:1907.05020 \[hep-ph\]](#). 1
- [37] A. Caputo, G. Raffelt, and E. Vitagliano, “Muonic boson limits: Supernova redux,” *Phys. Rev. D* **105** no. 3, (2022) 035022, [arXiv:2109.03244 \[hep-ph\]](#). 1, 4
- [38] P. F. Depta, M. Hufnagel, and K. Schmidt-Hoberg, “Robust cosmological constraints on axion-like particles,” *JCAP* **05** (2020) 009, [arXiv:2002.08370 \[hep-ph\]](#). 1, 4
- [39] P. Carenza, O. Straniero, B. Döbrich, M. Giannotti, G. Lucente, and A. Mirizzi, “Constraints on the coupling with photons of heavy axion-like-particles from Globular Clusters,” *Phys. Lett. B* **809** (2020) 135709, [arXiv:2004.08399 \[hep-ph\]](#). 1
- [40] J. Jaeckel, E. Masso, J. Redondo, A. Ringwald, and

- F. Takahashi, “The Need for purely laboratory-based axion-like particle searches,” *Phys. Rev. D* **75** (2007) 013004, [arXiv:hep-ph/0610203](https://arxiv.org/abs/hep-ph/0610203). 1
- [41] J. L. Feng, I. Galon, F. Kling, and S. Trojanowski, “ForwArd Search ExpeRiment at the LHC,” *Phys. Rev. D* **97** no. 3, (2018) 035001, [arXiv:1708.09389](https://arxiv.org/abs/1708.09389) [hep-ph]. 1, 2, 3
- [42] **FASER** Collaboration, A. Ariga *et al.*, “Letter of Intent for FASER: ForwArd Search ExpeRiment at the LHC,” [arXiv:1811.10243](https://arxiv.org/abs/1811.10243) [physics.ins-det].
- [43] **FASER** Collaboration, A. Ariga *et al.*, “Technical Proposal for FASER: ForwArd Search ExpeRiment at the LHC,” [arXiv:1812.09139](https://arxiv.org/abs/1812.09139) [physics.ins-det]. 1, 3, 6
- [44] F. Kling and L. J. Nevay, “Forward neutrino fluxes at the LHC,” *Phys. Rev. D* **104** no. 11, (2021) 113008, [arXiv:2105.08270](https://arxiv.org/abs/2105.08270) [hep-ph]. 2
- [45] I. Zurbano Fernandez *et al.*, “High-Luminosity Large Hadron Collider (HL-LHC): Technical design report,” . 2
- [46] L. A. Anchordoqui *et al.*, “The Forward Physics Facility: Sites, Experiments, and Physics Potential,” [arXiv:2109.10905](https://arxiv.org/abs/2109.10905) [hep-ph]. 1, 4
- [47] J. L. Feng *et al.*, “The Forward Physics Facility at the High-Luminosity LHC,” in *2022 Snowmass Summer Study*. 3, 2022. [arXiv:2203.05090](https://arxiv.org/abs/2203.05090) [hep-ex]. 1, 4
- [48] **ATLAS** Collaboration, G. Aad *et al.*, “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC,” *Phys. Lett. B* **716** (2012) 1–29, [arXiv:1207.7214](https://arxiv.org/abs/1207.7214) [hep-ex]. 2
- [49] **CMS** Collaboration, S. Chatrchyan *et al.*, “Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC,” *Phys. Lett. B* **716** (2012) 30–61, [arXiv:1207.7235](https://arxiv.org/abs/1207.7235) [hep-ex]. 2
- [50] F. Kling and S. Trojanowski, “Forward experiment sensitivity estimator for the LHC and future hadron colliders,” *Phys. Rev. D* **104** no. 3, (2021) 035012, [arXiv:2105.07077](https://arxiv.org/abs/2105.07077) [hep-ph]. 2
- [51] T. Pierog, I. Karpenko, J. M. Katzy, E. Yatsenko, and K. Werner, “EPOS LHC: Test of collective hadronization with data measured at the CERN Large Hadron Collider,” *Phys. Rev. C* **92** no. 3, (2015) 034906, [arXiv:1306.0121](https://arxiv.org/abs/1306.0121) [hep-ph]. 2
- [52] R. Ulrich, T. Pierog, and C. Baus, “Cosmic ray monte carlo package, crmc,” Aug., 2021. <https://doi.org/10.5281/zenodo.5270381>. 2
- [53] L. J. Nevay *et al.*, “BDSIM: An accelerator tracking code with particle–matter interactions,” *Comput. Phys. Commun.* **252** (2020) 107200, [arXiv:1808.10745](https://arxiv.org/abs/1808.10745) [physics.comp-ph]. 2
- [54] J. Redondo and A. Ringwald, “Light shining through walls,” *Contemp. Phys.* **52** (2011) 211–236, [arXiv:1011.3741](https://arxiv.org/abs/1011.3741) [hep-ph]. 2, 6
- [55] H. Primakoff, “Photoproduction of neutral mesons in nuclear electric fields and the mean life of the neutral meson,” *Phys. Rev.* **81** (1951) 899. 3
- [56] Y.-S. Tsai, “AXION BREMSSTRAHLUNG BY AN ELECTRON BEAM,” *Phys. Rev. D* **34** (1986) 1326. 3
- [57] J. L. Feng, I. Galon, F. Kling, and S. Trojanowski, “Axionlike particles at FASER: The LHC as a photon beam dump,” *Phys. Rev. D* **98** no. 5, (2018) 055021, [arXiv:1806.02348](https://arxiv.org/abs/1806.02348) [hep-ph]. 3
- [58] **FASER** Collaboration, A. Ariga *et al.*, “FASER: ForwArd Search ExpeRiment at the LHC,” [arXiv:1901.04468](https://arxiv.org/abs/1901.04468) [hep-ex]. 3
- [59] **FASER** Collaboration, H. Abreu *et al.*, “The trigger and data acquisition system of the FASER experiment,” *JINST* **16** no. 12, (2021) P12028, [arXiv:2110.15186](https://arxiv.org/abs/2110.15186) [physics.ins-det].
- [60] **FASER** Collaboration, H. Abreu *et al.*, “The tracking detector of the FASER experiment,” [arXiv:2112.01116](https://arxiv.org/abs/2112.01116) [physics.ins-det]. 3
- [61] J. L. Feng, I. Galon, F. Kling, and S. Trojanowski, “Dark Higgs bosons at the ForwArd Search ExpeRiment,” *Phys. Rev. D* **97** no. 5, (2018) 055034, [arXiv:1710.09387](https://arxiv.org/abs/1710.09387) [hep-ph]. 3
- [62] F. Kling and S. Trojanowski, “Heavy Neutral Leptons at FASER,” *Phys. Rev. D* **97** no. 9, (2018) 095016, [arXiv:1801.08947](https://arxiv.org/abs/1801.08947) [hep-ph].
- [63] J. L. Feng, I. Galon, F. Kling, and S. Trojanowski, “Axionlike particles at FASER: The LHC as a photon beam dump,” *Phys. Rev. D* **98** no. 5, (2018) 055021, [arXiv:1806.02348](https://arxiv.org/abs/1806.02348) [hep-ph]. 5
- [64] **FASER** Collaboration, A. Ariga *et al.*, “FASER’s physics reach for long-lived particles,” *Phys. Rev. D* **99** no. 9, (2019) 095011, [arXiv:1811.12522](https://arxiv.org/abs/1811.12522) [hep-ph]. 3, 5
- [65] **FASER** Collaboration, H. Abreu *et al.*, “Detecting and Studying High-Energy Collider Neutrinos with FASER at the LHC,” *Eur. Phys. J. C* **80** no. 1, (2020) 61, [arXiv:1908.02310](https://arxiv.org/abs/1908.02310) [hep-ex]. 3
- [66] **FASER** Collaboration, H. Abreu *et al.*, “Technical Proposal: FASERnu,” [arXiv:2001.03073](https://arxiv.org/abs/2001.03073) [physics.ins-det].
- [67] **FASER** Collaboration, H. Abreu *et al.*, “First neutrino interaction candidates at the LHC,” *Phys. Rev. D* **104** no. 9, (2021) L091101, [arXiv:2105.06197](https://arxiv.org/abs/2105.06197) [hep-ex]. 3
- [68] J. Boyd, “The FASER W-Si High Precision Preshower Technical Proposal,” tech. rep., CERN, Geneva, Mar, 2022. <https://cds.cern.ch/record/2803084>. 3
- [69] G. Grilli di Cortona, E. Hardy, J. Pardo Vega, and G. Villadoro, “The QCD axion, precisely,” *JHEP* **01** (2016) 034, [arXiv:1511.02867](https://arxiv.org/abs/1511.02867) [hep-ph]. 5
- [70] C. O’Hare, “cajohare/axionlimits: Axionlimits,” . <https://doi.org/10.5281/zenodo.3932430>. 5
- [71] K. Jodlowski and S. Trojanowski, “Neutrino beam-dump experiment with FASER at the LHC,” *JHEP* **05** (2021) 191, [arXiv:2011.04751](https://arxiv.org/abs/2011.04751) [hep-ph]. 4
- [72] **CAST** Collaboration, S. Andriamonje *et al.*, “An Improved limit on the axion-photon coupling from the CAST experiment,” *JCAP* **04** (2007) 010, [arXiv:hep-ex/0702006](https://arxiv.org/abs/hep-ex/0702006). 4
- [73] **CAST** Collaboration, V. Anastassopoulos *et al.*, “New CAST Limit on the Axion-Photon Interaction,” *Nature Phys.* **13** (2017) 584–590, [arXiv:1705.02290](https://arxiv.org/abs/1705.02290) [hep-ex].
- [74] I. Shilon, A. Dudarev, H. Silva, U. Wagner, and H. H. J. ten Kate, “The Superconducting Toroid for the New International AXion Observatory (IAXO),” *IEEE Trans. Appl. Supercond.* **24** no. 3, (2014) 4500104, [arXiv:1309.2117](https://arxiv.org/abs/1309.2117) [physics.ins-det].
- [75] S.-F. Ge, K. Hamaguchi, K. Ichimura, K. Ishidoshiro, Y. Kanazawa, Y. Kishimoto, N. Nagata, and J. Zheng, “Supernova-scope for the Direct Search of Supernova Axions,” *JCAP* **11** (2020) 059, [arXiv:2008.03924](https://arxiv.org/abs/2008.03924) [hep-ph]. 4

- [76] J. L. Ouellet *et al.*, “First Results from ABRACADABRA-10 cm: A Search for Sub- μ eV Axion Dark Matter,” *Phys. Rev. Lett.* **122** no. 12, (2019) 121802, [arXiv:1810.12257 \[hep-ex\]](#). 4
- [77] C. P. Salemi *et al.*, “Search for Low-Mass Axion Dark Matter with ABRACADABRA-10 cm,” *Phys. Rev. Lett.* **127** no. 8, (2021) 081801, [arXiv:2102.06722 \[hep-ex\]](#).
- [78] S. J. Asztalos, G. Carosi, C. Hagmann, D. Kinion, K. van Bibber, M. Hotz, L. J. Rosenberg, G. Rybka, J. Hoskins, J. Hwang, P. Sikivie, D. B. Tanner, R. Bradley, J. Clarke, and ADMX Collaboration, “SQUID-Based Microwave Cavity Search for Dark-Matter Axions,” *Phys. Rev. Lett.* **104** no. 4, (Jan., 2010) 041301, [arXiv:0910.5914 \[astro-ph.CO\]](#).
- [79] ADMX Collaboration, N. Du *et al.*, “A Search for Invisible Axion Dark Matter with the Axion Dark Matter Experiment,” *Phys. Rev. Lett.* **120** no. 15, (2018) 151301, [arXiv:1804.05750 \[hep-ex\]](#).
- [80] ADMX Collaboration, T. Braine *et al.*, “Extended Search for the Invisible Axion with the Axion Dark Matter Experiment,” *Phys. Rev. Lett.* **124** no. 10, (2020) 101303, [arXiv:1910.08638 \[hep-ex\]](#).
- [81] ADMX Collaboration, C. Bartram *et al.*, “Search for Invisible Axion Dark Matter in the 3.3–4.2 μ eV Mass Range,” *Phys. Rev. Lett.* **127** no. 26, (2021) 261803, [arXiv:2110.06096 \[hep-ex\]](#).
- [82] ADMX Collaboration, C. Boutan *et al.*, “Piezoelectrically Tuned Multimode Cavity Search for Axion Dark Matter,” *Phys. Rev. Lett.* **121** no. 26, (2018) 261302, [arXiv:1901.00920 \[hep-ex\]](#).
- [83] C. Bartram *et al.*, “Dark Matter Axion Search Using a Josephson Traveling Wave Parametric Amplifier,” [arXiv:2110.10262 \[hep-ex\]](#).
- [84] N. Crisosto, P. Sikivie, N. S. Sullivan, D. B. Tanner, J. Yang, and G. Rybka, “ADMX SLIC: Results from a Superconducting LC Circuit Investigating Cold Axions,” *Phys. Rev. Lett.* **124** no. 24, (2020) 241101, [arXiv:1911.05772 \[astro-ph.CO\]](#).
- [85] S. Lee, S. Ahn, J. Choi, B. R. Ko, and Y. K. Semertzidis, “Axion Dark Matter Search around 6.7 μ eV,” *Phys. Rev. Lett.* **124** no. 10, (2020) 101802, [arXiv:2001.05102 \[hep-ex\]](#).
- [86] J. Jeong, S. Youn, S. Bae, J. Kim, T. Seong, J. E. Kim, and Y. K. Semertzidis, “Search for Invisible Axion Dark Matter with a Multiple-Cell Haloscope,” *Phys. Rev. Lett.* **125** no. 22, (2020) 221302, [arXiv:2008.10141 \[hep-ex\]](#).
- [87] CAPP Collaboration, O. Kwon *et al.*, “First Results from an Axion Haloscope at CAPP around 10.7 μ eV,” *Phys. Rev. Lett.* **126** no. 19, (2021) 191802, [arXiv:2012.10764 \[hep-ex\]](#).
- [88] J. A. Devlin *et al.*, “Constraints on the Coupling between Axionlike Dark Matter and Photons Using an Antiproton Superconducting Tuned Detection Circuit in a Cryogenic Penning Trap,” *Phys. Rev. Lett.* **126** no. 4, (2021) 041301, [arXiv:2101.11290 \[astro-ph.CO\]](#).
- [89] T. Grenet, R. Ballou, Q. Basto, K. Martineau, P. Perrier, P. Pugnat, J. Quevillon, N. Roch, and C. Smith, “The Grenoble Axion Haloscope platform (GrAHal): development plan and first results,” [arXiv:2110.14406 \[hep-ex\]](#).
- [90] HAYSTAC Collaboration, L. Zhong *et al.*, “Results from phase 1 of the HAYSTAC microwave cavity axion experiment,” *Phys. Rev. D* **97** no. 9, (2018) 092001, [arXiv:1803.03690 \[hep-ex\]](#).
- [91] HAYSTAC Collaboration, K. M. Backes *et al.*, “A quantum-enhanced search for dark matter axions,” *Nature* **590** no. 7845, (2021) 238–242, [arXiv:2008.01853 \[quant-ph\]](#).
- [92] B. T. McAllister, G. Flower, E. N. Ivanov, M. Goryachev, J. Bourhill, and M. E. Tobar, “The ORGAN Experiment: An axion haloscope above 15 GHz,” *Phys. Dark Univ.* **18** (2017) 67–72, [arXiv:1706.00209 \[physics.ins-det\]](#).
- [93] D. Alesini *et al.*, “Galactic axions search with a superconducting resonant cavity,” *Phys. Rev. D* **99** no. 10, (2019) 101101, [arXiv:1903.06547 \[physics.ins-det\]](#).
- [94] D. Alesini *et al.*, “Search for invisible axion dark matter of mass $m_a = 43 \mu$ eV with the QUAX- $a\gamma$ experiment,” *Phys. Rev. D* **103** no. 10, (2021) 102004, [arXiv:2012.09498 \[hep-ex\]](#).
- [95] CAST Collaboration, A. A. Melcón *et al.*, “First results of the CAST-RADES haloscope search for axions at 34.67 μ eV,” *JHEP* **21** (2020) 075, [arXiv:2104.13798 \[hep-ex\]](#).
- [96] S. DePanfilis, A. C. Melissinos, B. E. Moskowitz, J. T. Rogers, Y. K. Semertzidis, W. U. Wuensch, H. J. Halama, A. G. Prodell, W. B. Fowler, and F. A. Nezrick, “Limits on the abundance and coupling of cosmic axions at $4.5 < m_a < 5.0 \mu$ eV,” *Phys. Rev. Lett.* **59** (Aug, 1987) 839–842. <https://link.aps.org/doi/10.1103/PhysRevLett.59.839>.
- [97] A. V. Gramolin, D. Aybas, D. Johnson, J. Adam, and A. O. Sushkov, “Search for axion-like dark matter with ferromagnets,” *Nature Phys.* **17** no. 1, (2021) 79–84, [arXiv:2003.03348 \[hep-ex\]](#).
- [98] A. Arza, M. A. Fedderke, P. W. Graham, D. F. Jackson Kimball, and S. Kalia, “Earth as a transducer for axion dark-matter detection,” [arXiv:2112.09620 \[hep-ph\]](#).
- [99] C. Hagmann, P. Sikivie, N. S. Sullivan, and D. B. Tanner, “Results from a search for cosmic axions,” *Phys. Rev. D* **42** (Aug, 1990) 1297–1300. <https://link.aps.org/doi/10.1103/PhysRevD.42.1297>.
- [100] C. A. Thomson, B. T. McAllister, M. Goryachev, E. N. Ivanov, and M. E. Tobar, “Upconversion Loop Oscillator Axion Detection Experiment: A Precision Frequency Interferometric Axion Dark Matter Search with a Cylindrical Microwave Cavity,” *Phys. Rev. Lett.* **126** no. 8, (2021) 081803, [arXiv:1912.07751 \[hep-ex\]](#). [Erratum: Phys.Rev.Lett. 127, 019901 (2021)]. 4
- [101] K. Ehret *et al.*, “New ALPS Results on Hidden-Sector Lightweights,” *Phys. Lett. B* **689** (2010) 149–155, [arXiv:1004.1313 \[hep-ex\]](#). 4, 5, 6
- [102] M. Betz, F. Caspers, M. Gasior, M. Thumm, and S. W. Rieger, “First results of the CERN Resonant Weakly Interacting sub-eV Particle Search (CROWS),” *Phys. Rev. D* **88** no. 7, (2013) 075014, [arXiv:1310.8098 \[physics.ins-det\]](#). 5
- [103] OSQAR Collaboration, R. Ballou *et al.*, “New exclusion limits on scalar and pseudoscalar axionlike particles from light shining through a wall,” *Phys. Rev. D* **92** no. 9, (2015) 092002, [arXiv:1506.08082](#).

- [hep-ex]. 5
- [104] F. Della Valle, A. Ejlli, U. Gastaldi, G. Messineo, E. Milotti, R. Pengo, G. Ruoso, and G. Zavattini, “The PVLAS experiment: measuring vacuum magnetic birefringence and dichroism with a birefringent Fabry–Perot cavity,” *Eur. Phys. J. C* **76** no. 1, (2016) 24, [arXiv:1510.08052 \[physics.optics\]](#). 5
- [105] **SAPPHIRE** Collaboration, K. Homma *et al.*, “Search for sub-eV axion-like resonance states via stimulated quasi-parallel laser collisions with the parameterization including fully asymmetric collisional geometry,” *JHEP* **12** (2021) 108, [arXiv:2105.01224 \[hep-ex\]](#).
- [106] M. D. Ortiz *et al.*, “Design of the ALPS II optical system,” *Phys. Dark Univ.* **35** (2022) 100968, [arXiv:2009.14294 \[physics.optics\]](#). 4, 5
- [107] G. Lucente, P. Carenza, T. Fischer, M. Giannotti, and A. Mirizzi, “Heavy axion-like particles and core-collapse supernovae: constraints and impact on the explosion mechanism,” *JCAP* **12** (2020) 008, [arXiv:2008.04918 \[hep-ph\]](#). 4
- [108] A. Caputo, H.-T. Janka, G. Raffelt, and E. Vitagliano, “Low-Energy Supernovae Severely Constrain Radiative Particle Decays,” [arXiv:2201.09890 \[astro-ph.HE\]](#). 4
- [109] M. Xiao, K. M. Perez, M. Giannotti, O. Straniero, A. Mirizzi, B. W. Grefenstette, B. M. Roach, and M. Nynka, “Constraints on Axionlike Particles from a Hard X-Ray Observation of Betelgeuse,” *Phys. Rev. Lett.* **126** no. 3, (2021) 031101, [arXiv:2009.09059 \[astro-ph.HE\]](#). 4
- [110] A. Keller, S. O’Brien, A. Kamdar, N. M. Rapidis, A. F. Leder, and K. van Bibber, “A Model-independent Radio Telescope Dark Matter Search,” *Astrophys. J.* **927** no. 1, (2022) 71, [arXiv:2112.03439 \[astro-ph.CO\]](#).
- [111] J. W. Foster, S. J. Witte, M. Lawson, T. Linden, V. Gajjar, C. Weniger, and B. R. Safdi, “Extraterrestrial Axion Search with the Breakthrough Listen Galactic Center Survey,” [arXiv:2202.08274 \[astro-ph.CO\]](#).
- [112] M. H. Chan, “Constraining the axion–photon coupling using radio data of the Bullet cluster,” *Sci. Rep.* **11** no. 1, (2021) 20087, [arXiv:2109.11734 \[astro-ph.CO\]](#).
- [113] D. Wouters and P. Brun, “Constraints on Axion-like Particles from X-Ray Observations of the Hydra Galaxy Cluster,” *Astrophys. J.* **772** (2013) 44, [arXiv:1304.0989 \[astro-ph.HE\]](#).
- [114] M. C. D. Marsh, H. R. Russell, A. C. Fabian, B. P. McNamara, P. Nulsen, and C. S. Reynolds, “A New Bound on Axion-Like Particles,” *JCAP* **12** (2017) 036, [arXiv:1703.07354 \[hep-ph\]](#).
- [115] C. S. Reynolds, M. C. D. Marsh, H. R. Russell, A. C. Fabian, R. Smith, F. Tombesi, and S. Veilleux, “Astrophysical limits on very light axion-like particles from Chandra grating spectroscopy of NGC 1275,” [arXiv:1907.05475 \[hep-ph\]](#).
- [116] J. S. Reynés, J. H. Matthews, C. S. Reynolds, H. R. Russell, R. N. Smith, and M. C. D. Marsh, “New constraints on light axion-like particles using Chandra transmission grating spectroscopy of the powerful cluster-hosted quasar H1821+643,” *Mon. Not. Roy. Astron. Soc.* **510** no. 1, (2021) 1264–1277, [arXiv:2109.03261 \[astro-ph.HE\]](#).
- [117] B. Bolliet, J. Chluba, and R. Battye, “Spectral distortion constraints on photon injection from low-mass decaying particles,” *Mon. Not. Roy. Astron. Soc.* **507** no. 3, (2021) 3148–3178, [arXiv:2012.07292 \[astro-ph.CO\]](#).
- [118] F. Calore, P. Carenza, C. Eckner, T. Fischer, M. Giannotti, J. Jaeckel, K. Kotake, T. Kuroda, A. Mirizzi, and F. Sivo, “3D template-based Fermi-LAT constraints on the diffuse supernova axion-like particle background,” *Phys. Rev. D* **105** no. 6, (2022) 063028, [arXiv:2110.03679 \[astro-ph.HE\]](#).
- [119] F. Calore, P. Carenza, M. Giannotti, J. Jaeckel, and A. Mirizzi, “Bounds on axionlike particles from the diffuse supernova flux,” *Phys. Rev. D* **102** no. 12, (2020) 123005, [arXiv:2008.11741 \[hep-ph\]](#).
- [120] M. A. Buen-Abad, J. Fan, and C. Sun, “Constraints on axions from cosmic distance measurements,” *JHEP* **02** (2022) 103, [arXiv:2011.05993 \[hep-ph\]](#).
- [121] **Fermi-LAT** Collaboration, M. Ajello *et al.*, “Search for Spectral Irregularities due to Photon–Axionlike-Particle Oscillations with the Fermi Large Area Telescope,” *Phys. Rev. Lett.* **116** no. 16, (2016) 161101, [arXiv:1603.06978 \[astro-ph.HE\]](#).
- [122] M. Meyer and T. Petrushevska, “Search for Axionlike-Particle-Induced Prompt γ -Ray Emission from Extragalactic Core-Collapse Supernovae with the *Fermi* Large Area Telescope,” *Phys. Rev. Lett.* **124** no. 23, (2020) 231101, [arXiv:2006.06722 \[astro-ph.HE\]](#). [Erratum: *Phys. Rev. Lett.* 125, 119901 (2020)].
- [123] S. Jacobsen, T. Linden, and K. Freese, “Constraining Axion-Like Particles with HAWC Observations of TeV Blazars,” [arXiv:2203.04332 \[hep-ph\]](#).
- [124] **H.E.S.S.** Collaboration, A. Abramowski *et al.*, “Constraints on axionlike particles with H.E.S.S. from the irregularity of the PKS 2155-304 energy spectrum,” *Phys. Rev. D* **88** no. 10, (2013) 102003, [arXiv:1311.3148 \[astro-ph.HE\]](#).
- [125] A. Ayala, I. Domínguez, M. Giannotti, A. Mirizzi, and O. Straniero, “Revisiting the bound on axion-photon coupling from Globular Clusters,” *Phys. Rev. Lett.* **113** no. 19, (2014) 191302, [arXiv:1406.6053 \[astro-ph.SR\]](#).
- [126] D. Wadekar and Z. Wang, “Strong constraints on decay and annihilation of dark matter from heating of gas-rich dwarf galaxies,” [arXiv:2111.08025 \[hep-ph\]](#).
- [127] C. Dessert, D. Dunsky, and B. R. Safdi, “Upper limit on the axion-photon coupling from magnetic white dwarf polarization,” [arXiv:2203.04319 \[hep-ph\]](#).
- [128] H.-J. Li, J.-G. Guo, X.-J. Bi, S.-J. Lin, and P.-F. Yin, “Limits on axion-like particles from Mrk 421 with 4.5-year period observations by ARGO-YBJ and Fermi-LAT,” *Phys. Rev. D* **103** no. 8, (2021) 083003, [arXiv:2008.09464 \[astro-ph.HE\]](#).
- [129] J. W. Foster, Y. Kahn, O. Macias, Z. Sun, R. P. Eatough, V. I. Kondratiev, W. M. Peters, C. Weniger, and B. R. Safdi, “Green Bank and Effelsberg Radio Telescope Searches for Axion Dark Matter Conversion in Neutron Star Magnetospheres,” *Phys. Rev. Lett.* **125** no. 17, (2020) 171301, [arXiv:2004.00011 \[astro-ph.CO\]](#).

- [130] J. Darling, “New Limits on Axionic Dark Matter from the Magnetar PSR J1745-2900,” *Astrophys. J. Lett.* **900** no. 2, (2020) L28, [arXiv:2008.11188 \[astro-ph.CO\]](#).
- [131] R. A. Battye, J. Darling, J. I. McDonald, and S. Srinivasan, “Towards robust constraints on axion dark matter using PSR J1745-2900,” *Phys. Rev. D* **105** no. 2, (2022) L021305, [arXiv:2107.01225 \[astro-ph.CO\]](#).
- [132] N. Vinyoles, A. Serenelli, F. L. Villante, S. Basu, J. Redondo, and J. Isern, “New axion and hidden photon constraints from a solar data global fit,” *JCAP* **2015** no. 10, (Oct., 2015) 015, [arXiv:1501.01639 \[astro-ph.SR\]](#).
- [133] J. Jaeckel, P. C. Malta, and J. Redondo, “Decay photons from the axionlike particles burst of type II supernovae,” *Phys. Rev. D* **98** no. 5, (2018) 055032, [arXiv:1702.02964 \[hep-ph\]](#).
- [134] A. Payez, C. Evoli, T. Fischer, M. Giannotti, A. Mirizzi, and A. Ringwald, “Revisiting the SN1987A gamma-ray limit on ultralight axion-like particles,” *JCAP* **02** (2015) 006, [arXiv:1410.3747 \[astro-ph.HE\]](#).
- [135] C. Dessert, J. W. Foster, and B. R. Safdi, “X-ray Searches for Axions from Super Star Clusters,” *Phys. Rev. Lett.* **125** no. 26, (2020) 261102, [arXiv:2008.03305 \[hep-ph\]](#).
- [136] B. D. Blout, E. J. Daw, M. P. Decowski, P. T. P. Ho, L. J. Rosenberg, and D. B. Yu, “A Radio telescope search for axions,” *Astrophys. J.* **546** (2001) 825–828, [arXiv:astro-ph/0006310](#).
- [137] M. Regis, M. Taoso, D. Vaz, J. Brinchmann, S. L. Zoutendijk, N. F. Bouché, and M. Steinmetz, “Searching for light in the darkness: Bounds on ALP dark matter with the optical MUSE-faint survey,” *Phys. Lett. B* **814** (2021) 136075, [arXiv:2009.01310 \[astro-ph.CO\]](#).
- [138] D. Grin, G. Covone, J.-P. Kneib, M. Kamionkowski, A. Blain, and E. Jullo, “A Telescope Search for Decaying Relic Axions,” *Phys. Rev. D* **75** (2007) 105018, [arXiv:astro-ph/0611502](#).
- [139] M. Meyer, M. Giannotti, A. Mirizzi, J. Conrad, and M. A. Sánchez-Conde, “Fermi Large Area Telescope as a Galactic Supernovae Axionscope,” *Phys. Rev. Lett.* **118** no. 1, (2017) 011103, [arXiv:1609.02350 \[astro-ph.HE\]](#).
- [140] C. Thorpe-Morgan, D. Malyshev, A. Santangelo, J. Jochum, B. Jäger, M. Sasaki, and S. Saeedi, “THESEUS insights into axionlike particles, dark photon, and sterile neutrino dark matter,” *Phys. Rev. D* **102** no. 12, (2020) 123003, [arXiv:2008.08306 \[astro-ph.HE\]](#).
- [141] A. Dekker, E. Peerbooms, F. Zimmer, K. C. Y. Ng, and S. Ando, “Searches for sterile neutrinos and axionlike particles from the Galactic halo with eROSITA,” *Phys. Rev. D* **104** no. 2, (2021) 023021, [arXiv:2103.13241 \[astro-ph.HE\]](#).
- [142] M. J. Dolan, F. J. Hiskens, and R. R. Volkas, “Constraining axion-like particles using the white dwarf initial-final mass relation,” *JCAP* **09** (2021) 010, [arXiv:2102.00379 \[hep-ph\]](#).
- [143] J. W. Foster, M. Kongsoore, C. Dessert, Y. Park, N. L. Rodd, K. Cranmer, and B. R. Safdi, “Deep Search for Decaying Dark Matter with XMM-Newton Blank-Sky Observations,” *Phys. Rev. Lett.* **127** no. 5, (2021) 051101, [arXiv:2102.02207 \[astro-ph.CO\]](#).
- [144] D. Cadamuro and J. Redondo, “Cosmological bounds on pseudo Nambu-Goldstone bosons,” *JCAP* **02** (2012) 032, [arXiv:1110.2895 \[hep-ph\]](#).
- [145] NOMAD Collaboration, P. Astier *et al.*, “Search for eV (pseudo)scalar penetrating particles in the SPS neutrino beam,” *Phys. Lett. B* **479** (2000) 371–380.
- [146] J. Jaeckel and M. Spannowsky, “Probing MeV to 90 GeV axion-like particles with LEP and LHC,” *Phys. Lett. B* **753** (2016) 482–487, [arXiv:1509.00476 \[hep-ph\]](#).
- [147] DELPHI Collaboration, P. Abreu *et al.*, “The reaction $e+e^- \rightarrow \gamma\gamma\gamma\gamma$ (gamma) at Z_0 energies,” *Phys. Lett. B* **268** (1991) 296–304.
- [148] DELPHI Collaboration, P. Abreu *et al.*, “Measurement of the $e+e^- \rightarrow \gamma\gamma\gamma\gamma$ (gamma) cross-section at LEP energies,” *Phys. Lett. B* **327** (1994) 386–396.
- [149] L3 Collaboration, M. Acciarri *et al.*, “Search for anomalous $Z \rightarrow \gamma\gamma\gamma\gamma$ events at LEP,” *Phys. Lett. B* **345** (1995) 609–616.
- [150] L3 Collaboration, M. Acciarri *et al.*, “Tests of QED at LEP energies using $e+e^- \rightarrow \gamma\gamma\gamma\gamma$ and $e+e^- \rightarrow \text{lepton}^+\text{lepton}^- \gamma\gamma\gamma\gamma$,” *Phys. Lett. B* **353** (1995) 136–144.
- [151] CHARM Collaboration, F. Bergsma *et al.*, “Search for Axion Like Particle Production in 400-GeV Proton - Copper Interactions,” *Phys. Lett. B* **157** (1985) 458–462.
- [152] E. M. Riordan *et al.*, “A Search for Short Lived Axions in an Electron Beam Dump Experiment,” *Phys. Rev. Lett.* **59** (1987) 755.
- [153] B. Döbrich, “Axion-like Particles from Primakov production in beam-dumps,” *CERN Proc. 1* (2018) 253, [arXiv:1708.05776 \[hep-ph\]](#).
- [154] J. D. Bjorken, S. Ecklund, W. R. Nelson, A. Abashian, C. Church, B. Lu, L. W. Mo, T. A. Nunamaker, and P. Rassmann, “Search for Neutral Metastable Penetrating Particles Produced in the SLAC Beam Dump,” *Phys. Rev. D* **38** (1988) 3375.
- [155] M. J. Dolan, T. Ferber, C. Hearty, F. Kahlhoefer, and K. Schmidt-Hoberg, “Revised constraints and Belle II sensitivity for visible and invisible axion-like particles,” *JHEP* **12** (2017) 094, [arXiv:1709.00009 \[hep-ph\]](#). [Erratum: JHEP 03, 190 (2021)].
- [156] J. Blumlein *et al.*, “Limits on neutral light scalar and pseudoscalar particles in a proton beam dump experiment,” *Z. Phys. C* **51** (1991) 341–350.
- [157] NA64 Collaboration, D. Banerjee *et al.*, “Search for Axionlike and Scalar Particles with the NA64 Experiment,” *Phys. Rev. Lett.* **125** no. 8, (2020) 081801, [arXiv:2005.02710 \[hep-ex\]](#).
- [158] Belle-II Collaboration, F. Abudinén *et al.*, “Search for Axion-Like Particles produced in e^+e^- collisions at Belle II,” *Phys. Rev. Lett.* **125** no. 16, (2020) 161806, [arXiv:2007.13071 \[hep-ex\]](#).
- [159] D. Aloni, C. Fanelli, Y. Soreq, and M. Williams, “Photoproduction of Axionlike Particles,” *Phys. Rev. Lett.* **123** no. 7, (2019) 071801, [arXiv:1903.03586 \[hep-ph\]](#).
- [160] J. Blumlein *et al.*, “Limits on the mass of light (pseudo)scalar particles from Bethe-Heitler $e+e^-$ and

- mu+ mu- pair production in a proton - iron beam dump experiment," *Int. J. Mod. Phys. A* **7** (1992) 3835–3850. 5
- [161] H. Fukuda, H. Otono, and S. Shirai, "QCD Axion Search with ILC Beam Facility," [arXiv:2203.06137 \[hep-ph\]](#). 5, 6
- [162] T. Behnke, J. E. Brau, B. Foster, J. Fuster, M. Harrison, J. M. Paterson, M. Peskin, M. Stanitzki, N. Walker, and H. Yamamoto, "The International Linear Collider Technical Design Report - Volume 1: Executive Summary," [arXiv:1306.6327 \[physics.acc-ph\]](#). 5
- [163] W. M. Bonivento, D. Kim, and K. Sinha, "PASSAT: Particle Accelerator helioScopes for Slim Axion-like-particle deTection," *Eur. Phys. J. C* **80** no. 2, (2020) 164, [arXiv:1909.03071 \[hep-ph\]](#). 6
- [164] P. S. B. Dev, D. Kim, K. Sinha, and Y. Zhang, "PASSAT at future neutrino experiments: Hybrid beam-dump-helioscope facilities to probe light axionlike particles," *Phys. Rev. D* **104** no. 3, (2021) 035037, [arXiv:2101.08781 \[hep-ph\]](#). 6
- [165] J. B. Dent, B. Dutta, D. Kim, S. Liao, R. Mahapatra, K. Sinha, and A. Thompson, "New Directions for Axion Searches via Scattering at Reactor Neutrino Experiments," *Phys. Rev. Lett.* **124** no. 21, (2020) 211804, [arXiv:1912.05733 \[hep-ph\]](#). 6
- [166] J. Beacham *et al.*, "Physics Beyond Colliders at CERN: Beyond the Standard Model Working Group Report," *J. Phys. G* **47** no. 1, (2020) 010501, [arXiv:1901.09966 \[hep-ex\]](#). 6
- [167] H. Georgi, D. B. Kaplan, and L. Randall, "Manifesting the Invisible Axion at Low-energies," *Phys. Lett.* **169B** (1986) 73–78. 6
- [168] L. Di Luzio, F. Mescia, and E. Nardi, "Redefining the Axion Window," *Phys. Rev. Lett.* **118** no. 3, (2017) 031801, [arXiv:1610.07593 \[hep-ph\]](#).
- [169] L. Di Luzio, F. Mescia, and E. Nardi, "Window for preferred axion models," *Phys. Rev. D* **96** no. 7, (2017) 075003, [1705.05370 \[hep-ph\]](#). 6
- [170] M. Farina, D. Pappadopulo, F. Rompineve, and A. Tesi, "The photo-philic QCD axion," *JHEP* **01** (2017) 095, [arXiv:1611.09855 \[hep-ph\]](#). 6
- [171] L. Darmé, L. Di Luzio, M. Giannotti, and E. Nardi, "Selective enhancement of the QCD axion couplings," [arXiv:2010.15846 \[hep-ph\]](#).
- [172] P. Agrawal *et al.*, "Feebly-interacting particles: FIPs 2020 workshop report," *Eur. Phys. J. C* **81** no. 11, (2021) 1015, [arXiv:2102.12143 \[hep-ph\]](#). 6
- [173] A. Hook, "Solving the Hierarchy Problem Discretely," *Phys. Rev. Lett.* **120** no. 26, (2018) 261802, [arXiv:1802.10093 \[hep-ph\]](#). 6
- [174] L. Di Luzio, B. Gavela, P. Quilez, and A. Ringwald, "An even lighter QCD axion," *JHEP* **05** (2021) 184, [arXiv:2102.00012 \[hep-ph\]](#).
- [175] L. Di Luzio, B. Gavela, P. Quilez, and A. Ringwald, "Dark matter from an even lighter QCD axion: trapped misalignment," *JCAP* **10** (2021) 001, [arXiv:2102.01082 \[hep-ph\]](#). 6
- [176] T. Dafni, C. A. J. O'Hare, B. Lakić, J. Galán, F. J. Iguañ, I. G. Irastorza, K. Jakovčić, G. Luzón, J. Redondo, and E. Ruiz Chóliz, "Weighing the solar axion," *Phys. Rev. D* **99** no. 3, (2019) 035037, [arXiv:1811.09290 \[hep-ph\]](#). 6
- [177] J. Jaeckel and L. J. Thormaehlen, "Distinguishing Axion Models with IAXO," *JCAP* **03** (2019) 039, [arXiv:1811.09278 \[hep-ph\]](#). 6
- [178] S. Gori, G. Perez, and K. Tobioka, "KOTO vs. NA62 Dark Scalar Searches," *JHEP* **08** (2020) 110, [arXiv:2005.05170 \[hep-ph\]](#). 6
- [179] F. Kling and S. Trojanowski, "Looking forward to test the KOTO anomaly with FASER," *Phys. Rev. D* **102** no. 1, (2020) 015032, [arXiv:2006.10630 \[hep-ph\]](#). 6
- [180] P. Arias, J. Jaeckel, J. Redondo, and A. Ringwald, "Optimizing Light-Shining-through-a-Wall Experiments for Axion and other WISP Searches," *Phys. Rev. D* **82** (2010) 115018, [arXiv:1009.4875 \[hep-ph\]](#). 6
- [181] CAST Collaboration, M. Arik *et al.*, "Search for Solar Axions by the CERN Axion Solar Telescope with ${}^3\text{He}$ Buffer Gas: Closing the Hot Dark Matter Gap," *Phys. Rev. Lett.* **112** no. 9, (2014) 091302, [arXiv:1307.1985 \[hep-ex\]](#). 6
- [182] ALPS Collaboration, K. Ehret *et al.*, "Resonant laser power build-up in ALPS: A 'Light-shining-through-walls' experiment," *Nucl. Instrum. Meth. A* **612** (2009) 83–96, [arXiv:0905.4159 \[physics.ins-det\]](#). 6