Variance-Aware Estimation of Kernel Mean Embedding

Geoffrey Wolfer †1 and Pierre Alquier ‡1

¹RIKEN AIP

Abstract

An important feature of kernel mean embeddings (KME) is that the rate of convergence of the empirical KME to the true distribution KME can be bounded independently of the dimension of the space, properties of the distribution and smoothness features of the kernel. We show how to speed-up convergence by leveraging variance information in the RKHS. Furthermore, we show that even when such information is a priori unknown, we can efficiently estimate it from the data, recovering the desiderata of a distribution agnostic bound that enjoys acceleration in fortuitous settings. We illustrate our methods in the context of hypothesis testing and robust parametric estimation.

Keywords— kernel mean embedding, maximum mean discrepancy, distribution learning, empirical Bernstein

Contents

[†] email: geoffrey.wolfer@riken.jp.

The author is supported by the Special Postdoctoral Researcher Program of RIKEN. ‡ email: pierrealain.alquier@riken.jp.

1 Introduction

Estimating a probability distribution $\mathbb P$ over a space $\mathcal X$ from an independent sample $X_1,\ldots,X_n\sim \mathbb P$ is a central problem in computer science and statistics [\[Kearns et al.,](#page-23-2) [1994,](#page-23-2) [Tsybakov,](#page-23-3) [2008\]](#page-23-3). To formalize the question, one selects a distance (or at least a contrast function) between distributions and oftentimes introduces assumptions on the underlying probability space (e.g. finitely supported, probability function is uniformly continuous with respect to the Lebesgue measure, Hölder continuous, \dots). Increasing the stringency of assumptions generally leads to substantially faster rates. For instance, in the finite support $|\mathcal{X}| < \infty$ case and with respect to total variation, whilst the expectation risk evolves roughly as $\sqrt{|\mathcal{X}|/n}$, it is known that this rate can be accelerated by replacing $|\mathcal{X}|$ with a bound on the "half-norm" $\|\mathbb{P}\|_{1/2}$ [\[Berend and Kontorovich,](#page-22-0) [2013\]](#page-22-0), which corresponds to some measure of entropy of the underlying distribution. Furthermore, even without prior knowledge of $\|\mathbb{P}\|_{1/2}$, one can effectively construct confidence intervals around $\mathbb P$ that depend on $\|\widehat{P}_n\|_{1/2}$, i.e. the half-norm of the empirical distribution [\[Cohen et al.,](#page-22-1) [2020,](#page-22-1) Theorem 2.1]. Namely, when **P** is supported on **N**, with probability at least 1 − *δ*, it holds that

$$
\left\|\widehat{\mathbf{P}}_n - \mathbf{P}\right\|_{\mathsf{TV}} \le \frac{1}{\sqrt{n}} \left(\left\|\widehat{\mathbf{P}}\right\|_{1/2}^{1/2} + 3\sqrt{\frac{1}{2}\log\frac{2}{\delta}} \right). \tag{1}
$$

The advantage of the above expression is twofold (*a*) we do not need make assumptions on $\|\mathbb{P}\|_{1/2}$, which for all we know could be infinite, and (*b*) in favorable cases where $\|\hat{P}\|_{1/2}$ is small, the intervals will be narrower. In this paper, we set out to explore the question of whether analogues of [\(1\)](#page-1-1) are possible for general probability spaces and with respect to maximum mean discrepancy.

Outline & summary of findings. In Section [2,](#page-3-0) for the problem of estimating a distribution with respect to maximum mean discrepancy, we give a convergence rate with a dominant term in $\mathcal{O}_P(n^{-1/2})$ involving a variance term *v* which depends on both the chosen kernel *k* and the underlying distribution **P**. We then illustrate how the rate can subdue minimax lower bounds when v is favorable. In particular, we show that for large collections of kernels, and when $\mathcal{X} \subset \mathbb{R}^d$, the variance v can be controlled by a quantity that decouples the influence of the kernel and a measure of total variance of **P**. In Section [3,](#page-6-0) we proceed to show that even if *v* is unknown, it is possible to efficiently estimate it from the data with an "empirical Bernstein" approach whenever the kernel is translation invariant, or at least enjoys a mildly varying diagonal. In Section [4.2,](#page-10-0) we put our methods into practice, first in the context of hypothesis testing, and second by improving the results of [Briol et al.](#page-22-2) [\[2019\]](#page-22-2) and Chérief-Abdellatif and Alquier [\[2022\]](#page-22-3) in the context of robust parametric MMD estimation. All proofs are deferred to Section [5](#page-15-0) for clarity of the exposition.

Notation & background. Let $\mathcal X$ be a separable topological space, and $\mathcal P(\mathcal X)$ the set of all Borel probability measures over X. For a bounded function $f: \overline{X} \to \mathbb{R}$, we define for convenience

$$
\overline{f} \doteq \sup_{x \in \mathcal{X}} f(x) \qquad \underline{f} \doteq \inf_{x \in \mathcal{X}} f(x), \qquad \Delta f \doteq \overline{f} - \underline{f}. \tag{2}
$$

Let $k: \mathcal{X} \times \mathcal{X} \to \mathbb{R}$ be a continuous positive definite kernel and \mathcal{H}_k be the associated reproducing kernel Hilbert space (RKHS). We assume that the kernel is bounded^{[1](#page-2-0)} in the sense that $\sup_{x\in\mathcal{X}} k(x,x) < \infty$. Letting **P** ∈ $\mathcal{P}(\overline{\mathcal{X}})$, the kernel mean embedding (KME) of **P** is defined as

$$
\mu_{\mathbb{P}} \doteq \mathbb{E}_{X \sim \mathbb{P}} \left[k(X, \cdot) \right] = \int_{\mathcal{X}} k(x, \cdot) d\mathbb{P}(x) \in \mathcal{H}_k,
$$

which is interpreted as a Bochner integral [\[Diestel and Uhl,](#page-22-4) [1977,](#page-22-4) Chapter 2]. Let $n \in \mathbb{N}$ and X_1, \ldots, X_n be a sequence of observations sampled independently from **P**. We write

$$
\widehat{\mu}_{\mathbb{P}}(X_1,\ldots,X_n)=\widehat{\mu}_{\mathbb{P}}\doteq \frac{1}{n}\sum_{t=1}^n k(X_t,\cdot)\in \mathcal{H}_k,
$$

for the KME of the empirical distribution $\hat{\mathbb{P}}_n = \frac{1}{n} \sum_{t=1}^n \delta_{X_t}$, and call the distance $\|\hat{\mu}_P - \mu_P\|_{\mathcal{H}_t}$ the maximum mean discrepancy (MMD) between the true mean embodding and its empirical estimator. mum mean discrepancy (MMD) between the true mean embedding and its empirical estimator. A kernel is called characteristic if the map $\mu\colon \mathbb{P}\to \mu_{\mathbb{P}}$ is injective. This property ensures that $\|\mu_{\mathbb{P}}-\mu_{\mathbb{P}'}\|_{\mathcal{H}_k}=0$ if and only if $\mathbb{P} = \mathbb{P}'$. A kernel is called translation invariant (TI) ^{[2](#page-2-1)} when there exists a positive definite function $\psi \colon \mathcal{X} \to \mathbb{R}$ such for any $x, x' \in \mathcal{X}$,

$$
k(x, x') = \psi(x' - x). \tag{3}
$$

In particular, $\overline{\psi}=\psi(0)=\bar{k}$. When $\mathcal{X}=\mathbb{R}^d$, a kernel k is said to be a radial basis function (RBF) when for any $x, x' \in \mathcal{X}$, $k(x, x') = \phi(\|x - x'\|_2)$ for some function $\phi \colon \mathbb{R}_+ \to \mathbb{R}$. Noticeably, *k* being positive definite does not preclude it from taking negative values. However, when *k* is RBF, the following lower bound on *φ* holds (see e.g. [Genton](#page-22-5) [\[2001\]](#page-22-5), [Stein](#page-23-4) [\[1999\]](#page-23-4)),

$$
\underline{\phi} \geq \overline{\phi} \inf_{t \geq 0} \left\{ \left(\frac{2}{t} \right)^{(d-2)/2} \Gamma(d/2) J_{(d-2)/2}(t) \right\},\,
$$

where Γ is the Gamma function and *J^β* is the Bessel function of the first kind of order *β*, showing that |*φ*| becomes evanescent as the dimension increases.

Related work. From an asymptotic standpoint, the weak law of large numbers asserts that $\hat{\mu}_P$ is guaranteed to converge to the true $\mu_{\mathbb{P}}$. Furthermore,

$$
\sqrt{n}\left(\widehat{\mu}_{\mathbb{P}}-\mu_{\mathbb{P}}\right)
$$

converges in distribution to a zero mean Gaussian process on \mathcal{H}_k [\[Berlinet and Thomas-Agnan,](#page-22-6) [2011,](#page-22-6) Section 9.1]. This work, however, is more concerned with the finite sample theory, and more specifically in the rate of convergence of $\hat{\mu}_P$ towards μ_P with respect to the MMD distance. Conveniently and perhaps surprisingly, it is possible to derive a rate that depends neither on the smoothness of the considered kernel *k*, nor the properties of the true distribution. To obtain a distribution independent rate at O*P*(*n* [−]1/2), a typical strategy (see e.g. [Lopez-Paz et al.](#page-23-5) [\[2015,](#page-23-5) Section B.1]) consists in first expressing the dual relationship between the norm in the RKHS and the uniform norm of an empirical process,

$$
\|\widehat{\mu}_{\mathbb{P}} - \mu_{\mathbb{P}}\|_{\mathcal{H}_k} = \sup_{\substack{f \in \mathcal{H}_k \\ \|f\|_{\mathcal{H}_k} \le 1}} \langle f, \widehat{\mu}_{\mathbb{P}} - \mu_{\mathbb{P}} \rangle_{\mathcal{H}_k} = \sup_{\substack{f \in \mathcal{H}_k \\ \|f\|_{\mathcal{H}_k} \le 1}} \left(\frac{1}{n} \sum_{t=1}^n f(X_t) - \mathbb{E}f\right). \tag{4}
$$

¹Note that the supremum of the kernel is always reached on the diagonal, that is, $\sup_{(x,y)\in\mathcal{X}^2} k(x,y)$ = sup*x*∈X *k*(*x*, *x*). ²Also sometimes referred to as an *anisotropic stationary* kernel.

A classical symmetrization argument [\[Mohri et al.,](#page-23-6) [2018\]](#page-23-6) followed by an application of McDiarmid's in-equality [\[McDiarmid et al.,](#page-23-7) [1989\]](#page-23-7) yield that with probability at least $1 - \delta$,

$$
\sup_{\substack{f \in \mathcal{H}_k \\ ||f||_{\mathcal{H}_k} \le 1}} \left(\frac{1}{n} \sum_{t=1}^n f(X_t) - \mathbb{E}f \right) \le 2\Re_n + \sqrt{\frac{2\bar{k} \log(1/\delta)}{n}}, \tag{5}
$$

where \mathfrak{R}_n is the Rademacher complexity [\[Mohri et al.,](#page-23-6) [2018,](#page-23-6) Definition 3.1] of the class of unit functions in the RKHS. The bound $\Re_n^2 \leq \bar{k}/n$ [\[Bartlett and Mendelson,](#page-22-7) [2002\]](#page-22-7), and an application of Jensen's inequality conclude the claim. With a more careful analysis, [Tolstikhin et al.](#page-23-8) $[2017,$ Proposition A.1] halved the constant of the first term in [\(5\)](#page-3-1), hence showed that with probability at least $1 - \delta$,

$$
\|\widehat{\mu}_{\mathbb{P}} - \mu_{\mathbb{P}}\|_{\mathcal{H}_k} \le \sqrt{\frac{\overline{k}}{n}} + \sqrt{\frac{2\overline{k}\log 1/\delta}{n}}.
$$
 (6)

What is more, [Tolstikhin et al.](#page-23-8) [\[2017,](#page-23-8) Theorems 1,6,8] provide corresponding lower bounds in $\Omega_P(n^{-1/2})$, showing that the embedding of the empirical measure achieves an unimprovable rate of convergence. Results similar to [\(6\)](#page-3-2) (with worse constants) can be derived when the observations are not independent, see Chérief-Abdellatif and Alquier [\[2022,](#page-22-3) Lemma 7.2].

2 Variance-aware convergence rates

The central quantity we propose to consider is the following variance term in the RKHS,

$$
v_k(\mathbb{P}) \doteq \mathbb{E}_{X \sim \mathbb{P}} \|k(X, \cdot) - \mu_{\mathbb{P}}\|_{\mathcal{H}_k}^2.
$$

It is clear that *v^k* (**P**) depends both on the choice of kernel *k* and on the underlying distribution, and we will simply write $v = v_k(\mathbb{P})$ to avoid encumbering notation. Simple calculations show that

$$
\mathbb{E} \left\| \widehat{\mu}_{\mathbb{P}} - \mu_{\mathbb{P}} \right\|_{\mathcal{H}_k}^2 = \frac{v}{n},
$$

where the expectation is taken with respect a sample $X_1, \ldots, X_n \sim \mathbb{P}$, thus by applications of Jensen's and Chebyshev's inequalities, we readily obtain a rate on the expected risk in terms of *v*,

$$
\mathbb{E} \|\widehat{\mu}_{\mathbb{P}} - \mu_{\mathbb{P}}\|_{\mathcal{H}_k} \leq \sqrt{\frac{v}{n}},
$$

and a deviation bound

$$
\mathbb{P}\left(\left\|\widehat{\mu}_{\mathbb{P}}-\mu_{\mathbb{P}}\right\|_{\mathcal{H}_k}>(1+\tau)\sqrt{\nu/n}\right)\leq 1/(1+\tau)^2, \tau\in(0,\infty).
$$

Another feature of v is linearity with respect the reproducing kernel. Reformulate

$$
v = \mathbb{E}_{X \sim \mathbb{P}} [k(X, X)] - \mathbb{E}_{X, X' \sim \mathbb{P}} [k(X, X')] ,
$$

and suppose that $k = \sum_{k_i \in \mathcal{K}} \alpha_i k_i$ where $\mathcal{K} = \left\{k_1,\ldots,k_{|\mathcal{K}|}\right\}$ is a collection of reproducing kernels and $\alpha \in \mathbb{R}^{|\mathcal{K}|}$. It then holds that

$$
v_k = \sum_{k_i \in \mathcal{K}} \alpha_i v_{k_i}.
$$

Moving on to high-probability confidence bounds, an application of Bernstein's inequality in Hilbert spaces [\[Pinelis and Sakhanenko,](#page-23-9) [1985,](#page-23-9) [Yurinsky,](#page-23-10) [1995\]](#page-23-10) not only recovers the rate of convergence, as alluded to by [Tolstikhin et al.](#page-23-8) [\[2017\]](#page-23-8), but in fact yields the following finer maximal inequality.

Theorem 2.1 (Variance-aware high-confidence convergence rate). Let $X_1, \ldots, X_n \sim \mathbb{P}$ *, k*: $\mathcal{X} \times \mathcal{X} \to \mathbb{R}$ *be a reproducing kernel, and let*

$$
v \doteq \mathbb{E}_{X \sim \mathbb{P}} \| k(X, \cdot) - \mu_{\mathbb{P}} \|_{\mathcal{H}_k}^2.
$$

With probability at least $1 - \delta$ *, it holds that*

$$
\max_{1\leq t\leq n}\left\{t\left\|\widehat{\mu}_{\mathbb{P}}(X_{1},\ldots,X_{t})-\mu_{\mathbb{P}}\right\|_{\mathcal{H}_{k}}\right\}\leq\sqrt{2vn\log 1/\delta}+\frac{4}{3}\sqrt{\bar{k}}\log 1/\delta.
$$

In particular, with probability at least $1 - \delta$ *, it holds that*

$$
\|\widehat{\mu}_{\mathbb{P}}(X_1,\ldots,X_n)-\mu_{\mathbb{P}}\|_{\mathcal{H}_k}\leq \mathcal{B}_{k,\delta}(\mathbb{P},n),
$$

with

$$
\mathcal{B}_{k,\delta}(\mathbb{P},n) = \mathcal{B}_{\delta} \doteq \sqrt{2v\frac{\log 1/\delta}{n}} + \frac{4}{3}\sqrt{k}\frac{\log 1/\delta}{n},
$$

(we remind that $\bar{k} = \sup_{x \in \mathcal{X}} k(x, x)$ *).*

Remark 2.1. *Since the reproducing kernel is bounded, it is always the case that*

$$
v \leq \mathbb{E}_{X \sim \mathbb{P}} \| k(X, \cdot) \|_{\mathcal{H}_k}^2 \leq \mathbb{E}_{X \sim \mathbb{P}} k(X, X) \leq \overline{k},
$$

where the first inequality can be found e.g. in [\[Ch´erief-Abdellatif and Alquier,](#page-22-3) [2022,](#page-22-3) Lemma 7.1, Proof]. As a result, Theorem [2.1](#page-4-0) *strictly supersedes* [\(6\)](#page-3-2) *at least when* $n > (16/9) \log^2 1/\delta$ *.*

Remark 2.2. *The Bernstein approach, in contrast to bounded differences, has the additional advantage of yielding a maximal inequality, further opening the door for early stopping methods.*

We immediately observe that:

- (*O*1) While the bound in [\(6\)](#page-3-2) depends solely on chosen quantities and is computable without any knowledge of \mathbb{P} , this is not the case for \mathcal{B}_{δ} in Theorem [2.1.](#page-4-0)
- (*O*2) Perhaps even more concerning, it is a priori unclear how *v* depends on properties of *k* and **P**, thus making it difficult to convert assumptions on **P** into an upper bound for *v*.

We defer (O1) to Section [3](#page-6-0) and first address (O2) by pointing out that when $\mathcal{X} = \mathbb{R}^d$ and for numerous hyper-parametrized families of kernels, it is possible to promptly obtain upper bounds on *v* that decouple the influence of the hyper-parameter and some measure of spread of the underlying distribution.

Gaussian kernel. For a TI kernel –see (3) – we can rewrite v as

$$
v = \overline{\psi} - \mathbb{E}_{X,X' \sim \mathbb{P}} \left[\psi(X - X') \right].
$$

The Gaussian kernel with lengthscale parameter $\gamma > 0$, defined by

$$
k_{\gamma}(x, x') = \exp\left(-\frac{\|x - x'\|_2^2}{2\gamma^2}\right),\tag{7}
$$

is the prototypical example of a characteristic translation invariant kernel, and satisfies $\overline{\psi}=1.$ For $x\in\mathbb{R}^d$, let x^i denote the *i*th component of *x*. The function $z \mapsto e^{-z}$ is convex, thus from Jensen's inequality

$$
v_{\gamma} = \mathbb{E}_{X \sim \mathbb{P}} ||k_{\gamma}(X, \cdot) - \mu_{\mathbb{P}}||_{\mathcal{H}_{k_{\gamma}}}^{2} \leq 1 - \exp\left(-\frac{\mathbb{E}_{X, X' \sim \mathbb{P}} ||X - X'||_{2}^{2}}{2\gamma^{2}}\right).
$$

We can further rewrite the expectation on the right side of this inequality as

$$
\mathbb{E}_{X,X'\sim P} \|X - X'\|_2^2 = \sum_{i=1}^d \mathbb{E}_{X,X'\sim P} \left[(X^i - X'^i)^2 \right] = \sum_{i=1}^d 2\mathbb{V}_{X\sim P} X^i = 2 \operatorname{Tr} \Sigma_P,
$$

where $\Sigma_{\mathbb{P}} \doteq \mathbb{V}_{X \sim \mathbb{P}} X = \mathbb{E}_{X \sim \mathbb{P}} [(X - \mathbb{E}X)(X - \mathbb{E}X)^*]$ is the covariance matrix of \mathbb{P} and its trace is interpreted as a measure a total variance, agnostic to correlations between distinct components. As a result, for any fixed $\gamma > 0$, we obtain from Theorem [2.1](#page-4-0) that with probability at least $1 - \delta$,

$$
\|\widehat{\mu}_{\mathbb{P}} - \mu_{\mathbb{P}}\|_{\mathcal{H}_{k_{\gamma}}} \leq \sqrt{2\left(1 - e^{-\operatorname{Tr}\Sigma_{\mathbb{P}}/\gamma^2}\right)}\frac{\log 1/\delta}{n} + \frac{4}{3}\frac{\log 1/\delta}{n}.\tag{8}
$$

While γ has no influence over the right-hand side in [\(6\)](#page-3-2), it is clear that for $\gamma \to \infty$, the rate of convergence will be accelerated in [\(8\)](#page-5-0).

Example 2.1 (Gaussian location model with known variance). Assume that $\mathbb{P} = \mathbb{P}_{\theta} = \mathcal{N}(\theta, \sigma^2 I_d)$, the *Gaussian distribution with unknown location parameter θ* ∈ **R***^d , but with known covariance matrix σ* 2 *Id . It holds that* $\text{Tr }\Sigma_{\mathbb{P}_{\theta}} = \sigma^2 d$.

- (*i*) *Fixing* γ *and taking* $\sigma \rightarrow 0$ *, the bound in* [\(8\)](#page-5-0) *vanishes, unlike* [\(6\)](#page-3-2)*.*
- (*ii*) *Setting* $\gamma^2 = \lambda \sigma^2 d$ with $\lambda > 0$, we readily obtain the variance upper bound $v \leq (1 e^{-1/\lambda}) \leq 1/\lambda$, *enabling uncomplicated tuning of the convergence rate by λ.*

Remark 2.3. *Example [2.1](#page-5-1) highlights that if we allow the lengthscale parameter to vary with the sample size n, Theorem* [2.1](#page-4-0) speeds up the convergence rate dramatically. For instance, setting $\lambda_n = -1/\log(1-1/n)$, we achieve *a rate in* O*P*(*n* −1)*, subduing the lower bounds of [Tolstikhin et al.](#page-23-8) [\[2017\]](#page-23-8). In fact, we can reach any prescribed rate* of convergence between $\mathcal{O}_P(n^{-1/2})$ and $\mathcal{O}_P(n^{-1})$ for a suitable choice of λ_n . This stands in sharp contrast with *bounds obtained through a bounded differences approach such as* [\(6\)](#page-3-2)*. However, it is important to remember that a larger lengthscale parameter will make the left hand side in* [\(8\)](#page-5-0) *less informative. Depending on the application, it is possible to achieve the optimal balance between these two effects, some examples are discussed in Section [4.](#page-9-0)*

Convex radial square basis functions. In fact, a technique similar to that for obtaining [\(8\)](#page-5-0) yields a bound on the variance for any (convex) radial kernel.

Lemma 2.1. Assume that for any $x, x' \in \mathcal{X}$, $k(x, x') = r(-||x - x'||_2^2)$ for some convex function $r: \mathbb{R}_+ \to \mathbb{R}$. *Then*

$$
v\leq \bar{r}-r\left(-\operatorname{Tr}\Sigma_{\mathbb{P}}\right) ,
$$

where $\Sigma_{\mathbb{P}}$ *is the covariance matrix that pertains to* \mathbb{P} *.*

Positive definitive matrix on the finite space. Consider $|\mathcal{X}| < \infty$ and a symmetric positive definite matrix *K* of size $|\mathcal{X}|$. Write $K = \sum_{x \in \mathcal{X}} \lambda_x v_x^T v_x$, where for any $x \in \mathcal{X}$, $Kv_x = \lambda_x v_x$, and by positive definiteness, $\lambda_x > 0$. Then the feature map can be expressed as

$$
K(x,\cdot)=\left(\sqrt{\lambda_{x'}}v_x(x')\right)_{x'\in\mathcal{X}},
$$

and by direct calculation,

$$
\|\widehat{\mu}_{\mathbb{P}} - \mu_{\mathbb{P}}\|_{\mathcal{H}_K}^2 = \sum_{x \in \mathcal{X}} \lambda_x \left(\frac{1}{n} \sum_{t=1}^n v_x(X_t) - \mathbb{E}[v_x(X)]\right)^2,
$$

$$
v = \sum_{x \in \mathcal{X}} \lambda_x \mathbb{V}[v_x(X)].
$$

In particular, for $K = I$,

$$
\|\widehat{\mu}_{\mathbb{P}} - \mu_{\mathbb{P}}\|_{\mathcal{H}_K} = \left\|\widehat{\mathbb{P}}_n - \mathbb{P}\right\|_2, \qquad v = 1 - \|\mathbb{P}\|_2^2,
$$

recovering that with probability at least $1 - \delta$,

$$
\left\|\widehat{\mathbb{P}}_n - \mathbb{P}\right\|_2 \leq \sqrt{2(1 - \|\mathbb{P}\|_2^2)\frac{\log 1/\delta}{n}} + \frac{4}{3}\frac{\log 1/\delta}{n}.
$$

So far, we have shown how to obtain for a natural class of kernels an upper bound on *v* that decouples the choice of *k* from some measure of total variance of **P**. Provided an upper bound on the latter is available (see Example [2.1\)](#page-5-1), we obtain an improved convergence rate. However, we understandably may not have a bound on the variance of **P**, or could have insight about its variance, but only have access to contaminated data (see later Section [4.2.1\)](#page-12-0). Recovering a bound that still enjoys the discussed speed-up without a priori knowledge on **P** is the problem we set out to solve in the next section.

3 Convergence rates with empirical variance proxy

We solve the issue (O1) of not knowing *v* by estimating it from the data, simultaneously to $\mu_{\rm P}$. A short heuristic analysis of the Epanechnikov function (which is not generally a kernel according to our defini-tion) that we conduct in Section [3.1](#page-6-1) hints at an "empirical Bernstein" [\[Audibert et al.,](#page-22-8) [2007,](#page-22-8) [Maurer and](#page-23-11) [Pontil,](#page-23-11) [2009\]](#page-23-11) approach –replacing the variance term by some empirical proxy– which we explore formally in Section [3.2.](#page-7-0) The argument is structured around the pivotal definition of a weakly self-bounding function that we first recall.

Definition 3.1 [\(Maurer](#page-23-12) [\[2006\]](#page-23-12), [Boucheron et al.](#page-22-9) [\[2009\]](#page-22-9)). *Let* $n \in \mathbb{N}$, $t_0 \in [n]$, $x = (x_1, ..., x_{t_0}, ..., x_n) \in \mathcal{X}^n$, $x'_{t_0} \in \mathcal{X}$, and write

$$
x^{(t_0)} = (x_1, \ldots, x_{t_0-1}, x'_{t_0}, x_{t_0+1}, \ldots, x_n),
$$

for the vector where x_{t_0} *has been replaced with* $x_{t'_0}$ *. Let* $f: \mathcal{X}^n \to \mathbb{R}$ *.*

(*i*) *The function f is called weakly* (α, β) -self-bounding when for all $x \in \mathcal{X}^n$,

$$
\sum_{t_0=1}^n \left(f(x) - \inf_{x'_{t_0} \in \mathcal{X}} f\left(x^{(t_0)}\right) \right)^2 \leq \alpha f(x) + \beta.
$$

(*ii*) The function f is said to have the bounded differences property when for all $x \in \mathcal{X}^n$ and all $t_0 \in [n]$,

$$
f(x) - \inf_{x'_{t_0} \in \mathcal{X}} f\left(x^{(t_0)}\right) \leq 1.
$$

3.1 Intuition in the hypercube

Let us take $\mathcal{X} = [0,1]^d$ the binary hypercube for $d \gg 1$, and consider the Epanechnikov (parabolic) function

$$
q(x, x') = 1 - ||x - x'||_2^2 / d.
$$

Note that *q* does not define a proper kernel [\[Cuturi,](#page-22-10) [2009,](#page-22-10) p.9]. Nevertheless, the function *q* is TI, with $\psi(t) = 1 - \|t\|_2^2 / d$, $\overline{\psi} = 1$, we can extend the definition of the variance term

$$
v = \overline{\psi} - \mathbb{E}_{X,X' \sim \mathbb{P}} \left[\psi(X - X') \right] = 2 \operatorname{Tr} \Sigma_{\mathbb{P}} / d.
$$

It is natural to define for $i \in [d]$,

$$
\widehat{v}^{i}(X_{1},...,X_{n}) \doteq \frac{1}{2n(n-1)}\sum_{s=1}^{n}\sum_{t=1}^{n} (X_{t}^{i}-X_{s}^{i})^{2},
$$

that will act as an unbiased empirical proxy for $v^i = V_{X \sim P} \left[X^i \right]$, and introduce $\text{Tr} \hat{\Sigma} = \sum_{i=1}^d \hat{v}^i$, as an estimator for the trace of the covariance matrix $\Sigma_{\mathbb{P}}$.

Lemma 3.1. *For* $x \in [0, 1]^d$, *the function* $\mathcal{X}^m \to \mathbb{R}_+$, $x \mapsto \text{Tr} \hat{\Sigma}(x)/d$ *is weakly* $(n/(n-1), 0)$ *-self-bounding and has the bounded differences property in the sense of Definition [3.1.](#page-6-3)*

As a consequence of Lemma [3.1,](#page-6-2) the technique of [Maurer and Pontil](#page-23-11) [\[2009,](#page-23-11) Theorem 10] provides the following deviations bounds on the square root of the total variance

$$
\mathbb{P}\left(b\left[\sqrt{\text{Tr}\Sigma_{\mathbb{P}}/d}-\sqrt{\text{Tr}\widehat{\Sigma}(X)/d}\right] > \sqrt{\frac{2\log 1/\delta}{n-1}}\right) \leq \delta, \qquad b \in \{-1,1\}.
$$
 (9)

In other words, with high confidence, we can replace the trace of the covariance matrix with its empirical proxy in the convergence bounds.

3.2 Systematic approach

Our goal is to rigorously develop the approach intuited in the previous section to include a large class of reproducing kernels. We propose the following empirical proxy for the variance term *v*,

$$
\widehat{v}(X_1, \ldots, X_n) \doteq \frac{1}{n-1} \sum_{t=1}^n \left(k(X_t, X_t) - \frac{1}{n} \sum_{s=1}^n k(X_t, X_s) \right), \tag{10}
$$

and promptly verify that \hat{v} is an unbiased estimator for v .

Lemma 3.2 (Unbiasedness)**.** *It holds that*

$$
\mathbb{E}\widehat{v}(X_1,\ldots,X_n)=v,
$$

 \mathbf{x} where the expectation is taken over the sample $X_1, \ldots, X_n \sim \mathbb{P}$.

The remainder of this section is devoted to analyzing the self-boundedness properties of \hat{v} under mild conditions on *k* and deriving corresponding empirical confidence intervals around μ **P**.

3.2.1 Translation invariant kernels

We first let *k* be a characteristic TI kernel defined by the positive definite function ψ . The expression of \hat{v} can be simplified as

$$
\widehat{v}(X_1,\ldots,X_n)\doteq\overline{\psi}-\frac{1}{(n-1)n}\sum_{t\neq s}\psi(X_t-X_s). \hspace{1cm} (11)
$$

Since the kernel is characteristic, *ψ* cannot be a constant function, i.e. ∆*ψ* > 0. Under our assumptions, we introduce a function involving *^v*^b and [∆]*^ψ* that is weak self-boundedness and has the bounded differences property.

Lemma 3.3. *Let ψ define a characteristic TI kernel. The function*

$$
\mathcal{X}^n \to \mathbb{R}, x \mapsto \frac{n}{2\Delta \psi} \widehat{v}(x),
$$

is weakly (2, 0)*-self-bounding and has bounded differences in the sense of Definition [3.1.](#page-6-3)*

This property leads to concentration of $\sqrt{\widehat{v}}$ around \sqrt{v} .

Lemma 3.4. *For b* \in {−1, 1}*, with probability at least* 1 – *δ,*

$$
b\left[\sqrt{\hat{v}} - \sqrt{v}\right] \le 2\sqrt{\frac{2\Delta\psi\log 1/\delta}{n}}.\tag{12}
$$

Theorem 3.1 (Empirical variance high-confidence convergence rate for translation invariant kernel)**.** *Let n* ∈ **N***, X*¹ , . . . , *Xⁿ* ∼ **P***, let k be a characteristic translation invariant kernel defined from a positive definite function* ψ *[see* [\(3\)](#page-2-2)*]*. Then with probability at least $1 - \delta$, it holds that

$$
\|\widehat{\mu}_{\mathbb{P}} - \mu_{\mathbb{P}}\|_{\mathcal{H}_k} \leq \widehat{\mathcal{B}}_{k,\delta}(X_1,\ldots,X_n),
$$

with

$$
\widehat{\mathcal{B}}_{k,\delta}(X_1,\ldots,X_n)=\widehat{\mathcal{B}}_{\delta}\doteq \sqrt{2\widehat{v}(X_1,\ldots,X_n)\frac{\log 2/\delta}{n}}+\frac{16}{3}\sqrt{\Delta \psi}\frac{\log 2/\delta}{n},
$$

and where \hat{v} *is the empirical variance proxy defined in* [\(11\)](#page-7-5)*.*

Example 3.1. *Theorem [3.1](#page-8-1) immediately holds for Gaussian kernels, regardless of the lengthscale parameter.*

3.2.2 General kernels

Let us define diag $k: \mathcal{X} \to \mathbb{R}$ by diag $k(x) = k(x, x)$, and use the shorthands diag k , diag k and Δ diag k introduced in (2) accordingly. We show that the properties obtained in Section $3.2.1$ can be extended to non TI kernel when ∆ diag *k* is controlled from above.

Lemma 3.5. *Let k be a characteristic reproducing kernel. The function*

$$
\mathcal{X}^n \to \mathbb{R}, x \mapsto \frac{n}{\Delta \operatorname{diag} k + 2\Delta k} \widehat{v}(x),
$$

is weakly

$$
\left(2, 2n \frac{\Delta \text{ diag } k}{\Delta \text{ diag } k + 2\Delta k}\right) \text{-self-bounding},
$$

and has bounded differences in the sense of Definition [3.1.](#page-6-3)

Lemma 3.6. *For b* \in {-1,1}*, with probability at least* 1 – *δ,*

$$
b\left[\sqrt{\widehat{v}(X) + \Delta \operatorname{diag} k} - \sqrt{v + \Delta \operatorname{diag} k}\right] \le 2\sqrt{\frac{(\Delta \operatorname{diag} k + 2\Delta k) \log 1/\delta}{n}}.\tag{13}
$$

Attempting to recover concentration of $\sqrt{\hat{v}}$ around \sqrt{v} with self-boundedness leads to an additional term in $\mathcal{O}(n^{-1/4})$ in [\(13\)](#page-8-5) when ∆ diag $k \neq 0$. We thus settle for concentration of $\sqrt{\hat{v} + \Delta \text{diag } k}$ around $\sqrt{v + \Delta \text{diag } k}$ instead.

Theorem 3.2 (Empirical variance high-confidence convergence rate for general kernel). *Let* $n \in \mathbb{N}$, $X_1, \ldots, X_n \sim$ **P***, let k be a characteristic reproducing kernel defined from a positive definite function ψ [see* [\(3\)](#page-2-2)*]. Then with probability at least* $1 - \delta$ *, it holds that*

$$
\|\widehat{\mu}_{\mathbb{P}} - \mu_{\mathbb{P}}\|_{\mathcal{H}_k} \leq \widehat{\mathcal{B}}_{k,\delta}(X_1,\ldots,X_n),
$$

with

$$
\hat{B}_{k,\delta}(X_1,\ldots,X_n) = \hat{B}_{\delta} \doteq \sqrt{2(\hat{v}(X_1,\ldots,X_n) + \Delta \operatorname{diag} k) \frac{\log 2/\delta}{n}} + \left(\frac{16}{3}\sqrt{\Delta k} + 2\sqrt{2}\sqrt{\Delta \operatorname{diag} k}\right) \frac{\log 2/\delta}{n},
$$

and where \hat{v} *is the empirical variance proxy defined in* [\(10\)](#page-7-6)*.*

Remark 3.1. *In particular, for a TI kernel* ∆ diag *k* = 0*, recovering the results of the previous section.*

Remark 3.2. *We make the following observations.*

- (*i*) It still almost surely holds that $\hat{v} \leq \Delta k \leq 2\bar{k}$, thus the fully empirical bound can never more than a constant *factor away from* [\(6\)](#page-3-2) *or Theorem [2.1.](#page-4-0)*
- (ii) *Crucially, self-boundedness of* \hat{v} *and being able to apply Theorem* [3.1](#page-8-1) *or Theorem* 3.1 *only depends on the choice of kernel, and not on the properties of the underlying distribution.*
- (*iii*) *Neither Theorem [3.1](#page-8-1) nor Theorem [3.2](#page-8-4) depends on smoothness properties of the kernel.*
- (*iv*) The proxy \hat{v} is computable from data with $\mathcal{O}(n^2)$ calls to the kernel function. In Section [4](#page-9-0) below, we discuss *how to use variance-aware bounds to improve confidence bounds and test procedures based on the minimum-MMD estimator. In general, this estimator is computed by stochastic gradient descent (SGD) [\[Dziugaite et al.,](#page-22-11) [2015,](#page-22-11) [Li et al.,](#page-23-13) [2015,](#page-23-13) [Ch´erief-Abdellatif and Alquier,](#page-22-3) [2022\]](#page-22-3) or variants like the stochastic natural gradient descent [\[Briol et al.,](#page-22-2) [2019\]](#page-22-2). A single step of SGD requires to sample m i.i.d. random variables from* **P***^θ where θ is the current estimate, and to compute the unbiased estimate of the gradient which requires (among others)* O(*mn*) *calls to the kernel function. In the natural version, one must also compute a Jacobian at each step, which requires* O(*n* 2) *calls to the kernel. Morever, in the discussion following Theorem 2 of [Briol et al.](#page-22-2) [\[2019\]](#page-22-2), it is argued that we should take m* \gtrsim *n in these algorithms. Thus, both SGD and its natural variant will require* at least $\mathcal{O}(n^2)$ calls to the kernel at each iteration. In this light, the computation of \widehat{v} does not significantly
affect the computational burden. Note however that, in a fern situations where the oradiant of *affect the computational burden. Note however that, in a few situations where the gradient of the MMD is available in close form, it is possible to use a non-stochastic gradient descent. Such examples include Gaussian mean estimation [\[Ch´erief-Abdellatif and Alquier,](#page-22-3) [2022\]](#page-22-3) and Gaussian copula estimation [\[Alquier et al.,](#page-22-12) [2022\]](#page-22-12). Each step of the gradient descent requires only* O(*n*) *calls to the kernel and convergence is typically very fast.* In these situations, the computation of \hat{v} can increase the computational cost when n is large.

4 Applications

We put the apparatus developed in Sections 2 and Section 3 to application in the context of hypothesis testing and robust parametric estimation. In this section, all considered kernels will be TI unless otherwise specified, and X_1, \ldots, X_n are i.i.d. from $\mathbb{P} \in \mathcal{P}(\mathcal{X})$. We introduce a statistical model $\mathcal{M} = \{\mathbb{P}_{\theta} : \theta \in \Theta\}$ indexed by the parameter space Θ. Examples of models studied in the literature on MMD include parametric models such as the Gaussian model $\mathbb{P}_{\theta} = \mathcal{N}(\theta, \sigma^2 I_d)$ (with σ^2 known) and mixture of Gaussians [\[Briol et al.,](#page-22-2) [2019,](#page-22-2) Chérief-Abdellatif and Alquier, [2022\]](#page-22-3), copulas [Alquier et al.](#page-22-12) [\[2022\]](#page-22-12) but also more com-plex models such as generative adversarial networks [\[Dziugaite et al.,](#page-22-11) [2015,](#page-22-11) [Li et al.,](#page-23-13) [2015\]](#page-23-13), stochastic volatility models and stochastic differential equations [\[Briol et al.,](#page-22-2) [2019\]](#page-22-2). Following [Briol et al.](#page-22-2) [\[2019\]](#page-22-2), our estimator will be the closest element^{[3](#page-9-2)} of the model M to the empirical measure obtained from *X*, where the distance is measured in the RKHS,

$$
\left\|\mu_{\mathbb{P}_{\hat{\theta}_n(X_1,\ldots,X_n)}}-\widehat{\mu}_{\mathbb{P}}(X_1,\ldots,X_n)\right\|_{\mathcal{H}_k}=\inf_{\theta\in\Theta}\left\|\mu_{\mathbb{P}_{\theta}}-\widehat{\mu}_{\mathbb{P}}(X_1,\ldots,X_n)\right\|_{\mathcal{H}_k}.
$$

The computation of $\theta_n(X_1, \ldots, X_n)$ is usually done via stochastic gradient descent and variants, see Remark [3.2](#page-9-3) above.

4.1 Hypothesis testing

In this subsection we wish to test the hypothesis $H_0: \mathbb{P} \in \mathcal{M} = {\mathbb{P}}_\theta: \theta \in \Theta$, under the alternative $H_1: \mathbb{P} \notin \{\mathbb{P}_{\theta} \colon \theta \in \Theta\}$. While "two-sample testing", performed on the data of two random samples, each

³Note that it might be that the infimum is reached by multiple elements of Θ , or is not reached. The first case does not lead to any difficulty, we must simply define a rule to break ties (for example, we can equip Θ with a total order and chose the smallest minimizer according to this order). While [Briol et al.](#page-22-2) [\[2019\]](#page-22-2) provide sufficient conditions to ensure that the infimum is reached, the non-existence of a minimizer is also not a problem in practice: all the non-asymptotic results in [Briol et al.](#page-22-2) [\[2019\]](#page-22-2) and Chérief-Abdellatif and Alquier [\[2022\]](#page-22-3) can easily be extended to any ϵ -minimizer, for ϵ small enough.

independently obtained from a different distribution, is more prevalent in the MMD literature [\[Gretton](#page-22-13) [et al.,](#page-22-13) [2009,](#page-22-13) [2012\]](#page-22-14), recent works also tackle goodness of fit testing [\[Chwialkowski et al.,](#page-22-15) [2016,](#page-22-15) [Jitkrittum](#page-22-16) [et al.,](#page-22-16) [2017\]](#page-22-16), albeit with an asymptotic treatment. Recall that we define the significance level *α* ∈ [0, 1] of a test as the probability of outputting **H**¹ when **H**⁰ is true. Taking advantage of a non-asymptotic bound $\mathcal{B}(n,\alpha)$ (for example $\mathcal{B}(n,\alpha) = \hat{\mathcal{B}}_{\alpha}$ given in Theorem [3.1\)](#page-8-1), we can design a test with prescribed level α for any *n* as follows. We introduce the test statistic

$$
T(X_1,\ldots,X_n)\doteq\inf_{\theta\in\Theta}\|\widehat{\mu}_{\mathbb{P}}-\mu_{\mathbb{P}_{\theta}}\|_{\mathcal{H}_k},
$$

and reject H_0 on the "critical set"

$$
\mathcal{C}(X_1,\ldots,X_n)\doteq \{T>\mathcal{B}(n,\alpha)\}.
$$

We show that the probability of rejection under the null hypothesis is at most α and that the test is consistent.

Theorem 4.1. Let $n \in \mathbb{N}$, $X_1, \ldots, X_n \sim \mathbb{P}$, and k a translation invariant kernel. Let $\mathcal{B}(n, \alpha)$ be such that

- *(a) for any fixed* α *,* $\mathcal{B}(n, \alpha) \xrightarrow[n \to \infty]{a.s.} 0$,
- *(b) there is a sequence* $\alpha_n \to 0$ *such that* $\mathcal{B}(n, \alpha_n) \xrightarrow[n \to \infty]{a.s.} 0$,
- *(c) for any α, with probability at least* $1 α$ *,* $\|\hat{\mu}_{\mathbf{P}} \mu_{\mathbf{P}}\|_{\mathcal{H}_k} \leq \mathcal{B}(n, α)$ *.*

Then

- $((i)$ $\mathbb{P}_{\mathbf{H}_0}(\mathcal{C}(X_1,\ldots,X_n)) \leq \alpha.$
- (*ii*) When the model M is closed with respect to the MMD metric $\frac{4}{7}$ $\frac{4}{7}$ $\frac{4}{7}$,

$$
\lim_{n\to\infty} \mathbb{P}_{\mathbf{H}_1} \left(\mathcal{C}(X_1,\ldots,X_n) \right) = 1,
$$

more precisely,

$$
1 - \mathbb{P}_{\mathbf{H}_1}(\mathcal{C}(X_1,\ldots,X_n)) = \mathcal{O}(\alpha_n).
$$

Obviously, the empirical bound \hat{B}_α of Theorem [3.1](#page-8-1) satisfies (*a*) and (*c*), and (*b*) with $\alpha_n = \exp(-n^{1-\epsilon})$ for any fixed $\epsilon > 0$. So does the McDiarmid-based bound in [\(6\)](#page-3-2). On the other hand, we claimed that the empirical bound is smaller than bounds that does not take the variance into account. In other words, the power of the test $\mathbb{P}_{H_1}(\mathcal{C}(X_1,\ldots,X_n))$ for a finite *n* will be larger if we use the variance aware bound.

We illustrate this on a toy example in \mathbb{R}^2 : here $\mathbb{P} = \mathcal{N}(0, \sigma^2 I_2)$ while $\mathbb{P}_{\theta} = \mathcal{N}(\theta, I_2)$. In other words, H_0 is true iff $\sigma = 1$. We sample X_1, \ldots, X_n from $\mathbb P$, perform the test based on the empirical bound and the test based on McDiarmid bound. This is repeated 100 times for each value $\sigma \in \{0, 1/50, 2/50, \ldots, 1\}$. We report the frequency of rejections in Figure [1.](#page-11-0) The kernel used is a Gaussian kernel with $\gamma = 1$, and we consider sample sized $n \in \{16, 40, 100, 250\}$. Both tests have a weak power (note that we did not try to optimize *γ* for now, this question will be tackled later). However, the test based on the empirical bound indeed rejects more often \mathbf{H}_0 when \mathbf{H}_1 is true. Note that the improvement is clearer for small sample sizes.

4.2 Robust parametric estimation under Hüber contamination

Chérief-Abdellatif and Alquier [\[2022,](#page-22-3) Proof of Theorem 3.1] show how to bound the above by the estimation error of the empirical measure in the RKHS and the approximation error of the model,

$$
\left\|\mu_{\mathbb{P}_{\hat{\theta}_n}} - \mu_{\mathbb{P}}\right\|_{\mathcal{H}_k} \leq \inf_{\theta \in \Theta} \left\|\mu_{\mathbb{P}_{\theta}} - \mu_{\mathbb{P}}\right\|_{\mathcal{H}_k} + 2\left\|\widehat{\mu}_{\mathbb{P}} - \mu_{\mathbb{P}}\right\|_{\mathcal{H}_k}.
$$
\n(14)

⁴For any Q, if there is a sequence $(\theta_h)_{h\in \mathbb N}$ of elements of Θ such that $\left\|\mu_\mathbb{Q}-\mu_{\mathbb{P}_{\theta_h}}\right\|_{\mathcal{H}_k} \xrightarrow[h\to\infty]{} 0$, then $\mathbb{Q}=\mathbb{P}_{\theta}$ for some *θ* ∈ Θ.

Figure 1: Comparison of the test based on the Bernstein empirical (EmpBer) bound, versus the test based on McDiarmid bound (McDia). Frequency of rejection of $H_0: \mathbb{P} \in \{\mathcal{N}(\theta, I_2), \theta \in \mathbb{R}^2\}$ as a function of σ with $\mathbb{P} = \mathcal{N}(0, \sigma^2 I_2)$.

Recall that in the Hüber contamination model [Hüber, [2011\]](#page-22-17), the observations X_1, \ldots, X_n are drawn independently from the mixture

$$
\mathbb{P} = (1 - \xi)\mathbb{P}_{\theta_0} + \xi \mathbb{H},
$$

where $\xi \in [0, 1/2)$ is the contamination rate, **H** is some unknown noise distribution and $\theta_0 \in \Theta$. In this setting, we can control the approximation error of the model: starting from [\(14\)](#page-10-3), and adapting the argument of Chérief-Abdellatif and Alquier [\[2022,](#page-22-3) Corollary 3.3], we obtain

$$
\left\|\mu_{\mathbb{P}_{\hat{\theta}_n}}-\mu_{\mathbb{P}_{\theta_0}}\right\|_{\mathcal{H}_k}\leq 4\xi\sqrt{\overline{\psi}}+2\left\|\widehat{\mu}_{\mathbb{P}}-\mu_{\mathbb{P}}\right\|_{\mathcal{H}_k}.
$$

We wish to apply Theorem [2.1](#page-4-0) to the second term. However, when the sample is contaminated, the variance term $v(\mathbb{P})$ we collect also depends on the properties of H, which are unknown –for all we know H does not even have finite moments. Fortunately, we now see how to bound $v(\mathbb{P})$ is terms of $v(\mathbb{P}_{\theta_0})$ and ξ .

4.2.1 Improved confidence bounds in the Hüber contamination model

Lemma 4.1. In the Hüber contamination model, i.e. $\mathbb{P} = (1 - \xi)\mathbb{P}_{\theta_0} + \xi\mathbb{H}$, writing $v = v(\mathbb{P})$ and $v_0 = v(\mathbb{P}_{\theta_0})$, *it holds that*

$$
v \le v_0 + 2\xi(\Delta \psi - v_0) + \xi^2(v_0 - \Delta \psi). \tag{15}
$$

The bound can be streamlined; when $\psi \geq 0$ *,*

$$
v\leq (1-2\xi)v_0+2\xi\overline{\psi},
$$

and otherwise

$$
v \le (1 - 2\xi)v_0 + 2\xi \Delta \psi.
$$

Remark 4.1. *In* [\(15\)](#page-12-3)*, observe that when* $\psi \ge 0$ *and* $\xi \to 1$ *, the first term vanishes, and* $v \to \overline{\psi}$ *, recovering the distribution independent rates. Conversely, when* $\xi \to 0$, we confirm that $v \to v_0$, which could be further bounded *using properties of the model. In practice, we will focus on small values* ζ ≤ 1/2, and the simplified bounds will be *sufficient.*

An application of Theorem [2.1](#page-4-0) yields the following.

Corollary 4.1. In the Hüber contamination model $\mathbb{P} = (1 - \xi)\mathbb{P}_{\theta_0} + \xi\mathbb{H}$, writing $v = v(\mathbb{P})$ and $v_0 = v(\mathbb{P}_{\theta_0})$, *with probability at least* $1 - \delta$ *,*

$$
\left\|\mu_{\mathbb{P}_{\widehat{\theta}_n}}-\mu_{\mathbb{P}_{\theta_0}}\right\|_{\mathcal{H}_k}\leq 4\xi\sqrt{\overline{\psi}}+2\sqrt{2[(1-2\xi)v_0+2\xi\overline{\psi}]\frac{\log 1/\delta}{n}}+2\frac{4\sqrt{\overline{\psi}}}{3}\frac{\log 1/\delta}{n}.
$$

When $\overline{\psi} = 1, v_0 \ll 1, \xi \ll 1$, the bound offers a significant improvement over Chérief-Abdellatif and [Alquier](#page-22-3) [\[2022,](#page-22-3) Corollary 3.4].

Example 4.1 (Gaussian location model)**.** *Assume that the model is*

$$
\mathcal{M}_{\sigma^2} = \left\{ \mathcal{N}(\theta, \text{diag}\,\sigma^2) \colon \theta \in \mathbb{R}^d \right\},\
$$

i.e. the collection of Gaussian random vectors with unknown location parameter θ ∈ **R***^d and known variance parameter* $\sigma^2 = (\sigma_1^2, \ldots, \sigma_d^2)$. Conducting estimation with respect to the Gaussian kernel k_γ [\(7\)](#page-4-1), the variance term *is bounded by*

$$
v \le (1-2\xi) \frac{\|\sigma\|_2^2}{\gamma^2} + 2\xi,
$$

which becomes small when $\|\sigma\|_2 \ll \gamma$ *.*

4.2.2 Improved confidence bounds in the parameter space

In this subsection we still employ the Gaussian kernel $k_γ$ [\(7\)](#page-4-1). It is instructive to analyze how robust estimation with respect to the MMD distance translates into what happens in the space of parameters. In fact, since we have freedom over the lengthscale parameter, we may understandably want to select *γ* such that the distance in the space of parameters is kept small as well.

Definition 4.1 (Link function *F*). Let $M = \{ \mathbb{P}_{\theta} : \theta \in \Theta \}$ *indexed by the parameter space* $\Theta \subset \mathcal{P}(\mathbb{R}^d)$ *. We* say that a non-decreasing function $F_k\colon\mathbb R_+\to\mathbb R_+$ is a link function for the model $\cal M$ and kernel k when for any $θ$ *,* $θ'$ ∈ $Θ$ *,*

$$
\|\theta-\theta'\|_2\leq F_k\left(\left\|\mu_{\mathbb{P}_{\theta}}-\mu_{\mathbb{P}_{\theta'}}\right\|_{\mathcal{H}_k}\right).
$$

Remark 4.2. Note that a link function F_k will only provide nontrivial information if $F_k(0) = 0$ and F_k is continuous *at* 0. The existence of such a nontrivial link function implies that the model is identifiable: if $\mu_{\mathbb{P}_\theta} = \mu_{\mathbb{P}_{\theta'}}$ then $0=F_k(\|\mu_{\mathbb{P}_\theta}-\mu_{\mathbb{P}_{\theta'}}\|_{\mathcal{H}_k})\geq \|\theta-\theta'\|_2$, hence $\theta=\theta'.$ In this case, there is a unique function $p:\mathcal{M}\to\tilde{\Theta}$ such *that* $p(\mathbb{P}_{\theta}) = \theta$, and F_k is a modulus of continuity of p .

A direct application of Corollary [4.1](#page-12-4) gives the following.

Corollary 4.2. *In the setting of Corollary [4.1,](#page-12-4) assume that* M *and k are such that a link function F^k exists. With probability at least* $1 - \delta$ *,*

$$
\left\|\widehat{\theta}_n-\theta_0\right\|_2 \leq F_k\left(4\xi\sqrt{\overline{\psi}}+2\sqrt{2[(1-2\xi)v_0+2\xi\overline{\psi}]\frac{\log 1/\delta}{n}}+2\frac{4\sqrt{\overline{\psi}}}{3}\frac{\log 1/\delta}{n}\right).
$$

Example 4.2 (Gaussian location model, continued)**.** *We continue Example [4.1.](#page-12-5) A direct computation (see e.g. [Ch´erief-Abdellatif and Alquier](#page-22-3) [\[2022\]](#page-22-3)) shows that*

$$
\left\|\mu_{\mathbb{P}_{\theta}}-\mu_{\mathbb{P}_{\theta'}}\right\|_{\mathcal{H}_{k_{\gamma}}}^{2}=2\left(\frac{\gamma^{2}}{2\sigma^{2}+\gamma^{2}}\right)^{d/2}\left[1-\exp\left(-\frac{\left\|\theta-\theta'\right\|_{2}^{2}}{4\sigma^{2}+2\gamma^{2}}\right)\right].
$$

In other words, a link function for the Gaussian location model is explicitly given by

$$
F_{k_{\gamma}}^{2}(h) = -2(2\sigma^{2} + \gamma^{2})\log\left(1 - \frac{h}{2}\left(1 + 2\frac{\sigma^{2}}{\gamma^{2}}\right)^{d/2}\right)
$$

An application of Corollary [4.2,](#page-13-0) and setting ^γ ⁼ *λσ*[√] *d, for some λ* > 0*, proves that, with probability at least* 1 − *δ,*

$$
\left\|\theta - \hat{\theta}_n\right\|_2^2 \le -2\sigma^2 (2 + d\lambda^2) \log \left\{ 1 - \left(1 + \frac{2}{d\lambda^2}\right)^{d/2} \left(4\xi + \sqrt{2\left[(1 - 2\xi)\left(1 - e^{-\frac{1}{\lambda^2}}\right) + 2\xi\right]} \frac{\log 1/\delta}{n} + \frac{4}{3} \frac{\log 1/\delta}{n}\right)^2 \right\}
$$
(16)

$$
\doteq G_{d,\sigma,n}(\lambda,\xi).
$$

.

For comparison, we provide the bound of [Ch´erief-Abdellatif and Alquier](#page-22-3) [\[2022,](#page-22-3) Proposition 4.1-(2)*] that is obtained by pluging* [\(6\)](#page-3-2) *into the link function:*

$$
\left\|\theta - \hat{\theta}_n\right\|_2^2 \le -2\sigma^2(2 + d\lambda^2)\log\left\{1 - \left(1 + \frac{2}{d\lambda^2}\right)^{d/2}\left(4\xi + \frac{1 + \sqrt{2\log 1/\delta}}{\sqrt{n}}\right)^2\right\}
$$
\n
$$
\stackrel{\text{d}{\rightarrow}}{=} H_{d,\sigma,n}(\lambda,\xi). \tag{17}
$$

We run experiments (see Figure [2](#page-14-0) & [3\)](#page-14-1), and make the following observations.

Figure 2: Comparison of the bounds in [\(16\)](#page-13-1) versus Chérief-Abdellatif and Alquier [\[2022,](#page-22-3) Proposition 4.1-(2)] as a function of $γ$.

Figure 3: Comparison of Chérief-Abdellatif and Alquier [\[2022\]](#page-22-3) and [\(16\)](#page-13-1) for the optimal hyperparameter γ as a function of *n*.

- (*i*) *The new bound is always tighter than the one of [Ch´erief-Abdellatif and Alquier](#page-22-3) [\[2022\]](#page-22-3).*
- (*ii*) *When the contamination level increases, the two bounds are getting closer together. This is expected since we are losing the benefit of the variance factor.*
- (*iii*) *Especially under weak contamination, the new bound is much flatter in the sense where overshooting for the value of γ does not lead to a catastrophic degradation of the bound.*
- (*iv*) *The new bound excels in the small sample setting.*

We conclude this section by showing heuristically that the bound in (16) can always be made smaller than the bound in [\(17\)](#page-13-2), at least asymptotically in *n*, with an adequate choice of λ . For the sake of simplicity, we work in the non-contaminated setting $\xi = 0$.

First,

$$
H_{d,\sigma,n}(\lambda,0) = 4\sigma^2 (2 + d\lambda^2) \left(1 + \frac{2}{d\lambda^2}\right)^{d/2} \frac{(1 + \sqrt{2\log 1/\delta})^2}{n} (1 + o(1))
$$

and the first order term is exactly minimized for $\lambda = 1$, it leads to

$$
H_{d,\sigma,n}(1,0) = 4\sigma^2(2+d)\left(1+\frac{2}{d}\right)^{d/2}\frac{(1+\sqrt{2\log 1/\delta})^2}{n}(1+o(1)).
$$

An exact minimization of the bound in [\(16\)](#page-13-1) is not feasible, but we will propose a choice of $\lambda = \lambda_n$ that will lead to an improvement on $H_{d,\sigma,n}(1,0)$. First, let us assume that $\lambda_1 > 0$ and that $(\lambda_n)_{n \in \mathbb{N}}$ is a non-decreasing sequence. We obtain:

$$
G_{d,\sigma,n}(\lambda_n,0) = 4\sigma^2 (2+d\lambda_n^2) \left(1 + \frac{2}{d\lambda_n^2}\right)^{d/2} \left(1 - e^{-\frac{1}{\lambda_n^2}}\right) \frac{\log 1/\delta}{n} (1+o(1))
$$

=
$$
4\sigma^2 (2+d\lambda_n^2) \left(1 + \frac{2}{d\lambda_n^2}\right)^{d/2} \frac{\log 1/\delta}{\lambda_n^2 n} (1+o(1))
$$

=
$$
4d\sigma^2 \left(1 + \frac{2}{d\lambda_n^2}\right)^{1+d/2} \frac{\log 1/\delta}{n} (1+o(1)).
$$

It is clear that chosing $\lambda_n \to \infty$ when $n \to \infty$ gives:

$$
G_{d,\sigma,n}(\lambda_n,0) = 4d\sigma^2 \frac{\log 1/\delta}{n} (1 + o(1)).
$$

Finally, observe that

$$
\frac{H_{d,\sigma,n}(1,0)}{G_{d,\sigma,n}(\lambda_n,0)} = 2\underbrace{\left(1+\frac{2}{d}\right)^{d/2+1}\underbrace{\frac{(1+\sqrt{2\log 1/\delta})^2}{(\sqrt{2\log 1/\delta})^2}(1+o(1))\geq 2e(1+o(1))}_{\geq 1}.
$$

Unfortunately, this first order analysis is too crude to provide an accurate recommendation on the choice of λ_n . On the other hand, it shows that our variance-aware bound leads to a significant improvement over the bound of Chérief-Abdellatif and Alquier [\[2022\]](#page-22-3) for Gaussian mean estimation. Moreover, the choice $\lambda_n \to \infty$ is in accordance with the considerations on the asymptotic variance in [Briol et al.](#page-22-2) [\[2019\]](#page-22-2) in this model.

We illustrate this with an experiment in Figure [4.](#page-16-1) We observe that: indeed, the variance-aware bound decrease with $\gamma \propto \lambda$, while the McDiarmid bound has a minimum. Interestingly, the true MSE also has a minimum. Note however how the MSE is actually very flat as a function of *γ*: in other words, in this experiment, the choice of γ does not matter so much on the performance of the MMD estimator. The possibility to use the variance-aware bound to calibrate γ in practice should be investigated further in other models.

Figure 4: Comparison of the bounds in [\(16\)](#page-13-1) versus Chérief-Abdellatif and Alquier [\[2022,](#page-22-3) Proposition 4.1-(2)] as a function of γ , versus the empirical MSE on 50 experiments, $n = 500$, log-scale, *ξ* = 0, *δ* = 0.05. Right panel: we zoom on the MSE.

5 Proofs

5.1 Proof of Theorem [2.1](#page-4-0)

In [Tolstikhin et al.](#page-23-8) [\[2017\]](#page-23-8), the authors mention but do not pursue the idea that Bernstein's inequality in separable Hilbert spaces yields the proper rate of $\mathcal{O}_P(n^{-1/2})$. We follow-up on their idea in this proof. First recall that

$$
\widehat{\mu}_{\mathbb{P}} \doteq \frac{1}{n} \sum_{t=1}^{n} k(X_t, \cdot), \qquad \mu_{\mathbb{P}} \doteq \mathbb{E}_{X \sim \mathbb{P}} \left[k(X, \cdot) \right],
$$

and that we can write

$$
\|\widehat{\mu}_{\mathbb{P}} - \mu_{\mathbb{P}}\|_{\mathcal{H}_k} = \frac{1}{n} \left\| \sum_{t=1}^n Z_t \right\|_{\mathcal{H}_k},
$$

where the

$$
Z_t \doteq k(X_t,\cdot) - \mathbb{E}_{X \sim \mathbb{P}} \left[k(X,\cdot) \right],
$$

are centered random variables. We are thus interested in bounding the probability of the following deviation,

$$
\mathbb{P}\left(\max_{1\leq s\leq n}\left\|\sum_{t=1}^s Z_t\right\|_{\mathcal{H}_k} > n\varepsilon\right).
$$

We rely on the following theorem, credited to [Pinelis and Sakhanenko](#page-23-9) [\[1985\]](#page-23-9).

Theorem 5.1 (Bernstein inequality in Hilbert space [see e.g. [Yurinsky](#page-23-10) [\[1995,](#page-23-10) Theorem 3.3.4])**.** *] Let Z*¹ , . . . , *Zn be independent random variables in a separable Hilbert space* H *that are centered, i.e.*

$$
\forall t \in [n], \mathbb{E}[Z_t] = 0,
$$

If there exists real numbers B, H \geq 0 *such that for any p* \geq 2*,*

$$
\sum_{t=1}^{n} \mathbb{E} \|Z_t\|_{\mathcal{H}}^p \le \frac{1}{2} p! B^2 H^{p-2},
$$

it holds that for any $\varepsilon > 0$ *,*

$$
\mathbb{P}\left(\max_{1\leq s\leq n}\left\|\sum_{t=1}^s Z_t\right\|_{\mathcal{H}} > \varepsilon\right) \leq 2\exp\left(-\frac{\varepsilon^2/2}{B^2+\varepsilon H}\right).
$$

 \mathcal{H}_k is separable by separability of the topological space $\mathcal X$ and continuity of k . For $p=2$,

$$
\sum_{t=1}^n \mathbb{E} \| k(X_t, \cdot) - \mu_{\mathbb{P}} \|_{\mathcal{H}_k}^2 = n v.
$$

Moving on to higher moments, for $p \geq 3$, we have that

$$
||Z_t||_{\mathcal{H}_k} \leq ||k(X_t, \cdot)||_{\mathcal{H}_k} + ||\mathbb{E}_{X \sim \mathbb{P}} [k(X, \cdot)]||_{\mathcal{H}_k}
$$

\n
$$
\leq \sqrt{k(X_t, X_t)} + \sup_{x \in \mathcal{X}} \sqrt{k(x, x)}
$$

\n
$$
\leq 2\sqrt{C_k}.
$$

As a result,

$$
\sum_{t=1}^{n} \mathbb{E} \left\| Z_t \right\|_{\mathcal{H}_k}^p \leq \left(\sum_{t=1}^{n} \mathbb{E} \left\| Z_t \right\|_{\mathcal{H}_k}^2 \right) \left(2\sqrt{C_k} \right)^{p-2}
$$

$$
\leq n v (2\sqrt{C_k})^{p-2} \leq \frac{p!}{2} n v \left(\frac{2\sqrt{C_k}}{3} \right)^{p-2}.
$$

Setting $B^2 = nv$, $H = 2\sqrt{C_k}/3$ and invoking Theorem [5.1,](#page-16-2) we obtain that

$$
\mathbb{P}\left(\max_{1\leq s\leq n}\left\|\sum_{t=1}^{s}Z_t\right\|_{\mathcal{H}_k}>n\varepsilon\right)\leq 2\exp\left(-\frac{(\varepsilon n)^2}{2(nv+(\varepsilon n)c\sqrt{C_k})}\right)
$$

$$
=\exp\left(-\frac{n\varepsilon^2}{2\left(v+2\varepsilon\sqrt{C_k}/3\right)}\right).
$$

 \Box

5.2 Proof of Lemma [3.2](#page-7-2)

We verify that \widehat{v} is an unbiased estimator for $v.$

$$
\mathbb{E}_{X_1,...,X_n \sim \mathbb{P}} [\hat{v}(X_1,...,X_n)]
$$
\n
$$
= \mathbb{E}_{X_1,...,X_n \sim \mathbb{P}} \left[\frac{1}{n-1} \sum_{t=1}^n \left(k(X_t, X_t) - \frac{1}{n} \sum_{s=1}^n k(X_t, X_s) \right) \right]
$$
\n
$$
= \frac{1}{n-1} \sum_{t=1}^n \left(\mathbb{E}_{X \sim \mathbb{P}} [k(X, X)] - \frac{1}{n} \sum_{s=1}^n \mathbb{E}_{X_t, X_s \sim \mathbb{P}} [k(X_t, X_s)] \right)
$$
\n
$$
= \frac{n}{n-1} \mathbb{E}_{X \sim \mathbb{P}} [k(X, X)] - \frac{1}{n(n-1)} \sum_{t=1}^n \sum_{s=1}^n \mathbb{E}_{X_t, X_s \sim \mathbb{P}} [k(X_t, X_s)]
$$
\n
$$
= \frac{n}{n-1} \mathbb{E}_{X \sim \mathbb{P}} [k(X, X)] - \frac{1}{(n-1)} \mathbb{E}_{X \sim \mathbb{P}} [k(X, X)]
$$
\n
$$
- \frac{n^2 - n}{n(n-1)} \mathbb{E}_{X, X' \sim \mathbb{P}} [k(X, X')]
$$
\n
$$
= \mathbb{E}_{X \sim \mathbb{P}} [k(X, X)] - \mathbb{E}_{X, X' \sim \mathbb{P}} [k(X, X')]
$$
\n
$$
= v.
$$

 \Box

5.3 Proof of Lemma [3.3](#page-7-3)

We first verify the bounded-differences property.

$$
\hat{v}(x) - \inf_{x'_{t_0} \in \mathcal{X}} \hat{v}(x^{(t_0)})
$$
\n
$$
= \overline{\psi} - \frac{1}{(n-1)n} \sum_{t \neq s} \psi(x_t - x_s) - \inf_{x'_{t_0} \in \mathcal{X}} \left[\overline{\psi} - \frac{1}{(n-1)n} \sum_{t \neq s} \psi(x_t^{(t_0)} - x_s^{(t_0)}) \right]
$$
\n
$$
= \frac{1}{(n-1)n} \left(\sup_{x'_{t_0} \in \mathcal{X}} \left\{ \sum_{t \neq s} \psi(x_t^{(t_0)} - x_s^{(t_0)}) \right\} - \sum_{t \neq s} \psi(x_t - x_s) \right)
$$
\n
$$
= \frac{1}{(n-1)n} \left(\sup_{x'_{t_0} \in \mathcal{X}} \left\{ \sum_{t \neq s} \psi(x_t^{(t_0)} - x_s^{(t_0)}) \right\} - \sum_{t \neq s} \psi(x_t - x_s) \right)
$$
\n
$$
= \frac{2}{(n-1)n} \left(\sup_{x'_{t_0} \in \mathcal{X}} \left\{ \sum_{t \in [n]} \psi(x_t^{(t_0)} - x_t) \right\} - \sum_{t \in [n]} \psi(x_t^{(t_0)} - x_t) \right\}
$$
\n
$$
\leq \frac{2}{(n-1)n} \left((n-1)\overline{\psi} - \sum_{t \in [n]} \psi(x_t^{(t_0)} - x_t) \right)
$$
\n
$$
\frac{(\star)}{\star} \frac{2}{\pi} \left(\overline{\psi} - \frac{1}{n-1} \sum_{t \in [n]} \psi(x_t^{(t_0)} - x_t) \right)
$$
\n
$$
\leq \frac{2}{n} \Delta \psi.
$$

It follows from (\star) that,

$$
\sum_{t_0=1}^n \left(\widehat{v}(x) - \inf_{x'_{t_0} \in \mathcal{X}} \widehat{v} \left(x^{(t_0)} \right) \right)^2 \le \sum_{t_0=1}^n \left(\frac{2}{n} \left(\overline{\psi} - \frac{1}{n-1} \sum_{\substack{t \in [n] \\ t \neq t_0}} \psi(x_{t_0} - x_t) \right) \right)^2
$$
\n
$$
= \frac{4}{n^2} \sum_{t_0=1}^n \left(\frac{1}{n-1} \sum_{\substack{t \in [n] \\ t \neq t_0}} \left(\overline{\psi} - \psi(x_{t_0} - x_t) \right) \right)^2
$$
\n
$$
\le \frac{4}{n^2} \sum_{t_0=1}^n \frac{1}{n-1} \sum_{\substack{t \in [n] \\ t \neq t_0}} \left(\overline{\psi} - \psi(x_{t_0} - x_t) \right)^2
$$
\n
$$
\le \frac{4\Delta \psi}{n} \sum_{t_0=1}^n \frac{1}{(n-1)n} \sum_{\substack{t \in [n] \\ t \neq t_0}} \left(\overline{\psi} - \psi(x_{t_0} - x_t) \right)^2
$$
\n
$$
= \frac{4\Delta \psi}{n} \widehat{v}
$$

where the second inequality is Jensen's. $\quad \Box$

5.4 Proof of Lemma [3.4](#page-7-4)

From Lemma [3.3,](#page-7-3) the function

$$
\mathcal{X}^n \to \mathbb{R}, x \mapsto \frac{n}{2\Delta\psi}\widehat{v}(x),
$$

is weakly (2, 0)-self-bounding and has the bounded differences property in the sense of Definition [3.1.](#page-6-3) From the second statement of [Maurer](#page-23-12) [\[2006,](#page-23-12) Theorem 13],

$$
\mathbb{P}\left(\mathbb{E}\left[\frac{n}{2\Delta\psi}\widehat{v}(X)\right] - \frac{n}{2\Delta\psi}\widehat{v}(X) > t\right) \le \exp\left(-\frac{t^2}{4\mathbb{E}\left[\frac{n}{2\Delta\psi}\widehat{v}(X)\right]}\right),
$$

that can be rewritten,

$$
\mathbb{P}\left(v-\widehat{v}(X)>t\right)\leq \exp\left(-\frac{nt^2}{8\Delta\psi v}\right).
$$

In other words, with probability at least $1 - \delta$,

$$
v - 2\sqrt{v} \sqrt{\frac{2\Delta \psi \log 1/\delta}{n}} \le \widehat{v}(X).
$$

Completing the square, and by sub-additivity of the square root function, we successively have with probability $1 - \delta$,

$$
\left(\sqrt{v} - \sqrt{\frac{2\Delta\psi\log 1/\delta}{n}}\right)^2 \le \widehat{v}(X) + \frac{2\Delta\psi\log 1/\delta}{n}
$$

$$
\sqrt{v} - \sqrt{\frac{2\Delta\psi\log 1/\delta}{n}} \le \sqrt{\widehat{v}(X) + \frac{2\Delta\psi\log 1/\delta}{n}}
$$

$$
\sqrt{v} \le \sqrt{\widehat{v}(X)} + 2\sqrt{\frac{2\Delta\psi\log 1/\delta}{n}},
$$

which proves the lemma for $b = -1$. On the other hand, from the first inequality in [Maurer](#page-23-12) [\[2006,](#page-23-12) Theorem 13],

$$
\mathbb{P}\left(\frac{n}{2\Delta \psi}\widehat{v}(X) - \mathbb{E}\left[\frac{n}{2\Delta \psi}\widehat{v}(X)\right] > t\right) \le \exp\left(-\frac{t^2}{4\mathbb{E}\left[\frac{n}{2\Delta \psi}\widehat{v}(X)\right] + 2t}\right),
$$

that can be rewritten as

$$
\mathbb{P}\left(\widehat{v}(X) - v > t\right) \le \exp\left(-\frac{nt^2}{8\Delta\psi(v+t/2)}\right).
$$

The positive solution t_+ for

$$
nt^2 = 8\Delta\psi(v + t/2)\log 1/\delta
$$

is readily given by

$$
t_{+} = \frac{2\Delta\psi\log 1/\delta}{n} + \frac{4}{n}\sqrt{\left(\frac{\Delta\psi}{2}\log 1/\delta\right)\left(\frac{\Delta\psi}{2}\log 1/\delta + nv\right)}.
$$

Thus, it holds with probability at least $1 - \delta$ that

$$
\widehat{v}(X) \leq v + t_+,
$$

and from sub-additivity of the square root, successively, with probability at least $1 - \delta$, we have

$$
\widehat{v}(X) \le v + \frac{4\Delta\psi\log 1/\delta}{n} + 2\sqrt{\frac{2v\Delta\psi\log 1/\delta}{n}},
$$

$$
\widehat{v}(X) \le \left(\sqrt{v} + \sqrt{\frac{2\Delta\psi\log 1/\delta}{n}}\right)^2 + \frac{2\Delta\psi\log 1/\delta}{n},
$$

$$
\sqrt{\widehat{v}(X)} \le \sqrt{\left(\sqrt{v} + \sqrt{\frac{2\Delta\psi\log 1/\delta}{n}}\right)^2 + \frac{2\Delta\psi\log 1/\delta}{n}},
$$

$$
\sqrt{\widehat{v}(X)} \le \sqrt{v} + 2\sqrt{\frac{2\Delta\psi\log 1/\delta}{n}},
$$

which finishes proving the lemma 5 for $b = 1$. \Box

5.5 Proof of Theorem [3.1](#page-8-1)

From Theorem [2.1,](#page-4-0) with probability at least $1 - \delta/2$, it holds that

$$
\|\widehat{\mu}_{\mathbb{P}}(X_1,\ldots,X_n)-\mu_{\mathbb{P}}\|_{\mathcal{H}_{\psi}}\leq \sqrt{2v\frac{\log 2/\delta}{n}}+\frac{4}{3}\overline{\psi}^{1/2}\frac{\log 2/\delta}{n}.\tag{18}
$$

Invoking Lemma [3.4](#page-7-4) for $b = -1$, at confidence $1 - \delta/2$ it holds that

$$
\sqrt{v} \le \sqrt{\widehat{v}(X)} + 2\sqrt{\frac{2\Delta\psi\log 2/\delta}{n}},\tag{19}
$$

thus by combining [\(18\)](#page-20-3) and [\(19\)](#page-20-4) and a union bound yields that with probability at least $1 - \delta$,

$$
\|\widehat{\mu}_{\mathbb{P}}(X_1,\ldots,X_n) - \mu_{\mathbb{P}}\|_{\mathcal{H}_{\psi}}\n\leq \left(\sqrt{\widehat{v}(X)} + 2\sqrt{\frac{2\Delta\psi\log 2/\delta}{n}}\right)\sqrt{\frac{2\log 2/\delta}{n}} + \frac{4}{3}\overline{\psi}^{1/2}\frac{\log 2/\delta}{n}\n= \sqrt{2\widehat{v}(X)\frac{\log 2/\delta}{n}} + 4\left(\Delta\psi^{1/2} + \frac{\overline{\psi}^{1/2}}{3}\right)\frac{\log 2/\delta}{n}\n\leq \sqrt{2\widehat{v}(X)\frac{\log 2/\delta}{n}} + \frac{16}{3}\Delta\psi^{1/2}\frac{\log 2/\delta}{n}.
$$
\n(20)

 \Box

5.6 Proof of Theorem [4.1](#page-10-1)

First, assume that H_0 is true, that is: there is a $\theta_0 \in \Theta$ such that $\mathbb{P} = \mathbb{P}_{\theta_0}$. Then,

$$
\mathbb{P}_{\mathbf{H}_0} \left(C \right) = \mathbb{P}_{\mathbf{H}_0} \left(T \geq \mathcal{B}(n, \alpha) \right)
$$
\n
$$
= \mathbb{P}_{\mathbf{H}_0} \left(\inf_{\theta \in \Theta} \left\| \widehat{\mu}_{\mathbb{P}}(X) - \mu_{\mathbb{P}_{\theta}} \right\|_{\mathcal{H}_k} \geq \mathcal{B}(n, \alpha) \right)
$$
\n
$$
\leq \mathbb{P}_{\mathbf{H}_0} \left(\left\| \widehat{\mu}_{\mathbb{P}}(X) - \mu_{\mathbb{P}_{\theta_0}} \right\|_{\mathcal{H}_k} \geq \mathcal{B}(\alpha) \right)
$$
\n
$$
\leq \alpha
$$

thanks to (*a*).

⁵In this proof [Maurer](#page-23-12) [\[2006\]](#page-23-12) is sufficient and we do need the stronger results of [Boucheron et al.](#page-22-9) [\[2009\]](#page-22-9).

Now, assume that H_0 is not true, that is $P \notin \{ \mathbb{P}_{\theta}, \theta \in \Theta \}$. Note that under the assumption that the model is closed, we have

$$
0<\Delta:=\inf_{\theta\in\Theta}\big\|\mu_{\mathbb{P}}-\mu_{\mathbb{P}_{\theta}}\big\|_{\mathcal{H}_k}.
$$

By the triangle inequality, for any *θ*,

$$
\|\mu_{\mathbb{P}_{\theta}} - \widehat{\mu}_{\mathbb{P}}(X)\|_{\mathcal{H}_k} \ge \|\mu_{\mathbb{P}_{\theta}} - \mu_{\mathbb{P}}\|_{\mathcal{H}_k} - \|\mu_{\mathbb{P}} - \widehat{\mu}_{\mathbb{P}}(X)\|_{\mathcal{H}_k}
$$

and thus, taking the infimum w.r.t *θ* on both sides,

$$
T \geq \Delta - \|\mu_{\mathbb{P}} - \widehat{\mu}_{\mathbb{P}}(X)\|_{\mathcal{H}_k}.
$$

We apply (c) with confidence level α_n and obtain

$$
\mathbb{P}_{\mathbf{H}_1}\left(\|\mu_{\mathbb{P}}-\widehat{\mu}_{\mathbb{P}}(X)\|_{\mathcal{H}_k}\geq\mathcal{B}(n,\alpha_n)\right)\leq\alpha_n
$$

and thus

$$
\mathbb{P}_{\mathbf{H}_1} \left(T \leq \Delta - \mathcal{B}(n, \alpha_n) \right) \leq \alpha_n.
$$

As $\mathcal{B}(n, \alpha_n) \to 0$ when $n \to \infty$ thanks to (b) , there is a *N* large enough such that, for any $n \ge N$, $\mathcal{B}(n, \alpha_n) \le$ ∆/2, and thus \overline{a}

$$
\mathbb{P}_{\mathbf{H}_1}\left(T\leq \frac{\Delta}{2}\right)\leq \alpha_n.
$$

Moreover, as we also have $\mathcal{B}(n,\alpha) \to 0$ when $n \to \infty$ thanks to (a), there is a N' large enough such that, for any $n \ge N'$, $\mathcal{B}(n, \alpha) \le \Delta/2$ and thus

$$
\mathbb{P}_{\mathbf{H}_1}(T \leq \mathcal{B}(n,\alpha)) \leq \alpha_n.
$$

Finally,

$$
\mathbb{P}_{\mathbf{H}_1}(\mathcal{C}) = 1 - \mathbb{P}_{\mathbf{H}_1}(T \leq \mathcal{B}(n, \alpha)) \geq 1 - \alpha_n
$$

as soon as $n \geq \max(N, N').$

5.7 Proof of Lemma [4.1](#page-12-2)

$$
v(\mathbb{P}) = \mathbb{E}_{X\sim\mathbb{P}} ||k(X,\cdot) - \mu_{\mathbb{P}}||_{\mathcal{H}_k}^2
$$

\n
$$
= \overline{\psi} - \int \int \psi(x'-x)d(\mathbb{P} \times \mathbb{P})(x,x')
$$

\n
$$
= \overline{\psi} - \int \int \psi(x'-x)d\mathbb{P}(x)d\mathbb{P}(x')
$$

\n
$$
= \overline{\psi} - \int \int \psi(x'-x)d((1-\xi)\mathbb{P}_{\theta_0}(x) + \xi \mathbb{H}(x))d((1-\xi)\mathbb{P}_{\theta_0}(x') + \xi \mathbb{H}(x'))
$$

\n
$$
= \overline{\psi} - (1-\xi)^2 \int \int \psi(x'-x)d\mathbb{P}_{\theta_0}(x)d\mathbb{P}_{\theta_0}(x')
$$

\n
$$
- 2\xi(1-\xi) \int \int \psi(x'-x)d\mathbb{P}_{\theta_0}(x)d\mathbb{H}(x')
$$

\n
$$
- \xi^2 \int \int \psi(x'-x)d\mathbb{H}(x)d\mathbb{H}(x')
$$

\n
$$
\leq \overline{\psi} - (1-\xi)^2(\overline{\psi}-v) - 2\xi(1-\xi)\underline{\psi} - \xi^2\underline{\psi}
$$

\n
$$
= \overline{\psi} - (1-2\xi+\xi^2)(\overline{\psi}-v) - \xi(2-\xi)\underline{\psi}
$$

\n
$$
= \overline{\psi} - (\overline{\psi}-v) + 2\xi(\overline{\psi}-v) - \xi^2(\overline{\psi}-v) - \xi(2-\xi)\underline{\psi}
$$

\n
$$
\leq (1-2\xi)v + 2\xi\Delta\psi + \xi^2\underline{\psi},
$$

\n
$$
= (1-2\xi)v + 2\xi\Delta\psi + \xi^2\underline{\psi},
$$

and the lemma follows by disjunction of cases. \Box

References

- P. Alquier, B.-E. Cherief-Abdellatif, A. Derumigny, and J.-D. Fermanian. Estimation of copulas via maxi- ´ mum mean discrepancy. Journal of the American Statistical Association (to appear), 2022.
- J.-Y. Audibert, R. Munos, and C. Szepesvári. Tuning bandit algorithms in stochastic environments. In *International conference on algorithmic learning theory*, pages 150–165. Springer, 2007.
- P. L. Bartlett and S. Mendelson. Rademacher and gaussian complexities: Risk bounds and structural results. *Journal of Machine Learning Research*, 3(Nov):463–482, 2002.
- D. Berend and A. Kontorovich. A sharp estimate of the binomial mean absolute deviation with applications. *Statistics & Probability Letters*, 83(4):1254–1259, 2013.
- A. Berlinet and C. Thomas-Agnan. *Reproducing kernel Hilbert spaces in probability and statistics*. Springer Science & Business Media, 2011.
- S. Boucheron, G. Lugosi, and P. Massart. On concentration of self-bounding functions. *Electronic Journal of Probability*, 14:1884–1899, 2009.
- F.-X. Briol, A. Barp, A. B. Duncan, and M. Girolami. Statistical inference for generative models with maximum mean discrepancy. *arXiv preprint arXiv:1906.05944*, 2019.
- B.-E. Cherief-Abdellatif and P. Alquier. Finite sample properties of parametric mmd estimation: robustness ´ to misspecification and dependence. *Bernoulli*, 28(1):181–213, 2022.
- K. Chwialkowski, H. Strathmann, and A. Gretton. A kernel test of goodness of fit. In M. F. Balcan and K. Q. Weinberger, editors, *Proceedings of The 33rd International Conference on Machine Learning*, volume 48 of *Proceedings of Machine Learning Research*, pages 2606–2615, New York, New York, USA, 20–22 Jun 2016. PMLR.
- D. Cohen, A. Kontorovich, and G. Wolfer. Learning discrete distributions with infinite support. In *Advances in Neural Information Processing Systems*, volume 33, pages 3942–3951, 2020.
- M. Cuturi. Positive definite kernels in machine learning. *arXiv preprint arXiv:0911.5367*, 2009.
- J. Diestel and J. Uhl. Vector measures, vol. 15 of mathematical surveys. *American Mathematical Society, Providence, RI, USA*, 1977.
- G. K. Dziugaite, D. M. Roy, and Z. Ghahramani. Training generative neural networks via maximum mean discrepancy optimization. *arXiv preprint arXiv:1505.03906*, 2015.
- M. G. Genton. Classes of kernels for machine learning: a statistics perspective. *Journal of machine learning research*, 2(Dec):299–312, 2001.
- A. Gretton, K. Fukumizu, Z. Harchaoui, and B. K. Sriperumbudur. A fast, consistent kernel two-sample test. *Advances in neural information processing systems*, 22, 2009.
- A. Gretton, K. M. Borgwardt, M. J. Rasch, B. Scholkopf, and A. Smola. A kernel two-sample test. ¨ *The Journal of Machine Learning Research*, 13(1):723–773, 2012.
- P. J. Hüber. Robust statistics. In *International encyclopedia of statistical science*, pages 1248–1251. Springer, 2011.
- W. Jitkrittum, W. Xu, Z. Szabó, K. Fukumizu, and A. Gretton. A linear-time kernel goodness-of-fit test. *Advances in Neural Information Processing Systems*, 30, 2017.
- M. Kearns, Y. Mansour, D. Ron, R. Rubinfeld, R. E. Schapire, and L. Sellie. On the learnability of discrete distributions. In *Proceedings of the twenty-sixth annual ACM symposium on Theory of computing*, pages 273–282, 1994.
- Y. Li, K. Swersky, and R. Zemel. Generative moment matching networks. In *International conference on machine learning*, pages 1718–1727. PMLR, 2015.
- D. Lopez-Paz, K. Muandet, B. Schölkopf, and I. Tolstikhin. Towards a learning theory of cause-effect inference. In *International Conference on Machine Learning*, pages 1452–1461. PMLR, 2015.
- A. Maurer. Concentration inequalities for functions of independent variables. *Random Structures & Algorithms*, 29(2):121–138, 2006.
- A. Maurer and M. Pontil. Empirical Bernstein bounds and sample variance penalization. *arXiv preprint arXiv:0907.3740*, 2009.
- C. McDiarmid et al. On the method of bounded differences. *Surveys in combinatorics*, 141(1):148–188, 1989.
- M. Mohri, A. Rostamizadeh, and A. Talwalkar. *Foundations of machine learning*. MIT press, 2018.
- I. F. Pinelis and A. I. Sakhanenko. Remarks on inequalities for the probabilities of large deviations. *Teoriya Veroyatnostei i ee Primeneniya*, 30(1):127–131, 1985.
- M. L. Stein. *Interpolation of spatial data: some theory for kriging*. Springer Science & Business Media, 1999.
- I. Tolstikhin, B. K. Sriperumbudur, and K. Muandet. Minimax estimation of kernel mean embeddings. *The Journal of Machine Learning Research*, 18(1):3002–3048, 2017.
- A. B. Tsybakov. *Introduction to nonparametric estimation*. Springer Science & Business Media, 2008.
- V. Yurinsky. *Sums and Gaussian Vectors. Lecture Notes in Mathematics*. Springer Berlin, Heidelberg, 1995.

A Additional proofs

A.1 Proof of Lemma [3.1](#page-6-2)

For some t_0 , we let $x^{(t_0)} = (x_1, \ldots, x_{t_0-1}, x_{t_0}', x_{t_0+1}, \ldots, x_n)$ where x_{t_0} was replaced with x_{t_0}' .

$$
\frac{1}{d} \operatorname{Tr} \hat{\Sigma}(x) - \frac{1}{d} \operatorname{Tr} \hat{\Sigma} \left(x^{(t_0)} \right) \n= \frac{1}{d} \sum_{i=1}^{d} \frac{1}{2n(n-1)} \left(\sum_{s=1}^{n} \sum_{t=1}^{n} \left(x_t^i - x_s^i \right)^2 - \sum_{s=1}^{n} \sum_{t=1}^{n} \left(x_t^{(t_0)i} - x_s^{(t_0)i} \right)^2 \right) \n\leq \frac{1}{d} \sum_{i=1}^{d} \left(\frac{1}{n-1} \sum_{s=1}^{n} \left(x_{t_0}^i - x_s^i \right)^2 \right) \leq 1,
$$
\n(21)

thus

$$
\frac{1}{d}\operatorname{Tr}\widehat{\Sigma}(x) - \inf_{x'_{t_0}\in\mathcal{X}}\frac{1}{d}\operatorname{Tr}\widehat{\Sigma}\left(x^{(t_0)}\right) \leq 1.
$$

Furthermore,

$$
\sum_{t_0=1}^n \left(\frac{1}{d} \operatorname{Tr} \widehat{\Sigma}(x) - \frac{1}{d} \operatorname{Tr} \widehat{\Sigma}\left(x^{(t_0)}\right)\right)^2 \stackrel{(i)}{\leq} \sum_{t_0=1}^n \left(\frac{1}{d} \sum_{i=1}^d \left(\frac{1}{n-1} \sum_{s=1}^n \left(x_{t_0}^i - x_s^i\right)^2\right)\right)^2
$$
\n
$$
\stackrel{(ii)}{\leq} \frac{1}{d} \sum_{i=1}^d \sum_{t_0=1}^n \left(\frac{1}{n-1} \sum_{s=1}^n \left(x_{t_0}^i - x_s^i\right)^2\right)^2
$$
\n
$$
\stackrel{(iii)}{\leq} \frac{1}{d} \sum_{i=1}^d \frac{n}{n-1} \widehat{v}^i(X)
$$
\n
$$
= \frac{n}{n-1} \frac{1}{d} \operatorname{Tr} \widehat{\Sigma}(X),
$$

where (*i*) follows from [\(21\)](#page-23-14), (*ii*) is Jensen's inequality, and (*iii*) stems from [Maurer and Pontil](#page-23-11) [\[2009,](#page-23-11) Corollary 9].

A.2 Proof of Lemma [A.1](#page-25-0)

$$
\hat{v}(x) - \hat{v}(x^{(t)})
$$
\n
$$
= \frac{1}{n-1} \sum_{r=1}^{n} \left(k(x_r, x_r) - \frac{1}{n} \sum_{s=1}^{n} k(x_r, x_s) \right)
$$
\n
$$
- \frac{1}{n-1} \sum_{r=1}^{n} \left(k(x_r^{(t)}, x_r^{(t)}) - \frac{1}{n} \sum_{s=1}^{n} k(x_r^{(t)}, x_s^{(t)}) \right)
$$
\n
$$
= \frac{1}{n-1} \sum_{r=1}^{n} \left(k(x_r, x_r) - k(x_r^{(t)}, x_r^{(t)}) \right)
$$
\n
$$
- \frac{1}{(n-1)n} \sum_{(r,s) \in [n]^2} \left(k(x_r, x_s) - k(x_r^{(t)}, x_s^{(t)}) \right)
$$
\n
$$
= \frac{1}{n-1} \left(k(x_t, x_t) - k(x_t', x_t') \right) - \frac{1}{(n-1)n} \sum_{(r,s) \in [n]^2, r=s} \left(k(x_r, x_s) - k(x_r^{(t)}, x_s^{(t)}) \right)
$$
\n
$$
- \frac{1}{(n-1)n} \sum_{(r,s) \in [n]^2, r \neq s} \left(k(x_r, x_s) - k(x_r^{(t)}, x_s^{(t)}) \right)
$$
\n
$$
= \frac{1}{n-1} \left(k(x_t, x_t) - k(x_t', x_t') \right) - \frac{1}{(n-1)n} \sum_{r=1}^{n} \left(k(x_r, x_r) - k(x_r^{(t)}, x_r^{(t)}) \right)
$$
\n
$$
- \frac{1}{(n-1)n} \sum_{(r,s) \in [n]^2, r \neq s, r=t} \left(k(x_r, x_s) - k(x_r^{(t)}, x_s^{(t)}) \right)
$$
\n
$$
= \frac{1}{n} \left(k(x_t, x_t) - k(x_t', x_t') \right)
$$
\n
$$
= \frac{1}{n} \left(k(x_t, x_t) - k(x_t', x_t') \right)
$$
\n
$$
= \frac{1}{n} \left(k(x_t, x_t) - k(x_t', x_t') \right)
$$
\n
$$
= \frac{1}{n-1} \left(k(x_t, x_s) - k(x_t', x_s) \right
$$

 \Box

A.3 Proof of Lemma [3.5](#page-8-2)

We will rely on the following property for general kernels.

Lemma A.1. *Let* $x = (x_1, \ldots, x_{t_0}, \ldots, x_n) \in \mathcal{X}^n$ and $x'_{t_0} \in \mathcal{X}$. Writing

$$
x^{(t)} = (x_1, \ldots, x_{t_0-1}, x'_{t_0}, x_{t_0+1}, \ldots, x_n),
$$

it holds that

$$
\widehat{v}(x) - \widehat{v}\left(x^{(t_0)}\right) = \frac{1}{n} \left(k \left(x_{t_0}, x_{t_0}\right) - k \left(x'_{t_0}, x'_{t_0}\right) \right) \n+ \frac{2}{(n-1)n} \sum_{t \in [n], t \neq t_0} \left(k \left(x'_{t_0}, x_t\right) - k \left(x_{t_0}, x_t\right) \right).
$$

From Lemma [A.1,](#page-25-0) if follows that

$$
\widehat{v}(x) - \inf_{x'_{t_0} \in \mathcal{X}} \widehat{v}\left(x^{(t_0)}\right) \stackrel{(+)}{\leq} \frac{1}{n} \left(k\left(x_{t_0}, x_{t_0}\right) - \frac{\text{diag }k}{n}\right) + \frac{2}{(n-1)n} \sum_{t \in [n], t \neq t_0} \left(\overline{k} - k\left(x_{t_0}, x_t\right)\right) + \frac{2}{(n-1)n} \sum_{t \in [n], t \neq t_0} \left(\overline{k} - k\left(x_{t_0}, x_t\right)\right) + \frac{2}{(n-1)n} \sum_{t \in [n], t \neq t_0} \left(\overline{k} - k\left(x_{t_0}, x_t\right)\right)
$$

and the bounded differences property follows directly from (\ddagger) . Furthermore, as a result of (\dagger) ,

$$
\sum_{t_{0}=1}^{n} \left(\hat{v}(x) - \inf_{x'_{t_{0}} \in \mathcal{X}} \hat{v}(x^{(t_{0})}) \right)^{2}
$$
\n
$$
\leq \sum_{t_{0}=1}^{n} \left(\frac{1}{n} \left(k(x_{t_{0}}, x_{t_{0}}) - \frac{\text{diag }k}{n} \right) + \frac{2}{(n-1)n} \sum_{t \in [n], t \neq t_{0}} \left(\overline{k} - k(x_{t_{0}}, x_{t}) \right) \right)^{2}
$$
\n
$$
= \frac{1}{n^{2}} \sum_{t_{0}=1}^{n} \left(\frac{1}{(n-1)} \sum_{t \in [n], t \neq t_{0}} k(x_{t_{0}}, x_{t_{0}}) - \frac{\text{diag }k}{n} + 2\overline{k} - 2k(x_{t_{0}}, x_{t}) \right)^{2}
$$
\n
$$
\leq \frac{1}{n^{2}} \sum_{t_{0}=1}^{n} \frac{1}{(n-1)} \sum_{t \in [n], t \neq t_{0}} \left(k(x_{t_{0}}, x_{t_{0}}) - \frac{\text{diag }k}{n} + 2\overline{k} - 2k(x_{t_{0}}, x_{t}) \right)^{2}
$$
\n
$$
\leq \frac{\Delta \text{diag }k + 2\Delta k}{n^{2}} \sum_{t_{0}=1}^{n} \frac{1}{(n-1)} \sum_{t \in [n], t \neq t_{0}} \left(k(x_{t_{0}}, x_{t_{0}}) - \frac{\text{diag }k}{n} + 2\overline{k} - 2k(x_{t_{0}}, x_{t}) \right)
$$
\n
$$
\leq \frac{\Delta \text{diag }k + 2\Delta k}{n^{2}} \sum_{t_{0}=1}^{n} \frac{1}{(n-1)} \sum_{t \in [n], t \neq t_{0}} \left(2k(x_{t_{0}}, x_{t_{0}}) - 2\frac{\text{diag }k}{n} + 2\overline{k} - 2k(x_{t_{0}}, x_{t}) \right)
$$
\n
$$
= 2 \frac{\Delta \text{diag }k + 2\Delta k}{n^{2}} \sum_{t_{0}=1}^{n} \frac{1}{(n-1)} \sum_{t \in [n], t \neq t_{0}} \left(k(x_{t_{0}}, x_{t_{0}}) - k(x
$$

where for (\star) we relied on $\overline{k} = \overline{\mathrm{diag}\,k}$ for a positive definite kernel. \Box

A.4 Proof of Lemma [3.6](#page-8-3)

Lemma [3.5](#page-8-2) established that the function

$$
\mathcal{X}^n \to \mathbb{R}, x \mapsto \frac{n}{\Delta \operatorname{diag} k + 2\Delta k} \widehat{v}(x),
$$

is weakly self-bounding and has the bounded differences property. From an application of [Boucheron](#page-22-9) [et al.](#page-22-9) [\[2009,](#page-22-9) Theorem 1],

$$
\mathbb{P}(\hat{v} - v > t) \le \exp\left(-\frac{nt^2}{4(\Delta \operatorname{diag} k + 2\Delta k)(v + \Delta \operatorname{diag} k + t/2)}\right),
$$

$$
\mathbb{P}(v - \hat{v} > t) \le \exp\left(-\frac{nt^2}{4(\Delta \operatorname{diag} k + 2\Delta k)(v + \Delta \operatorname{diag} k)}\right).
$$

The claim follows from rewriting

 $\hat{v} - v = [\hat{v} + \Delta \text{diag } k] - [v + \Delta \text{diag } k],$

and solving quadratic inequalities as in Lemma $3.4.$ \Box

A.5 Proof of Theorem [3.2](#page-8-4)

From Theorem [2.1,](#page-4-0) and since ∆ diag *k* > 0, with probability at least 1 − *δ*/2, it holds that

$$
\|\widehat{\mu}_{\mathbb{P}}(X_1,\ldots,X_n)-\mu_{\mathbb{P}}\|_{\mathcal{H}_k}\leq \sqrt{2(v+\Delta \operatorname{diag} k)\frac{\log 2/\delta}{n}}+\frac{4}{3}\overline{k}^{1/2}\frac{\log 2/\delta}{n}.\tag{22}
$$

By Lemma [3.6](#page-8-3) for $b = -1$, at confidence $1 - \delta/2$ it holds that

$$
\sqrt{v + \Delta \text{diag} k} \le \sqrt{\hat{v} + \Delta \text{diag} k} + 2\sqrt{\frac{(\Delta \text{diag} k + 2\Delta k) \log 2/\delta}{n}},\tag{23}
$$

thus by combining [\(22\)](#page-27-1) and [\(23\)](#page-27-2) and a union bound yields that with probability at least $1 - \delta$,

$$
\|\widehat{\mu}_{\mathbb{P}} - \mu_{\mathbb{P}}\|_{\mathcal{H}_k} \leq \left(\sqrt{\widehat{v} + \Delta \text{diag } k} + 2\sqrt{\frac{(\Delta \text{diag } k + 2\Delta k) \log 2/\delta}{n}}\right) \sqrt{\frac{2 \log 2/\delta}{n}} + \frac{4}{3} \bar{k}^{1/2} \frac{\log 2/\delta}{n}
$$

$$
= \sqrt{2(\widehat{v} + \Delta \text{diag } k) \frac{\log 2/\delta}{n}} + \left(2\sqrt{2}\sqrt{\Delta \text{diag } k + 2\Delta k} + \frac{4}{3} \bar{k}^{1/2}\right) \frac{\log 2/\delta}{n}
$$

$$
\leq \sqrt{2(\widehat{v} + \Delta \text{diag } k) \frac{\log 2/\delta}{n}} + \left(\frac{16}{3}\sqrt{\Delta k} + 2\sqrt{2}\sqrt{\Delta \text{diag } k}\right) \frac{\log 2/\delta}{n},\tag{24}
$$

where the last inequality is by sub-additivity of the square root. \Box