# Stability for the Helmholtz equation in deterministic and random periodic structures

Gang Bao<sup>1</sup>, Yiwen  $\text{Lin}^2$  and Xiang  $\text{Xu}^1$ 

<sup>1</sup>School of Mathematical Sciences, Zhejiang University, Hangzhou, 310027, China.

<sup>2</sup>School of Mathematical Sciences and Institute of Natural Sciences, Shanghai Jiao Tong University, Shanghai, 200240, China.

#### Abstract

Stability results for the Helmholtz equations in both deterministic and random periodic structures are proved in this paper. Under the assumption of excluding resonances, by a variational method and Fourier analysis in the energy space, the stability estimate for the Helmholtz equation in a deterministic periodic structure is established. For the stochastic case, by introducing a variable transform, the variational formulation of the scattering problem in a random domain is reduced to that in a definite domain with random medium. Combining the stability result for the deteministic case with regularity and stochastic regularity of the scattering surface, Pettis measurability theorem and Bochner's Theorem further yield the stability result for the scattering problem by random periodic structures. Both stability estimates are explicit with respect to the wavenumber.

Keywords: Helmholtz equation, periodic structure, high wavenumber, stability, estimate explicit on wavenumber

## 1 Introduction

This paper is concerned with scattering of time-harmonic electromagnetic plane waves by periodic structures, which is known as gratings in optics. The goal is to investigate stability properties of the Helmholtz equation in both deterministic and random periodic structures.

For the scattering by a periodic structure, considerable progress has been made mathematically in literatures. Bao, Dobson and Cox  $[4]$  reduced the scattering problem into a bounded domain problem by introducing a transparent boundary condition and proved that there exists a unique solution at all but a sequence of countable frequencies. Lord and Mulholland [\[20\]](#page-26-0) used the variational formula to derive a priori estimate for periodic structures, which can be viewed as an extension of [\[2\]](#page-24-1). More results on the Helmholtz and Maxwell equations in periodic structures can be found in [\[3](#page-24-2), [5](#page-24-3), [25\]](#page-26-1).

Recently, there is an increasing interest in the study of wavenumber-explicit bounds for scattering problems. Chandler-Wilde and Monk [\[10\]](#page-25-0) obtained a priori bounds explicit with the wavenumber for the scattering problem by unbounded rough surfaces. Hetmaniuk [\[16\]](#page-25-1) established stability estimates for the Helmholtz equation with mixed boundary conditions. Esterhazy and Melenk [\[13\]](#page-25-2) further established an estimate for bounded Lipschitz domains with a Robin boundary condition. The stability for the scattering problem by a large rectangular cavity was obtained by Bao, Yun and Zhou [\[8](#page-25-3)] in transverse electric polarization and was improved and extended to transverse magnetic polarization in [\[7\]](#page-25-4). Du, Li and Sun [\[12\]](#page-25-5) presented a numerical study of the stability estimate of the scattering by a rectangular cavity. Wavenumber-explicit stability on the scattering problem by an obstacle in homogeneous media case [\[28\]](#page-26-2) and heterogeneous media case [\[15](#page-25-6)] were obtained under the assumption that the scatterer  $D$  is a star-shaped domain satisfying nontrapping conditions. However, for periodic structures, little is known about stability analysis with explicit dependence on the wavenumber, which will be derived in this paper. As noted in [\[7,](#page-25-4) [8,](#page-25-3) [10\]](#page-25-0), both the geometry and the type of boundary condition strongly affect the wavenumber-dependent stability. The transparent boundary conditions (TBC) of periodic structures are different from those of open cavities and unbounded rough surfaces, which leads to additional difficulties on resonances. To overcome the difficulties, a uniform distance  $\varepsilon$  is introduced in this paper and a k-explicit stability for periodic structures is established by a combination of the variational method and Fourier analysis.

We also analyze the stability for scattering by random periodic structures in this paper. Progress has been made recently in the exterior scattering problem by an obstacle with randomness[\[17](#page-25-7), [28\]](#page-26-2) and scattering problems in random media [\[15](#page-25-6), [24\]](#page-26-3). However, no stability result is available for scattering by random periodic structures, which has many applications in diffractive optics [\[26\]](#page-26-4). Numerical methods for the scattering by random periodic structures have been developed recently in  $[6, 14]$  $[6, 14]$ . In this work, we focus on the stability for gratings with uncertainty. One difficulty is the lack of compactness. In fact, since both the deterministic and the stochastic Helmholtz equations are not coercive and the necessary compactness results are not valid any more in Bochner spaces, Fredholm theory cannot be used to the stochastic Helmholtz equation to compensate for the lack of coercivity as for the deterministic Helmholtz equation. Consequently, it is difficult to obtain the well-posedness for the random case directly as that for the deterministics case. In this work, we employ

Pettis measurability theorem and Bochner's theorem as in [\[24](#page-26-3)] to obtain the stability estimate explicit on the wavenumber for the random case based on the stability result for the deterministic case. However, the randomness of the integral domain for the scattering by random periodic structures prevents a direct application of the general framework in [\[24](#page-26-3)], which leads to the other difficulty. To overcome this difficulty, a variable transform is introduced here so that the transformed random variational form is defined on a deterministic domain with random coefficients. Similar transformation idea has also been used for other types of random differential equations [\[29\]](#page-26-5). Therefore, for the stochastic case, by integrating the deterministic result over the probability space and introducing a transform to change the stochastic integral area into a definite one, regularity and stochastic regularity of the scattering surface, Pettis measurability theorem and Bochner's theorem yield the well-posedness and a stability estimate of our model problem.

The rest of the paper is as follows. In Section [2,](#page-2-0) the model problem is introduced, the variational formulation is described, and the results (i.e., Theorems [2.1-](#page-5-0)[2.2\)](#page-6-0) are presented. Theorem [2.1](#page-5-0) concerns the stability for the Helmholtz equation in a deterministic periodic structure. Theorem [2.2](#page-6-0) gives the stability estimate for large wavenumber for the Helmholtz equation in a random periodic structure. The following sections are devoted to the proofs of the two results. Section [3](#page-6-1) and Section [4](#page-18-0) give the detailed proofs of Theorem [2.1](#page-5-0) and Theorem [2.2](#page-6-0) respectively, followed by conclusion given in Section [5.](#page-24-4)

## <span id="page-2-0"></span>2 Main results

#### 2.1 Model problem

Consider a plane wave incident on a random periodic structure

$$
S := \{ x \in \mathbb{R}^2 : x_2 = f(\omega; x_1), \omega \in \Omega, x_1 \in [0, \Lambda] \},
$$

which is characterized by the wavenumber  $k$  ruled on a perfect conductor. Here,  $\omega \in \Omega$  denotes the random sample in a complete probability space  $(\Omega, \mathcal{F}, \mu)$ ,  $x = (x_1, x_2) \in \mathbb{R}^2$  are the spatial variables, and the random surface  $f : \Omega \times \mathcal{X} \to$ R is a stationary Gaussian process, each of which is a Lipschitz function, i.e.  $f \in L^2(\Omega; Lip)$ , where  $Lip$  is the space of all Lipschitz functions. The medium and material are assumed to be invariant in the  $x_3$  direction and  $\Lambda$ -periodic in the  $x_1$  direction. There are two fundamental polarizations for the electromagnetic fields: transverse-electric (TE) and transverse-magnetic (TM) polarization. Here, we consider the random periodic perfectly conducting grating problem for TE polarization.

As shown in Figure [1,](#page-3-0) the grating is illuminated from above by a timeharmonic plane wave  $u^i = e^{i\alpha x_1 - i\beta x_2}$ , where  $\alpha = k \sin \theta$ ,  $\beta = k \cos \theta$ ,  $\theta \in$  $(-\pi/2, \pi/2)$  is the incident angle with respect to the positive  $x_2$ -axis. Denote

$$
D^{+} = \{x \in \mathbb{R}^{2} : x_{2} > f(\omega; x_{1}), \ \omega \in \Omega, x_{1} \in [0, \Lambda]\}.
$$



<span id="page-3-0"></span>Fig. 1 Problem geometry

Since the medium below  $S$  is perfectly electric conducting, the scattering problem for TE polarization can be modeled by the following two-dimensional Helmholtz equation with the homogeneous boundary condition:

<span id="page-3-1"></span>
$$
\begin{cases} \Delta u(\omega; \cdot) + k^2 u(\omega; \cdot) = 0 & \text{in } \Omega \times D^+, \\ u(\omega; \cdot) = 0 & \text{on } \Omega \times S, \end{cases}
$$
 (1)

where  $u(\omega; \cdot)$  is the total field.

#### <span id="page-3-3"></span>2.2 Variational form

In order to obtain a stability estimate, an important step is to reduce the infinite scattering problem into a bounded domain problem by introducing a transparent boundary condition [\[1\]](#page-24-5).

Since the total field can be decomposed into  $u(\omega; \cdot) = u^i + u^s(\omega; \cdot)$  where the incident wave satisfies

$$
\Delta u^i + k^2 u^i = 0 \quad \text{in } \Omega \times D^+,
$$

the scattered field  $u^s(\omega; \cdot)$  satisfies

$$
\begin{cases} \Delta u^s(\omega; \cdot) + k^2 u^s(\omega; \cdot) = 0 & \text{in } \Omega \times D^+, \\ u^s(\omega; \cdot) + u^i = 0 & \text{on } \Omega \times S. \end{cases}
$$
 (2)

Denote by  $\Gamma = \{ (x_1, x_2) \in \mathbb{R}^2 : x_2 = b, x_1 \in [0, \Lambda] \}$  with  $b >$  $\max_{\omega \in \Omega, x_1 \in (0,\Lambda)} f(\omega; x_1)$ . The scattered field  $u^s$  above  $\Gamma$  admits the Rayleigh expansion

$$
u^{s}(\omega; x_{1}, x_{2}) = \sum_{n \in Z} A_{n} e^{i(\alpha_{n} x_{1} + \beta_{n} x_{2})}, \qquad (3)
$$

where

$$
\alpha_n = \alpha + \frac{2\pi n}{\Lambda}, \beta_n^2 = k^2 - \alpha_n^2 \text{ with } \Im \beta_n \ge 0.
$$
 (4)

<span id="page-3-2"></span>**Assumption 2.1** (Exclusion of resonances). Assume there exists  $\varepsilon > 0$ , such that  $|k - \alpha_n| \geq \varepsilon$ ,  $\forall n \in \mathbb{Z}$ .

**Remark 2.1** A constant  $\varepsilon$  is introduced here to exclude all possible resonances ( $\kappa =$  $\alpha_n$ ) uniformly and thus a stability explicit with the wavenumber k is obtained by a combination of the variational method and Fourier analysis.

Define a boundary operator  $T: H^{\frac{1}{2}}(\Gamma) \to H^{-\frac{1}{2}}(\Gamma)$  by

$$
(Tv)(x_1) = \sum_{n \in \mathbb{Z}} i\beta_n v_n e^{i\alpha_n x_1},
$$

where  $v_n(\omega) = \frac{1}{\Lambda} \int_0^{\Lambda} v(\omega; x_1, b) e^{-i\alpha_n x_1} dx_1$ , which yields a transparent boundary condition on Γ:

$$
\partial_{\nu}u = Tu + g(x_1) \quad \text{on } \Gamma,
$$

and  $g(x_1) = \partial_\nu u^i - Tu^i = -2i\beta e^{i\alpha x_1 - i\beta b}$ .

Denote by  $D = \{x \in \mathbb{R}^2 : f(\omega; x_1) < x_2 < b, \ \omega \in \Omega, x_1 \in [0, \Lambda] \}.$ The scattering problem [\(1\)](#page-3-1) in an unbounded domain may be reduced to the following problem in the bounded domain:

<span id="page-4-2"></span>
$$
\begin{cases}\n\Delta u(\omega; \cdot) + k^2 u(\omega; \cdot) = 0 & \text{in } \Omega \times D, \\
u(\omega; \cdot) = 0 & \text{on } \Omega \times S, \\
\partial_{\nu} u(\omega; \cdot) = Tu(\omega; \cdot) + g(x) & \text{on } \Omega \times \Gamma.\n\end{cases}
$$
\n(5)

Motivated by uniqueness, we seek for quasi-periodic solutions of  $u(\omega; \cdot)$ , that is,  $u(\omega; x_1 + \Lambda, x_2) = e^{i\alpha\Lambda}u(\omega; x_1, x_2)$ . Let  $\tau$  denote a trace operator from D or S to  $\Gamma$ . Define the function spaces inside D by

$$
H1(D) = \{v \in L2(D) : \nabla v \in L2(D)2\},
$$
  
\n
$$
H1S(D) = \{v \in H1(D) : \tau v = 0 \text{ on } S\},
$$
  
\n
$$
H1S,qp(D) = \{v \in H1S(D) : v(0, y) = v(\Lambda, y)e^{-i\alpha\Lambda}\}.
$$

Let  $c : \Omega \to \mathcal{C}$  be defined by

<span id="page-4-1"></span>
$$
c(\omega) = f(\omega),\tag{6}
$$

where  $\mathcal{C} := Lip$  is the space of all Lipschitz functions.

Variational form. Define the sesquilinear forms A on  $L^2(\Omega; H^1_{S,qp}(D)) \times$  $L^2(\Omega; H^1_{S,qp}(D))$  and  $\tilde{a}_{c(\omega)}$  on  $H^1_{S,qp}(D) \times H^1_{S,qp}(D)$  by

<span id="page-4-3"></span>
$$
\mathcal{A}(u,v) := \int_{\Omega} \tilde{a}_{c(\omega)}(u,v) \, d\mathbb{P}(\omega),\tag{7}
$$

and

<span id="page-4-0"></span>
$$
\tilde{a}_{c(\omega)}(u,v) := \int_D \nabla u \cdot \nabla \bar{v} - k^2 \int_D u \bar{v} - \int_{\Gamma} T \tau u \overline{\tau v}.
$$
\n(8)

Define the antilinear functionals  $\mathcal G$  on  $L^2(\Omega; H^1_{S,qp}(D))$  and  $\tilde{G}_{c(\omega)}$  on  $H^1_{S,qp}(D)$ by

<span id="page-5-5"></span>
$$
\mathcal{G}(v) := \int_{\Omega} \tilde{G}_{c(\omega)}(v) d\mathbb{P}(\omega), \tag{9}
$$

and

<span id="page-5-3"></span>
$$
\tilde{G}_{c(\omega)}(v) := \int_{\Gamma} g \overline{\tau v}.
$$
\n(10)

Consider the following variational form of the Helmholtz equation in a random periodic structure:

<span id="page-5-2"></span>**Problem 1** Find  $u \in L^2(\Omega; H^1_{S,qp}(D))$  such that

<span id="page-5-4"></span>
$$
\mathcal{A}(u,v) = \mathcal{G}(v), \ \forall v \in L^{2}(\Omega; H^{1}_{S,qp}(D)).
$$
\n(11)

### 2.3 Stability results

We first present a stability result explicit with respect to the wavenumber  $k$ for the scattering problem by a deterministic periodic structure is given in Theorem [2.1.](#page-5-0)

<span id="page-5-0"></span>**Theorem 2.1** (Stability for the deterministic case). Fix one sample  $\omega_0$ . Let  $\tilde{u} =$  $u(\omega_0) \in H^1_{S,qp}(D)$  be the solution of the deterministic scattering problem. Under Assumption [2.1,](#page-3-2) there exists a constant C independent of  $b > 1$  and large k, such that

$$
\|\nabla \tilde{u}\|_{L^{2}(D)} + k\|\tilde{u}\|_{L^{2}(D)} \leq C \max\left\{\frac{b^{2}k^{2}}{\sqrt{\varepsilon}}, b^{3}k^{\frac{5}{2}}\right\} \|g\|_{L^{2}(\Gamma)}.
$$
 (12)

**Remark 2.2** For the estimate,  $\varepsilon$  is independent of the wavenumber k and the measured height b. Note that for small  $\varepsilon$ , wavenumber is close to resonances and then the corresponding bound is very large. If  $\varepsilon \geq \frac{1}{b^2k}$ , then the estimate becomes

$$
\|\nabla \tilde{u}\|_{L^{2}(D)} + k\|\tilde{u}\|_{L^{2}(D)} \leq Cb^{3}k^{\frac{5}{2}}\|g\|_{L^{2}(\Gamma)}.
$$
\n(13)

Next, the first stability result in a random periodic structure is shown in Theorem [2.2](#page-6-0) under Assumption [2.1](#page-3-2) and Assumption [2.2.](#page-5-1)

<span id="page-5-1"></span>**Assumption 2.2** (Regularity and stochastic regularity of  $f$ ). Assume the random structure  $f \in L^2(\Omega; Lip)$ .

Remark 2.3 (Random surfaces satisfying Assumption [2.2\)](#page-5-1). Motivated by the  $uncertainty$  quantification such as Karhunen-Loève expansion and other similar expansions, it makes sense to consider f as series expansions around the known deterministic surface  $f_0$  satisfying the Lipschitz condition.

<span id="page-6-0"></span>Theorem 2.2 (Stability for the random case). Under Assumption [2.1](#page-3-2) and Assump-tion [2.2,](#page-5-1) there exists a unique solution  $u \in L^2(\Omega; H^1_{S,qp}(D))$  to Problem [1.](#page-5-2) Moreover, there exists a constant C independent of  $b > 1$  and large k, such that

<span id="page-6-4"></span>
$$
\|\nabla u\|_{L^{2}(\Omega; L^{2}(D))} + k\|u\|_{L^{2}(\Omega; L^{2}(D))} \leq C \max\left\{\frac{b^{2}k^{2}}{\sqrt{\varepsilon}}, b^{3}k^{\frac{5}{2}}\right\} \|g\|_{L^{2}(\Omega; L^{2}(\Gamma))}. (14)
$$

## <span id="page-6-1"></span>3 Stability for the deterministic case

This section is devoted to the proof of Theorem [2.1.](#page-5-0) Our approach based on the variational method and Fourier analysis consists of two steps: the first step is to estimate the norms of u in the domain  $D$  by the norms of u on the boundary Γ and the second step is to estimate the norms of u on Γ by the norm of g.

Although our approach is related to [\[7](#page-25-4), [8](#page-25-3)] for the scattering by a large rectangular cavity, additional difficulties arise: (1) Because of the different model geometry, quasiperiodic solutions have to be considered for our model here. (2) Instead of the Dirichlet or Neumann boundary condition for the cavity problem, the (nonlocal) transparent boundary conditions must be dealt with. (3) Unlike the cavity problem, our model problem may not have unique solutions due to the resonances. Therefore, new techniques must be developed to prove the stability result.

Before proceeding to the detailed proof, some useful notations and preliminary results are introduced first.

#### 3.1 Notations and preliminary results

Let  $\tilde{u}(x_1, x_2) = u(\omega_0; x_1, x_2)$  be the solution to the deterministic problem for a given sample  $\omega_0$ . Then the total field  $\tilde{u}$  satisfies

<span id="page-6-3"></span>
$$
\begin{cases}\n\Delta \tilde{u} + k^2 \tilde{u} = 0 & \text{in } D, \\
\tilde{u} = 0 & \text{on } S, \\
\partial_{\nu} \tilde{u} = T \tilde{u} + g & \text{on } \Gamma.\n\end{cases}
$$
\n(15)

Let C denote positive constants which are independent of large k and  $b > 1$ . Denote by  $\alpha \simeq \beta$  if there exists a positive constant C independent of large k and  $b > 1$  such that  $\frac{1}{C}\alpha \leq \beta \leq C\alpha$ . Without loss of generality, assume the period  $\Lambda = 2\pi$  and denote  $\tilde{u}$  as u for simplicity.

The scattered field  $u$  can be written in the single-layer potential representation:

<span id="page-6-2"></span>
$$
u(x_1, x_2) = \int_0^{\Lambda} \phi(s) G(x_1, x_2; s, 0) \, ds,\tag{16}
$$

where  $\phi \in L^2(0,\Lambda)$  is an unknown periodic density function and G is a quasiperiodic Green function given explicitly as

$$
G(x_1, x_2; s, t) = \frac{i}{2\pi} \sum_{n \in \mathbb{Z}} \frac{1}{\beta_n} e^{i\alpha_n(x_1 - s) + i\beta_n |x_2 - t|}, \quad (x_1, x_2) \neq (s, t).
$$

Without loss of generality, it is assumed that the normal incidence is with  $\sqrt{k^2 - n^2}$ , and thus 2π-periodicity, i.e.  $\theta = 0, \alpha = 0, \beta = k, \Lambda = 2\pi$ . Therefore  $\alpha_n = n$  and  $\beta_n =$ 

<span id="page-7-0"></span>
$$
G(x_1, x_2; s, t) = \frac{i}{2\pi} \sum_{n \in \mathbb{Z}} \frac{1}{\sqrt{k^2 - n^2}} e^{in(x - s) + i\sqrt{k^2 - n^2}|y - t|} \text{ and } \phi(s) = \sum_{n \in \mathbb{Z}} \phi_n e^{ins}.
$$
\n(17)

Substituting  $(17)$  into  $(16)$  yields

$$
u^{s}(x_{1}, x_{2}) = \sum_{n \in \mathbb{Z}} \psi_{n} e^{inx_{1} + i\sqrt{k^{2} - n^{2}}x_{2}}.
$$

Since  $u = u^{inc} + u^s$ , the solution u can be expressed as

$$
u(x_1, x_2) = \sum_{n=0}^{\infty} a_n \sin\left(\sqrt{k^2 - n^2} x_2\right) \sin(nx_1) + \sum_{n=0}^{\infty} b_n \cos\left(\sqrt{k^2 - n^2} x_2\right) \cos(nx_1) + i \left[ \sum_{n=0}^{\infty} c_n \cos\left(\sqrt{k^2 - n^2} x_2\right) \sin(nx_1) + \sum_{n=0}^{\infty} d_n \sin\left(\sqrt{k^2 - n^2} x_2\right) \cos(nx_1) \right]
$$
  

$$
\triangleq \mu + i\nu.
$$

where  $a_n = -\psi_n + \psi_{-n}, b_n = \psi_n + \psi_{-n}, n \in \mathbb{N}_+, a_0 = -\psi_0 + 1, b_0 = \psi_0 + 1;$  $c_n = -a_n, d_n = b_n, n \in \mathbb{N}_+, c_0 = \psi_0 - 1, d_0 = \psi_0 + 1.$  For the case of non-normal cases, we only need to replace *n* with  $\alpha_n$  and  $\sqrt{k^2 - n^2}$  with  $\sqrt{k^2 - \alpha_n^2}$ . Since  $\mu$  and  $\nu$  can be studied similarly, only detailed study on  $\mu$  is provided here. Also, for convenience, in the following proof, the notation  $\mu$  is denoted as u.

Let  $u_n$ ,  $(\partial_\nu u)_n$ ,  $(T u)_n$  be the sine coefficients, and  $v_n$ ,  $(\partial_\nu v)_n$ ,  $(T v)_n$  be the cosine coefficients of the expansions of u,  $\partial_{\nu}u$ , Tu, on Γ, respectively. That is,

<span id="page-7-1"></span>
$$
u_n := a_n \sin\left(\sqrt{k^2 - n^2}b\right), v_n := b_n \cos\left(\sqrt{k^2 - n^2}b\right),\tag{18}
$$

$$
\left(\partial_{\nu}u\right)_n := a_n\sqrt{k^2 - n^2}\cos\left(\sqrt{k^2 - n^2}b\right),\tag{19}
$$

<span id="page-7-2"></span>
$$
\left(\partial_{\nu}v\right)_n := -b_n\sqrt{k^2 - n^2}\sin\left(\sqrt{k^2 - n^2}b\right). \tag{20}
$$

Hence

$$
u|_{\Gamma}(x_1) = \sum_{n=0}^{\infty} u_n \sin(nx_1) + \sum_{n=0}^{\infty} v_n \cos(nx_1),
$$

$$
\partial_{\nu} u|_{\Gamma}(x_1) = \sum_{n=0}^{\infty} (\partial_{\nu} u)_n \sin(nx_1) + \sum_{n=0}^{\infty} (\partial_{\nu} v)_n \cos(nx_1),
$$

and

$$
Tu|_{\Gamma}(x_1) = \sum_{n=0}^{\infty} (Tu)_n \sin(nx_1) + \sum_{n=0}^{\infty} (Tv)_n \cos(nx_1).
$$

<span id="page-8-1"></span>**Definition 3.1** For 
$$
u = \sum_{n=0}^{\infty} u_n \sin(nx_1) + \sum_{n=0}^{\infty} v_n \cos(nx_1)
$$
, define  

$$
||u||_A = \sqrt{\sum_{n=0}^{\infty} (|u_n|^2 + |v_n|^2) \frac{1}{\sqrt{|k^2 - n^2|} + \frac{1}{b}}},
$$

and

$$
||u||_B = \sqrt{\sum_{n=0}^{\infty} (|u_n|^2 + |v_n|^2) \left(\sqrt{|k^2 - n^2|} + \frac{1}{b}\right)}.
$$

Here, the norms A and B are formerly dual to each other and behave like  $H^{-1/2}$ ,  $H^{1/2}$ , respectively. Moreover, a simple calculation gives that

$$
\int_{\Gamma} u_1(x_1) \overline{u_2(x_1)} dx_1 \le \pi \|u_1\|_{A} \|u_2\|_{B}.
$$
 (21)

**Definition 3.2** Define the lower frequency set **L** and the higher frequency set **H** by  $\mathbf{L} = \{n \in \mathbb{N} \mid n < k\} \text{ and } \mathbf{H} = \{n \in \mathbb{N} \mid n > k\}.$ 

Then, u,  $\partial_{\nu}u$ ,  $T(u)$  on  $\Gamma$  defined above can be separated into the lower frequency part and the higher frequency part:

$$
u(x_1) = \mathcal{V}_\mathbf{L} + \mathcal{V}_\mathbf{H}, \quad \partial_\nu u(x_1) = \mathcal{P}_\mathbf{L} + \mathcal{P}_\mathbf{H}, \quad (T(u))(x_1) = \mathcal{T}_\mathbf{L} + \mathcal{T}_\mathbf{H},
$$

with

$$
\mathcal{V}_{\mathbf{L}} = \sum_{n \in \mathbf{L}} u_n \sin(nx_1) + \sum_{n \in \mathbf{L}} v_n \cos(nx_1),
$$
  
\n
$$
\mathcal{V}_{\mathbf{H}} = \sum_{n \in \mathbf{H}} u_n \sin(nx_1) + \sum_{n \in \mathbf{H}} v_n \cos(nx_1),
$$
  
\n
$$
\mathcal{P}_{\mathbf{L}} = \sum_{n \in \mathbf{L}} (\partial_\nu u)_n \sin(nx_1) + \sum_{n \in \mathbf{L}} (\partial_\nu v)_n \cos(nx_1),
$$
  
\n
$$
\mathcal{P}_{\mathbf{H}} = \sum_{n \in \mathbf{H}} (\partial_\nu u)_n \sin(nx_1) + \sum_{n \in \mathbf{H}} (\partial_\nu v)_n \cos(nx_1),
$$
  
\n
$$
\mathcal{T}_{\mathbf{L}} = \sum_{n \in \mathbf{L}} (Tu)_n \sin(nx_1) + \sum_{n \in \mathbf{L}} (Tv)_n \cos(nx_1),
$$
  
\n
$$
\mathcal{T}_{\mathbf{H}} = \sum_{n \in \mathbf{H}} (Tu)_n \sin(nx_1) + \sum_{n \in \mathbf{H}} (Tv)_n \cos(nx_1).
$$

The higher frequency terms have useful properties presented in Lemma [3.1.](#page-8-0)

<span id="page-8-0"></span>**Lemma 3.1** If  $u(x_1, x_2)$  is the solution to the scattering problem [\(15\)](#page-6-3), then

$$
\|\mathcal{P}_{\mathbf{H}}\|_{A}^{2} \simeq \sum_{n\in\mathbf{H}} \left(\partial_{\nu} u\right)_{n} \overline{u_{n}} + \sum_{n\in\mathbf{H}} \left(\partial_{\nu} v\right)_{n} \overline{v_{n}} \simeq \|\mathcal{V}_{\mathbf{H}}\|_{B}^{2}.
$$

Proof If 
$$
n \in \mathbb{H}
$$
,  
\n
$$
\left( |a_n \sqrt{k^2 - n^2} \cos \left( \sqrt{k^2 - n^2} b \right)|^2 + |-b_n \sqrt{k^2 - n^2} \sin \left( \sqrt{k^2 - n^2} b \right)|^2 \right) \frac{1}{\sqrt{|k^2 - n^2|} + \frac{1}{b^2|}} \approx \left( |a_n \sin \left( \sqrt{k^2 - n^2} b \right)|^2 + |b_n \cos \left( \sqrt{k^2 - n^2} b \right)|^2 \right) \left( \sqrt{|k^2 - n^2|} + \frac{1}{b} \right).
$$
\nHence  
\n
$$
\left( |(\partial_\nu u)_n|^2 + |(\partial_\nu v)_n|^2 \right) = \frac{1}{\sqrt{|k^2 - n^2|}} \approx \left( |u_n|^2 + |v_n|^2 \right) \left( \sqrt{|k^2 - n^2|} + \frac{1}{b} \right).
$$

$$
\left( |(\partial_{\nu} u)_n|^2 + |(\partial_{\nu} v)_n|^2 \right) \frac{1}{\sqrt{|k^2 - n^2|} + \frac{1}{b}} \simeq \left( |u_n|^2 + |v_n|^2 \right) \left( \sqrt{|k^2 - n^2|} + \frac{1}{b} \right).
$$
\n
$$
\text{where } (18), (20), \text{ we can show that for every } 5 \text{ H.} \text{ (3a)} \implies 0, (3a) \implies 0 \leq |u_n|^2 + |v_n|^2.
$$

From [\(18\)](#page-7-1)-[\(20\)](#page-7-2), one can show that for any  $n \in \mathbf{H}$ ,  $(\partial_{\nu}u)_n \overline{u_n} \geq 0$ ,  $(\partial_{\nu}v)_n \overline{v_n} \geq 0$ . Therefore,

$$
\|\mathcal{P}_{\mathbf{H}}\|_{A}^{2} \simeq \sum_{n \in \mathbf{H}} (\partial_{\nu} u)_{n} \overline{u_{n}} + \sum_{n \in \mathbf{H}} (\partial_{\nu} v)_{n} \overline{v_{n}}.
$$

The next result provides a quantitative estimate of the function  $u$  and its normal derivative.

<span id="page-9-1"></span>**Lemma 3.2** If  $u(x_1, x_2)$  is the solution to the scattering problem [\(15\)](#page-6-3), then

<span id="page-9-0"></span>
$$
\Lambda \sum_{n \in \mathbb{Z}} \sqrt{|k^2 - \alpha_n^2|} |\hat{u}_n|^2 \ge \sqrt{\varepsilon} \sqrt{k} \|u\|_{L^2(\Gamma)}^2.
$$
 (22)

Proof Since  $\Lambda \sum_{n\in\mathbb{Z}}|\hat{u}_n|^2 = ||u||^2_{L^2(\Gamma)}$  and  $\sqrt{|k^2 - \alpha_n^2|} \geq \sqrt{\varepsilon}\sqrt{k}$  for  $|k - \alpha_n| \geq$ ε, <sup>∀</sup><sup>n</sup> <sup>∈</sup> <sup>Z</sup>, Λ X <sup>n</sup>∈<sup>Z</sup>  $\sqrt{|k^2 - \alpha_n^2|} |\hat{u}_n|^2 \ge \sqrt{\varepsilon} \sqrt{k} \|u\|_{L^2(\Gamma)}^2,$ which completes the proof.  $\Box$ 

**Remark 3.1** Different from Lemma 3.5 in [\[8](#page-25-3)] for the scattering by a cavity, TBC of gratings yield additional difficulties on resonances and stability. Hence, in this paper, a uniform distance  $\varepsilon$  is introduced to exclude resonances and thus obtain the estimate [\(22\)](#page-9-0) in Lemma [3.2,](#page-9-1) which is essential for the proof procedure of the k-explicit stability for periodic structures.

Next, the relation between the norm of A (or B) and the  $L^2$  norm is established in Remark [3.2.](#page-9-2)

<span id="page-9-2"></span>**Remark 3.2** For the norm A, if there exists a natural number  $n^*$  such that  $\sqrt{|k^2-n^{*2}|} \leq \frac{1}{b}$ , then  $n^*$  is very close to k so that  $|k-n^*| \leq \frac{1}{b^2k}$ . Since  $\sqrt{|k^2 - n^2|} \ge \sqrt{\frac{k}{2}}$ , the norm A can be simplified as  $||u||_A \simeq \sqrt{\sqrt{\left(\frac{1}{2}\right)^2+1}}$  $\left| \right|$  $n∈\mathbb{N}\backslash\{n^*\}$  $(|u_n|^2 + |v_n|^2) \frac{1}{\sqrt{|v_n^2|}}$  $\sqrt{|k^2 - n^2|}$  $\setminus$  $+ (|u_{n^*}|^2 + |v_{n^*}|^2) b.$ 

If such  $n^*$  does not exist, then the norm A has no relation to b since

$$
||u||_A \simeq \sqrt{\sum_{n \in \mathbb{N}} (|u_n|^2 + |u_n|^2) \frac{1}{\sqrt{|k^2 - n^2|}}}.
$$

Hence, whether  $n^*$  exists or not, there always exists a constant C independent of k and b such that

$$
||u||_A^2 \leq C \left( \frac{1}{\sqrt{k}} ||u||_{L^2(\Gamma)}^2 + b \left( |u_{k^*}|^2 + |v_{k^*}|^2 \right) \right),
$$

where  $k^* = \arg \min_{n \in \mathbb{N}} |n - k|, u_{k^*} = \frac{1}{\pi} \int_{\Gamma} u(x_1) \sin (k^* x_1) dx_1$  and  $v_{k^*} =$  $\frac{1}{\pi} \int_{\Gamma} u(x_1) \cos(k^*x_1) dx_1$ . It follows immediately that

$$
||u||_A^2 \le Cb||u||_{L^2(\Gamma)}^2.
$$
\n(23)

The similar approximation can be deduced for the norm B:

<span id="page-10-2"></span>
$$
b||u||_B^2 \le C||u||_{L^2(\Gamma)}^2.
$$
\n(24)

As for the proof of Theorem [2.1,](#page-5-0) the first step given in Lemma [3.3](#page-10-0) is to estimate the norm of u in D by the norms of u on  $\Gamma$ , i.e., the norms A and B, and the second step given in Lemma [3.4](#page-10-1) is to estimate  $\|\partial_\nu u\|_A$  and  $\|u\|_B$ by  $||g||_A$ . Finally, with the correlation between the norm A and the  $L^2$  norm discussed in Remark [3.2,](#page-9-2) the stability estimate of Theorem [2.1](#page-5-0) is proved.

<span id="page-10-0"></span>**Lemma 3.3** Let u be a solution to the scattering problem  $(15)$ . Under Assumption [2.1,](#page-3-2) there exists a constant C independent of  $b > 1$  and large k, such that

$$
\|\nabla u\|_{L^{2}(D)} + k\|u\|_{L^{2}(D)} \leq Cbk \left(\|\partial_{\nu} u\|_{A} + \|u\|_{B}\right),\tag{25}
$$

where the norms  $A$  and  $B$  are defined in Definition [3.1.](#page-8-1)

<span id="page-10-1"></span>**Lemma 3.4** Let u be a solution to the scattering problem  $(15)$ . Under Assumption [2.1,](#page-3-2) there exists a constant C independent of  $b > 1$  and large k, such that

$$
\|\partial_{\nu}u\|_{A} + \|u\|_{B} \le C \max\left\{\frac{b^{\frac{1}{2}}k}{\sqrt{\varepsilon}}, b^{\frac{3}{2}}k^{\frac{3}{2}}\right\} \|g\|_{A},\tag{26}
$$

where the norms A and B are defined in Definition [3.1.](#page-8-1)

### 3.2 Proof of Lemma [3.3](#page-10-0) and Lemma [3.4](#page-10-1)

The proof of the bound in Lemma [3.3](#page-10-0) is mainly based on the definition of the norms and the representation of the series, which means that a stability estimate for the solution  $u$  in  $D$  can be given in terms of its Dirichlet and Neumann data on Γ.

*Proof of Lemma* [3.3](#page-10-0) From the equation  $\Delta u + k^2 u = 0$  in D and its boundary condition on  $S$  as well as the quasi-periodicity of the solution  $u$ , one has

$$
\int_{\Gamma} u \overline{\partial_{\nu} u} \mathrm{d}x_1 = \int_{D} \left( |\nabla u|^2 - k^2 |u|^2 \right) \mathrm{d}x_1 \mathrm{d}x_2.
$$

The left-hand side of the equality above is obviously bounded by  $\|\partial_\nu u\|_A^2 + \|u\|_B^2$ . However, the right-hand side has a negative sign for the term  $|u|^2$ . In order to prove Lemma [3.3,](#page-10-0) it suffices to show that

<span id="page-11-0"></span>
$$
||u||_{L^{2}(D)}^{2} \leq Cb^{2} \left( ||\partial_{\nu}u||_{A}^{2} + ||u||_{B}^{2} \right).
$$
 (27)

Note that  $\partial_\nu u$ , u on Γ, and u in D can be expressed in terms of sine and cosine functions in the direction of  $x_1$ . Because of the orthogonality of sine and cosine functions, one may assume that

$$
u(x_1, x_2) = \eta \sin\left(\sqrt{k^2 - n^2} x_2\right) \sin(nx_1) + \zeta \cos\left(\sqrt{k^2 - n^2} x_2\right) \cos(nx_1) \text{ in } D, n \neq k.
$$

For  $n \neq k$ , consider three cases: (i)  $n < k$  and  $b\sqrt{k^2 - n^2} \geq 1$ ; (ii)  $b\sqrt{|k^2 - n^2|} < 1$ ; and (iii)  $n > k$  and  $b\sqrt{|k^2 - n^2|} \geq 1$ .

For Case (i), it is easy to see that  $||u||_{L^2(D)}^2 \simeq b$ . On the boundary  $\Gamma$ ,

$$
\partial_{\nu}u = \eta\sqrt{k^2 - n^2}\cos\left(\sqrt{k^2 - n^2}b\right)\sin(nx_1) - \zeta\sqrt{k^2 - n^2}\sin\left(\sqrt{k^2 - n^2}b\right)\cos(nx_1),
$$
  

$$
u = \eta\sin\left(\sqrt{k^2 - n^2}b\right)\sin(nx_1) + \zeta\cos\left(\sqrt{k^2 - n^2}b\right)\cos(nx_1).
$$

From the definition of the norms A and B, one has

$$
\|\partial_{\nu}u\|_{A}^{2} + \|u\|_{B}^{2}
$$
\n
$$
= |\eta|^{2} \left( \frac{k^{2} - n^{2}}{\sqrt{k^{2} - n^{2} + \frac{1}{b}}} |\cos \left( \sqrt{k^{2} - n^{2}b} \right)|^{2} + \left( \sqrt{k^{2} - n^{2} + \frac{1}{b}} \right) |\sin \left( \sqrt{k^{2} - n^{2}b} \right)|^{2} \right)
$$
\n
$$
+ |\zeta|^{2} \left( \frac{k^{2} - n^{2}}{\sqrt{k^{2} - n^{2} + \frac{1}{b}}} |\sin \left( \sqrt{k^{2} - n^{2}b} \right)|^{2} + \left( \sqrt{k^{2} - n^{2} + \frac{1}{b}} \right) |\cos \left( \sqrt{k^{2} - n^{2}b} \right)|^{2} \right)
$$
\n
$$
\geq \frac{1}{2} (|\eta|^{2} + |\zeta|^{2}) \sqrt{k^{2} - n^{2}} \geq \frac{(|\eta|^{2} + |\zeta|^{2})}{2b} \geq C \frac{1}{b^{2}} \|u\|_{L^{2}(D)}^{2}.
$$

For Case (ii), one has

$$
|\sin\left(\sqrt{k^2 - n^2}x_2\right)| \simeq \sqrt{|k^2 - n^2|}x_2 \text{ and } |\cos\left(\sqrt{k^2 - n^2}x_2\right)| \simeq 1 - \frac{1}{2}|k^2 - n^2|x_2^2|
$$
  
from  $\sqrt{|k^2 - n^2|}b < 1$ . Therefore,  $||u||_{L^2(D)}^2 \simeq |k^2 - n^2|b^3$  and  

$$
||u(x, b)||_B^2 \simeq \left(\sqrt{|k^2 - n^2|} + \frac{1}{b}\right)|k^2 - n^2|b^2 \simeq b|k^2 - n^2|,
$$

which implies the inequality [\(27\)](#page-11-0).

Finally, for Case (iii),

$$
u = \eta \sin\left(\sqrt{k^2 - n^2}x_2\right) \sin(nx_1) + \zeta \cos\left(\sqrt{k^2 - n^2}x_2\right) \cos(nx_1)
$$
  
=  $\eta i \sinh\left(\sqrt{n^2 - k^2}x_2\right) \sin(nx_1) + \zeta \cosh\left(\sqrt{n^2 - k^2}x_2\right) \cos(nx_1)$  in *D*.

The representation of u yields

$$
||u||_{L^{2}(D)}^{2} \leq C \frac{1}{\sqrt{n^{2}-k^{2}}} e^{2\sqrt{n^{2}-k^{2}}b} \leq Cb^{2} \left(\sqrt{n^{2}-k^{2}}+\frac{1}{b}\right) e^{2\sqrt{n^{2}-k^{2}}b} \leq Cb^{2} ||u(\cdot,b)||_{B}^{2}.
$$

For the proof of Lemma [3.4,](#page-10-1) the main difficulty is due to the fact that the boundary data  $\partial_{\nu}u = Tu + g$  does not yield direct estimates on  $\|\partial_{\nu}u\|_{A}$  and  $||u||_B$ . To be specific, let us consider the key term  $-\mathfrak{Im} \int_{\Gamma} g \bar{u} dx_1$ . Obviously,

<span id="page-12-0"></span>
$$
-\Im \mathfrak{m} \int_{\Gamma} g \bar{u} \mathrm{d} x_1 = \Lambda \sum_{|\alpha_n| < k} \sqrt{k^2 - \alpha_n^2} |\hat{u}_n|^2 \ge 0. \tag{28}
$$

However, the integral in [\(28\)](#page-12-0) does not contain information on  $\hat{u}$  for  $|\alpha_n| > k$ . In order to obtain the stability, we have to analyze  $\hat{u}$  for  $|\alpha_n| > k$ , and thus to generate a bound on u. Next, we give the proof of Lemma [3.4.](#page-10-1)

Proof of Lemma [3.4](#page-10-1) From the definition,

$$
\int_{\Gamma} g \bar{u} dx_1 = \int_{\Gamma} (\partial_{\nu} u - T(u)) \bar{u} dx_1
$$
\n
$$
= \int_{\Gamma} \partial_{\mathbf{n}} u \bar{u} dx_1 - i \Lambda \sum_{|n| < k} \sqrt{k^2 - n^2} |\hat{u}_n|^2 + \Lambda \sum_{|n| > k} \sqrt{n^2 - k^2} |\hat{u}_n|^2
$$
\n
$$
\triangleq \mathfrak{Re} + i \mathfrak{Im},
$$

with  $\Re$  =  $\int_{\Gamma} \partial_{\mathbf{n}} u \bar{u} \mathrm{d} x_1 + \Lambda \sum_{|n|>k}$  $\sqrt{n^2 - k^2} |\hat{u}_n|^2$  and  $\mathfrak{Im}$  =  $-\Lambda \sum_{|n|< k}$  $\sqrt{k^2 - n^2} |\hat{u}_n|^2$ . Obviously,

<span id="page-12-3"></span>
$$
\Lambda \sum_{|n| (29)
$$

As for the real part, since

$$
|\Re\mathfrak{e}| \leq |\int_{\Gamma} g\bar{u}\mathrm{d}x_1| \leq \pi ||g||_A ||u||_B \leq \gamma\pi ||\partial_{\nu}u||_A ||u||_B
$$

provieded that

<span id="page-12-2"></span>
$$
||g||_A \le \gamma ||\partial_\nu u||_A,\tag{30}
$$

and

$$
\mathfrak{Re} = \sum_{n \in \mathbf{L}} (\partial_{\nu} u)_n \bar{u}_n + \sum_{n \in \mathbf{L}} (\partial_{\nu} v)_n \bar{v}_n + \sum_{n \in \mathbf{H}} (\partial_{\nu} u)_n \bar{u}_n + \sum_{n \in \mathbf{H}} (\partial_{\nu} v)_n \bar{v}_n
$$
  
+ 
$$
\Lambda \sum_{|n| > k} \sqrt{n^2 - k^2} |\hat{u}_n|^2,
$$

one obtains

<span id="page-12-1"></span>
$$
\gamma \pi \left\| \partial_{\nu} u \right\|_{A} \| u \|_{B} + \left| \sum_{n \in \mathbf{L}} (\partial_{\nu} u)_{n} \bar{u}_{n} + \sum_{n \in \mathbf{L}} (\partial_{\nu} v)_{n} \bar{v}_{n} \right|
$$
  
\n
$$
\geq \sum_{n \in \mathbf{H}} (\partial_{\nu} u)_{n} \overline{u}_{n} + \sum_{n \in \mathbf{H}} (\partial_{\nu} v)_{n} \overline{v}_{n} + \Lambda \sum_{|n| > k} \sqrt{n^{2} - k^{2}} |\hat{u}_{n}|^{2},
$$
\n(31)

with

$$
\sum_{n \in \mathbf{H}} (\partial_{\nu} u)_n \overline{u_n} + \sum_{n \in \mathbf{H}} (\partial_{\nu} v)_n \overline{v_n} \simeq ||\mathcal{P}_\mathbf{H}||_A^2 \ge 0
$$

and

$$
\Lambda \sum_{|n|>k} \sqrt{n^2 - k^2} |\hat{u}_n|^2 \ge 0.
$$

The estimate [\(31\)](#page-12-1) implies that  $\sum (\partial_\nu u)_n \overline{u_n} + \sum (\partial_\nu v)_n \overline{v_n}$  and  $n \in H$   $n \in H$ Λ X  $|n|>k$  $\sqrt{n^2 - k^2 |\hat{u}_n|^2}$  can be bounded by  $|\sum$  $n \in \mathbf{L}$  $(\partial_\nu u)_n \bar{u}_n + \sum$  $n \in \mathbf{L}$  $(\partial_\nu v)_n \bar{v}_n$  since  $\gamma \pi \left\| (\partial_\nu u)_n \right\|_A \|u\|_B$  may be absorbed by choosing an appropriate  $\gamma$ .

Evidently, in order to estimate  $\Lambda$  $|n|>k$  $\sqrt{n^2 - k^2} |\hat{u}_n|^2$ , it is sufficient to estimate

 $|\sum$  $n \in \mathbf{L}$  $\left(\partial_\nu u\right)_n \bar{u}_n + \sum$ <sup>n</sup>∈<sup>L</sup>  $(\partial_\nu v)_n \bar{v}_n$ . Since the estimate [\(31\)](#page-12-1) is based on the condition

[\(30\)](#page-12-2), we will consider  $||g||_A \leq \gamma ||\partial_\nu u||_A$  and  $||g||_A \geq \gamma ||\partial_\nu u||_A$  separately. For the case  $||g||_A \leq \gamma ||\partial_\nu u||_A$ , since the term  $|\sum$  $n \in \mathbf{L}$  $(\partial_\nu u)_n\,\bar u_n + \sum$  $n \in \mathbf{L}$  $(\partial_\nu v)_n \bar{v}_n$  concerns

the low frequency part for  $u_n$  and  $(\partial_\nu u)_n$ , to characterize which term is bigger,  $\delta \|\mathcal{P}_{\mathbf{L}}\|_{A} \geq \|\mathcal{V}_{\mathbf{L}}\|_{B}$  and  $\delta \|\mathcal{P}_{\mathbf{L}}\|_{A} \leq \|\mathcal{V}_{\mathbf{L}}\|_{B}$  are considered separately. To sum up, we consider the following three cases:

 $\bullet$  Case (i)

$$
||g||_A \le \gamma ||\partial_\nu u||_A, \quad \delta ||\mathcal{P}_L||_A \ge ||\mathcal{V}_L||_B; \tag{32}
$$

 $\bullet$  Case (ii)

$$
||g||_A \le \gamma ||\partial_\nu u||_A, \quad \delta ||\mathcal{P}_L||_A \le ||\mathcal{V}_L||_B.
$$
 (33)

 $\bullet$  Case (iii)

<span id="page-13-5"></span>
$$
||g||_A \ge \gamma ||\partial_\nu u||_A;\tag{34}
$$

The constant  $\gamma$  and  $\delta$  are small numbers which will be chosen later.

For Case (i), it yields from [\(31\)](#page-12-1) that

<span id="page-13-0"></span>
$$
\delta \|\mathcal{P}_{\mathbf{L}}\|_{A}^{2} + \gamma \pi \left\| (\partial_{\nu} u)_{n} \right\|_{A} \|u\|_{B}
$$
\n
$$
\geq \left( \sum_{n \in \mathbf{H}} (\partial_{\nu} u)_{n} \bar{u}_{n} + \sum_{n \in \mathbf{H}} (\partial_{\nu} v)_{n} \overline{v_{n}} \right) + \Lambda \sum_{|n| > k} \sqrt{n^{2} - k^{2}} |\hat{u}_{n}|^{2}.
$$
\n
$$
(35)
$$

For the second term in [\(35\)](#page-13-0), Case (i) and Lemma [3.1](#page-8-0) yield

<span id="page-13-1"></span>
$$
\gamma \pi \left\| (\partial_{\nu} u)_n \right\|_A \|u\|_B \leq \gamma \pi \left( \|\mathcal{P}_L\|_A + C \|\mathcal{V}_H\|_B \right) \times \left( \delta \|\mathcal{P}_L\|_A + C \|\mathcal{V}_H\|_B \right)
$$
  

$$
\leq C' \left( \gamma \delta \|\mathcal{P}_L\|_A^2 + \gamma^2 \|\mathcal{P}_L\|_A^2 + \frac{1}{2} \|\mathcal{V}_H\|_B^2 \right).
$$
 (36)

Since

<span id="page-13-4"></span>
$$
\gamma^2 \|\partial_\nu u\|_A^2 \ge \|\partial_\nu u - T(u)\|_A^2 \ge \|\mathcal{P}_\mathbf{L} - \mathcal{T}_\mathbf{L}\|_A^2 \ge \frac{1}{2} \|\mathcal{P}_\mathbf{L}\|_A^2 - \|\mathcal{T}_\mathbf{L}\|_A^2, \qquad (37)
$$

one has

<span id="page-13-2"></span>
$$
\frac{1}{2} \|\mathcal{P}_{\mathbf{L}}\|_{A}^{2} \le \|\mathcal{T}_{\mathbf{L}}\|_{A}^{2} + \gamma^{2} \|\partial_{\nu}u\|_{A}^{2} \le \|\mathcal{T}_{\mathbf{L}}\|_{A}^{2} + \gamma^{2} \left( \|\mathcal{P}_{\mathbf{L}}\|_{A}^{2} + C \|\mathcal{V}_{\mathbf{H}}\|_{B}^{2} \right). \tag{38}
$$

By  $(35)$ ,  $(36)$  and  $(38)$ , choosing

<span id="page-13-6"></span>
$$
\gamma = C_{(i)} \sqrt{\delta},\tag{39}
$$

where  $C_{(i)}$  is an arbitrarily sufficiently small constant independent of k and b, it can be deduced that

<span id="page-13-3"></span>
$$
\|\mathcal{T}_{\mathbf{L}}\|_{A}^{2} \ge \frac{C_{1}}{\delta} \left( \|\mathcal{V}_{\mathbf{H}}\|_{B}^{2} + \Lambda \sum_{|n|>k} \sqrt{n^{2}-k^{2}} |\hat{u}_{n}|^{2} \right). \tag{40}
$$

In order to get a bound for  $||\mathcal{T}_{\mathbf{L}}||_A^2$ , it is sufficient to estimate  $(Tu)_n$  and  $(Tv)_n$ for  $n \leq k$  due to the definition of **L**. By the Cauchy-Schwarz inequality, one has

$$
\sum_{|n|k} \sqrt{n^2 - k^2} |\hat{u}_n|^2
$$
\n
$$
\geq |\sum_{n\in\mathbb{Z}} \sqrt{k^2 - n^2} \hat{u}_n \overline{\sin(nx_1)}|^2 \left( \sum_{n\in\mathbb{Z}} \sqrt{|k^2 - n^2| |\sin(nx_1)|^2} \right)^{-1}
$$
\n
$$
\geq |(Tu)_n|^2 \left( \sum_{n\in\mathbb{Z}} \sqrt{|k^2 - n^2| |\sin(nx_1)|^2} \right)^{-1}
$$
\n
$$
\geq \frac{1}{2k} |(Tu)_n|^2 \quad n < k,
$$
\n(41)

and

<span id="page-14-1"></span>
$$
\sum_{|n|k} \sqrt{n^2 - k^2} |\hat{u}_n|^2
$$
\n
$$
\geq |\sum_{n\in\mathbb{Z}} \sqrt{k^2 - n^2} \hat{u}_n \overline{\cos(nx_1)}|^2 \left(\sum_{n\in\mathbb{Z}} \sqrt{|k^2 - n^2| |\cos(nx_1)|^2}\right)^{-1}
$$
\n
$$
\geq |(Tv)_n|^2 \left(\sum_{n\in\mathbb{Z}} \sqrt{|k^2 - n^2| |\cos(nx_1)|^2}\right)^{-1}
$$
\n
$$
\geq \frac{1}{2k} |(Tv)_n|^2 \quad n < k.
$$
\n(42)

Note that

<span id="page-14-0"></span>
$$
\sum_{n \in \mathbf{L}} \frac{1}{\sqrt{|k^2 - n^2|} + \frac{1}{b}} = \sum_{n=0}^{\lfloor k \rfloor - 1} \frac{1}{\sqrt{|k^2 - n^2|} + \frac{1}{b}} + \frac{1}{\sqrt{|k^2 - \lfloor k \rfloor^2|} + \frac{1}{b}}
$$
\n
$$
\leq \int_0^{\lfloor k \rfloor} \frac{1}{\sqrt{|k^2 - x^2|} + \frac{1}{b}} dx + b
$$
\n
$$
\leq C + b \leq C_2 b
$$
\n(43)

since  $b > 1$ . A combination of the above estimate [\(43\)](#page-14-0) with [\(40\)](#page-13-3)-[\(42\)](#page-14-1) yields

<span id="page-14-2"></span>
$$
C_2bk\left(\Lambda \sum_{|n|k} \sqrt{n^2 - k^2} |\hat{u}_n|^2\right)
$$
  
\n
$$
\geq \sum_{n\in\mathbf{L}} \frac{1}{\sqrt{|k^2 - n^2|} + \frac{1}{b}} \left(|(Tu)_n|^2 + |(Tv)_n|^2\right)
$$
  
\n
$$
= ||\mathcal{T}_{\mathbf{L}}||_A^2
$$
  
\n
$$
\geq \frac{C_1}{\delta} \left(||V_{\mathbf{H}}||_B^2 + \Lambda \sum_{|n|>k} \sqrt{n^2 - k^2} |\hat{u}_n|^2\right)
$$
  
\n
$$
\geq \frac{C_1}{\delta} \Lambda \sum_{|n|>k} \sqrt{n^2 - k^2} |\hat{u}_n|^2.
$$
 (44)

By choosing

<span id="page-15-1"></span>
$$
\delta = \frac{C_{(ii)}}{bk},\tag{45}
$$

with  $C_{(ii)} = \frac{2C_1}{1+2C_2} > 0$ , which is independent of k and  $\varepsilon$ , one has

<span id="page-15-0"></span>
$$
C \sum_{|n|k} \sqrt{n^2 - k^2} |\hat{u}_n|^2. \tag{46}
$$

Applying the above estimate [\(46\)](#page-15-0) and the choice of  $\delta$  [\(45\)](#page-15-1) to [\(44\)](#page-14-2), one has

<span id="page-15-2"></span>
$$
C \sum_{|n|k} \sqrt{n^2 - k^2} |\hat{u}_n|^2. \tag{47}
$$

By combining  $(29)$ ,  $(47)$ , Lemma [3.2](#page-9-1) and the proposition  $(24)$  of norm B, one gets  $Ck$  $\int_\Gamma g \bar{u} \mathrm{d} x_1 |\geq k\, \|\mathcal{V}_{\mathbf{H}}\|_B^2 + C'' \sqrt{\varepsilon} k \sqrt{k} \|u\|_{L^2(\Gamma)}^2 \geq k\, \|\mathcal{V}_{\mathbf{H}}\|_B^2 + \sqrt{\varepsilon} \sqrt{k}\, \|\mathcal{V}_{\mathbf{L}}\|_B^2 \geq \sqrt{\varepsilon} \sqrt{k} \|u\|_B^2.$ Hence,

<span id="page-15-5"></span>
$$
\frac{C\sqrt{k}}{\sqrt{\varepsilon}}\|g\|_A \ge \|u\|_B. \tag{48}
$$

Moreover, since Lemma [3.1](#page-8-0) indicates

<span id="page-15-3"></span>
$$
\left\| \mathcal{V}_{\mathbf{H}} \right\|_{B}^{2} \ge C \left\| \mathcal{P}_{\mathbf{H}} \right\|_{A}^{2},\tag{49}
$$

and [\(37\)](#page-13-4) implies

<span id="page-15-4"></span>
$$
\|\mathcal{T}_{\mathbf{L}}\|_{A}^{2} \ge \frac{1}{2} \|\mathcal{P}_{\mathbf{L}}\|_{A}^{2} - \gamma^{2} \|\partial_{\nu}u\|_{A}^{2},\tag{50}
$$

a combination of the above  $(49)-(50)$  $(49)-(50)$  with  $(29)$ ,  $(44)$  and Lemma [3.1](#page-8-0) yields

$$
Cbk\pi\|g\|_A\|u\|_B \ge Cbk|\int_\Gamma g\bar{u}\mathrm{d} x_1| \ge Cbk\Lambda \sum_{|n|
$$

Therefore, using [\(48\)](#page-15-5), one has

$$
\frac{Cbk^{\frac{3}{2}}}{\varepsilon} \|g\|_A^2 \ge \|\partial_\nu u\|_A^2.
$$

Hence,

<span id="page-15-6"></span>
$$
\frac{Cb^{\frac{1}{2}}k^{\frac{3}{4}}}{\sqrt{\varepsilon}}\|g\|_{A} \ge \|\partial_{\nu}u\|_{A}.
$$
\n(51)

The estimate  $(48)$  and  $(51)$  indicate that

<span id="page-15-9"></span>
$$
\frac{Cb^{\frac{1}{2}}k^{\frac{3}{4}}}{\sqrt{\varepsilon}}\|g\|_{A} \ge \|\partial_{\nu}u\|_{A} + \|u\|_{B}.
$$
 (52)

For Case (ii), Applying Lemma [3.1](#page-8-0) to [\(31\)](#page-12-1), one has

<span id="page-15-7"></span>
$$
\frac{1}{\delta} \|\mathcal{V}_{\mathbf{L}}\|_{B}^{2} + \gamma \pi \|\partial_{\nu} u\|_{A} \|u\|_{B} \ge C \|\mathcal{V}_{\mathbf{H}}\|_{B}^{2} + \Lambda \sum_{|n|>k} \sqrt{n^{2}-k^{2}} |\hat{u}_{n}|^{2}.
$$
 (53)

For the second term in [\(53\)](#page-15-7), Case (ii) and Lemma [3.1](#page-8-0) yield

<span id="page-15-8"></span>
$$
\gamma \pi \|\partial_{\nu} u\|_{A} \|u\|_{B} \leq \gamma \pi \left(\frac{1}{\delta} \|\mathcal{V}_{\mathbf{L}}\|_{B} + C \|\mathcal{V}_{\mathbf{H}}\|_{B}\right) \times \left(\|\mathcal{V}_{\mathbf{L}}\|_{B} + \|\mathcal{V}_{\mathbf{H}}\|_{B}\right)
$$
  

$$
\leq C \frac{\gamma}{\delta} \|\mathcal{V}_{\mathbf{L}}\|_{B}^{2} + C \gamma^{2} \|\mathcal{V}_{\mathbf{L}}\|_{B}^{2} + \frac{1}{2} \|\mathcal{V}_{\mathbf{H}}\|_{B}^{2}.
$$
 (54)

Since  $\gamma = C_{(i)}\sqrt{\delta}$  is sufficiently small, the estimate [\(53\)](#page-15-7) and [\(54\)](#page-15-8) indicate that

<span id="page-16-0"></span>
$$
\frac{C}{\delta} \|\mathcal{V}_{\mathbf{L}}\|_{B}^{2} \geq \|\mathcal{V}_{\mathbf{H}}\|_{B}^{2} + \Lambda \sum_{|n|>k} \sqrt{n^{2}-k^{2}} |\hat{u}_{n}|^{2} \geq \|\mathcal{V}_{\mathbf{H}}\|_{B}^{2}.
$$
 (55)

On the other hand, for the lower frequency, it is obvious that

$$
C\Lambda \sum_{|n|
$$

Hence, it follows from [\(29\)](#page-12-3) that

$$
C|\int_{\Gamma} g\bar{u}\mathrm{d}x_1|\geq C\Lambda \sum_{|n|
$$

Thus

<span id="page-16-2"></span>
$$
C||g||_A||u||_B \ge ||\mathcal{V}_{\mathbf{L}}||_B^2.
$$
\n(56)

By [\(55\)](#page-16-0) and  $\delta = \frac{C_{(ii)}}{bk}$ , one has

$$
Cbk||g||_A||u||_B \ge ||u||_B^2.
$$

Thus

<span id="page-16-4"></span>
$$
Cbk||g||_A \ge ||u||_B.
$$
\n
$$
(57)
$$

Note that

<span id="page-16-1"></span>
$$
||u||_B \ge ||\mathcal{V}_{\mathbf{L}}||_B \ge \frac{C_{(ii)}}{bk} ||\mathcal{P}_{\mathbf{L}}||_A, \tag{58}
$$

due to the assumption of Case (ii). Applying  $(58)$  to  $(56)$ , one gets

$$
Cb^2k^2||g||_A||u||_B \ge ||\mathcal{P}_{\mathbf{L}}||_A^2.
$$

Using [\(55\)](#page-16-0) and Lemma [3.1,](#page-8-0) this implies that

<span id="page-16-3"></span>
$$
Cb^2k^2\|g\|_A\|u\|_B \ge \|\partial_\nu u\|_A^2. \tag{59}
$$

Combining the above estimate  $(59)$  with  $(57)$  yields

<span id="page-16-5"></span>
$$
Cb^{\frac{3}{2}}k^{\frac{3}{2}}\|g\|_{A} \ge \|\partial_{\nu}u\|_{A}.\tag{60}
$$

The estimate  $(57)$  and  $(60)$  indicate that

<span id="page-16-8"></span>
$$
Cb^{\frac{3}{2}}k^{\frac{3}{2}}\|g\|_A \ge \|\partial_\nu u\|_A + \|u\|_B. \tag{61}
$$

For Case (iii), it suffices to estimate  $||u||_B$ . Using the definition of  $T(u)$ , the property of the norm defined above and Lemma [3.2,](#page-9-1) one obtains

<span id="page-16-7"></span>
$$
\left(1+\frac{1}{\gamma}\right)\sqrt{k}\pi\|g\|_{A}\|u\|_{B} \ge \sqrt{k}\pi\|T(u)\|_{A}\|u\|_{B}
$$

$$
\ge \sqrt{k}|\Lambda \sum_{n\in\mathbb{Z}}\sqrt{k^{2}-n^{2}}|\hat{u}_{n}|^{2}|\left(\ge \sqrt{k}\Big|\frac{\Lambda}{2}\sum_{n\in\mathbb{Z}}\sqrt{|k^{2}-n^{2}}|\hat{u}_{n}|^{2}|\right)
$$

$$
\ge \frac{\sqrt{\varepsilon}}{2}k\|u\|_{L^{2}(\Gamma)}^{2}.
$$
(62)

Since  $k \geq \sqrt{k^2 - n^2}$  for  $n \in \mathbf{L}$ , it is obvious that there exists a constant C such that

<span id="page-16-6"></span>
$$
Ck \ge \sqrt{|k^2 - n^2|} + \frac{1}{b}.\tag{63}
$$

Applying [\(63\)](#page-16-6) to [\(62\)](#page-16-7), the bound for  $n \in L$  can be deduced:

<span id="page-17-0"></span>
$$
C\frac{\sqrt{k}}{\sqrt{\varepsilon}\gamma}||g||_A||u||_B \ge \sum_{n\in\mathbf{L}} \left(\sqrt{|k^2 - n^2|} + \frac{1}{b}\right) \left(|u_n|^2 + |v_n|^2\right) = ||\mathcal{V}_\mathbf{L}||_B^2. \tag{64}
$$

Using Lemma [3.1,](#page-8-0) the bound for  $n \in H$  can be deduced:

<span id="page-17-1"></span>
$$
\frac{1}{\gamma^2} \|g\|_A^2 \ge \|\partial_\nu u\|_A^2 \ge \|\mathcal{P}_\mathbf{H}\|_A^2 \simeq \|\mathcal{V}_\mathbf{H}\|_B^2. \tag{65}
$$

Combining  $(64)$  and  $(65)$  yields

$$
C\left(\frac{1}{\gamma^2}||g||_A^2 + \frac{\sqrt{k}}{\sqrt{\varepsilon}\gamma}||g||_A||u||_B\right) \ge ||u||_B^2.
$$

Solving the second order polynomial above, one has

<span id="page-17-2"></span>
$$
C\frac{\sqrt{k}}{\sqrt{\varepsilon}\gamma}||g||_A \ge ||u||_B.
$$
\n(66)

Therefore, for Case  $(i)$ , one obtains from  $(34)$  and  $(66)$  that

<span id="page-17-3"></span>
$$
C\frac{\sqrt{k}}{\sqrt{\varepsilon}\gamma}||g||_A \ge ||\partial_\nu u||_A + ||u||_B.
$$
 (67)

Applying choice of  $\gamma$  [\(39\)](#page-13-6) and  $\delta$  [\(45\)](#page-15-1) to the estimate [\(67\)](#page-17-3) immediately yields that

<span id="page-17-4"></span>
$$
C\frac{b^{\frac{1}{2}}k}{\sqrt{\varepsilon}}\|g\|_{A} \ge \|\partial_{\mathbf{n}}u\|_{A} + \|u\|_{B}.
$$
 (68)

Finally, we complete the proof of Lemma [3.4](#page-10-1) based on the above proofs for Case  $(i)$ - $(iii)$ . Combining the bounds  $(52)$ ,  $(61)$  and  $(68)$  for Case  $(i)$ - $(iii)$  respectively, one arrives

$$
\|\partial_{\mathbf{n}}u\|_{A} + \|u\|_{B} \le C \max\left\{\frac{b^{\frac{1}{2}}k}{\sqrt{\varepsilon}}, b^{\frac{3}{2}}k^{\frac{3}{2}}\right\} \|g\|_{A},
$$
\nwhich completes the proof.

### 3.3 Proof of Theorem [2.1](#page-5-0)

Now we are ready to prove the bounds in Theorem [2.1.](#page-5-0)

Proof Lemma [3.4](#page-10-1) yields that: Under Assumption [2.1,](#page-3-2) there exists a constant C independent of  $b > 1$  and large k, such that

$$
\|\partial_{\nu}u\|_{A} + \|u\|_{B} \leq C \max\left\{\frac{b^{\frac{1}{2}}k}{\sqrt{\varepsilon}}, b^{\frac{3}{2}}k^{\frac{3}{2}}\right\} \|g\|_{A}.
$$

Furthermore, Lemma [3.3](#page-10-0) gives

$$
\|\nabla u\|_{L^2(D)} + k\|u\|_{L^2(D)} \le Cbk \left( \|\partial_\nu u\|_A + \|u\|_B \right).
$$

Therefore,

<span id="page-17-6"></span>
$$
\|\partial_{\nu}u\|_{A} + \|u\|_{B} \le C \max\left\{\frac{b^{\frac{3}{2}}k^{2}}{\sqrt{\varepsilon}}, b^{\frac{5}{2}}k^{\frac{5}{2}}\right\} \|g\|_{A}.
$$
 (69)

From the definition and the property of the norm A mentioned in Definition [3.1](#page-8-1) and Remark [3.2,](#page-9-2)

<span id="page-17-5"></span>
$$
||g||_A \le C\sqrt{b}||g||_{L^2(\Gamma)}.\tag{70}
$$

Applying [\(70\)](#page-17-5) to [\(69\)](#page-17-6), one derives the bounds in Theorem [2.1,](#page-5-0) which completes the proof.  $\Box$ 

## <span id="page-18-0"></span>4 Stability for the random case

In this section, Theorem [2.2](#page-6-0) is proved by utilizing the general framework developed in [\[24](#page-26-3)] based upon the stability in Theorem [2.1](#page-5-0) for the deterministic case. Note that the randomness of integral domain prevents direct application of general framework in [\[24\]](#page-26-3). Thus the difficulty lies in that the scattering surface in model problem is with randomness, which implies the stochastic domain for the scattering problem. For this, a variable transform, which changes the random variational form with stochastic domains into a transformed one with a definite domain and random medium, is introduced to prove all necessary propositions, including the continuity of sesquilinear form  $\tilde{a}$  and antilinear functional  $\tilde{G}$ , and stability and uniqueness which hold almost surely, which implies the stability for the random case.

#### 4.1 Prerequisites

Denote  $\tilde{a}_{c(\omega)}$  and  $\tilde{G}_{c(\omega)}$  (defined by [\(8\)](#page-4-0) and [\(10\)](#page-5-3) in Sec [2.2\)](#page-3-3) by  $(\tilde{a} \circ c)(\omega)$  and  $(\tilde{G} \circ c)(\omega)$ , respectively. Define the norm  $||v||_{1,k}^2 := ||\nabla v||_{L^2(D)}^2 + k^2 ||v||_{L^2(D)}^2$ and  $||v||_{1,\infty} := ||v||_{\infty} + ||v'||_{\infty}$  on  $H^1_{S,qp}(D)$ . Necessary propositions required in the general framework [\[24](#page-26-3)] include the continuity of  $\tilde{a}$  and  $\tilde{G}$ , regularity of  $\tilde{a} \circ c$  and  $\tilde{G} \circ c$ , measurability and  $\mu$ -essentially separability of c, stability a.s. and uniqueness a.s., described in Proposition [4.1](#page-18-1) and Proposition [4.2.](#page-18-2)

<span id="page-18-1"></span>**Proposition 4.1**  $\tilde{a}$ ,  $\tilde{G}$  and c in the variational form [\(11\)](#page-5-4) of the scattering problem in a random periodic structure satisfies the following properties:

- (i) The function  $c : \Omega \to \mathcal{C}$  defined by [\(6\)](#page-4-1) is measurable and  $\mu$ -essentially separably valued.
- (ii) Let  $B(H_{S,qp}^1(D), H_{S,qp}^1(D))$  denote the space of bounded linear maps  $H^1_{S,qp}(D) \rightarrow H^1_{S,qp}(D)$ . The functions  $\tilde{a}$  :  $C \rightarrow B(H^1_{S,qp}(D), H^1_{S,qp}(D))$ and  $\tilde{G}: \mathcal{C} \to H^1_{S,qp}(D)$  defined by [\(8\)](#page-4-0) and [\(10\)](#page-5-3) are continuous, the maps  $\tilde{a} \circ c \in L^{\infty}(\Omega; B(H^1_{S,qp}(D), H^1_{S,qp}(D))), \ \tilde{G} \circ c \in L^2(\Omega; H^1_{S,qp}(D)).$

**Definition 4.1** (The solution operator U) Define  $\mathcal{U}: \mathcal{C} \to H^1_{S,qp}(D)$  by letting  $\mathcal{U}(f_0) \;\in\; H^1_{S, qp}(D)$  be the solution of the deterministic Helmholtz problem with sampling  $f_0 \in \mathcal{C}$ .

<span id="page-18-2"></span>**Proposition 4.2** For  $f_0 \in Lip$ , the solution of the variational form [\(11\)](#page-5-4) exists and is unique. Let  $v_0 = \mathcal{U}(f_0)$  be the solution. Under Assumption [2.1,](#page-3-2) there exists a constant C independent of  $b > 1$  and large k, such that

<span id="page-18-3"></span>
$$
\|\nabla v_0\|_{L^2(D)} + k\|v_0\|_{L^2(D)} \le C \max\left\{\frac{b^2 k^2}{\sqrt{\varepsilon}}, b^3 k^{\frac{5}{2}}\right\} \|g\|_{L^2(\Gamma)}.
$$
 (71)

Moreover, the scattering problem  $(5)$  in a random periodic structure satisfies the following stability and uniqueness properties:

- (i) (Uniqueness almost surely)  $\text{ker}(\tilde{a}_{c(\omega)}) = \{0\}$   $\mu$ -almost surely.
- (ii) (Stability almost surely) There exist  $C_1, h_1 : \Omega \to \mathbb{R}$  such that  $C_1 h_1 \in L^1(\Omega)$ and the bound

$$
||u(\omega)||_{1,k}^2 \le C_1(\omega)h_1(\omega) \tag{72}
$$

holds almost surely.

**Remark 4.1** From the general framework in  $[24]$ , one can conclude that (i) with properties of  $\tilde{a}$ , G and c stated in Proposition [4.1,](#page-18-1) the maps A and G defined by [\(7\)](#page-4-3) and [\(9\)](#page-5-5) are well-defined; (ii) with the almost surely uniqueness property in Proposition [4.2,](#page-18-2) the solution to the Helmholtz equation in the stochastic case is unique in  $L^2(\Omega; H^1_{S,qp}(D));$  (iii) with the almost surely stability property in Proposition [4.2,](#page-18-2) integrability and measurability implies the stochastic a priori bound.

In order to verify Proposition [4.1](#page-18-1) and Proposition [4.2,](#page-18-2) we give the following four lemmas. Lemma [4.1](#page-19-0) means that the solution to the scattering problem depends continuously on the scattering surface. Lemma [4.2](#page-19-1) shows the equivalence between strong measurability and measurability as well as  $\mu$ -essentially separability. Lemma [4.3](#page-19-2) shows the necessary and sufficient condition for Bochner integrability. Lemma [4.4](#page-19-3) shows that the solution operator  $U$  is well defined.

<span id="page-19-0"></span>**Lemma 4.1** (continuous dependence on the boundary curve f [\[19\]](#page-26-6)) Let  $f \in C^2(\mathbb{R})$ be  $2\pi$ -periodic and  $u \in H_{S,qp}^1$  be the unique solution of the scattering problem [\(15\)](#page-6-3). Let  $K \subset D$  be compact. Then there exists  $\gamma > 0$  and  $C > 0$  both depending on  $k, u^i, f$ and K, such that for all  $2\pi$ -periodic  $r \in C^1(\mathbb{R})$  with  $||r - f||_{1,\infty} \leq \gamma$ , the unique solution  $u_r$  of the scattering problem corresponding to r satisfies

$$
||u_r - u||_{H^1(K)} \leq C||f - r||_{1,\infty}.
$$

Here,  $||q||_{1,\infty} := ||q||_{\infty} + ||q'||_{\infty}$  denotes the norm in  $C^1[0,\Lambda]$ , where  $\Lambda$  is the period of the scattered structures.

<span id="page-19-1"></span>**Lemma 4.2** (Pettis measurability theorem (Proposition 2.15 in [\[27\]](#page-26-7))). Let  $(\Omega, \mathcal{F}, \mu)$ be a complete  $\sigma$ -finite measure space. The following are equivalent for a function  $f : \Omega \to X$  (i) f is strongly measurable, (ii) f is measurable and  $\mu$ -essentially separably valued.

<span id="page-19-2"></span>**Lemma 4.3** (Bochner's Theorem (Proposition 2.16 in [\[27](#page-26-7)])). If  $f : \Omega \to X$  is a strongly measurable function, then f is Bochner integrable if and only if the scalar function  $\|f\|$  is integrable.

<span id="page-19-3"></span>**Lemma 4.4** (*U* is well defined) For  $f_0 \in \mathcal{C}$ , the solution U of the scattering problem exists, is unique, and depends continuously on  $f_0$ .

*Proof* For  $f_0 \in \mathcal{C}$ , the scattering problem is a deterministic case satisfying the condition in Theorem [2.1.](#page-5-0) Based on the previous work in [\[4\]](#page-24-0) on the existence and uniqueness, it follows from Theorem [2.1](#page-5-0) and Lemma [4.1](#page-19-0) that the solution  $U$  is well defined. defined.  $\Box$ 

Here the basic propositions on measure theory and Bochner spaces needed for the proof are omitted. See [\[9\]](#page-25-10) and [\[11\]](#page-25-11) for more details.

#### 4.2 Proof of Proposition [4.1](#page-18-1) and Proposition [4.2](#page-18-2)

In this section, we verify that the random periodic structure scattering problem has all necessary propositions required as in the general framework [\[24\]](#page-26-3). Proposition [4.1](#page-18-1) includes measurability and  $\mu$ -essentially separability of c, continuity of  $\tilde{a}$  and  $\tilde{G}$  and regularity of  $\tilde{a} \circ c$  and  $\tilde{G} \circ c$ . Proposition [4.2](#page-18-2) includes the necessary stability and uniqueness which hold almost surely. To be specific, under the assumpiton of resonance exclusion and stochastic regularity of the scattering surface, a variable transform is introduced and theory of Bochner spaces such as Pettis measurability theorm and Bochner's theorem is used to complete the proof.

First, we give the proof of Proposition [4.1.](#page-18-1)

*Proof of Proposition* [4.1](#page-18-1) Since each of f is a Lipschitz function, f is strongly measurable. By Pettis measurability theorem (Lemma  $4.2$ ), it follows that c defined by Definition [6](#page-4-1) is measurable and  $\mu$ -essentially separably valued, so properties of c in Proposition [4.1](#page-18-1) are satisfied.

In order to prove the continuity of  $\tilde{a}$  and  $\tilde{G}$ , we need to show that if  $(f_m) \to (f_0)$ in C then  $\tilde{a}(f_m) \to \tilde{a}(f_0)$  in  $B(H_{S,qp}^1(D), H_{S,qp}^1(D))$ , and similarly for  $\tilde{G}$ . Since

<span id="page-20-0"></span>
$$
\tilde{a}_{f_m}(v_m, \phi) = \tilde{G}_{f_m}(\phi),\tag{73}
$$

where

$$
\tilde{a}_{f_m} = \int_{D_{f_m}} \nabla v_m \cdot \nabla \bar{\phi} - k^2 \int_{D_{f_m}} v_m \bar{\phi} - \int_{\Gamma} T \tau v_m \overline{\tau \phi}, \quad \tilde{G}_{f_m} = \int_{\Gamma} g \overline{\tau \phi}, \tag{74}
$$

and

<span id="page-20-1"></span>
$$
\tilde{a}_{f_0}(v_0, \phi) = \tilde{G}_{f_0}(\phi),\tag{75}
$$

where

<span id="page-20-2"></span>
$$
\tilde{a}_{f_0} = \int_{D_{f_0}} \nabla v_0 \cdot \nabla \bar{\phi} - k^2 \int_{D_{f_0}} v_0 \bar{\phi} - \int_{\Gamma} T \tau v_0 \overline{\tau \phi}, \quad \tilde{G}_{f_0} = \int_{\Gamma} g \overline{\tau \phi}.
$$
 (76)

Our goal is to transform the volume integral in  $D_{f_m}$  into an integral over  $D_{f_0}$  by a suitable change of the variable  $x$ .

Choose  $\gamma_0 > 0$  such that

<span id="page-20-3"></span>
$$
\gamma_0 < \min\{\min\{x_2 : (x_1, x_2) \in D_{f_0}\} - f_0(x_1) : x_1 \in \mathbb{R}\},\tag{77}
$$

and a function  $\alpha \in C^1(\mathbb{R})$  with

<span id="page-20-4"></span>
$$
\alpha(t) = \begin{cases} 1, \ t \le \gamma_0/2 \\ 0, \ t \ge \gamma_0. \end{cases} . \tag{78}
$$

For  $f_m$  such that  $||f_m - f_0||_{1,\infty} \leq \gamma_0$ , define  $\mathcal{F} : D_{f_0} \to D_{f_m}$  by

$$
\mathcal{F}(y)=y_2+\alpha(y_2-f_0(y_1))[f_m(y_1)-f_0(y_1)]\hat{e}_2,\quad y\in D_{f_0},
$$

where  $\hat{e}_2 = (0, 1)^T$ . For  $||f_m - f_0||_{1,\infty} \leq \gamma$  with sufficiently small  $\gamma \leq \gamma_0$ , F is a diffeomorphism and  $\mathcal{F} = \mathcal{I}$  on K. The Jacobian  $J_{\mathcal{F}}$  of F satisfies  $J_{\mathcal{F}}(y) = \mathcal{I}$  +  $\mathcal{O}(\|f_m - f_0\|_{1,\infty})$  uniformly in  $y \in D_{f_0}$ . For the inverse Q of F, one also has  $J_{\mathcal{Q}}(y) =$  $\mathcal{I} + \mathcal{O}(\Vert f_m - f_0 \Vert_{1,\infty})$ . The change of variable  $x = \mathcal{F}(y)$  then transforms [\(73\)](#page-20-0) into

<span id="page-21-0"></span>
$$
\int_{D_{f_0}} \left[ \sum_{i,j=1}^2 b_{ij} \frac{\partial \hat{v}_m}{\partial y_i} \frac{\partial \overline{\hat{\phi}}}{\partial y_j} - k^2 \hat{v}_m \overline{\hat{\phi}} \right] \det(J_{\mathcal{F}}) dy - \int_{\Gamma} \overline{\tau \hat{\phi}} T \tau \hat{v}_m ds = \int_{\Gamma} g \overline{\tau \hat{\phi}} ds, \quad (79)
$$

where  $\hat{v}_m = v_m \circ \mathcal{F}, \hat{\phi} = \phi \circ \mathcal{F}$  both in  $H^1_{S,qp}(D_{f_0}),$  and

<span id="page-21-2"></span>
$$
b_{ij}(y) = \sum_{l=1}^{2} \frac{\partial Q_i(x)}{\partial x_l} \frac{\partial Q_j(x)}{\partial x_l} \Big|_{x = \mathcal{F}(y)}, \quad i, j = 1, 2. \tag{80}
$$

The left hand side of [\(79\)](#page-21-0) defines a sesquilinear form  $\tilde{a}_{f_m}(v, \phi)$  on  $H^1_{S,qp}(D_{f_0})$ . Since  $\det(J_{\mathcal{F}}) = 1+\mathcal{O}(\Vert f_m - f_0 \Vert_{1,\infty})$  and  $b_{ij} = \delta_{ij} + \mathcal{O}(\Vert f_m - f_0 \Vert_{1,\infty})$ , one concludes that

$$
|\tilde{a}_{f_m}(v,\phi)-\tilde{a}_{f_0}(v,\phi)| \le C \|f_m - f_0\|_{1,\infty} \|v\|_{H^1(D_{f_0})} \|\phi\|_{H^1(D_{f_0})} \text{ for all } v,\phi \in H^1_{S,qp}(D_{f_0}).
$$

Hence if  $(f_m) \to (f_0)$  in C, then  $\tilde{a}_{f_m} \to \tilde{a}_{f_0}$  in  $B(H_{S,qp}^1(D), H_{S,qp}^1(D))$ . For  $\tilde{G}$ , it follows from the definition [\(75\)](#page-20-1)-[\(76\)](#page-20-2), the right hand side of [\(79\)](#page-21-0),  $\hat{\phi} = \phi \circ \mathcal{F}$  and  $\phi = \phi \circ \mathcal{I}$  that if  $(f_m) \to (f_0)$  in C, then  $\tilde{G}_{f_m} \to \tilde{G}_{f_0}$  in  $H^1_{S,qp}(D)$ .

Since c is strongly measurable and the map  $\tilde{a}$  is continuous,  $\tilde{a} \circ c$  is strongly measurable. Recall that the operator T is continuous from  $H^{1/2}(\Gamma)$  to  $H^{-1/2}(\Gamma)$ [\[22\]](#page-26-8). For  $v \in H_{S,qp}^1(D)$ ,  $\phi \in H_{S,qp}^1(D)$ , one has

<span id="page-21-1"></span>
$$
\tilde{a}_{f(\omega)}(v,\phi) = \int_{D_{f(\omega)}} \nabla v \cdot \nabla \bar{\phi} - k^2 \int_{D_{f(\omega)}} v \bar{\phi} - \int_{\Gamma} T \tau v \overline{\tau \phi}.
$$
\n(81)

Same as above, transform the volume integral in  $D_{f(\omega)}$  into an integral over  $D_{f_0}$ . For  $f_1 = f(\omega)$  such that  $||f_1 - f_0||_{1,\infty} \leq \gamma_0$  (defined same as [\(77\)](#page-20-3)), define  $\mathcal{F} : D_{f_0} \to D_{f_1}$ by

$$
\mathcal{F}(y) = y_2 + \alpha(y_2 - f_0(y_1)) [f_1(y_1) - f_0(y_1)] \hat{e}_2, \quad y \in D_{f_0},
$$

where  $\hat{e}_2 = (0, 1)^T$  and  $\alpha$  is defined by [\(78\)](#page-20-4). For  $||f_1 - f_0||_{1, \infty} \leq \gamma$  with sufficiently small  $\gamma \leq \gamma_0$ , F is a diffeomorphism and  $\mathcal{F} = \mathcal{I}$  on K. The change of variable  $x = \mathcal{F}(y)$  then transforms [\(81\)](#page-21-1) into

$$
\tilde{a}_{f(\omega)}(\hat{v}, \phi) = \int_{D} \left[ \nabla \hat{v} \det(J_{\mathcal{F}}) B \nabla \bar{\phi} - k^2 (\det(J_{\mathcal{F}}) \hat{v} \bar{\phi}) \right] dy, \tag{82}
$$

where  $\hat{v} = v \circ \mathcal{F}, \hat{\phi} = \phi \circ \mathcal{F}$  both in  $H^1_{S,qp}(D_{f_0})$ , and  $B = (b_{ij})_{i,j=1,2}$  from [\(80\)](#page-21-2). Since  $\det(J_{\mathcal{F}}) = 1 + \mathcal{O}(\Vert f_m - f_0 \Vert_{1,\infty})$  and  $b_{ij} = \delta_{ij} + \mathcal{O}(\Vert f_m - f_0 \Vert_{1,\infty})$ , observe that the Cauchy-Schwarz inequality and properties of T imply that there exists  $C > 0$  such that

$$
|\tilde{a}_{f(\omega)}(\hat{v},\phi)| \le C \|f(\omega) - f_0\|_{1,\infty}^2 \|v\|_{H^1(D)} \|\phi\|_{H^1(D)}
$$
 for all  $v, \phi \in H^1_{S,qp}(D_{f_0}),$ 

and hence  $\tilde{a} \circ c \in L^{\infty}(\Omega; B(H^1_{S,qp}(D), H^1_{S,qp}(D))).$ 

Since c is strongly measurable and the map  $\tilde{G}$  is continuous,  $\tilde{G} \circ c$  is strongly measurable. It is clear that  $\|(\tilde{G} \circ c)(\omega)\|_{H^1_{S,qp}(D)} \leq \|g\|_{L^2(D)}$ , and thus  $\tilde{G} \circ c \in$  $L^2(\Omega; H^1_{S,qp}(D))$  since  $g \in L^2(\Omega; L^2(D)).$ 

Now prove Proposition [4.2.](#page-18-2)

*Proof of Proposition* [4.2](#page-18-2) For  $f_0 \in \mathcal{C}$ , it follows from Theorem [2.1](#page-5-0) and Lemma [4.1](#page-19-0) that the solution  $\mathcal{U}(f_0)$  of the variational problem exists, is unique, and has the kexplicit stability estimate [\(71\)](#page-18-3). Uniqueness almost surely holds immediately from Lemma [4.4.](#page-19-3)

For stability almost surely, choose  $C_1 = C'$  max  $\left\{\frac{b^2 k^2}{\sqrt{2}}\right\}$  $\left\{\frac{\partial^2 k^2}{\sqrt{\varepsilon}}, b^3 k^{\frac{5}{2}}\right\}$  with  $C' = \sup_{\Omega} C$ , C being the previous constant in [\(71\)](#page-18-3), and  $h_1 = ||g||_{L^2(\Gamma)}$ . It remains to show that  $C_1 h_1 \in L^1(\Omega)$ . We first show that  $C_1 h_1$  is measurable and then show that it lies in  $L^1(\Omega)$ .

To conclude  $C_1$  is measurable, use the fact that the product, sum and maximum of two measurable functions are measurable. Since g is measurable and the map  $g \mapsto$  $||g||_{L^2(\Gamma)}^2$  is clearly continuous,  $h_1$  is measurable. As the product of two measurable functions is measurable, it follows that  $C_1h_1$  is measurable.

Now show that  $C_1 h_1 \in L^1(\Omega)$ . Using the Cauchy-Schwarz inequality yields

<span id="page-22-0"></span>
$$
||C_1h_1||_{L^1(\Omega)} = \int_{\Omega} C_1(\omega)h_1(\omega) \, d\mathbb{P}(\omega) \le ||C_1||_{L^1(\Omega)} \, ||||g||_{L^2(\Gamma)}^2 ||_{L^1(\Omega)} < \infty. \tag{83}
$$

Therefore  $C_1h_1 \in L^1(\Omega)$  as required. Integrating [\(71\)](#page-18-3) in the probability space and using  $(83)$  obviously yield  $(14)$ .

#### 4.3 Proof of Theorem [2.2](#page-6-0)

Before the proof, the continuity of the solution operator  $\mathcal U$  is given in the following lemma.

<span id="page-22-1"></span>**Lemma 4.5** (Continuity of  $U$ ) For the scattering problem by a random periodic structure, the solution operator  $\mathcal{U}: \mathcal{C} \to H^1_{S,qp}(D)$  is continuous.

*Proof* Let  $f_0, f_1 \in \mathcal{C}$ , with  $\mathcal{U}(f_0) = u_0$  and  $\mathcal{U}(f_1) = u_1$ . Then for any  $v \in H^1_{S,qp}(D)$ ,

$$
\tilde{a}_{f_j}(u_j, v) = \tilde{G}_{f_j}(v), \ j = 0, 1.
$$

Since one has

$$
|\tilde{a}_{f_1}(u,\phi) - \tilde{a}_{f_0}(u,\phi)| \le C||f_1 - f_0||_{1,\infty} ||u||_{H^1(D_{f_0})} ||\phi||_{H^1(D_{f_0})} \text{ for all } u,\phi \in H^1_{S,qp}(D_{f_0})
$$

from the proof of Proposition [4.1,](#page-18-1) where both  $\tilde{a}_{f_1}(u, \phi)$  and  $\tilde{a}_{f_0}(u, \phi)$  are sesquilinear forms, one can applies the general perturbation theory of variational equation [\[18\]](#page-26-9) which yields

$$
||u_1 \circ \mathcal{F} - u_0||_{H^1(D_{f_0})} = ||\hat{u}_1 - u_0||_{H^1(D_{f_0})} \leq C||f_m - f_0||_{1,\infty},
$$

and thus, since  $\mathcal{F} = \mathcal{I}$  on  $D_{f_0}$ ,

$$
||u_1 - u_0||_{H^1(K)} \le C||f_1 - f_0||_{1,\infty}.
$$

Let  $u_d := u_0 - u_1$ . It can be deduced that  $u_d \to 0$  in  $H^1_{S,qp}(D_{f_0})$  as  $f_1 \to f_0$  in C. This ends the proof of this lemma.

Now we are ready to prove Theorem [2.2.](#page-6-0)

*Proof* Let  $u = U \circ c$  (which is well-defined by Lemma [4.4\)](#page-19-3). By construction,  $a_{c(\omega)}(u(\omega),v) = G_{c(\omega)}(v)$  for all  $v \in H^1_{S,qp}(D)$  almost surely. Since it follows from Assumption [2.2](#page-5-1) and Lemma [4.5](#page-22-1) that  $u$  is measurable, u solves the variational problem.

Under Assumption [2.2,](#page-5-1) continuity of  $\tilde{a}$  and  $\tilde{G}$ , regularity of  $\tilde{a} \circ c$  and  $\tilde{G} \circ c$ , and measurability and  $\mu$ -essentially separability of c hold by Proposition [4.1;](#page-18-1) stability a.s. and uniqueness a.s. hold by Proposition [4.2.](#page-18-2) Moreover, since there is no trapping cases for the scattering problem by a Lipschitz perfectly conductor, the nontrapping condition required in [\[24\]](#page-26-3) is naturally satisfied. Therefore the maps  $A$  and  $G$  (defined by [\(7\)](#page-4-3) and [\(9\)](#page-5-5)) are well-defined; there exists  $u \in L^2(\Omega; H^1_{S,qp}(D))$  being the solution to Problem [1;](#page-5-2) the solution to Problem [1](#page-5-2) is unique in  $L^2(\Omega; H^1_{S,qp}(D)).$ 

Combining the stability for the scattering problem by deterministic periodic structures in Theorem [2.1](#page-5-0) with integrability and measurability properties on stochastic quantities verified by Proposition [4.1](#page-18-1) and stability and uniqueness which hold almost surely given by Proposition [4.2](#page-18-2) yields the well-posedness and the k-explicit stability for the scattering problem by random periodic structures, which completes the proof of Theorem [2.2.](#page-6-0)

**Remark 4.2** If the random structure is (uncertainty) quantified using the Karhunen-Loève (KL) expansion as in [\[6\]](#page-25-8), then the random structure can be represented by the following Karhunen-Loève expansion

$$
f(\omega; x_1) = \tilde{f}(x_1) + \sum_{j=0}^{\infty} \sqrt{\lambda_j} \xi_j(\omega) \varphi_j(x_1),
$$

where  $\tilde{f}(x_1)$  is a  $\Lambda$ -periodic deterministic function, eigenvalues  $\{\lambda_j\}_{j=0}^{\infty}$  arranged in a descending order are corresponding to the orthonormal eigenfunctions  $\{\varphi_j\}_{j=0}^{\infty}$  of the covariance operator  $C_f$  and  $\{\xi_j\}_{j=0}^{\infty}$  is a random variable with zero mean and unit covariance. The covariance function [\[21](#page-26-10), [23](#page-26-11)] takes the following form

$$
c(x_1 - y_1) = \sigma^2 \exp(-\frac{|x_1 - y_1|^2}{l^2}), 0 < l \ll \Lambda,
$$

where  $\sigma$  is the root mean square of the surface and l is the correlation length. It follows that the KL expansion of the random process  $f(\omega; x_1)$  may be written as

$$
f(\omega; x_1) = \tilde{f}(x_1) + \sqrt{\lambda_0} \xi_0(\omega) \sqrt{\frac{1}{\Lambda}} + \sum_{j=1}^{\infty} \sqrt{\lambda_j} \left( \xi_{j,s}(\omega) \sqrt{\frac{2}{\Lambda}} \sin\left(\frac{2j\pi x_1}{\Lambda}\right) + \xi_{j,c}(\omega) \sqrt{\frac{2}{\Lambda}} \cos\left(\frac{2j\pi x_1}{\Lambda}\right) \right),
$$

where  $\xi_0$ ,  $\xi_{j,s}$  and  $\xi_{j,c}$  are independent and identically distributed random variables with zero mean and unit covariance.

Since the eigenvalues  $\{\lambda_j\}_{j=0}^{\infty}$  decrease exponentially, it is easy to show that the derivative of the random surface  $f(\omega; x_1)$  respect to  $x_1$  is bounded and there exists a positive constant C such that  $|f(\omega; x_1) - f(\omega; x_2)| \leq C |x_1 - x_2|$ . Thus the random surface  $f(\omega; x_1)$  in KL expansion is Lipschitz. Therefore, we have the results on the well-posedness of variational formulations of the Helmholtz equation on a random periodic structure proved in this paper, which verifies the well-posedness of the direct problem in  $[6]$ .

## <span id="page-24-4"></span>5 Conclusions

In this work, we have established stability results for the Helmholtz equation on deterministic and random periodic structures, respectively. Both estimates are explicit with respect to the wavenumber. To the authors' best knowledge, these are the first stability results explicit with respect to the wavenumber for the scattering problem in periodic structures. An interesting future direction is to apply the stability results for conducting convergence analysis of numerical methods for solving the scattering problems. Our techniques may be applicable to other (stochastic) scattering problems. In particular, another future direction is to conduct stability analysis for the electromagnetic scattering problems.

Acknowledgments. This work was supported in part by National Natural Science Foundation of China (11621101, U21A20425, 12071430, 12201404), a Key Laboratory of Zhejiang Province, and Postdoctoral Science Foundation of China (2021TQ0203).

## Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Data availability Data sharing is not applicable to this article as obviously no datasets were generated or analyzed during the current study.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## References

- <span id="page-24-5"></span>[1] Bao, G.: Finite element approximation of time harmonic waves in periodic structures. SIAM J. Numer. Anal.  $32(4)$ , 1155–1169 (1995). [https://doi.](https://doi.org/10.1137/0732053) [org/10.1137/0732053](https://doi.org/10.1137/0732053)
- <span id="page-24-1"></span>[2] Bao, G.: Numerical analysis of diffraction by periodic structures: TM polarization. Numer. Math. 75(1), 1–16 (1996). [https://doi.org/10.1007/](https://doi.org/10.1007/s002110050227) [s002110050227](https://doi.org/10.1007/s002110050227)
- <span id="page-24-2"></span>[3] Bao, G., Cowsar, L., Masters, W.: Mathematical Modeling in Optical Science. SIAM, Philadelphia (2001). [https://doi.org/10.1137/1.](https://doi.org/10.1137/1.9780898717594) [9780898717594](https://doi.org/10.1137/1.9780898717594)
- <span id="page-24-0"></span>[4] Bao, G., Dobson, D.C., Cox, J.A.: Mathematical studies in rigorous grating theory. J. Opt. Soc. Am. A 12(5), 1029–1042 (1995). [https://doi.org/](https://doi.org/10.1364/JOSAA.12.001029) [10.1364/JOSAA.12.001029](https://doi.org/10.1364/JOSAA.12.001029)
- <span id="page-24-3"></span>[5] Bao, G., Li, P.: Maxwell's Equations in Periodic Structures. Springer, Singapore (2021). <https://doi.org/10.1007/978-981-16-0061-6>

- <span id="page-25-8"></span>[6] Bao, G., Lin, Y., Xu, X.: Inverse scattering by a random periodic structure. SIAM J. Numer. Anal. 58(5), 2934–2952 (2020). [https://doi.org/10.](https://doi.org/10.1137/20M132167X) [1137/20M132167X](https://doi.org/10.1137/20M132167X)
- <span id="page-25-4"></span>[7] Bao, G., Yun, K.H.: Stability for the electromagnetic scattering from large cavities. Arch. Ration. Mech. Anal.  $220(3)$ , 1003-1044 (2016). [https://](https://doi.org/10.1007/s00205-015-0947-x) [doi.org/10.1007/s00205-015-0947-x](https://doi.org/10.1007/s00205-015-0947-x)
- <span id="page-25-3"></span>[8] Bao, G., Yun, K.H., Zhou, Z.: Stability of the scattering from a large electromagnetic cavity in two dimensions. SIAM J. Math. Anal. 44(1), 383–404 (2012). <https://doi.org/10.1137/110823791>
- <span id="page-25-10"></span>[9] Bogachev, V.I.: Measure Theory. Springer, Berlin (2007).
- <span id="page-25-0"></span>[10] Chandler-Wilde, S.N., Monk, P.: Existence, uniqueness, and variational methods for scattering by unbounded rough surfaces. SIAM Journal on Mathematical Analysis 37(2), 598–618 (2005). [https://doi.org/10.1137/](https://doi.org/10.1137/040615523) [040615523](https://doi.org/10.1137/040615523)
- <span id="page-25-11"></span>[11] Diestel, J., Uhl, J.J.: Vector Measures. American Mathematical Society, Providence (1977).
- <span id="page-25-5"></span>[12] Du, K., Li, B., Sun, W.: A numerical study on the stability of a class of Helmholtz problems. J. Comput. Phys. 287, 46–59 (2015). [https://doi.](https://doi.org/10.1016/j.jcp.2015.02.008) [org/10.1016/j.jcp.2015.02.008](https://doi.org/10.1016/j.jcp.2015.02.008)
- <span id="page-25-2"></span>[13] Esterhazy, S., Melenk, J.M.: On Stability of Discretizations of the Helmholtz Equation. In: I. Graham, T. Hou, O. Lakkis, R. Scheichl (Ed.) Numerical Analysis of Multiscale Problems. Lecture Notes in Computational Science and Engineering, Vol 83. Springer, Berlin (1998).
- <span id="page-25-9"></span>[14] Feng, X., Lin, J., Nicholls, D.P.: An efficient monte carlo-transformed field expansion method for electromagnetic wave scattering by random rough surfaces. Commun. Comput. Phys. 23, 685–705 (2018). [https://doi.org/](https://doi.org/10.4208/cicp.OA-2017-0041) [10.4208/cicp.OA-2017-0041](https://doi.org/10.4208/cicp.OA-2017-0041)
- <span id="page-25-6"></span>[15] Graham, I.G., Pembery, O.R., Spence, E.A.: The Helmholtz equation in heterogeneous media: A priori bounds, well-posedness, and resonances. J. Differ. Equ. 266(6), 2869–2923 (2019). [https://doi.org/10.1016/j.jde.](https://doi.org/10.1016/j.jde.2018.08.048) [2018.08.048](https://doi.org/10.1016/j.jde.2018.08.048)
- <span id="page-25-1"></span>[16] Hetmaniuk, U.: Stability estimates for a class of Helmholtz problems. Commun. Math. Sci. 5(3), 665–678 (2007). [https://doi.org/10.4310/CMS.](https://doi.org/10.4310/CMS.2007.v5.n3.a8) [2007.v5.n3.a8](https://doi.org/10.4310/CMS.2007.v5.n3.a8)
- <span id="page-25-7"></span>[17] Hiptmair, R., Scarabosio, L., Schillings, C., Schwab, C.: Large deformation shape uncertainty quantification in acoustic scattering. Adv. Comput.

Math. 44, 1475–1518 (2018). <https://doi.org/10.1007/s10444-018-9594-8>

- <span id="page-26-9"></span>[18] Kato, T.: Perturbation Theory for Linear Operators. Springer, Berlin (1976).
- <span id="page-26-6"></span>[19] Kirsch, A.: Diffraction by Periodic Structures. In: L. Pävärinta, E. Somersalo (Ed.) Inverse Problems in Mathematical Physics, 87-102. Springer, Berlin (1993).
- <span id="page-26-0"></span>[20] Lord, N.H., Mulholland, A.J.: Explicit error bounds for the  $\alpha$ -quasiperiodic Helmholtz problem. J. Opt. Soc. Am. A 30(10), 2111–2123 (2013). <https://doi.org/10.1364/JOSAA.30.002111>
- <span id="page-26-10"></span>[21] Maradudin, A.A. (ed.): Light Scattering and Nanoscale Surface Roughness. Springer, New York (2007).
- <span id="page-26-8"></span>[22] Nedelec, J.C.: Acoustic and Electromagnetic Equations: Integral Representations for Harmonic Problems. Springer, New York (2001).
- <span id="page-26-11"></span>[23] Ogilvy, J.A.: Theory of Wave Scattering from Random Rough Surfaces. Adam Hilger, Bristol (1991).
- <span id="page-26-3"></span>[24] Pembery, O.R., Spence, E.A.: The Helmholtz equation in random media: Well-posedness and a priori bounds. SIAM/ASA J. Uncertain. Quantif. 8(1), 58–87 (2020). <https://doi.org/10.1137/18M119327X>
- <span id="page-26-1"></span>[25] Petit, R.: Electromagnetic Theory of Gratings. Springer, Heidelberg (1980). <https://doi.org/10.1007/978-3-642-81500-3>
- <span id="page-26-4"></span>[26] Rico-García, J.M., Sanchez-Brea, L.M.: Binary gratings with random heights. Appl. Opt. 48(16), 3062–3069 (2009). [https://doi.org/10.1364/](https://doi.org/10.1364/AO.48.003062) [AO.48.003062](https://doi.org/10.1364/AO.48.003062)
- <span id="page-26-7"></span>[27] Ryan, R.A.: Introduction to Tensor Products of Banach Spaces. Springer, London (2002).
- <span id="page-26-2"></span>[28] Spence, E.A.: Wavenumber-explicit bounds in time-harmonic acoustic scattering. SIAM J. Math. Anal.  $46(4)$ , 2987–3024 (2014). [https://doi.](https://doi.org/10.1137/060662575) [org/10.1137/060662575](https://doi.org/10.1137/060662575)
- <span id="page-26-5"></span>[29] Xiu, D., Tartakovsky, D.M.: Numerical methods for differential equations in random domains. SIAM J. Sci. Comput. 28(3), 1167–1185 (2006). <https://doi.org/10.1137/040613160>