

Regularity for a geometrically nonlinear flat Cosserat micropolar membrane shell with curvature

Andreas Gastel¹ and Patrizio Neff²

November 22, 2022

Abstract

We consider the rigorously derived thin shell membrane Γ -limit of a three-dimensional isotropic geometrically nonlinear Cosserat micropolar model and deduce full interior regularity of both the midsurface deformation $m: \omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$ and the orthogonal microrotation tensor field $R: \omega \subset \mathbb{R}^2 \rightarrow \text{SO}(3)$. The only further structural assumption is that the curvature energy depends solely on the uni-constant isotropic Dirichlet type energy term $|DR|^2$. We use Rivière's regularity techniques of harmonic map type systems for our system which couples harmonic maps to $\text{SO}(3)$ with a linear equation for m . The additional coupling term in the harmonic map equation is of critical integrability and can only be handled because of its special structure.

Key words: flat shell, membrane, Γ -limit, Hölder regularity, Cosserat surface, Cosserat shell, micropolar shell, harmonic maps, $\text{SO}(3)$, graphen, generalized continua.

AMS 2010 subject classification: 58E20, 74G40, 74B20, 35J35

Contents

1	Introduction	2
1.1	Regularity background and setting of the problem	2
1.2	Engineering background and application	3
2	Notation	5
3	Three-dimensional geometrically nonlinear isotropic Cosserat model	6
3.1	Connections to the Oseen-Frank energy in nematic liquid crystals	7
4	Dimensional descent towards a membrane model	8
4.1	Membrane Γ -limit	8
4.2	Alternative engineering ad-hoc dimensional descent	11
5	The two-dimensional Euler-Lagrange equations	12
6	Regularity	15
6.1	Hölder regularity	15
6.2	Higher regularity	20
6.3	Body forces	21
6.4	Remarks on a special case	23

¹Corresponding author: Andreas Gastel, Head of Lehrstuhl für Geometrische Analysis, Fakultät für Mathematik, Universität Duisburg-Essen, Thea-Leymann Str. 9, 45127 Essen, Germany; email: andreas.gastel@uni-due.de

²Patrizio Neff, Head of Lehrstuhl für Nichtlineare Analysis und Modellierung, Fakultät für Mathematik, Universität Duisburg-Essen, Thea-Leymann Str. 9, 45127 Essen, Germany; email: patrizio.neff@uni-due.de

7	Conclusion and open problems	24
	References	27
A	Appendix	27
A.1	Three-dimensional Euler-Lagrange equations in dislocation tensor format	27
A.2	Two-dimensional Euler-Lagrange equations: alternative derivation	28
A.3	Lifting to the Δ -operator	30
A.4	A glimpse on a Reissner-Mindlin type flat membrane shell model	31
A.5	Numerical experiments	32

1 Introduction

1.1 Regularity background and setting of the problem

This paper is a contribution to the wide field of regularity theory of harmonic map type equations. Driven by the application to geometrically nonlinear flat Cosserat shell models, we extend known regularity results to a system that couples a harmonic map equation with another uniformly elliptic equation. The system we consider is of the form

$$\operatorname{Div} S(Dm, R) = 0, \tag{1.1}$$

$$\Delta R - \Omega_R \cdot DR - \operatorname{skew}(Dm \circ S(Dm, R))R = 0. \tag{1.2}$$

The unknown functions here are the midsurface deformation $m \in W^{1,2}(\omega, \mathbb{R}^3)$ and the microrotation $R \in W^{1,2}(\omega, \operatorname{SO}(3))$, while $\omega \subset \mathbb{R}^2$ is a smooth domain. Moreover, there are functions $\Omega_R \in L^2(\omega, (\mathbb{R}^2)^* \otimes \mathfrak{so}(3))$ and the force stress tensor $S(Dm, R) \in L^2(\omega, \mathbb{R}^{3 \times 2})$ involved, and \circ is some bilinear product explained later. The function Ω_R is the same that makes

$$\Delta R - \Omega_R \cdot DR = 0 \tag{1.3}$$

the harmonic map equation for harmonic mappings to $\operatorname{SO}(3) \subset \mathbb{R}^{3 \times 3}$. The theory of harmonic map equations of 2-dimensional domains (to any sufficiently smooth compact target manifold, here $\operatorname{SO}(3)$) has a long history. It has been proven in 1948 by Morrey [44] that minimizing weakly harmonic maps are smooth. In 1981, Grüter [30] generalized that to conformal weakly harmonic maps, and then in 1984 Schoen [69] to stationary ones. The regularity proof for general weakly harmonic maps was then found in 1990 by Hélein [32] [33]. (Note that in our case, the target manifold $\operatorname{SO}(3)$ is a Lie group, and in this case the harmonic map problem has a lot of interesting extra structure, many aspects of which are covered in Hélein's book [34].) Later, in 2007, Rivière [63] revisited harmonic map type equations and asked for which Ω_R all weak solutions of (1.3) on a two-dimensional domain are smooth. It turned out that Ω_R need not come from the harmonic map equation (in which case it can be seen as the anti-symmetrized tensor derived from the second fundamental form of the target manifold), but for the regularity result only the skew-symmetry of Ω_R is needed. This gave deeper insight in the structures necessary to have regularity results, and it is Rivière's philosophy that we rely upon.

It should be pointed out that, with Ω_R and DR being in L^2 , the nonlinear term $\Omega_R \cdot DR$ in (1.3) is only in L^1 , and if it would not have any further structure, it would be difficult to start with any regularity theory, due to the lack of an L^p -theory working for $p = 1$. But it turns out that the product $\Omega_R \cdot DR$, after a suitable gauge transformation, is the sum of products of divergence-free vector fields and gradients in L^2 , which is known to be in the Hardy space \mathcal{H}^1 rather than L^1 . This little bit of extra regularity is enough to perform regularity theory.

Now let us have a look at our equation (1.2). Compared with (1.3), it has an extra term $\operatorname{skew}(Dm \circ S(Dm, R))R$, and again, with $DR \in L^2$, $S(Dm, R) \in L^2$, and $R \in L^\infty$, this has only L^1 -integrability. But once more, DR is a gradient, and $S(Dm, R)$ is divergence free due to equation (1.1). This time, we have the product of a gradient Dm , a divergence free vector field $S(Dm, R)$, and a bounded function R . Based on a

crucial estimate by Coifman, Lion, Meyer and Semmes [14], Rivière and Struwe [64] were able to handle such products in their work on partial regularity in dimensions ≥ 3 . They encountered such products in the course of their proof for the equation (1.3) without any extra terms, and we can modify their arguments to handle our extra term from the coupling. The handling of the first equation, which is linear in m with some right-hand side, is easier, in principle. But we have to do the iteration procedure for both equations simultaneously in the proof of Hölder continuity, resulting in some technicalities. Once we have that, classical Schauder theory helps with the higher regularity of m , while for the second equation controlling the smoothness of R , we still need some machinery.

1.2 Engineering background and application

The Cosserat model is one of the best known generalized continuum models [13]. It assumes that material points can undergo translation, described by the standard deformation map $\varphi: \mathcal{U} \rightarrow \mathbb{R}^3$ and independent micro rotations described by the orthogonal tensor field $R: \mathcal{U} \rightarrow \text{SO}(3)$, where $\mathcal{U} \subset \mathbb{R}^3$ describes the smooth reference configuration of the material. Therefore, the geometrically nonlinear Cosserat model induces immediately the Lie-group structure on the configuration space $\mathbb{R}^3 \times \text{SO}(3)$.

Both fields are coupled in the assumed elastic energy $W = W(D\varphi, R, DR)$ and the static Cosserat model appears as a two-field minimization problem which is automatically geometrically nonlinear due to the presence of the non-abelian rotation group $\text{SO}(3)$. Material frame-indifference (objectivity) dictates left-invariance of the Lagrangian W under the action of $\text{SO}(3)$ and material symmetry (here isotropy) implies right-invariance under action of $\text{SO}(3)$.

In the early 20th century the Cosserat brothers E. and F. Cosserat introduced this model in its full geometrically nonlinear splendor [17] in a bold attempt to unify field theories embracing mechanics, optics and electrodynamics through a common principal of least action. They used the invariance of the energy under Euclidean transformations [15, 4] to deduce the correct form of the energy $W = W(R^T D\varphi, R^T \partial_x R, R^T \partial_y R, R^T \partial_z R)$ and to derive the equations of balance of forces (variations w.r.t the deformation φ , the force-stress tensor may lose symmetry [56]) and balance of angular momentum (variations w.r.t. rotations R). The Cosserat brothers did not provide, however, any specific constitutive form of the energy since they were not interested in applications.

While the appearance of an additional rotational field R for describing the elastic response of bulk material is requiring getting used to, such an appearance is most natural in the case of shell-theory. There, the Frenet-Darboux trièdre [16] (trièdre caché in the terminology of the Cosserats, trihedron) naturally plays a role and it is no big step to assume that this orthogonal field is supposed to be kinematically independent of the former (trièdre mobile). Hence the Cosserat approach [16]; the independent rotation field R describes the rotations of the cross-sections of the shell (including in-plane drill rotations about the normal n_m to the midsurface m) and these cross-sections are all allowed to shear with respect to the normal of the midsurface ($Re_3 \neq n_m$).

On this basis, very efficient ad-hoc Cosserat shell-models have been introduced, see e.g. [2, 3]. A special case of these shell models is the family of Reissner-Mindlin shells in which the in-plane rotations are discarded (no drill energy) [37] and one is left with a one director theory [39]¹. Upon identifying/constraining the trièdre mobile with the trièdre caché (microrotation equals continuum rotation, Cosserat couple modulus $\mu_c \rightarrow \infty$), canonical shell models of Kirchhoff-Love type emerge [27]. However, engineers would often prefer the Cosserat shell models since these yield nonlinear balance equations of second order [67, 72, 73, 35, 62, 7].

The precise derivation of Cosserat shell models may proceed in several different ways: integration of equilibrium equations through the thickness [18, 62], direct modeling as a two-dimensional directed surface [2, 3, 29], or the derivation approach, which starts from a three-dimensional variational problem and introduces certain assumptions for the deformation behavior through the thickness. The second author has introduced this derivation procedure based on the geometrically nonlinear Cosserat model in his habilitation thesis [50,

¹One director geometrically nonlinear, physically linear Reissner-Mindlin shells are typically not well-posed, since the membrane stretch energy part depends quadratically on $Dm^T Dm - \mathbb{1}_2$, which is not rank-one elliptic in the compression regime, for a detailed exposition, see the Appendix and [57].




Figure 1: The mapping $m: \omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$ describes the deformation of the flat midsurface $\omega \subset \mathbb{R}^2$. The Frenet-Darboux frame (in blue, trièdre caché) is tangent to the midsurface. The independent frame mapped by $R \in \text{SO}(3)$ is the trièdre mobile (in red, not necessary tangent to the midsurface). Both fields m and R are coupled in the variational problem.

48]. Lastly, there is the "ansatz-free" method of Γ -convergence [12, 11, 8] (while letting the thickness h tend to zero) to perform the dimensional descent.

In this method, one needs to choose an energy scaling regime and obtains typically either membrane or bending like theories [21, 39, 40, 41] when starting from classical finite strain elasticity [21, 22, 23]. However, the Γ -limit membrane model [39, 40] has a serious shortcoming which is connected to the necessary relaxation step: it does not predict any resistance against compression and averages out the expected fine scale wrinkling response. The situation is strikingly different when starting from a three-dimensional Cosserat model, as done in [51]. This is true since the bulk-Cosserat model already features a curvature term (derivatives of R) which "survives" the membrane scaling.

The Cosserat membrane Γ -limit with remaining curvature effects can be used as an effective surrogate model to describe ultra thin graphen mono-layers. Graphen is the name given to a single atomic layer of carbon atoms tightly packed into a two-dimensional honeycomb lattice (see Figure 2). It can be wrapped up to form fullerenes, rolled into nanotubes [74] or stacked into graphite. It's stiffness properties are extreme. Such a graphen layer has resistance against in-plane stretch and curvature changes but it's thickness is so small, that a classical membrane-bending model (where the bending terms scale with h^3 while the membrane terms with h) is clearly insufficient. It is simply impossible to speak about the "thickness" of graphen in a classical continuum framework. Researchers then usually resort to introducing an "effective bending rigidity" in order to apply concepts from classical shell theory. This can be completely avoided in the Cosserat membrane model.

In this paper we will consider, for the first time, the challenging regularity questions for the flat shell Cosserat membrane Γ -limit. To the best knowledge of the authors, such a regularity investigation for the flat Cosserat membrane shell has never been undertaken. Two recent previous contributions consider the regularity issue for the geometrically isotropic nonlinear Cosserat bulk equations [24, 42], both times restricting attention to the uni-constant Dirichlet curvature energy $|DR|^2$, leading to a ΔR -term in the Euler-Lagrange equations and allowing the sophisticated techniques for harmonic map type systems to be used.

This paper is now structured as follows. After this introduction and the introduction of our notation, in Section 3 we will introduce the three-dimensional isotropic Cosserat model, together with a short discussion of suitable representations for the curvature term. Following, in Section 4, we briefly describe the dimensional descent towards a membrane shell, juxtaposing the result of the Γ -limit procedure and a formal engineering approach. In Section 5 we introduce the final two-dimensional Cosserat membrane shell model together with some pertinent notations and simplifications. The remainder of the paper is devoted to showing the interior Hölder regularity of these weak solutions. In the appendix we gather further useful calculations like the three-dimensional Euler-Lagrange equations in dislocation tensor format, we present a more engineering oriented derivation of the two-dimensional Euler-Lagrange equations and give a glimpse on a related Reissner-




Figure 2: Deformed graphen mono atomic-layer resisting against in-plane stretches (membrane effects) and curvature. Classical continuum models are not any more suitable, since there is no tangible thickness, c.f. [74]. Graphen is thought to be the strongest among all known materials. Nevertheless it is soft in the sense that it can be easily bent due to its one atom thin nature.

Mindlin model. Finally, we show some numerical experiments of the flat Cosserat membrane shell model in compression.

2 Notation

Let $a, b \in \mathbb{R}^3$. We denote the scalar product on \mathbb{R}^3 with $\langle a, b \rangle_{\mathbb{R}^3}$ and the associated vector norm with $|a|_{\mathbb{R}^3}^2 = \langle a, a \rangle_{\mathbb{R}^3}$. The set of real-valued 3×3 second order tensors is denoted by $\mathbb{R}^{3 \times 3}$. The standard Euclidean scalar product on $\mathbb{R}^{3 \times 3}$ is given by $\langle X, Y \rangle_{\mathbb{R}^{3 \times 3}} = \text{tr}(XY^T)$, and the associated norm is $|X|^2 = \langle X, X \rangle_{\mathbb{R}^{3 \times 3}}$. If $\mathbb{1}_3$ denotes the identity matrix in $\mathbb{R}^{3 \times 3}$, we have $\text{tr}(X) = \langle X, \mathbb{1}_3 \rangle$. For an arbitrary matrix $X \in \mathbb{R}^{3 \times 3}$ we define $\text{sym}(X) = \frac{1}{2}(X + X^T)$ and $\text{skew}(X) = \frac{1}{2}(X - X^T)$ as the symmetric and skew-symmetric parts, respectively and the trace free deviatoric part is defined as $\text{dev } X = X - \frac{1}{n} \text{tr}(X) \mathbb{1}_n$, for all $X \in \mathbb{R}^{n \times n}$. We let $\text{Sym}(n)$ and $\text{Sym}^+(n)$ denote the symmetric and positive definite symmetric tensors, respectively. The Lie-algebra of skewsymmetric matrices is denoted by $\mathfrak{so}(3) := \{X \in \mathbb{R}^{3 \times 3} \mid X^T = -X\}$ and the Lie-algebra of traceless tensors is defined by $\mathfrak{sl}(3) := \{X \in \mathbb{R}^{3 \times 3} \mid \text{tr}(X) = 0\}$. We consider the orthogonal decomposition $X = \text{dev sym } X + \text{skew } X + \frac{1}{3} \text{tr}(X) \cdot \mathbb{1}_3 = \text{sym } X + \text{skew } X$. The *canonical identification* of $\mathfrak{so}(3)$ and \mathbb{R}^3 is given by $\text{axl}: \mathfrak{so}(3) \rightarrow \mathbb{R}^3$ and its inverse $\text{Anti}: \mathbb{R}^3 \rightarrow \mathfrak{so}(3)$. We note the following properties

$$\underbrace{\text{axl} \begin{pmatrix} 0 & \alpha & \beta \\ -\alpha & 0 & \gamma \\ -\beta & -\gamma & 0 \end{pmatrix}}_{=A} := \begin{pmatrix} -\gamma \\ \beta \\ -\alpha \end{pmatrix}, \quad |A|_{\mathbb{R}^{3 \times 3}}^2 = 2 | \text{axl } A |_{\mathbb{R}^3}^2, \quad A v = \text{axl}(A) \times v. \quad (2.1)$$

and

$$\text{Anti} \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} := \begin{pmatrix} 0 & -v_3 & v_2 \\ v_3 & 0 & -v_1 \\ -v_2 & v_1 & 0 \end{pmatrix} \in \mathfrak{so}(3). \quad (2.2)$$

A matrix having the three column vectors R_1, R_2, R_3 will be written sometimes as $R = (R_1 \mid R_2 \mid R_3) \in \mathbb{R}^{3 \times 3}$. The matrix Curl and matrix Div are defined row-wise as

$$\text{Curl } R = \begin{pmatrix} \text{curl}(R^T e_1) \\ \text{curl}(R^T e_2) \\ \text{curl}(R^T e_3) \end{pmatrix} \in \mathbb{R}^{3 \times 3}, \quad \text{Div } R = \begin{pmatrix} \text{div}(R^T e_1) \\ \text{div}(R^T e_2) \\ \text{div}(R^T e_3) \end{pmatrix}. \quad (2.3)$$

For $\varphi \in C^1(\mathcal{U}, \mathbb{R}^3)$ and for every vector $(x, y, z) \in \mathbb{R}^3$, we write

$$D\varphi = \begin{pmatrix} \varphi_{1,x} & \varphi_{1,y} & \varphi_{1,z} \\ \varphi_{2,x} & \varphi_{2,y} & \varphi_{2,z} \\ \varphi_{3,x} & \varphi_{3,y} & \varphi_{3,z} \end{pmatrix} = (\partial_x \varphi | \partial_y \varphi | \partial_z \varphi). \quad (2.4)$$

The mapping $m: \omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$ will always denote the deformation of the midsurface ω and we write

$$Dm = \begin{pmatrix} m_{1,x} & m_{1,y} \\ m_{2,x} & m_{2,y} \\ m_{3,x} & m_{3,y} \end{pmatrix} = (\partial_x m | \partial_y m), \quad D^\perp m = \begin{pmatrix} -m_{1,y} & m_{1,x} \\ -m_{2,y} & m_{2,x} \\ -m_{3,y} & m_{3,x} \end{pmatrix} = (-\partial_y m | \partial_x m). \quad (2.5)$$

Moreover, we will use the notations

$$\text{Div}(A_1 | A_2) = \partial_x A_1 + \partial_y A_2, \quad \text{Div}^\perp(A_1 | A_2) = \partial_x A_2 - \partial_y A_1, \quad (2.6)$$

where A_1, A_2 may be numbers-, vector-, or matrix-valued functions on ω of the same type. Note that it is also customary to write Curl instead of Div^\perp , but the latter underscores the symmetry of (D, Div) with $(D^\perp, \text{Div}^\perp)$, hence we reserve Curl for three-dimensional domains.

We assume that $h > 0$ with $h \ll 1$. The three-dimensional flat thin domain $\mathcal{U}_h \subset \mathbb{R}^3$ is introduced as

$$\mathcal{U}_h := \omega \times \left[-\frac{h}{2}, \frac{h}{2}\right], \quad \omega \subset \mathbb{R}^2. \quad (2.7)$$

We also need to define the projection operator on the first two columns

$$\pi_{12}: \mathbb{R}^{3 \times 3} \rightarrow \mathbb{R}^{3 \times 2}, \quad \pi_{12}(X) = \pi_{12}(X_1 | X_2 | X_3) = (X_1 | X_2) = \begin{pmatrix} X_{11} & X_{12} \\ X_{21} & X_{22} \\ X_{31} & X_{32} \end{pmatrix}, \quad (2.8)$$

and the operator

$$\begin{aligned} \mathbb{P}_{\mu, \mu_c, \kappa}: \mathbb{R}^{3 \times 3} &\rightarrow \{\mathfrak{sl}(3) \cap \text{Sym}(3)\} \oplus \mathfrak{so}(3) \oplus \mathbb{R} \cdot \mathbb{1}_3, \\ \mathbb{P}_{\mu, \mu_c, \kappa}(X) &= \mathbb{P}(X) = \sqrt{\mu} \text{dev sym } X + \sqrt{\mu_c} \text{skew } X + \frac{\sqrt{\kappa}}{3} \text{tr}(X) \cdot \mathbb{1}_3, \\ \mathbb{P}^* \mathbb{P}(X) &= \mathbb{P}^2(X) = \mu \text{dev sym } X + \mu_c \text{skew } X + \frac{\sqrt{\kappa}}{3} \text{tr}(X) \mathbb{1}_3, \quad \mathbb{P}^* = \mathbb{P}. \end{aligned} \quad (2.9)$$

3 Three-dimensional geometrically nonlinear isotropic Cosserat model

The underlying three-dimensional isotropic Cosserat model can be described in terms of the standard deformation mapping $\varphi: \mathcal{U} \subset \mathbb{R}^3 \rightarrow \mathbb{R}^3$ and an additional orthogonal microrotation tensor $R: \mathcal{U} \subset \mathbb{R}^3 \rightarrow \text{SO}(3)$.

The goal is to find a minimiser of the following isotropic energy

$$\begin{aligned} E^{3D}(\varphi, R) &= \int_{\mathcal{U}} \mu |\text{dev sym}(R^T D\varphi - \mathbb{1}_3)|^2 + \mu_c |\text{skew}(R^T D\varphi - \mathbb{1}_3)|^2 + \frac{\kappa_{3D}}{2} \text{tr}(R^T D\varphi - \mathbb{1}_3)^2 \\ &\quad + \mu \frac{L_c^2}{2} \left(a_1 |\text{dev sym } R^T \text{Curl } R|^2 + a_2 |\text{skew } R^T \text{Curl } R|^2 + \frac{a_3}{3} \text{tr}(R^T \text{Curl } R)^2 \right) dx \\ &= \int_{\mathcal{U}} \mu |\text{sym}(R^T D\varphi - \mathbb{1}_3)|^2 + \mu_c |\text{skew}(R^T D\varphi - \mathbb{1}_3)|^2 + \frac{\lambda}{2} \text{tr}(R^T D\varphi - \mathbb{1}_3)^2 \\ &\quad + \mu \frac{L_c^2}{2} \left(a_1 |\text{dev sym } R^T \text{Curl } R|^2 + a_2 |\text{skew } R^T \text{Curl } R|^2 + \frac{a_3}{3} \text{tr}(R^T \text{Curl } R)^2 \right) dx \\ &= \int_{\mathcal{U}} W_{\text{mp}}(R^T D\varphi) + W_{\text{disloc}}^{3D}(R^T \text{Curl } R) dx \rightarrow \min \quad \text{w.r.t } (\varphi, R). \end{aligned} \quad (3.1)$$

The problem will be supplemented by Dirichlet boundary conditions for the deformation φ but the microrotations R can be left free. Here, $\mu > 0$ is the standard elastic shear modulus, $\kappa_{3D} = \frac{3\lambda+2\mu}{3} > 0$ is the three dimensional elastic bulk modulus (with λ the second elastic Lamé parameter) and $\mu_c \geq 0$ is the so-called Cosserat couple modulus, a_1, a_2, a_3 are non-dimensional non-negative weights and $L_c > 0$ is a characteristic length. The energy (3.1) is the most general isotropic quadratic representation for the Cosserat model in terms of the nonsymmetric Biot type stretch tensor $\bar{U} = R^T D\varphi$ (first Cosserat deformation tensor [17]) and the curvature measure $R^T \text{Curl } R$ (physically linear, small strain, but geometrically nonlinear). We call

$$\boldsymbol{\alpha} := R^T \text{Curl } R, \quad (3.2)$$

the *second order* dislocation density tensor [10]. Due to the orthogonality of dev sym, skew and $\text{tr}(\cdot)\mathbb{1}_3$, the curvature energy provides a complete control of

$$|\boldsymbol{\alpha}|^2 = |R^T \text{Curl } R|^2 \quad \text{provided} \quad a_1, a_2, a_3 > 0. \quad (3.3)$$

For example, we can express the uni-constant isotropic curvature term

$$\begin{aligned} |DR|_{\mathbb{R}^{3 \times 3 \times 3}}^2 &= |R^T DR|_{\mathbb{R}^{3 \times 3 \times 3}}^2 = |R^T \partial_x R|_{\mathbb{R}^{3 \times 3}}^2 + |R^T \partial_y R|_{\mathbb{R}^{3 \times 3}}^2 + |R^T \partial_z R|_{\mathbb{R}^{3 \times 3}}^2 \\ &= 1 \cdot |\text{dev sym } R^T \text{Curl } R|_{\mathbb{R}^{3 \times 3}}^2 + 1 \cdot |\text{skew } R^T \text{Curl } R|_{\mathbb{R}^{3 \times 3}}^2 + \frac{1}{12} \cdot \text{tr}(R^T \text{Curl } R)^2 = |\mathbb{P}_{1,1,\frac{1}{12}}(\boldsymbol{\alpha})|^2, \end{aligned} \quad (3.4)$$

where we have used (4.4) and $|\boldsymbol{\Gamma}|^2 = |\text{axl}(R^T \partial_x R)|^2 + |\text{axl}(R^T \partial_y R)|^2 + |\text{axl}(R^T \partial_z R)|^2$ together with $2|\text{axl}(A)|_{\mathbb{R}^3}^2 = |A|_{\mathbb{R}^{3 \times 3}}^2$. Using the result in [60]

$$|\text{Curl } R|_{\mathbb{R}^{3 \times 3}}^2 \geq c^+ |DR|_{\mathbb{R}^{3 \times 3 \times 3}}^2, \quad (3.5)$$

shows that (3.1) controls DR in $L^2(\mathcal{U}, \mathbb{R}^{3 \times 3 \times 3})$.

In this setting, the minimization problem is strictly convex in the strain and curvature measures $(\bar{U}, \boldsymbol{\alpha})$ but highly non-convex w.r.t (φ, R) . Existence of minimizers for (3.1) with $\mu_c > 0$ has been shown first in [49], see also [19, 43, 53, 49, 38, 10, 51]. The partial regularity of minimizers/stationary solutions is investigated in [24, 42] under additional assumptions. Note also that in [24], the first author gives an example of a solution that exhibits a point singularity.

The Cosserat couple modulus μ_c controls the deviation of the microrotation R from the continuum rotation $\text{polar}(D\varphi)$ in the polar decomposition of $D\varphi = \text{polar}(D\varphi) \cdot \sqrt{D\varphi^T D\varphi}$, cf. [59].

For $\mu_c \rightarrow \infty$ the constraint $R = \text{polar}(D\varphi)$ is generated and the model would turn into a Toupin couple stress model.

3.1 Connections to the Oseen-Frank energy in nematic liquid crystals

In nematic liquid crystals one considers the unit-director field $n: \mathcal{U} \subset \mathbb{R}^3 \rightarrow \mathbb{S}^2$, minimizing the three-parameter frame-indifferent "curvature energy" [71]

$$\int_{\mathcal{U}} \frac{1}{2} K_1 |\text{div } n|^2 + \frac{1}{2} K_2 |\langle n, \text{curl } n \rangle|^2 + \frac{1}{2} K_3 |n \times \text{curl } n|^2 \, dx. \quad (3.6)$$

The uni-constant approximation $K_1 = K_2 = K_3$ leads to the Dirichlet type integral²

$$\int_{\mathcal{U}} \frac{1}{2} K_1 |Dn|^2 \, dx. \quad (3.8)$$

²For this, we note the identity (see, [5] eq (2.5) and [1] eq (2.6))

$$\text{tr}(Dv)^2 + \langle v, \text{curl } v \rangle^2 + |v \times \text{curl } v|^2 = |Dv|^2 + (|v|^2 - 1) |\text{curl } v|^2, \quad (3.7)$$

valid for all sufficiently smooth vector fields $v: \mathcal{U} \subset \mathbb{R}^3 \rightarrow \mathbb{R}^3$.

The corresponding Euler-Lagrange equations for the uni-constant case are (see e.g. [1])

$$\Delta n = -|Dn|^2 \cdot n, \quad (3.9)$$

see equation (A.56) for a self-contained derivation. Since (3.8) and (3.9) are just the energy and Euler-Lagrange equation for harmonic maps to spheres, all regularity theorems for harmonic maps apply. In the three dimensional case, minimizers are smooth up to a discrete set of singularities. Stationary solutions have a co-dimension 1 singular set. In the two dimensional case, all weak solutions of (3.9) are smooth, see Section (1.1) for the literature on this.

For K_1, K_2, K_3 positive and different, any minimizer to (3.6) is smooth except for a closed set of Hausdorff dimension strictly less than 1, cf. [31]. Ball and Bedford [6] consider the sublinear regime $|Dn|^q$, $1 < q < 2$.

4 Dimensional descent towards a membrane model

4.1 Membrane Γ -limit

We are interested in a situation, where the reference configuration is flat with uniform shell thickness $h > 0$, i.e. the reference configuration is taken to be of the form (see Figure 3)

$$\mathcal{U}_h = \omega \times \left[-\frac{h}{2}, \frac{h}{2}\right], \quad \omega \subset \mathbb{R}^2. \quad (4.1)$$

The goal is to derive a limit two-dimensional problem, posed over the referential midsurface $\omega \subset \mathbb{R}^2$, as




Figure 3: Process of dimensional reduction. Flat reference configuration with height h and deformed configuration.

$h \rightarrow 0$. This has been achieved in [58] based on Γ -convergence arguments and using the nonlinear membrane scaling. We say that the dimensionally reduced model is a membrane, since no dedicated bending terms appear in the problem.

However, since the Cosserat model already includes curvature terms (those depending on space derivatives DR), these curvature terms "survive" in the Γ -limit procedure and scale with h , while canonical bending terms scale with h^3 . This sets the Cosserat membrane model apart from more canonical membrane models [54].

For the Γ -limit procedure it is useful to re-express the curvature energy from (3.1)

$$\begin{aligned} & \mu \frac{L_c^2}{2} \left(a_1 |\operatorname{dev} \operatorname{sym}(R^T \operatorname{Curl} R)|^2 + a_2 |\operatorname{skew}(R^T \operatorname{Curl} R)|^2 + \frac{a_3}{3} \operatorname{tr}(R^T \operatorname{Curl} R)^2 \right) \\ & = \mu \frac{L_c^2}{2} \left(a_1 |\operatorname{dev} \operatorname{sym} \boldsymbol{\alpha}|^2 + a_2 |\operatorname{skew} \boldsymbol{\alpha}|^2 + \frac{a_3}{3} \operatorname{tr}(\boldsymbol{\alpha})^2 \right) = \mu \frac{L_c^2}{2} |\mathbb{P}_{a_1, a_2, a_3}(\boldsymbol{\alpha})|^2 =: W_{\text{disloc}}^{3D}(\boldsymbol{\alpha}), \quad (4.2) \end{aligned}$$

in terms of the so-called second order *wryness tensor* [60, 18] (second Cosserat deformation tensor [17])

$$\mathbf{\Gamma} := \left(\underbrace{\text{axl}(R^T \partial_x R)}_{\in \mathfrak{so}(3)} \mid \text{axl}(R^T \partial_y R) \mid \text{axl}(R^T \partial_z R) \right) = (\Gamma_1 \mid \Gamma_2 \mid \Gamma_3) \in \mathbb{R}^{3 \times 3}. \quad (4.3)$$

Since $R^T \partial_{x_i} R \in \mathfrak{so}(3)$, $i = 1, 2, 3$ is skew-symmetric, we have the following relation [25, 26, 61]

$$\mathbf{\Gamma} = -\boldsymbol{\alpha}^T + \frac{1}{2} \text{tr}(\boldsymbol{\alpha}) \mathbb{1}_3, \quad \boldsymbol{\alpha} = -\mathbf{\Gamma}^T + \text{tr}(\mathbf{\Gamma}) \mathbb{1}_3. \quad (4.4)$$

By using these formulas we note

$$\text{dev sym } \boldsymbol{\alpha} = -\text{dev sym } \mathbf{\Gamma}, \quad \text{skew } \boldsymbol{\alpha} = \text{skew } \mathbf{\Gamma}, \quad \text{tr}(\boldsymbol{\alpha}) = 2 \text{tr}(\mathbf{\Gamma}). \quad (4.5)$$

Now using (4.2), we obtain

$$\begin{aligned} W_{\text{disloc}}^{\text{3D}}(\boldsymbol{\alpha}) &= \mu \frac{L_c^2}{2} \left(a_1 |\text{dev sym } \boldsymbol{\alpha}|^2 + a_2 |\text{skew } \boldsymbol{\alpha}|^2 + \frac{a_3}{3} \text{tr}(\boldsymbol{\alpha})^2 \right) \\ &= \mu \frac{L_c^2}{2} \left(a_1 |\text{dev sym } \mathbf{\Gamma}|^2 + a_2 |\text{skew } \mathbf{\Gamma}|^2 + 4a_3 \text{tr}(\mathbf{\Gamma})^2 \right) \\ &= \mu \frac{L_c^2}{2} \left(\tilde{a}_1 |\text{dev sym } \mathbf{\Gamma}|^2 + \tilde{a}_2 |\text{skew } \mathbf{\Gamma}|^2 + \tilde{a}_3 \text{tr}(\mathbf{\Gamma})^2 \right) =: W_{\text{curv}}^{\text{3D}}(\mathbf{\Gamma}), \end{aligned} \quad (4.6)$$

where $\tilde{a}_1 = a_1, \tilde{a}_2 = a_2$ and $\tilde{a}_3 = 4a_3$. Altogether we get

$$\begin{aligned} W_{\text{disloc}}^{\text{3D}}(\boldsymbol{\alpha}) &= W_{\text{curv}}^{\text{3D}}(\mathbf{\Gamma}) = \mu \frac{L_c^2}{2} \left(\tilde{a}_1 |\text{dev sym } \mathbf{\Gamma}|^2 + \tilde{a}_2 |\text{skew } \mathbf{\Gamma}|^2 + \tilde{a}_3 \text{tr}(\mathbf{\Gamma})^2 \right), \quad \text{with } \tilde{a}_1, \tilde{a}_2, \tilde{a}_3 > 0, \\ &= \mu \frac{L_c^2}{2} \left(\tilde{b}_1 |\text{sym } \mathbf{\Gamma}| + \tilde{b}_2 |\text{skew } \mathbf{\Gamma}|^2 + \tilde{b}_3 \text{tr}(\mathbf{\Gamma})^2 \right), \end{aligned} \quad (4.7)$$

with $\tilde{a}_1 = \tilde{b}_1 > 0$, $\tilde{a}_2 = \tilde{b}_2 > 0$ and $\tilde{b}_3 = \frac{\tilde{a}_1}{3} + \tilde{a}_3 > 0$. Thus, the variational problem (3.1) can be equivalently expressed as

$$E^{\text{3D}}(\varphi, R) = \int_{\mathcal{U}_h} W_{\text{mp}}(R^T \text{D}\varphi) + W_{\text{curv}}^{\text{3D}}(\mathbf{\Gamma}) \, dx \rightarrow \min \quad \text{w.r.t } (\varphi, R). \quad (4.8)$$

Applying the nonlinear scaling [20], allows to rewrite the problem on a domain $\mathcal{U}_1 = \omega \times [-\frac{1}{2}, \frac{1}{2}]$ with unit thickness in terms of properly scaled variables $\varphi^{\natural}, R^{\natural}$ in (thickness) z -direction

$$E_h^{\text{3D}}(\varphi^{\natural}, R^{\natural}) = \int_{\mathcal{U}_1} W_{\text{mp}}(R^{\natural, T} \text{D}\varphi^{\natural}) + W_{\text{curv}}^{\text{3D}}(\mathbf{\Gamma}^{\natural}) \, dx. \quad (4.9)$$

The descaled Γ -limit of E_h^{3D} as $h \rightarrow 0$ is then given by [51]

$$E^{\text{2D}}(m, R) = \int_{\omega} h \left(W_{\text{mp}}^{\text{hom}}(R^T (\text{D}m \mid R_3)) + W_{\text{curv}}^{\text{hom}}(\widehat{\mathbf{\Gamma}}) \right) \, dx \rightarrow \min \quad \text{w.r.t } (m, R), \quad (4.10)$$

where $m: \omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$ describes the deformation of the midsurface, $R: \omega \subset \mathbb{R}^2 \rightarrow \text{SO}(3)$ and

$$\begin{aligned}
W_{\text{mp}}^{\text{hom}}(R^T(Dm|R_3)) &:= \inf_{d \in \mathbb{R}^3} W_{\text{mp}}(R^T(Dm|d)) \\
&= \mu |\text{sym}((R_1|R_2)^T Dm - \mathbb{1}_2)|^2 + \mu_c |\text{skew}((R_1|R_2)^T Dm - \mathbb{1}_2)|^2 \\
&\quad + \frac{2\mu\mu_c}{\mu + \mu_c} (\langle R_3, \partial_x m \rangle^2 + \langle R_3, \partial_y m \rangle^2) + \frac{\mu\lambda}{2\mu + \lambda} \text{tr}((R_1|R_2)^T Dm - \mathbb{1}_2)^2, \\
W_{\text{curv}}^{\text{hom}}(\widehat{\Gamma}) &:= \inf_{A \in \mathfrak{so}(3)} W_{\text{curv}}^{3\text{D}}(\text{axl}(R^T \partial_x R), \text{axl}(R^T \partial_y R), \text{axl}(A)) \\
&= \mu \frac{L_c^2}{2} \left(\widetilde{b}_1 \left| \text{sym} \begin{pmatrix} \widehat{\Gamma}_{11} & \widehat{\Gamma}_{12} \\ \widehat{\Gamma}_{21} & \widehat{\Gamma}_{22} \end{pmatrix} \right|^2 + \widetilde{b}_2 \left| \text{skew} \begin{pmatrix} \widehat{\Gamma}_{11} & \widehat{\Gamma}_{12} \\ \widehat{\Gamma}_{21} & \widehat{\Gamma}_{22} \end{pmatrix} \right|^2 \right. \\
&\quad \left. + \frac{\widetilde{b}_1 \widetilde{b}_3}{\widetilde{b}_1 + \widetilde{b}_3} \text{tr} \begin{pmatrix} \widehat{\Gamma}_{11} & \widehat{\Gamma}_{12} \\ \widehat{\Gamma}_{21} & \widehat{\Gamma}_{22} \end{pmatrix}^2 + 2 \frac{\widetilde{b}_1 \widetilde{b}_2}{\widetilde{b}_1 + \widetilde{b}_2} \left| \begin{pmatrix} \widehat{\Gamma}_{31} \\ \widehat{\Gamma}_{32} \end{pmatrix} \right|^2 \right), \tag{4.11}
\end{aligned}$$

where the matrix $\widehat{\Gamma} = (\text{axl}(R^T \partial_x R) | \text{axl}(R^T \partial_y R)) = \pi_{12}(\Gamma)$ is in the form (see [27])

$$\widehat{\Gamma} = (\text{axl}(R^T \partial_x R) | \text{axl}(R^T \partial_y R)) = \begin{pmatrix} \widehat{\Gamma}_{11} & \widehat{\Gamma}_{12} \\ \widehat{\Gamma}_{21} & \widehat{\Gamma}_{22} \\ \widehat{\Gamma}_{31} & \widehat{\Gamma}_{32} \end{pmatrix} \in \mathbb{R}^{3 \times 2}. \tag{4.12}$$

We set $\Gamma_{\square} = \begin{pmatrix} \widehat{\Gamma}_{11} & \widehat{\Gamma}_{12} \\ \widehat{\Gamma}_{21} & \widehat{\Gamma}_{22} \end{pmatrix}$ and $\Gamma_{\perp} = \begin{pmatrix} \widehat{\Gamma}_{31} \\ \widehat{\Gamma}_{32} \end{pmatrix}$. Thus we can write the Γ -limit minimization problem as³

$$\begin{aligned}
E_{\Gamma\text{-lim}}^{2\text{D}}(m, R) &= \int_{\omega} h \left\{ \mu |\text{sym}((R_1|R_2)^T Dm - \mathbb{1}_2)|^2 + \mu_c |\text{skew}((R_1|R_2)^T Dm - \mathbb{1}_2)|^2 \right. \\
&\quad + \frac{2\mu\mu_c}{\mu + \mu_c} (\langle R_3, \partial_x m \rangle^2 + \langle R_3, \partial_y m \rangle^2) + \frac{\mu\lambda}{2\mu + \lambda} \text{tr}((R_1|R_2)^T Dm - \mathbb{1}_2)^2 \\
&\quad \left. + \mu \frac{L_c^2}{2} \left(\widetilde{b}_1 |\text{sym} \widehat{\Gamma}_{\square}|^2 + \widetilde{b}_2 |\text{skew} \widehat{\Gamma}_{\square}|^2 + \frac{\widetilde{b}_1 \widetilde{b}_3}{\widetilde{b}_1 + \widetilde{b}_3} \text{tr}(\widehat{\Gamma}_{\square})^2 + 2 \frac{\widetilde{b}_1 \widetilde{b}_2}{\widetilde{b}_1 + \widetilde{b}_2} |\widehat{\Gamma}_{\perp}|^2 \right) \right\} dx. \tag{4.14}
\end{aligned}$$

If we assume that in the underlying Cosserat bulk curvature energy we have the uni-constant expression

$$\begin{aligned}
W_{\text{curv}}^{3\text{D}}(R^T DR) &= \frac{\mu L_c^2}{2} |R^T DR|^2 = \frac{\mu L_c^2}{2} |DR|^2 = \mu L_c^2 \left(|\text{axl}(R^T \partial_x R)|^2 + |\text{axl}(R^T \partial_y R)|^2 + |\text{axl}(R^T \partial_z R)|^2 \right) \\
&= \frac{\mu L_c^2}{2} \left(|(R^T \partial_x R)|^2 + |(R^T \partial_y R)|^2 + |(R^T \partial_z R)|^2 \right), \tag{4.15}
\end{aligned}$$

then the homogenized curvature energy is given by [10, 20]

$$\begin{aligned}
W_{\text{curv}}^{\text{hom}}(R^T DR) &= \inf_{A \in \mathfrak{so}(3)} W_{\text{curv}}^{3\text{D}}(\text{axl}(R^T \partial_x R), \text{axl}(R^T \partial_y R), \text{axl}(A)) \\
&= \mu \frac{L_c^2}{2} \left(|R^T \partial_x R|^2 + |R^T \partial_y R|^2 \right) = \mu \frac{L_c^2}{2} |R^T DR|^2 = \mu L_c^2 |\widehat{\Gamma}|_{\mathbb{R}^{3 \times 2}}^2 = \mu L_c^2 |\pi_{12}(\Gamma)|_{\mathbb{R}^{3 \times 2}}^2. \tag{4.16}
\end{aligned}$$

³Note the four fold appearance of the harmonic mean \mathcal{H} , i.e.

$$\frac{2\mu\mu_c}{\mu + \mu_c} = \mathcal{H}(\mu, \mu_c), \quad \frac{\mu\lambda}{2\mu + \lambda} = \frac{1}{2} \mathcal{H}\left(\mu, \frac{\lambda}{2}\right), \quad \frac{\widetilde{b}_1 \widetilde{b}_3}{\widetilde{b}_1 + \widetilde{b}_3} = \frac{1}{2} \mathcal{H}(\widetilde{b}_1, \widetilde{b}_3), \quad \frac{2\widetilde{b}_1 \widetilde{b}_2}{\widetilde{b}_1 + \widetilde{b}_2} = \mathcal{H}(\widetilde{b}_1, \widetilde{b}_2). \tag{4.13}$$

4.2 Alternative engineering ad-hoc dimensional descent

In [48] the three-dimensional Cosserat model has been reduced to a flat shell problem by proposing an engineering ansatz for the deformation φ and the microrotation R over the shell thickness. We let again $m: \omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$ denote the midsurface deformation, $\bar{U} := R^T(Dm|_{R_3})$ the non-symmetric membrane stretch tensor and $R: \mathcal{U}_h \subset \mathbb{R}^2 \rightarrow \text{SO}(3)$ denotes the microrotation tensor field with $R^T DR \cong (R^T \partial_x R, R^T \partial_y R)$. Since we are only interested in the membrane like response, we will neglect terms related to bending effects right away while keeping the curvature change⁴ scaling with h .

The dimensionally reduced energy reads then [48, (4.5)]

$$\begin{aligned}
E_{\text{eng}}^{2\text{D}} &= \int_{\omega} h \left\{ \mu |\text{sym}(\bar{U} - \mathbb{1}_3)|^2 + \mu_c |\text{skew}(\bar{U} - \mathbb{1}_3)|^2 + \frac{\mu\lambda}{2\mu + \lambda} \text{tr}(\bar{U} - \mathbb{1}_3)^2 + \mu \frac{L_c^2}{2} |R^T DR|^2 \right\} dx \\
&= \int_{\omega} h \left\{ \mu |\text{dev sym}(\bar{U} - \mathbb{1}_3)|^2 + \mu_c |\text{skew}(\bar{U} - \mathbb{1}_3)|^2 + \underbrace{\left(\frac{\mu\lambda}{2\mu + 3\lambda} + \frac{\mu}{3} \mu \right)}_{=: \kappa^{\text{hom}}} \text{tr}(\bar{U} - \mathbb{1}_3)^2 + \mu \frac{L_c^2}{2} |R^T DR|^2 \right\} dx \\
&= \int_{\omega} h \left\{ \mu |\text{dev sym}(R^T(Dm|_{R_3}) - \mathbb{1}_3)|^2 + \mu_c |\text{skew}(R^T(Dm|_{R_3}) - \mathbb{1}_3)|^2 \right. \\
&\quad \left. + \frac{\kappa^{\text{hom}}}{2} \text{tr}(R^T(Dm|_{R_3}) - \mathbb{1}_3)^2 + \mu \frac{L_c^2}{2} |DR|^2 \right\} dx \\
&= \int_{\omega} h \left\{ \underbrace{\mu |\text{sym}((R_1|R_2)^T Dm - \mathbb{1}_2)|^2}_{\text{shear-stretch energy}} + \underbrace{\mu_c |\text{skew}((R_1|R_2)^T Dm - \mathbb{1}_2)|^2}_{\text{drill-energy}} \right. \\
&\quad \left. + \frac{\mu + \mu_c}{2} \underbrace{(\langle R_3, \partial_x m \rangle^2 + \langle R_3, \partial_y m \rangle^2)}_{\text{transverse shear energy}} + \frac{\mu\lambda}{2\mu + \lambda} \underbrace{\text{tr}((R_1|R_2)^T Dm - \mathbb{1}_2)^2}_{\text{elogational stretch energy}} + \mu \underbrace{\frac{L_c^2}{2} |R^T DR|^2}_{\text{curvature energy}} \right\} dx.
\end{aligned} \tag{4.18}$$

Letting $\mu_c \rightarrow \infty$ in the reduced membrane model implies on the one hand that $R_3 = n_m$ is normal to the midsurface m and on the other hand $\text{skew}(R^T(Dm|_{n_m})) = 0$ implies $R = \text{polar}(Dm|_{n_m})$ (trièdre caché).

In contrast to the representation of the energy in (4.18) the rigorously derived Γ -limit membrane model [54] has the energy (see equation (4.14))

$$\begin{aligned}
E_{\Gamma\text{-lim}}^{2\text{D}}(m, R) &= \int_{\omega} h \left\{ \mu |\text{sym}((R_1|R_2)^T Dm - \mathbb{1}_2)|^2 + \mu_c |\text{skew}((R_1|R_2)^T Dm - \mathbb{1}_2)|^2 \right. \\
&\quad \left. + \frac{2\mu\mu_c}{\mu + \mu_c} (\langle R_3, \partial_x m \rangle^2 + \langle R_3, \partial_y m \rangle^2) + \frac{\mu\lambda}{2\mu + \lambda} \text{tr}((R_1|R_2)^T Dm - \mathbb{1}_2)^2 + W_{\text{curv}}^{\text{hom}}(\hat{\Gamma}) \right\} dx,
\end{aligned} \tag{4.19}$$

where $\hat{\Gamma} = (\text{axl}(R^T \partial_x R) | \text{axl}(R^T \partial_y R))$. Thus, the engineering formulation in (4.18) coincides with the membrane Γ -limit if and only if

$$\mathcal{A}(\mu, \mu_c) = \frac{\mu + \mu_c}{2} = \frac{2\mu\mu_c}{\mu + \mu_c} = \mathcal{H}(\mu, \mu_c) \iff \mu = \mu_c, \tag{4.20}$$

and

$$|R^T DR|^2 = W_{\text{curv}}^{\text{hom}}(\hat{\Gamma}) \iff \tilde{b}_1 = \tilde{b}_2 = 1, \quad \tilde{b}_3 = 0. \tag{4.21}$$

⁴The missing Cosserat bending terms scaling with h^3 are of the type [48, (4.5)]

$$\frac{h^3}{12} \left\{ \mu |\text{sym}(R^T(DR_3|_0))|^2 + \mu_c |\text{skew}(R^T(DR_3|_0))|^2 + \frac{\mu\lambda}{2\mu + \lambda} \text{tr}(\text{sym}(R^T(DR_3|_0)))^2 \right\}, \tag{4.17}$$

and the uni-constant case would appear for $\mu = \mu_c, \lambda = 0$.

In (4.18)₂, we are also led to define the appropriate modified bulk modulus κ^{hom} via⁵

$$\frac{\kappa^{\text{hom}}}{2} := \frac{\mu\lambda}{2\mu + \lambda} + \frac{\mu}{3} = \frac{2\mu}{3} \frac{2\lambda + \mu}{2\mu + \lambda} \quad (\text{effective two-dimensional bulk modulus}). \quad (4.22)$$

Since we will need $\kappa^{\text{hom}} > 0$ for our subsequent regularity analysis, (4.20)₂ implies $2\mu + \lambda > 0$ and $2\lambda + \mu > 0$. One can show that the latter implies for the engineering Poisson number $\nu := \frac{\lambda}{2(\mu + \lambda)}$ the bound $\nu > -\frac{1}{2}$ (instead of $\nu > -1$ for three-dimensional linear elasticity).⁶

5 The two-dimensional Euler-Lagrange equations

Henceforth, we skip all unnecessary material parameters in (4.18) in order to arrive at a compact representation. Again, we consider the midsurface deformation $m: \omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$ and the orthogonal microrotation tensor $R: \omega \subset \mathbb{R}^2 \rightarrow \text{SO}(3)$. We set $h = 1$ and assume the normalization $\mu \frac{L_c^2}{2} = 1$. Moreover, we set $\kappa = \frac{3\kappa^{\text{hom}}}{2}$. Thus, the corresponding energy function describing the two-dimensional membrane shell problem is

$$E(m, R) := \int_{\omega} \mu |\text{dev sym}(R^T(Dm|R_3) - \mathbb{1}_3)|^2 + \mu_c |\text{skew}(R^T(Dm|R_3) - \mathbb{1}_3)|^2 + \frac{\kappa}{3} \text{tr}(R^T(Dm|R_3) - \mathbb{1}_3)^2 + |DR|^2 \, dx. \quad (5.1)$$

We assume μ, μ_c, κ to be positive. Remember that we have defined a linear operator $\mathbb{P}: \mathbb{R}^{3 \times 3} \rightarrow \mathbb{R}^{3 \times 3}$ by

$$\mathbb{P}_{\mu, \mu_c, \kappa}(X) = \mathbb{P}(X) = \sqrt{\mu} \text{dev sym } X + \sqrt{\mu_c} \text{skew } X + \frac{\sqrt{\kappa}}{3} (\text{tr } X) \mathbb{1}_3. \quad (5.2)$$

Using the mutual orthogonality of $\text{dev sym } X$, $\text{skew } X$, and $(\text{tr } X) \mathbb{1}_3$, we can write down the functional in a simplified form, it reads

$$E(m, R) = \int_{\omega} |\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3)|^2 + |DR|^2 \, dx. \quad (5.3)$$

Now we are going to calculate the Euler-Lagrange equations for the dimensionally reduced problem based on E . The first variation of E in the direction of $(\vartheta, 0): \mathcal{U} \rightarrow \mathbb{R}^3 \times \mathbb{R}^{3 \times 3}$ is

$$\delta E(m, R; \vartheta, 0) = 2 \int_{\omega} \langle \mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3), \mathbb{P}(R^T(D\vartheta|0)) \rangle \, dx = 2 \int_{\omega} \langle \mathbb{P}(R^T(Dm|0) - (\mathbb{1}_2|0)), \mathbb{P}(R^T(D\vartheta|0)) \rangle \, dx, \quad (5.4)$$

and the first variation in the direction of $(0, Q): \mathcal{U} \rightarrow \mathbb{R}^3 \times \mathbb{R}^{3 \times 3}$ with $Q(x) \in T_R \text{SO}(3)$ for almost all $x \in \mathcal{U}$ is

$$\begin{aligned} \delta E(m, R; 0, Q) &= 2 \int_{\omega} [\langle \mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3), \mathbb{P}(Q^T(Dm|0)) \rangle + \langle DR, DQ \rangle] \, dx \\ &= 2 \int_{\omega} [\langle \mathbb{P}(R^T(Dm|0) - (\mathbb{1}_2|0)), \mathbb{P}(Q^T(Dm|0)) \rangle + \langle DR, DQ \rangle] \, dx. \end{aligned} \quad (5.5)$$

⁵In linear elasticity theory for the displacement $u: \mathcal{U} \subset \mathbb{R}^3 \rightarrow \mathbb{R}^3$, the common bulk modulus κ appears in the form $\mu |\text{dev sym } Du|^2 + \frac{\kappa}{2} \text{tr}(Du)^2$ and not as $\mu |\text{dev sym } Du|^2 + \frac{\kappa}{3} \text{tr}(Du)^2$, which would be more natural from the perspective of orthogonality of $\text{dev sym } Du$ and $\text{tr}(Du) \cdot \mathbb{1}_3$.

⁶ $2\lambda + \mu > 0$ and $\mu > 0$ implies $2\lambda + 2\mu = 2(\lambda + \mu) > 0$. Therefore, $\nu = \frac{\lambda}{2(\mu + \lambda)} > -\frac{1}{2} \Leftrightarrow \frac{\lambda}{\mu + \lambda} > -1 \Leftrightarrow \lambda > -(\mu + \lambda) \Leftrightarrow 2\lambda + \mu > 0$.

Using $\mathbb{P}^* = \mathbb{P}$ and $\mathbb{P}(X^T) = \mathbb{P}(X)^T$ such that $\mathbb{P}^*\mathbb{P} = \mathbb{P}^2$, and observing $\pi_{12}(v_1|v_2|v_3) = (v_1|v_2)$, we rewrite these as

$$\begin{aligned}\delta E(m, R; \vartheta, 0) &= 2 \int_{\omega} \langle R \mathbb{P}^2(R^T(Dm|0) - (\mathbb{1}_2|0)), (D\vartheta|0) \rangle_{\mathbb{R}^{3 \times 3}} dx \\ &= 2 \int_{\omega} \langle \pi_{12}(R \mathbb{P}^2(R^T(Dm|0) - (\mathbb{1}_2|0))), D\vartheta \rangle_{\mathbb{R}^{3 \times 2}} dx, \\ \delta E(m, R; 0, Q) &= 2 \int_{\omega} [\langle \mathbb{P}((Dm|0)^T R - (\mathbb{1}_2|0)), \mathbb{P}((Dm|0)^T Q) \rangle_{\mathbb{R}^{3 \times 3}} + \langle DR, DQ \rangle] dx \\ &= 2 \int_{\omega} [\langle (Dm|0) \mathbb{P}^2((Dm|0)^T R - (\mathbb{1}_2|0)), Q \rangle + \langle DR, DQ \rangle] dx.\end{aligned}\tag{5.6}$$

The pair of Euler-Lagrange equations then consists of

$$\text{Div}[\pi_{12}(2R \mathbb{P}^2(R^T(Dm|0) - (\mathbb{1}_2|0)))] = 0,\tag{5.7}$$

and

$$\Delta R - (Dm|0) \mathbb{P}^2((Dm|0)^T R - (\mathbb{1}_2|0)) \perp T_R \text{SO}(3).\tag{5.8}$$

Note that it is *not* true that $X^T \mathbb{P}^2(X) = X^T \mathbb{P}^* \mathbb{P} X$ is symmetric for all matrices X ; this is because \mathbb{P} is not a matrix. Therefore, $(Dm|0) \mathbb{P}^2((Dm|0)^T R - (\mathbb{1}_2|0))$ is not automatically orthogonal to $T_R \text{SO}(3)$. And this term, being formally only in L^1 due to Dm being in L^2 , makes the structure of the equation interesting, as explained in Section 1.1.

For readability, we introduce a product which shares aspects of scalar products and matrix multiplication. We define $\circ : \mathbb{R}^{3 \times 2} \times \mathbb{R}^{3 \times 2} \rightarrow \mathbb{R}^{3 \times 3}$ by

$$B \circ C := \frac{1}{2} B C^T = \frac{1}{2} (B|0) \begin{pmatrix} C^T \\ 0 \end{pmatrix}.\tag{5.9}$$

Defining

$$S(Dm, R) := \pi_{12}(2R \mathbb{P}^2(R^T(Dm|0) - (\mathbb{1}_2|0))),\tag{5.10}$$

we rewrite the second term of (5.8) as

$$\begin{aligned}(Dm|0) \mathbb{P}^2((Dm|0)^T R - (\mathbb{1}_2|0)) &= Dm \circ \pi_{12}(2\mathbb{P}^2(R^T(Dm|0) - (\mathbb{1}_2|0))) \\ &= Dm \circ R^T \pi_{12}(2R \mathbb{P}^2(R^T(Dm|0) - (\mathbb{1}_2|0))) \\ &= Dm \circ R^T S(Dm, R) = (Dm \circ S(Dm, R))R.\end{aligned}\tag{5.11}$$

Noting that the projection of any matrix $X \in \mathbb{R}^{3 \times 3}$ to $T_R \text{SO}(3)$ is $R \text{skew}(R^T X)$, we find that the projection of $(Dm \circ S(Dm, R))R$ is $R \text{skew}(R^T(Dm \circ S(Dm, R))R) = \text{skew}(Dm \circ S(Dm, R))R$. This means that the pair of Euler-Lagrange equations (5.7)–(5.8) can be rewritten as

$$\text{Div} S(Dm, R) = 0,\tag{5.12}$$

$$\Delta R - \text{skew}(Dm \circ S(Dm, R))R \perp T_R \text{SO}(3).\tag{5.13}$$

The latter is a relation rather than an equation, but we can rewrite it as an equation. In geometric analysis, this is usually done using the second fundamental form of $\text{SO}(3)$, but we present the calculation in a more elementary way. Our aim is to calculate the tangential part $(\Delta R)^\top$ of ΔR .

Differentiating $R R^T \equiv \mathbb{1}_3$ gives

$$0 = \partial_i(R R^T) = (\partial_i R) R^T + R \partial_i R^T = 2 \operatorname{sym}(R \partial_i R^T). \quad (5.14)$$

Differentiating $R^T R \equiv \mathbb{1}_3$ twice and summing over i , we find

$$\begin{aligned} 0 &= (\Delta R^T) R + R^T \Delta R + 2 \sum_i \partial_i R^T \partial_i R = (\Delta R)^T R + R^T \Delta R + 2 \sum_i \partial_i R^T \partial_i R \\ &= 2 \operatorname{sym}(R^T \Delta R) + 2 \sum_i \partial_i R^T \partial_i R, \end{aligned} \quad (5.15)$$

implying

$$\operatorname{sym}(R^T \Delta R) = - \sum_i \partial_i R^T \partial_i R. \quad (5.16)$$

For any fixed matrix $R \in \operatorname{SO}(3)$, we have $\mathbb{T}_R \operatorname{SO}(3) = R \mathfrak{so}(3)$, where $\mathfrak{so}(3)$ is the space of skew-symmetric matrices in $\mathbb{R}^{3 \times 3}$. The projections of any $X \in \mathbb{R}^{3 \times 3}$ to $\mathbb{T}_R \operatorname{SO}(3)$ or its orthogonal complement $[\mathbb{T}_R \operatorname{SO}(3)]^\perp$ therefore are

$$X^\top = R \operatorname{skew}(R^T X), \quad X^\perp = R \operatorname{sym}(R^T X). \quad (5.17)$$

Therefore, we can calculate the orthogonal component of ΔR as

$$(\Delta R)^\perp = R \operatorname{sym}(R^T \Delta R) = - \sum_i R \partial_i R^T \partial_i R = - \sum_i \operatorname{skew}(R \partial_i R^T) \partial_i R. \quad (5.18)$$

We have used (5.16) in the second “=”, and (5.14) in the third. We now abbreviate

$$\Omega_R := \begin{pmatrix} (\Omega_R)_1 \\ (\Omega_R)_2 \end{pmatrix} = - \begin{pmatrix} R \partial_x R^T \\ R \partial_y R^T \end{pmatrix} = - \begin{pmatrix} \operatorname{skew}(R \partial_x R^T) \\ \operatorname{skew}(R \partial_y R^T) \end{pmatrix}, \quad \Omega_R \cdot DR := \sum_{i=1}^2 (\Omega_R)_i \partial_i R \in \mathbb{R}^{3 \times 3}, \quad (5.19)$$

and hence have

$$(\Delta R)^\top = \Delta R - (\Delta R)^\perp = \Delta R - \Omega_R \cdot DR.$$

Combining with the result of (5.13), we have calculated the tangential part of the left-hand side of (5.8) as

$$\Delta R - \Omega_R \cdot DR - \operatorname{skew}(Dm \circ S(Dm, R)) R,$$

and thus have derived the Euler-Lagrange equations in their final form. We summarize

$$\begin{aligned} \operatorname{Div} S(Dm, R) &= 0, & (5.20) \\ \Delta R - \Omega_R \cdot DR - \operatorname{skew}(Dm \circ S(Dm, R)) R &= 0, & (5.21) \end{aligned}$$

where here

$$S(Dm, R) := \pi_{12}(2R \mathbb{P}^2(R^T(Dm|0) - (\mathbb{1}_2|0))), \quad \Omega_R := - \begin{pmatrix} \operatorname{skew}(R \partial_x R^T) \\ \operatorname{skew}(R \partial_y R^T) \end{pmatrix}.$$

Remark 5.1. In engineering language, (5.20) is the balance of forces, while 5.21 is the balance of angular momentum equation. The tensor

$$T(Dm, R) := 2 \mathbb{P}^2(R^T(Dm|0) - (\mathbb{1}_2|0)) \quad (5.22)$$

is the *non-symmetric Biot-type stress tensor* (symmetric if $\mu_c = 0$), while

$$S(Dm, R) = \pi_{12}(RT(Dm, R)) \quad (5.23)$$

is the *first Piola-Kirchhoff type force-stress tensor*. Note the analogy to the corresponding tensors in the 3D-Cosserat model presented in (A.7) and (A.8).

6 Regularity

The objective of this section is to prove our main theorem.

Theorem 6.1 (interior regularity). *Every weak solution $(m, R) \in W^{1,2}(\omega, \mathbb{R}^3 \times \text{SO}(3))$ of (5.20)-(5.21) is smooth on the interior of ω .*

Remark 6.2. Due to the results in [48, 52, 54], we know that energy minimizers to problem (5.1) exist and these are weak solutions $(m, R) \in W^{1,2}(\omega, \mathbb{R}^3 \times \text{SO}(3))$ of (5.20)-(5.21). Since the problem is highly nonlinear, uniqueness can neither be shown nor is it expected.

6.1 Hölder regularity

We observe that the last term in (5.21) is, up to “skew” and the harmless factor R , the product of a “gradient” Dm with a divergence-free quantity $S(Dm, R)$, with both factors in L^2 . As we know from [14], such a product is in the Hardy space \mathcal{H}^1 rather than just in L^1 , and we will use arguments from [64] that tell us how to handle the additional R factor. A standard source for the Hardy space \mathcal{H}^1 is in Chapter III of Stein’s book [70]. Note that [64](see also [24]) is about harmonic maps in ≥ 3 dimensions, and it is Rivière’s paper [63] about two-dimensional harmonic maps that is mostly the basis of what we are doing here. Schikorra [68] found some simplification to the arguments of [63] and [64], and the most accessible account of all these arguments to date is the textbook [28] which allows us to handle the Euler-Lagrange equation (5.21) quite flexibly. Note that our equation (5.21) is more general than the equations of the form $\Delta R - \Omega \cdot DR = 0$ studied in those papers and the book [28], since we have the extra term $-R \text{skew}(Dm \circ S(Dm, R))$ of order 0 in R . We are lucky that we have the additional structure coming from $S(Dm, R)$ being divergence-free, again implying that up to a bounded factor the extra term is in \mathcal{H}^1 . Without that additional information, we would not know how to incorporate that into the existing regularity theory.

It will be crucial to use Morrey norms, at least locally. We say that $u \in L^p(U)$ is in the Morrey space $M^{p,s}(U)$ if

$$[u]_{M^{p,s}(U)}^p := \sup \left\{ r^{-s} \int_{B_r(x_0) \cap U} |u|^p \, dx \mid x_0 \in U, r \in (0, 1) \right\} < \infty. \quad (6.1)$$

Having this, we define the Morrey norm by $\|u\|_{M^{p,s}(U)} := [u]_{M^{p,s}(U)} + \|u\|_{L^p(U)}$.

We need the following lemmas. The first one is a special case of Lemma A.1 in [68], in the spirit of similar estimates from [14]. This is where Hardy-BMO duality comes in as a hidden ingredient of our proof.

Lemma 6.3. *There is a constant C such that for all choices of $x_0 \in \mathbb{R}^2$, $r > 0$, and functions $a \in W^{1,2}(B_{2r}(x_0))$, $\Gamma \in L^2(B_r(x_0), (\mathbb{R}^2)^*)$, $b \in W_0^{1,2} \cap L^\infty(B_r(x_0))$ with $\text{Div } \Gamma = 0$ in the weak sense on $B_r(x_0)$, we have*

$$\left| \int_{B_r(x_0)} \langle Da, \Gamma \rangle b \, dx \right| \leq C \|\Gamma\|_{L^2(B_r(x_0))} \|Db\|_{L^2(B_r(x_0))} \|Da\|_{M^{3/2,1/2}(B_{2r}(x_0))}.$$

Another one, due to Rivière [63] and Schikorra [68], can be found as a special case of Theorem 10.57 in [28].

Lemma 6.4. *For every $\Omega \in L^2(B^2, (\mathbb{R}^2)^* \otimes \mathfrak{so}(3))$, there exists $G \in W^{1,2}(B^2, \text{SO}(3))$ such that*

$$\text{Div}(G^{-1}\Omega G - G^{-1}DG) = 0 \quad \text{in } B^2 \quad (6.2)$$

and ⁷

$$\|DG\|_{L^2(B^2)} + \|G^{-1}\Omega G - G^{-1}DG\|_{L^2(B^2)} \leq 3 \|\Omega\|_{L^2(B^2)}. \quad (6.3)$$

⁷ $(\mathbb{R}^2)^* \otimes \mathfrak{so}(3)$ is isomorphic to $\mathfrak{so}(3) \times \mathfrak{so}(3)$.

We also need a version of the Hodge decomposition theorem. This one is a special case of [36, Corollary 10.5.1], adapted from the differential forms version to 2-dimensional vector calculus as in [28, Corollary 10.70]

Lemma 6.5. *Let $p \in (1, \infty)$. On $B_r(x_0) \subset \mathbb{R}^2$, every 1-form $V \in L^p(B_r(x_0), (\mathbb{R}^2)^*)$ can be decomposed uniquely as*

$$V = D\alpha + D^\perp\beta + h,$$

where $\alpha \in W^{1,p}(B_r(x_0))$, $\beta \in W_0^{1,p}(B_r(x_0))$, and $h \in C^\infty(B_r(x_0), (\mathbb{R}^2)^*)$ is harmonic. Moreover, there is a constant C depending only on p , such that

$$\|\alpha\|_{W^{1,p}(B_r(x_0))} + \|\beta\|_{W_0^{1,p}(B_r(x_0))} + \|h\|_{L^p(B_r(x_0))} \leq C \|V\|_{L^p(B_r(x_0))}. \quad (6.4)$$

We now start our regularity proof. Our first step is local Hölder continuity.

Proposition 6.6. *Assume that $(m, R) \in W^{1,2}(\omega, \mathbb{R}^3 \times \text{SO}(3))$ is a weak solution of (5.20)–(5.21). Then there is $\beta > 0$ such that m and R are $C^{0,\beta}$ -Hölder continuous locally on ω .*

Proof. We write B_ρ for any ball $B_\rho(x_0) \subset \omega$. We assume r to be small enough such that $B_{2r}(x_0) \subset \omega$. We will collect more smallness conditions on r during the proof.

We choose G according to Lemma 6.4 and find, abbreviating $\Omega^G := G^{-1}\Omega_R G - G^{-1}DG$,

$$\begin{aligned} \text{Div}(G^{-1}DR) &= D(G^{-1}) \cdot DR + G^{-1}\Delta R \\ &= -G^{-1}(DG)G^{-1} \cdot DR + G^{-1}\Omega \cdot DR + G^{-1} \text{skew}(Dm \circ S(Dm, R))R \\ &= \Omega^G \cdot G^{-1}DR + G^{-1} \text{skew}(Dm \circ S(Dm, R))R. \end{aligned} \quad (6.5)$$

Now we Hodge-decompose $G^{-1}DR$ according to Lemma 6.5. We find $f \in W^{1,2}(B_r, \mathbb{R}^{3 \times 3})$, $g \in W_0^{1,2}(B_r, \mathbb{R}^{3 \times 3})$ with $dg = 0$, and a component-wise harmonic 1-form $h \in C^\infty(B_r, L(\mathbb{R}^2, \mathbb{R}^{3 \times 3}))$ such that

$$G^{-1}DR = Df + D^\perp g + h \quad (6.6)$$

almost everywhere in B_r . Using the well-known relations $\text{Div} D = \text{Div}^\perp D^\perp = \Delta$ and $\text{Div} D^\perp = \text{Div}^\perp D = 0$, we calculate

$$\Delta f = \text{Div} Df = \text{Div}(G^{-1}DR) = \Omega^G \cdot G^{-1}DR + G^{-1} \text{skew}(Dm \circ S(Dm, R))R, \quad (6.7)$$

and

$$\Delta g = \text{Div}^\perp D^\perp g = \text{Div}^\perp(G^{-1}DR) = D^\perp G^{-1} \cdot DR = \text{Div}((D^\perp G^{-1})(R - R_0)), \quad (6.8)$$

for any constant $R_0 \in \mathbb{R}^{3 \times 3}$ (not necessarily a rotation). Both terms on the right-hand side, multiplied with some $\varphi \in W_0^{1,3}(B_r, \mathbb{R}^{3 \times 3})$, can be estimated using Lemma 6.3. Choosing $a := R^{k\ell}$, $b := (G^{-1})^{jk} \varphi^{i\ell}$, $\Gamma^s := (\Omega^G)_{ij}^s$, we find

$$\int_{B_r} \langle \Omega^G \cdot G^{-1}DR, \varphi \rangle dx \leq C \|\Omega^G\|_{L^2(B_r)} (\|DG\|_{L^2(B_r)} \|\varphi\|_{L^\infty(B_r)} + \|D\varphi\|_{L^2(B_r)}) \|DR\|_{M^{3/2,1/2}(B_r)}, \quad (6.9)$$

and choosing $a := m^j$, $b := (G_{ij}^{-1}) R^{k\ell} \varphi^{i\ell}$, $\Gamma^s := S(Dm, R)_s^k$, we have

$$\begin{aligned} &\int_{B_r} \langle (G^{-1} \text{skew}(Dm \circ S(Dm, R))R), \varphi \rangle dx \\ &\leq C \|S(Dm, R)\|_{L^2(B_r)} \left(\|DG\|_{L^2(B_r)} \|\varphi\|_{L^\infty(B_r)} \|DR\|_{L^2(B_r)} \|\varphi\|_{L^\infty(B_r)} + \|D\varphi\|_{L^2(B_r)} \right) \|Dm\|_{M^{3/2,1/2}(B_r)}. \end{aligned} \quad (6.10)$$

We assume $\varepsilon \in (0, \varepsilon_0)$ with some $\varepsilon_0 > 0$ to be determined. Choosing $r > 0$ small enough, we may assume $\|\Omega_R\|_{L^2(B_r)} \leq \varepsilon$ and $\|Dm\|_{L^2(B_r)} \leq \varepsilon$.

We let $T := \{\varphi \in C_0^\infty(B_r, \mathbb{R}^{3 \times 3}) \mid \|\mathbf{D}\varphi\|_{L^3(B_r)} \leq 1\}$. Combining the duality of $L^{3/2}$ and L^3 and (6.7) with (6.9) and (6.10), we find, using $\varphi = 0$ on ∂B_r ,

$$\begin{aligned}
\|\mathbf{D}f\|_{L^{3/2}(B_r)} &\leq C \sup_{\varphi \in T} \int_{B_r} \langle \mathbf{D}f, \mathbf{D}\varphi \rangle \, dx \\
&= C \sup_{\varphi \in T} \int_{B_r} \langle \Omega^G \cdot G^{-1} \mathbf{D}R + G^{-1} \text{skew}(\mathbf{D}m \circ S(\mathbf{D}m, R))R, \varphi \rangle \, dx \\
&\leq C \sup_{\varphi \in T} \|\Omega^G\|_{L^2(B_r)} (\|\mathbf{D}G\|_{L^2} \|\varphi\|_{L^\infty(B_r)} + \|\mathbf{D}\varphi\|_{L^2(B_r)}) \|\mathbf{D}R\|_{M^{3/2,1/2}(B_{2r})} \\
&\quad + C \sup_{\varphi \in T} \|S(\mathbf{D}m, R)\|_{L^2(B_r)} \left(\|\mathbf{D}G\|_{L^2(B_r)} \|\varphi\|_{L^\infty(B_r)} + \|\mathbf{D}R\|_{L^2(B_r)} \|\varphi\|_{L^\infty(B_r)} \right. \\
&\quad \left. + \|\mathbf{D}\varphi\|_{L^2(B_r)} \right) \|\mathbf{D}m\|_{M^{3/2,1/2}(B_{2r})} \\
&\leq C \sup_{\varphi \in T} \|\Omega_R\|_{L^2(B_r)} (\|\Omega_R\|_{L^2(B_r)} \|\varphi\|_{L^\infty(B_r)} + \|\mathbf{D}\varphi\|_{L^2(B_r)}) \|\mathbf{D}R\|_{M^{3/2,1/2}(B_{2r})} \\
&\quad + C \sup_{\varphi \in T} (\|\mathbf{D}m\|_{L^2(B_r)} + r) \left(\|\Omega_R\|_{L^2(B_r)} \|\varphi\|_{L^\infty(B_r)} + \|\mathbf{D}R\|_{L^2(B_r)} \|\varphi\|_{L^\infty(B_r)} \right. \\
&\quad \left. + \|\mathbf{D}\varphi\|_{L^2(B_r)} \right) \|\mathbf{D}m\|_{M^{3/2,1/2}(B_{2r})} \\
&\leq C(\varepsilon + r) r^{1/3} (\|\mathbf{D}R\|_{M^{3/2,1/2}(B_{2r})} + \|\mathbf{D}m\|_{M^{3/2,1/2}(B_{2r})}).
\end{aligned} \tag{6.11}$$

Here, in the second “ \leq ”, we have used Lemma 6.3. And in the fourth “ \leq ”, we have used $\|\varphi\|_{L^\infty(B_r)} \leq Cr^{1/3} \|\mathbf{D}\varphi\|_{L^3(B_r)} \leq Cr^{1/3}$, $\|\mathbf{D}\varphi\|_{L^2(B_r)} \leq Cr^{1/3} \|\mathbf{D}\varphi\|_{L^3(B_r)} \leq Cr^{1/3}$, $\|\Omega_R\|_{L^2(B_r)} \leq \varepsilon$, and $\|\mathbf{D}m\|_{L^2(B_r)} \leq \varepsilon$.

Using (6.8), we can also estimate the $L^{3/2}$ -norm of $\mathbf{D}^\perp g$. We find

$$\begin{aligned}
\|\mathbf{D}^\perp g\|_{L^{3/2}(B_r)} &\leq C \sup_{\varphi \in T} \int_{B_r} \langle \mathbf{D}^\perp g, \mathbf{D}^\perp \varphi \rangle \, dx \\
&= C \sup_{\varphi \in T} \int_{B_r} \langle \Delta g, \varphi \rangle \, dx \\
&= C \sup_{\varphi \in T} \int_{B_r} \langle \text{Div}((\mathbf{D}^\perp G^{-1})(R - R_{B_r})), \varphi \rangle \, dx \\
&= C \sup_{\varphi \in T} \int_{B_r} \langle (\mathbf{D}^\perp G^{-1})(R - R_{B_r}), \mathbf{D}\varphi \rangle \, dx \\
&\leq C \sup_{\varphi \in T} \|\mathbf{D}\varphi\|_{L^3(B_r)} \|\mathbf{D}G\|_{L^2(B_r)} \|R - R_{B_r}\|_{L^6(B_r)} \\
&\leq C\varepsilon r^{1/3} \|\mathbf{D}R\|_{M^{3/2,1/2}(B_{2r})}.
\end{aligned} \tag{6.12}$$

This time, we have used $\|\mathbf{D}G\|_{L^2(B_r)} \leq 3 \|\Omega_R\|_{L^2(B_r)} \leq 3\varepsilon$, and the Sobolev embedding $W^{1,2/3} \hookrightarrow L^6$ for R .

For h , being harmonic, we have the standard estimate

$$\int_{B_\rho} |h|^{3/2} \, dx \leq C \left(\frac{\rho}{r}\right)^2 \int_{B_r} |h|^{3/2} \, dx, \tag{6.13}$$

for any $0 < \rho < r$. From (6.6), and then (6.11) and (6.12), we hence infer

$$\begin{aligned}
\|DR\|_{L^{3/2}(B_\rho)} &= \|G^{-1}DR\|_{L^{3/2}(B_\rho)} \\
&\leq \|h\|_{L^{3/2}(B_\rho)} + \|Df\|_{L^{3/2}(B_\rho)} + \|D^\perp g\|_{L^{3/2}(B_\rho)} \\
&\leq C\left(\frac{\rho}{r}\right)^{4/3} \|h\|_{L^{3/2}(B_r)} + \|Df\|_{L^{3/2}(B_\rho)} + \|D^\perp g\|_{L^{3/2}(B_\rho)} \\
&\leq C\left(\frac{\rho}{r}\right)^{4/3} \|DR\|_{L^{3/2}(B_r)} + C(\|Df\|_{L^{3/2}(B_r)} + \|D^\perp g\|_{L^{3/2}(B_r)}) \\
&\leq C\left(\frac{\rho}{r}\right)^{4/3} \|DR\|_{L^{3/2}(B_r)} + C(\varepsilon + r)r^{1/3}(\|DR\|_{M^{3/2,1/2}(B_{2r})} + \|Dm\|_{M^{3/2,1/2}(B_{2r})}).
\end{aligned} \tag{6.14}$$

Now we are going to derive a similar estimate for $\|Dm\|_{L^{3/2}}$. Hodge-decompose $S(Dm, R)$, i.e.

$$\pi_{12}(2R\mathbb{P}^2(R^T(Dm|0) - (\mathbb{1}_2|0))) = D^\perp\alpha + \chi, \tag{6.15}$$

with $\alpha \in W_0^{1,2}(B_r, \mathbb{R}^{3 \times 2})$, and $\chi \in W^{1,2}(B_r, L(\mathbb{R}^2, \mathbb{R}^{3 \times 2}))$ harmonic. This time, there is no term of the form $D\zeta$, since Div of the left-hand side is 0. This would imply that ζ is harmonic, and so would be $D\zeta$, which hence can be absorbed into χ . We have, abbreviating \mathbb{P}_R for the linear mapping $\xi \mapsto 2R\mathbb{P}^2(R^T(\xi))$,

$$\begin{aligned}
\Delta\alpha &= \text{Div}^\perp D^\perp\alpha \\
&= \text{Div}^\perp[\pi_{12}(2R\mathbb{P}^2(R^T(Dm|0) - (\mathbb{1}_2|0)))] \\
&= \text{Div}^\perp[\pi_{12}(\mathbb{P}_R(Dm|0) - 2R\mathbb{P}^2(\mathbb{1}_2|0))] \\
&= (D^\perp|0) \cdot [\mathbb{P}_R(Dm|0)] - \text{Div}^\perp[\pi_{12}(2R\mathbb{P}^2(\mathbb{1}_2|0))] \\
&= D^\perp\mathbb{P}_R \cdot Dm - \text{Div}^\perp[\pi_{12}(2R\mathbb{P}^2(\mathbb{1}_2|0))] \\
&= \text{Div}[(D^\perp\mathbb{P}_R)(m - m_{B_r})] - \text{Div}^\perp[\pi_{12}(2(R - R_{B_r})\mathbb{P}^2(\mathbb{1}_2|0))].
\end{aligned} \tag{6.16}$$

Using the same ideas as before, and defining $U := \{\psi \in C_0^\infty(B_r, \mathbb{R}^{3 \times 2}) \mid \|D^\perp\psi\|_{L^3(B_r)} \leq 1\}$, we estimate

$$\begin{aligned}
\|D^\perp\alpha\|_{L^{3/2}(B_r)} &\leq C \sup_{\psi \in U} \int_{B_r} \langle D^\perp\alpha, D^\perp\psi \rangle dx \\
&= C \sup_{\psi \in U} \int_{B_r} \left(\langle (D^\perp\mathbb{P}_R)(m - m_{B_r}), D\psi \rangle - \langle \pi_{12}(2(R - R_{B_r})\mathbb{P}^2(\mathbb{1}_2|0)), D^\perp\psi \rangle \right) dx \\
&\leq C \sup_{\psi \in U} (\|D\psi\|_{L^3(B_r)} \|D\mathbb{P}_R\|_{L^2(B_r)} \|m - m_{B_r}\|_{L^6(B_r)} + \|D\psi\|_{L^3(B_r)} \|R - R_{B_r}\|_{L^{3/2}(B_r)}) \\
&\leq C \sup_{\psi \in U} (\|D\psi\|_{L^3(B_r)} \|DR\|_{L^2(B_r)} \|Dm\|_{L^{3/2}(B_r)} + r \|D\psi\|_{L^3(B_r)} \|DR\|_{L^{3/2}(B_r)}) \\
&\leq C(\varepsilon + r)r^{1/3}(\|DR\|_{M^{3/2,1/2}(B_{2r})} + \|Dm\|_{M^{3/2,1/2}(B_{2r})}).
\end{aligned} \tag{6.17}$$

Proceeding exactly as above, we find

$$\begin{aligned}
\|Dm\|_{L^{3/2}(B_\rho)} &\leq C(\|\chi\|_{L^{3/2}(B_\rho)} + \|D^\perp\alpha\|_{L^{3/2}(B_\rho)} + \rho^{4/3}) \\
&\leq C\left(\frac{\rho}{r}\right)^{4/3} \|\chi\|_{L^{3/2}(B_r)} + C(\|D^\perp\alpha\|_{L^{3/2}(B_\rho)} + \rho^{4/3}) \\
&\leq C\left(\frac{\rho}{r}\right)^{4/3} \|Dm\|_{L^{3/2}(B_r)} + C(\|D^\perp\alpha\|_{L^{3/2}(B_r)} + \rho^{4/3}) \\
&\leq C\left(\frac{\rho}{r}\right)^{4/3} \|Dm\|_{L^{3/2}(B_r)} + C(\varepsilon + r)r^{1/3}(\|DR\|_{M^{3/2,1/2}(B_{2r})} + \|Dm\|_{M^{3/2,1/2}(B_{2r})}) + C\rho^{4/3}.
\end{aligned} \tag{6.18}$$

In order to do so, we have used

$$C^{-1}\|Dm\|_{L^{3/2}(B_s)} - Cs^{4/3} \leq \|\pi_{12}(2R\mathbb{P}^2(R^T(Dm|0) - (\mathbb{1}_2|0)))\|_{L^{3/2}(B_s)} \leq C(\|Dm\|_{L^{3/2}(B_s)} + s^{4/3}). \quad (6.19)$$

We divide (6.14) and (6.18) by $\rho^{1/3}$ and combine them into

$$\begin{aligned} & \rho^{-1/3}(\|DR\|_{L^{3/2}(B_\rho)} + \|Dm\|_{L^{3/2}(B_\rho)}) \\ & \leq C \frac{\rho}{r^{4/3}} (\|DR\|_{L^{3/2}(B_r)} + \|Dm\|_{L^{3/2}(B_r)}) \\ & \quad + C(\varepsilon + r) \left(\frac{r}{\rho}\right)^{1/3} (\|DR\|_{M^{3/2,1/2}(B_{2r})} + \|Dm\|_{M^{3/2,1/2}(B_{2r})}) + C\rho \\ & \leq C \left(\frac{\rho}{r} + (\varepsilon + r) \left(\frac{r}{\rho}\right)^{1/3}\right) (\|DR\|_{M^{3/2,1/2}(B_{2r})} + \|Dm\|_{M^{3/2,1/2}(B_{2r})}) + C\rho. \end{aligned} \quad (6.20)$$

We now assume $r \leq \varepsilon$, where $\varepsilon > 0$ is yet to be determined. For formal reasons, we also add ρ on both sides, which gives

$$\begin{aligned} & \rho^{-1/3}(\|DR\|_{L^{3/2}(B_\rho)} + \|Dm\|_{L^{3/2}(B_\rho)}) + \rho \\ & \leq C_0 \left(\frac{\rho}{r} + (\varepsilon + r) \left(\frac{r}{\rho}\right)^{1/3}\right) (\|DR\|_{M^{3/2,1/2}(B_{2r})} + \|Dm\|_{M^{3/2,1/2}(B_{2r})} + 2r) \end{aligned} \quad (6.21)$$

for some suitable constant C_0 . Now we fix $\rho := \frac{r}{12C_0}$ and $\varepsilon := (12C_0)^{-4/3}$, making $C_0(\frac{\rho}{r} + (\varepsilon + r)(\frac{r}{\rho})^{1/3}) = \frac{1}{6}$. Abbreviating $\theta := \frac{1}{12C_0}$, we thus have

$$(\theta r)^{-1/3}(\|DR\|_{L^{3/2}(B_{\theta r})} + \|Dm\|_{L^{3/2}(B_{\theta r})}) + \theta r \leq \frac{1}{6} (\|DR\|_{M^{3/2,1/2}(B_{2r})} + \|Dm\|_{M^{3/2,1/2}(B_{2r})} + 2r).$$

This holds for all $B_{\theta r}(x_0)$ and $B_{2r}(x_0) \subset \omega$ which share the same center x_0 . But clearly, we can replace $B_{2r}(x_0)$ with any ball $B_s(y_0) \supset B_{2r}(x_0)$ which is still in ω . All smallness assumptions made so far for $B_r(X_0)$ will now also be assumed for s , that is $s \leq \varepsilon$, $\|\Omega_R\|_{L^2(B_s(y_0))} \leq \varepsilon$, and $\|Dm\|_{L^2(B_s(y_0))} \leq \varepsilon$. We then have

$$(\theta r)^{-1/3}(\|DR\|_{L^{3/2}(B_{\theta r}(x_0))} + \|Dm\|_{L^{3/2}(B_{\theta r}(x_0))}) + \theta r \leq \frac{1}{6} (\|DR\|_{M^{3/2,1/2}(B_s(y_0))} + \|Dm\|_{M^{3/2,1/2}(B_s(y_0))} + s),$$

which is valid for all r, s, x_0, y_0 such that $B_{2r}(x_0) \subset B_s(y_0) \subset \omega$. Then the $B_{\theta r}(x_0)$ cover all of $B_{\theta s/2}(y_0)$. Hence, on the left-hand-side, we can take the infimum over all feasible r and x_0 , and find

$$\begin{aligned} & \|DR\|_{M^{3/2,1/2}(B_{\theta s/2}(y_0))} + \|Dm\|_{M^{3/2,1/2}(B_{\theta s/2}(y_0))} + \frac{\theta s}{2} \\ & \leq \frac{1}{2} (\|DR\|_{M^{3/2,1/2}(B_s(y_0))} + \|Dm\|_{M^{3/2,1/2}(B_s(y_0))} + s). \end{aligned} \quad (6.22)$$

We may replace s by $\frac{\theta}{2}s$ and iterate this, finding

$$\begin{aligned} & \|DR\|_{M^{3/2,1/2}(B_{(\theta/2)^k s}(y_0))} + \|Dm\|_{M^{3/2,1/2}(B_{(\theta/2)^k s}(y_0))} \\ & \leq 2^{-k} (\|DR\|_{M^{3/2,1/2}(B_s(y_0))} + \|Dm\|_{M^{3/2,1/2}(B_s(y_0))} + s) \end{aligned} \quad (6.23)$$

for all $k \in \mathbb{N}$. Now, for $r \approx (\theta/2)^k s$, we have $k \approx \frac{\log r/s}{\log(\theta/2)}$, and therefore $2^{-k} \approx (r/s)^{\frac{\log 2}{\log(\theta/2)}} =: (r/s)^\beta$. Hence we have proven that, for all $r \leq s$, the estimate

$$\|DR\|_{M^{3/2,1/2}(B_r(y_0))} + \|Dm\|_{M^{3/2,1/2}(B_r(y_0))} \leq Cr^\beta (\|DR\|_{M^{3/2,1/2}(B_s(y_0))} + \|Dm\|_{M^{3/2,1/2}(B_s(y_0))} + s)$$

holds. For $x_0 \in B_{s/2}(y_0)$ and $r \leq s/2$, we can apply the same with $B_s(y_0)$ replaced by $B_{2r}(y_0) \subset B_s(y_0)$, and hence find

$$\|DR\|_{M^{3/2,1/2}(B_r(x_0))} + \|Dm\|_{M^{3/2,1/2}(B_r(x_0))} \leq Cr^\beta (\|DR\|_{M^{3/2,1/2}(B_s(y_0))} + \|Dm\|_{M^{3/2,1/2}(B_s(y_0))} + s),$$

which implies

$$\|DR\|_{M^{3/2,1/2+3\beta/2}(B_{s/2}(y_0))} + \|Dm\|_{M^{3/2,1/2+3\beta/2}(B_{s/2}(y_0))} \leq C (\|DR\|_{M^{3/2,1/2}(B_s(y_0))} + \|Dm\|_{M^{3/2,1/2}(B_s(y_0))} + s).$$

This means

$$DR, Dm \in M_{\text{loc}}^{3/2,1/2+3\beta/2}(\omega). \quad (6.24)$$

We now use the following well-known fact, which can be found in [28, Theorem 5.7], for example.

Lemma 6.7 (Morrey' Dirichlet growth criterion). *Assume $U \subset \mathbb{R}^n$ to be open, $u \in W_{\text{loc}}^{1,p}(U)$, $Du \in M_{\text{loc}}^{p,n-p+\varepsilon}(U)$ for some $\varepsilon > 0$. Then $u \in C^{0,\varepsilon/p}$.*

With $p = \frac{3}{2}$, $n = 2$, the last estimate (6.24) and Lemma 6.7 imply $R, m \in C_{\text{loc}}^{0,\beta}(\omega)$, which is the Hölder regularity asserted in Proposition 6.6. \blacksquare

Remark 6.8. It is essential that we are working in the critical dimension $n = 2$ here, even though this may not be too obvious in the preceding proof which uses methods developed for supercritical dimensions. But the arithmetic of the exponents crucially uses $n = 2$. In particular, Lemma 6.3 for $n > 2$ is only available with exponents adding up to n instead of $(\frac{3}{2}, \frac{1}{2})$. But we would not succeed in finding similarly good estimates in the corresponding Morrey spaces.

6.2 Higher regularity

In this subsection, we are going to complete the proof of Theorem 6.1.

Proof. Remember we have the equations

$$\begin{aligned} \text{Div } S(Dm, R) &= 0, \\ \Delta R - \Omega_R \cdot DR - \text{skew}(Dm \circ S(Dm, R))R &= 0, \end{aligned} \quad (6.25)$$

where for $\xi \in \mathbb{R}^{3 \times 2}$ we have defined

$$S(\xi, R) = \pi_{12}(2R\mathbb{P}^2(R^T(\xi, 0) - (\mathbb{1}_2|0))) = \pi_{12}(2R\mathbb{P}^T\mathbb{P}(R^T(\xi|R_3) - \mathbb{1}_3)), \quad (6.26)$$

and

$$|\Omega_R| \leq C |DR|.$$

Abbreviating $L_R(\xi) := \pi_{12}(2R\mathbb{P}^2(R^T(\xi, 0)))$, we rewrite the first equation (6.25) as

$$\text{Div } L_R(Dm) = \text{Div}(\pi_{12}(2R\mathbb{P}^2(R^T(\mathbb{1}_2|0)))). \quad (6.27)$$

For every $R \in \text{SO}(3)$, $L_R : \mathbb{R}^{3 \times 2} \rightarrow \mathbb{R}^{3 \times 2}$ is a linear mapping satisfying the Legendre condition (uniform positivity) because of

$$\begin{aligned} \langle L_R(\xi), \xi \rangle &= \langle \pi_{12}(2R\mathbb{P}^2(R^T(\xi, 0))), \xi \rangle = \langle 2R\mathbb{P}^2(R^T(\xi, 0)), (\xi, 0) \rangle \\ &= \langle 2\mathbb{P}(R^T(\xi, 0)), \mathbb{P}(R^T(\xi, 0)) \rangle \geq 2\hat{\lambda}|R^T(\xi, 0)|^2 = 2\hat{\lambda}|\xi|^2, \end{aligned} \quad (6.28)$$

where here $\hat{\lambda} := \min\{\mu, \mu_c, \kappa\}$ is independent of R , hence we have a uniformly elliptic operator $m \mapsto \text{Div } L_R(Dm)$. For this operator, classical Schauder theory applies once it depends Hölder continuously on x through $R(x)$. And it does, because we already know $R \in C_{\text{loc}}^{0,\beta}$ for some $\beta > 0$.

We use the following version of Schauder theory. The proof is well known, a good reference is [28, Theorem 5.19] which reads as follows.

Lemma 6.9. *Let $u \in W_{\text{loc}}^{1,2}(\mathcal{U}, \mathbb{R}^m)$ be a solution to*

$$\text{Div}(A(x) \cdot Du) = -\text{Div } \mathcal{F},$$

with A satisfying the Legendre-Hadamard condition and having its components $A_{ij}^{\alpha\beta}$ in $C_{\text{loc}}^{0,\sigma}(\mathcal{U})$ for some $\sigma \in (0, 1)$. If $\mathcal{F}_i^\alpha \in C_{\text{loc}}^{0,\sigma}(\mathcal{U})$, then also Du is of class $C_{\text{loc}}^{0,\sigma}(\mathcal{U})$.

From what was proven in the last section, we know that both L_R and the right-hand side of (6.27) are in $C_{\text{loc}}^{0,\beta}$ locally, hence Lemma 6.9 implies that $m \in C_{\text{loc}}^{1,\beta}$ for some $\beta > 0$.

This simplifies the discussion of the regularity of R , because the Dm -terms in the equation for ΔR are now locally bounded. We can therefore rewrite it as

$$\Delta R + a(x, DR) = 0, \tag{6.29}$$

where the function a depends on R and Dm additionally, but those are locally bounded. The function satisfies

$$|a(x, DR)| \leq C(|DR|^2 + 1). \tag{6.30}$$

Since $DR \in L^2$, this means that ΔR is in L^1 , but L^1 is just not enough to perform regularity theory for R . However, the structure of the equation almost allows to apply the higher regularity theory for harmonic maps, where we could deal with $C|DR|^2$ instead of $C(|DR|^2 + 1)$ on the right-hand side. A simple formal trick will care for that condition. Let

$$u(x) = (u_0(x), u_1(x)) := (R(x), x_1) \tag{6.31}$$

with values in $\text{SO}(3) \times \mathbb{R}$. Then, letting $\tilde{a}(x, Du) := (a(x, Du_0), 0)$, we have

$$\Delta u + \tilde{a}(x, Du) = 0, \tag{6.32}$$

where here

$$|\tilde{a}(x, Du)| = |a(x, DR)| \leq C(|DR|^2 + 1) = C(|Du_0|^2 + |Du_1|^2) = C|Du|^2. \tag{6.33}$$

Now we can follow the regularity theory for harmonic maps for a while. Note that in [45], Lemma 3.7 and Proposition 3.2 assume u to be a harmonic map, but the proof uses only $|\Delta u| \leq C|Du|^2$ instead of the full harmonic map equation. We therefore can apply Lemmas 3.6 and 3.7, Proposition 3.2 in [45] to our u and find that $Du \in L_{\text{loc}}^\infty$. This means that the second term in (6.29) is in L_{loc}^p for all $p > 1$, and standard L^p -theory gives us $u \in W_{\text{loc}}^{2,p}$ for all $p > 1$. The Sobolev embedding $W^{1,p} \hookrightarrow C^{0,1-2/p}$ for $p > 2$ then gives us $Du \in C_{\text{loc}}^{0,\beta}$ with $\beta > 0$. Together with the result for m , we now have

$$(m, R) \in C_{\text{loc}}^{1,\beta} \quad \text{for some } \beta > 0.$$

Once we have this, we can iterate the Schauder estimates, i.e. differentiate the equations and apply Lemma 6.9 to partial derivatives of m, R instead of m and R alone. Thus we find that $(m, R) \in C_{\text{loc}}^{k,\beta}$ for our $\beta > 0$ and all $k \in \mathbb{N}$, which means we have proven that m and R are smooth on the interior of the domain. \blacksquare

6.3 Body forces

It is physically reasonable to consider the equations with an additional external body force term in the first equation of balance of forces,

$$\begin{aligned} \text{Div } S(Dm, R) &= f, \\ \Delta R - \Omega_R \cdot DR - \text{skew}(Dm \circ S(Dm, R))R &= 0 \end{aligned} \tag{6.34}$$

with $f \in W^{1,2}(\omega, \mathbb{R}^3)$. By integrating in one direction and setting

$$\mathcal{F}(x_1, x_2) := \left(\int_{(x_0)_1}^{x_1} f(t, x_2) dt \mid 0_{\mathbb{R}^3} \right) \in \mathbb{R}^{3 \times 2}, \quad (6.35)$$

we can always assume $f = \text{Div } \mathcal{F}$. Note that \mathcal{F} depends on the first component $(x_0)_1$ of the center of the ball $B_r(x_0)$ on which we are momentarily working. We have $\mathcal{F} \in W^{1,2}(\omega, \mathbb{R}^{3 \times 2})$, implying $\mathcal{F} \in L^p(\omega, \mathbb{R}^{3 \times 2})$ for all $p \in [1, \infty)$. Now we may rewrite the first equation as

$$\text{Div}(S(\text{D}m, R) - \mathcal{F}) = 0. \quad (6.36)$$

We will need to estimate $\text{D}\mathcal{F}$, which we calculate via $\partial_1 \mathcal{F} = (f, 0)$ and $\partial_2 \mathcal{F} = (\int_{(x_0)_1}^{x_1} \partial_2 f(t, x_2) dt, 0)$. The latter gives

$$\begin{aligned} \int_{B_r} |\partial_2 \mathcal{F}|^{3/2} dx &= \int_{B_r} \left| \int_{(x_0)_1}^{x_1} \partial_2 f(t, x_2) dt \right|^{3/2} dx \\ &\leq \int_{B_r} \left(\int_{(x_0)_1 - \sqrt{r^2 - x_2^2}}^{(x_0)_1 + \sqrt{r^2 - x_2^2}} |\partial_2 f(t, x_2)| dt \right)^{3/2} dx \\ &\leq Cr^{1/3} \int_{B_r} \int_{(x_0)_1 - \sqrt{r^2 - x_2^2}}^{(x_0)_1 + \sqrt{r^2 - x_2^2}} |\partial_2 f(t, x_2)|^{3/2} dt dx \\ &\leq Cr^{4/3} \int_{B_r} |\partial_2 f|^{3/2} dx. \end{aligned} \quad (6.37)$$

Since we can always assume $r \leq 1$, we have proven

$$\|\text{D}\mathcal{F}\|_{L^{3/2}} \leq C \|f\|_{W^{1,3/2}}. \quad (6.38)$$

The regularity theory for the more general equation including forces goes pretty much along the lines of the $\mathcal{F} = 0$ case presented in Section 6.1. We only indicate the necessary modifications. We rewrite (6.7) as

$$\Delta f = \Omega^G \cdot G^{-1} \text{D}R + G^{-1} \text{skew}(\text{D}m \circ (S(\text{D}m, R) - \mathcal{F}))R + G^{-1} \text{skew}(\text{D}m \circ \mathcal{F})R. \quad (6.39)$$

In (6.11), we replace $\|S(\text{D}m, R)\|_{L^2(B_r)}$ by $\|S(\text{D}m, R) - \mathcal{F}\|_{L^2(B_r)}$. Choosing the radius of B_r sufficiently small, we can also assume that $\|\mathcal{F}\|_{L^2(B_r)} \leq \varepsilon$, hence we can estimate $\|S(\text{D}m, R) - \mathcal{F}\|_{L^2(B_r)}$ by $C(\varepsilon + r)$ just as we did for $\|S(\text{D}m, R)\|_{L^2(B_r)}$ in (6.11). But we also have an additional term on the right-hand side of that estimate. Using the boundedness of G^{-1} and R , it is estimated as follows, assuming also $\|\mathcal{F}\|_{L^3(B_r)} \leq \varepsilon$. We have

$$\begin{aligned} \sup_{\varphi \in T} \int_{B_r} \langle G^{-1} \text{skew}(\text{D}m \circ \mathcal{F})R, \varphi \rangle dx &\leq C \sup_{\varphi \in T} r^{-1/3} \|\text{D}m\|_{L^{3/2}(B_r)} \|\mathcal{F}\|_{L^3(B_r)} r^{1/3} \|\varphi\|_{L^\infty(B_r)} \\ &\leq C \sup_{\varphi \in T} \|\text{D}m\|_{M^{3/2,1/2}} \|\mathcal{F}\|_{L^3(B_r)} r^{2/3} \|\text{D}\varphi\|_{L^3(B_r)} \\ &\leq C \varepsilon r^{1/3} \|\text{D}m\|_{M^{3/2,1/2}}, \end{aligned} \quad (6.40)$$

which can be absorbed in the right-hand side of (6.11). Hence the conclusion of (6.11) continues to hold also in the $\mathcal{F} \neq 0$ case.

The second modification we have to make is that we now Hodge-decompose $S(\text{D}m, R) - \mathcal{F}$, which means

$$\pi_{12}(2R\mathbb{P}^2(R^T(\text{D}m|0) - (\mathbb{1}_2|0))) - \mathcal{F} = \text{D}^\perp \alpha + \chi. \quad (6.41)$$

The additional term involving \mathcal{F} on the right-hand side of (6.16) can be written as $-\text{Div}^\perp(\mathcal{F} - \mathcal{F}_{B_r})$. In (6.17), $(\mathcal{F} - \mathcal{F}_{B_r})$ can be processed exactly like $(R - R_{B_r})$, resulting in an additional $C\tau \|\text{D}\psi\|_{L^3(B_r)} \|\text{D}\mathcal{F}\|_{L^{3/2}(B_r)}$,

which can be estimated using (6.38) and $\psi \in U$ as follows, making the additional smallness assumption $\|f\|_{W^{1,2}(B_r)} \leq \varepsilon$ for r ,

$$Cr \|D\psi\|_{L^3(B_r)} \|D\mathcal{F}\|_{L^{3/2}(B_r)} \leq Cr \|f\|_{W^{1,3/2}(B_r)} \leq Cr^{4/3} \|f\|_{W^{1,2}(B_r)} \leq \varepsilon r^{4/3}. \quad (6.42)$$

This additional term in (6.17) now contributes to the right-hand side of (6.18), but here enlarges only the $r^{4/3}$ and $\rho^{4/3}$ terms that are there, anyway. By the same argument, taking \mathcal{F} into account also contributes only to more of $s^{4/3}$ terms in

$$\begin{aligned} C^{-1} \|Dm\|_{L^{3/2}(B_s)} - Cs^{4/3} &\leq \|\pi_{12}(2R\mathbb{P}^2(R^T(Dm|0) - (\mathbb{1}_2|0))) - \mathcal{F}\|_{L^{3/2}(B_s)} \\ &\leq C(\|Dm\|_{L^{3/2}(B_s)} + s^{4/3}), \end{aligned} \quad (6.43)$$

which updates (6.19). Hence the contributions of the modified versions of both (6.17) and (6.19) do not change the conclusion of (6.18).

Now that we have adapted (6.11) and (6.18) to nonvanishing body forces, we can conclude Hölder continuity just as in the end of Section 6.1, under the weak assumption of f being in $W^{1,2}$. If we assume $f \in C^\infty$ instead, both f and \mathcal{F} are bounded, and the higher regularity proof from Section 6.2 goes through with hardly any modification. Note, for example, that (6.30) continues to hold.

6.4 Remarks on a special case

Our system simplifies considerably when $\mu = \mu_c = \kappa$, which makes \mathbb{P} the identity⁸. Even though this assumption is not too natural from the point of applications, we would like to comment briefly on that case.

The simplified variational functional reads now

$$\begin{aligned} E(m, R) &:= \int_{\omega} \mu |R^T(Dm|R_3) - \mathbb{1}_3|^2 + |DR|^2 \, dx = \int_{\omega} \mu |R^T(Dm|R_3) - R^T R|^2 + |DR|^2 \, dx \\ &= \int_{\omega} \mu |(Dm|0) - (R_1|R_2|0)|^2 + |DR|^2 \, dx, \end{aligned} \quad (6.44)$$

which has the Euler-Lagrange equations (c.f. (5.7))

$$\Delta m - \text{Div}(R_1|R_2|0) = 0, \quad (6.45)$$

and

$$\Delta R - \Omega_R \cdot DR + \mu R \text{skew}(R^T(Dm|0)) = \Delta R - \Omega_R \cdot DR + \mu \text{skew}((Dm|0)R^T) \cdot R = 0. \quad (6.46)$$

The point here is that the last term in the second equation now depends on Dm only linearly, making it an L^2 -term instead of L^1 (the L^1 -part is cancelled by the skew-operator). But harmonic map type equations with a right-hand side in L^2 have been studied by Moser in quite some generality, see the book [45] for an excellent exposition of the methods.

In particular, Moser has two theorems that help us here. Here, $N \subset \mathbb{R}^n$ is a compact manifold, $\mathcal{U} \subset \mathbb{R}^d$ a domain, and Π is the second fundamental form of the target manifold, which corresponds to our term quadratic in DR , i.e. $\sum_i \Pi(u)(\partial_i u, \partial_i u) = \Omega_u \cdot Du$ in our case.

Theorem 6.10. [45, Theorem 4.1] *Suppose $u \in W^{1,2}(\mathcal{U}, N)$ is a stationary solution of*

$$\Delta u - \sum_i \Pi(u)(\partial_i u, \partial_i u) = f,$$

in \mathcal{U} , for a function $f \in L^p(\mathcal{U}, \mathbb{R}^n)$, where $p > \frac{d}{2}$ and $p \geq 2$. Then there exists a relatively closed set $\Sigma \subset \mathcal{U}$ of vanishing $(d-2)$ -dimensional Hausdorff measure, such that $u \in C_{\text{loc}}^{0,\alpha}(\mathcal{U} \setminus \Sigma, N)$ for a number $\alpha > 0$ that depends only on m, N , and p .

⁸This case corresponds to $\mu = \lambda = \mu_c$ in the Cosserat bulk model and Poisson number $\nu = \frac{\lambda}{2(\mu+\lambda)} = \frac{1}{4}$ (nearly satisfied for magnesium).

Theorem 6.11. [45, Theorem 4.2] *Under the assumptions of the previous theorem, if $n \leq 4$ and $p = 2$, we also have $u \in W_{\text{loc}}^{2,2} \cap W_{\text{loc}}^{1,4}(\mathcal{U} \setminus \Sigma, N)$.*

While those theorems are highly nontrivial, it is standard to deduce regularity of the solutions to our model in the special case considered here.

Theorem 6.12 (interior regularity for $\mu = \mu_c = \kappa$). *Any solution $(m, R) \in W^{1,2}(\omega, \mathbb{R}^3 \times \text{SO}(3))$ of the simplified problem (6.45)-(6.46) is smooth on the interior of the domain ω .*

Proof. We first consider eq. (6.46). Since $-R \text{skew}(R^T(Dm|_0)) \in L^2$, we can apply Theorem 6.10 and Theorem 6.11 to find $R \in C_{\text{loc}}^{0,\alpha} \cap W_{\text{loc}}^{2,2}(\omega, \text{SO}(3))$. Note that $\Sigma = \emptyset$ here, since its 0-dimensional Hausdorff measure vanishes. Similarly, by L^2 -theory for (6.45), we have $m \in W_{\text{loc}}^{2,2}(\omega, \mathbb{R}^3)$. By the embedding $W^{2,2} \hookrightarrow W^{1,q}$ for all $q \in [2, \infty)$, we find that Δm and ΔR are in L_{loc}^q for every $q < \infty$, hence $(m, R) \in W_{\text{loc}}^{2,q}(\omega, \mathbb{R}^3 \times \text{SO}(3))$ for all $q < \infty$. This, in turn, embeds into $C_{\text{loc}}^{1,\alpha}$ for all $\alpha \in (0, 1)$, and hence the right-hand sides are Hölder continuous. From here, we can use Schauder estimates to show that (m, R) is C_{loc}^∞ on ω . ■

7 Conclusion and open problems

We have deduced interior Hölder regularity for a Dirichlet type geometrically nonlinear Cosserat flat membrane shell. The model is objective and isotropic but highly nonconvex. Therefore, our regularity result is astonishing and shows again the great versatility of the Cosserat approach compared to other more classical models. At present, we are limited to treating the uni-constant curvature case $|DR|^2$ since only then can sophisticated methods for harmonic functions with values in $\text{SO}(3)$ be employed. This calls for more effort of researchers to generalize the foregoing. Progress in this direction would also allow to consider the full Cosserat membrane-bending flat shell [48, 49, 50, 9]. Another case warrants further attention: taking the Cosserat couple modulus $\mu_c = 0$ in the model (in-plane drill allowed, but no energy connected to it) may still allow for regular minimizers. However, even the existence of minimizers remains unclear at present since it hinges on some sort of a priori regularity for the rotation field R (the non-quadratic curvature term $|DR|^{2+\varepsilon}$, $\varepsilon > 0$, together with zero Cosserat couple modulus $\mu_c = 0$ allows for minimizers [52, 47]). Finally, it is interesting to understand regularity properties of Cosserat shell models with curved initial geometry [25, 26, 27].

We expect some boundary regularity to hold, too. On the geometric analysis side, an adaptation of Rivière's boundary methods to problems with continuous Dirichlet boundary data has been performed in [46], which one could try to use. But with a view on applications, partially free boundary problems would probably be more interesting.

Acknowledgements: The authors are indebted to Oliver Sander and Lisa Julia Nebel (Chair of Numerical Mathematics, Technical University of Dresden) for preparing the calculations leading to Figures 4,5 and 6. The first author thanks Armin Schikorra (University of Pittsburgh) for an interesting discussion. The second author is grateful to Maryam Mohammadi Saem and Peter Lewintan (Faculty of Mathematics, University of Duisburg-Essen) for help in preparing the manuscript and the pictures.

Andreas Gastel acknowledges support in the framework of the DFG-Priority Programm 2256 "Variational Methods for Predicting Complex Phenomena in Engineering Structures and Materials" with the project title "Very singular solutions of a nonlinear Cosserat elasticity model for solids" with Project-no. 441380936 and Patrizio Neff acknowledges support in the framework of the DFG-Priority Programm 2256 "Variational Methods for Predicting Complex Phenomena in Engineering Structures and Materials" with the project title "A variational scale-dependent transition scheme: From Cauchy elasticity to the relaxed micromorphic continuum" with Project-no. 440935806 and also the DFG research grant Neff 902/8-1 Project-no. 415894848 with the title "Modelling and mathematical analysis of geometrically nonlinear Cosserat shells with higher order and residual effects".

References

- [1] F. Alouges and J.-M. Ghidaglia. “Minimizing Oseen-Frank energy for nematic liquid crystals: algorithms and numerical results”. *Annales de l’Institut Henri Poincaré Physique Théorique*. Vol. 66. 4. 1997, pp. 411–447.
- [2] H. Altenbach and V. Eremeyev. *Shell-like Structures: Non-Classical Theories and Applications*. Vol. 15. Advanced Structured Materials. Springer-Verlag, 2011.
- [3] J. Altenbach, H. Altenbach, and V. Eremeyev. “On generalized Cosserat-type theories of plates and shells: a short review and bibliography.” *Archive of Applied Mechanics* 80 (2010). Pp. 73–92.
- [4] P. Appell. “Traité de Mécanique Rationnelle”. *Gauthier-Villars, Paris* (1909).
- [5] J. M. Ball. “Mathematics and liquid crystals”. *Molecular Crystals and Liquid Crystals* 647.1 (2017). Pp. 1–27.
- [6] J. M. Ball and S. J. Bedford. “Discontinuous order parameters in liquid crystal theories”. *Molecular Crystals and Liquid Crystals* 612.1 (2015). Pp. 1–23.
- [7] Y. Başar. “A consistent theory of geometrically non-linear shells with an independent rotation vector”. *International Journal of Solids and Structures* 23.10 (1987). Pp. 1401–1415.
- [8] K. Bhattacharya and R. D. James. “A theory of thin films of martensitic materials with applications to microactuators”. *Journal of the Mechanics and Physics of Solids* 47.3 (1999). Pp. 531–576.
- [9] M. Birsan and P. Neff. “Existence theorems in the geometrically non-linear 6-parameter theory of elastic plates”. *Journal of Elasticity* 112.2 (2013). Pp. 185–198.
- [10] M. Birsan and P. Neff. “On the dislocation density tensor in the Cosserat theory of elastic shells”. *Advanced Methods of Continuum Mechanics for Materials and Structures, Advances Structures Materials, 60*, K. Naumenko and M. Aßmus (editors), Springer, 2016, pp. 391–413.
- [11] A. Braides. *Gamma-Convergence for Beginners*. Vol. 22. Clarendon Press, 2002.
- [12] A. Braides. “A Handbook of Γ -convergence”. *Handbook of Differential Equations: Stationary Partial Differential Equations*. Vol. 3. Elsevier, 2006, pp. 101–213.
- [13] G. Capriz. *Continua with Microstructure*. Springer, 1989.
- [14] R. R. Coifman. “Compensated compactness and Hardy spaces”. *Journal de Mathématiques Pures et Appliquées* 72.9 (1993). Pp. 247–286.
- [15] E. Cosserat and F. Cosserat. *Note sur la théorie de l’action euclidienne*. Appendix in [4], pp. 557–629.
- [16] E. Cosserat and F. Cosserat. “Sur la théorie des corps minces”. *Comptes Rendus* 146 (1908). Pp. 169–172.
- [17] E. Cosserat and F. Cosserat. *Théorie des corps déformables*. A. Hermann et fils, http://www.uni-due.de/~hm0014/Cosserat_files/Cosserat09_eng.pdf, English translation by D. Delphenich 2007, <https://www.nature.com/articles/081067a0.pdf>, 1909.
- [18] V. A. Eremeyev and W. Pietraszkiewicz. “The nonlinear theory of elastic shells with phase transitions”. *Journal of Elasticity* 74.1 (2004). Pp. 67–86.
- [19] M. Focardi, P. M. Mariano, and E. Spadaro. “Multi-value microstructural descriptors for complex materials: analysis of ground states”. *Archive for Rational Mechanics and Analysis* 217.3 (2015). Pp. 899–933.
- [20] I. Fonseca and G. Francfort. “On the inadequacy of the scaling of linear elasticity for 3D–2D asymptotics in a nonlinear setting”. *Journal de Mathématiques Pures et Appliquées* 80.5 (2001). Pp. 547–562.
- [21] G. Friesecke, R. James, and S. Müller. “A theorem on geometric rigidity and the derivation of nonlinear plate theory from three-dimensional elasticity”. *Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences* 55.11 (2002). Pp. 1461–1506.
- [22] G. Friesecke, R. D. James, M. G. Mora, and S. Müller. “Derivation of nonlinear bending theory for shells from three-dimensional nonlinear elasticity by Gamma-convergence”. *Comptes Rendus Mathématique* 336.8 (2003). Pp. 697–702.
- [23] G. Friesecke, R. D. James, and S. Müller. “A hierarchy of plate models derived from nonlinear elasticity by Γ -convergence”. *Archive for Rational Mechanics and Analysis* 180.2 (2006). Pp. 183–236.
- [24] A. Gastel. “Regularity issues for Cosserat continua and p-harmonic maps”. *SIAM Journal on Mathematical Analysis* 51.6 (2019). Pp. 4287–4310.
- [25] I. Ghiba, M. Birsan, P. Lewintan, and P. Neff. “The isotropic Cosserat shell model including terms up to $O(h^5)$. Part I: Derivation in matrix notation”. *Journal of Elasticity* 142 (2020). Pp. 201–262.
- [26] I. Ghiba, M. Birsan, P. Lewintan, and P. Neff. “The isotropic elastic Cosserat shell model including terms up to order $O(h^5)$ in the shell thickness. Part II: Existence of minimizers”. *Journal of Elasticity* 142 (2020). 263–290.
- [27] I. Ghiba, M. Mohammadi Saem, and P. Neff. “On the choice of third order curvature tensors in the geometrically nonlinear Cosserat micropolar model and applications”. *in preparation* ().

- [28] M. Giaquinta and L. Martinazzi. *An Introduction to the Regularity Theory for Elliptic Systems, Harmonic Maps and Minimal Graphs*. Springer -Verlag Business Media, 2013.
- [29] A. E. Green, P. M. Naghdi, and W. Wainwright. “A general theory of a Cosserat surface”. *Archive for Rational Mechanics and Analysis* 20.4 (1965). Pp. 287–308.
- [30] M. Grüter. “Regularity of weak H -surfaces”. *Journal für die reine und angewandte Mathematik* 329 (1981). Pp. 1–15.
- [31] R. Hardt, D. Kinderlehrer, and F.-H. Lin. “Existence and partial regularity of static liquid crystal configurations”. *Communications in Mathematical Physics* 105.4 (1986). Pp. 547–570.
- [32] F. Hélein. “Régularité des application faiblement harmoniques entre une surface et une sphère”. *Comptes Rendus de l’Académie des Sciences Paris. Série I. Mathématique* 311 (1990). Pp. 519–524.
- [33] F. Hélein. “Régularité des application faiblement harmoniques entre une surface et une variété riemannienne”. *Comptes Rendus de l’Académie des Sciences Paris. Série I. Mathématique* 312 (1991). Pp. 591–596.
- [34] F. Hélein. *Harmonic Maps, Conservation Laws and Moving Frames*. Cambridge Tracts in Mathematics. Cambridge University Press, 2002.
- [35] A. Ibrahimbegović. “Stress resultant geometrically nonlinear shell theory with drilling rotations—Part I. A consistent formulation”. *Computer Methods in Applied Mechanics and Engineering* 118.3-4 (1994). Pp. 265–284.
- [36] T. Iwaniec and G. Martin. *Geometric Function Theory and Non-Linear Analysis*. Oxford Mathematics. The Clarendon Press, Oxford University Press, New York, 2001.
- [37] G. Kikis and S. Klinkel. “Two-field formulations for isogeometric Reissner–Mindlin plates and shells with global and local condensation”. *Computational Mechanics* 69.1 (2022). Pp. 1–21.
- [38] J. Lankeit, P. Neff, and F. Osterbrink. “Integrability conditions between the first and second Cosserat deformation tensor in geometrically nonlinear micropolar models and existence of minimizers”. *Zeitschrift für angewandte Mathematik und Physik* 68.1 (2017). Pp. 1–19.
- [39] H. Le Dret and A. Raoult. “The nonlinear membrane model as variational limit of nonlinear three-dimensional elasticity”. *Journal de Mathématiques Pures et Appliquées* 74.6 (1995). Pp. 549–578.
- [40] H. Le Dret and A. Raoult. “The membrane shell model in nonlinear elasticity: a variational asymptotic derivation”. *Journal of Nonlinear Science* 6.1 (1996). Pp. 59–84.
- [41] M. Lewicka, M. G. Mora, and M. R. Pakzad. “Shell theories arising as low energy Γ -limit of 3d nonlinear elasticity”. *Annali della Scuola Normale Superiore di Pisa - Classe di Scienze* Ser. 5, 9.2 (2010). Pp. 253–295.
- [42] Y. Li and C. Wang. “Regularity of weak solution of variational problems modeling the Cosserat micropolar elasticity”. *International Mathematics Research Notices* 2022.6 (2022). Pp. 4620–4658.
- [43] P. M. Mariano and G. Modica. “Ground states in complex bodies”. *ESAIM: Control, Optimisation and Calculus of Variations* 15.2 (2009). Pp. 377–402.
- [44] C. B. Morrey Jr. “The problem of Plateau on a Riemannian manifold”. *Annals of Mathematics* 49 (1948). Pp. 807–951.
- [45] R. Moser. *Partial Regularity for Harmonic Maps and Related Problems*. World Scientific, 2005.
- [46] F. Müller and A. Schikorra. “Boundary regularity via Uhlenbeck–Rivière decomposition”. *Analysis (Munich)* 29.2 (2009). Pp. 199–220.
- [47] P. Neff. “On Korn’s first inequality with non-constant coefficients”. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics* 132.1 (2002). Pp. 221–243.
- [48] P. Neff. “A geometrically exact Cosserat shell-model including size effects, avoiding degeneracy in the thin shell limit. Part I: Formal dimensional reduction for elastic plates and existence of minimizers for positive Cosserat couple modulus”. *Continuum Mechanics and Thermodynamics* 16.6 (2004). Pp. 577–628.
- [49] P. Neff. “Existence of minimizers for a geometrically exact Cosserat solid”. *Proceedings in Applied Mathematics and Mechanics*. Vol. 4. 1. Wiley Online Library. 2004, pp. 548–549.
- [50] P. Neff. “Geometrically Exact Cosserat Theory for Bulk Behaviour and Thin Structures: Modelling and Mathematical Analysis”. *Habilitation Thesis, TU-Darmstadt* (2004).
- [51] P. Neff. “The Γ -limit of a finite-strain Cosserat model for asymptotically thin domains and a consequence for the Cosserat couple modulus μ_c ”. *PAMM: Proceedings in Applied Mathematics and Mechanics*. Vol. 5. 1. Wiley Online Library. 2005, pp. 629–630.
- [52] P. Neff. “A geometrically exact planar Cosserat shell-model with microstructure: Existence of minimizers for zero Cosserat couple modulus”. *Mathematical Models and Methods in Applied Sciences* 17.03 (2007). Pp. 363–392.
- [53] P. Neff, M. Birsan, and F. Osterbrink. “Existence theorem for geometrically nonlinear Cosserat micropolar model under uniform convexity requirements”. *Journal of Elasticity* 121.1 (2015). Pp. 119–141.
- [54] P. Neff and K. Chelminski. “A geometrically exact Cosserat shell-model for defective elastic crystals. Justification via Γ -convergence”. *Interfaces and Free Boundaries* 9.4 (2007). Pp. 455–492.

- [55] P. Neff, A. Fischle, and L. Borisov. “Explicit global minimization of the symmetrized Euclidean distance by a characterization of real matrices with symmetric square”. *SIAM Journal on Applied Algebra and Geometry* 3.1 (2019). Pp. 31–43.
- [56] P. Neff, A. Fischle, and I. Münch. “Symmetric Cauchy stresses do not imply symmetric Biot strains in weak formulations of isotropic hyperelasticity with rotational degrees of freedom”. *Acta Mechanica* 197.1 (2008). Pp. 19–30.
- [57] P. Neff, I. Ghiba, O. Sander, and L. J. Nebel. “The classical geometrically nonlinear, physically linear Reissner–Mindlin and Kirchhoff–Love membrane-bending model is ill-posed”. *in preparation* ().
- [58] P. Neff, K.-I. Hong, and J. Jeong. “The Reissner–Mindlin plate is the Γ -limit of Cosserat elasticity”. *Mathematical Models and Methods in Applied Sciences* 20.09 (2010). Pp. 1553–1590.
- [59] P. Neff, J. Lankeit, and A. Madeo. “On Grioli’s minimum property and its relation to Cauchy’s polar decomposition”. *International Journal of Engineering Science* 80 (2014). Pp. 209–217.
- [60] P. Neff and I. Münch. “Curl bounds Grad on $SO(3)$ ”. *ESAIM: Control, Optimisation and Calculus of Variations* 14.1 (2008). Pp. 148–159.
- [61] J. F. Nye. “Some geometrical relations in dislocated crystals”. *Acta Metallurgica* 1.2 (1953). Pp. 153–162.
- [62] W. Pietraszkiewicz and V. Konopińska. “Drilling couples and refined constitutive equations in the resultant geometrically non-linear theory of elastic shells”. *International Journal of Solids and Structures* 51.11-12 (2014). Pp. 2133–2143.
- [63] T. Riviere. “Conservation laws for conformally invariant variational problems”. *Inventiones Mathematicae* 168.1 (2007). Pp. 1–22.
- [64] T. Riviere and M. Struwe. “Partial regularity for harmonic maps and related problems”. *Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences* 61.4 (2008). Pp. 451–463.
- [65] O. Sander. “Geodesic finite elements of higher order”. *IMA Journal of Numerical Analysis* 36.1 (2016). Pp. 238–266.
- [66] O. Sander. *DUNE—The Distributed and Unified Numerics Environment*. Vol. 140. Springer Nature, 2020.
- [67] O. Sander, P. Neff, and M. Birsan. “Numerical treatment of a geometrically nonlinear planar Cosserat shell model”. *Computational Mechanics* 57.5 (2016). Pp. 817–841.
- [68] A. Schikorra. “A remark on gauge transformations and the moving frame method”. *Annales de l’Institut Henri Poincaré Analyse Non Linéaire*. Vol. 27. 2. 2010, pp. 503–515.
- [69] R. Schoen. “Analytic aspects of the harmonic map problem”. *Seminar on Nonlinear Partial Differential Equations*. Ed. by S. Chern. Vol. 2. MSRI Publications. Springer-Verlag, New York, 1984.
- [70] E. M. Stein and T. S. Murphy. *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*. Vol. 3. Princeton University Press, 1993.
- [71] E. G. Virga. *Variational Theories for Liquid Crystals*. Chapman and Hall/CRC, 2018.
- [72] K. Wisniewski. “A shell theory with independent rotations for relaxed Biot stress and right stretch strain”. *Computational Mechanics* 21.2 (1998). Pp. 101–122.
- [73] K. Wisniewski and E. Turska. “Kinematics of finite rotation shells with in-plane twist parameter”. *Computer Methods in Applied Mechanics and Engineering* 190.8-10 (2000). Pp. 1117–1135.
- [74] Y. Zhang, C. Sansour, and C. Bingham. “Single-walled carbon nanotube modelling based on Cosserat surface theory”. *Recent Advances in Electrical and Computer Engineering* (2013). Pp. 32–37.

A Appendix

A.1 Three-dimensional Euler-Lagrange equations in dislocation tensor format

Here, for the convenience of the reader we derive the three-dimensional Euler-Lagrange equations based on the curvature expressed in the dislocation tensor $\boldsymbol{\alpha} = R^T \text{Curl} R$. We can write the bulk elastic energy as

$$E^{3D}(\varphi, R) = \int_{\mathcal{U}} W_{\text{mp}}(\bar{U}) + W_{\text{disloc}}(\boldsymbol{\alpha}) \, dx, \quad \bar{U} = R^T D\varphi, \quad \boldsymbol{\alpha} = R^T \text{Curl} R. \quad (\text{A.1})$$

Taking variations of (A.1) w.r.t. the deformation φ leads to

$$\begin{aligned} \delta E^{3D}(\varphi, R) \cdot \delta\varphi &= \int_{\mathcal{U}} \langle DW_{\text{mp}}(\bar{U}), R^T D\delta\varphi \rangle_{\mathbb{R}^{3 \times 3}} \, dx = 0, \quad \forall \delta\varphi \in C_0^\infty(\mathcal{U}, \mathbb{R}^3) \\ &\Leftrightarrow \int_{\mathcal{U}} \langle R DW_{\text{mp}}(\bar{U}), D\delta\varphi \rangle_{\mathbb{R}^{3 \times 3}} \, dx = \int_{\mathcal{U}} \langle \text{Div}[R \cdot DW_{\text{mp}}(\bar{U})], \delta\varphi \rangle_{\mathbb{R}^3} \, dx = 0. \end{aligned} \quad (\text{A.2})$$

Taking variation w.r.t. $R \in \text{SO}(3)$ results in (abbreviate $F := D\varphi$)

$$\begin{aligned} \delta E^{3\text{D}}(\varphi, R) \cdot \delta R &= \int_{\mathcal{U}} \langle DW_{\text{mp}}(\bar{U}), \delta R^T F \rangle + \langle DW_{\text{disloc}}(\boldsymbol{\alpha}), \delta R^T \text{Curl} R + R^T \text{Curl} \delta R \rangle \, dx \\ &= \int_{\mathcal{U}} \langle DW_{\text{mp}}(\bar{U}), \delta R^T R \cdot R^T F \rangle + \langle DW_{\text{disloc}}(\boldsymbol{\alpha}), \delta R^T R \cdot R^T \text{Curl} R + R^T \text{Curl} \delta R \rangle \, dx \\ &= \int_{\mathcal{U}} \langle DW_{\text{mp}}(\bar{U}) \cdot \bar{U}^T, \delta R^T R \rangle + \langle DW_{\text{disloc}}(\boldsymbol{\alpha}), \delta R^T R \cdot \boldsymbol{\alpha} + R^T \text{Curl} \delta R \rangle \, dx = 0. \end{aligned} \quad (\text{A.3})$$

Since $R^T R = \mathbb{1}_3$, it follows that $\delta R^T R + R^T \delta R = 0$ and $\delta R^T R = A \in \mathfrak{so}(3)$ is arbitrary. Therefore, (A.3) can be written as

$$0 = \int_{\mathcal{U}} \langle DW_{\text{mp}}(\bar{U}) \cdot \bar{U}^T, A \rangle + \langle DW_{\text{disloc}}(\boldsymbol{\alpha}) \cdot \boldsymbol{\alpha}^T, A \rangle + \langle DW_{\text{disloc}}(\boldsymbol{\alpha}), R^T \text{Curl}(R A^T) \rangle \, dx \quad (\text{A.4})$$

for all $A \in C_0^\infty(\mathcal{U}, \mathfrak{so}(3))$. Using that Curl is a self-adjoint operator, this is equal to

$$\begin{aligned} 0 &= \int_{\mathcal{U}} \langle DW_{\text{mp}}(\bar{U}) \cdot \bar{U}^T + DW_{\text{disloc}}(\boldsymbol{\alpha}) \boldsymbol{\alpha}^T, A \rangle + \langle \text{Curl}(R DW_{\text{disloc}}(\boldsymbol{\alpha})), R A^T \rangle \, dx \\ &= \int_{\mathcal{U}} \langle DW_{\text{mp}}(\bar{U}) \cdot \bar{U}^T + DW_{\text{disloc}}(\boldsymbol{\alpha}) \boldsymbol{\alpha}^T - R^T \text{Curl}(R DW_{\text{disloc}}(\boldsymbol{\alpha})), A \rangle \, dx \quad \forall A \in C_0^\infty(\mathcal{U}, \mathfrak{so}(3)). \end{aligned} \quad (\text{A.5})$$

Thus, the strong form of the Euler-Lagrange equations reads

$$\begin{aligned} \text{Div}[R DW_{\text{mp}}(\bar{U})] &= 0, & \text{"balance of forces"}, \\ \text{skew}[R^T \text{Curl}(R DW_{\text{disloc}}(\boldsymbol{\alpha}))] &= \text{skew}(DW_{\text{mp}}(\bar{U}) \cdot \bar{U}^T + DW_{\text{disloc}}(\boldsymbol{\alpha}) \cdot \boldsymbol{\alpha}^T), \\ & \text{"balance of angular momentum"}. \end{aligned} \quad (\text{A.6})$$

Defining the first Piola-Kirchhoff stress tensor

$$S_1(D\varphi, R) := D_F[W_{\text{mp}}(\bar{U})] = R DW_{\text{mp}}(\bar{U}) = R \cdot T_{\text{Biot}}(\bar{U}), \quad (\text{A.7})$$

where the non-symmetric Biot type stress tensor is given by

$$T_{\text{Biot}} := DW_{\text{mp}}(\bar{U}), \quad (\text{A.8})$$

allows to rewrite the system (A.6) as

$$\begin{aligned} \text{Div} S_1(D\varphi, R) &= 0 \\ \text{skew}[R^T \text{Curl}(R DW_{\text{disloc}}(\boldsymbol{\alpha}))] &= \text{skew}(T_{\text{Biot}}(\bar{U}) \cdot \bar{U}^T + DW_{\text{disloc}}(\boldsymbol{\alpha}) \cdot \boldsymbol{\alpha}^T). \end{aligned} \quad (\text{A.9})$$

Observe that (A.9)₁ is a uniformly elliptic linear system for φ at given R . It is clear that global minimizers $\varphi \in W^{1,2}(\mathcal{U}, \mathbb{R}^3)$ and $R \in W^{1,2}(\mathcal{U}, \text{SO}(3))$ are weak solutions of the Euler-Lagrange equations.

If $DW_{\text{disloc}}(\boldsymbol{\alpha}) \equiv 0$ (no moment stresses) then balance of angular momentum turns into the symmetry constraint

$$DW_{\text{mp}}(\bar{U}) \cdot \bar{U}^T \in \text{Sym}(3). \quad (\text{A.10})$$

A complete discussion of the solutions to this constraint can be found in [55, 56].

A.2 Two-dimensional Euler-Lagrange equations: alternative derivation

$$\begin{aligned} E^{2\text{D}}(m, R) &= \int_{\omega} \mu |\text{dev sym}(R^T(Dm|R_3) - \mathbb{1}_3)|^2 + \mu_c |\text{skew}(R^T(Dm|R_3) - \mathbb{1}_3)|^2 + \frac{\kappa}{3} \text{tr}(R^T(Dm|R_3) - \mathbb{1}_3)^2 + |DR|^2 \, dx \\ &= \int_{\omega} |\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3)|^2 + |DR|^2 \, dx = \int_{\omega} |\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3)|^2 + |\partial_x R|^2 + |\partial_y R|^2 \, dx. \end{aligned} \quad (\text{A.11})$$

Taking free variations w.r.t the midsurface deformation m leads to

$$\begin{aligned} \delta E^{2\text{D}}(m, R) \cdot \delta \vartheta &= \int_{\omega} 2 \langle \mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3), \mathbb{P}(R^T(D\vartheta|0)) \rangle_{\mathbb{R}^{3 \times 3}} \, dx \quad \forall \vartheta \in C_0^\infty(\omega, \mathbb{R}^3) \\ &= \int_{\omega} 2 \langle \mathbb{P}^T \mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3), (R^T(D\vartheta|0)) \rangle_{\mathbb{R}^{3 \times 3}} \, dx = \int_{\omega} \langle 2R \mathbb{P}^T \mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3), (D\vartheta|0) \rangle_{\mathbb{R}^{3 \times 3}} \, dx \\ &= \int_{\omega} \langle \pi_{12}(2R \mathbb{P}^T \mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3)), D\vartheta \rangle_{\mathbb{R}^{3 \times 2}} \, dx = \int_{\omega} \langle \text{Div} \pi_{12}(2R \mathbb{P}^T \mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3)), \vartheta \rangle_{\mathbb{R}^3} \, dx = 0. \end{aligned} \quad (\text{A.12})$$

Thus the strong form of balance of forces can be expressed as

$$\text{Div} S(Dm, R) = 0, \quad (\text{A.13})$$

where

$$S(Dm, R) = \pi_{12}(2R\mathbb{P}^T\mathbb{P}(R^T(Dm|R_3)) - \mathbb{1}_3) = \pi_{12}(2R\mathbb{P}^T\mathbb{P}(R^T(Dm|0) - (\mathbb{1}_2|0))), \quad (\text{A.14})$$

is the *first Piola-Kirchhoff type force-stress tensor* and

$$\begin{aligned} T(Dm, R) &= 2\mathbb{P}^T\mathbb{P}(R^T(Dm|0) - (\mathbb{1}_2|0)) = 2\mathbb{P}^T\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3) \\ &= 2\mu \operatorname{dev} \operatorname{sym}(\bar{U} - \mathbb{1}_3) + 2\mu_c \operatorname{skew}(\bar{U} - \mathbb{1}_3) + \frac{2\kappa}{3} \operatorname{tr}(\bar{U} - \mathbb{1}_3) \cdot \mathbb{1}_3, \quad \bar{U} = R^T(Dm|R_3), \end{aligned} \quad (\text{A.15})$$

is the *non-symmetric Biot-type stress tensor* (symmetric if $\mu_c = 0$). We note the relation

$$S(Dm, R) = \pi_{12}(R \cdot T(Dm, R)). \quad (\text{A.16})$$

resembling relation (A.7).

For balance of angular momentum we proceed similarly, but need some preparation. It is clear that

$$(R + \delta R)^T(Dm|(R + \delta R)e_3) - \mathbb{1}_3 = R^T(Dm|R_3) - \mathbb{1}_3 + \underbrace{R^T(0|0|\delta R_3) + (\delta R^T(Dm|R_3))}_{\text{linear increment}} + \delta R^T(0|0|\delta R_3). \quad (\text{A.17})$$

Therefore, taking variations of the energy w.r.t R leads to

$$\delta E^{2D}(m, R) \cdot \delta R = \int_{\omega} 2\langle \mathbb{P}^T\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3), \mathbb{P}(R^T(0|0|\delta R_3) + \delta R^T(Dm|R_3)) \rangle + 2\langle \partial_x R, \partial_x \delta R \rangle + 2\langle \partial_y R, \partial_y \delta R \rangle \, dx = 0. \quad (\text{A.18})$$

Since $R^T R = \mathbb{1}_3$, we have $\delta R^T R + R^T \delta R = 0$, hence $\delta R = RA$ for $A \in \mathfrak{so}(3)$ arbitrary. Therefore the latter turns into

$$\begin{aligned} &\int_{\omega} 2\langle \mathbb{P}^T\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3), R^T(0|0|(RA)e_3) + (RA)^T(Dm|R_3) \rangle - 2\langle \partial_x^2 R, RA \rangle - 2\langle \partial_y^2 R, RA \rangle \, dx \\ &= \int_{\omega} 2\langle \mathbb{P}^T\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3), A(0|0|e_3) - AR^T(Dm|R_3) \rangle - \langle 2\Delta R, RA \rangle \, dx \\ &= \int_{\omega} 2\langle \mathbb{P}^T\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3), -A(R^T(Dm|R_3) - (0|0|e_3)) \rangle - \langle 2R^T \Delta R, A \rangle \, dx, \quad \forall A \in C_0^\infty(\omega, \mathfrak{so}(3)) \\ &= - \int_{\omega} 2\langle \mathbb{P}^T\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3), AR^T(Dm|0) \rangle + \langle 2R^T \Delta R, A \rangle \, dx \\ &= - \int_{\omega} \langle 2\mathbb{P}^T\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3)(Dm|0)^T R, A \rangle + \langle 2R^T \Delta R, A \rangle \, dx = 0. \end{aligned} \quad (\text{A.19})$$

This implies the stationary condition in strong form

$$\begin{aligned} \operatorname{skew}(2R^T \Delta R) &= -\operatorname{skew}(2\mathbb{P}^T\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3) \cdot (Dm|0)^T R) = -\operatorname{skew}(2\mathbb{P}^T\mathbb{P}(R^T(Dm|0) - (\mathbb{1}_2|0)) \cdot (Dm|0)^T R) \\ &= -\operatorname{skew}(T(Dm|R) \cdot (Dm|0)^T R), \end{aligned} \quad (\text{A.20})$$

where T is defined in (A.15)₁.

For $\mathbb{P}^T\mathbb{P} = \mu \cdot \mathbb{1}$ the last equation simplifies to

$$\begin{aligned} \operatorname{skew}(2R^T \Delta R) &= -\mu \operatorname{skew} \left(2R^T(Dm|0)(Dm|0)^T R - (\mathbb{1}_2|0)(Dm|0)^T R \right) = 2\mu \operatorname{skew} \left((\mathbb{1}_2|0)(Dm|0)^T R \right) \\ &= 2\mu \operatorname{skew}((Dm|0)^T R) = -2\mu \operatorname{skew}(R^T(Dm|0)). \end{aligned} \quad (\text{A.21})$$

We can also rewrite (A.19)₅ as

$$\begin{aligned} 0 &= \int_{\omega} 2\langle \mathbb{P}^T\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3)(Dm|0)^T, AR^T \rangle + \langle 2\Delta R, RA \rangle \, dx \\ &= \int_{\omega} 2\left\langle \left(\mathbb{P}^T\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3)(Dm|0)^T \right)^T, RA^T \right\rangle + \langle 2\Delta R, RA \rangle \, dx \\ &= \int_{\omega} 2\left\langle (Dm|0) \left(\mathbb{P}^T\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3) \right)^T, RA \right\rangle + \langle 2\Delta R, RA \rangle \, dx, \quad \forall A \in C_0^\infty(\omega, \mathfrak{so}(3)) \end{aligned} \quad (\text{A.22})$$

being equivalent to

$$\Delta R - (Dm|0) \left(\mathbb{P}^T\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3) \right)^T \perp \mathbb{T}_R \operatorname{SO}(3). \quad (\text{A.23})$$

Since

$$\left(\mathbb{P}^T\mathbb{P}(R^T(Dm|R_3) - \mathbb{1}_3) \right)^T = \left(\mathbb{P}^T\mathbb{P}(R^T(Dm|0) - (\mathbb{1}_2|0)) \right)^T = \mathbb{P}^T\mathbb{P}((R^T(Dm|0)(\mathbb{1}_2|0))^T),$$

we may express (A.23) also as

$$\Delta R - (Dm|0)\mathbb{P}^T\mathbb{P}((Dm|0)^T R - (\mathbb{1}_2|0)) \perp \mathbb{T}_R \operatorname{SO}(3). \quad (\text{A.24})$$

This is the form for balance of angular momentum given in equation (5.8).

A.3 Lifting to the Δ -operator

This last equation (A.24) is not, however, the final form of the balance of angular momentum equation that we will consider. Indeed, since $R^T R = \mathbb{1}_3$, we can differentiate once to obtain

$$(\partial_x R)^T R + R^T \partial_x R = 0, \quad (\partial_y R)^T R + R^T \partial_y R = 0. \quad (\text{A.25})$$

Taking second partial derivatives, we get

$$\begin{aligned} (\partial_x^2 R)^T R + (\partial_x R)^T \partial_x R + (\partial_x R)^T \partial_x R + R^T \partial_x^2 R &= 0, \\ (\partial_y^2 R)^T R + (\partial_y R)^T \partial_y R + (\partial_y R)^T \partial_y R + R^T \partial_y^2 R &= 0. \end{aligned} \quad (\text{A.26})$$

Summing up, shows

$$\begin{aligned} (\Delta R)^T R + R^T \Delta R + 2((\partial_x R)^T \partial_x R + (\partial_y R)^T \partial_y R) &= 0, \\ \iff 2 \text{sym}(R^T \Delta R) + 2[(\partial_x R)^T \partial_x R + (\partial_y R)^T \partial_y R] &= 0, \\ \text{sym}(R^T \Delta R) &= -[(\partial_x R)^T \partial_x R + (\partial_y R)^T \partial_y R]. \end{aligned} \quad (\text{A.27})$$

From (A.20) we have

$$\text{skew}(2R^T \Delta R) = -\text{skew}(T(Dm, R)(Dm|0)^T R). \quad (\text{A.28})$$

Adding (A.27)₂ and (A.28) yields, due to the orthogonality of sym and skew

$$2R^T \Delta R = \text{sym}(2R^T \Delta R) + \text{skew}(2R^T \Delta R) = -2[(\partial_x R)^T \partial_x R + (\partial_y R)^T \partial_y R] - \text{skew}(T(Dm, R)(Dm|0)^T R). \quad (\text{A.29})$$

Hence using the isotropy of skew we obtain

$$\begin{aligned} 2\Delta R &= -2R[(\partial_x R)^T \partial_x R + (\partial_y R)^T \partial_y R] - R \text{skew}(T(Dm, R)(Dm|0)^T R) \\ &= -2R[(\partial_x R)^T \partial_x R + (\partial_y R)^T \partial_y R] - R \text{skew}(R^T R T(Dm, R)(Dm|0)^T R) \\ &= -2R[(\partial_x R)^T \partial_x R + (\partial_y R)^T \partial_y R] - R R^T \text{skew}(R T(Dm, R)(Dm|0)^T) \cdot R \\ &= -2R[(\partial_x R)^T \partial_x R + (\partial_y R)^T \partial_y R] + \text{skew}((Dm|0)(R T(Dm, R))^T) \cdot R \\ &= -2R[(\partial_x R)^T \partial_x R + (\partial_y R)^T \partial_y R] + \text{skew}((Dm|0)[(\pi_{12}(R T(Dm, R)))]^T) \cdot R \\ &= -2R[(\partial_x R)^T \partial_x R + (\partial_y R)^T \partial_y R] + \text{skew}(Dm \cdot S(Dm, R)^T) \cdot R, \end{aligned} \quad (\text{A.30})$$

giving

$$\begin{aligned} \Delta R &= -R[(\partial_x R)^T \partial_x R + (\partial_y R)^T \partial_y R] + \frac{1}{2} \text{skew}(Dm \cdot S(Dm, R)^T) \cdot R \\ &= -R[(\partial_x R)^T \partial_x R + (\partial_y R)^T \partial_y R] + \text{skew}(Dm \circ S(Dm, R)) \cdot R. \end{aligned} \quad (\text{A.31})$$

where we used the definition of \circ given in equation (5.9).

We set

$$-R[(\partial_x R)^T \partial_x R + (\partial_y R)^T \partial_y R] = -R \partial_x R^T \cdot \partial_x R - R \partial_y R^T \partial_y R =: \Omega_R \cdot DR. \quad (\text{A.32})$$

$$(\Omega_R)_1 := -R \partial_x R^T \in \mathfrak{so}(3), \quad (\Omega_R)_2 := -R \partial_y R^T \in \mathfrak{so}(3). \quad (\text{A.33})$$

With this definition, (A.30) can be written as

$$\Delta R = \underbrace{\Omega_R \cdot DR}_{\in L^1(\omega)} - \underbrace{R \text{skew}(T(Dm, R)(Dm|0)^T R)}_{\in L^1(\omega)}. \quad (\text{A.34})$$

Considering the special case $\mathbb{P}^T \mathbb{P} = \mu \cdot \mathbb{1}$, equation (A.34) turns into

$$\Delta R = \Omega_R \cdot DR + \mu R \text{skew}((Dm|0)^T R) = \Omega_R \cdot DR - \underbrace{\mu R \text{skew}(R^T (Dm|0))}_{\in L^2(\omega)}. \quad (\text{A.35})$$

We observe finally, that

$$R^T (\Omega_R)_i R = -(\partial_i R)^T R = R^T \partial_i R, \quad i = 1, 2, \quad (\text{A.36})$$

which implies for $\Gamma_i = \text{axl}(R^T \partial_i R)$

$$R^T (\Omega_R)_i R = \text{Anti}(\Gamma_i), \quad \text{axl}(R^T (\Omega_R)_i R) = \Gamma_i, \quad (\text{A.37})$$

where Γ is the wryness tensor from eq. (4.3).

A.4 A glimpse on a Reissner-Mindlin type flat membrane shell model

It is interesting to compare our Cosserat flat membrane shell model (allowing for existence of minimizers and their full regularity) with one that would appear closer to classical approaches. For this sake we consider a Reissner-Mindlin flat membrane shell model next.

In case of the one-director geometrically nonlinear, physically linear Reissner-Mindlin flat membrane shell model without independent drilling rotations, the problem can be described as a two-field minimization for the midsurface $m: \omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$ and the unit-director field $d: \omega \subset \mathbb{R}^2 \rightarrow \mathbb{S}^2$ of the elastic energy⁹

$$E_{\text{Reissner}}^{2\text{D}}(m, d) = \int_{\omega} h \left\{ \underbrace{\frac{\mu}{4} |\text{D}m^T \text{D}m - \mathbb{1}_2|^2}_{\text{in-plane stretch}} + \frac{1}{8} \frac{2\mu\lambda}{2\mu + \lambda} \text{tr} \left(\text{D}m^T \text{D}m - \mathbb{1}_2 \right)^2}_{\text{elongational stretch}} + \underbrace{\mu \left(\langle d, \partial_x m \rangle^2 + \langle d, \partial_y m \rangle^2 \right)}_{\text{transverse shear}} + \underbrace{\mu \frac{L_c^2}{2} |\nabla d|^2}_{\text{curvature}} \right\} dx. \quad (\text{A.39})$$

Here, the membrane energy part is not rank-one elliptic due to the presence of the membrane strain $\text{D}m^T \text{D}m - \mathbb{1}_2$. The uni-constant curvature energy could be generalized to the Oseen-Frank form, cf. subsection 3.1. We note that $\langle d, \partial_x m \rangle^2 + \langle d, \partial_y m \rangle^2 = |\text{D}m^T d|_{\mathbb{R}^2}^2$, and look for simplicity at the energy

$$\int_{\omega} \frac{1}{2} |\text{D}m^T \text{D}m - \mathbb{1}_2|_{\mathbb{R}^{2 \times 2}}^2 + \frac{1}{2} |\text{D}m^T d|_{\mathbb{R}^2}^2 + \frac{1}{2} |\nabla d|^2 = \int_{\omega} \frac{1}{2} |(\text{D}m|n_m)^T (\text{D}m|n_m) - \mathbb{1}_3|^2 + \frac{1}{2} |\text{D}m^T d|^2 + \frac{1}{2} |\nabla d|^2 dx. \quad (\text{A.40})$$

The Euler-Lagrange equations are then given by

$$\begin{aligned} \delta E_{\text{Reissner}}^{2\text{D}}(m, d) \cdot \delta m &= \int_{\omega} 2 \langle (\text{D}m|n_m)^T (\text{D}m|n_m) - \mathbb{1}_3, (\text{D}\delta m|0) \rangle_{\mathbb{R}^{3 \times 3}} + \langle \text{D}m^T d, (\text{D}\delta m)^T d \rangle_{\mathbb{R}^2} dx \\ &= \int_{\omega} \langle (\text{D}m|n_m) \cdot 2 \langle (\text{D}m|n_m)^T (\text{D}m|n_m) - \mathbb{1}_3, (\text{D}\delta m|0) \rangle_{\mathbb{R}^{3 \times 3}} \rangle_{\mathbb{R}^3} dx + \langle \text{D}m^T d \otimes \text{D}\delta m^T d, \mathbb{1}_2 \rangle_{\mathbb{R}^{2 \times 2}} dx \\ &= \int_{\omega} \langle (\text{D}m|n_m) \cdot 2 \langle (\text{D}m|n_m)^T (\text{D}m|n_m) - \mathbb{1}_3, (\text{D}\delta m|0) \rangle_{\mathbb{R}^{3 \times 3}} \rangle_{\mathbb{R}^3} dx + \langle (\text{D}m^T d \otimes d) \text{D}\delta m, \mathbb{1}_2 \rangle_{\mathbb{R}^{2 \times 2}} dx \\ &= \int_{\omega} \langle \pi_{12}((\text{D}m|n_m) \cdot 2 \langle (\text{D}m|n_m)^T (\text{D}m|n_m) - \mathbb{1}_3 \rangle) + d \otimes \text{D}m^T d, \text{D}\delta m \rangle_{\mathbb{R}^{3 \times 2}} dx \\ &= - \int_{\omega} \langle \text{Div} \left[\pi_{12}((\text{D}m|n_m) \cdot 2 \langle (\text{D}m|n_m)^T (\text{D}m|n_m) - \mathbb{1}_3 \rangle) + d \otimes \text{D}m^T d \right], \delta m \rangle_{\mathbb{R}^3} dx = 0 \quad \forall \delta m \in C_0^\infty(\omega, \mathbb{R}^3). \end{aligned} \quad (\text{A.41})$$

Here, we can define the first Piola-Kirchhoff type stress tensor

$$S(\text{D}m, d) = \pi_{12}((\text{D}m|n_m) \cdot 2 \langle (\text{D}m|n_m)^T (\text{D}m|n_m) - \mathbb{1}_3 \rangle + d \otimes \text{D}m^T d). \quad (\text{A.42})$$

For variations w.r.t $d \in \mathbb{S}^2$ we note that

$$|d + \delta d|^2 = 1 \iff |d|^2 + 2\langle d, \delta d \rangle + |\delta d|^2 = 1. \quad (\text{A.43})$$

Hence, the variation δd is orthogonal to d , i.e. $\langle d, \delta d \rangle = 0$. Without loss of generality, we express δd as $\delta d = d \times \delta v$ for some $\delta v \in C_0^\infty(\omega, \mathbb{R}^3)$. Therefore, taking variations w.r.t. d gives

$$\begin{aligned} \delta E_{\text{Reissner}}^{2\text{D}}(m, d) \cdot \delta d &= \int_{\omega} \langle \nabla d, \nabla \delta d \rangle + \langle \text{D}m^T d, \text{D}m^T \delta d \rangle dx = 0 \iff - \int_{\omega} \langle \Delta d, d \times \delta v \rangle + \langle \text{D}m \text{D}m^T d, d \times \delta v \rangle dx = 0 \\ &\iff \int_{\omega} - \langle \Delta d, \text{Anti}(d) \delta v \rangle + \langle \text{D}m \text{D}m^T d, \text{Anti}(d) \delta v \rangle dx = 0 \iff \int_{\omega} \langle \text{Anti}(d) \Delta d, \delta v \rangle_{\mathbb{R}^3} - \langle \text{Anti}(d) \text{D}m \text{D}m^T d, \delta v \rangle dx = 0, \\ &\quad \forall \delta v \in C_0^\infty(\omega, \mathbb{R}^3). \end{aligned} \quad (\text{A.44})$$

The latter leads to the strong form

$$\text{Anti}(d) \left(\Delta d - \text{D}m \text{D}m^T d \right) = 0 \iff d \times (\Delta d - \text{D}m \text{D}m^T d) = 0. \quad (\text{A.45})$$

However, since $d \in \mathbb{S}^2$, we know $|d|^2 = 1$. Therefore, in addition, taking partial derivatives, we obtain

$$\langle \partial_x d, d \rangle = 0, \quad \langle \partial_y d, d \rangle = 0. \quad (\text{A.46})$$

Taking second partial derivatives yields

$$\langle \partial_x^2 d, d \rangle + \langle \partial_x d, \partial_y d \rangle = 0, \quad \langle \partial_y^2 d, d \rangle + \langle \partial_y d, \partial_x d \rangle = 0. \quad (\text{A.47})$$

⁹The missing Reissner-Mindlin bending contribution scaling with h^3 would be of the form [48, (7.25)]

$$\frac{h^3}{12} \left\{ \mu |\text{sym} \left((\text{D}m|d)^T (\text{D}d|0) \right)|^2 + \frac{\mu\lambda}{2\mu + \lambda} \text{tr} \left(\text{sym} \left((\text{D}m|d)^T (\text{D}d|0) \right) \right)^2 \right\}. \quad (\text{A.38})$$

Here, no choice of constitutive parameters reduces the bending energy to the uni-constant case.

Summing up shows

$$\langle \Delta d, d \rangle + |\partial_x d|^2 + |\partial_y d|^2 = \langle \Delta d, d \rangle + |Dd|^2 = 0. \quad (\text{A.48})$$

Adding (A.45) and (A.48) shows

$$d * \Delta d := \underbrace{d \times \Delta d + \langle d, \Delta d \rangle}_{\text{geometric product, Clifford product}} = \underbrace{d \times (Dm Dm^T d)}_{\in \mathbb{R}^3} - \underbrace{|Dd|^2}_{\in \mathbb{R}}. \quad (\text{A.49})$$

Formally, this implies

$$\Delta d = \frac{d}{|d|^2} * \left[d \times (Dm Dm^T d) - |Dd|^2 \right]. \quad (\text{A.50})$$

In terms of equations, we have altogether from (A.45)₁ and (A.48), respectively

$$\text{Anti}(d) \Delta d = Dm Dm^T d, \quad \langle d, \Delta d \rangle = -|Dd|^2, \quad (\text{A.51})$$

equivalently

$$\underbrace{\begin{pmatrix} & \text{Anti}(d) & \\ d_1 & d_2 & d_3 \end{pmatrix}}_{=: \widehat{A} \in \mathbb{R}^{4 \times 3}} \begin{pmatrix} \Delta d \\ | \end{pmatrix} = \begin{pmatrix} Dm Dm^T d \\ -|Dd|^2 \end{pmatrix}_{\mathbb{R}^4}. \quad (\text{A.52})$$

We multiply (A.52) with \widehat{A}^T to get

$$\widehat{A}^T \widehat{A} \Delta d = \widehat{A}^T \begin{pmatrix} Dm Dm^T d \\ -|Dd|^2 \end{pmatrix} \in \mathbb{R}^3, \quad |d| = 1. \quad (\text{A.53})$$

Since in fact (sic)

$$\widehat{A}^T \widehat{A} = |d|^2 \cdot \mathbb{1}_3 = \mathbb{1}_3,$$

we obtain the system of Euler-Lagrange equations

$$\text{Div } S(Dm, d) = 0, \quad \text{"balance of forces"} \quad (\text{A.54})$$

with a first Piola-Kirchhoff type stress tensor

$$S(Dm, d) = \pi_{12}((Dm|n_m) \cdot 2((Dm|n_m)^T(Dm|n_m) - \mathbb{1}_3) + d \otimes Dm^T d), \quad (\text{A.55})$$

and

$$\begin{aligned} \Delta d &= \begin{pmatrix} -\text{Anti}(d)|d \end{pmatrix}_{\mathbb{R}^{3 \times 4}} \begin{pmatrix} Dm Dm^T d \\ -|Dd|^2 \end{pmatrix} \\ &= -\text{Anti}(d) (Dm Dm^T d) - |Dd|^2 \cdot d, \quad \text{"balance of director equilibrium"} \\ &= -\underbrace{d \times (Dm Dm^T d)}_{\in L^2(\omega, \mathbb{R}^3)} - |Dd|^2 \cdot d. \end{aligned} \quad (\text{A.56})$$

We observe that (A.54) constitutes a nonlinear, nonconvex problem for the midsurface m once the unit director d is determined. Therefore, existence to (A.54,A.56) is not yet known and likely not true. We note that the right-hand side in (A.56) contains an $L^2(\omega)$ -term, since $Dm \in L^4(\omega)$ instead of our $L^1(\omega)$ -term in equation (A.34). For $Dm \equiv 0$ we recover from (A.56) the director equilibrium for the uni-constant liquid crystal problem equation (3.9).

A.5 Numerical experiments

We present a sequence of numerical experiments for the problem (A.11). In these experiments we compare the dimensionally reduced energy (4.18) (where the transverse shear energy is multiplied by the arithmetic mean of μ and μ_c) to the energy (4.19) of the rigorously derived Γ -limit membrane model (where the transverse shear energy is multiplied by the harmonic mean of μ and μ_c). We set the Lamé parameters to $\mu = 2.7191 \cdot 10^4$, $\lambda = 4.4364 \cdot 10^4$, and vary μ_c and L_c .

For the domain we choose the unit disk, which we discretized by $6 \cdot 4^6 = 24576$ triangular elements. We used Lagrange finite elements of second order for the midsurface deformation m and geodesic finite elements of second order for the microrotation field R [65, 67].

To trigger the deformation process, we radially compressed the membrane to a new radius $r < 1$ by Dirichlet boundary conditions for the deformation on the entire domain boundary. The microrotation field was not subject to Dirichlet boundary conditions at all. We minimized the discrete energy using a trust-region method [67] starting from the cap function $m_0(x, y) = (x, y, 0.1 - 0.1\sqrt{x^2 + y^2})$ for all (x, y) in the interior of the unit disk and $m_0(x, y) = (rx, ry, 0)$ on the boundary. The initial microrotation was $R = \mathbb{1}$. We conducted several simulations resulting in different wrinkle patterns depending on the Cosserat couple modulus μ_c and the characteristic length L_c as shown in Figures 4, 5 and 6. The numerical algorithms were implemented in C++ using the DUNE libraries (www.dune-project.org) [66].

From the figures one can see that wrinkling only happens if the characteristic length L_c is small enough. Indeed, if $L_c = 10^{-3}$ then the deformation is largely bending-dominated, with small wrinkles only appearing next to the boundary, if μ_c is large enough. With smaller values for L_c one can see wrinkling in larger parts of the domain, even if the radial compression factor r is much smaller. Note that the choice $\mu_c = 0$ does not lead to a well-posed problem when used in the energy (4.19), because there it makes the transverse shear energy term disappear.







	Arithmetic Mean (4.18)	Harmonic Mean (4.19)
$\mu_c = 0$		not well-posed
$\mu_c = 10^{-5}\mu$		
$\mu_c = 10^{-2}\mu$		
$\mu_c = \mu$		

Figure 4: Deformation of a radially compressed shell with $L_c = 10^{-3}$. Simulations by Lisa Julia Nebel and Oliver Sander (TU Dresden).





	Arithmetic Mean (4.18)	Harmonic Mean (4.19)
$\mu_c = 0$	 $r = 0.98$	not well-posed
$\mu_c = 10^{-2}\mu$	 $r = 0.91$	 $r = 0.94$
$\mu_c = \mu$	 $r = 0.9$	

Figure 5: Deformation of a radially compressed shell with $L_c = 10^{-5}$. Simulations by Lisa Julia Nebel and Oliver Sander (TU Dresden).





	Arithmetic Mean (4.18)	Harmonic Mean (4.19)
$\mu_c = 0$	 $r = 0.99$	not well-posed
$\mu_c = 10^{-2}\mu$	 $r = 0.99$	 $r = 0.99$
$\mu_c = \mu$	 $r = 0.99$	

Figure 6: Deformation of a radially compressed shell with $L_c = 10^{-8}$. Simulations by Lisa Julia Nebel and Oliver Sander (TU Dresden).