

Signless Laplacian energies of non-commuting graphs of finite groups and related results

Monalisha Sharma and Rajat Kanti Nath*

*Department of Mathematical Sciences, Tezpur University,
Napaam-784028, Sonitpur, Assam, India.*

Emails: monalishasharma2013@gmail.com and rajatkantinath@yahoo.com

Abstract

The non-commuting graph of a non-abelian group G with center $Z(G)$ is a simple undirected graph whose vertex set is $G \setminus Z(G)$ and two vertices x, y are adjacent if $xy \neq yx$. In this study, we compute Signless Laplacian spectrum and Signless Laplacian energy of non-commuting graphs of finite groups. We obtain several conditions such that the non-commuting graph of G is Q-integral and observe relations between energy, Signless Laplacian energy and Laplacian energy. In addition, we look into the energetic hyper- and hypo-properties of non-commuting graphs of finite groups. We also assess whether the same graphs are Q-hyperenergetic and L-hyperenergetic.

Key words: Non-commuting graph, Spectrum, Energy.

2010 Mathematics Subject Classification: 20D60, 05C50, 15A18, 05C25.

1 Introduction

Recall that for a graph \mathcal{G} , its spectrum denoted by $\text{Spec}(\mathcal{G})$ is $\{(\lambda_1)^{k_1}, (\lambda_2)^{k_2}, \dots, (\lambda_n)^{k_n}\}$, where λ_i 's are the eigenvalues of the adjacency matrix of \mathcal{G} with multiplicities k_i for $1 \leq i \leq n$, respectively. Let $A(\mathcal{G})$ and $D(\mathcal{G})$ be the adjacency matrix and degree matrix of \mathcal{G} , respectively. Consequently, $L(\mathcal{G}) = D(\mathcal{G}) - A(\mathcal{G})$ gives the Laplacian matrix of \mathcal{G} . The set $\{(\beta_1)^{b_1}, (\beta_2)^{b_2}, \dots, (\beta_m)^{b_m}\}$, where β_j 's are the eigenvalues of $L(\mathcal{G})$ with multiplicities b_j for $1 \leq j \leq m$, respectively is referred to as the Laplacian spectrum of \mathcal{G} and is denoted by $\text{L-spec}(\mathcal{G})$. While, $Q(\mathcal{G}) = A(\mathcal{G}) - D(\mathcal{G})$ provides the Signless Laplacian matrix of \mathcal{G} . The set $\{(\gamma_1)^{d_1}, (\gamma_2)^{d_2}, \dots, (\gamma_l)^{d_l}\}$, where γ_r 's are the eigenvalues of $Q(\mathcal{G})$ with multiplicities d_r for $1 \leq r \leq l$, respectively is known as the Signless Laplacian spectrum of \mathcal{G} and is denoted by $\text{Q-spec}(\mathcal{G})$. A graph \mathcal{G} is called integral, L-integral and Q-integral if eigenvalues of $A(\mathcal{G})$, $L(\mathcal{G})$ and $Q(\mathcal{G})$ are integers.

The energy, Laplacian energy and Signless Laplacian energy of \mathcal{G} are given by

$$E(\mathcal{G}) = \sum_{\lambda \in \text{Spec}(\mathcal{G})} |\lambda|, \quad LE(\mathcal{G}) = \sum_{\beta \in \text{L-spec}(\mathcal{G})} \left| \beta - \frac{2|e(\mathcal{G})|}{|v(\mathcal{G})|} \right| \quad \text{and} \quad LE^+(\mathcal{G}) = \sum_{\gamma \in \text{Q-spec}(\mathcal{G})} \left| \gamma - \frac{2|e(\mathcal{G})|}{|v(\mathcal{G})|} \right|$$

respectively, where $v(\mathcal{G})$ and $e(\mathcal{G})$ are the set of vertices and edges of \mathcal{G} respectively. A finite graph \mathcal{G} is hyperenergetic if $E(\mathcal{G}) > E(K_{|v(\mathcal{G})|})$ and is hypoenergetic if $E(\mathcal{G}) < |v(\mathcal{G})|$. Likewise, the graph \mathcal{G} is L-hyperenergetic if $LE(\mathcal{G}) > LE(K_{|v(\mathcal{G})|})$ and is Q-hyperenergetic if $LE^+(\mathcal{G}) > LE^+(K_{|v(\mathcal{G})|})$. Gutman [20] and Walikar et. al [31] pioneered the research of hyperenergetic graphs in 1999. Gutman [23] first presented the hypoenergetic graph in 2007. L-hyperenergetic and Q-hyperenergetic graphs were considered in [15].

Assume that G is a finite non-abelian group and that $Z(G) = \{z \in G : zx = xz, \forall x \in G\}$ is the centre of G . The non-commuting graph of G , represented by Γ_G , is a basic undirected network connecting two unique vertices x and y whenever $xy \neq yx$ with $G \setminus Z(G)$ as the vertex set. Erdős and Neumann's study [27] considered this graph for the first time in 1976. [1, 4, 6, 16, 25, 30] are some references on non-commuting graphs of finite groups. Spectral aspects of this graph have also grabbed attention of numerous mathematicians. In [18], Ghorbani et al. calculated spectrum of Γ_G for certain groups. The energy of the same was later explored by Ghorbani and Gharavi-Alkhansari in [17]. In [8], Dutta et al. computed the Laplacian spectrum of Γ_G following which Dutta and Nath [12] worked on Laplacian energy of the same. Signless Laplacian spectrum and energy of Γ_G are not yet computed. At present, [2] is the only paper where Abdussakir et al. investigated the Signless Laplacian spectrum of non-commuting graphs of dihedral groups. However, various spectra and energies (including Signless Laplacian spectrum and energy) of the complement of Γ_G , known as commuting

*Corresponding author

graph of G , are already computed in [7, 9, 10, 11, 13, 14, 28]. Various spectra and energies of non-commuting conjugacy class graphs of several families of finite groups are computed in [26]. In this paper, we consider the following questions and answer them up to some extent.

Question 1.1. *Which finite non-abelian groups give Q -integral non-commuting graphs?*

Question 1.2. *Are there any finite non-abelian group G such that Γ_G is hypoenergetic, hyperenergetic, L -hyperenergetic and Q -hyperenergetic?*

Question 1.3. *Which finite non-abelian groups satisfy the following inequalities?*

- (a) $E(\Gamma_G) \leq LE(\Gamma_G)$.
- (b) $LE(\Gamma_G) \leq LE^+(\Gamma_G)$.

We compute $Q\text{-spec}(\Gamma_G)$ and determine several conditions such that Γ_G is Q -integral. We also compute $LE^+(\Gamma_G)$ for many families of finite non-abelian groups and compare various energies of Γ_G . These comparisons are interesting because mathematicians want to know about the graphs \mathcal{G} such that $E(\mathcal{G}) \leq LE(\mathcal{G})$ (see [7, 22, 24, 29]) and $LE(\mathcal{G}) \leq LE^+(\mathcal{G})$ (see [5, 7]). Using energies of Γ_G for various classes of finite groups we determine whether they are hyperenergetic or hypoenergetic. In [19], Gutman posed Conjecture 1.4 which was disproved by different mathematicians providing counter examples (see [21]).

Conjecture 1.4. *Any finite graph $\mathcal{G} \not\cong K_{|v(\mathcal{G})|}$ is non-hyperenergetic.*

It is worth mentioning that we also disprove Conjecture 1.4 by considering non-commuting graphs of finite groups (see Theorem 6.3(b)). We shall also determine finite groups such that their non-commuting graphs are Q -hyperenergetic and L -hyperenergetic.

We conclude this section with the following results which will help us to compute spectrum, Laplacian spectrum and Signless Laplacian spectrum of Γ_G in the succeeding sections.

Theorem 1.5 ([8, Corollary 2.3]). *Let \mathcal{G} be a graph and $\mathcal{G} = l_1 K_{m_1} \cup l_2 K_{m_2} \cup \dots \cup l_k K_{m_k}$, where $l_i K_{m_i}$ denotes the disjoint union of l_i copies of K_{m_i} for $1 \leq i \leq k$ and $m_1 < m_2 < \dots < m_k$. Then*

$$\text{L-spec}(\mathcal{G}^c) = \left\{ (0)^1, \left(\sum_{i=1}^k l_i m_i - m_k \right)^{l_k(m_k-1)}, \left(\sum_{i=1}^k l_i m_i - m_{k-1} \right)^{l_{k-1}(m_{k-1}-1)}, \dots, \right. \\ \left. \left(\sum_{i=1}^k l_i m_i - m_1 \right)^{l_1(m_1-1)}, \left(\sum_{i=1}^k l_i m_i \right)^{\sum_{i=1}^k l_i - 1} \right\}.$$

Theorem 1.6 ([32, Corollary 2.3] and [33, Corollary 2.2]). *Let \mathcal{G} be the complete r -partite graph $K_{\underbrace{p_1, \dots, p_1}_{a_1\text{-times}}, \underbrace{p_2, \dots, p_2}_{a_2\text{-times}}, \dots, \underbrace{p_s, \dots, p_s}_{a_s\text{-times}}}$ on n vertices, where $r = a_1 + a_2 + \dots + a_s$. Then*

- (a) *the characteristic polynomial of $A(\mathcal{G})$ is*

$$P_{\mathcal{G}}(x) := x^{n-r} \prod_{i=1}^s (x + p_i)^{a_i - 1} \left(\prod_{i=1}^s (x + p_i) - \sum_{j=1}^s a_j p_j \prod_{i=1, i \neq j}^s (x + p_i) \right).$$

- (b) *the characteristic polynomial of $Q(\mathcal{G})$ (Q -polynomial) is*

$$Q_{\mathcal{G}}(x) := \prod_{i=1}^s (x - n + p_i)^{a_i(p_i-1)} \prod_{i=1}^s (x - n + 2p_i)^{a_i} \left(1 - \sum_{i=1}^s \frac{a_i p_i}{x - n + 2p_i} \right).$$

2 $\frac{G}{Z(G)}$ is isomorphic to D_{2m}

Here, we primarily compute the Signless Laplacian spectrum and Signless Laplacian energy of Γ_G , where G is isomorphic to D_{2m} , QD_{2n} , M_{2rs} , Q_{4n} and U_{6n} . Further we compare different energies of Γ_G and look into the hyper- and hypo-energetic properties of Γ_G for each of the above-mentioned groups. The energy and Laplacian energy of Γ_G for each of the aforementioned groups have already been determined and it is crucial to note.

2.1 The dihedral groups, D_{2m}

We consider $D_{2m} := \langle a, b : a^m = b^2 = 1; bab^{-1} = a^{-1} \rangle$, the dihedral groups of order $2m$ (where $m > 2$). Results regarding different energies of non-commuting graphs of D_{2m} are given below.

Theorem 2.1 ([14, Corollary 4.1.7 and (4.3.e)]). *Let G be isomorphic to D_{2m} .*

(a) *If m is odd then*

$$E(\Gamma_{D_{2m}}) = (m-1) + \sqrt{(m-1)(5m-1)} \quad \text{and} \quad LE(\Gamma_{D_{2m}}) = \frac{2m(m-1)(m-2) + 2m(2m-1)}{2m-1}.$$

(b) *If m is even then*

$$E(\Gamma_{D_{2m}}) = (m-2) + \sqrt{(m-2)(5m-2)} \quad \text{and} \quad LE(\Gamma_{D_{2m}}) = \frac{m(m-2)(m-4) + 2m(m-1)}{m-1}.$$

Theorem 2.2. *Let G be isomorphic to D_{2m} , where m is odd. Then*

$$\text{Q-spec}(\Gamma_{D_{2m}}) = \left\{ (m)^{m-2}, (2m-3)^{m-1}, \left(\frac{4m-3 + \sqrt{8m^2-16m+9}}{2} \right)^1, \left(\frac{4m-3 - \sqrt{8m^2-16m+9}}{2} \right)^1 \right\}$$

and

$$LE^+(\Gamma_{D_{2m}}) = \begin{cases} \frac{9}{5} + \sqrt{33}, & \text{if } m = 3 \\ \frac{2m^3-10m^2+12m-3}{2m-1} + \sqrt{8m^2-16m+9}, & \text{if } m \geq 5. \end{cases}$$

Proof. If $G \cong D_{2m}$ and m is odd then $|v(\Gamma_{D_{2m}})| = 2m-1$ and $\Gamma_{D_{2m}} = K_{m,1,1,(m-1)}$. Using Theorem 1.6(b), we have

$$\begin{aligned} Q_{\Gamma_{D_{2m}}}(x) &= \prod_{i=1}^2 (x - (2m-1) + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - (2m-1) + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - (2m-1) + 2p_i} \right) \\ &= (x - (2m-2))^0 (x - m)^{m-2} (x - 2m+3)^m (x-1) \left(1 - \frac{m}{x-2m+3} - \frac{m-1}{x-1} \right) \\ &= (x-m)^{m-2} (x-(2m-3))^{m-1} (x^2 - (4m-3)x + 2m^2 - 2m). \end{aligned}$$

Thus, $\text{Q-spec}(\Gamma_{D_{2m}}) = \left\{ (m)^{m-2}, (2m-3)^{m-1}, \left(\frac{4m-3 + \sqrt{8m^2-16m+9}}{2} \right)^1, \left(\frac{4m-3 - \sqrt{8m^2-16m+9}}{2} \right)^1 \right\}$.

Number of edges in $\Gamma_{D_{2m}}^c$ is $\frac{m^2-3m+2}{2}$. Thus, $|e(\Gamma_{D_{2m}})| = \frac{(2m-1)(2m-1-1)}{2} - \frac{m^2-3m+2}{2} = \frac{3m(m-1)}{2}$. Now

$$\left| m - \frac{2|e(\Gamma_{D_{2m}})|}{|v(\Gamma_{D_{2m}})|} \right| = \left| \frac{-m(m-2)}{2m-1} \right| = \frac{m(m-2)}{2m-1},$$

$$\left| 2m-3 - \frac{2|e(\Gamma_{D_{2m}})|}{|v(\Gamma_{D_{2m}})|} \right| = \left| \frac{m^2-5m+3}{2m-1} \right| = \begin{cases} \frac{3}{5}, & \text{if } m = 3 \\ \frac{m^2-5m+3}{2m-1}, & \text{if } m \geq 5, \end{cases}$$

$$\begin{aligned} \left| \frac{1}{2} \left(4m-3 + \sqrt{8m^2-16m+9} \right) - \frac{2|e(\Gamma_{D_{2m}})|}{|v(\Gamma_{D_{2m}})|} \right| &= \left| \frac{1}{2} \left(\sqrt{8m^2-16m+9} + m - \frac{3}{2} + \frac{3}{4m-2} \right) \right| \\ &= \frac{1}{2} \left(\sqrt{8m^2-16m+9} + m - \frac{3}{2} + \frac{3}{4m-2} \right) \end{aligned}$$

and

$$\begin{aligned} \left| \frac{1}{2} \left(4m-3 - \sqrt{8m^2-16m+9} \right) - \frac{2|e(\Gamma_{D_{2m}})|}{|v(\Gamma_{D_{2m}})|} \right| &= \left| \frac{1}{2} \left(-\sqrt{8m^2-16m+9} + m - \frac{3}{2} + \frac{3}{4m-2} \right) \right| \\ &= \frac{1}{2} \left(\sqrt{8m^2-16m+9} - m + \frac{3}{2} - \frac{3}{4m-2} \right). \end{aligned}$$

Therefore, for $m = 3$ we have $LE^+(\Gamma_{D_{2m}}) = \frac{9}{5} + \sqrt{33}$. For $m \geq 5$ we have

$$\begin{aligned} LE^+(\Gamma_{D_{2m}}) &= (m-2) \times \frac{m(m-2)}{2m-1} + (m-1) \times \frac{m^2-5m+3}{2m-1} + \\ &\quad \frac{1}{2} \left(\sqrt{8m^2-16m+9} + m - \frac{3}{2} + \frac{3}{4m-2} \right) + \frac{1}{2} \left(\sqrt{8m^2-16m+9} - m + \frac{3}{2} - \frac{3}{4m-2} \right) \end{aligned}$$

and the result follows on simplification. \square

Theorem 2.3. *Let G be isomorphic to D_{2m} , where m is even. Then*

$\text{Q-spec}(\Gamma_{D_{2m}})$

$$= \left\{ (2m-4)^{\frac{m}{2}}, (m)^{m-3}, (2m-6)^{\frac{m}{2}-1}, \left(2m-3+\sqrt{2m^2-8m+9}\right)^1, \left(2m-3-\sqrt{2m^2-8m+9}\right)^1 \right\}$$

and

$$LE^+(\Gamma_{D_{2m}}) = \begin{cases} \frac{m^3-4m^2+12}{2m-2} + 2\sqrt{2m^2-8m+9}, & \text{if } 4 \leq m \leq 8 \\ \frac{m^3-8m^2+16m-6}{m-1} + 2\sqrt{2m^2-8m+9}, & \text{if } m \geq 10. \end{cases}$$

Proof. If $G \cong D_{2m}$ and m is even then $|v(\Gamma_{D_{2m}})| = 2m-2$ and $\Gamma_{D_{2m}} = K_{\frac{m}{2}, 2, 1, (m-2)}$. Using Theorem 1.6(b), we have

$$\begin{aligned} Q_{\Gamma_{D_{2m}}}(x) &= \prod_{i=1}^2 (x - (2m-2) + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - (2m-2) + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - (2m-2) + 2p_i} \right) \\ &= (x - (2m-4))^{\frac{m}{2}} (x-m)^{m-3} (x-2m+6)^{\frac{m}{2}} (x-2) \left(1 - \frac{m}{x-2m+6} - \frac{m-2}{x-2} \right) \\ &= (x - (2m-4))^{\frac{m}{2}} (x-m)^{m-3} (x - (2m-6))^{\frac{m}{2}-1} (x^2 - (4m-6)x + 2m^2 - 4m). \end{aligned}$$

Therefore

$\text{Q-spec}(\Gamma_{D_{2m}})$

$$= \left\{ (2m-4)^{\frac{m}{2}}, (m)^{m-3}, (2m-6)^{\frac{m}{2}-1}, \left(2m-3+\sqrt{2m^2-8m+9}\right)^1, \left(2m-3-\sqrt{2m^2-8m+9}\right)^1 \right\}.$$

Number of edges in $\Gamma_{D_{2m}}^c$ is $\frac{m^2-4m+6}{2}$ and so $|e(\Gamma_{D_{2m}})| = \frac{(2m-2)(2m-2-1)}{2} - \frac{m^2-4m+6}{2} = \frac{3m(m-2)}{2}$. Now

$$\left| 2m-4 - \frac{2|e(\Gamma_{D_{2m}})|}{|v(\Gamma_{D_{2m}})|} \right| = \left| \frac{(m-2)(m-4)}{2m-2} \right| = \frac{(m-2)(m-4)}{2m-2},$$

$$\left| m - \frac{2|e(\Gamma_{D_{2m}})|}{|v(\Gamma_{D_{2m}})|} \right| = \left| \frac{(-m^2+4m)}{2m-2} \right| = \frac{(m^2-4m)}{2m-2},$$

$$\left| 2m-6 - \frac{2|e(\Gamma_{D_{2m}})|}{|v(\Gamma_{D_{2m}})|} \right| = \left| \frac{(m^2-10m+12)}{2m-2} \right| = \begin{cases} \frac{(-m^2+10m-12)}{2m-2}, & \text{if } m \leq 8 \\ \frac{(m^2-10m+12)}{2m-2}, & \text{if } m \geq 10, \end{cases}$$

$$\begin{aligned} \left| 2m-3+\sqrt{2m^2-8m+9} - \frac{2|e(\Gamma_{D_{2m}})|}{|v(\Gamma_{D_{2m}})|} \right| &= \left| \sqrt{2m^2-8m+9} + \frac{m}{2} - \frac{3}{2} + \frac{3}{2m-2} \right| \\ &= \sqrt{2m^2-8m+9} + \frac{m}{2} - \frac{3}{2} + \frac{3}{2m-2} \end{aligned}$$

and

$$\begin{aligned} \left| 2m-3-\sqrt{2m^2-8m+9} - \frac{2|e(\Gamma_{D_{2m}})|}{|v(\Gamma_{D_{2m}})|} \right| &= \left| -\sqrt{2m^2-8m+9} + \frac{m}{2} - \frac{3}{2} + \frac{3}{2m-2} \right| \\ &= \sqrt{2m^2-8m+9} - \frac{m}{2} + \frac{3}{2} - \frac{3}{2m-2}. \end{aligned}$$

Therefore, for $4 \leq m \leq 8$, we have

$$\begin{aligned} LE^+(\Gamma_{D_{2m}}) &= \frac{m}{2} \times \frac{(m-2)(m-4)}{2m-2} + (m-3) \times \frac{(m^2-4m)}{2m-2} + \left(\frac{m}{2}-1\right) \times \frac{-(m^2-10m+12)}{2m-2} \\ &\quad + \sqrt{2m^2-8m+9} + \frac{m}{2} - \frac{3}{2} + \frac{3}{2m-2} + \sqrt{2m^2-8m+9} - \frac{m}{2} + \frac{3}{2} - \frac{3}{2m-2} \end{aligned}$$

and for $m \geq 10$, we have

$$\begin{aligned} LE^+(\Gamma_{D_{2m}}) &= \frac{m}{2} \times \frac{(m-2)(m-4)}{2m-2} + (m-3) \times \frac{(m^2-4m)}{2m-2} + \left(\frac{m}{2}-1\right) \times \frac{(m^2-10m+12)}{2m-2} \\ &\quad + \sqrt{2m^2-8m+9} + \frac{m}{2} - \frac{3}{2} + \frac{3}{2m-2} + \sqrt{2m^2-8m+9} - \frac{m}{2} + \frac{3}{2} - \frac{3}{2m-2}. \end{aligned}$$

The required expressions for $LE^+(\Gamma_{D_{2m}})$ can be obtained on simplification. \square

Theorem 2.4. *If G is isomorphic to D_{2m} then*

- (a) $E(\Gamma_{D_{2m}}) \leq LE^+(\Gamma_{D_{2m}}) \leq LE(\Gamma_{D_{2m}})$, equality holds if and only if $G \cong D_8$.
- (b) $\Gamma_{D_{2m}}$ is non-hypoenergetic as well as non-hyperenergetic.
- (c) Γ_{D_6} is L -hyperenergetic but not Q -hyperenergetic. Γ_{D_8} is not L -hyperenergetic and not Q -hyperenergetic. If $m \neq 3, 4$ then $\Gamma_{D_{2m}}$ is Q -hyperenergetic and L -hyperenergetic.

Proof. (a) **Case 1:** m is odd

For $m = 3$, using Theorems 2.1 and 2.2, we have $E(\Gamma_{D_6}) = 2 + 2\sqrt{7}$, $LE(\Gamma_{D_6}) = \frac{42}{5}$ and $LE^+(\Gamma_{D_6}) = \frac{9}{5} + \sqrt{33}$. Clearly, $E(\Gamma_{D_6}) < LE^+(\Gamma_{D_6}) < LE(\Gamma_{D_6})$.

For $m \geq 5$, using Theorems 2.1 and 2.2, we have

$$LE(\Gamma_{D_{2m}}) - LE^+(\Gamma_{D_{2m}}) = \frac{8m^2 - 10m + 3}{2m - 1} - \sqrt{8m^2 - 16m + 9} \quad (2.1)$$

and

$$LE^+(\Gamma_{D_{2m}}) - E(\Gamma_{D_{2m}}) = \frac{2m^2(m - 6) + 15m - 4}{2m - 1} + \sqrt{8m^2 - 16m + 9} - \sqrt{5m^2 - 6m + 1}. \quad (2.2)$$

Since $8m^2 - 10m + 3 > 0$, $(2m - 1)\sqrt{8m^2 - 16m + 9} > 0$ and $(8m^2 - 10m + 3)^2 - (\sqrt{8m^2 - 16m + 9})^2 (2m - 1)^2 = 32m^3(m - 2) + 8m(5m - 1) > 0$ we have $8m^2 - 10m + 3 - (2m - 1)\sqrt{8m^2 - 16m + 9} > 0$. Therefore, by (2.1), $(2m - 1)(LE(\Gamma_{D_{2m}}) - LE^+(\Gamma_{D_{2m}})) > 0$. Hence, $LE(\Gamma_{D_{2m}}) > LE^+(\Gamma_{D_{2m}})$.

Again, $\sqrt{8m^2 - 16m + 9} > 0$, $\sqrt{5m^2 - 6m + 1} > 0$ and $(\sqrt{8m^2 - 16m + 9})^2 - (\sqrt{5m^2 - 6m + 1})^2 = m(3m - 10) + 8 > 0$. Therefore, $\sqrt{8m^2 - 16m + 9} - \sqrt{5m^2 - 6m + 1} > 0$. Since $2m^2(m - 6) + 15m - 4 > 0$ we have $\frac{2m^2(m - 6) + 15m - 4}{2m - 1} + \sqrt{8m^2 - 16m + 9} - \sqrt{5m^2 - 6m + 1} > 0$. Therefore, by (2.2), $LE^+(\Gamma_{D_{2m}}) > E(\Gamma_{D_{2m}})$. Hence, $E(\Gamma_{D_{2m}}) < LE^+(\Gamma_{D_{2m}}) < LE(\Gamma_{D_{2m}})$.

Case 2: m is even

For $4 \leq m \leq 8$, using Theorems 2.1 and 2.3, we have

$$LE(\Gamma_{D_{2m}}) - LE^+(\Gamma_{D_{2m}}) = \frac{m^3 - 4m^2 + 12m - 12}{2m - 2} - 2\sqrt{2m^2 - 8m + 9} \quad (2.3)$$

and

$$LE^+(\Gamma_{D_{2m}}) - E(\Gamma_{D_{2m}}) = \frac{(m - 4)(m^2 - 2m - 2)}{m - 1} + 2\sqrt{2m^2 - 8m + 9} - \sqrt{5m^2 - 12m + 4}. \quad (2.4)$$

Since $m^3 - 4m^2 + 12m - 12 > 0$, $2(2m - 2)\sqrt{2m^2 - 8m + 9} > 0$ and

$$(m^3 - 4m^2 + 12m - 12)^2 - (2\sqrt{2m^2 - 8m + 9})^2 (2m - 2)^2 = m(m - 4)^2(m - 2)(m^2 + 2m - 4) \geq 0$$

(equality holds if and only if $m = 4$). It follows that $m^3 - 4m^2 + 12m - 12 - 2(2m - 2)\sqrt{2m^2 - 8m + 9} \geq 0$. Therefore, by (2.3), $(2m - 2)(LE(\Gamma_{D_{2m}}) - LE^+(\Gamma_{D_{2m}})) \geq 0$. Hence, $LE(\Gamma_{D_{2m}}) \geq LE^+(\Gamma_{D_{2m}})$ equality holds if and only if $G \cong D_8$.

Again, $2\sqrt{2m^2 - 8m + 9} > 0$, $\sqrt{5m^2 - 12m + 4} > 0$ and $(2\sqrt{2m^2 - 8m + 9})^2 - (\sqrt{5m^2 - 12m + 4})^2 = (m - 4)(3m - 8) \geq 0$. Therefore, $2\sqrt{2m^2 - 8m + 9} - \sqrt{5m^2 - 12m + 4} \geq 0$ (equality holds if and only if $m = 4$). Since $(m - 4)(m^2 - 2m - 2) \geq 0$ we have $\frac{(m - 4)(m^2 - 2m - 2)}{m - 1} + 2\sqrt{2m^2 - 8m + 9} - \sqrt{5m^2 - 12m + 4} \geq 0$ (equality holds if and only if $m = 4$). Therefore, by (2.4), $LE^+(\Gamma_{D_{2m}}) \geq E(\Gamma_{D_{2m}})$. Hence, $E(\Gamma_{D_{2m}}) \leq LE^+(\Gamma_{D_{2m}}) \leq LE(\Gamma_{D_{2m}})$ equality holds if and only if $G \cong D_8$.

For $m \geq 10$, using Theorems 2.1 and 2.3, we have

$$LE(\Gamma_{D_{2m}}) - LE^+(\Gamma_{D_{2m}}) = \frac{4m^2 - 10m + 6}{m - 1} - 2\sqrt{2m^2 - 8m + 9} \quad (2.5)$$

and

$$LE^+(\Gamma_{D_{2m}}) - E(\Gamma_{D_{2m}}) = \frac{m^3 - 9m^2 + 19m - 8}{m - 1} + 2\sqrt{2m^2 - 8m + 9} - \sqrt{5m^2 - 12m + 4}. \quad (2.6)$$

Since $4m^2 - 10m + 6 > 0$, $2(m - 1)\sqrt{2m^2 - 8m + 9} > 0$ and $(4m^2 - 10m + 6)^2 - (2\sqrt{2m^2 - 8m + 9})^2 (m - 1)^2 = 8m^3(m - 4) + 8m(5m - 2) > 0$ we have $4m^2 - 10m + 6 - 2(m - 1)\sqrt{2m^2 - 8m + 9} > 0$. Therefore, by (2.5), $(m - 1)(LE(\Gamma_{D_{2m}}) - LE^+(\Gamma_{D_{2m}})) > 0$. Hence, $LE(\Gamma_{D_{2m}}) > LE^+(\Gamma_{D_{2m}})$.

Again, $2\sqrt{2m^2 - 8m + 9} > 0$, $\sqrt{5m^2 - 12m + 4} > 0$ and $(2\sqrt{2m^2 - 8m + 9})^2 - (\sqrt{5m^2 - 12m + 4})^2 = m(3m - 10) + 8 > 0$. Therefore, $2\sqrt{2m^2 - 8m + 9} - \sqrt{5m^2 - 12m + 4} > 0$. Since $m^3 - 9m^2 + 19m - 8 > 0$ we

have $\frac{m^3-9m^2+19m-8}{m-1} + 2\sqrt{2m^2-8m+9} - \sqrt{5m^2-12m+4} > 0$. Therefore, by (2.6), $LE^+(\Gamma_{D_{2m}}) > E(\Gamma_{D_{2m}})$. Hence, $E(\Gamma_{D_{2m}}) < LE^+(\Gamma_{D_{2m}}) < LE(\Gamma_{D_{2m}})$.

(b) **Case 1:** m is odd

Here, $|v(\Gamma_{D_{2m}})| = 2m - 1$ and $E(K_{|v(\Gamma_{D_{2m}})|}) = LE(K_{|v(\Gamma_{D_{2m}})|}) = LE^+(K_{|v(\Gamma_{D_{2m}})|}) = 4m - 4$. Using Theorem 2.1, we have

$$E(\Gamma_{D_{2m}}) - |v(\Gamma_{D_{2m}})| = \sqrt{(m-1)(5m-1)} - m \quad (2.7)$$

and

$$E(K_{|v(\Gamma_{D_{2m}})|}) - E(\Gamma_{D_{2m}}) = 3(m-1) - \sqrt{(m-1)(5m-1)}. \quad (2.8)$$

Since $\sqrt{(m-1)(5m-1)} > 0$, $m > 0$ and $(\sqrt{(m-1)(5m-1)})^2 - m^2 = 4m^2 - 6m + 1 > 0$ we have $\sqrt{(m-1)(5m-1)} - m > 0$. Therefore, by (2.7), $E(\Gamma_{D_{2m}}) > |v(\Gamma_{D_{2m}})|$.

Again, $\sqrt{(m-1)(5m-1)} > 0$, $3(m-1) > 0$ and $(3(m-1))^2 - (\sqrt{(m-1)(5m-1)})^2 = 4(m^2 - 3m + 2) > 0$ and so $3(m-1) - \sqrt{(m-1)(5m-1)} > 0$. Therefore, by (2.8), $E(K_{|v(\Gamma_{D_{2m}})|}) > E(\Gamma_{D_{2m}})$.

Case 2: m is even

Here, $|v(\Gamma_{D_{2m}})| = 2m - 2$ and $E(K_{|v(\Gamma_{D_{2m}})|}) = LE(K_{|v(\Gamma_{D_{2m}})|}) = LE^+(K_{|v(\Gamma_{D_{2m}})|}) = 4m - 6$. Using Theorem 2.1, we have

$$E(\Gamma_{D_{2m}}) - |v(\Gamma_{D_{2m}})| = \sqrt{(m-2)(5m-2)} - m \quad (2.9)$$

and

$$E(K_{|v(\Gamma_{D_{2m}})|}) - E(\Gamma_{D_{2m}}) = 3(m-2) + 2 - \sqrt{(m-2)(5m-2)}. \quad (2.10)$$

Since $\sqrt{(m-2)(5m-2)} > 0$, $m > 0$ and $(\sqrt{(m-2)(5m-2)})^2 - m^2 = 4(m^2 - 3m + 1) > 0$ we have $\sqrt{(m-2)(5m-2)} - m > 0$. Therefore, by (2.9), $E(\Gamma_{D_{2m}}) > |v(\Gamma_{D_{2m}})|$.

Again, $\sqrt{(m-2)(5m-2)} > 0$, $3(m-2) + 2 > 0$ and

$$(3(m-2) + 2)^2 - (\sqrt{(m-2)(5m-2)})^2 = 4(m^2 - 3m + 3) > 0$$

and so $3(m-2) + 2 - \sqrt{(m-2)(5m-2)} > 0$. Therefore, by (2.10), $E(K_{|v(\Gamma_{D_{2m}})|}) > E(\Gamma_{D_{2m}})$.

(c) **Case 1:** m is odd

For $m = 3$, using Theorems 2.1 and 2.2, $LE(\Gamma(D_6)) = \frac{42}{5}$, $LE^+(\Gamma(D_6)) = \frac{9}{5} + \sqrt{33}$ and $LE^+(K_{|v(\Gamma(D_6))|}) = LE(K_{|v(\Gamma(D_6))|}) = 8$. Clearly,

$$LE^+(\Gamma(D_6)) < LE^+(K_{|v(\Gamma(D_6))|}) = LE(K_{|v(\Gamma(D_6))|}) < LE(\Gamma(D_6)).$$

For $m \geq 5$, using Theorem 2.2, we have

$$LE^+(\Gamma_{D_{2m}}) - LE^+(K_{|v(\Gamma_{D_{2m}})|}) = \frac{2m^2(m-9) + 24m - 7}{2m-1} + \sqrt{8m^2 - 16m + 9} > 0.$$

Therefore, $LE^+(\Gamma_{D_{2m}}) > LE^+(K_{|v(\Gamma_{D_{2m}})|})$ which implies $\Gamma_{D_{2m}}$ is Q-hyperenergetic and consequently part (a) implies $\Gamma_{D_{2m}}$ is L-hyperenergetic.

Case 2: m is even

For $m = 4$, using Theorem 2.1, we have $LE(\Gamma(D_8)) = 8$ and $LE(K_{|v(\Gamma(D_8))|}) = 10$. Clearly, $LE(\Gamma(D_8)) < LE(K_{|v(\Gamma(D_8))|})$. Therefore, Γ_{D_8} is not L-hyperenergetic and not Q-hyperenergetic.

Using Theorem 2.3, for $m = 6$ and 8, we have

$$LE^+(\Gamma_{D_{2m}}) - LE^+(K_{|v(\Gamma_{D_{2m}})|}) = \frac{m^2(m-12) + 20m}{2m-2} + 2\sqrt{2m^2 - 8m + 9} > 0$$

and for $m \geq 10$, $LE^+(\Gamma_{D_{2m}}) - LE^+(K_{|v(\Gamma_{D_{2m}})|}) = \frac{m^2(m-12) + 26m - 9}{m-1} + 2\sqrt{2m^2 - 8m + 9} > 0$. Therefore, $LE^+(\Gamma_{D_{2m}}) > LE^+(K_{|v(\Gamma_{D_{2m}})|})$ which implies $\Gamma_{D_{2m}}$ is Q-hyperenergetic and consequently part (a) implies $\Gamma_{D_{2m}}$ is L-hyperenergetic. \square

In Theorem 2.4, we compare $E(\Gamma_{D_{2m}})$, $LE(\Gamma_{D_{2m}})$ and $LE^+(\Gamma_{D_{2m}})$. However, in the following figures, we show how close are they.




Figure 1: Energies of $\Gamma_{D_{2m}}$, m is odd




Figure 2: Energies of $\Gamma_{D_{2m}}$, m is even

2.2 The Quasidihedral groups, QD_{2^n}

We consider $QD_{2^n} := \langle a, b : a^{2^{n-1}} = b^2 = 1, bab^{-1} = a^{2^{n-2}-1} \rangle$, the quasidihedral groups of order 2^n (where $n \geq 4$). Results regarding different energies of non-commuting graphs of QD_{2^n} are given below.

Theorem 2.5 ([14, Result 1.2.16 (d)] and [12, Proposition 3.2]). *Let G be isomorphic to QD_{2^n} . Then*

$$E(\Gamma_{QD_{2^n}}) = (2^{n-1} - 2) + 2\sqrt{(5 \times 2^{n-2} - 1)(2^{n-2} - 1)} \text{ and } LE(\Gamma_{QD_{2^n}}) = \frac{2^{3n-3} - 2^{2n} + 3 \times 2^n}{2^{n-1} - 1}.$$

Theorem 2.6. *Let G be isomorphic to QD_{2^n} . Then*

$$\text{Q-spec}(\Gamma_{QD_{2^n}}) = \left\{ (2^n - 4)^{2^{n-2}}, (2^n - 2^{n-1})^{2^{n-1}-3}, (2^n - 6)^{2^{n-2}-1}, \left(2^n - 3 + \sqrt{2^{2n-1} - 2^{n+2} + 9}\right)^1, \right. \\ \left. \left(2^n - 3 - \sqrt{2^{2n-1} - 2^{n+2} + 9}\right)^1 \right\}$$

and

$$LE^+(\Gamma_{QD_{2^n}}) = \begin{cases} \frac{134}{7} + 2\sqrt{73}, & \text{if } n = 4 \\ \frac{2^{3n-2} + 2^{n+4} - 2^{2n+2} - 12}{2^{n-2}} + 2\sqrt{2^{2n-1} - 2^{n+2} + 9}, & \text{if } n \geq 5. \end{cases}$$

Proof. If $G \cong QD_{2^n}$ then $|v(\Gamma_{QD_{2^n}})| = 2^n - 2$ and $\Gamma_{QD_{2^n}} = K_{2^{n-2}, 2, 1, (2^{n-1}-2)}$. Using Theorem 1.6(b), we have

$$Q_{\Gamma_{QD_{2^n}}}(x) = \prod_{i=1}^2 (x - (2^n - 2) + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - (2^n - 2) + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - (2^n - 2) + 2p_i} \right) \\ = (x - 2^n + 4)^{2^{n-2}} (x - 2^n + 2^{n-1})^{2^{n-1}-3} (x - 2^n + 6)^{2^{n-2}-1} (x - 2) \left(1 - \frac{2^{n-1}}{x - 2^n + 6} - \frac{2^{n-1} - 2}{x - 2} \right) \\ = (x - (2^n - 4))^{2^{n-2}} (x - (2^n - 2^{n-1}))^{2^{n-1}-3} (x - (2^n - 6))^{2^{n-2}-1} (x^2 - (2^{n+1} - 6)x + 2^{2n-1} - 2^{n+1}).$$

Thus,

$$\text{Q-spec}(\Gamma_{QD_{2^n}}) = \left\{ (2^n - 4)^{2^{n-2}}, (2^n - 2^{n-1})^{2^{n-1}-3}, (2^n - 6)^{2^{n-2}-1}, \left(2^n - 3 + \sqrt{2^{2n-1} - 2^{n+2} + 9}\right)^1, \right. \\ \left. \left(2^n - 3 - \sqrt{2^{2n-1} - 2^{n+2} + 9}\right)^1 \right\}.$$

Number of edges of $\Gamma_{QD_{2^n}}^c$ is $2^{2n-3} - 2^n + 3$. Thus, $|e(\Gamma_{QD_{2^n}})| = \frac{(2^n-2)(2^{2n-2}-1)}{2} - (2^{2n-3} - 2^n + 3) = 3(2^{2n-3} - 2^{n-1})$. Now

$$\left| 2^n - 4 - \frac{2|e(\Gamma_{QD_{2^n}})|}{|v(\Gamma_{QD_{2^n}})|} \right| = \left| \frac{8 + 2^{2n-2} - 3 \times 2^n}{2^n - 2} \right| = \frac{8 + 2^{2n-2} - 3 \times 2^n}{2^n - 2}, \\ \left| 2^n - 2^{n-1} - \frac{2|e(\Gamma_{QD_{2^n}})|}{|v(\Gamma_{QD_{2^n}})|} \right| = \left| \frac{2^{n+1} - 2^{2n-2}}{2^n - 2} \right| = \frac{2^{2n-2} - 2^{n+1}}{2^n - 2},$$

$$\left| 2^n - 6 - \frac{2|e(\Gamma_{QD_{2^n}})|}{|v(\Gamma_{QD_{2^n}})|} \right| = \left| \frac{12 + 2^{2n-2} - 5 \times 2^n}{2^n - 2} \right| = \begin{cases} \frac{2}{7}, & \text{if } n = 4 \\ \frac{12 + 2^{2n-2} - 5 \times 2^n}{2^n - 2}, & \text{if } n \geq 5, \end{cases}$$

$$\begin{aligned} \left| 2^n - 3 + \sqrt{2^{2n-1} - 2^{n+2} + 9} - \frac{2|e(\Gamma_{QD_{2^n}})|}{|v(\Gamma_{QD_{2^n}})|} \right| &= \left| \sqrt{2^{2n-1} - 2^{n+2} + 9} + \frac{2^{2n-2} - 2^{n+1} + 6}{2^n - 2} \right| \\ &= \sqrt{2^{2n-1} - 2^{n+2} + 9} + \frac{2^{2n-2} - 2^{n+1} + 6}{2^n - 2} \end{aligned}$$

and

$$\begin{aligned} \left| 2^n - 3 - \sqrt{2^{2n-1} - 2^{n+2} + 9} - \frac{2|e(\Gamma_{QD_{2^n}})|}{|v(\Gamma_{QD_{2^n}})|} \right| &= \left| -\sqrt{2^{2n-1} - 2^{n+2} + 9} + \frac{2^{2n-2} - 2^{n+1} + 6}{2^n - 2} \right| \\ &= \sqrt{2^{2n-1} - 2^{n+2} + 9} - \frac{2^{2n-2} - 2^{n+1} + 6}{2^n - 2}. \end{aligned}$$

Therefore, for $n = 4$ we have $LE^+(\Gamma_{QD_{2^n}}) = \frac{134}{7} + 2\sqrt{73}$. For $n \geq 5$ we have

$$\begin{aligned} LE^+(\Gamma_{QD_{2^n}}) &= (2^{n-2}) \times \frac{8 + 2^n(2^{n-2} - 3)}{2^n - 2} + (2^{n-1} - 3) \times \frac{2^{n+1}(2^{n-3} - 1)}{2^n - 2} + (2^{n-2} - 1) \times \frac{12 + 2^n(2^{n-2} - 5)}{2^n - 2} \\ &\quad + \sqrt{2^{2n-1} - 2^{n+2} + 9} + \frac{2^{2n-2} - 2^{n+1} + 6}{2^n - 2} + \sqrt{2^{2n-1} - 2^{n+2} + 9} - \frac{2^{2n-2} - 2^{n+1} + 6}{2^n - 2} \end{aligned}$$

and the result follows on simplification. \square

Theorem 2.7. *If G is isomorphic to QD_{2^n} then*

- (a) $E(\Gamma_{QD_{2^n}}) < LE^+(\Gamma_{QD_{2^n}}) < LE(\Gamma_{QD_{2^n}})$.
- (b) $\Gamma_{QD_{2^n}}$ is non-hypoenergetic as well as non-hyperenergetic.
- (c) $\Gamma_{QD_{2^n}}$ is Q -hyperenergetic and L -hyperenergetic.

Proof. (a) For $n = 4$, using Theorems 2.5 and 2.6, we have $E(\Gamma_{QD_{2^n}}) = 6 + 2\sqrt{57}$, $LE(\Gamma_{QD_{2^n}}) = \frac{304}{7}$ and $LE^+(\Gamma_{QD_{2^n}}) = \frac{134}{7} + 2\sqrt{73}$. Clearly, $E(\Gamma_{QD_{16}}) < LE^+(\Gamma_{QD_{16}}) < LE(\Gamma_{QD_{16}})$.

For $n \geq 5$, using Theorems 2.5 and 2.6, we have

$$LE(\Gamma_{QD_{2^n}}) - LE^+(\Gamma_{QD_{2^n}}) = \frac{12 + 2^{2n+1} - 5 \times 2^{n+1}}{2^n - 2} - 2\sqrt{2^{2n-1} - 2^{n+2} + 9} \quad (2.11)$$

and

$$LE^+(\Gamma_{QD_{2^n}}) - E(\Gamma_{QD_{2^n}}) = \frac{2^{2n-2}(2^n - 18) + 19 \times 2^n - 16}{2^n - 2} + 2\sqrt{2^{2n-1} - 2^{n+2} + 9} - 2\sqrt{5 \times 2^{2n-4} - 3 \times 2^{n-1} + 1}. \quad (2.12)$$

Since $12 + 2^{2n+1} - 5 \times 2^{n+1} > 0$, $2\sqrt{2^{2n-1} - 2^{n+2} + 9}(2^n - 2) > 0$ and

$$(12 + 2^{2n+1} - 5 \times 2^{n+1})^2 - (2\sqrt{2^{2n-1} - 2^{n+2} + 9})^2 (2^n - 2)^2 = 2^{3n+1}(2^n - 8) + 2^{n+3}(5 \times 2^n - 4) > 0$$

we have $12 + 2^{2n+1} - 5 \times 2^{n+1} - 2(2^n - 2)\sqrt{2^{2n-1} - 2^{n+2} + 9} > 0$. Therefore, by (2.11), $(2^n - 2)(LE(\Gamma_{QD_{2^n}}) - LE^+(\Gamma_{QD_{2^n}})) > 0$. Hence, $LE(\Gamma_{QD_{2^n}}) > LE^+(\Gamma_{QD_{2^n}})$.

Again, $\sqrt{2^{2n-1} - 2^{n+2} + 9} > 0$, $\sqrt{5 \times 2^{2n-4} - 3 \times 2^{n-1} + 1} > 0$ and

$$(\sqrt{2^{2n-1} - 2^{n+2} + 9})^2 - (\sqrt{5 \times 2^{2n-4} - 3 \times 2^{n-1} + 1})^2 = 2^{n-4}(3 \times 2^n - 40) + 8 > 0.$$

Therefore, we have $\sqrt{2^{2n-1} - 2^{n+2} + 9} - \sqrt{5 \times 2^{2n-4} - 3 \times 2^{n-1} + 1} > 0$. Since $2^{2n-2}(2^n - 18) + 19 \times 2^n - 16 > 0$ we have $\frac{2^{2n-2}(2^n - 18) + 19 \times 2^n - 16}{2^n - 2} + 2\sqrt{2^{2n-1} - 2^{n+2} + 9} - 2\sqrt{5 \times 2^{2n-4} - 3 \times 2^{n-1} + 1} > 0$. Therefore, by (2.12), $LE^+(\Gamma_{QD_{2^n}}) \geq E(\Gamma_{QD_{2^n}})$. Hence, $E(\Gamma_{QD_{2^n}}) < LE^+(\Gamma_{QD_{2^n}}) < LE(\Gamma_{QD_{2^n}})$.

(b) Here, $|v(\Gamma_{QD_{2^n}})| = 2^n - 2$ and $E(K_{|v(\Gamma_{QD_{2^n}})|}) = LE(K_{|v(\Gamma_{QD_{2^n}})|}) = LE^+(K_{|v(\Gamma_{QD_{2^n}})|}) = 2^{n+1} - 6$. Using Theorem 2.5, we have

$$E(\Gamma_{QD_{2^n}}) - |v(\Gamma_{QD_{2^n}})| = 2 \left(\sqrt{(5 \times 2^{n-2} - 1)(2^{n-2} - 1)} - (2^{n-1} - 2^{n-2}) \right) \quad (2.13)$$

and

$$E(K_{|v(\Gamma_{QD_{2n}})|}) - E(\Gamma_{QD_{2n}}) = 2 \left(3 \times 2^{n-2} - 2 - \sqrt{(5 \times 2^{n-2} - 1)(2^{n-2} - 1)} \right). \quad (2.14)$$

Since $\sqrt{(5 \times 2^{n-2} - 1)(2^{n-2} - 1)} > 0$, $2^{n-1} - 2^{n-2} > 0$ and $\left(\sqrt{(5 \times 2^{n-2} - 1)(2^{n-2} - 1)} \right)^2 - (2^{n-1} - 2^{n-2})^2 = 2^{n-2}(4 \times 2^{n-2} - 6) + 1 > 0$ we have $\sqrt{(5 \times 2^{n-2} - 1)(2^{n-2} - 1)} - (2^{n-1} - 2^{n-2}) > 0$. Therefore, by (2.13), $E(\Gamma_{QD_{2n}}) > |v(\Gamma_{QD_{2n}})|$.

Again, $\sqrt{(5 \times 2^{n-2} - 1)(2^{n-2} - 1)} > 0$, $3 \times 2^{n-2} - 2 > 0$ and $(3 \times 2^{n-2} - 2)^2 - \left(\sqrt{(5 \times 2^{n-2} - 1)(2^{n-2} - 1)} \right)^2 = (2^{n-2} - 3)(4 \times 2^{n-2} + 6) + 21 > 0$ and so $3 \times 2^{n-2} - 2 - \sqrt{(5 \times 2^{n-2} - 1)(2^{n-2} - 1)} > 0$. Therefore, by (2.14), $E(K_{|v(\Gamma_{QD_{2n}})|}) > E(\Gamma_{QD_{2n}})$.

(c) For $n = 4$, using Theorem 2.6, $LE^+(\Gamma_{QD_{16}}) - LE^+(K_{|v(\Gamma_{QD_{16}})|}) = 2\sqrt{73} - \frac{48}{7} > 0$. Therefore, $LE^+(\Gamma_{QD_{16}}) > LE^+(K_{|v(\Gamma_{QD_{16}})|})$ which implies $\Gamma_{QD_{16}}$ is Q-hyperenergetic and consequently part (a) implies $\Gamma_{QD_{16}}$ is L-hyperenergetic.

For $n \geq 5$, using Theorem 2.6,

$$LE^+(\Gamma_{QD_{2n}}) - LE^+(K_{|v(\Gamma_{QD_{2n}})|}) = \frac{2^{2n-2}(2^{2n}-24)+2(13 \times 2^n-12)}{2^{n-2}} + 2\sqrt{2^{2n-1}-2^{n+2}+9} > 0.$$

Therefore, $LE^+(\Gamma_{QD_{2n}}) > LE^+(K_{|v(\Gamma_{QD_{2n}})|})$ which implies $\Gamma_{QD_{2n}}$ is Q-hyperenergetic and consequently part (a) implies $\Gamma_{QD_{2n}}$ is L-hyperenergetic. \square

In Theorem 2.7, we compare $E(\Gamma_{QD_{2n}})$, $LE(\Gamma_{QD_{2n}})$ and $LE^+(\Gamma_{QD_{2n}})$. However, in the following figures, we show how close are they.




Figure 3: Energies of $\Gamma_{QD_{2n}}$




Figure 4: A close up view of Figure 3

2.3 The groups M_{2rs}

We consider $M_{2rs} := \langle a, b : a^r = b^{2s} = 1, bab^{-1} = a^{-1} \rangle$, the groups of order $2rs$ (where $r \geq 3$ and $s \geq 1$). Results regarding different energies of non-commuting graphs of M_{2rs} are given below.

Theorem 2.8 ([14, Corollary 4.1.6 and (4.3.d)]). *Let G be isomorphic to M_{2rs} .*

(a) *If m is odd then*

$$E(\Gamma_{M_{2rs}}) = s(r-1) + s\sqrt{(r-1)(5r-1)} \text{ and } LE(\Gamma_{M_{2rs}}) = \frac{s}{2r-1} (2r^3s - 6r^2s + 4rs + 4r^2 - 2r).$$

(b) *If m is even then*

$$E(\Gamma_{M_{2rs}}) = s(r-2) + s\sqrt{(r-2)(5r-2)} \text{ and } LE(\Gamma_{M_{2rs}}) = \frac{s}{r-1} (r^3s - 6r^2s + 8rs + 2r^2 - 2r).$$

Theorem 2.9. *Let G be isomorphic to M_{2rs} , where r is odd. Then*

$$\text{Q-spec}(\Gamma_{M_{2rs}}) = \left\{ (2s(r-1))^{r(s-1)}, (rs)^{(r-1)s-1}, ((2r-3)s)^{r-1}, \left(\frac{s(4r-3 + \sqrt{8r^2 - 16r + 9})}{2} \right)^1, \left(\frac{s(4r-3 + \sqrt{8r^2 - 16r + 9})}{2} \right)^1 \right\}$$

and

$$LE^+(\Gamma_{M_{2rs}}) = \begin{cases} \frac{3s(4s-1)}{5} + s\sqrt{33}, & \text{if } r = 3 \\ s \left(\frac{(2r(r-1)(r-2))s}{2r-1} - (2r-3) + \sqrt{8r^2 - 16r + 9} \right), & \text{if } r \geq 5. \end{cases}$$

Proof. If $G \cong M_{2rs}$, where r is odd, then $|v(\Gamma_{M_{2rs}})| = (2r-1)s$ and $\Gamma_{M_{2rs}} = K_{r,s,1,((r-1)s)}$. Using Theorem 1.6(b), we have

$$\begin{aligned} Q_{\Gamma_{M_{2rs}}}(x) &= \prod_{i=1}^2 (x - (2rs - s) + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - (2rs - s) + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - (2rs - s) + 2p_i} \right) \\ &= (x - 2s(r-1))^{r(s-1)} (x - rs)^{(r-1)s-1} (x - (2r-3)s)^r (x - s) \left(1 - \frac{rs}{x - (2r-3)s} - \frac{(r-1)s}{x-s} \right) \\ &= (x - 2s(r-1))^{r(s-1)} (x - rs)^{(r-1)s-1} (x - (2r-3)s)^{r-1} (x^2 - (4r-3)sx + (2r^2 - 2r)s^2). \end{aligned}$$

$$\text{Thus, } Q\text{-spec}(\Gamma_{M_{2rs}}) = \left\{ (2s(r-1))^{r(s-1)}, (rs)^{(r-1)s-1}, ((2r-3)s)^{r-1}, \left(\frac{s(4r-3+\sqrt{8r^2-16r+9})}{2} \right)^1, \left(\frac{s(4r-3-\sqrt{8r^2-16r+9})}{2} \right)^1 \right\}.$$

Number of edges of $\Gamma_{M_{2rs}}^c$ is $\frac{(r^2-r+1)s^2-(2r-1)s}{2}$. Therefore,

$$|e(\Gamma_{M_{2rs}})| = \frac{(2r-1)^2 s^2 - (2r-1)s}{2} - \frac{(r^2-r+1)s^2 - (2r-1)s}{2} = \frac{3r(r-1)s^2}{2}.$$

Now

$$\left| (2r-2)s - \frac{2|e(\Gamma_{M_{2rs}})|}{|v(\Gamma_{M_{2rs}})|} \right| = \left| \frac{(r-1)(r-2)s}{2r-1} \right| = \frac{(r-1)(r-2)s}{2r-1},$$

$$\left| rs - \frac{2|e(\Gamma_{M_{2rs}})|}{|v(\Gamma_{M_{2rs}})|} \right| = \left| \frac{-r(r-2)s}{2r-1} \right| = \frac{r(r-2)s}{2r-1},$$

$$\left| (2r-3)s - \frac{2|e(\Gamma_{M_{2rs}})|}{|v(\Gamma_{M_{2rs}})|} \right| = \left| \frac{(r^2-5r+3)s}{2r-1} \right| = \begin{cases} \frac{3s}{5}, & \text{if } r = 3 \\ \frac{(r^2-5r+3)s}{2r-1}, & \text{if } r \geq 5, \end{cases}$$

$$\begin{aligned} \left| \frac{s}{2} (4r-3 + \sqrt{8r^2-16r+9}) - \frac{2|e(\Gamma_{M_{2rs}})|}{|v(\Gamma_{M_{2rs}})|} \right| &= \left| \frac{s}{2} \left(\sqrt{8r^2-16r+9} + r - \frac{3}{2} + \frac{3}{4r-2} \right) \right| \\ &= \frac{s}{2} \left(\sqrt{8r^2-16r+9} + r - \frac{3}{2} + \frac{3}{4r-2} \right) \end{aligned}$$

and

$$\begin{aligned} \left| \frac{s}{2} (4r-3 - \sqrt{8r^2-16r+9}) - \frac{2|e(\Gamma_{M_{2rs}})|}{|v(\Gamma_{M_{2rs}})|} \right| &= \left| \frac{s}{2} \left(-\sqrt{8r^2-16r+9} + r - \frac{3}{2} + \frac{3}{4r-2} \right) \right| \\ &= \frac{s}{2} \left(\sqrt{8r^2-16r+9} - r + \frac{3}{2} - \frac{3}{4r-2} \right). \end{aligned}$$

Therefore, for $n = 3$, we have $LE^+(\Gamma_{M_{2rs}}) = \frac{3s(4s-1)}{5} + s\sqrt{33}$. For $r \geq 5$, we have

$$\begin{aligned} LE^+(\Gamma_{M_{2rs}}) &= r(s-1) \times \frac{(r-1)(r-2)s}{2r-1} + ((r-1)s-1) \times \frac{r(r-2)s}{2r-1} + (r-1) \times \frac{(r^2-5r+3)s}{2r-1} + \\ &\quad \frac{s}{2} \left(\sqrt{8r^2-16r+9} + r - \frac{3}{2} + \frac{3}{4r-2} \right) + \frac{s}{2} \left(\sqrt{8r^2-16r+9} - r + \frac{3}{2} - \frac{3}{4r-2} \right) \end{aligned}$$

and the result follows on simplification. \square

Theorem 2.10. *Let G be isomorphic to M_{2rs} , where r is even. Then*

$$\begin{aligned} Q\text{-spec}(\Gamma_{M_{2rs}}) &= \left\{ (2s(r-2))^{rs-\frac{r}{2}}, (rs)^{rs-2s-1}, (2s(r-3))^{\frac{r}{2}-1}, \left(4rs - 6s + 2s\sqrt{2r^2-8r+9} \right)^1, \right. \\ &\quad \left. \left(4rs - 6s - 2s\sqrt{2r^2-8r+9} \right)^1 \right\} \end{aligned}$$

and

$$LE^+(\Gamma_{M_{2rs}}) = \begin{cases} \frac{s(r^3s-6r^2s+8rs-\frac{r^3}{2}+4r^2-8r+6)}{r-1} + 2s\sqrt{2r^2-8r+9}, & \text{if } 4 \leq r \leq 8 \\ \frac{s(r^3s-6r^2s+8rs-2r^2+8r-6)}{r-1} + 2s\sqrt{2r^2-8r+9}, & \text{if } r \geq 10. \end{cases}$$

Proof. If $G \cong M_{2rs}$ and r is even then $|v(\Gamma_{M_{2rs}})| = 2s(r-1)$ and $\Gamma_{M_{2rs}} = K_{\frac{r}{2}, (2s), 1, ((\frac{r}{2}-1)2s)}$. Using Theorem 1.6(b), we have

$$\begin{aligned} Q_{\Gamma_{M_{2rs}}}(x) &= \prod_{i=1}^2 (x - 2(rs-s) + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - 2(rs-s) + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - 2(rs-s) + 2p_i} \right) \\ &= (x - 2s(r-2))^{rs-\frac{r}{2}} (x-rs)^{rs-2s-1} (x-2s(r-3))^{\frac{r}{2}} (x-2s) \left(1 - \frac{rs}{x-2s(r-3)} - \frac{rs-2s}{x-2s} \right) \\ &= (x - 2s(r-2))^{rs-\frac{r}{2}} (x-rs)^{rs-2s-1} (x-2s(r-3))^{\frac{r}{2}-1} (x^2 - (4r-6)sx + rs^2(2r-4)). \end{aligned}$$

Thus,

$$\text{Q-spec}(\Gamma_{M_{2rs}}) = \left\{ (2s(r-2))^{rs-\frac{r}{2}}, (rs)^{rs-2s-1}, (2s(r-3))^{\frac{r}{2}-1}, (4rs-6s+2s\sqrt{2r^2-8r+9})^1, (4rs-6s-2s\sqrt{2r^2-8r+9})^1 \right\}.$$

Number of edges of $\Gamma_{M_{2rs}}^c$ is $\frac{(r^2-2r+4)s^2-2(r-1)s}{2}$. Thus, $|e(\Gamma_{M_{2rs}})| = \frac{2(r-1)s(2(r-1)s-1)}{2} - \frac{(r^2-r+1)s^2-2(r-1)s}{2} = \frac{3r(r-2)s^2}{2}$. Now

$$\begin{aligned} \left| 2s(r-2) - \frac{2|e(\Gamma_{M_{2rs}})|}{|v(\Gamma_{M_{2rs}})|} \right| &= \left| \frac{(r-2)(r-4)s}{2r-2} \right| = \frac{(r-2)(r-4)s}{2r-2}, \\ \left| rs - \frac{2|e(\Gamma_{M_{2rs}})|}{|v(\Gamma_{M_{2rs}})|} \right| &= \left| \frac{-r(r-4)s}{2r-2} \right| = \frac{r(r-4)s}{2r-2}, \\ \left| (2r-3)s - \frac{2|e(\Gamma_{M_{2rs}})|}{|v(\Gamma_{M_{2rs}})|} \right| &= \left| \frac{(r^2-10r+12)s}{2r-2} \right| = \begin{cases} \frac{(-r^2+10r-12)s}{2r-2}, & \text{if } 4 \leq r \leq 8 \\ \frac{(r^2-10r+12)s}{2r-2}, & \text{if } r \geq 10, \end{cases} \end{aligned}$$

$$\begin{aligned} \left| 4rs-6s+2s\sqrt{2r^2-8r+9} - \frac{2|e(\Gamma_{M_{2rs}})|}{|v(\Gamma_{M_{2rs}})|} \right| &= \left| \frac{rs}{2} - \frac{3s}{2} + \frac{3s}{2r-2} + s\sqrt{2r^2-8r+9} \right| \\ &= \frac{rs}{2} - \frac{3s}{2} + \frac{3s}{2r-2} + s\sqrt{2r^2-8r+9} \end{aligned}$$

and

$$\begin{aligned} \left| 4rs-6s-2s\sqrt{2r^2-8r+9} - \frac{2|e(\Gamma_{M_{2rs}})|}{|v(\Gamma_{M_{2rs}})|} \right| &= \left| \frac{rs}{2} - \frac{3s}{2} + \frac{3s}{2r-2} - s\sqrt{2r^2-8r+9} \right| \\ &= -\frac{rs}{2} + \frac{3s}{2} - \frac{3s}{2r-2} + s\sqrt{2r^2-8r+9}. \end{aligned}$$

Therefore, for $4 \leq r \leq 8$, we have

$$\begin{aligned} LE^+(\Gamma_{M_{2rs}}) &= \left(rs - \frac{r}{2} \right) \times \frac{(r-2)(r-4)s}{2r-2} + (rs-2s-1) \times \frac{r(r-4)s}{2r-2} + \left(\frac{r}{2} - 1 \right) \times \frac{(-r^2+10r-12)s}{2r-2} \\ &\quad + \frac{rs}{2} - \frac{3s}{2} + \frac{3s}{2r-2} + s\sqrt{2r^2-8r+9} - \frac{rs}{2} + \frac{3s}{2} - \frac{3s}{2r-2} + s\sqrt{2r^2-8r+9} \end{aligned}$$

and for $r \geq 10$, we have

$$\begin{aligned} LE^+(\Gamma_{M_{2rs}}) &= \left(rs - \frac{r}{2} \right) \times \frac{(r-2)(r-4)s}{2r-2} + (rs-2s-1) \times \frac{r(r-4)s}{2r-2} + \left(\frac{r}{2} - 1 \right) \times \frac{(r^2-10r+12)s}{2r-2} \\ &\quad + \frac{rs}{2} - \frac{3s}{2} + \frac{3s}{2r-2} + s\sqrt{2r^2-8r+9} - \frac{rs}{2} + \frac{3s}{2} - \frac{3s}{2r-2} + s\sqrt{2r^2-8r+9}. \end{aligned}$$

Hence, the results follow on simplification. \square

Theorem 2.11. *If G is isomorphic to M_{2rs} then*

- (a) $E(\Gamma_{M_{2rs}}) \leq LE^+(\Gamma_{M_{2rs}}) \leq LE(\Gamma_{M_{2rs}})$, equality holds if and only if $G \cong M_{8s}$.
- (b) $\Gamma_{M_{2rs}}$ is non-hypoenergetic as well as non-hyperenergetic.
- (c) Γ_{M_6} is L -hyperenergetic but not Q -hyperenergetic. $\Gamma_{M_{8s}}$ is not L -hyperenergetic and not Q -hyperenergetic. If $2rs \neq 6$ and $8s$ then $\Gamma_{M_{2rs}}$ is Q -hyperenergetic and L -hyperenergetic.

Proof. (a) **Case 1:** r is odd

Using Theorems 2.8 and 2.9, for $r = 3$, $LE(\Gamma_{M_{2rs}}) - LE^+(\Gamma_{M_{2rs}}) = \frac{33s}{5} - s\sqrt{33} > 0$ and $LE^+(\Gamma_{M_{2rs}}) - E(\Gamma_{M_{2rs}}) = \frac{12s^2 - 13s}{5} + (\sqrt{33} - 2\sqrt{7})s > 0$. For $r \geq 5$, using Theorems 2.8 and 2.9, we have

$$LE(\Gamma_{M_{2rs}}) - LE^+(\Gamma_{M_{2rs}}) = \frac{s(8r^2 - 10r + 3)}{2r - 1} - s\sqrt{8r^2 - 16r + 9} \quad (2.15)$$

and

$$LE^+(\Gamma_{M_{2rs}}) - E(\Gamma_{M_{2rs}}) = s \left(\frac{(2r^3 - 6r^2 + 4r)s - 6r^2 + 11r - 4}{2r - 1} + \sqrt{8r^2 - 16r + 9} - \sqrt{5r^2 - 6r + 1} \right). \quad (2.16)$$

Since $8r^2 - 10r + 3 > 0$, $(2r - 1)\sqrt{8r^2 - 16r + 9} > 0$ and $(8r^2 - 10r + 3)^2 - (2r - 1)^2(8r^2 - 16r + 9) = 32r^4 - 64r^3 + 40r^2 - 8r > 0$ we have $8r^2 - 10r + 3 - (2r - 1)\sqrt{8r^2 - 16r + 9} > 0$. Therefore, by (2.15), $(2r - 1)(LE(\Gamma_{M_{2rs}}) - LE^+(\Gamma_{M_{2rs}})) > 0$. Hence, $LE(\Gamma_{M_{2rs}}) > LE^+(\Gamma_{M_{2rs}})$.

Again, $\sqrt{8r^2 - 16r + 9} > 0$, $\sqrt{5r^2 - 6r + 1} > 0$ and $(\sqrt{8r^2 - 16r + 9})^2 - (\sqrt{5r^2 - 6r + 1})^2 = r(3r - 10) + 8 > 0$. Therefore, $\sqrt{8r^2 - 16r + 9} - \sqrt{5r^2 - 6r + 1} > 0$. Since $2r^3 - 6r^2 + 4r > 6r^2 - 11r + 4$ we have $\frac{(2r^3 - 6r^2 + 4r)s - 6r^2 + 11r - 4}{2r - 1} + \sqrt{8r^2 - 16r + 9} - \sqrt{5r^2 - 6r + 1} > 0$. Therefore, by (2.16), $LE^+(\Gamma_{M_{2rs}}) > E(\Gamma_{M_{2rs}})$. Hence, $E(\Gamma_{M_{2rs}}) < LE^+(\Gamma_{M_{2rs}}) < LE(\Gamma_{M_{2rs}})$.

Case 2: r is even

For $4 \leq r \leq 8$, using Theorems 2.8 and 2.10, we have

$$LE(\Gamma_{M_{2rs}}) - LE^+(\Gamma_{M_{2rs}}) = \frac{s}{r - 1} \left(\frac{r^3}{2} - 2r^2 + 6r - 6 \right) - 2s\sqrt{2r^2 - 8r + 9} \quad (2.17)$$

and

$$LE^+(\Gamma_{M_{2rs}}) - E(\Gamma_{M_{2rs}}) = \frac{(r^3 - 6r^2 + 8r)s^2 - \frac{r^3s}{2} + 3r^2s - 5rs + 4s}{r - 1} + 2s\sqrt{2r^2 - 8r + 9} - s\sqrt{5r^2 - 12r + 4}. \quad (2.18)$$

Since $\frac{r^3}{2} - 2r^2 + 6r - 6 > 0$, $2(r - 1)\sqrt{2r^2 - 8r + 9} > 0$ and $(\frac{r^3}{2} - 2r^2 + 6r - 6)^2 - 4(r - 1)^2(2r^2 - 8r + 9) = \frac{r^5(r - 8)}{4} + 2r^4 + 6r^2(3r - 8) + 32r \geq 0$ (equality holds if and only if $r = 4$) we have $\frac{r^3}{2} - 2r^2 + 6r - 6 - 2(r - 1)\sqrt{2r^2 - 8r + 9} \geq 0$. Therefore, by (2.17), $(r - 1)(LE(\Gamma_{M_{2rs}}) - LE^+(\Gamma_{M_{2rs}})) \geq 0$. Hence, $LE(\Gamma_{M_{2rs}}) \geq LE^+(\Gamma_{M_{2rs}})$ equality holds if and only if $G \cong M_{8s}$.

Again, $2\sqrt{2r^2 - 8r + 9} > 0$, $\sqrt{5r^2 - 12r + 4} > 0$ and $(2\sqrt{2r^2 - 8r + 9})^2 - (\sqrt{5r^2 - 12r + 4})^2 = (r - 4)(3r - 8) \geq 0$ (equality holds if and only if $r = 4$). Therefore, $2\sqrt{2r^2 - 8r + 9} - \sqrt{5r^2 - 12r + 4} \geq 0$. Since $r^3 - 6r^2 + 8r \geq \frac{r^3}{2} - 3r^2 + 5r - 4$ we have $\frac{(r^3 - 6r^2 + 8r)s^2 - \frac{r^3s}{2} + 3r^2s - 5rs + 4s}{r - 1} + 2s\sqrt{2r^2 - 8r + 9} - s\sqrt{5r^2 - 12r + 4} \geq 0$. Therefore, by (2.18), $LE^+(\Gamma_{M_{2rs}}) \geq E(\Gamma_{M_{2rs}})$. Hence, $E(\Gamma_{M_{2rs}}) \leq LE^+(\Gamma_{M_{2rs}}) \leq LE(\Gamma_{M_{2rs}})$ equality holds if and only if $G \cong M_{8s}$.

For $r \geq 10$, using Theorems 2.8 and 2.10, we have

$$LE(\Gamma_{M_{2rs}}) - LE^+(\Gamma_{M_{2rs}}) = 2s \left(\frac{2r^2 - 5r + 3}{r - 1} - \sqrt{2r^2 - 8r + 9} \right) \quad (2.19)$$

and

$$LE^+(\Gamma_{M_{2rs}}) - E(\Gamma_{M_{2rs}}) = s \left(\frac{(r^3 - 6r^2 + 8r)s - 3r^2 + 11r - 8}{r - 1} + 2\sqrt{2r^2 - 8r + 9} - \sqrt{5r^2 - 12r + 4} \right). \quad (2.20)$$

Since $2r^2 - 5r + 3 > 0$, $(r - 1)\sqrt{2r^2 - 8r + 9} > 0$ and $(2r^2 - 5r + 3)^2 - (r - 1)^2(2r^2 - 8r + 9) = 2r(r - 2)(r - 1)^2 > 0$ we have $2r^2 - 5r + 3 - (r - 1)\sqrt{2r^2 - 8r + 9} > 0$. Therefore, by (2.19), $(r - 1)(LE(\Gamma_{M_{2rs}}) - LE^+(\Gamma_{M_{2rs}})) > 0$. Hence, $LE(\Gamma_{M_{2rs}}) > LE^+(\Gamma_{M_{2rs}})$.

Again, $2\sqrt{2r^2 - 8r + 9} > 0$, $\sqrt{5r^2 - 12r + 4} > 0$ and $(2\sqrt{2r^2 - 8r + 9})^2 - (\sqrt{5r^2 - 12r + 4})^2 = (r - 4)(3r - 8) > 0$. Therefore, $2\sqrt{2r^2 - 8r + 9} - \sqrt{5r^2 - 12r + 4} > 0$. Since $r^3 - 6r^2 + 8r > 3r^2 - 11r + 8$ we have

$\frac{(r^3-6r^2+8r)s-3r^2+11r-8}{r-1} + 2\sqrt{2r^2-8r+9} - \sqrt{5r^2-12r+4} > 0$. Therefore, by (2.20), $LE^+(\Gamma_{M_{2rs}}) > E(\Gamma_{M_{2rs}})$. Hence, $E(\Gamma_{M_{2rs}}) < LE^+(\Gamma_{M_{2rs}}) < LE(\Gamma_{M_{2rs}})$.

(b) **Case 1:** r is odd

Here, $|v(\Gamma_{M_{2rs}})| = 2rs - s$ and $E(K_{|v(\Gamma_{M_{2rs}})|}) = LE(K_{|v(\Gamma_{M_{2rs}})|}) = LE^+(K_{|v(\Gamma_{M_{2rs}})|}) = 4rs - 2s - 2$. Using Theorem 2.8, we have

$$E(\Gamma_{M_{2rs}}) - |v(\Gamma_{M_{2rs}})| = s(\sqrt{(r-1)(5r-1)} - r) \quad (2.21)$$

and

$$E(K_{|v(\Gamma_{M_{2rs}})|}) - E(\Gamma_{M_{2rs}}) = 3rs - s - 2 - s\sqrt{(r-1)(5r-1)}. \quad (2.22)$$

Since $\sqrt{(r-1)(5r-1)} > 0$, $r > 0$ and $(\sqrt{(r-1)(5r-1)})^2 - (r)^2 = 2r(2r-3)+1 > 0$ we have $\sqrt{(r-1)(5r-1)} - r > 0$. Therefore, by (2.21), $E(\Gamma_{M_{2rs}}) > |v(\Gamma_{M_{2rs}})|$.

Again, $s\sqrt{(r-1)(5r-1)} > 0$, $3rs - s - 2 > 0$ and $(3rs - s - 2)^2 - (s\sqrt{(r-1)(5r-1)})^2 = 4rs(rs-3) + 4(s+1) > 0$ and so $3rs - s - 2 - s\sqrt{(r-1)(5r-1)} > 0$. Therefore, by (2.22), $E(K_{|v(\Gamma_{M_{2rs}})|}) > E(\Gamma_{M_{2rs}})$.

Case 2: r is even

Here, $|v(\Gamma_{M_{2rs}})| = 2rs - 2s$ and $E(K_{|v(\Gamma_{M_{2rs}})|}) = LE(K_{|v(\Gamma_{M_{2rs}})|}) = LE^+(K_{|v(\Gamma_{M_{2rs}})|}) = 4rs - 4s - 2$. Using Theorem 2.8, we have

$$E(\Gamma_{M_{2rs}}) - |v(\Gamma_{M_{2rs}})| = s(\sqrt{(r-2)(5r-2)} - r) \quad (2.23)$$

and

$$E(K_{|v(\Gamma_{M_{2rs}})|}) - E(\Gamma_{M_{2rs}}) = 3rs - 2s - 2 - s\sqrt{(r-2)(5r-2)}. \quad (2.24)$$

Since $\sqrt{(r-2)(5r-2)} > 0$, $r > 0$ and $(\sqrt{(r-2)(5r-2)})^2 - r^2 = 4(r(r-3)+1) > 0$ we have $\sqrt{(r-2)(5r-2)} - r > 0$. Therefore, by (2.23), $E(\Gamma_{M_{2rs}}) > |v(\Gamma_{M_{2rs}})|$.

Again, $s\sqrt{(r-2)(5r-2)} > 0$, $3rs - 2s - 2 > 0$ and $(3rs - 2s - 2)^2 - (s\sqrt{(r-2)(5r-2)})^2 = 4rs(rs-3) + 4(2s+1) > 0$ and so $3rs - 2s - 2 - s\sqrt{(r-2)(5r-2)} > 0$. Therefore, by (2.24), $E(K_{|v(\Gamma_{M_{2rs}})|}) > E(\Gamma_{M_{2rs}})$.

(c) **Case 1:** r is odd

For $r = 3$, using Theorem 2.9, we have $LE^+(\Gamma_{M_{2rs}}) - LE^+(K_{|v(\Gamma_{M_{2rs}})|}) = \frac{12s^2-53s}{5} + 2 + s\sqrt{33} > 0$ for all $s \neq 1$. Therefore, for $r = 3$ and $s \neq 1$, $LE^+(\Gamma_{M_{2rs}}) > LE^+(K_{|v(\Gamma_{M_{2rs}})|})$ which implies $\Gamma_{M_{2rs}}$ is Q-hyperenergetic and consequently part (a) implies $\Gamma_{M_{2rs}}$ is L-hyperenergetic. If $r = 3$ and $s = 1$, then $G \cong D_6$ so result follows from Theorem 2.4(c).

For $r \geq 5$, using Theorem 2.9, we have $LE^+(\Gamma_{M_{2rs}}) - LE^+(K_{|v(\Gamma_{M_{2rs}})|}) = \frac{(2r^3-6r^2+4r)s^2-(12r^2-16r+5)s}{2r-1} + s\sqrt{8r^2-16r+9} + 2 > 0$. Therefore, $LE^+(\Gamma_{M_{2rs}}) > LE^+(K_{|v(\Gamma_{M_{2rs}})|})$ which implies $\Gamma_{M_{2rs}}$ is Q-hyperenergetic and consequently part (a) implies $\Gamma_{M_{2rs}}$ is L-hyperenergetic.

Case 2: r is even

For $r = 4$ and $s \neq 1$, using Theorem 2.8, we have $LE(K_{|v(\Gamma_{M_{2rs}})|}) - LE(\Gamma_{M_{2rs}}) = \frac{12s}{7} - 2 > 0$. Therefore, $\Gamma_{M_{8s}}$ is not L-hyperenergetic and consequently part (a) implies that it is not Q-hyperenergetic. If $r = 4$ and $s = 1$, then $G \cong D_8$ so result follows from Theorem 2.4(c).

Using Theorem 2.10, for $4 < r \leq 8$, we get

$$LE^+(\Gamma_{M_{2rs}}) - LE^+(K_{|v(\Gamma_{M_{2rs}})|}) = \frac{(r^3-6r^2+8r)s^2-(\frac{r^3}{2}-2)s}{r-1} + 2s\sqrt{2r^2-8r+9} + 2 > 0.$$

Therefore, $LE^+(\Gamma_{M_{2rs}}) > LE^+(K_{|v(\Gamma_{M_{2rs}})|})$ which implies $\Gamma_{M_{2rs}}$ is Q-hyperenergetic and consequently part (a) implies $\Gamma_{M_{2rs}}$ is L-hyperenergetic.

Using Theorem 2.10, for $r \geq 10$, we get

$$LE^+(\Gamma_{M_{2rs}}) - LE^+(K_{|v(\Gamma_{M_{2rs}})|}) = \frac{(r^3-6r^2+8r)s^2-(6r^2-16r+10)s}{r-1} + 2s\sqrt{2r^2-8r+9} + 2 > 0.$$

Therefore, $LE^+(\Gamma_{M_{2rs}}) > LE^+(K_{|v(\Gamma_{M_{2rs}})|})$ which implies $\Gamma_{M_{2rs}}$ is Q-hyperenergetic and consequently part (a) implies $\Gamma_{M_{2rs}}$ is L-hyperenergetic. \square

In Theorem 2.11, we compare $E(\Gamma_{M_{2rs}})$, $LE(\Gamma_{M_{2rs}})$ and $LE^+(\Gamma_{M_{2rs}})$. However, in the following figures, we show how close are they.




Figure 5: Energies of $\Gamma_{M_{8r}}$




Figure 6: Energies of $\Gamma_{M_{10r}}$

2.4 The dicyclic groups, Q_{4n}

We consider $Q_{4n} := \langle x, y : x^{2n} = y^4 = 1; x^n = y^2; y^{-1}xy = x^{-1} \rangle$, the dicyclic groups of order $4n$ (where $n \geq 2$). Results regarding different energies of non-commuting graphs of Q_{4n} are given below.

Theorem 2.12 ([14, Corollary 4.1.8 and (4.3.f)]). *Let G be isomorphic to Q_{4n} . Then*

$$E(\Gamma_{Q_{4n}}) = 2 \left((n-1) + \sqrt{(n-1)(5n-1)} \right) \quad \text{and} \quad LE(\Gamma_{Q_{4n}}) = \frac{8n(n-1)(n-2) + 4n(2n-1)}{2n-1}.$$

Theorem 2.13. *Let G be isomorphic to Q_{4n} . Then*

$$\text{Q-spec}(\Gamma_{Q_{4n}}) = \left\{ (4n-4)^n, (2n)^{2n-3}, (4n-6)^{n-1}, (4n-3 + \sqrt{8n^2 - 16n + 9})^1, (4n-3 - \sqrt{8n^2 - 16n + 9})^1 \right\}$$

and

$$LE^+(\Gamma_{Q_{4n}}) = \begin{cases} \frac{4n^3 - 8n^2 + 6}{2n-1} + 2\sqrt{8n^2 - 16n + 9}, & \text{if } n \leq 4 \\ \frac{8n^3 - 32n^2 + 32n - 6}{2n-1} + 2\sqrt{8n^2 - 16n + 9}, & \text{if } n \geq 5. \end{cases}$$

Proof. If $G \cong Q_{4n}$ then $|v(\Gamma_{Q_{4n}})| = 4n - 2$ and $\Gamma_{Q_{4n}} = K_{n,2,1,(2n-2)}$. Using Theorem 1.6(b), we have

$$\begin{aligned} Q_{\Gamma_{Q_{4n}}}(x) &= \prod_{i=1}^2 (x - (4n-2) + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - (4n-2) + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - (4n-2) + 2p_i} \right) \\ &= (x - (4n-4))^n (x - 2n)^{2n-3} (x - 4n + 6)^n (x - 2) \left(1 - \frac{2n}{x - 4n + 6} - \frac{2n-2}{x-2} \right) \\ &= (x - (4n-4))^n (x - 2n)^{2n-3} (x - (4n-6))^{n-1} (x^2 - (8n-6)x + 8n^2 - 8n). \end{aligned}$$

Thus,

$$\text{Q-spec}(\Gamma_{Q_{4n}}) = \left\{ (4n-4)^n, (2n)^{2n-3}, (4n-6)^{n-1}, (4n-3 + \sqrt{8n^2 - 16n + 9})^1, (4n-3 - \sqrt{8n^2 - 16n + 9})^1 \right\}.$$

Number of edges of $\Gamma_{Q_{4n}}^c$ is $2n^2 - 4n + 3$. Thus, $|e(\Gamma_{Q_{4n}})| = \frac{(4n-2)(4n-2-1)}{2} - (2n^2 - 4n + 3) = 6n(n-1)$.

Now

$$\begin{aligned} \left| 4n-4 - \frac{2|e(\Gamma_{Q_{4n}})|}{|v(\Gamma_{Q_{4n}})|} \right| &= \left| \frac{(2n-4)(n-1)}{2n-1} \right| = \frac{(2n-4)(n-1)}{2n-1}, \\ \left| 2n - \frac{2|e(\Gamma_{Q_{4n}})|}{|v(\Gamma_{Q_{4n}})|} \right| &= \left| \frac{-2n(n-2)}{2n-1} \right| = \frac{2n(n-2)}{2n-1}, \\ \left| 4n-6 - \frac{2|e(\Gamma_{Q_{4n}})|}{|v(\Gamma_{Q_{4n}})|} \right| &= \left| \frac{2(n^2 - 5n + 3)}{2n-1} \right| = \begin{cases} \frac{-2(n^2 - 5n + 3)}{2n-1}, & \text{if } n \leq 4 \\ \frac{2(n^2 - 5n + 3)}{2n-1}, & \text{if } n \geq 5, \end{cases} \\ \left| 4n-3 + \sqrt{8n^2 - 16n + 9} - \frac{2|e(\Gamma_{Q_{4n}})|}{|v(\Gamma_{Q_{4n}})|} \right| &= \left| \sqrt{8n^2 - 16n + 9} + n - \frac{3}{2} + \frac{3}{4n-2} \right| \\ &= \sqrt{8n^2 - 16n + 9} + n - \frac{3}{2} + \frac{3}{4n-2} \end{aligned}$$

and

$$\begin{aligned} \left| 4n - 3 - \sqrt{8n^2 - 16n + 9} - \frac{2|e(\Gamma_{Q_{4n}})|}{|v(\Gamma_{Q_{4n}})|} \right| &= \left| -\sqrt{8n^2 - 16n + 9} + n - \frac{3}{2} + \frac{3}{4n-2} \right| \\ &= \sqrt{8n^2 - 16n + 9} - n + \frac{3}{2} - \frac{3}{4n-2}. \end{aligned}$$

Therefore, for $n \leq 4$ we have

$$\begin{aligned} LE^+(\Gamma_{Q_{4n}}) &= n \times \frac{(2n-4)(n-1)}{2n-1} + (2n-3) \times \frac{2n(n-2)}{2n-1} + (n-1) \times \frac{-2(n^2-5n+3)}{2n-1} + \\ &\quad \sqrt{8n^2 - 16n + 9} + n - \frac{3}{2} + \frac{3}{4n-2} + \sqrt{8n^2 - 16n + 9} - n + \frac{3}{2} - \frac{3}{4n-2} \end{aligned}$$

and for $n \geq 5$ we have

$$\begin{aligned} LE^+(\Gamma_{Q_{4n}}) &= n \times \frac{(2n-4)(n-1)}{2n-1} + (2n-3) \times \frac{2n(n-2)}{2n-1} + (n-1) \times \frac{2(n^2-5n+3)}{2n-1} + \\ &\quad \sqrt{8n^2 - 16n + 9} + n - \frac{3}{2} + \frac{3}{4n-2} + \sqrt{8n^2 - 16n + 9} - n + \frac{3}{2} - \frac{3}{4n-2} \end{aligned}$$

Hence, the results follow on simplification. \square

Theorem 2.14. *If G is isomorphic to Q_{4n} then*

- (a) $E(\Gamma_{Q_{4n}}) \leq LE^+(\Gamma_{Q_{4n}}) \leq LE(\Gamma_{Q_{4n}})$, equality holds if and only if $G \cong Q_8$.
- (b) $\Gamma_{Q_{4n}}$ is non-hypoenergetic as well as non-hyperenergetic.
- (c) Γ_{Q_8} is not L -hyperenergetic and not Q -hyperenergetic. If $n \neq 2$ then $\Gamma_{Q_{4n}}$ is Q -hyperenergetic and L -hyperenergetic.

Proof. (a) For $n \leq 4$, using Theorems 2.12 and 2.13, we have

$$LE(\Gamma_{Q_{4n}}) - LE^+(\Gamma_{Q_{4n}}) = 2 \left(\frac{2n^3 - 4n^2 + 6n - 3}{2n-1} - \sqrt{8n^2 - 16n + 9} \right) \quad (2.25)$$

and

$$LE^+(\Gamma_{Q_{4n}}) - E(\Gamma_{Q_{4n}}) = \frac{2(n-2)(2n^2-2n-1)}{2n-1} + 2\sqrt{8n^2-16n+9} - 2\sqrt{5n^2-6n+1}. \quad (2.26)$$

Since $2n^3 - 4n^2 + 6n - 3 > 0$, $(2n-1)\sqrt{8n^2-16n+9} > 0$ and $(2n^3 - 4n^2 + 6n - 3)^2 - (\sqrt{8n^2-16n+9})^2(2n-1)^2 = 8n^5(n-4) + 16n^4 + 24n^2(3n-4) + 32n \geq 0$ (equality holds if and only if $n=2$) we have $2n^3 - 4n^2 + 6n - 3 - (2n-1)\sqrt{8n^2-16n+9} \geq 0$. Therefore, by (2.25), $(2n-1)(LE(\Gamma_{Q_{4n}}) - LE^+(\Gamma_{Q_{4n}})) \geq 0$. Hence, $LE(\Gamma_{Q_{4n}}) \geq LE^+(\Gamma_{Q_{4n}})$ equality holds if and only if $G \cong Q_8$.

Again, $\sqrt{8n^2-16n+9} > 0$, $\sqrt{5n^2-6n+1} > 0$ and $(\sqrt{8n^2-16n+9})^2 - (\sqrt{5n^2-6n+1})^2 = n(3n-10) + 8 \geq 0$ (equality holds if and only if $n=2$). Therefore, $\sqrt{8n^2-16n+9} - \sqrt{5n^2-6n+1} \geq 0$. Since $(n-2)(2n^2-2n-1) \geq 0$ we have $\frac{2(n-2)(2n^2-2n-1)}{2n-1} + 2\sqrt{8n^2-16n+9} - 2\sqrt{5n^2-6n+1} \geq 0$ (equality holds if and only if $n=2$). Therefore, by (2.26), $LE^+(\Gamma_{Q_{4n}}) \geq E(\Gamma_{Q_{4n}})$. Hence, $E(\Gamma_{Q_{4n}}) \leq LE^+(\Gamma_{Q_{4n}}) \leq LE(\Gamma_{Q_{4n}})$ equality holds if and only if $G \cong Q_8$.

For $n \geq 5$, using Theorems 2.12 and 2.13, we have

$$LE(\Gamma_{Q_{4n}}) - LE^+(\Gamma_{Q_{4n}}) = 2 \left(\frac{8n^2 - 10n + 3}{2n-1} - \sqrt{8n^2 - 16n + 9} \right) \quad (2.27)$$

and

$$LE^+(\Gamma_{Q_{4n}}) - E(\Gamma_{Q_{4n}}) = \frac{2(2n^2(4n-9) + 19n - 4)}{2n-1} + 2\sqrt{8n^2-16n+9} - 2\sqrt{5n^2-6n+1}. \quad (2.28)$$

Since $8n^2 - 10n + 3 > 0$, $(2n-1)\sqrt{8n^2-16n+9} > 0$ and $(8n^2 - 10n + 3)^2 - (\sqrt{8n^2-16n+9})^2(2n-1)^2 = 32n^3(n-2) + 8n(5n-1) > 0$ we have $8n^2 - 10n + 3 - (2n-1)\sqrt{8n^2-16n+9} > 0$. Therefore, by (2.27), $(2n-1)(LE(\Gamma_{Q_{4n}}) - LE^+(\Gamma_{Q_{4n}})) > 0$. Hence, $LE(\Gamma_{Q_{4n}}) > LE^+(\Gamma_{Q_{4n}})$.

Again, $\sqrt{8n^2-16m+9} > 0$, $\sqrt{5n^2-6n+1} > 0$ and $(\sqrt{8n^2-16n+9})^2 - (\sqrt{5n^2-6n+1})^2 = n(3n-10) + 8 > 0$. Therefore, $\sqrt{8n^2-16n+9} - \sqrt{5n^2-6n+1} > 0$. Since $2n^2(4n-9) + 19n - 4 > 0$ we have

$\frac{2(2n^2(4n-9)+19n-4)}{2n-1} + 2\sqrt{8n^2 - 16n + 9} - 2\sqrt{5n^2 - 6n + 1} > 0$. Therefore, by (2.28), $LE^+(\Gamma_{Q_{4n}}) > E(\Gamma_{Q_{4n}})$. Hence, $E(\Gamma_{Q_{4n}}) < LE^+(\Gamma_{Q_{4n}}) < LE(\Gamma_{Q_{4n}})$.

(b) Here, $|v(\Gamma_{Q_{4n}})| = 4n - 2 = 2(2n - 1)$ and $E(K_{|v(\Gamma_{Q_{4n}})|}) = LE(K_{|v(\Gamma_{Q_{4n}})|}) = LE^+(K_{|v(\Gamma_{Q_{4n}})|}) = 8n - 6$. Using Theorem 2.12,

$$E(\Gamma_{Q_{4n}}) - |v(\Gamma_{Q_{4n}})| = 2(\sqrt{(n-1)(5n-1)} - n) \quad (2.29)$$

and

$$E(K_{|v(\Gamma_{Q_{4n}})|}) - E(\Gamma_{Q_{4n}}) = 2(3(n-1) + 1 - \sqrt{(n-1)(5n-1)}). \quad (2.30)$$

Since $\sqrt{(n-1)(5n-1)} > 0$, $n > 0$ and $(\sqrt{(n-1)(5n-1)})^2 - n^2 = 2n(2n-3)+1 > 0$ we have $\sqrt{(n-1)(5n-1)} - n > 0$. Therefore, by (2.29), $E(\Gamma_{Q_{4n}}) > |v(\Gamma_{Q_{4n}})|$.

Again, $\sqrt{(n-1)(5n-1)} > 0$, $3(n-1)+1 > 0$ and $(3(n-1)+1)^2 - (\sqrt{(n-1)(5n-1)})^2 = 2n(2n-3)+3 > 0$ and so $3(n-1) + 1 - \sqrt{(n-1)(5n-1)} > 0$. Therefore, by (2.30), $E(K_{|v(\Gamma_{Q_{4n}})|}) > E(\Gamma_{Q_{4n}})$.

(c) For $n = 2$, using Theorem 2.12, $LE(\Gamma_{Q_8}) = 8$ and $LE(K_{|v(\Gamma_{Q_8})|}) = 10$. Clearly, $LE(\Gamma_{Q_8}) < LE(K_{|v(\Gamma_{Q_8})|})$. Therefore, Γ_{Q_8} is not L-hyperenergetic and consequently part (a) implies that Γ_{Q_8} is not Q-hyperenergetic.

Using Theorem 2.13, for $2 < n \leq 4$,

$$LE^+(\Gamma_{Q_{4n}}) - LE^+(K_{|v(\Gamma_{Q_{4n}})|}) = \frac{4n(n-1)(n-5)}{2n-1} + 2\sqrt{8n^2 - 16n + 9} > 0.$$

Also, for $n \geq 5$, $LE^+(\Gamma_{Q_{4n}}) - LE^+(K_{|v(\Gamma_{Q_{4n}})|}) = \frac{8n^2(n-6)+52n-12}{2n-1} + 2\sqrt{8n^2 - 16n + 9} > 0$. Therefore, $LE^+(\Gamma_{Q_{4n}}) > LE^+(K_{|v(\Gamma_{Q_{4n}})|})$ which implies $\Gamma_{Q_{4n}}$ is Q-hyperenergetic and consequently part (a) implies that $\Gamma_{Q_{4n}}$ is L-hyperenergetic. Hence, the result holds. \square

2.5 The groups U_{6n}

We consider $U_{6n} := \langle x, y : x^{2n} = y^3 = 1; x^{-1}yx = y^{-1} \rangle$, the groups of order $6n$. Results regarding different energies of non-commuting graphs of U_{6n} are given below.

Theorem 2.15 ([14, Corollary 4.1.9 and putting $m = 3$ in (4.3.c)]). *Let G be isomorphic to U_{6n} . Then $E(\Gamma_{U_{6n}}) = 2n(1 + \sqrt{7})$ and $LE(\Gamma_{U_{6n}}) = \frac{12n^2+30n}{5}$.*

Theorem 2.16. *Let G be isomorphic to U_{6n} . Then*

$$\text{Q-spec}(\Gamma_{U_{6n}}) = \left\{ (3n)^{2n+1}, (4n)^{3n-3}, \left(\frac{(9 + \sqrt{33})n}{2} \right)^1, \left(\frac{(9 - \sqrt{33})n}{2} \right)^1 \right\}$$

and

$$LE^+(\Gamma_{U_{6n}}) = \frac{12n^2 - 3n}{5} + \sqrt{33}n.$$

Proof. If $G \cong U_{6n}$ then $|v(\Gamma_{U_{6n}})| = 5n$ and $\Gamma_{U_{6n}} = K_{1,2n,3,n}$. Using Theorem 1.6(b), we have

$$\begin{aligned} Q_{\Gamma_{U_{6n}}}(x) &= \prod_{i=1}^2 (x - 5n + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - 5n + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - 5n + 2p_i} \right) \\ &= (x - 3n)^{2n-1} (x - 4n)^{3n-3} (x - n)(x - 3n)^3 \left(1 - \frac{2n}{x - n} - \frac{3n}{x - 3n} \right) \\ &= (x - 3n)^{2n+1} (x - 4n)^{3n-3} (x^2 - 9nx + 12n^2). \end{aligned}$$

Thus, $\text{Q-spec}(\Gamma_{U_{6n}}) = \left\{ (3n)^{2n+1}, (4n)^{3n-3}, \left(\frac{(9+\sqrt{33})n}{2} \right)^1, \left(\frac{(9-\sqrt{33})n}{2} \right)^1 \right\}$.

Number of edges of $\Gamma_{U_{6n}}^c$ is $\frac{7n^2-5n}{2}$. Thus, $|e(\Gamma_{U_{6n}})| = \frac{5n(5n-1)}{2} - \frac{7n^2-5n}{2} = \frac{18n^2}{2}$. Now

$$\begin{aligned} \left| 3n - \frac{2|e(\Gamma_{U_{6n}})|}{|v(\Gamma_{U_{6n}})|} \right| &= \left| \frac{-3n}{5} \right| = \frac{3n}{5}, \quad \left| 4n - \frac{2|e(\Gamma_{U_{6n}})|}{|v(\Gamma_{U_{6n}})|} \right| = \left| \frac{2n}{5} \right| = \frac{2n}{5}, \\ \left| \frac{(9 + \sqrt{33})n}{5} - \frac{2|e(\Gamma_{U_{6n}})|}{|v(\Gamma_{U_{6n}})|} \right| &= \left| \frac{(9 + 5\sqrt{33})n}{10} \right| = \frac{(9 + 5\sqrt{33})n}{10} \end{aligned}$$

and

$$\left| \frac{(9 - \sqrt{33})n}{5} - \frac{2|e(\Gamma_{U_{6n}})|}{|v(\Gamma_{U_{6n}})|} \right| = \left| \frac{(9 - 5\sqrt{33})n}{10} \right| = \frac{(5\sqrt{33} - 9)n}{10}.$$

Therefore, $LE^+(\Gamma_{U_{6n}}) = (2n + 1) \times \frac{3n}{5} + (3n - 3) \times \frac{2n}{5} + \frac{(9+5\sqrt{33})n}{10} + \frac{(5\sqrt{33}-9)n}{10}$ and the result follows on simplification. \square

Theorem 2.17. *If G is isomorphic to U_{6n} then*

- (a) $E(\Gamma_{U_{6n}}) < LE^+(\Gamma_{U_{6n}}) < LE(\Gamma_{U_{6n}})$.
- (b) $\Gamma_{U_{6n}}$ is non-hypoenergetic as well as non-hyperenergetic.
- (c) $\Gamma_{U_{6n}}$ is Q -hyperenergetic and L -hyperenergetic.

Proof. (a) Using Theorems 2.15 and 2.16, we have

$$LE(\Gamma_{U_{6n}}) - LE^+(\Gamma_{U_{6n}}) = \frac{33n}{5} - \sqrt{33}n > 0 \quad (2.31)$$

and

$$LE^+(\Gamma_{U_{6n}}) - E(\Gamma_{U_{6n}}) = \frac{12n^2 - 13n}{5} + (\sqrt{33} - 2\sqrt{7})n > 0. \quad (2.32)$$

Thus, the conclusion is drawn from equations (2.31) and (2.32).

(b) Here, $|v(\Gamma_{U_{6n}})| = 5n$ and $E(K_{|v(\Gamma_{U_{6n}})|}) = LE(K_{|v(\Gamma_{U_{6n}})|}) = LE^+(K_{|v(\Gamma_{U_{6n}})|}) = 10n - 2$. Using Theorem 2.15, we have

$$E(\Gamma_{U_{6n}}) - |v(\Gamma_{U_{6n}})| = (2\sqrt{7} - 3)n > 0 \quad (2.33)$$

and

$$E(K_{|v(\Gamma_{U_{6n}})|}) - E(\Gamma_{U_{6n}}) = (8 - 2\sqrt{7})n - 2 > 0. \quad (2.34)$$

Thus, the conclusion is drawn from equations (2.33) and (2.34).

(c) Using Theorem 2.16, we have $LE^+(\Gamma_{U_{6n}}) - LE^+(K_{|v(\Gamma_{U_{6n}})|}) = \frac{12n^2 - 53n + 10}{5} + n\sqrt{33} > 0$. Therefore, $LE^+(\Gamma_{U_{6n}}) > LE^+(K_{|v(\Gamma_{U_{6n}})|})$ which implies $\Gamma_{U_{6n}}$ is Q -hyperenergetic and consequently part (a) implies $\Gamma_{U_{6n}}$ is L -hyperenergetic. \square

In Theorems 2.14 and 2.17, we compare $E(\Gamma_G)$, $LE(\Gamma_G)$ and $LE^+(\Gamma_G)$ if $G \cong Q_{4n}$ and U_{6n} respectively. However, in the following figures, we show how close are they for both the groups.




Figure 7: Energies of $\Gamma_{Q_{4n}}$




Figure 8: Energies of $\Gamma_{U_{6n}}$

It can be seen that if G is isomorphic to D_{2m} , QD_{2^n} , M_{2rs} , Q_{4n} or U_{6n} , then the central quotient of G is also isomorphic to some dihedral group. Therefore, we conclude this section with the following theorems for the non-commuting graphs of the groups G such that $\frac{G}{Z(G)} \cong D_{2m}$.

Theorem 2.18 ([14, Theorem 4.1.5]). *Let $\frac{G}{Z(G)}$ be isomorphic to D_{2m} ($m \geq 3$) and $|Z(G)| = n$. Then*

$$E(\Gamma_G) = n \left((m - 1) + \sqrt{(m - 1)(5m - 1)} \right) \quad \text{and} \quad LE(\Gamma_G) = \frac{n}{2m - 1} \left((2m^3 - 6m^2 + 4m)n + 4m^2 - 2m \right).$$

Theorem 2.19. Let $\frac{G}{Z(G)}$ be isomorphic to D_{2m} ($m \geq 3$) and $|Z(G)| = n$. Then

$$\text{Q-spec}(\Gamma_G) = \left\{ \left((2m-2)n^{m(n-1)}, (mn)^{(m-1)n-1}, ((2m-3)n)^{m-1}, \left(\frac{n(4m-3+\sqrt{8m^2-16m+9})}{2} \right)^1, \right. \right. \\ \left. \left. \left(\frac{n(4m-3-\sqrt{8m^2-16m+9})}{2} \right)^1 \right\}$$

and

$$LE^+(\Gamma_G) = \begin{cases} \frac{12n^2-3n}{5} + n\sqrt{33}, & \text{if } m = 3 \\ \frac{48n^2-29n}{7} + n\sqrt{73}, & \text{if } m = 4 \\ \frac{(2m^3-6m^2+4m)n^2-(4m^2-8m+3)n}{2m-1} + n\sqrt{8m^2-16m+9}, & \text{if } m \geq 5. \end{cases}$$

Proof. If $\frac{G}{Z(G)} \cong D_{2m}$ then $|v(\Gamma_G)| = (2m-1)n$ and $\Gamma_G = K_{m,n,1,((m-1)n)}$. Using Theorem 1.6(b), we have

$$Q_{\Gamma_G}(x) = \prod_{i=1}^2 (x - (2m-1)n + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - (2m-1)n + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - (2m-1)n + 2p_i} \right) \\ = (x - (2m-2)n)^{m(n-1)} (x - mn)^{(m-1)n-1} (x - (2m-3)n)^m (x - n) \left(1 - \frac{mn}{x - (2m-3)n} - \frac{(m-1)n}{x - n} \right) \\ = (x - (2m-2)n)^{m(n-1)} (x - mn)^{(m-1)n-1} (x - (2m-3)n)^{m-1} (x^2 - (4m-3)nx + (2m^2-2m)n^2).$$

Thus,

$$\text{Q-spec}(\Gamma_G) = \left\{ \left((2m-2)n^{m(n-1)}, (mn)^{(m-1)n-1}, ((2m-3)n)^{m-1}, \left(\frac{n(4m-3+\sqrt{8m^2-16m+9})}{2} \right)^1, \right. \right. \\ \left. \left. \left(\frac{n(4m-3-\sqrt{8m^2-16m+9})}{2} \right)^1 \right\}.$$

Number of edges of Γ_G^c is $\frac{(m^2-m+1)n^2-(2m-1)n}{2}$. Thus, $|e(\Gamma_G)| = \frac{(2m-1)^2n^2-(2m-1)n}{2} - \frac{(m^2-m+1)n^2-(2m-1)n}{2} = \frac{3m(m-1)n^2}{2}$. Now

$$\left| (2m-2)n - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{(m-1)(m-2)n}{2m-1} \right| = \frac{(m-1)(m-2)n}{2m-1},$$

$$\left| mn - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{-m(m-2)n}{2m-1} \right| = \frac{m(m-2)n}{2m-1},$$

$$\left| (2m-3)n - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{(m^2-5m+3)n}{2m-1} \right| = \begin{cases} \frac{(-m^2+5m-3)n}{2m-1}, & \text{if } m \leq 4 \\ \frac{(m^2-5m+3)n}{2m-1}, & \text{if } m \geq 5, \end{cases}$$

$$\left| \frac{n}{2} (4m-3+\sqrt{8m^2-16m+9}) - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{n}{2} \left(\sqrt{8m^2-16m+9} + m - \frac{3}{2} + \frac{3}{4m-2} \right) \right| \\ = \frac{n}{2} \left(\sqrt{8m^2-16m+9} + m - \frac{3}{2} + \frac{3}{4m-2} \right)$$

and

$$\left| \frac{n}{2} (4m-3-\sqrt{8m^2-16m+9}) - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{n}{2} \left(-\sqrt{8m^2-16m+9} + m - \frac{3}{2} + \frac{3}{4m-2} \right) \right| \\ = \frac{n}{2} \left(\sqrt{8m^2-16m+9} - m + \frac{3}{2} - \frac{3}{4m-2} \right).$$

Therefore, for $m \leq 4$ we have

$$LE^+(\Gamma_G) = m(n-1) \times \frac{(m-1)(m-2)n}{2m-1} + ((m-1)n-1) \times \frac{m(m-2)n}{2m-1} + (m-1) \times \frac{(-m^2+5m-3)n}{2m-1} + \frac{n}{2} \left(\sqrt{8m^2-16m+9} + m - \frac{3}{2} + \frac{3}{4m-2} \right) + \frac{n}{2} \left(\sqrt{8m^2-16m+9} - m + \frac{3}{2} - \frac{3}{4m-2} \right)$$

and for $m \geq 5$ we have

$$LE^+(\Gamma_G) = m(n-1) \times \frac{(m-1)(m-2)n}{2m-1} + ((m-1)n-1) \times \frac{m(m-2)n}{2m-1} + (m-1) \times \frac{(m^2-5m+3)n}{2m-1} + \frac{n}{2} \left(\sqrt{8m^2-16m+9} + m - \frac{3}{2} + \frac{3}{4m-2} \right) + \frac{n}{2} \left(\sqrt{8m^2-16m+9} - m + \frac{3}{2} - \frac{3}{4m-2} \right)$$

Hence, the results follow on simplification. \square

Theorem 2.20. *If $\frac{G}{Z(G)}$ is isomorphic to D_{2m} ($m \geq 3$) and $|Z(G)| = n$ then*

- (a) $E(\Gamma_G) < LE^+(\Gamma_G) < LE(\Gamma_G)$.
- (b) Γ_G is non-hypoenergetic as well as non-hyperenergetic.
- (c) Γ_G is L-hyperenergetic but not Q-hyperenergetic if $m = 3$ and $|Z(G)| = 1$. For $m = 3, 4$ and $|Z(G)| \neq 1$ or $m \geq 5$ and $|Z(G)| \geq 1$, Γ_G is Q-hyperenergetic and L-hyperenergetic.

Proof. (a) For $m = 3$, using Theorems 2.18 and 2.19, we have $LE(\Gamma_G) - LE^+(\Gamma_G) = \frac{33n}{5} - n\sqrt{33} > 0$ and $LE^+(\Gamma_G) - E(\Gamma_G) = \frac{12n^2-13n}{5} + (\sqrt{33} - 2\sqrt{7})n > 0$.

For $m = 4$, using Theorems 2.18 and 2.19, we have $LE(\Gamma_G) - LE^+(\Gamma_G) = \frac{85n}{7} - n\sqrt{73} > 0$ and $LE^+(\Gamma_G) - E(\Gamma_G) = \frac{48n^2-50n}{5} + (\sqrt{73} - \sqrt{57})n > 0$.

For $m \geq 5$, using Theorems 2.18 and 2.19, we have

$$LE(\Gamma_G) - LE^+(\Gamma_G) = \frac{n(8m^2 - 10m + 3)}{2m-1} - n\sqrt{8m^2 - 16m + 9} \quad (2.35)$$

and

$$LE^+(\Gamma_G) - E(\Gamma_G) = n \left(\frac{(2m^3 - 6m^2 + 4m)n - 6m^2 + 11m - 4}{2m-1} + \sqrt{8m^2 - 16m + 9} - \sqrt{5m^2 - 6m + 1} \right). \quad (2.36)$$

Since $8m^2 - 10m + 3 > 0$, $(2m-1)\sqrt{8m^2 - 16m + 9} > 0$ and $(8m^2 - 10m + 3)^2 - (2m-1)^2(8m^2 - 16m + 9) = 32m^4 - 64m^3 + 40m^2 - 8m > 0$ we have $8m^2 - 10m + 3 - (2m-1)\sqrt{8m^2 - 16m + 9} > 0$. Therefore, by (2.35), $(2m-1)(LE(\Gamma_G) - LE^+(\Gamma_G)) > 0$. Hence, $LE(\Gamma_G) > LE^+(\Gamma_G)$.

Again, $\sqrt{8m^2 - 16m + 9} > 0$, $\sqrt{5m^2 - 6m + 1} > 0$ and $(\sqrt{8m^2 - 16m + 9})^2 - (\sqrt{5m^2 - 6m + 1})^2 = m(3m - 10) + 8 > 0$. Therefore, $\sqrt{8m^2 - 16m + 9} - \sqrt{5m^2 - 6m + 1} > 0$. Since $2m^3 - 6m^2 + 4m > 6m^2 - 11m + 4$ we have $\frac{(2m^3 - 6m^2 + 4m)n - 6m^2 + 11m - 4}{2m-1} + \sqrt{8m^2 - 16m + 9} - \sqrt{5m^2 - 6m + 1} > 0$. Therefore, by (2.36), $LE^+(\Gamma_G) > E(\Gamma_G)$. Hence, $E(\Gamma_G) < LE^+(\Gamma_G) < LE(\Gamma_G)$.

(b) Here, $|v(\Gamma_G)| = 2mn - n$ and $E(K_{|v(\Gamma_G)|}) = LE(K_{|v(\Gamma_G)|}) = LE^+(K_{|v(\Gamma_G)|}) = 4mn - 2n - 2$. Using Theorem 2.18, we have

$$E(\Gamma_G) - |v(\Gamma_G)| = n(\sqrt{(m-1)(5m-1)} - m) \quad (2.37)$$

and

$$E(K_{|v(\Gamma_G)|}) - E(\Gamma_G) = 3mn - n - 2 - n\sqrt{(m-1)(5m-1)}. \quad (2.38)$$

Since $\sqrt{(m-1)(5m-1)} > 0$, $m > 0$ and $(\sqrt{(m-1)(5m-1)})^2 - m^2 = 2m(2m-3) + 1 > 0$ we have $\sqrt{(m-1)(5m-1)} - m > 0$. Therefore, by (2.37), $E(\Gamma_G) > |v(\Gamma_G)|$.

Again, $n\sqrt{(m-1)(5m-1)} > 0$, $3mn - n - 2 > 0$ and $(3mn - n - 2)^2 - (n\sqrt{(m-1)(5m-1)})^2 = 4mn(mn - 3) + 4(n+1) > 0$ and so $3mn - n - 2 - n\sqrt{(m-1)(5m-1)} > 0$. Therefore, by (2.38), $E(K_{|v(\Gamma_G)|}) > E(\Gamma_G)$.

(c) Using Theorem 2.19, for $m = 3$, we have $LE^+(\Gamma_G) - LE^+(K_{|v(\Gamma_G)|}) = \frac{12n^2-53n}{7} + 2 + n\sqrt{33} > 0$ for all $n \neq 1$. Therefore, for $m = 3$ and $n \neq 1$, $LE^+(\Gamma_G) > LE^+(K_{|v(\Gamma_G)|})$ which implies Γ_G is Q-hyperenergetic and consequently part (a) implies Γ_G is L-hyperenergetic. If $m = 3$ and $n = 1$, then $G \cong D_6$ so result follows from Theorem 2.4(c). Using Theorem 2.19, for $m = 4$, we have $LE^+(\Gamma_G) - LE^+(K_{|v(\Gamma_G)|}) = \frac{48n^2-127n}{7} + 2 + n\sqrt{73} > 0$

0 for all $n \neq 1$. Therefore, for $m = 4$ and $n \neq 1$, $LE^+(\Gamma_G) > LE^+(K_{|v(\Gamma_G)|})$ which implies Γ_G is Q-hyperenergetic and consequently part (a) implies Γ_G is L-hyperenergetic. The case $m = 4$ and $n = 1$ does not arise since $|Z(D_8)| = 2$.

For $r \geq 5$, using Theorem 2.19, we have $LE^+(\Gamma_G) - LE^+(K_{|v(\Gamma_G)|}) = \frac{(2m^3 - 6m^2 + 4m)n^2 - (12m^2 - 16m + 5)n}{2m - 1} + n\sqrt{8m^2 - 16m + 9} + 2 > 0$. Therefore, $LE^+(\Gamma_G) > LE^+(K_{|v(\Gamma_G)|})$ which implies Γ_G is Q-hyperenergetic and consequently part (a) implies Γ_G is L-hyperenergetic. \square

In Theorem 2.20, we compare $E(\Gamma_G)$, $LE(\Gamma_G)$ and $LE^+(\Gamma_G)$. However, in the following figures, we show how close are they.




Figure 9: Energies of Γ_G , $\frac{G}{Z(G)} \cong D_{2m}$, $|Z(G)| = 3$




Figure 10: Energies of Γ_G , $\frac{G}{Z(G)} \cong D_{2m}$, $|Z(G)| = 4$

3 $\frac{G}{Z(G)}$ is isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$

We compute the Signless Laplacian spectrum and Signless Laplacian energy of Γ_G considering the group G whose central quotient is isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$, where p is a prime. Further we compare energy, Laplacian energy and Signless Laplacian energy of Γ_G and look into the hyper- and hypo-properties of Γ_G .

Theorem 3.1 ([12, Theorem 2.2] and [14, Theorem 4.1.1 (c)]). *Let $\frac{G}{Z(G)}$ be isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$. Then $E(\Gamma_G) = LE(\Gamma_G) = 2p(p-1)|Z(G)|$. In particular, if G is non-abelian and $|G| = p^3$ then $E(\Gamma_G) = LE(\Gamma_G) = 2p^2(p-1)$.*

Theorem 3.2. *Let $\frac{G}{Z(G)}$ be isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$. Then*

$$\text{Q-spec}(\Gamma_G) = \left\{ (pn(p-1))^{(p^2-1)n-(p+1)}, (n(p-1)^2)^p, (2pn(p-1))^1 \right\}, \text{ where } |Z(G)| = n$$

and $LE^+(\Gamma_G) = 2p(p-1)|Z(G)|$. In particular, if G is non-abelian and $|G| = p^3$ then $LE^+(\Gamma_G) = 2p^2(p-1)$.

Proof. If $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$ then $|v(\Gamma_G)| = (p^2-1)n$ and $\Gamma_G = K_{(p+1).(p-1)n}$, where $|Z(G)| = n$. Using Theorem 1.6(b), we have

$$Q_{\Gamma_G}(x)$$

$$\begin{aligned} &= (x - (p^2-1)n + (p-1)n)^{(p+1)((p-1)n-1)} (x - (p^2-1)n + 2(p-1)n)^{p+1} \left(1 - \frac{(p^2-1)n}{x - (p^2-1)n + 2(p-1)n} \right) \\ &= (x - pn(p-1))^{(p^2-1)n-(p+1)} (x - n(p-1)^2)^p (x - 2pn(p-1)). \end{aligned}$$

Thus, $\text{Q-spec}(\Gamma_G) = \left\{ (pn(p-1))^{(p^2-1)n-(p+1)}, (n(p-1)^2)^p, (2pn(p-1))^1 \right\}$.

Number of edges of Γ_G^c is $\frac{n(p^2-1)(pn-n-1)}{2}$. Therefore,

$$|e(\Gamma_G)| = \frac{n^2(p^2-1)^2 - n(p^2-1)}{2} - \frac{n(p^2-1)(pn-n-1)}{2} = \frac{(p^2-p)(p^2-1)n^2}{2}.$$

Now

$$\left| pn(p-1) - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = |pn(p-1) - (p^2-p)n| = 0, \quad \left| n(p-1)^2 - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = |n - pn| = pn - n$$

and

$$\left| 2pn(p-1) - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = |p^2n - pn| = p^2n - pn.$$

Therefore, $LE^+(\Gamma_G) = ((p^2 - 1)n - (p + 1)) \times 0 + p \times (pn - n) + p^2n - pn = 2pn(p - 1)$. In particular, if G is non-abelian and $|G| = p^3$ then $n = p$. Therefore, $LE^+(\Gamma_G) = 2p^2(p - 1)$. \square

Theorem 3.3. *If $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$ then*

(a) $E(\Gamma_G) = LE(\Gamma_G) = LE^+(\Gamma_G)$.

(b) Γ_G is non-hypoenergetic, non-hyperenergetic, not L -hyperenergetic and not Q -hyperenergetic.

In particular, if G is non-abelian and $|G| = p^3$ then $E(\Gamma_G) = LE(\Gamma_G) = LE^+(\Gamma_G)$ and Γ_G is non-hyperenergetic, non-hypoenergetic, not L -hyperenergetic as well as not Q -hyperenergetic.

Proof. (a) For $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$, from Theorems 3.1 and 3.2, we have $E(\Gamma_G) = LE(\Gamma_G) = LE^+(\Gamma_G) = 2p^{2n}(p^n - 1)$ and hence follows.

(b) Here, $|v(\Gamma_G)| = (p^2 - 1)|Z(G)|$. Thus, $E(K_{|v(\Gamma_G)|}) = LE(K_{|v(\Gamma_G)|}) = LE^+(K_{|v(\Gamma_G)|}) = 2p(p - 1)|Z(G)| + 2(p - 1)|Z(G)| - 2$. Therefore, by Theorems 3.1 and 3.2, we have $E(K_{|v(\Gamma_G)|}) - E(\Gamma_G) = LE(K_{|v(\Gamma_G)|}) - LE(\Gamma_G) = LE^+(K_{|v(\Gamma_G)|}) - LE^+(\Gamma_G) = 2(p - 1)|Z(G)| - 2 > 0$. Also, $E(\Gamma_G) - |v(\Gamma_G)| = (p - 1)^2|Z(G)| > 0$. Hence, the results follow.

In particular, if G is non-abelian and $|G| = p^3$, then $|Z(G)| = p$ in the above cases so the results hold. \square

4 $\frac{G}{Z(G)}$ is isomorphic to $Sz(2)$

We compute spectrum, energy, Signless Laplacian Spectrum and Signless Laplacian energy of Γ_G considering the group G whose central quotient is isomorphic to the Suzuki group of order 20 denoted by $Sz(2)$. Further we compare energy, Laplacian energy and Signless Laplacian energy of Γ_G and look into the hyper- and hypo-properties of Γ_G .

Theorem 4.1 ([12, Theorem 2.1]). *Let $\frac{G}{Z(G)} \cong Sz(2)$. Then $LE(\Gamma_G) = (\frac{120}{19}n + 30)n$, where $|Z(G)| = n$.*

Theorem 4.2. *Let $\frac{G}{Z(G)} \cong Sz(2)$. Then*

(a) $\text{Spec}(\Gamma_G) = \left\{ 0^{19n-6}, (-3n)^4, (2n(3 + 2\sqrt{6}))^1, (2n(3 - 2\sqrt{6}))^1 \right\}$ and $E(\Gamma_G) = 4n(3 + 2\sqrt{6})$, where $n = |Z(G)|$.

(b) $Q\text{-spec}(\Gamma_G) = \left\{ (16n)^{15n-5}, (15n)^{4n-1}, (13n)^4, \left(\frac{n(43+\sqrt{409})}{2}\right)^1, \left(\frac{n(43-\sqrt{409})}{2}\right)^1 \right\}$ and $LE^+(\Gamma_G) = \frac{120n^2+177n}{19} + \sqrt{409}n$, where $n = |Z(G)|$.

Proof. If $\frac{G}{Z(G)} \cong Sz(2)$ and $|Z(G)| = n$ then $\Gamma_G = K_{5,3n,1,4n}$ and it is a complete 6-partite graph with $19n$ vertices.

(a) Using Theorem 1.6(a), the characteristic polynomial of Γ_G is

$$P_{\Gamma_G}(x) = x^{19n-6}(x + 3n)^4(x^2 - 12nx - 60n^2).$$

Therefore, $\text{Spec}(\Gamma_G) = \left\{ 0^{19n-6}, (-3n)^4, (2n(3 + 2\sqrt{6}))^1, (2n(3 - 2\sqrt{6}))^1 \right\}$ and $E(\Gamma_G) = 4n(3 + 2\sqrt{6})$.

(b) Using Theorem 1.6(b), we have

$$\begin{aligned} Q_{\Gamma_G}(x) &= \prod_{i=1}^2 (x - 19n + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - 19n + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - 19n + 2p_i} \right) \\ &= (x - 19n + 3n)^{5(3n-1)} (x - 19n + 4n)^{4n-1} (x - 19n + 2 \times 3n)^5 (x - 19n + 2 \times 4n)^1 \\ &\quad \times \left(1 - \frac{5 \times 3n}{x - 19n + 2 \times 3n} - \frac{4n}{x - 19n + 2 \times 4n} \right) \\ &= (x - 16n)^{15n-5} (x - 15n)^{4n-1} (x - 13n)^4 (x^2 - 43nx + 360n^2). \end{aligned}$$

Thus, $Q\text{-spec}(\Gamma_G) = \left\{ (16n)^{15n-5}, (15n)^{4n-1}, (13n)^4, \left(\frac{n(43+\sqrt{409})}{2}\right)^1, \left(\frac{n(43-\sqrt{409})}{2}\right)^1 \right\}$.

Number of edges of Γ_G^c is $\frac{61n^2-19n}{2}$. Thus, $|e(\Gamma_G)| = \frac{19n(19n-1)}{2} - \frac{61n^2-19n}{2} = 150n^2$. Now

$$\left| 16n - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{4n}{19} \right| = \frac{4n}{19}, \quad \left| 15n - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{-15n}{19} \right| = \frac{15n}{19},$$

$$\left| 13n - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{-53n}{19} \right| = \frac{53n}{19},$$

$$\left| \frac{(43 + \sqrt{409})n}{2} - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{(217 + 19\sqrt{409})n}{38} \right| = \frac{(217 + 19\sqrt{409})n}{38}$$

and

$$\left| \frac{(43 - \sqrt{409})n}{2} - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{(217 - 19\sqrt{409})n}{38} \right| = \frac{(19\sqrt{409} - 217)n}{38}.$$

Therefore, $LE^+(\Gamma_G) = (15n - 5) \times \frac{4n}{19} + (4n - 1) \times \frac{15n}{19} + 4 \times \frac{53n}{19} + \frac{(217+19\sqrt{409})n}{38} + \frac{(19\sqrt{409}-217)n}{38}$ and the result follows on simplification. \square

Theorem 4.3. *If $\frac{G}{Z(G)}$ is isomorphic to $Sz(2)$ then*

- (a) $E(\Gamma_G) < LE^+(\Gamma_G) < LE(\Gamma_G)$.
- (b) Γ_G is non-hypoenergetic as well as non-hyperenergetic.
- (c) $\Gamma_{Sz(2)}$ is not Q-hyperenergetic but is L-hyperenergetic. If $G \not\cong Sz(2)$ then Γ_G is Q-hyperenergetic and L-hyperenergetic.

Proof. (a) Using Theorems 4.2 and 4.1, we have $LE(\Gamma_G) - LE^+(\Gamma_G) = \left(\frac{393}{19} - \sqrt{409}\right)n > 0$ and $LE^+(\Gamma_G) - E(\Gamma_G) = \frac{3n(40n-17)}{19} + (\sqrt{409} - 8\sqrt{6})n > 0$, where $n = |Z(G)|$. Hence, the result follows.

(b) Here, $|v(\Gamma_G)| = 19n$, $n = |Z(G)|$ and $E(K_{|v(\Gamma_G)|}) = LE(K_{|v(\Gamma_G)|}) = LE^+(K_{|v(\Gamma_G)|}) = 38n - 2$. Using Theorem 4.2(a), we have $E(\Gamma_G) - |v(\Gamma_G)| = (8\sqrt{6} - 7)n > 0$ and also $E(K_{|v(\Gamma_G)|}) - E(\Gamma_G) = 2(13 - 4\sqrt{6})n - 2 > 0$. Hence, the result follows.

(c) For $n = |Z(G)| = 1$, from Proposition 4.3.13 of [14], we have $LE(\Gamma_{Sz(2)}) = \frac{690}{19} > 36 = LE(K_{|v(\Gamma_{Sz(2)})})$. Hence, for $n = 1$, Γ_G is L-hyperenergetic.

For $n = |Z(G)| > 1$, using Theorem 4.2(b), we have $LE^+(\Gamma_G) - LE^+(K_{|v(\Gamma_G)|}) = \frac{5n(24n-109)+38}{19} + n\sqrt{409} > 0$. Therefore, $LE^+(\Gamma_G) > LE^+(K_{|v(\Gamma_G)|})$ which implies Γ_G is Q-hyperenergetic and consequently part (a) implies Γ_G is L-hyperenergetic. \square

For $\frac{G}{Z(G)} \cong Sz(2)$, the following figures also demonstrate that among the three energies, $E(\Gamma_G)$ is the least and the fact that although $LE^+(\Gamma_G) < LE(\Gamma_G)$ but these two energies are very close to each other.




Figure 11: Energies of Γ_G where $\frac{G}{Z(G)} \cong Sz(2)$




Figure 12: A close up view of Figure 11

5 Some more classes of groups

In this section we discuss results on energy, Laplacian energy and Signless Laplacian energy of non-commuting graph of certain well-known classes of finite groups.

5.1 The Hanaki groups

We consider the Hanaki groups

$$A(n, \mathcal{V}) = \left\{ U(a, b) = \begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ b & \mathcal{V}(a) & 1 \end{bmatrix} : a, b \in GF(2^n) \right\} \quad (n \geq 2),$$

under matrix multiplication given by $U(a, b)U(a', b') = U(a + a', b + b' + a'\mathcal{V}(a))$ (here \mathcal{V} be the Frobenius automorphism of $GF(2^n)$, i.e., $\mathcal{V}(x) = x^2$ for all $x \in GF(2^n)$) and

$$A(n, p) = \left\{ V(a, b, c) = \begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ b & c & 1 \end{bmatrix} : a, b, c \in GF(p^n) \right\} \quad (p \text{ is any prime})$$

under matrix multiplication $V(a, b, c)V(a', b', c') = V(a + a', b + b' + ca', c + c')$. In this section, we compute Signless Laplacian spectrum and Signless Laplacian energy of non-commuting graph of the groups $A(n, \mathcal{V})$ and $A(n, p)$. Further we compare Signless Laplacian energy of Γ_G with its predetermined energy, Laplacian energy and look into the hyper- and hypo-properties of Γ_G if G is isomorphic to $A(n, \mathcal{V})$ and $A(n, p)$.

Theorem 5.1 ([12, Proposition 3.5] and [14, Theorem 4.1.3]). *If G denotes the Hanaki group $A(n, \mathcal{V})$, then*

$$E(\Gamma_G) = LE(\Gamma_G) = 2^{2n+1} - 2^{n+2}.$$

Theorem 5.2. *If G is isomorphic to the Hanaki group $A(n, \mathcal{V})$ then $Q\text{-spec}(\Gamma_G) = \{(2^{2n} - 2^{n+1})^{2^{2n} - 2^{n+1} + 1}, (2^{2n} - 3 \times 2^n)^{2^n - 2}, (2^{2n+1} - 2^{n+2})^1\}$ and $LE^+(\Gamma_G) = 2^{2n+1} - 2^{n+2}$.*

Proof. If G is isomorphic to the Hanaki group $A(n, \mathcal{V})$ then $|v(\Gamma_G)| = 2^{2n} - 2^n$ and $\Gamma_G = K_{(2^n-1), 2^n}$. Using Theorem 1.6(b), we have

$$\begin{aligned} Q_{\Gamma_G}(x) &= (x - (2^{2n} - 2^n) + 2^n)^{(2^n-1)(2^n-1)} (x - (2^{2n} - 2^n) + 2 \times 2^n)^{(2^n-1)} \left(1 - \frac{(2^n - 1) \cdot 2^n}{x - (2^{2n} - 2^n) + 2 \times 2^n} \right) \\ &= (x - (2^{2n} - 2^{n+1}))^{(2^n-1)^2} (x - (2^{2n} - 3 \times 2^n))^{2^n-2} (x - (2^{2n+1} - 2^{n+2})). \end{aligned}$$

Thus, $Q\text{-spec}(\Gamma_G) = \{(2^{2n} - 2^{n+1})^{2^{2n} - 2^{n+1} + 1}, (2^{2n} - 3 \times 2^n)^{2^n - 2}, (2^{2n+1} - 2^{n+2})^1\}$.

Number of edges of Γ_G^c is $2^{n-2}(2^{2n} - 2^{n+1} + 1)$. Thus, $|e(\Gamma_G)| = \frac{(2^{2n} - 2^n)(2^{2n} - 2^{n+1})}{2} - 2^{n-2}(2^{2n} - 2^{n+1} + 1) = 2^{4n-1} - 3 \times 2^{3n-1} + 2^{2n}$. Now,

$$\left| 2^{2n} - 2^{n+1} - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{2^{3n+1} - 2^{3n+1}}{2^{2n} - 2^n} \right| = 0, \quad \left| 2^{2n} - 3 \times 2^n - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{2^{2n} - 2^{3n}}{2^{2n} - 2^n} \right| = \frac{2^{3n} - 2^{2n}}{2^{2n} - 2^n}$$

and

$$\left| 2^{2n+1} - 2^{n+2} - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{2^{4n} - 3 \times 2^{3n} + 2^{2n+1}}{2^{2n} - 2^n} \right| = \frac{2^{4n} - 3 \times 2^{3n} + 2^{2n+1}}{2^{2n} - 2^n}.$$

Therefore, $LE^+(\Gamma_G) = (2^{2n} - 2^{n+1} + 1) \times 0 + (2^n - 2) \times \frac{2^{3n} - 2^{2n}}{2^{2n} - 2^n} + \frac{2^{4n} - 3 \times 2^{3n} + 2^{2n+1}}{2^{2n} - 2^n}$ and the result follows on simplification. \square

Theorem 5.3. *If G is isomorphic to the Hanaki group $A(n, \mathcal{V})$ then*

- (a) $E(\Gamma_G) = LE(\Gamma_G) = LE^+(\Gamma_G)$.
- (b) Γ_G is non-hypoenergetic, non-hyperenergetic, not L -hyperenergetic and not Q -hyperenergetic.

Proof. (a) Using Theorems 5.1 and 5.2, we have $E(\Gamma_G) = LE(\Gamma_G) = LE^+(\Gamma_G) = 2^{n+1}(2^n - 2)$ and hence the result follows.

(b) Here, $|v(\Gamma_G)| = 2^n(2^n - 1)$ and $E(K_{|v(\Gamma_G)|}) = LE(K_{|v(\Gamma_G)|}) = LE^+(K_{|v(\Gamma_G)|}) = 2^{n+1}(2^n - 1) - 2$. Therefore, by Theorems 5.1 and 5.2, we have $E(\Gamma_G) - |v(\Gamma_G)| = 2^n(2^n - 3) > 0$, for $n > 1$ and $E(K_{|v(\Gamma_G)|}) - E(\Gamma_G) = LE(K_{|v(\Gamma_G)|}) - LE(\Gamma_G) = LE^+(K_{|v(\Gamma_G)|}) - LE^+(\Gamma_G) = 2(2^n - 1) > 0$. Hence, the results follow. \square

Theorem 5.4 ([12, Proposition 3.6] and [14, Theorem 4.1.4]). *If G denotes the Hanaki group $A(n, p)$ then*

$$E(\Gamma_G) = LE(\Gamma_G) = 2(p^{3n} - p^{2n}).$$

Theorem 5.5. *If G is isomorphic to the Hanaki group $A(n, p)$ then $\text{Q-spec}(\Gamma_G) = \{(p^{3n} - p^{2n})^{(p^n+1)(p^{2n}-p^n-1)}, (p^{3n} - 2p^{2n} + p^n)^{p^n}, (2p^{3n} - 2p^{2n})^1\}$ and $LE^+(\Gamma_G) = 2p^{2n}(p^n - 1)$.*

Proof. If G is isomorphic to the Hanaki group $A(n, p)$ then $|v(\Gamma_G)| = p^{3n} - p^n$ and $\Gamma_G = K_{(p^n+1), (p^{2n}-p^n)}$. Using Theorem 1.6(b), we have

$$\begin{aligned} Q_{\Gamma_G}(x) &= (x - (p^{3n} - p^n) + (p^{2n} - p^n))^{(p^n+1)(p^{2n}-p^n-1)} (x - (p^{3n} - p^n) + 2(p^{2n} - p^n))^{(p^n+1)} \\ &\quad \times \left(1 - \frac{p^{3n} - p^n}{x - (p^{3n} - p^n) + 2(p^{2n} - p^n)}\right) \\ &= (x - (p^{3n} - p^{2n}))^{(p^n+1)(p^{2n}-p^n-1)} (x - (p^{3n} - 2p^{2n} + p^n))^{p^n} (x - (2p^{3n} - 2p^{2n})). \end{aligned}$$

Thus, $\text{Q-spec}(\Gamma_G) = \{(p^{3n} - p^{2n})^{(p^n+1)(p^{2n}-p^n-1)}, (p^{3n} - 2p^{2n} + p^n)^{p^n}, (2p^{3n} - 2p^{2n})^1\}$.

Number of edges of Γ_G is $\frac{(p^{3n}-p^n)(p^{2n}-p^n-1)}{2}$. Thus, $|e(\Gamma_G)| = \frac{(p^{3n}-p^n)(p^{2n}-p^n-1)}{2} - \frac{(p^{3n}-p^n)(p^{2n}-p^n-1)}{2} = \frac{p^{6n}-p^{5n}-p^{4n}+p^{3n}}{2}$. Now

$$\begin{aligned} \left|p^{3n} - p^{2n} - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|}\right| &= \left|\frac{0}{p^{3n} - p^n}\right| = 0, \\ \left|p^{3n} - 2p^{2n} + p^n - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|}\right| &= \left|\frac{-p^{5n} + p^{4n} + p^{3n} - p^{2n}}{p^{3n} - p^n}\right| = \frac{p^{5n} - p^{4n} - p^{3n} + p^{2n}}{p^{3n} - p^n} \end{aligned}$$

and

$$\left|2p^{3n} - 2p^{2n} - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|}\right| = \left|\frac{p^{6n} - p^{5n} - p^{4n} + p^{3n}}{p^{3n} - p^n}\right| = \frac{p^{6n} - p^{5n} - p^{4n} + p^{3n}}{p^{3n} - p^n}.$$

Therefore, $LE^+(\Gamma_G) = (p^n + 1)(p^{2n} - p^n - 1) \times 0 + p^n \times \frac{p^{5n} - p^{4n} - p^{3n} + p^{2n}}{p^{3n} - p^n} + \frac{p^{6n} - p^{5n} - p^{4n} + p^{3n}}{p^{3n} - p^n}$ and the result follows on simplification. \square

Theorem 5.6. *If G is isomorphic to the Hanaki group $A(n, p)$ then*

(a) $E(\Gamma_G) = LE(\Gamma_G) = LE^+(\Gamma_G)$.

(b) Γ_G is non-hypoenergetic, non-hyperenergetic, not L -hyperenergetic and not Q -hyperenergetic.

Proof. (a) Using Theorems 5.4 and 5.5 we have $E(\Gamma_G) = LE(\Gamma_G) = LE^+(\Gamma_G) = 2p^{2n}(p^n - 1)$ and hence the result follows.

(b) Here, $|v(\Gamma_G)| = p^{3n} - p^n$ and $E(K_{|v(\Gamma_G)|}) = LE(K_{|v(\Gamma_G)|}) = LE^+(K_{|v(\Gamma_G)|}) = 2(p^{2n}(p^n - 1) + p^{2n} - p^n - 1)$. Therefore, by Theorems 5.4 and 5.5, we have $E(\Gamma_G) - |v(\Gamma_G)| = p^n(p^n - 1)^2 > 0$ and $E(K_{|v(\Gamma_G)|}) - E(\Gamma_G) = LE(K_{|v(\Gamma_G)|}) - LE(\Gamma_G) = LE^+(K_{|v(\Gamma_G)|}) - LE^+(\Gamma_G) = 2(p^n(p^n - 1) - 1) > 0$. Hence, the results follow. \square

5.2 The semi-dihedral groups, SD_{8n}

We consider $SD_{8n} := \langle a, b : a^{4n} = b^2 = 1, bab^{-1} = a^{2n-1} \rangle$, the semi-dihedral groups of order $8n$ (where $n > 1$). Results regarding different energies of non-commuting graph of SD_{8n} are given below.

Theorem 5.7 ([14, Theorem 4.1.1 (b) and Theorem 4.3.2]). *Let G be isomorphic to SD_{8n} .*

(a) *If n is odd then*

$$E(\Gamma_{SD_{8n}}) = 4(n-1) + 4\sqrt{(n-1)(5n-1)} \quad \text{and} \quad LE(\Gamma_{SD_{8n}}) = \frac{8n(4n^2 - 10n + 7)}{2n-1}.$$

(b) *If n is even then*

$$E(\Gamma_{SD_{8n}}) = 2(2n-1) + 2\sqrt{(2n-1)(10n-1)} \quad \text{and} \quad LE(\Gamma_{SD_{8n}}) = \frac{8n(8n^2 - 8n + 3)}{4n-1}.$$

Theorem 5.8. *Let G be isomorphic to SD_{8n} , where n is odd. Then*

$$\begin{aligned} \text{Q-spec}(\Gamma_{SD_{8n}}) &= \left\{ (8n-8)^{3n}, (4n)^{4n-5}, (8n-12)^{n-1}, \left(8n-6+2\sqrt{8n^2-16n+9}\right)^1, \right. \\ &\quad \left. \left(8n-6-2\sqrt{8n^2-16n+9}\right)^1 \right\} \end{aligned}$$

and

$$LE^+(\Gamma_{SD_{8n}}) = \begin{cases} 36 + 4\sqrt{33}, & \text{if } n = 3 \\ \frac{32n^3 - 112n^2 + 96n - 12}{2n-1} + 4\sqrt{8n^2 - 16n + 9}, & \text{if } n \geq 5. \end{cases}$$

Proof. If $G \cong SD_{8n}$ and n is odd then $|v(\Gamma_{SD_{8n}})| = 8n - 4$ and $\Gamma_{SD_{8n}} = K_{n,4,1,(4n-4)}$. Using Theorem 1.6(b), we have

$$\begin{aligned} Q_{\Gamma_{SD_{8n}}}(x) &= \prod_{i=1}^2 (x - (8n - 4) + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - (8n - 4) + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - (8n - 4) + 2p_i} \right) \\ &= (x - 8n + 8)^{3n} (x - 4n)^{4n-5} (x - 8n + 12)^n (x - 4) \left(1 - \frac{4n}{x - 8n + 12} - \frac{4n - 4}{x - 4} \right) \\ &= (x - (8n - 8))^{3n} (x - 4n)^{4n-5} (x - (8n - 12))^{n-1} (x^2 - (16n - 12)x + 32n^2 - 32n). \end{aligned}$$

Thus, $\text{Q-spec}(\Gamma_{SD_{8n}}) = \left\{ (8n - 8)^{3n}, (4n)^{4n-5}, (8n - 12)^{n-1}, (8n - 6 + 2\sqrt{8n^2 - 16n + 9})^1, (8n - 6 - 2\sqrt{8n^2 - 16n + 9})^1 \right\}$.

Number of edges of $\Gamma_{SD_{8n}}^c$ is $8n^2 - 12n + 10$. Thus, $|e(\Gamma_{SD_{8n}})| = \frac{(8n-4)(8n-4-1)}{2} - (8n^2 - 12n + 10) = 24n(n-1)$.

Now

$$\begin{aligned} \left| 8n - 8 - \frac{2|e(\Gamma_{SD_{8n}})|}{|v(\Gamma_{SD_{8n}})|} \right| &= \left| \frac{4(n-1)(n-2)}{2n-1} \right| = \frac{4(n-1)(n-2)}{2n-1}, \\ \left| 4n - \frac{2|e(\Gamma_{SD_{8n}})|}{|v(\Gamma_{SD_{8n}})|} \right| &= \left| \frac{4n(2-n)}{2n-1} \right| = \frac{4n(n-2)}{2n-1}, \\ \left| 8n - 12 - \frac{2|e(\Gamma_{SD_{8n}})|}{|v(\Gamma_{SD_{8n}})|} \right| &= \left| \frac{4(n^2 - 5n + 3)}{2n-1} \right| = \begin{cases} \frac{4(-n^2+5n-3)}{2n-1}, & \text{if } n \leq 3 \\ \frac{4(n^2-5n+3)}{2n-1}, & \text{if } n \geq 5, \end{cases} \end{aligned}$$

$$\begin{aligned} \left| 8n - 6 + 2\sqrt{8n^2 - 16n + 9} - \frac{2|e(\Gamma_{SD_{8n}})|}{|v(\Gamma_{SD_{8n}})|} \right| &= \left| 2\sqrt{8n^2 - 16n + 9} + (2n - 3) + \frac{3}{2n-1} \right| \\ &= 2\sqrt{8n^2 - 16n + 9} + (2n - 3) + \frac{3}{2n-1} \end{aligned}$$

and

$$\begin{aligned} \left| 8n - 6 - 2\sqrt{8n^2 - 16n + 9} - \frac{2|e(\Gamma_{SD_{8n}})|}{|v(\Gamma_{SD_{8n}})|} \right| &= \left| -2\sqrt{8n^2 - 16n + 9} + (2n - 3) + \frac{3}{2n-1} \right| \\ &= 2\sqrt{8n^2 - 16n + 9} - (2n - 3) - \frac{3}{2n-1}. \end{aligned}$$

Therefore, for $n = 3$ we have $LE^+(\Gamma_{SD_{8n}}) = 36 + 4\sqrt{33}$. For $n \geq 5$ we have

$$\begin{aligned} LE^+(\Gamma_{SD_{8n}}) &= 3n \times \frac{4(n-1)(n-2)}{2n-1} + (4n-5) \times \frac{4n(n-2)}{2n-1} + (n-1) \times \frac{4(n^2-5n+3)}{2n-1} \\ &\quad + 2\sqrt{8n^2 - 16n + 9} - 2n - 3 + \frac{3}{2n-1} + 2\sqrt{8n^2 - 16n + 9} + 2n + 3 - \frac{3}{2n-1} \end{aligned}$$

and the result follows on simplification. \square

Theorem 5.9. *Let G be isomorphic to SD_{8n} , where n is even. Then*

$$\begin{aligned} \text{Q-spec}(\Gamma_{SD_{8n}}) &= \left\{ (8n - 4)^{2n}, (4n)^{4n-3}, (8n - 6)^{2n-1}, (8n - 3 + \sqrt{32n^2 - 32n + 9})^1, \right. \\ &\quad \left. (8n - 3 - \sqrt{32n^2 - 32n + 9})^1 \right\} \end{aligned}$$

and

$$LE^+(\Gamma_{SD_{8n}}) = \begin{cases} \frac{134}{7} + 2\sqrt{73}, & \text{if } n = 2 \\ \frac{64n^3 - 128n^2 + 64n - 6}{4n-1} + 2\sqrt{32n^2 - 32n + 9}, & \text{if } n \geq 4. \end{cases}$$

Proof. If $G \cong SD_{8n}$ and n is even then $|v(\Gamma_{SD_{8n}})| = 8n - 2$ and $\Gamma_{SD_{8n}} = K_{2n,2,1,(4n-2)}$. Using Theorem 1.6(b), we have

$$\begin{aligned} Q_{\Gamma_{SD_{8n}}}(x) &= \prod_{i=1}^2 (x - (8n - 2) + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - (8n - 2) + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - (8n - 2) + 2p_i} \right) \\ &= (x - 8n + 4)^{2n} (x - 4n)^{4n-3} (x - 8n + 6)^{2n} (x - 2) \left(1 - \frac{4n}{x - 8n + 6} - \frac{4n - 2}{x - 2} \right) \\ &= (x - (8n - 4))^{2n} (x - 4n)^{4n-3} (x - (8n - 6))^{2n-1} (x^2 - (16n - 6)x + 32n^2 - 16n). \end{aligned}$$

Thus, $Q\text{-spec}(\Gamma_{SD_{8n}}) = \left\{ (8n-4)^{2n}, (4n)^{4n-3}, (8n-6)^{2n-1}, (8n-3 + \sqrt{32n^2 - 32n + 9})^1, (8n-3 - \sqrt{32n^2 - 32n + 9})^1 \right\}$.

Number of edges of $\Gamma_{SD_{8n}}^c$ is $8n^2 - 8n + 3$. Therefore, $|e(\Gamma_{SD_{8n}})| = \frac{(8n-2)(8n-2-1)}{2} - (8n^2 - 8n + 3) = 12n(2n-1)$. Now

$$\begin{aligned} \left| 8n-4 - \frac{2|e(\Gamma_{SD_{8n}})|}{|v(\Gamma_{SD_{8n}})|} \right| &= \left| \frac{(8n-4)(n-1)}{4n-1} \right| = \frac{(8n-4)(n-1)}{4n-1}, \\ \left| 4n - \frac{2|e(\Gamma_{SD_{8n}})|}{|v(\Gamma_{SD_{8n}})|} \right| &= \left| \frac{-8n(n-1)}{4n-1} \right| = \frac{8n(n-1)}{4n-1}, \\ \left| 8n-6 - \frac{2|e(\Gamma_{SD_{8n}})|}{|v(\Gamma_{SD_{8n}})|} \right| &= \left| \frac{(8n^2 - 20n + 6)}{4n-1} \right| = \begin{cases} \frac{2}{7}, & \text{if } n=2 \\ \frac{(8n^2-20n+6)}{4n-1}, & \text{if } n \geq 4, \end{cases} \end{aligned}$$

$$\begin{aligned} \left| 8n-3 + \sqrt{32n^2 - 32n + 9} - \frac{2|e(\Gamma_{SD_{8n}})|}{|v(\Gamma_{SD_{8n}})|} \right| &= \left| \sqrt{32n^2 - 32n + 9} + 2n - \frac{3}{2} + \frac{3}{8n-2} \right| \\ &= \sqrt{32n^2 - 32n + 9} + 2n - \frac{3}{2} + \frac{3}{8n-2} \end{aligned}$$

and

$$\begin{aligned} \left| 8n-3 - \sqrt{32n^2 - 32n + 9} - \frac{2|e(\Gamma_{SD_{8n}})|}{|v(\Gamma_{SD_{8n}})|} \right| &= \left| -\sqrt{32n^2 - 32n + 9} + 2n - \frac{3}{2} + \frac{3}{8n-2} \right| \\ &= \sqrt{32n^2 - 32n + 9} - 2n + \frac{3}{2} - \frac{3}{8n-2}. \end{aligned}$$

Therefore, for $n=2$, we have $LE^+(\Gamma_{SD_{8n}}) = \frac{134}{7} + 2\sqrt{73}$. For $n \geq 4$, we have

$$\begin{aligned} LE^+(\Gamma_{SD_{8n}}) &= 2n \times \frac{(8n-4)(n-1)}{4n-1} + (4n-3) \times \frac{8n(n-1)}{4n-1} + (2n-1) \times \frac{(8n^2-20n+6)}{4n-1} + \\ &\quad \sqrt{32n^2 - 32n + 9} + 2n - \frac{3}{2} + \frac{3}{8n-2} + \sqrt{32n^2 - 32n + 9} - 2n + \frac{3}{2} - \frac{3}{8n-2} \end{aligned}$$

and the result follows on simplification. \square

Theorem 5.10. *If G is isomorphic to SD_{8n} then*

- (a) $E(\Gamma_{SD_{8n}}) < LE^+(\Gamma_{SD_{8n}}) < LE(\Gamma_{SD_{8n}})$.
- (b) $\Gamma_{SD_{8n}}$ is non-hypoenergetic as well as non-hyperenergetic.
- (c) $\Gamma_{SD_{8n}}$ is Q -hyperenergetic and L -hyperenergetic.

Proof. (a) **Case 1:** n is odd

For $n=3$, using Theorems 5.7 and 5.8, we have $E(\Gamma_{SD_{24}}) = 8 + 8\sqrt{7}$, $LE(\Gamma_{SD_{24}}) = \frac{312}{5}$ and $LE^+(\Gamma_{SD_{24}}) = 36 + 4\sqrt{33}$. Clearly, $E(\Gamma_{SD_{24}}) < LE^+(\Gamma_{SD_{24}}) < LE(\Gamma_{SD_{24}})$.

For $n \geq 5$, using Theorems 5.7 and 5.8, we have

$$LE(\Gamma_{SD_{8n}}) - LE^+(\Gamma_{SD_{8n}}) = \frac{32n^2 - 40n + 12}{2n-1} - 4\sqrt{8n^2 - 16n + 9} \quad (5.1)$$

and

$$LE^+(\Gamma_{SD_{8n}}) - E(\Gamma_{SD_{8n}}) = \frac{32n^3 - 116n^2 + 102n - 14}{2n-1} + 4\sqrt{8n^2 - 16n + 9} - 4\sqrt{5n^2 - 6n + 1}. \quad (5.2)$$

Since $32n^2 - 40n + 12 > 0$, $4(2n-1)\sqrt{8n^2 - 16n + 9} > 0$ and $(32n^2 - 40n + 12)^2 - (4\sqrt{8n^2 - 16n + 9})^2 (2n-1)^2 = 512n^3(n-2) + 128n(5n-1) > 0$ we have

$$32n^2 - 40n + 12 - 4(2n-1)\sqrt{8n^2 - 16n + 9} > 0.$$

Therefore, by (5.1), $(2n-1)(LE(\Gamma_{SD_{8n}}) - LE^+(\Gamma_{SD_{8n}})) > 0$. Hence, $LE(\Gamma_{SD_{8n}}) > LE^+(\Gamma_{SD_{8n}})$.

Again, $\sqrt{8n^2 - 16n + 9} > 0$, $\sqrt{5n^2 - 6n + 1} > 0$ and $(\sqrt{8n^2 - 16n + 9})^2 - (\sqrt{5n^2 - 6n + 1})^2 = n(3n - 10) + 8 > 0$. Therefore, $\sqrt{8n^2 - 16n + 9} - \sqrt{5n^2 - 6n + 1} > 0$. Since $32n^3 - 116n^2 + 102n - 14 > 0$ we have $\frac{32n^3 - 116n^2 + 102n - 14}{2n-1} + 4\sqrt{8n^2 - 16n + 9} - 4\sqrt{5n^2 - 6n + 1} > 0$. Therefore, by (5.2), $LE^+(\Gamma_{SD_{8n}}) > E(\Gamma_{SD_{8n}})$. Hence, $E(\Gamma_{SD_{8n}}) < LE^+(\Gamma_{SD_{8n}}) < LE(\Gamma_{SD_{8n}})$.

Case 2: n is even

For $n = 2$, using Theorems 5.7 and 5.9, we have $E(\Gamma_{SD_{16}}) = 6 + 2\sqrt{57}$, $LE(\Gamma_{SD_{16}}) = \frac{304}{7}$ and $LE^+(\Gamma_{SD_{16}}) = \frac{134}{7} + 2\sqrt{73}$. Clearly, $E(\Gamma_{SD_{16}}) < LE^+(\Gamma_{SD_{16}}) < LE(\Gamma_{SD_{16}})$.

For $n \geq 4$, using Theorems 5.7 and 5.9, we have

$$LE(\Gamma_{SD_{8n}}) - LE^+(\Gamma_{SD_{8n}}) = \frac{64n^2 - 40n + 6}{4n - 1} - 2\sqrt{32n^2 - 32n + 9} \quad (5.3)$$

and

$$LE^+(\Gamma_{SD_{8n}}) - E(\Gamma_{SD_{8n}}) = \frac{16n^2(4n - 9) + 76n - 8}{4n - 1} + 2\sqrt{32n^2 - 32n + 9} - 2\sqrt{20n^2 - 12n + 1}. \quad (5.4)$$

Since $64n^2 - 40n + 6 > 0$, $2(4n - 1)\sqrt{32n^2 - 32n + 9} > 0$ and $(64n^2 - 40n + 6)^2 - (2\sqrt{32n^2 - 32n + 9})^2(4n - 1)^2 = 2048n^3(n - 1) + 64n(10n - 1) > 0$ we have $64n^2 - 40n + 6 - 2(4n - 1)\sqrt{32n^2 - 32n + 9} > 0$. Therefore, by (5.3), $(4n - 1)(LE^+(\Gamma_{SD_{8n}}) - LE(\Gamma_{SD_{8n}})) > 0$. Hence, $LE(\Gamma_{SD_{8n}}) > LE^+(\Gamma_{SD_{8n}})$.

Again, $\sqrt{32n^2 - 32n + 9} > 0$, $\sqrt{20n^2 - 12n + 1} > 0$ and $(\sqrt{32n^2 - 32n + 9})^2 - (\sqrt{20n^2 - 12n + 1})^2 = 4n(3n - 5) + 8 > 0$. Therefore, $\sqrt{32n^2 - 32n + 9} - \sqrt{20n^2 - 12n + 1} > 0$. Since $16n^2(4n - 9) + 76n - 8 > 0$ we have $\frac{16n^2(4n - 9) + 76n - 8}{2n - 1} + 2\sqrt{32n^2 - 32n + 9} - 2\sqrt{20n^2 - 12n + 1} > 0$. Therefore, by (5.4), $LE^+(\Gamma_{SD_{8n}}) > E(\Gamma_{SD_{8n}})$. Hence, $E(\Gamma_{SD_{8n}}) < LE^+(\Gamma_{SD_{8n}}) < LE(\Gamma_{SD_{8n}})$.

(b) **Case 1:** n is odd

Here $|v(\Gamma_{SD_{8n}})| = 8n - 4$ and $E(K_{|v(\Gamma_{SD_{8n}})|}) = LE(K_{|v(\Gamma_{SD_{8n}})|}) = LE^+(K_{|v(\Gamma_{SD_{8n}})|}) = 16n - 10$. Using Theorem 5.7, we have

$$E(\Gamma_{SD_{8n}}) - |v(\Gamma_{SD_{8n}})| = 4\sqrt{(n - 1)(5n - 1)} - 4n \quad (5.5)$$

and

$$E(K_{|v(\Gamma_{SD_{8n}})|}) - E(\Gamma_{SD_{8n}}) = 12n - 6 - 4\sqrt{(n - 1)(5n - 1)}. \quad (5.6)$$

Since $4\sqrt{(n - 1)(5n - 1)} > 0$, $4n > 0$ and $(4\sqrt{(n - 1)(5n - 1)})^2 - (4n)^2 = 16(4n^2 - 6n + 1) > 0$ we have $4\sqrt{(n - 1)(5n - 1)} - 4n > 0$. Therefore, by (5.5), $E(\Gamma_{SD_{8n}}) > |v(\Gamma_{SD_{8n}})|$.

Again, $4\sqrt{(n - 1)(5n - 1)} > 0$, $12n - 6 > 0$ and $(12n - 6)^2 - (4\sqrt{(n - 1)(5n - 1)})^2 = 4(16n^2 - 12n + 5) > 0$ and so $12n - 6 - 4\sqrt{(n - 1)(5n - 1)} > 0$. Therefore, by (5.6), $E(K_{|v(\Gamma_{SD_{8n}})|}) > E(\Gamma_{SD_{8n}})$.

Case 2: n is even

Here $|v(\Gamma_{SD_{8n}})| = 8n - 2$ and $E(K_{|v(\Gamma_{SD_{8n}})|}) = LE(K_{|v(\Gamma_{SD_{8n}})|}) = LE^+(K_{|v(\Gamma_{SD_{8n}})|}) = 16n - 16$. Using Theorem 5.7, we have

$$E(\Gamma_{SD_{8n}}) - |v(\Gamma_{SD_{8n}})| = 2\left(\sqrt{(2n - 1)(10n - 1)} - 2n\right) \quad (5.7)$$

and

$$E(K_{|v(\Gamma_{SD_{8n}})|}) - E(\Gamma_{SD_{8n}}) = 2\left(3(2n - 1) + 1 - \sqrt{(2n - 1)(10n - 1)}\right). \quad (5.8)$$

Since $\sqrt{(2n - 1)(10n - 1)} > 0$, $2n > 0$ and $(\sqrt{(2n - 1)(10n - 1)})^2 - (2n)^2 = 4n(4n - 3) + 1 > 0$ we have $\sqrt{(2n - 1)(10n - 1)} - 2n > 0$. Therefore, by (5.7), $E(\Gamma_{SD_{8n}}) > |v(\Gamma_{SD_{8n}})|$.

Again, $\sqrt{(2n - 1)(10n - 1)} > 0$, $3(2n - 1) + 1 > 0$ and $(3(2n - 1) + 1)^2 - (\sqrt{(2n - 1)(10n - 1)})^2 = 4n(4n - 3) + 3 > 0$ and so $3(2n - 1) + 1 - \sqrt{(2n - 1)(10n - 1)} > 0$. Therefore, by (5.8), $E(K_{|v(\Gamma_{SD_{8n}})|}) > E(\Gamma_{SD_{8n}})$.

(c) **Case 1:** n is odd

Using Theorem 5.8, for $n = 3$ we have $LE^+(\Gamma_{SD_{24}}) = 36 + 4\sqrt{33}$ and $LE^+(K_{|v(\Gamma_{SD_{24}})|}) = 38$. Also, for $n \geq 5$ we have

$$LE^+(\Gamma_{SD_{8n}}) - LE^+(K_{|v(\Gamma_{SD_{8n}})|}) = \frac{2(8n^2(2n - 9) + 66n - 11)}{2n - 1} + 4\sqrt{8n^2 - 14n + 9} > 0.$$

Therefore, $LE^+(\Gamma_{SD_{8n}}) > LE^+(K_{|v(\Gamma_{SD_{8n}})|})$ which implies $\Gamma_{SD_{8n}}$ is Q-hyperenergetic and consequently part (a) implies $\Gamma_{SD_{8n}}$ is L-hyperenergetic.

Case 2: n is even

For $n = 2$ we have $LE^+(K_{|v(\Gamma_{SD_{16}})|}) = 16$ and using Theorem 5.9, $LE^+(\Gamma_{SD_{16}}) = \frac{134}{7} + 2\sqrt{73}$. Therefore, $\Gamma_{SD_{16}}$ is Q-hyperenergetic and consequently part (a) implies $\Gamma_{SD_{16}}$ is L-hyperenergetic.

For $n \geq 4$, using Theorem 5.9, we have

$$LE^+(\Gamma_{SD_{8n}}) - LE^+(K_{|v(\Gamma_{SD_{8n}})|}) = \frac{64n^2(n-3) + 144n - 22}{4n-1} + 2\sqrt{32n^2 - 32n + 9} > 0.$$

Therefore, $LE^+(\Gamma_{SD_{8n}}) > LE^+(K_{|v(\Gamma_{SD_{8n}})|})$ which implies $\Gamma_{SD_{8n}}$ is Q-hyperenergetic and consequently part (a) implies $\Gamma_{SD_{8n}}$ is L-hyperenergetic. \square

In Theorem 5.10, we compare $E(\Gamma_{SD_{8n}})$, $LE(\Gamma_{SD_{8n}})$ and $LE^+(\Gamma_{SD_{8n}})$. However, in the following figures, we show how close are they.




Figure 13: Energies of $\Gamma_{SD_{8n}}$, where n is odd




Figure 14: Energies of $\Gamma_{SD_{8n}}$, where n is even

5.3 The groups, V_{8n}

We consider $V_{8n} := \langle a, b : a^{4n} = b^4 = 1, b^{-1}ab^{-1} = bab = a^{-1} \rangle$, the groups of order $8n$ (where $n > 1$). We compute Signless Laplacian spectrum, Signless Laplacian energy, Laplacian spectrum, Laplacian energy and spectrum and energy of $\Gamma_{V_{8n}}$. If n is odd then energy and Laplacian energy of $\Gamma_{V_{8n}}$ are as given in the following theorem.

Theorem 5.11 ([14, Theorem 4.1.1 (a) and Theorem 4.3.1]). *Let G be isomorphic to V_{8n} , where n is odd. Then*

$$E(\Gamma_{V_{8n}}) = 2(2n-1) + 2\sqrt{(2n-1)(10n-1)} \text{ and } LE(\Gamma_{V_{8n}}) = \frac{8n(8n^2 - 8n + 3)}{4n-1}.$$

Theorem 5.12. *Let G be isomorphic to V_{8n} , where n is odd. Then*

$$\text{Q-spec}(\Gamma_{V_{8n}}) = \left\{ (8n-4)^{2n}, (4n)^{4n-3}, (8n-6)^{2n-1}, \left(8n-3 + \sqrt{32n^2 - 32n + 9}\right)^1, \left(8n-3 - \sqrt{32n^2 - 32n + 9}\right)^1 \right\}$$

$$\text{and } LE^+(\Gamma_{V_{8n}}) = \frac{64n^3 - 128n^2 + 64n - 6}{4n-1} + 2\sqrt{32n^2 - 32n + 9}.$$

Proof. If $G \cong V_{8n}$ and n is odd then $|v(\Gamma_{V_{8n}})| = 8n - 2$ and $\Gamma_{V_{8n}} = K_{2n, 2, 1, (4n-2)}$. Using Theorem 1.6(b), we have

$$\begin{aligned} Q_{\Gamma_{V_{8n}}}(x) &= \prod_{i=1}^2 (x - (8n-2) + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - (8n-2) + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - (8n-2) + 2p_i} \right) \\ &= (x - 8n + 4)^{2n} (x - 4n)^{4n-3} (x - 8n + 6)^{2n} (x - 2) \left(1 - \frac{4n}{x - 8n + 6} - \frac{4n-2}{x-2} \right) \\ &= (x - (8n-4))^{2n} (x - 4n)^{4n-3} (x - (8n-6))^{2n-1} (x^2 - (16n-6)x + 32n^2 - 16n). \end{aligned}$$

$$\text{Thus, } \text{Q-spec}(\Gamma_{V_{8n}}) = \left\{ (8n-4)^{2n}, (4n)^{4n-3}, (8n-6)^{2n-1}, (8n-3 + \sqrt{32n^2 - 32n + 9})^1, \right. \\ \left. (8n-3 - \sqrt{32n^2 - 32n + 9})^1 \right\}.$$

Number of edges of $\Gamma_{V_{8n}}^c$ is $8n^2 - 8n + 3$. Therefore, $|e(\Gamma_{V_{8n}})| = \frac{(8n-2)(8n-2-1)}{2} - (8n^2 - 8n + 3) = 12n(2n-1)$.

Now

$$\left| 8n-4 - \frac{2|e(\Gamma_{V_{8n}})|}{|v(\Gamma_{V_{8n}})|} \right| = \left| \frac{(8n-4)(n-1)}{4n-1} \right| = \frac{(8n-4)(n-1)}{4n-1}, \\ \left| 4n - \frac{2|e(\Gamma_{V_{8n}})|}{|v(\Gamma_{V_{8n}})|} \right| = \left| \frac{-8n(n-1)}{4n-1} \right| = \frac{8n(n-1)}{4n-1}, \\ \left| 8n-6 - \frac{2|e(\Gamma_{V_{8n}})|}{|v(\Gamma_{V_{8n}})|} \right| = \left| \frac{(8n^2 - 20n + 6)}{4n-1} \right| = \frac{8n^2 - 20n + 6}{4n-1},$$

$$\left| 8n-3 + \sqrt{32n^2 - 32n + 9} - \frac{2|e(\Gamma_{SD_{8n}})|}{|v(\Gamma_{SD_{8n}})|} \right| = \left| \sqrt{32n^2 - 32n + 9} + 2n - \frac{3}{2} + \frac{3}{8n-2} \right| \\ = \sqrt{32n^2 - 32n + 9} + 2n - \frac{3}{2} + \frac{3}{8n-2}$$

and

$$\left| 8n-3 - \sqrt{32n^2 - 32n + 9} - \frac{2|e(\Gamma_{SD_{8n}})|}{|v(\Gamma_{SD_{8n}})|} \right| = \left| -\sqrt{32n^2 - 32n + 9} + 2n - \frac{3}{2} + \frac{3}{8n-2} \right| \\ = \sqrt{32n^2 - 32n + 9} - 2n + \frac{3}{2} - \frac{3}{8n-2}.$$

Therefore, for $n \geq 3$, we have

$$LE^+(\Gamma_{V_{8n}}) = 2n \times \frac{(8n-4)(n-1)}{4n-1} + (4n-3) \times \frac{8n(n-1)}{4n-1} + (2n-1) \times \frac{8n^2 - 20n + 6}{4n-1} + \\ \sqrt{32n^2 - 32n + 9} + 2n - \frac{3}{2} + \frac{3}{8n-2} + \sqrt{32n^2 - 32n + 9} - 2n + \frac{3}{2} - \frac{3}{8n-2}$$

and the result follows on simplification. \square

Theorem 5.13. *Let G be isomorphic to V_{8n} , where n is even. Then*

$$\text{Q-spec}(\Gamma_{V_{8n}}) \\ = \left\{ (8n-8)^{3n}, (4n)^{4n-5}, (8n-12)^{n-1}, \left(8n-6 + 2\sqrt{8n^2 - 16n + 9} \right)^1, \left(8n-6 - 2\sqrt{8n^2 - 16n + 9} \right)^1 \right\},$$

$$\text{L-spec}(\Gamma_{V_{8n}}) = \{0, (4n)^{4n-5}, (8n-8)^{3n}, (8n-4)^n\} \text{ and}$$

$$\text{Spec}(\Gamma_{V_{8n}}) = \left\{ 0^{7n-5}, (-4)^{n-1}, \left(2(n-1) + 2\sqrt{(n-1)(5n-1)} \right)^1, \left(2(n-1) + 2\sqrt{(n-1)(5n-1)} \right)^1 \right\}.$$

Further

$$LE^+(\Gamma_{V_{8n}}) = \begin{cases} \frac{24n^3 - 64n^2 + 32n + 12}{2n-1} + 4\sqrt{8n^2 - 16n + 9}, & \text{if } n \leq 4 \\ \frac{32n^3 - 112n^2 + 96n - 12}{2n-1} + 4\sqrt{8n^2 - 16n + 9}, & \text{if } n \geq 6, \end{cases}$$

$$LE(\Gamma_{V_{8n}}) = \frac{8n(4n^2 - 10n + 7)}{2n-1} \text{ and } E(\Gamma_{V_{8n}}) = 4(n-1) + 4\sqrt{(n-1)(5n-1)}.$$

Proof. If $G \cong V_{8n}$ and n is even then $|v(\Gamma_{V_{8n}})| = 8n-4$ and $\Gamma_{V_{8n}} = K_{n,4,1.(4n-4)}$. Using Theorem 1.6(b), we have

$$Q_{\Gamma_{V_{8n}}}(x) = \prod_{i=1}^2 (x - (8n-4) + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - (8n-4) + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - (8n-4) + 2p_i} \right) \\ = (x-8n+8)^{3n} (x-4n)^{4n-5} (x-8n+12)^n (x-4) \left(1 - \frac{4n}{x-8n+12} - \frac{4n-4}{x-4} \right) \\ = (x-(8n-8))^{3n} (x-4n)^{4n-5} (x-(8n-12))^{n-1} (x^2 - (16n-12)x + 32n^2 - 32n).$$

$$\text{Thus, } Q\text{-spec}(\Gamma_{V_{8n}}) = \left\{ (8n-8)^{3n}, (4n)^{4n-5}, (8n-12)^{n-1}, (8n-6+2\sqrt{8n^2-16n+9})^1, \right. \\ \left. (8n-6-2\sqrt{8n^2-16n+9})^1 \right\}.$$

Number of edges of $\Gamma_{V_{8n}}^c$ is $8n^2-12n+10$ and so $|e(\Gamma_{V_{8n}})| = \frac{(8n-4)(8n-4-1)}{2} - (8n^2-12n+10) = 24n(n-1)$.

Now

$$\left| 8n-8 - \frac{2|e(\Gamma_{V_{8n}})|}{|v(\Gamma_{V_{8n}})|} \right| = \left| \frac{4(n-1)(n-2)}{2n-1} \right| = \frac{4(n-1)(n-2)}{2n-1},$$

$$\left| 4n - \frac{2|e(\Gamma_{V_{8n}})|}{|v(\Gamma_{V_{8n}})|} \right| = \left| \frac{4n(2-n)}{2n-1} \right| = \frac{4n(n-2)}{2n-1},$$

$$\left| 8n-12 - \frac{2|e(\Gamma_{V_{8n}})|}{|v(\Gamma_{V_{8n}})|} \right| = \left| \frac{4(n^2-5n+3)}{2n-1} \right| = \begin{cases} \frac{4(-n^2+5n-3)}{2n-1}, & \text{if } n \leq 4 \\ \frac{4(n^2-5n+3)}{2n-1}, & \text{if } n \geq 6, \end{cases}$$

$$\left| 8n-6+2\sqrt{8n^2-16n+9} - \frac{2|e(\Gamma_{V_{8n}})|}{|v(\Gamma_{V_{8n}})|} \right| = \left| 2\sqrt{8n^2-16n+9} + (2n-3) + \frac{3}{2n-1} \right| \\ = 2\sqrt{8n^2-16n+9} + (2n-3) + \frac{3}{2n-1}$$

and

$$\left| 8n-6-2\sqrt{8n^2-16n+9} - \frac{2|e(\Gamma_{V_{8n}})|}{|v(\Gamma_{V_{8n}})|} \right| = \left| -2\sqrt{8n^2-16n+9} + (2n-3) + \frac{3}{2n-1} \right| \\ = 2\sqrt{8n^2-16n+9} - (2n-3) - \frac{3}{2n-1}.$$

Therefore, for $n \leq 4$, we have

$$LE^+(\Gamma_{V_{8n}}) = 3n \times \frac{4(n-1)(n-2)}{2n-1} + (4n-5) \times \frac{4n(n-2)}{2n-1} + (n-1) \times \frac{4(-n^2+5n-3)}{2n-1} + \\ 2\sqrt{8n^2-16n+9} - 2n - 3 + \frac{3}{2n-1} + 2\sqrt{8n^2-16n+9} + 2n + 3 - \frac{3}{2n-1}.$$

For $n \geq 6$, we have

$$LE^+(\Gamma_{V_{8n}}) = 3n \times \frac{4(n-1)(n-2)}{2n-1} + (4n-5) \times \frac{4n(n-2)}{2n-1} + (n-1) \times \frac{4(n^2-5n+3)}{2n-1} + \\ 2\sqrt{8n^2-16n+9} - 2n - 3 + \frac{3}{2n-1} + 2\sqrt{8n^2-16n+9} + 2n + 3 - \frac{3}{2n-1}.$$

Thus we get the required expressions for $LE^+(\Gamma_{V_{8n}})$ on simplification.

Since $\Gamma_{V_{8n}}^c = nK_4 \cup K_{4n-4}$, using Theorem 1.5, we have

$$L\text{-spec}(\Gamma_{V_{8n}}) = \left\{ (0)^1, \left(\sum_{i=1}^2 l_i m_i - m_2 \right)^{l_2(m_2-1)}, \left(\sum_{i=1}^2 l_i m_i - m_1 \right)^{l_1(m_1-1)}, \left(\sum_{i=1}^2 l_i m_i \right)^{\sum_{i=1}^2 l_i - 1} \right\} \\ = \{(0)^1, (n \cdot 4 + 4n - 4 - 4n + 4)^{1(4n-4-1)}, (n \cdot 4 + 4n - 4 - 4)^{n(4-1)}, (n \cdot 4 + 4n - 4)^{n+1-1}\} \\ = \{(0)^1, (4n)^{4n-5}, (8n-8)^{3n}, (8n-4)^n\}.$$

Now

$$\left| 0 - \frac{2|e(\Gamma_{V_{8n}})|}{|v(\Gamma_{V_{8n}})|} \right| = \left| \frac{-12n(n-1)}{2n-1} \right| = \frac{12n(n-1)}{2n-1}, \quad \left| 4n - \frac{2|e(\Gamma_{V_{8n}})|}{|v(\Gamma_{V_{8n}})|} \right| = \left| \frac{4n(2-n)}{2n-1} \right| = \frac{4n(n-2)}{2n-1},$$

$$\left| 8n-8 - \frac{2|e(\Gamma_{V_{8n}})|}{|v(\Gamma_{V_{8n}})|} \right| = \left| \frac{4(n-1)(n-2)}{2n-1} \right| = \frac{4(n-1)(n-2)}{2n-1}$$

and

$$\left| 8n-4 - \frac{2|e(\Gamma_{V_{8n}})|}{|v(\Gamma_{V_{8n}})|} \right| = \left| \frac{4(n^2-n+1)}{2n-1} \right| = \frac{4(n^2-n+1)}{2n-1}.$$

Therefore, $LE(\Gamma_{V_{8n}}) = 1 \times \frac{12n(n-1)}{2n-1} + (4n-5) \times \frac{4n(n-2)}{2n-1} + 3n \times \frac{4(n-1)(n-2)}{2n-1} + n \times \frac{4(n^2-n+1)}{2n-1}$ and we get the required expression for $LE(\Gamma_{V_{8n}})$ on simplification.

Since $\Gamma_{V_{8n}}$ is a complete $(n+1)$ -partite graph with $8n-4$ vertices and $\Gamma_{V_{8n}} = K_{n,4,1,(4n-4)}$. Therefore, using Theorem 1.6(a), the characteristic polynomial of $\Gamma_{V_{8n}}$ is

$$\begin{aligned} P_{\Gamma_{V_{8n}}}(x) &= x^{(8n-4)-(n+1)}(x+4)^{n-1}(x+4n-4)^{1-1}(x^2 - (4n-4)x - 16n^2 - 16n) \\ &= x^{7n-5}(x+4)^{n-1}(x^2 - (4n-4)x - 16n^2 - 16n). \end{aligned}$$

Thus, $\text{Spec}(\Gamma_{V_{8n}}) = \left\{ (0)^{7n-5}, (-4)^{n-1}, \left(2(n-1) + 2\sqrt{(n-1)(5n-1)}\right)^1, \left(2(n-1) + 2\sqrt{(n-1)(5n-1)}\right)^1 \right\}$.

Therefore, $E(\Gamma_{V_{8n}}) = (7n-5) \times |0| + (n-1) \times |-4| + |2(n-1) + 2\sqrt{(n-1)(5n-1)}| + |2(n-1) - 2\sqrt{(n-1)(5n-1)}|$ and we get the required expression for $E(\Gamma_{V_{8n}})$ on simplification. \square

Theorem 5.14. *If G is isomorphic to V_{8n} then*

- (a) $E(\Gamma_{V_{8n}}) \leq LE^+(\Gamma_{V_{8n}}) \leq LE(\Gamma_{V_{8n}})$; equality holds if and only if $G \cong V_{16}$.
- (b) $\Gamma_{V_{8n}}$ is non-hypoenergetic as well as non-hyperenergetic.
- (c) $\Gamma_{V_{16}}$ is not L -hyperenergetic and not Q -hyperenergetic. If $n \neq 2$ then $\Gamma_{V_{8n}}$ is Q -hyperenergetic and L -hyperenergetic.

Proof. (a) **Case 1:** n is odd

Using Theorems 5.11 and 5.12, we have

$$LE(\Gamma_{V_{8n}}) - LE^+(\Gamma_{V_{8n}}) = \frac{64n^2 - 40n + 6}{4n-1} - 2\sqrt{32n^2 - 32n + 9} \quad (5.9)$$

and

$$LE^+(\Gamma_{V_{8n}}) - E(\Gamma_{V_{8n}}) = \frac{16n^2(4n-9) + 76n - 8}{4n-1} + 2\sqrt{32n^2 - 32n + 9} - 2\sqrt{20n^2 - 12n + 1}. \quad (5.10)$$

Since $64n^2 - 40n + 6 > 0$, $2(4n-1)\sqrt{32n^2 - 32n + 9} > 0$ and $(64n^2 - 40n + 6)^2 - (2\sqrt{32n^2 - 32n + 9})^2(4n-1)^2 = 2048n^3(n-1) + 64n(10n-1) > 0$ we have $64n^2 - 40n + 6 - 2(4n-1)\sqrt{32n^2 - 32n + 9} > 0$. Therefore, by (5.9), $(4n-1)(LE^+(\Gamma_{V_{8n}}) - LE(\Gamma_{V_{8n}})) > 0$. Hence, $LE(\Gamma_{V_{8n}}) > LE^+(\Gamma_{V_{8n}})$.

Again, $\sqrt{32n^2 - 32n + 9} > 0$, $\sqrt{20n^2 - 12n + 1} > 0$ and $(\sqrt{32n^2 - 32n + 9})^2 - (\sqrt{20n^2 - 12n + 1})^2 = 4n(3n-5) + 8 > 0$. Therefore, $\sqrt{32n^2 - 32n + 9} - \sqrt{20n^2 - 12n + 1} > 0$. Since $16n^2(4n-9) + 76n - 8 > 0$ we have $\frac{16n^2(4n-9) + 76n - 8}{2n-1} + 2\sqrt{32n^2 - 32n + 9} - 2\sqrt{20n^2 - 12n + 1} > 0$. Therefore, by (5.10), $LE^+(\Gamma_{V_{8n}}) > E(\Gamma_{V_{8n}})$. Hence, $E(\Gamma_{V_{8n}}) < LE^+(\Gamma_{V_{8n}}) < LE(\Gamma_{V_{8n}})$.

Case 2: n is even

Using Theorems 5.11 and 5.13, for $n \leq 4$, we have

$$LE(\Gamma_{V_{8n}}) - LE^+(\Gamma_{V_{8n}}) = \frac{8n^3 - 16n^2 + 24n - 12}{2n-1} - 4\sqrt{8n^2 - 16n + 9} \quad (5.11)$$

and

$$LE^+(\Gamma_{V_{8n}}) - E(\Gamma_{V_{8n}}) = \frac{4(n-2)(6n^2 - 6n - 1)}{2n-1} + 4\sqrt{8n^2 - 16n + 9} - 4\sqrt{5n^2 - 6n + 1}. \quad (5.12)$$

Since $8n^3 - 16n^2 + 24n - 12 > 0$, $4(2n-1)\sqrt{8n^2 - 16n + 9} > 0$ and

$$(8n^3 - 16n^2 + 24n - 12)^2 - (4\sqrt{8n^2 - 16n + 9})^2(2n-1)^2 = 64n(n-2)^2(n-1)(n^2 + n - 1) \geq 0$$

(equality holds if and only if $n = 2$) we have $8n^3 - 16n^2 + 24n - 12 - 4(2n-1)\sqrt{8n^2 - 16n + 9} \geq 0$. Therefore, by (5.11), $(2n-1)(LE(\Gamma_{V_{8n}}) - LE^+(\Gamma_{V_{8n}})) \geq 0$. Hence, $LE(\Gamma_{V_{8n}}) \geq LE^+(\Gamma_{V_{8n}})$ equality holds if and only if $G \cong V_{16}$.

Again, $\sqrt{8n^2 - 16n + 9} > 0$, $\sqrt{5n^2 - 6n + 1} > 0$ and $(\sqrt{8n^2 - 16n + 9})^2 - (\sqrt{5n^2 - 6n + 1})^2 = n(3n-10) + 8 \geq 0$ (equality holds if and only if $n = 2$). Therefore, $\sqrt{8n^2 - 16n + 9} - \sqrt{5n^2 - 6n + 1} \geq 0$. Since $4(n-2)(6n^2 - 6n - 1) \geq 0$ we have $\frac{4(n-2)(6n^2 - 6n - 1)}{2n-1} + 4\sqrt{8n^2 - 16n + 9} - 4\sqrt{5n^2 - 6n + 1} \geq 0$. Therefore, by (5.12), $LE^+(\Gamma_{V_{8n}}) \geq E(\Gamma_{V_{8n}})$. Hence, $E(\Gamma_{V_{8n}}) \leq LE^+(\Gamma_{V_{8n}}) \leq LE(\Gamma_{V_{8n}})$ equality holds if and only if $G \cong V_{16}$.

Using Theorems 5.11 and 5.13, for $n \geq 6$, we have

$$LE(\Gamma_{V_{8n}}) - LE^+(\Gamma_{V_{8n}}) = \frac{32n^2 - 40n + 12}{2n - 1} - 4\sqrt{8n^2 - 16n + 9} \quad (5.13)$$

and

$$LE^+(\Gamma_{V_{8n}}) - E(\Gamma_{V_{8n}}) = \frac{32n^3 - 116n^2 + 102n - 14}{2n - 1} + 4\sqrt{8n^2 - 16n + 9} - 4\sqrt{5n^2 - 6n + 1}. \quad (5.14)$$

Since $32n^2 - 40n + 12 > 0$, $4(2n - 1)\sqrt{8n^2 - 16n + 9} > 0$ and $(32n^2 - 40n + 12)^2 - (4\sqrt{8n^2 - 16n + 9})^2 (2n - 1)^2 = 512n^3(n - 2) + 128n(5n - 1) > 0$ we have $32n^2 - 40n + 12 - 4(2n - 1)\sqrt{8n^2 - 16n + 9} > 0$. Therefore, by (5.13), $(2n - 1)(LE(\Gamma_{V_{8n}}) - LE^+(\Gamma_{V_{8n}})) > 0$. Hence, $LE(\Gamma_{V_{8n}}) > LE^+(\Gamma_{V_{8n}})$.

Again, $\sqrt{8n^2 - 16n + 9} > 0$, $\sqrt{5n^2 - 6n + 1} > 0$ and $(\sqrt{8n^2 - 16n + 9})^2 - (\sqrt{5n^2 - 6n + 1})^2 = n(3n - 10) + 8 > 0$. Therefore, $\sqrt{8n^2 - 16n + 9} - \sqrt{5n^2 - 6n + 1} > 0$. Since $32n^3 - 116n^2 + 102n - 14 > 0$ we have $\frac{32n^3 - 116n^2 + 102n - 14}{2n - 1} + 4\sqrt{8n^2 - 16n + 9} - 4\sqrt{5n^2 - 6n + 1} > 0$. Therefore, by (5.14), $LE^+(\Gamma_{V_{8n}}) > E(\Gamma_{V_{8n}})$. Hence, $E(\Gamma_{V_{8n}}) < LE^+(\Gamma_{V_{8n}}) < LE(\Gamma_{V_{8n}})$.

(b) **Case 1:** n is odd

Here $|v(\Gamma_{V_{8n}})| = 8n - 2$ and $E(K_{|v(\Gamma_{V_{8n}})|}) = LE(K_{|v(\Gamma_{V_{8n}})|}) = LE^+(K_{|v(\Gamma_{V_{8n}})|}) = 16n - 16$. Using Theorem 5.11, we have

$$E(\Gamma_{V_{8n}}) - |v(\Gamma_{V_{8n}})| = 2 \left(\sqrt{(2n - 1)(10n - 1)} - 2n \right) \quad (5.15)$$

and

$$E(K_{|v(\Gamma_{V_{8n}})|}) - E(\Gamma_{V_{8n}}) = 2 \left(3(2n - 1) + 1 - \sqrt{(2n - 1)(10n - 1)} \right). \quad (5.16)$$

Since $\sqrt{(2n - 1)(10n - 1)} > 0$, $2n > 0$ and $(\sqrt{(2n - 1)(10n - 1)})^2 - (2n)^2 = 4n(4n - 3) + 1 > 0$ we have $\sqrt{(2n - 1)(10n - 1)} - 2n > 0$. Therefore, by (5.15), $E(\Gamma_{V_{8n}}) > |v(\Gamma_{V_{8n}})|$.

Again, $\sqrt{(2n - 1)(10n - 1)} > 0$, $3(2n - 1) + 1 > 0$ and $(3(2n - 1) + 1)^2 - (\sqrt{(2n - 1)(10n - 1)})^2 = 4n(4n - 3) + 3 > 0$ and so $3(2n - 1) + 1 - \sqrt{(2n - 1)(10n - 1)} > 0$. Therefore, by (5.16), $E(K_{|v(\Gamma_{V_{8n}})|}) > E(\Gamma_{V_{8n}})$.

Case 2: n is even

Here $|v(\Gamma_{V_{8n}})| = 8n - 4$ and $E(K_{|v(\Gamma_{V_{8n}})|}) = LE(K_{|v(\Gamma_{V_{8n}})|}) = LE^+(K_{|v(\Gamma_{V_{8n}})|}) = 16n - 10$. Using Theorem 5.11, we have

$$E(\Gamma_{V_{8n}}) - |v(\Gamma_{V_{8n}})| = 4\sqrt{(n - 1)(5n - 1)} - 4n \quad (5.17)$$

and

$$E(K_{|v(\Gamma_{V_{8n}})|}) - E(\Gamma_{V_{8n}}) = 12n - 6 - 4\sqrt{(n - 1)(5n - 1)}. \quad (5.18)$$

Since $4\sqrt{(n - 1)(5n - 1)} > 0$, $4n > 0$ and $(4\sqrt{(n - 1)(5n - 1)})^2 - (4n)^2 = 16(4n^2 - 6n + 1) > 0$ we have $4\sqrt{(n - 1)(5n - 1)} - 4n > 0$. Therefore, by (5.17), $E(\Gamma_{V_{8n}}) > |v(\Gamma_{V_{8n}})|$.

Again, $4\sqrt{(n - 1)(5n - 1)} > 0$, $12n - 6 > 0$ and $(12n - 6)^2 - (4\sqrt{(n - 1)(5n - 1)})^2 = 4(16n^2 - 12n + 5) > 0$ and so $12n - 6 - 4\sqrt{(n - 1)(5n - 1)} > 0$. Therefore, by (5.18), $E(K_{|v(\Gamma_{V_{8n}})|}) > E(\Gamma_{V_{8n}})$.

(c) **Case 1:** n is odd

Using Theorem 5.12 we have $LE^+(\Gamma_{V_{8n}}) - LE^+(K_{|v(\Gamma_{V_{8n}})|}) = \frac{64n^2(n - 3) + 144n - 22}{4n - 1} + 2\sqrt{32n^2 - 32n + 9} > 0$. Therefore, $LE^+(\Gamma_{V_{8n}}) > LE^+(K_{|v(\Gamma_{V_{8n}})|})$ which implies $\Gamma_{V_{8n}}$ is Q-hyperenergetic and consequently part (a) implies $\Gamma_{V_{8n}}$ is L-hyperenergetic.

Case 2: n is even

Using Theorem 5.13, for $n = 2$, we have $LE(\Gamma_{V_{8n}}) = 16$ and $LE(K_{|v(\Gamma_{V_{8n}})|}) = 22$. Clearly, $LE(\Gamma_{V_{8n}}) < LE(K_{|v(\Gamma_{V_{8n}})|})$. For $n \leq 4$,

$$LE^+(\Gamma_{V_{8n}}) - LE^+(K_{|v(\Gamma_{V_{8n}})|}) = \frac{4(6n^2(n - 4) + 20n - 1)}{2n - 1} + 4\sqrt{8n^2 - 16n + 9} > 0$$

for all $n \neq 2$. Therefore, for all $n \neq 2$, $LE^+(\Gamma_{V_{8n}}) > LE^+(K_{|v(\Gamma_{V_{8n}})|})$ which implies $\Gamma_{V_{8n}}$ is Q-hyperenergetic and consequently part (a) implies $\Gamma_{V_{8n}}$ is L-hyperenergetic. For $n \geq 6$,

$$LE^+(\Gamma_{V_{8n}}) - LE^+(K_{|v(\Gamma_{V_{8n}})|}) = \frac{2(8n^2(2n - 9) + 66n - 11)}{2n - 1} + 4\sqrt{8n^2 - 16n + 9} > 0.$$

Hence, the result holds. \square

In Theorem 5.14, we compare $E(\Gamma_{V_{8n}})$, $LE(\Gamma_{V_{8n}})$ and $LE^+(\Gamma_{V_{8n}})$. However, in the following figures, we show how close are they.




Figure 15: Energies of $\Gamma_{V_{8n}}$, where n is odd




Figure 16: Energies of $\Gamma_{V_{8n}}$, where n is even

5.4 The Frobenious groups of order pq

We consider $F_{p,q} := \langle a, b : a^p = b^q = 1; b^{-1}ab = a^u \rangle$, the Frobenious groups of order pq , where p and q are two primes such that $q|(p-1)$ and u is an integer such that $\bar{u} \in \mathbb{Z}_p \setminus \{\bar{0}\}$ having order q . Results regarding different energies of non-commuting graph of $F_{p,q}$ are given below.

Theorem 5.15 ([14, (4.1.e) and (4.3.b)]). *Let G be isomorphic to $F_{p,q}$. Then*

$$E(\Gamma_G) = \alpha + \sqrt{\alpha^2 + 4p\alpha} \text{ and } LE(\Gamma_G) = \frac{2p^2\alpha + 2p(q-1)^2}{pq-1}, \text{ where } \alpha = (p-1)(q-1).$$

Theorem 5.16. *Let G be isomorphic to $F_{p,q}$. Then*

$$\text{Q-spec}(\Gamma_G) = \left\{ (pq-p)^{p-2}, (pq-q)^{pq-2p}, (pq-2q+1)^{p-1}, \left(\frac{A}{2}\right)^1, \left(\frac{B}{2}\right)^1 \right\},$$

where $A = 3pq - 2p - 2q + 1 + \sqrt{pq(pq-2) + 4(p-q)(pq-p-q+1) + 1}$ and $B = 3pq - 2p - 2q + 1 - \sqrt{pq(pq-2) + 4(p-q)(pq-p-q+1) + 1}$ and

$$LE^+(\Gamma_G) = \frac{2p^3q - p^2q^2 - 2pq^2 - 6pq - 4p^3 + 6p^2 + 2q - 1}{pq-1} + \sqrt{pq(pq-2) + 4(p-q)(pq-p-q+1) + 1}.$$

Proof. If $G \cong F_{p,q}$ then $|v(\Gamma_G)| = pq-1$ and $\Gamma_G = K_{1,(p-1),p,(q-1)}$. Using Theorem 1.6(b), we have

$$\begin{aligned} Q_{\Gamma_G}(x) &= \prod_{i=1}^2 (x - (pq-1) + p_i)^{a_i(p_i-1)} \prod_{i=1}^2 (x - (pq-1) + 2p_i)^{a_i} \left(1 - \sum_{i=1}^2 \frac{a_i p_i}{x - (pq-1) + 2p_i} \right) \\ &= (x - pq + p)^{p-2} (x - pq + q)^{pq-2p} (x - pq + 2p - 1)(x - pq + 2q - 1)^p \\ &\quad \times \left(1 - \frac{p-1}{x - pq + 2p - 1} - \frac{pq-p}{x - pq + 2q - 1} \right) \\ &= (x - (pq-p))^{p-2} (x - (pq-q))^{pq-2p} (x - (pq-2q+1))^p \\ &\quad \times (x^2 - (3pq - 2p - 2q + 1)x + 2p^2q^2 - 4p^2q + 2pq - 2pq^2 + 2p^2 - 2p + 2q - 1). \end{aligned}$$

Thus, $\text{Q-spec}(\Gamma_G) = \left\{ (pq-p)^{p-2}, (pq-q)^{pq-2p}, (pq-2q+1)^{p-1}, \left(\frac{A}{2}\right)^1, \left(\frac{B}{2}\right)^1 \right\}$.

Number of edges of Γ_G^c is $\frac{p^2-3p+2+pq^2-3pq+2p}{2}$. Thus, $|e(\Gamma_G)| = \frac{(pq-1)(pq-2)}{2} - \frac{p^2-3p+2+pq^2-3pq+2p}{2} = \frac{(p^2-p)(q^1-1)}{2}$. Now,

$$\begin{aligned} \left| pq - p - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| &= \left| \frac{(q-1)(pq-p^2)}{pq-1} \right| = \frac{(q-1)(p^2-pq)}{pq-1}, \\ \left| pq - q - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| &= \left| \frac{(p-q)(p-1)}{pq-1} \right| = \frac{(p-q)(p-1)}{pq-1}, \\ \left| pq - 2q + 1 - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| &= \left| \frac{(p^2+2q) - (pq^2+p+1)}{pq-1} \right| = \frac{-(p^2+2q) + (pq^2+p+1)}{pq-1}, \end{aligned}$$

$$\left| A - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{p^2q^2 + 2p^2 + p + 2q - 2p^2q - 2pq - 1}{2(pq - 1)} + \frac{C}{2} \right| = \frac{p^2q^2 + 2p^2 + p + 2q - 2p^2q - 2pq - 1}{2(pq - 1)} + \frac{C}{2}$$

and

$$\left| B - \frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} \right| = \left| \frac{p^2q^2 + 2p^2 + p + 2q - 2p^2q - 2pq - 1}{2(pq - 1)} - \frac{C}{2} \right| = \frac{-(p^2q^2 + 2p^2 + p + 2q - 2p^2q - 2pq - 1)}{2(pq - 1)} + \frac{C}{2},$$

where $C = \sqrt{pq(pq - 2) + 4(p - q)(pq - p - q + 1) + 1}$. Therefore,

$$LE^+(\Gamma_G) = (p - 2) \times \frac{(q - 1)(p^2 - pq)}{pq - 1} + (pq - 2p) \times \frac{(p - q)(p - 1)}{pq - 1} + (p - 1) \times \frac{(pq^2 + p + 1) - (p^2 + 2q)}{pq - 1} + \frac{p^2q^2 + 2p^2 + p + 2q - 2p^2q - 2pq - 1}{2(pq - 1)} + \frac{C}{2} - \left(\frac{p^2q^2 + 2p^2 + p + 2q - 2p^2q - 2pq - 1}{2(pq - 1)} \right) + \frac{C}{2}$$

and the result follows on simplification. \square

6 Some implications of the preceding findings

It was shown in [8] that the non-commuting graphs of the groups considered in this paper are L-integral. Also, in [14, Chapter 4], several conditions were obtained such that the non-commuting graphs of these groups are integral. In view of Theorems 2.16 and 4.2, it follows that Γ_G is not Q-integral if $G \cong U_{6n}$ or $\frac{G}{Z(G)} \cong Sz(2)$. However, Γ_G is Q-integral if $G \cong A(n, \mathcal{V})$, $A(n, p)$ or $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$ (follows from Theorems 3.2, 5.2 and 5.5). As a consequence of our results we also have the following theorem related to Question 1.1.

Theorem 6.1. Γ_G is Q-integral if

- (a) $G \cong D_{2r}, M_{2rs}$ or SD_{8r} , r is odd and $8r^2 - 16r + 9$ is a perfect square.
- (b) $G \cong V_{8n}$, n is even and $8n^2 - 16n + 9$ is a perfect square.
- (c) $G \cong Q_{4n}$ or $\frac{G}{Z(G)} \cong D_{2n}$ and $8n^2 - 16n + 9$ is a perfect square.
- (d) $G \cong D_{2r}$ or M_{2rs} , r is even and $2r^2 - 8r + 9$ is a perfect square.
- (e) $G \cong V_{8n}$, n is odd and $32n^2 - 32n + 9$ is a perfect square.
- (f) $G \cong SD_{8n}$, n is even and $32n^2 - 32n + 9$ is a perfect square.
- (g) $G \cong QD_{2n}$ and $2^{2n-1} - 2^{n+2} + 9$ is a perfect square.

In the following table we give some positive integers such that $8n^2 - 16n + 9$, $2n^2 - 8n + 9$ and $32n^2 - 32n + 9$ are perfect squares. It may be interesting to obtain general terms of such sequences of positive integers.

n	$\sqrt{8n^2 - 16n + 9}$	n	$\sqrt{2n^2 - 8n + 9}$	n	$\sqrt{32n^2 - 32n + 9}$
1	1	2	1	1	3
2	3	4	3	18	99
7	17	14	17	595	3363
36	99	72	99	20196	114243
205	577	410	577	686053	3880899
1190	3363	2380	3363	23305590	131836323
6931	19601	13862	19601		
40392	114243	80784	114243		
235417	665857	470834	665857		
1372106	3880899	2744212	3880899		
7997215	22619537	15994430	22619537		
46611180	131836323	93222360	131836323		
271669861	768398401	543339722	768398401		

Table 1

As consequences of our results we also have the following theorems related to Questions 1.2 and 1.3.

Theorem 6.2. *Let G be a finite non-abelian group. Then*

- (a) $E(\Gamma_G) = LE(\Gamma_G) = LE^+(\Gamma_G)$ if $G \cong D_8, Q_8, M_{8s}, A(n, \mathcal{V}), A(n, p), V_{16}$ a non-abelian group of order p^3 . Also, if $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$, where p is a prime, then $E(\Gamma_G) = LE(\Gamma_G) = LE^+(\Gamma_G)$.
- (b) $E(\Gamma_G) < LE^+(\Gamma_G) < LE(\Gamma_G)$ if $G \cong D_{2m}(m \neq 4), QD_{2^n}, M_{2rs}(r \neq 4), Q_{4n}(n \neq 2), U_{6n}, SD_{8n}$ and $V_{8n}(n \neq 2)$. Also, if $\frac{G}{Z(G)} \cong D_{2m}(m \geq 3)$ and $Sz(2)$ then $E(\Gamma_G) < LE^+(\Gamma_G) < LE(\Gamma_G)$.
- (c) Γ_G is non-hypoenergetic as well as non-hyperenergetic if $G \cong D_{2m}, QD_{2^n}, M_{2rs}, Q_{4n}, U_{6n}, A(n, \mathcal{V}), A(n, p), SD_{8n}$ and V_{8n} . Also, if $\frac{G}{Z(G)} \cong D_{2m}, \mathbb{Z}_p \times \mathbb{Z}_p$ and $Sz(2)$, where $m \geq 3$ and p is a prime, then Γ_G is non-hypoenergetic as well as non-hyperenergetic.
- (d) Γ_G is L-hyperenergetic but not Q-hyperenergetic if $G \cong D_6, M_6$ and $Sz(2)$.
- (e) Γ_G is neither L-hyperenergetic nor Q-hyperenergetic if $G \cong D_8, M_{8s}, Q_8, A(n, \mathcal{V}), A(n, p), V_{16}$. Also, if $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$ then Γ_G is neither L-hyperenergetic nor Q-hyperenergetic.
- (f) Γ_G is L-hyperenergetic as well as Q-hyperenergetic if $G \cong D_{2m}(m \neq 3, 4), QD_{2^n}, M_{2rs}(2rs \neq 6, 8s), Q_{4n}(n \neq 2), U_{6n}, SD_{8n}$ and $V_{8n}(n \neq 2)$. Also, if $\frac{G}{Z(G)} \cong Sz(2)$ ($G \not\cong Sz(2)$) and D_{2m} ($m = 3, 4$ and $|Z(G)| \neq 1$ or $m \geq 5$ and $|Z(G)| \geq 1$) then Γ_G is L-hyperenergetic as well as Q-hyperenergetic.

In the following theorem we get an example of a graph (non-commuting graph of the symmetric group of degree 4) disproving Conjecture 1.4.

Theorem 6.3. *Let the commuting graph of a finite group G be planar. Then*

- (a) Γ_G is Q-integral if $G \cong D_8, Q_8, \mathbb{Z}_2 \times D_8, \mathbb{Z}_2 \times Q_8, \mathcal{M}_{16}, \mathbb{Z}_4 \rtimes \mathbb{Z}_4, D_8 * \mathbb{Z}_4$ or $SG(16, 3)$, otherwise not Q-integral.
- (b) Γ_G is non-hypoenergetic.
- (c) Γ_G is hyperenergetic if $G \cong S_4$, otherwise non-hyperenergetic.
- (d) Γ_G is Q-hyperenergetic if $G \cong D_{10}, D_{12}, Q_{12}, A_4, A_5, S_4$ or $SL(2, 3)$, otherwise not Q-hyperenergetic
- (e) Γ_G is L-hyperenergetic if $G \cong D_6, D_{10}, D_{12}, Q_{12}, A_4, A_5, S_4, SL(2, 3)$ or $Sz(2)$, otherwise not L-hyperenergetic.

Proof. If the commuting graph of G is planar then, by [3, Theorem 2.2], $G \cong D_6, D_8, D_{10}, D_{12}, Q_8, Q_{12}, \mathbb{Z}_2 \times D_8, \mathbb{Z}_2 \times Q_8, \mathcal{M}_{16}, \mathbb{Z}_4 \rtimes \mathbb{Z}_4, D_8 * \mathbb{Z}_4, SG(16, 3), A_4, A_5, S_4, SL(2, 3), Sz(2)$.

By Theorems 2.2, 2.3 and 2.13, Γ_G is Q-integral if $G \cong D_8, Q_8$ and not Q-integral if $G \cong D_6, D_{10}, D_{12}, Q_{12}$. If $G \cong D_6$, then from Theorem 2.4, Γ_G is non-hypoenergetic, non-hyperenergetic, not Q-hyperenergetic but is L-hyperenergetic. If $G \cong D_{10}, D_{12}$, then from Theorem 2.4, Γ_G is non-hypoenergetic, non-hyperenergetic but is Q-hyperenergetic and L-hyperenergetic. If $G \cong D_8$, then from Theorem 2.4, Γ_G is non-hypoenergetic, non-hyperenergetic, not Q-hyperenergetic and not L-hyperenergetic.

If $G \cong Q_8$, then from Theorem 2.14, Γ_G is non-hypoenergetic, non-hyperenergetic, not Q-hyperenergetic and not L-hyperenergetic. If $G \cong Q_{12}$, then from Theorem 2.14 Γ_G is non-hypoenergetic, non-hyperenergetic but is Q-hyperenergetic and L-hyperenergetic.

If $G \cong \mathbb{Z}_2 \times D_8, \mathbb{Z}_2 \times Q_8, \mathcal{M}_{16}, \mathbb{Z}_4 \rtimes \mathbb{Z}_4, D_8 * \mathbb{Z}_4, SG(16, 3)$, then $\frac{G}{Z(G)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. Using Theorems 3.2 and 3.3, for $p = 2$, we get Γ_G is Q-integral but not hypoenergetic, hyperenergetic, Q-hyperenergetic as well as L-hyperenergetic.

If $G \cong A_4$ then from Proposition 4.3.13 of [14], we have $E(\Gamma_{A_4}) = 6 + 2\sqrt{33}$ and $LE(\Gamma_{A_4}) = \frac{224}{11}$. Now, $|v(\Gamma_{A_4})| = 11$ so $E(K_{|v(\Gamma_{A_4})|}) = LE^+(K_{|v(\Gamma_{A_4})|}) = LE(K_{|v(\Gamma_{A_4})|}) = 20$. Here, $\Gamma_{A_4} = K_{4.2.1.3}$ so using Theorem 1.6(b) we get

$$\text{Q-spec}(\Gamma_{A_4}) = \left\{ (9)^4, (8)^2, (7)^3, \left(\frac{23 + \sqrt{145}}{2} \right)^1, \left(\frac{23 - \sqrt{145}}{2} \right)^1 \right\}.$$

It follows that Γ_{A_4} is not Q-integral. We have $|e(\Gamma_{A_4})| = 48$ and so $\frac{2|e(\Gamma_{A_4})|}{|v(\Gamma_{A_4})|} = \frac{96}{11}$. Therefore, $|9 - \frac{96}{11}| = \frac{3}{11}$, $|8 - \frac{96}{11}| = \frac{8}{11}$, $|7 - \frac{96}{11}| = \frac{19}{11}$, $\left| \frac{23 + \sqrt{145}}{2} - \frac{96}{11} \right| = \frac{61}{22} + \frac{\sqrt{145}}{2}$ and $\left| \frac{23 - \sqrt{145}}{2} - \frac{96}{11} \right| = -\frac{61}{22} + \frac{\sqrt{145}}{2}$. Thus,

$$LE^+(\Gamma_{A_4}) = 4 \times \frac{3}{11} + 2 \times \frac{8}{11} + 3 \times \frac{19}{11} + \frac{61}{22} + \frac{\sqrt{145}}{2} - \frac{61}{22} + \frac{\sqrt{145}}{2} = \frac{85}{11} + \sqrt{145} > 20.$$

Hence, Γ_{A_4} is non-hypoenergetic, non-hyperenergetic but is Q-hyperenergetic and L-hyperenergetic.

If $G \cong A_5$ then from Proposition 4.3.13 of [14], we have $E(\Gamma_{A_5}) = 111.89$ and $LE(\Gamma_{A_5}) = \frac{8580}{59}$. Now, $|v(\Gamma_{A_5})| = 59$ so $E(K_{|v(\Gamma_{A_5})|}) = LE^+(K_{|v(\Gamma_{A_5})|}) = LE(K_{|v(\Gamma_{A_5})|}) = 116$. Here, $\Gamma_{A_5} = K_{5,3,10,2,6,4}$ so using Theorem 1.6(b) we get

$$\text{Q-spec}(\Gamma_{A_5}) = \{(56)^{10}, (57)^{10}, (55)^{27}, (53)^4, (51)^5, (x_1)^1, (x_2)^1, (x_3)^1\},$$

where $x_1 \approx 52.03252, x_2 \approx 54.05266$ and $x_3 \approx 111.91482$ are the roots of the equation $x^3 - 218x^2 + 14685x - 314760 = 0$. It follows that Γ_{A_5} is not Q-integral. We have $|e(\Gamma_{A_5})| = 1650$ and so $\frac{2|e(\Gamma_{A_5})|}{|v(\Gamma_{A_5})|} = \frac{3300}{59}$. Therefore, $|57 - \frac{3300}{59}| = \frac{63}{59}, |56 - \frac{3300}{59}| = \frac{4}{59}, |55 - \frac{3300}{59}| = \frac{55}{59}, |53 - \frac{3300}{59}| = \frac{173}{59}, |51 - \frac{3300}{59}| = -\frac{291}{59} + \frac{\sqrt{145}}{2}, |x_1 - \frac{3300}{59}| = -(x_1 - \frac{3300}{59}), |x_2 - \frac{3300}{59}| = -(x_2 - \frac{3300}{59})$ and $|x_3 - \frac{3300}{59}| = x_3 - \frac{3300}{59}$. Thus,

$$LE^+(\Gamma_{A_5}) = \frac{7602}{59} - x_1 - x_2 + x_3 > 116.$$

Hence, Γ_{A_5} is non-hypoenergetic, non-hyperenergetic but is Q-hyperenergetic and L-hyperenergetic.

If $G \cong S_4$ then from Proposition 4.3.13 of [14], we have $E(\Gamma_{S_4}) = 35.866 + 4\sqrt{5}$ and $LE(\Gamma_{S_4}) = \frac{1072}{23} + 4\sqrt{13}$. Now, $|v(\Gamma_{S_4})| = 23$ so $E(K_{|v(\Gamma_{S_4})|}) = LE^+(K_{|v(\Gamma_{S_4})|}) = LE(K_{|v(\Gamma_{S_4})|}) = 44$. Using GAP [34], the characteristic polynomial of $Q(\Gamma_{S_4})$ is

$$Q_{\Gamma_{S_4}}(x) = x(x+20)^4(x+21)^7(x+23)^7(x^2+40x+394)^2$$

and so

$$\text{Q-spec}(\Gamma_{S_4}) = \left\{ (0)^1, (-20)^4, (-21)^7, (-23)^7, \left(-20 + \sqrt{6}\right)^2, \left(-20 - \sqrt{6}\right)^2 \right\}.$$

It follows that Γ_{S_4} is not Q-integral. We have $|e(\Gamma_{S_4})| = 228$ and so $\frac{2|e(\Gamma_{S_4})|}{|v(\Gamma_{S_4})|} = \frac{456}{23}$. Therefore, $|0 - \frac{456}{23}| = \frac{456}{23}, |20 - \frac{456}{23}| = \frac{4}{23}, |21 - \frac{456}{23}| = \frac{27}{23}, |23 - \frac{456}{23}| = \frac{73}{23}, |-20 + \sqrt{6} - \frac{456}{23}| = \frac{916}{23} - \sqrt{6}$ and $|-20 - \sqrt{6} - \frac{456}{23}| = \frac{916}{23} + \sqrt{6}$. Thus,

$$LE^+(\Gamma_{S_4}) = \frac{456}{23} + 4 \times \frac{4}{23} + 7 \times \frac{27}{23} + 7 \times \frac{73}{23} + 2 \times \left(\frac{916}{23} - \sqrt{6}\right) + 2 \times \left(\frac{916}{23} + \sqrt{6}\right) = \frac{4836}{23}.$$

Hence, Γ_{S_4} is non-hypoenergetic, hyperenergetic but is Q-hyperenergetic and L-hyperenergetic.

If $G \cong SL(2, 3)$ then from Proposition 4.3.13 of [14], we have $E(\Gamma_{SL(2,3)}) = 16 + 8\sqrt{7}$ and $LE(\Gamma_{SL(2,3)}) = \frac{552}{11}$. Now, $|v(\Gamma_{SL(2,3)})| = 22$ so $E(K_{|v(\Gamma_{SL(2,3)})|}) = LE^+(K_{|v(\Gamma_{SL(2,3)})|}) = LE(K_{|v(\Gamma_{SL(2,3)})|}) = 42$. Here, $\Gamma_{SL(2,3)} = K_{3,2,4,4}$ so using Theorem 1.6(b) we get

$$\text{Q-spec}(\Gamma_{SL(2,3)}) = \left\{ (20)^3, (18)^{14}, (14)^3, \left(\frac{54 + \sqrt{420}}{2}\right)^1, \left(\frac{54 - \sqrt{420}}{2}\right)^1 \right\}.$$

It follows that $\Gamma_{SL(2,3)}$ is not Q-integral. We have $|e(\Gamma_{SL(2,3)})| = 204$ and so $\frac{2|e(\Gamma_{SL(2,3)})|}{|v(\Gamma_{SL(2,3)})|} = \frac{204}{11}$. Therefore, $|20 - \frac{204}{11}| = \frac{16}{11}, |18 - \frac{204}{11}| = \frac{6}{11}, |14 - \frac{204}{11}| = \frac{50}{11}, \left|\frac{54 + \sqrt{420}}{2} - \frac{204}{11}\right| = \frac{93}{11} + \frac{\sqrt{420}}{2}$ and $\left|\frac{54 - \sqrt{420}}{2} - \frac{204}{11}\right| = -\frac{93}{11} + \frac{\sqrt{420}}{2}$. Thus,

$$LE^+(\Gamma_{SL(2,3)}) = 3 \times \frac{16}{11} + 14 \times \frac{6}{11} + 3 \times \frac{50}{11} + \frac{93}{11} + \frac{\sqrt{420}}{2} - \frac{93}{11} + \frac{\sqrt{420}}{2} = \frac{282}{11} + \sqrt{420}.$$

Hence, $\Gamma_{SL(2,3)}$ is non-hypoenergetic, non-hyperenergetic but is Q-hyperenergetic and L-hyperenergetic.

If $G \cong Sz(2)$ then, by Theorem 4.2, we have Γ_G is not Q-integral. Also, Theorem 4.3 gives that Γ_G is non-hypoenergetic, non-hyperenergetic, not Q-hyperenergetic but is L-hyperenergetic. \square

Theorem 6.4. *Let G be a finite group and the commuting graph of G is toroidal. Then*

- (a) Γ_G is Q-integral if $G \cong D_{14}$ or $A_4 \times \mathbb{Z}_2$, otherwise not Q-integral.
- (b) Γ_G is non-hypoenergetic, non-hyperenergetic but is Q-hyperenergetic and L-hyperenergetic.

Proof. If commuting graph of G is toroidal then, by [9, Theorem 3.3] $G \cong D_{14}, D_{16}, QD_{16}, \mathbb{Z}_7 \rtimes \mathbb{Z}_3, D_6 \times \mathbb{Z}_3, A_4 \times \mathbb{Z}_2$. By Theorems 2.2, 2.6 and 2.13, Γ_G is Q-integral if $G \cong D_{14}$ and not Q-integral if $G \cong D_{16}, QD_{16}$ or QD_{16} . If $G \cong D_{14}, D_{16}$, then from Theorem 2.4, Γ_G is non-hypoenergetic, non-hyperenergetic but is Q-hyperenergetic and L-hyperenergetic. If $G \cong Q_{16}$, then from Theorem 2.14, Γ_G is non-hypoenergetic, non-hyperenergetic but is Q-hyperenergetic and L-hyperenergetic. If $G \cong QD_{16}$, then from Theorem 2.7, Γ_G is non-hypoenergetic, non-hyperenergetic but is Q-hyperenergetic and L-hyperenergetic.

If $G \cong \mathbb{Z}_7 \rtimes \mathbb{Z}_3$ then, by Theorem 5.16, we have

$$\text{Q-spec}(\Gamma_G) = \left\{ (14)^5, (18)^7, (16)^6, \left(22 + 2\sqrt{37}\right)^1, \left(22 - 2\sqrt{37}\right)^1 \right\}.$$

Thus Γ_G is not Q-integral. By Theorem 5.15, we also have $E(\Gamma_G) = 12 + 4\sqrt{30}$ and $LE(\Gamma_G) = \frac{308}{5}$ and from Theorem 5.16, we have $LE^+(\Gamma_G) = \frac{292}{20} + 4\sqrt{37}$. Now, $|v(\Gamma_G)| = 20$ so $E(K_{|v(\Gamma_G)|}) = LE^+(K_{|v(\Gamma_G)|}) = LE(K_{|v(\Gamma_G)|}) = 38$. Hence, Γ_G is non-hypoenergetic, non-hyperenergetic but is Q-hyperenergetic and L-hyperenergetic.

If $G \cong D_6 \times \mathbb{Z}_3$, then from Proposition 4.3.14 of [14], we have $E(\Gamma_G) = 6 + 6\sqrt{7}$ and $LE(\Gamma_G) = \frac{594}{15}$. Now, $|v(\Gamma_G)| = 15$ so $E(K_{|v(\Gamma_G)|}) = LE^+(K_{|v(\Gamma_G)|}) = LE(K_{|v(\Gamma_G)|}) = 28$. Here, $\Gamma_G = K_{3,3,1.6}$ so using Theorem 1.6(b) we get

$$\text{Q-spec}(\Gamma_G) = \left\{ (12)^6, (9)^7, \left(\frac{27 + \sqrt{297}}{2}\right)^1, \left(\frac{27 - \sqrt{297}}{2}\right)^1 \right\}.$$

It follows that Γ_G is not Q-integral. We have $|e(\Gamma_G)| = 81$ and so $\frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} = \frac{162}{15}$. Therefore, $|12 - \frac{162}{15}| = \frac{18}{15}$, $|9 - \frac{162}{15}| = \frac{27}{15}$, $\left|\frac{27 + \sqrt{297}}{2} - \frac{162}{15}\right| = \frac{81}{30} + \frac{\sqrt{297}}{2}$ and $\left|\frac{27 - \sqrt{145}}{2} - \frac{162}{15}\right| = -\frac{81}{30} + \frac{\sqrt{297}}{2}$. Thus,

$$LE^+(\Gamma_G) = 6 \times \frac{18}{15} + 7 \times \frac{189}{11} + \frac{81}{30} + \frac{\sqrt{297}}{2} - \frac{81}{30} + \frac{\sqrt{297}}{2} = \frac{99}{5} + 3\sqrt{33}.$$

Hence, Γ_G is non-hypoenergetic, non-hyperenergetic but is Q-hyperenergetic and L-hyperenergetic.

If $G \cong A_4 \times \mathbb{Z}_2$, then from Proposition 4.3.14 of [14], we have $E(\Gamma_G) = 12 + 4\sqrt{33}$ and $LE(\Gamma_G) = \frac{544}{11}$. Now, $|v(\Gamma_G)| = 22$ so $E(K_{|v(\Gamma_G)|}) = LE^+(K_{|v(\Gamma_G)|}) = LE(K_{|v(\Gamma_G)|}) = 42$. Here, $\Gamma_G = K_{4,4,1.6}$ so using Theorem 1.6(b) we get

$$\text{Q-spec}(\Gamma_G) = \{(18)^{12}, (16)^5, (14)^3, (36)^1, (10)^1\}.$$

Clearly, Γ_G is Q-integral. We have $|e(\Gamma_G)| = 192$ and so $\frac{2|e(\Gamma_G)|}{|v(\Gamma_G)|} = \frac{192}{11}$. Therefore, $|18 - \frac{192}{11}| = \frac{6}{11}$, $|16 - \frac{192}{11}| = \frac{16}{11}$, $|14 - \frac{192}{11}| = \frac{38}{11}$, $|36 - \frac{192}{11}| = \frac{204}{11}$ and $|10 - \frac{192}{11}| = \frac{82}{11}$. Thus,

$$LE^+(\Gamma_G) = 12 \times \frac{6}{11} + 5 \times \frac{16}{11} + 3 \times \frac{38}{11} + \frac{204}{11} + \frac{82}{11} = \frac{552}{11}.$$

Hence, Γ_G is non-hypoenergetic, non-hyperenergetic but is Q-hyperenergetic and L-hyperenergetic. \square

If non-commuting graph of G is planar then, by [1, Proposition 2.3] $G \cong D_6, D_8, Q_8$. Therefore, we have the following theorem.

Theorem 6.5. *Let G be a finite group whose non-commuting graph is planar. Then*

- (a) Γ_G is non-hypoenergetic, non-hyperenergetic and not Q-hyperenergetic.
- (b) Γ_G is not L-hyperenergetic but Q-integral if $G \not\cong D_6$.

Acknowledgements

The authors would like to thank Mr. Nabin Pokhrel and Mr. Kallol Ray for their help in drawing Figures and computing perfect squares in Table 1. Ms. M. Sharma expresses gratitude to DST for the INSPIRE fellowship.

Funding

No funding was received by the authors.

Data Availability

No data was used in the preparation of this manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- [1] Abdollahi, A., Akbari, S. and Maimani, H. R. Non-commuting graph of a group, *Journal of Algebra*, **298**, 468–492, 2006.
- [2] Abdussakir, Elvierayani, R. R. and Nafisah, M. On the spectra of commuting and non-commuting graph on dihedral group, *Cauchy-Jurnal Matematika Murni dan Aplikasi*, **4**(4), 176–182, 2017.
- [3] Afkhami, M., Farrokhi, D. G. M. and Khashyarmanesh, K. Planar, toroidal, and projective commuting and non-commuting graphs, *Communications in Algebra*, **43**(7), 2964–2970, 2015.
- [4] Barati, Z., Erfanian, A., Khashyarmanesh, K. and Nafar, K. A generalization of non-commuting graph via automorphisms of a group, *Communications in Algebra*, **42**, 174–185, 2014.
- [5] Bhowal, P. and Nath, R. K. Spectral aspects of commuting conjugacy class graph of finite groups, *Algebraic Structures and Their Applications*, **8**(2), 67–118, 2021.
- [6] Darafshah, M. R., Bigdely, H., Bahrami, A. and Monfared, M. D. Some results on non-commuting graph of a finite group, *Italian Journal of Pure and Applied Mathematics*, **27**, 107–118, 2010.
- [7] Dutta, P., Bagchi, B. and Nath, R. K. Various energies of commuting graphs of finite nonabelian groups, *Khayyam Journal of Mathematics*, **6**(1), 27–45, 2020.
- [8] Dutta, P., Dutta, J. and Nath, R. K. Laplacian spectrum of noncommuting graphs of finite groups, *Indian Journal of Pure and Applied Mathematics*, **49**(2), 205–216, 2018.
- [9] Dutta, J. and Nath, R. K. Spectrum of commuting graphs of some classes of finite groups, *Matematika*, **33**(1), 87–95, 2017.
- [10] Dutta, J. and Nath, R. K. Finite groups whose commuting graphs are integral, *Matematički Vesnik*, **69**(3), 226–230, 2017.
- [11] Dutta, J. and Nath, R. K. Laplacian and signless Laplacian spectrum of commuting graphs of finite groups, *Khayyam Journal of Mathematics*, **4**(1), 77–87, 2018.
- [12] Dutta, P. and Nath, R. K. On Laplacian energy of non-commuting graphs of finite groups, *Journal of Linear and Topological Algebra*, **7**(2), 121–132, 2018.
- [13] Dutta, P. and Nath, R. K. Various energies of commuting graphs of some super integral groups, *Indian Journal of Pure and Applied Mathematics*, **52**(1), 1–10, 2021.
- [14] Fafous, W. N. T., *Various spectra and energies of commuting graphs of finite groups and rings*, PhD thesis, Department of Mathematical Sciences, Tezpur University, Assam, 2021.
- [15] Fafous, W. N. T., Nath, R.K. and Sharafadini, R. Various spectra and energies of commuting graphs of finite rings, *Haceteppe Journal of Mathematics and Statistics*, **49**(6), 1915–1925, 2020.
- [16] Ghayekhloo, S., Erfanian, A. and Tolue, B. The generalised non-commuting graph of a finite group, *Proceedings of the Bulgarian Academy of Sciences*, **67**, 1037–1044, 2014.
- [17] Ghorbani, M. and Gharavi-Alkhansari, Z. On the energy of non-commuting graphs, *Journal of Linear and Topological Algebra*, **6**(2), 135–146, 2017.
- [18] Ghorbani, M., Gharavi-Alkhansari, Z. and Bashi, A. Z. On the eigenvalues of non-commuting graphs, *Journal of Algebraic Structures and Their Applications*, **4**(2), 27–38, 2017.
- [19] Gutman, I. The energy of a graph, *Ber. Math-Statist. Sect. Forschungsz. Graz*, **103**, 1–22, 1978, 1–22.
- [20] Gutman, I. Hyperenergetic molecular graphs, *Journal of the Serbian Chemical Society*, **64**, 199–205, 1999.
- [21] Gutman, I. Hyperenergetic and Hypoenergetic Graphs, *Zbornik Radova* **22**, 113–135, 2011.
- [22] Gutman, I., Abreu, N. M. M., Vinagre, C. T. M., Bonifácio, A. S. and Radenković, S. Relation between energy and Laplacian energy, *MATCH Communications in Mathematical and in Computer Chemistry*, **59**, 343–354, 2008.
- [23] Gutman, I. and Radenković, S. Hypoenergetic molecular graphs, *Indian Journal of Chemistry*, **46**(A), 1733–1736, 2007.
- [24] Liu, J. and Liu, B. On relation between energy and Laplacian energy, *MATCH Communications in Mathematical and in Computer Chemistry*, **61**, 403–406, 2009.

- [25] Moghaddamfar, A. R., Shi, W. J., Zhou, W. and Zokayi, A. R. On the non-commuting graph associated with a finite group, *Siberian Mathematical Journal*, **46**(2), 325–332, 2005.
- [26] Nath, R. K. et al. Various spectra and energies of non-commuting conjugacy class graphs of finite groups, *Preprint*.
- [27] Neumann, B. H. A problem of Paul Erdős on groups, *Journal of Australian Mathematical Society*, **21**, 467–472, 1976.
- [28] Sharafdini, R., Nath, R. K. and Darbandi, R. Energy of commuting graph of finite AC-groups, *Proyecciones Journal of Mathematics*, **41**(1), 263–273, 2022.
- [29] Stevanovic, D., Stankovic, I. and Milosević, M. More on the relation between energy and Laplacian energy of graphs, *MATCH Communications in Mathematical and in Computer Chemistry*, **61**, 395–401, 2009.
- [30] Tolve, B., Erfanian, A. and Jafarzadeh, A. A kind of non-commuting graph of finite groups, *Journal of Science Islamic Republic of Iran*, **25**(4), 379–384, 2014.
- [31] Walikar, H. B., Ramane, H. S. and Hampiholi, P. R. On the energy of a graph, *Graph Connections, Allied Publishers, New Delhi*, 120–123, 1999.
- [32] Wang, L., Li, X. and Hoede, C. Integral complete r -partite graphs, *Discrete Mathematics*, **283**, 231–241, 2004.
- [33] Zhao, G., Wang, L. and Li, K. Q -integral complete r -partite graphs, *Linear Algebra and its Applications*, **438**, 1067–1077, 2013.
- [34] The GAP Group, GAP – Groups, Algorithms, and Programming, Version 4.12.2; 2022. (<http://www.gap-system.org>).