PROBING THE SOLAR INTERIOR WITH LENSED GRAVITATIONAL WAVES FROM KNOWN PULSARS

Ryuichi Takahashi¹, Soichiro Morisaki² and Teruaki Suyama³

¹ Faculty of Science and Technology, Hirosaki University, 3 Bunkyo-cho, Hirosaki, Aomori 036-8561, Japan

² Institute for Cosmic Ray Research (ICRR), KAGRA Observatory, The University of Tokyo, Kashiwa City, Chiba 277-8582, Japan

³ Department of Physics, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8551, Japan

Draft version October 9, 2023

ABSTRACT

When gravitational waves (GWs) from a spinning neutron star arrive from behind the Sun, they are subjected to gravitational lensing that imprints a frequency-dependent modulation on the waveform. This modulation traces the projected solar density and gravitational potential along the path as the Sun passes in front of the neutron star. We calculate how accurately the solar density profile can be extracted from the lensed GWs using a Fisher analysis. For this purpose, we selected three promising candidates (the highly spinning pulsars J1022+1001, J1730-2304, and J1745-23) from the pulsar catalog of the Australia Telescope National Facility. The lensing signature can be measured with 3σ confidence when the signal-to-noise ratio (SNR) of the GW detection reaches 100 (f/300Hz)⁻¹ over a one-year observation period (where f is the GW frequency). The solar density profile can be plotted as a function of radius when the SNR improves to $\gtrsim 10^4$.

Subject headings: gravitational lensing: weak — gravitational waves — Sun: general

1. INTRODUCTION

the direct detection of a gravitational-Since wave (GW) signal from a merging black hole binary (GW150914; Abbott et al. 2016), GW astronomy has attracted increasing interest. Currently, all GW sources detected by the ground-based detectors of the Laser Interferometer Gravitational-Wave Observatory (LIGO), the Virgo interferometer, and the Kamioka Gravitational Wave Detector (KAGRA) are compact binary coalescences (CBCs) of stellar-mass black holes and neutron stars. The LIGO-Virgo-KAGRA (LVK) collaboration reported ~ 90 candidates of CBC events (The LIGO Scientific Collaboration et al. 2021). However, the ground-based detectors are also expected to detect spinning neutron stars (e.g., Abbott et al. 2022a). A neutron star that is non-axisymmetric around its spin axis, resulting in a so-called mountainous profile, emits continuous GWs. Non-axisymmetry may be caused by crustal deformation, magnetic fields, and mass accretion from the star's companion (e.g., see recent reviews by Glampedakis & Gualtieri (2018) and Riles (2022)). Although the LVK collaboration has been searching for continuous GW signals, no event has yet been reported (Abbott et al. 2021, 2022a,b; The LIGO Scientific Collaboration et al. 2022).

If a GW signal encounters the Sun along its path, gravitational lensing imprints a frequency-dependent modulation on the waveform. In geometrical optics (i.e., the zero-wavelength limit of GWs), the solar density modulates the amplitude with a magnification effect, while the gravitational potential modulates the phase by imposing a potential (or Shapiro) time delay. These modulations can be obtained along the transversal path of the Sun moving in front of the source (the duration of this movement is approximately half a day). Therefore, in principle, one can probe the solar interior using the lensed signal. GW lensing by the Sun has been studied as a tool for amplifying the strain amplitude of GWs from a distant source and probing the solar structure (e.g., Cyranski & Lubkin 1974; Sonnabend 1979; Patla & Nemiroff 2008; Marchant et al. 2020). Before the 1980s, these studies were based on geometrical optics. Bontz & Haugan (1981) first demonstrated that the diffraction effect caused by the finite wavelength of GWs suppresses magnification near the focal point at lower frequencies $(f \lesssim 10^4 \,\mathrm{Hz})$. Recently, Marchant et al. (2020) proposed that the solar structure can be probed through the lensed GW signal from a pulsar behind the Sun. However, they only roughly estimated the detectability of the lensing signature, without calculating the measurement accuracy of the density profile. Very recently, Jung & Kim (2022) reported that the Fresnel scale is comparable to the solar radius within the frequency band of the ground-based detectors; therefore, the density profile can (in principle) be probed with the lensing modulation on a chirp signal from a CBC.

In this study, we investigate the accuracy of measuring the solar density profile using the lensed signals of known pulsars. Because ground-based detectors detect GW wavelengths longer than (or comparable to) the solar Schwarzschild radius, gravitational lensing should employ wave optics (e.g., Ohanian 1974; Bliokh & Minakov 1975; Schneider et al. 1992; Nakamura & Deguchi 1999; Takahashi & Nakamura 2003; Dai et al. 2018; Oguri 2019; Suvorov 2022; Liao et al. 2022). We first calculate the lensed waveform based on wave optics. We discuss the effects of frequency and impact parameter on the waveform (Section 2). Then we extract the known pulsars




FIG. 1.— Configuration of the detector, Sun, and source. $D_{\rm L}$, $D_{\rm LS}$, and $D_{\rm S}$ denote the distances between them. Solid line is the GW path. Angular coordinates θ and $\theta_{\rm s}$ describe the incoming direction of the GWs and source position, respectively, with respect to the solar center.

crossing behind the Sun from the Australia Telescope National Facility (ATNF) pulsar catalog (Manchester et al. 2005). From the extracted list, we select suitable candidates by calculating the lensing modulations of these pulsars (Section 3). Using these candidates, we calculate the detectability of the lensing signature and the accuracy of measuring the solar density profile through a Fisher analysis (Section 4). We roughly estimate the number of Galactic millisecond pulsars (MSPs) behind the Sun, which are potentially detectable by near-future radio surveys (Section 5). We discuss the lensing by a Galactic star, which may be confused with the lensing by the Sun (Section 6). The study findings are summarized in Section 7.

2. AMPLITUDE AND PHASE MODULATIONS OF SOLAR LENSING

This section introduces the amplitude and phase modulations of a lensed waveform, based on wave optics.

2.1. Lensed waveform

Figure 1 shows the configuration of the detector, Sun, and source. The solid line describes a single GW path¹. We assume the thin-lens approximation, in which the solar density and its gravitational potential are projected onto a plane perpendicular to the line of sight. The thin-lens approximation is valid within an error less than 0.5% ($\simeq R_{\odot}/D_{\rm L}$) on the lensed waveform (Appendix A; Suyama et al. 2005). Furthermore, we assume the flat-sky approximation, in which $|\boldsymbol{\theta}|, |\boldsymbol{\theta}_{\rm s}| \ll 1$. As the coordinate system is fixed at the solar position, the background source (pulsar) moves along the ecliptic longitude, following the annual solar motion. The distance to the Sun is $D_{\rm L} = 1$ au and the solar angular radius is $\theta_{\odot} = \arctan[R_{\odot}/(1 \text{ au})] \simeq 16$ arcmin. The annual modulation of $D_{\rm L}$ is ignored because it changes by < 2% as the Earth moves elliptically around the Sun.

The lensed waveform $\tilde{h}^{L}(f)$ in the frequency domain is obtained by multiplying the unlensed waveform $\tilde{h}(f)$ by a function $F(f; \boldsymbol{\theta}_{s})$ (e.g., Nakamura 1998; Takahashi & Nakamura 2003):

$$\tilde{h}^{\mathrm{L}}(f;\boldsymbol{\theta}_{\mathrm{s}}) = F(f;\boldsymbol{\theta}_{\mathrm{s}})\tilde{h}(f),$$

= $[1 + A_{F}(f;\boldsymbol{\theta}_{\mathrm{s}})] \,\mathrm{e}^{\mathrm{i}\boldsymbol{\Phi}_{F}(f;\boldsymbol{\theta}_{\mathrm{s}})}\tilde{h}(f).$ (1)

In the second line, $A_F (\equiv |F| - 1)$ and $\Phi_F (\equiv \ln(F/|F|))$ represent the amplitude and phase modulations, respectively. The GW polarization is ignored because the polarization rotation due to lensing is negligibly small (e.g., Hou et al. 2019; Ezquiaga et al. 2021; Dalang et al. 2022). The function F, called the amplification factor or transmission factor, is given by the following diffraction integral (e.g., Schneider et al. 1992):

$$F(f;\boldsymbol{\theta}_{\rm s}) = \frac{D_{\rm L} D_{\rm S}}{c D_{\rm LS}} \frac{f}{{\rm i}} \int {\rm d}^2 \boldsymbol{\theta} \exp\left[2\pi {\rm i} f t_{\rm d}(\boldsymbol{\theta},\boldsymbol{\theta}_{\rm s})\right], \qquad (2)$$

where the time delay is given by

$$t_{\rm d}(\boldsymbol{\theta}, \boldsymbol{\theta}_{\rm s}) = \frac{1}{c} \left[\frac{D_{\rm L} D_{\rm S}}{2D_{\rm LS}} \left| \boldsymbol{\theta} - \boldsymbol{\theta}_{\rm s} \right|^2 - \frac{\hat{\psi}(\boldsymbol{\theta})}{c^2} \right].$$
(3)

The first and second terms are the geometrical and potential (or Shapiro) time delays, respectively. Because the distance to the Sun is much shorter than the distance to the source, hereafter we assume $D_{\rm S}/D_{\rm LS} \simeq 1$. The two-dimensional lens potential $\hat{\psi}(\boldsymbol{\theta})$ is determined from the solar-projected density profile $\Sigma(\boldsymbol{\theta})$ using the Poisson equation:

$$\nabla_{\theta}^2 \hat{\psi}(\boldsymbol{\theta}) = 8\pi G D_{\rm L}^2 \Sigma(\boldsymbol{\theta}). \tag{4}$$

Outside the Sun, the potential is identical to that of a point mass:

$$\hat{\psi}(\theta) = 4GM_{\odot} \ln\left(\frac{\theta}{\theta_{\rm E}}\right) \quad \text{for } \theta > \theta_{\odot},$$
 (5)

where $\theta_{\rm E}$ is an arbitrary constant, here set to the angular Einstein radius; i.e., $\theta_{\rm E} = [4GM_{\odot}/(c^2D_{\rm L})]^{1/2} \simeq 0.043 \,\theta_{\odot}$. The Sun is modeled using the BS05(OP) spherical density model² in Bahcall et al. (2005). The model density agrees within 2% of the helioseismological results across the whole radius (their Fig. 1). It also agrees within 3% and 4% of the recent solar density models of B16-GS98 and B16-AGSS09met, respectively, in Vinyoles et al. (2017). As the density profile is spherical, the integration (2) can be performed over the azimuth and F reduces to

$$F(f;\theta_{\rm s}) = \frac{2\pi f D_{\rm L}}{{\rm i}c} \int_0^\infty {\rm d}\theta \,\theta J_0\left(\frac{2\pi f D_{\rm L}}{c}\theta\theta_{\rm s}\right) \\ \times \exp\left[2\pi {\rm i}f\left\{\frac{1}{2c}D_{\rm L}\left(\theta^2 + \theta_{\rm s}^2\right) - \frac{\hat{\psi}(\theta)}{c^3}\right\}\right], \quad (6)$$

where J_0 is the zero-order Bessel function. Equation (6) is numerically integrated using integration by parts (Appendix A of Takahashi (2004); Guo & Lu (2020)).

Let us comment on the validity of the spherical density model of the Sun. The spin rotation makes the equatorial radius larger than the polar radius, but the observed relative difference between these surface radii

 $^{^1}$ In wave optics, the lensed waveform is the superposition of many GW paths from the source to the detector (Eq. (2)). The solid line represents one of them.

² The numerical table can be downloaded from http://www.sns. ias.edu/~jnb/SNdata/sndata.html#bs2005.

is very small $\approx 10^{-5}$ (Chapter 7 of Stix 2004). The quadrupole mass moment of the Sun $(J_2)^3$ generates the precession of Mercury's orbital perihelion (which is much smaller than the known general-relativistic effect). The precession was precisely measured by the MESSENGER (MErcury Surface, Space ENviroment, GEochemistry, and Ranging) spacecraft⁴, which gave a constraint on $J_2 \simeq 2 \times 10^{-7}$ (Park et al. 2017). These observations suggest that F for the spherical model would be valid with an error of $\lesssim 10^{-5}$.

2.2. High- and low-frequency limits

This subsection presents the high- and low-frequency limits of the amplification factor. In the high-frequency limit (i.e., the geometrical-optics limit), because the exponential term of F in Eq. (2) oscillates violently, a stationary point of $t_d(\theta, \theta_s)$ contributes to the integral (e.g., Schneider et al. 1992). This point is a solution of $\nabla_{\theta} t_d(\theta, \theta_s) = 0$:

$$\boldsymbol{\theta}_{\rm s} = \boldsymbol{\theta} - \hat{\boldsymbol{\alpha}}(\boldsymbol{\theta}), \tag{7}$$

from Eq. (3), where $\hat{\alpha} = \nabla_{\theta} \hat{\psi}/(c^2 D_{\rm L})$ is the deflection angle. Equation (7) is the lens equation. The image position $\theta_{\rm i}$ is obtained by solving Eq. (7). Note that only a single image is formed irrespective of $\theta_{\rm s}$. In the high-frequency limit, F reduces to the following simple form:

$$\lim_{f \to \infty} F(f; \boldsymbol{\theta}_{s}) = |\mu(\boldsymbol{\theta}_{i}, \boldsymbol{\theta}_{s})|^{1/2} \exp\left[2\pi i f t_{d}(\boldsymbol{\theta}_{i}, \boldsymbol{\theta}_{s})\right], \quad (8)$$

where $\mu = [\det(\partial \theta_s / \partial \theta_i)]^{-1}$ is the magnification of the image.

In the weak-gravitational-field limit, the magnification can be approximated as $\mu \simeq 1 + 2\kappa$, where $\kappa = \Sigma/\Sigma_{\rm cr}$ is the convergence and $\Sigma_{\rm cr} = c^2/(4\pi G D_{\rm L})$ is the critical density (e.g., Bartelmann & Schneider 2001). Accordingly, A_F and Φ_F reduce to

$$\lim_{f \to \infty} A_F(f; \boldsymbol{\theta}_{\rm s}) \simeq \kappa(\boldsymbol{\theta}_{\rm s}) = \frac{4\pi G D_{\rm L}}{c^2} \Sigma(\boldsymbol{\theta}_{\rm s}),$$
$$\lim_{f \to \infty} \Phi_F(f; \boldsymbol{\theta}_{\rm s}) = 2\pi f t_{\rm d}(\boldsymbol{\theta}_{\rm i}, \boldsymbol{\theta}_{\rm s}) \simeq -2\pi f \frac{\hat{\psi}(\boldsymbol{\theta}_{\rm s})}{c^3}.$$
 (9)

Therefore, A_F and Φ_F trace the projected density profile and gravitational potential, respectively.

Figure 2 plots the deflection angle, convergence, and gravitational potential as functions of θ (as these variables are functions of θ , we plot θ rather than $\theta_{\rm s}$ along the x-axis). The deflection angle is consistent with the known result $\hat{\alpha} = 4GM_{\odot}/(c^2R_{\odot}) \simeq 1.75$ arcsec at the surface. The maximum deflection is $\hat{\alpha} \simeq 4.57$ arcsec at $\theta \simeq 0.23 \theta_{\odot}$. Because $\hat{\alpha}$ is much smaller than the solar radius ($\theta_{\odot} \simeq 16$ arcmin), we can safely set $\theta_{\rm i} \simeq \theta_{\rm s}$ from Eq. (7). Therefore, the second term of $t_{\rm d}$ in Eq. (3) exceeds the first term in the geometrical-optics limit. Meanwhile, the convergence κ is maximized at

~ 0.04 near the center. The profile is flat at the core $(\theta \leq 0.03 \theta_{\odot})$, but it drops steeply at $\theta \geq 0.1 \theta_{\odot}$. The typical gravitational time delay is $\approx 10^{-5}$ s. Near the center, the potential in Eq. (4) can be approximated as $\hat{\psi}(\theta) = \text{const.} + (c^2 D_{\text{L}}/2) \kappa(\theta = 0) \theta^2 + \mathcal{O}(\theta^3)$. The results of Fig. 2 are consistent with previous works on lensing by the transparent Sun (Bontz & Haugan 1981; Patla & Nemiroff 2008; Marchant et al. 2020; Jung & Kim 2022).

In the large-angle limit $\theta_{\rm s} \gg \theta_{\odot}$, the geometrical-optics result (9) is recovered because $2\pi f t_{\rm d} \gg 1$. As the gravitational potential in Eq. (5) is valid only for $\theta_{\odot} < \theta \ll 1$, it must be replaced with (e.g., Backer & Hellings 1986)

$$\hat{\psi}(\theta) = 2GM_{\odot} \ln\left\{\frac{2(1-\cos\theta)}{\theta_{\rm E}^2}\right\},\tag{10}$$

when $\theta \gtrsim 1$.

We now investigate the low-frequency limit. Changing the variables to $\theta' = \sqrt{f}\theta$ and $\theta'_{s} = \sqrt{f}\theta_{s}$ in Eq. (2), F is rewritten as

$$F(f;\boldsymbol{\theta}_{\rm s}) = \frac{D_{\rm L}}{\mathrm{i}c} \int \mathrm{d}^2 \boldsymbol{\theta}' \exp\left[\frac{2\pi\mathrm{i}}{c} \left\{\frac{D_{\rm L}}{2} \left|\boldsymbol{\theta}' - \boldsymbol{\theta}_{\rm s}'\right|^2 - \frac{f}{c^2} \hat{\psi}\left(\frac{\boldsymbol{\theta}'}{\sqrt{f}}\right)\right\}\right].$$
(11)

In the low-frequency limit, $\hat{\psi}$ can be replaced with its value at infinity:

$$\lim_{f \to 0} \hat{\psi} \left(\frac{\theta'}{\sqrt{f}} \right) = 4GM_{\odot} \ln \left(\frac{\theta'}{\sqrt{f}\theta_{\rm E}} \right),$$

from Eq. (5). Inserting this $\hat{\psi}$ into Eq. (11) and expanding f as a Taylor series, we have

$$\lim_{f \to 0} A_F(f; \boldsymbol{\theta}_{\rm s}) = \frac{2\pi^2 G M_{\odot}}{c^3} f,$$
$$\lim_{f \to 0} \Phi_F(f; \boldsymbol{\theta}_{\rm s}) = \frac{4\pi G M_{\odot}}{c^3} f \left[\gamma + \ln\left(\frac{4\pi G M_{\odot}}{c^3} f\right) \right],$$
(12)

where $\gamma = 0.5772 \cdots$ is Euler's constant.

2.3. Amplitude and phase modulations

Figure 3 plots A_F and Φ_F as functions of f for various θ_s . In the high- and low-frequency limits, the plots are consistent with the analytical results of Eqs. (9) and (12), respectively. As the frequency of a chirp signal from an inspiral binary sweeps from low to high, A_F increases proportionally to f in the wave-optics regime but approaches a constant value of κ in the geometrical-optics regime. Similarly, $-\Phi_F/(2\pi f)$ decreases logarithmically, but approaches a constant value of the potential time delay $(\hat{\psi}/c^3)$. These modulations are observed even outside the Sun $(\theta_s = 2\theta_{\odot})$. Therefore, within the frequency range of ground-based detectors, the solar lensing imprints unique frequency-dependent modulations on both the amplitude and phase. The solar structure can (in principle) be extracted from the chirp signal (see the recent study by Jung & Kim 2022). However, in the high-frequency limit, solar structure extraction from the signal (even from a chirp signal) is impossible at a single θ_s because the constant A_F is degenerate with an

³ Modeling the Sun as a spheroidal body about its spin axis (z axis), J_2 is simply $J_2 = (I_{zz} - I_{xx})/(M_{\odot}R_{\odot}^2)$ with the moment of inertia I_{ij} . A positive (negative) J_2 means an oblate (prolate) shape.

⁴ https://solarsystem.nasa.gov/missions/messenger/ in-depth/




FIG. 2.— Solar deflection angle, convergence, and potential (or Shapiro) time delay as functions of angular radius θ (normalized by the solar radius θ_{\odot}). The convergence κ in the middle panel is the dimensionless projected density profile. The potential time delay in the right panel is the projected gravitational potential divided by c^3 . Near and outside the surface ($\theta/\theta_{\odot} \gtrsim 1$), the result approaches that of Eq. (5) for a point mass (the dashed line).




FIG. 3.— Amplitude and phase modulations as functions of frequency at various source positions $\theta_s/\theta_{\odot} = 0-2$. Horizontal dashed lines and dot-dashed orange line represent the analytical results in the high- and low-frequency limits, respectively, given by Eqs. (9) and (12). Up arrows indicate the rough boundary between the wave- and geometrical-optics regions for each θ_s (see the description after Eq. (13) in the text). In the left panel, dotted curves denote negative values ($A_F < 0$). In the right panel, the phase modulation is divided by $2\pi f$, which corresponds to an arrival time delay.

intrinsic amplitude of GWs. Similarly, the constants Φ_F and $\Phi_F/(2\pi f)$ are degenerate with an intrinsic phase and arrival time, respectively. Therefore, in geometrical optics, the lensing modulations are indistinguishable from the intrinsic source properties (unless lensed signals with various θ_s from a single source are available). Figure 3 is consistent with Fig. 2 of Jung & Kim (2022).

To discuss the behavior of these modulations in more detail, let us introduce the angular Fresnel scale (Macquart 2004; Takahashi 2006):

$$\theta_{\rm F} = \left(\frac{c}{2\pi f} \frac{1}{D_{\rm L}}\right)^{1/2},$$
$$\simeq 0.38 \,\theta_{\odot} \left(\frac{f}{100 \,\mathrm{Hz}}\right)^{-1/2}.$$
(13)

In the low-frequency limit, the lensed signal probes a circle of radius $\theta_{\rm F}$ around the source position (Section 2 and Fig. 1 of Choi et al. 2021). In other words, the Fresnel scale can be interpreted as the effective source




FIG. 4.— Same as Fig.3, but plotted as functions of θ_s/θ_{\odot} for various f. Red curve overlaps the purple curve everywhere except near the center ($\theta_s/\theta_{\odot} \leq 0.2$).

radius (Oguri & Takahashi 2020). The effective source radius shrinks with increasing f in the chirp signal. In the high-frequency limit, the lensing probes a small region around the image position. As the modulations at different f can probe different regions, one can in principle probe the density profile (Jung & Kim 2022). Here, $\theta_{\rm F} = \theta_{\rm s}$ (unless $\theta_{\rm s} = 0$) roughly represents the boundary between the wave- and geometrical-optics regions. This boundary condition is derived from $2\pi f t_{\rm d} = 1$ with neglecting the potential term in Eq. (3) (Section II.B of Choi et al. (2021) and Section 3 of Jung & Kim (2022))⁵. The boundary is plotted as a series of up arrows in Figs. 3 and 4.

Figure 4 plots A_F and Φ_F as functions of θ_s for various fixed f. As the GWs from a pulsar are continuous, one can (in principle) measure A_F and Φ_F as functions of θ_s for a pulsar moving behind the Sun. For smaller (larger) θ_s , the exponential term of the amplification factor (= $2\pi f t_d$) is smaller (larger) and the results approach the low-frequency (high-frequency) limit in Eq. (12) (Eq. (9)). At f = 1000 Hz, the result almost matches that of geometrical optics, except in the near-center region ($\theta_s/\theta_{\odot} \leq 0.2$). As Φ_F reflects the gravitational potential, it can be measured even outside the Sun. However, A_F , which reflects the mass density, falls steeply near the surface. Figure 4 is consistent with Figs. 6 and 7 of Marchant et al. (2020).

The LVK analysis accounts for the Shapiro time delay imposed by the Sun. In searches of continuous GWs (Abbott et al. 2017, 2019, 2022a), the time delay has been calculated using the TEMPO (Nice et al. 2015) and TEMPO2 (Edwards et al. 2006; Hobbs et al. 2006) packages developed for timing analyses of pulsar radio signals. However, as shown in Figs. 3 and 4, the Shapiro time delay is less accurate at lower f and smaller θ_s . To clarify the accuracy of the Shapiro time delay, Fig. 5 plots a contour map of the difference between the phase




FIG. 5.— Contour plot of the difference between the phase modulation and its geometrical-optics limit $(\Phi_F - 2\pi f t_d)$. Red, blue, and white regions correspond to positive, negative, and zero differences, respectively. Dotted line plots the rough boundary between the wave- and geometrical-optics regions, calculated from $\theta_F = \theta_s$ (the description after Eq. (13) in the text).

modulation and its geometrical-optics limit. The difference ranges from -0.015 to 0.002 rad and is most significant at $f \simeq 100-1000$ Hz near the center ($\theta_s \leq 0.2 \theta_{\odot}$). At lower frequencies ($f \leq 10$ Hz), the difference reduces because both Φ_F and $2\pi f t_d$ are proportional to f. As expected, the Shapiro time delay is recovered at high f or large θ_s . The dotted line, predicted by setting $\theta_F = \theta_s$, is apparently consistent with the boundary. Although the Shapiro time delay is less accurate inside the Sun, the error is small (0.015 rad at most).

3. KNOWN PULSARS

This section describes some known pulsars moving behind the Sun. From the results in Subsection 2.3, a pulsar with a higher frequency and a smaller impact parameter to the Sun is more suitable for probing the solar structure. We searched the ATNF pulsar catalog⁶

⁵ This condition can also be derived from the intersection of the low- and high-frequency limits of the phase modulation. Setting $\lim_{f\to 0} \Phi_F = \lim_{f\to\infty} \Phi_F$ in Eqs. (9) and (12) with the point-mass potential (5) (which is valid only for $\theta_{\rm s} \gtrsim 0.3 \,\theta_{\odot}$ in the right panel of Fig. 2), one obtains $\theta_{\rm F} \simeq 0.94 \,\theta_{\rm s}$.

⁶ http://www.atnf.csiro.au/research/pulsar/psrcat/

BUT NOT BEHIND, THE SUN (LOWER EIGHT)						
name	$f(\mathrm{Hz})$	$ heta_{ m LAT}/ heta_{\odot}$	$D_{\rm s}({\rm kpc})$	$h_0^{95\%}$	$\epsilon^{95\%}$	references
J1745-23	369	-0.07	7.94			C20
J1022+1001	122	-0.24	0.64	7.7×10^{-27}	3.2×10^{-7}	C96,R21
J1809 - 2332	14	-0.45	0.88			A09,R11
J1730 - 2304	246	0.71	0.47	5.0×10^{-27}	$3.7 imes 10^{-8}$	L95, R21
J1858-2216	839	1.80	0.92	7.8×10^{-27}	9.6×10^{-9}	S16
J1142 + 0119	394	-1.84	2.17	7.4×10^{-27}	9.8×10^{-8}	S16
J2310-0555	766	-1.99	1.56			S16
J1756-2251	70	2.14	0.73	6.1×10^{-27}	8.6×10^{-7}	F04,F14
J1646-2142	342	2.44	0.97			R12,R13
J1811-2405	752	-2.52	1.83	1.5×10^{-26}	4.5×10^{-8}	K10,N20
J1836-2354B	619	-2.74	3.20			L11
J1836-2354A	596	-2.78	3.20			L11

TABLE 1 PULSARS WITH f > 10 Hz crossing behind the Sun (upper four) and close to, BUT NOT BEHIND, THE SUN (LOWER EIGHT)

NOTE. — Second column: GW frequency. Third column: ecliptic latitude $\theta_{\rm LAT}$ (i.e., the minimum angular separation to the solar center). Pulsars are listed in order of increasing $|\theta_{\rm LAT}|$. Fourth column: distance $D_{\rm s}$, measured by parallax for J1022+1001, J1730-2304, and J1756-2251 and estimated from the dispersion measures with the Galactic free-electron distribution model YMW16 (Yao et al. 2017) for the other pulsars. Fifth and sixth columns: 95% upper limits on the strain amplitude and ellipticity, respectively, taken from the LVK results (Abbott et al. 2022a). Last column: references A09 (Abdo et al. 2009), C20 (Cameron et al. 2020), C96 (Camilo et al. 1996), F04 (Faulkner et al. 2004), F14 (Ferdman et al. 2014), K10 (Keith et al. 2010), L95 (Lorimer et al. 1995), L11 (Lynch et al. 2011), N20 (Ng et al. 2020), R11 (Ray et al. 2011), R12 (Ray et al. 2012), R21 (Reardon et al. 2021), R16 (Roy & Bhattacharyya 2013), and S16 (Sanpa-Arsa 2016).

for pulsars satisfying the following criteria: i) $f > 10 \,\text{Hz}$ and ii) closest distance to the solar center is within $3 \theta_{\odot}$. Twelve samples met the criteria: four pulsars crossing behind the Sun and eight pulsars passing near but not behind the solar surface. The information of these pulsars is summarized in Table 1. The GW frequency f is twice the spin frequency for the GW emission from the mass-quadrupole moment. These frequencies are covered by the sensitivity range of the current ground-based detectors. The LVK collaboration obtained 95% credible upper limits on the strain amplitudes of some of these samples from the O2 and O3 runs (Abbott et al. 2022a). They also presented the corresponding upper bounds on the ellipticity of the mass distribution (estimated from Eq. (15)). The collaboration also performed an all-sky search for continuous GWs coming from any (known and unknown) spinning neutron stars in isolated and binary systems (Abbott et al. 2021; The LIGO Scientific Collaboration et al. 2022). Their upper bound was $h_0^{95\%} \sim 1.1 (2.0) \times 10^{-25}$ at f = 100-200 (50-700) Hz for isolated systems (and similarly for binary systems).

Figure 6 shows the trajectories of the upper four pulsars in Table 1. These pulsars move along the horizontal axis with a velocity of $\dot{\theta}_{\rm LON} \simeq -2\pi/(1{\rm yr}) = -2.5 \, {\rm arcmin/hour} = -0.15 \, \theta_{\odot}/{\rm hour}$. The typical crossing time is $2\theta_{\odot}/|\dot{\theta}_{\rm LON}| \simeq 13$ hours. Marchant et al. (2020) included J1022+1001 and J1730-2304 as candidates, but excluded the very recently discovered pulsar J1745-23 (Cameron et al. 2020).

Figure 7 plots A_F and Φ_F as functions of θ_{LON} for the four pulsars. J1745-23, with the highest frequency and closest impact parameter, yields the maximum A_F (~ 2%). The highest $|\Phi_F|$ is ~ 0.1 rad for all pulsars except J1809-2332. The anomalous result for J1809-2332 is attributable to the low frequency (14 Hz) of this pulsar. The typical $|\Phi_F|$ is estimated as $|\Phi_F| \simeq 2\pi f |t_d| \simeq$





FIG. 6.— Trajectories of pulsars moving behind the Sun (represented by an orange circle of radius θ_{\odot}). θ_{LAT} is the ecliptic latitude and θ_{LON} is the ecliptic longitude with origin set at the solar center.

0.1 rad $(f/200 \text{ Hz}) (|t_{\rm d}|/100 \,\mu\text{s})$. The right panel clearly shows the potential well around $\theta_{\rm LON} = 0$. The effect is especially noticeable in the curves of J1745-23 and J1022+1001. As evidenced in the figure, $|A_F| \ll |\Phi_F|$.

Figure 8 plots Φ_F as a function of θ_{LON} for the lower eight pulsars in Table 1. Because these pulsars (except for J1756-2251) have relatively high frequencies, their $|\Phi_F|$ are relatively large (> 0.1 rad). As these pulsars do not pass behind the Sun, their $|A_F|$ are negligibly small.

4. PARAMETER EXTRACTION FROM THE LENSED WAVEFORM

This section evaluates the accuracy to which the solar density profile can be extracted from the lensed GW signal through a Fisher analysis.




-2

0.01

6

FIG. 7.— Amplitude and phase modulations of the four pulsars moving behind the Sun. The lower x axis is the pulsar position $\theta_{\text{LON}}/\theta_{\odot}$ along the ecliptic longitude (Fig. 6), while the upper x axis represents the corresponding crossing time. Dotted curves in the left panel represent negative values.



0

 $\theta_{\rm LON}/\theta_{\odot}$

1

2

10

10⁻²

10⁻³

10

-2

-1

 A_F

FIG. 8.— Phase modulations of the eight pulsars passing close to (but not behind) the Sun.

4.1. Fisher matrix analysis

We consider a highly spinning neutron star with a small non-axisymmetry around the spin axis as the GW source. Neglecting spin down, the source is assumed to emit continuous monochromatic GWs during the observational period. A waveform of frequency f is then described as

$$h(t;f) = h_0 \cos(2\pi f t + \phi_0), \qquad (14)$$

where ϕ_0 is a constant phase. The strain amplitude h_0 is determined by the non-axisymmetry as (e.g., Riles 2022)

$$h_0 = \frac{4\pi^2 \epsilon G I f^2}{c^4 D_{\rm s}},\tag{15}$$

where I is the moment of inertia about the spin axis and ϵ is the ellipticity of the moment of inertia. A gravitationally lensed waveform $h^{L}(t; f)$ can be obtained from

Eqs. (1) and (14) as⁷ $b^{L}(t, f) = b [1 + A (t, f)] \cos [2\pi ft + \phi + f]$

-1

$$h^{\rm L}(t;f) = h_0 \left[1 + A_F(t;f) \right] \cos \left[2\pi f t + \phi_0 + \Phi_F(t;f) \right].$$
(16)

0

 $\theta_{\rm LON}/\theta_{\odot}$

1

2

Here, A_F and Φ_F are numerically obtained from Eqs. (1) and (6). When θ_s is large ($\theta_s > 20 \theta_{\odot}$), we can apply the geometrical-optics results (Eqs. (9) and (10)), which are fully valid at such large angles (Fig. 4).

Suppose that the pulsar is observed from time t = -T/2 to T/2, where T is the observational period and t = 0 is the time at which the pulsar is closest to the solar center. The signal-to-noise ratio (SNR) is calculated as

$$SNR^{2} = \frac{1}{S_{n}(f)} \int_{-T/2}^{T/2} dt \left[h^{L}(t;f) \right]^{2},$$
$$\simeq \frac{h_{0}^{2}}{2S_{n}(f)} T, \qquad (17)$$

where S_n is the noise spectrum of the detector. Throughout this paper, the measurement accuracies of the fitting parameters are normalized by the SNR. Therefore, the results do not depend on a specific form of S_n . In the second equality of Eq. (17), we neglect A_F and use the approximation $\cos^2(2\pi ft + \phi_0 + \Phi_F) \simeq 1/2$ obtained by long-time averaging. We comment on the validity of the approximation in Eq. (17). The left panel of Fig. 7 shows $A_F \approx \mathcal{O}(0.01)$ for a few hours just behind the Sun; therefore, its average over one year is $\langle A_F \rangle_{1\mathrm{yr}} \approx \mathcal{O}(0.01) \times (\text{a few hours})/(1\mathrm{yr}) \approx \mathcal{O}(10^{-6}).$ Similarly, in the phase of Eq. (16), the first term, $2\pi ft \approx 6 \times 10^{10} (f/300 \,\mathrm{Hz})(t/1 \,\mathrm{yr})$, is ten orders of magnitude larger than the third term, $\Phi_F \approx \mathcal{O}(1)$ rad from Figs. 7 and 8. This suggests that A_F and Φ_F give negligible contributions to the SNR (but note that A_F does increase the SNR during the crossing). This also suggests

⁷ Here, we perform a Fourier transform to obtain the unlensed waveform in the frequency domain, then calculate the lensed waveform using Eq. (1), and finally repeat the Fourier transform to obtain the lensed waveform (16) in the time domain.

that $h_L(t; f)$ is rapidly oscillating during the observation and thus the approximation $\cos^2(2\pi ft + \phi_0 + \Phi_F) \simeq 1/2$ is valid with a high precision.

The lensed waveform (16) depends on the solar density profile, the constant amplitude h_0 , and the constant phase ϕ_0 . Let the solar density profile be characterized by N parameters A_i (i = 1, 2, ..., N). The total set of fitting parameters is $p_{\mu} = (A_i, \ln h_0, \phi_0)$ with $\mu = 1, 2, \dots, N+2$. Let us consider a GW signal as a sum of the theoretical template (16) characterized by the fiducial parameters \bar{p}_{μ} and a Gaussian noise. Here, our fiducial solar model is the BS05(OP) in Bahcall et al. (2005). In a likelihood analysis, one can obtain the best-fit parameters by fitting the theoretical template to the signal. In the limit of high SNR, the posterior distribution of \boldsymbol{p} is multivariate Gaussian: $P(\mathbf{p}) \propto \exp\left[-\sum_{\mu,\nu} (p_{\mu} - \bar{p}_{\mu}) \Gamma_{\mu\nu} (p_{\nu} - \bar{p}_{\nu})/2\right]$ (for a detailed discussion, see Cutler & Flanagan 1994). The measurement accuracy of p_{μ} is given by the inverse of the Fisher matrix: $\Delta p_{\mu} = [(\Gamma^{-1})_{\mu\mu}]^{1/2}$. Neglecting the degeneracy with the other parameters, $\Delta p_{\mu} = (\Gamma_{\mu\mu})^{-1/2}$. The Fisher matrix is given by (e.g., Cutler 1998)

$$\Gamma_{\mu\nu} = \frac{1}{S_n(f)} \int_{-T/2}^{T/2} dt \, \frac{\partial h^{\rm L}(t;f)}{\partial p_{\mu}} \frac{\partial h^{\rm L}(t;f)}{\partial p_{\nu}}.$$
 (18)

Using Eqs. (16) and (17), all components of Γ are given as follows:

$$\Gamma_{A_{i}A_{j}} = \int_{t} \left[\frac{\partial A_{F}(t;f)}{\partial A_{i}} \frac{\partial A_{F}(t;f)}{\partial A_{j}} + \left\{ 1 + A_{F}(t;f) \right\}^{2} \frac{\partial \Phi_{F}(t;f)}{\partial A_{i}} \frac{\partial \Phi_{F}(t;f)}{\partial A_{j}} \right],$$

$$\Gamma_{A_{i}\lnh_{0}} = \int_{t} \left\{ 1 + A_{F}(t;f) \right\} \frac{\partial A_{F}(t;f)}{\partial A_{i}},$$

$$\Gamma_{A_{i}\phi_{0}} = \int_{t} \left\{ 1 + A_{F}(t;f) \right\}^{2} \frac{\partial \Phi_{F}(t;f)}{\partial A_{i}},$$

$$\Gamma_{\lnh_{0}\lnh_{0}} = \Gamma_{\phi_{0}\phi_{0}} = \int_{t} \left\{ 1 + A_{F}(t;f) \right\}^{2},$$

$$\Gamma_{\lnh_{0}\phi_{0}} = 0,$$
(19)

with

$$\int_t \equiv \mathrm{SNR}^2 \, \frac{1}{T} \int_{-T/2}^{T/2} dt.$$

Similar to Eq. (17), we here applied the approximations $\sin^2 \operatorname{or} \cos^2(2\pi ft + \phi_0 + \Phi_F) \simeq 1/2$ and $\sin(2\pi ft + \phi_0 + \Phi_F) \cos(2\pi ft + \phi_0 + \Phi_F) \simeq 0$. From Eq. (19), Δp_{μ} simply scales as $\Delta p_{\mu} \propto \mathrm{SNR}^{-1}$.

The simple waveform (14) neglects the detector response which depends on other source parameters such as the inclination of the spin axis and GW polarization. Owing to Earth's spin and orbital motion, the detector response causes daily and yearly periodic modulations on both amplitude and phase. Unlike the solar modulation (which is temporary but repeats annually), the response modulations are trigonometric functions of time. We expect that the solar modulation can be distinguishable from the response modulations and thus the response does not significantly affect the measurement accuracies of the solar model parameters.

TABLE 2MEASUREMENT ACCURACIES OF THEPOTENTIAL AMPLITUDE A_{ψ} (THE FIDUCIALVALUE = 1) OVER ONE YEAR OFOBSERVATION WITH SNR = 100.

name	ΔA_{ψ}	$f\Delta A_{\psi}$ (Hz)
J1745-23	0.24 (0.026)	90
J1022 + 1001	$0.74 \ (0.077)$	90
J1809 - 2332	6.6(0.69)	90
J1730 - 2304	0.37 (0.038)	91
J1858-2216	$0.11 \ (0.011)$	92
J1142 + 0119	0.23 (0.024)	92
J2310-0555	$0.12 \ (0.012)$	92
J1756-2251	1.3(0.13)	92
J1646-2142	0.27 (0.028)	93
J1811-2405	0.12(0.013)	93
J1836-2354B	0.16(0.016)	93
J1836-2354A	0.15 (0.015)	93

NOTE. — Values in parentheses are calculated without the parameter degeneracy.

4.2. Constraint on the potential amplitude

This subsection presents the measurement accuracy of the overall amplitude of the solar gravitational potential. Replacing the potential as

$$\hat{\psi}(\theta) \to A_{\psi}\hat{\psi}(\theta),$$
 (20)

we calculate the measurement error in A_{ψ} . The derivatives in the Fisher matrix with respect to A_{ψ} are numerically obtained by changing A_{ψ} by ± 1 %.

Figure 9 plots the measurement accuracies of A_{ψ} , $\ln h_0$ and ϕ_0 for the pulsars moving behind the Sun at f >100 Hz. The results are normalized with SNR = 100 overa one-year observation period. Because $|A_F| \ll |\Phi_F|$, A_{ψ} is mainly determined by the phase modulation. The slope of ΔA_{ψ} changes slightly at the surface (corresponding to $T \approx 10^{-3}$ yr); the slope is steeper (shallower) inside (outside) the Sun. These results can be explained as follows. In the Sun's interior, $|\Phi_F|$ increases with θ (in contrast to the constant ϕ_0), thus A_{ψ} and ϕ_0 can be determined almost independently. Outside the Sun, Φ_F is nearly constant at large $\theta \gg \theta_{\odot}$, thus A_{ψ} and ϕ_0 are degenerate to some extent. The dashed red curve plots ΔA_{ψ} without the parameter degeneracy (i.e., $\Delta A_{\psi} = (\Gamma_{A_{\psi}A_{\psi}})^{-1/2}$, which is ~ 10 times better than the solid red curve. Previously, Marchant et al. (2020) roughly estimated the detectability of lensing signatures. As they ignored the parameter degeneracy, they underestimated the measurement error. Inside the Sun, h_0 cannot be determined, because h_0 and A_F are highly degenerate, but outside the Sun, the degeneracy is broken and $\Delta \ln h_0 \simeq \text{SNR}^{-1}$.

We caution that $\Delta \phi_0$ shown in Fig. 9 depends on an arbitrary constant in the gravitational potential. If a constant term is added to the potential, i.e., $\hat{\psi}(\theta) \rightarrow \hat{\psi}(\theta) + \hat{\psi}_0$, the phase modulation changes as $\Phi_F \rightarrow \Phi_F - 2\pi f \hat{\psi}_0/c^3$ using Eqs. (1)–(3). Two components of the Fisher matrix change accordingly⁸. After some algebra, one finds that ΔA_{ψ} , $\Delta \ln h_0$, and $(\Gamma^{-1})_{A_{\psi} \ln h_0}$ are

 $\label{eq:Gamma-star} \begin{array}{l} ^8 \ \Gamma_{A_\psi \phi_0} \rightarrow \Gamma_{A_\psi \phi_0} - (2\pi f \hat{\psi}_0/c^3) \Gamma_{\phi_0 \phi_0} \ \text{and} \ \Gamma_{A_\psi A_\psi} \rightarrow \Gamma_{A_\psi A_\psi} - (4\pi f \hat{\psi}_0/c^3) \Gamma_{A_\psi \phi_0} + (2\pi f \hat{\psi}_0/c^3)^2 \Gamma_{\phi_0 \phi_0}. \end{array}$




FIG. 9.— Measurement accuracies of the solar-potential amplitude A_{ψ} defined in Eq. (20), amplitude h_0 , and phase ϕ_0 of the pulsars crossing behind the Sun. Dashed red curve plots ΔA_{ψ} neglecting the parameter degeneracy. The horizontal axis T denotes the observational period. The signal-to-noise ratio (SNR) is set to $100 (T/\text{yr})^{1/2}$ and the results simply scale as $\Delta p \propto \text{SNR}^{-1}$. Thick gray line denotes SNR^{-1} .

TABLE 3 Measurement accuracies of the density amplitudes in two annuli over a one-year observation period with $SNR = 10^4$.

name	$\begin{array}{c} \text{Case (I)} \\ \Delta A_{\Sigma 1,2} \end{array}$	Case (II) $\Delta A_{\Sigma 1}, \Delta A_{\Sigma 2}$
J1745-23	0.40	0.11, 1.8
J1022+1001 J1730-2304	$2.0 \\ 2.9$	$\begin{array}{ccc} 0.47, \ 8.0 \\ 1.3, \ 22 \end{array}$
<u> </u>		

NOTE. — In Case (I), $\Delta A_{\Sigma 1} = \Delta A_{\Sigma 2}$.

independent of $\hat{\psi}_0$ but $\Delta \phi_0$ and its cross correlations, $(\Gamma^{-1})_{A_\psi \phi_0}$ and $(\Gamma^{-1})_{\ln h_0 \phi_0}$, do depend on $\hat{\psi}_0$. Table 2 lists the ΔA_ψ results of all pulsars. Although

Table 2 lists the ΔA_{ψ} results of all pulsars. Although ΔA_{ψ} varies among the pulsars, $f \Delta A_{\psi}$ is almost constant; i.e., it is independent of the impact parameter to the Sun. ΔA_{ψ} can then be approximately fitted as

$$\Delta A_{\psi} \approx 0.3 \left(\frac{f}{300 \,\mathrm{Hz}}\right)^{-1} \left(\frac{\mathrm{SNR}}{100}\right)^{-1}, \qquad (21)$$

indicating that a higher-frequency pulsar is more promising for detecting A_{ψ} . According to Eq. (21), A_{ψ} can be detected at the 3σ confidence level when SNR $\approx 100 (f/300 \text{Hz})^{-1}$.

4.3. Measurement of the solar density profile

This subsection presents our main result, namely the measurement accuracy of the solar density profile. Let the surface density be divided into N annuli (Fig. 10). We write the surface mass density of the i-th annulus as $\Sigma_i(\theta)$ and the mass as M_i ($i = 1, \dots, N$), where a smaller i corresponds to an inner annulus. We consider the following two cases:

Case (I): Each annulus has the same mass
$$(= M_{\odot}/N)$$
.




FIG. 10.— Division of the solar surface into N annuli. Each annulus has the same mass in Case (I) and the same radial width in Case (II).

TABLE 4 SAME AS TABLE 3, BUT FOR THREE ANNULI AND A ONE-YEAR OBSERVATION PERIOD WITH $SNR = 10^5$.

name	Case (I) $\Delta A_{\Sigma 1}, \Delta A_{\Sigma 2}, \Delta A_{\Sigma 3}$	Case (II) $\Delta A_{\Sigma 1}, \Delta A_{\Sigma 2}, \Delta A_{\Sigma 3}$
$\begin{array}{c} J1745{-}23\\ J1022{+}1001\\ J1730{-}2304 \end{array}$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$

Case (II): Each annulus has the same radial thickness $(= \theta_{\odot}/N)$.

We first consider N = 2 and 3. In Case (I), the boundary radius between the annuli is $0.20 \theta_{\odot}$ for N = 2. The




FIG. 11.— Measurement accuracies of the solar-density amplitudes in the i-th annulus $A_{\Sigma i}$ for the pulsars crossing behind the Sun. Upper and lower panels plot the results in Case (I) and (II), respectively. In the upper panels, $\Delta A_{\Sigma 1} = \Delta A_{\Sigma 2}$. The horizontal axis T denotes the observational period. SNR is set to $10^4 (T/yr)^{1/2}$ and the results simply scale as $\Delta A_{\Sigma i} \propto \text{SNR}^{-1}$.

boundary radii are $0.14 \theta_{\odot}$ and $0.26 \theta_{\odot}$ between annulus 1 and 2 and between annulus 2 and 3, respectively, for N = 3. In Case (II), the mass ratios are $M_1 : M_2 \simeq 17 : 1$ for N = 2 and $M_1 : M_2 : M_3 \simeq 57 : 13 : 1$ for N = 3 (i.e., the mass reduces from the innermost to the outermost annulus).

We change the amplitude of the i-th density as

$$\Sigma_{\rm i}(\theta) \to A_{\Sigma \rm i} \Sigma_{\rm i}(\theta),$$
 (22)

while fixing the total mass, i.e., $\sum_{i=1}^{N} A_{\Sigma i} M_i = M_{\odot}$. Under these settings, we calculate the measurement accuracies of $A_{\Sigma i}$, $\ln h_0$, and ϕ_0 . When N = 2, the density profile is solely characterized by the fitting parameter $A_{\Sigma 1}$ because the total mass is fixed, i.e., $M_1 \Delta A_{\Sigma 1} = M_2 \Delta A_{\Sigma 2}$. Similarly, when N = 3, the fitting parameters are $A_{\Sigma 1}$ and $A_{\Sigma 2}$. The other parameter, $A_{\Sigma 3}$, is determined via $(M_3 \Delta A_{\Sigma 3})^2 = (M_1 \Delta A_{\Sigma 1})^2 + (M_2 \Delta A_{\Sigma 2})^2 + 2M_1 M_2 (\Gamma^{-1})_{A_{\Sigma 1} A_{\Sigma 2}}$. To obtain the derivative with respect to $A_{\Sigma i}$ in the Fisher matrix, we numerically calculate the derivatives of the potential in Eq. (4) and the amplification factor in Eq. (6) by changing $A_{\Sigma i}$ by ± 1 %.

Figure 11 plots the measurement accuracies of $A_{\Sigma i}$ for N = 2. As the total mass is fixed, the lensed signal is insensitive to the density profile when the pulsar is outside the Sun. Therefore, the accuracies are not improved at $T \gtrsim 10$ hours. J1745-23 gives the best accuracy, because it has the highest frequency and smallest impact parameter. J1730-2304 has a higher frequency than J1022+1001, but its larger impact parameter lessens the constraint. In Case (II), $\Delta A_{\Sigma 2}$ is 17 times larger than

 $\Delta A_{\Sigma 1}$, reflecting the mass ratio $M_1/M_2 \simeq 17$. According to the figure, at least SNR $\approx 10^4 (T/\text{yr})^{1/2}$ is required for probing the solar density profile, but a lower SNR may be sufficient for probing with J1745-23. We comment that $\Delta \ln h_0$ and $\Delta \phi_0$ agree with SNR⁻¹ outside the Sun (corresponding to T > 10 hours; these results are not plotted in the figure). The numerical values of $\Delta A_{\Sigma i}$ are listed in Table 3.

Figure 12 plots the measurement accuracies of $A_{\Sigma i}$ for N = 3. In Case (I), $\Delta A_{\Sigma 3}$ shows the highest accuracy for most pulsars (the exception is J1730-2304), because the third annulus (with radius ranging from $0.26 \theta_{\odot}$ to θ_{\odot}) has the largest area. Therefore, the pulsar remains longer in the third annulus than in the other annuli. In Case (II), $\Delta A_{\Sigma 1} < \Delta A_{\Sigma 2} < \Delta A_{\Sigma 3}$ simply because the inner annulus is more massive than the outer one. The numerical values of $\Delta A_{\Sigma i}$ are listed in Table 4. According to Fig. 12, at least SNR $\approx 10^5 (T/\text{yr})^{1/2}$ is required for measuring the density profile with N = 3. The SNR must be further increased for larger N (> 3).

In geometrical optics, Φ_F is fully determined by the mass enclosed within a given radius. Therefore, a pulsar can probe the enclosed mass down to the impact parameter. In wave optics, Φ_F also depends on the outer density profile, owing to diffraction (Subsection 2.3). Therefore, the accuracy of $A_{\Sigma i}$ depends on the frequency and impact parameter in a complicated manner. Combining the results of different pulsars improves the measurement accuracy of the solar density and increases the radial range of the estimation.




FIG. 12.— Same as Fig. 11, but for three annuli with SNR = $10^5 (T/yr)^{1/2}$.

Figures 11 and 12 suggest that the accuracies strongly depend on the chosen boundary. In Case (I), the accuracies are similar in different annuli, but in Case (II), the accuracy is better in the inner annulus. This implies that the annulus mass mainly determines the accuracy. The boundary that maximizes the accuracy will depend on the frequency and impact parameter of the pulsar.

The solar density can (in principle) be plotted as a function of radius in the limit of large N (but a very high SNR is needed). As a demonstration, Fig. 13 plots the measurement accuracies for N = 6 and 10. Here, we follow the same procedure as done for N = 2and 3. As expected, J1745-23 gives the best accuracy. Though J1730-2304 has a large impact parameter (denoted by the down blue arrow), it can constrain the inner density due to the diffraction. The density of the outermost annulus is determined most precisely, because it has the largest area and the pulsars remain there for the longest time. The density of the inter-mediate annuli $(0.1 \lesssim \theta/\theta_{\odot} \lesssim 0.3)$ is poorly determined, because the errors of nearby annuli are strongly correlated. For instance, the cross-correlation coefficient of the accuracies in between the i- and j-th annuli, $r_{ij} \equiv (\Gamma^{-1})_{A_{\Sigma i}A_{\Sigma j}}/[(\Gamma^{-1})_{A_{\Sigma i}A_{\Sigma i}}(\Gamma^{-1})_{A_{\Sigma j}A_{\Sigma j}}]^{1/2}$, is large for a close pair of the annuli, $0.6 < |r_{ij}| < 1$ for |i-j| = 1.

The strain sensitivities of the planned detectors of the Cosmic Explorer $(CE)^9$ and the Einstein Telescope $(ET)^{10}$ will be approximately 100 times better than those

of the LIGO O3 run. The current upper limits in Table 1 were determined from observations taken over approximately one year in the O3 run. These limits correspond to an upper limit of SNR ≤ 10 (Appendix C of Abbott et al. 2019). Therefore, in CE and ET observations, the best SNR for these known pulsars is estimated as $\approx 10^3 (T/\text{yr})^{1/2}$ (a similar discussion is given in Section V of Marchant et al. 2020). Judging from these estimates, measuring the density profile using these pulsars will be a difficult task in the near future. To alleviate this problem, we require more pulsars than those detected by radio telescopes to date (as discussed in the next section). Moreover, these pulsars must have high frequencies and small impact parameters.

5. MILLISECOND PULSARS

This section roughly estimates the number of MSPs moving behind the Sun that will be found in the near future.

The present ATNF catalog lists approximately 3300 pulsars, including around 500 highly spinning pulsars with a spin period of < 10 ms (corresponding to f > 200 Hz). These pulsars are the so-called MSPs. The Milky Way is estimated to host 40000–90000 MSPs (e.g., Bhattacharyya & Roy 2022). As the sky fraction of the zodiac belt swept by the Sun is $2\pi \times 2\theta_{\odot}/(4\pi) \simeq 0.5\%$, the belt probably holds 200–400 MSPs. Almost none of these MSPs have been found, but tens of them will be identified in ongoing and upcoming radio surveys. For instance, the Canadian Hydrogen Intensity

⁹ https://cosmicexplorer.org/

¹⁰ https://www.et-gw.eu/




FIG. 13.— Measurement accuracies of the projected solar density as functions of radius. Solid black curve is the fiducial solar model and error bars are the accuracies for each pulsar. In the left and right panels, the density is divided into six and ten annuli in Case (I) with $SNR = 2 \times 10^6 (T/yr)^{1/2}$ and $10^7 (T/yr)^{1/2}$, respectively. Vertical dashed lines are the boundaries between the annuli. Down arrows indicate the ecliptic latitude of each pulsar.

TABLE 5GLOBULAR CLUSTERS CROSSING BEHIND THE SUN

name	$ heta_{ m LAT}/ heta_{\odot}$	$ heta_{ m h}/ heta_{\odot}$	$D_{\rm s}({\rm kpc})$	mass (M_{\odot})
NGC 6287 NGC 6717 NGC 6642	$0.49 \\ 0.50 \\ -0.89$	$\begin{array}{c} 0.046 \\ 0.043 \\ 0.046 \end{array}$	$7.9 \\ 7.5 \\ 8.1$	$\begin{array}{c} 1.5\times 10^5 \\ 3.6\times 10^4 \\ 3.4\times 10^4 \end{array}$

NOTE. — Listed are the ecliptic latitude of the cluster center θ_{LAT} , the half-light angular radius θ_{h} , the distance D_{s} , and the mass. Here, the distance and mass are taken from Baumgardt's catalog of globular clusters^a (e.g., Baumgardt & Hilker 2018).

^ahttps://people.smp.uq.edu.au/HolgerBaumgardt/globular/

Mapping Experiment (CHIME)¹¹, MeerKAT¹², and the Five-hundred-meter Aperture Spherical radio Telescope (FAST)¹³ will detect ≈ 1000 MSPs over the sky (Lorimer et al. 2019, their Table 1). These performances will be surpassed by the Square Kilometer Array (SKA) Phase 1, which will detect 1500 MSPs (Keane et al. 2015).

At present, approximately 220 MSPs have been found in 33 globular clusters¹⁴; \sim seven MSPs per cluster on average. Per unit mass, pulsars are two or three orders of magnitude more abundant in clusters than in the Galactic disk (Freire 2013). Therefore, many MSPs probably exist in clusters. Among the 157 clusters listed in the Mc-Master catalog of Milky Way globular clusters¹⁵ (version 2010, Harris 1996, 2010), three globular clusters move behind the Sun. The data are summarized in Table 5. Although no MSPs have been discovered in these clusters¹⁶, tens of MSPs are expected in each cluster.

6. LENSING BY A GALACTIC STAR

This section briefly discusses GW lensing by a Milky Way star along the line of sight to the pulsar. As a star crosses in front of the pulsar, lensing imprints a time-dependent modulation on the waveform. This event might be confused with the solar lensing. We comment that these two events can be distinguished due to the following reasons. First, the stellar lensing is observed only once, but the solar lensing is observed every year on the same day. In addition, the probability of the stellar lensing is not high, $\approx 10^{-6} \text{ yr}^{-1}$, on average over the full sky (e.g., Han 2008). If the pulsar is in a globular cluster, the probability of lensing by a cluster star (so-called self-lensing) can reach $\approx 0.1 \text{ yr}^{-1}$ (Kıroğlu et al. 2022). Second, the lensing modulation by the star (modeled as a point mass) differs from the solar modulation (e.g., Figs. 3 and 5 of Marchant et al. 2020). For example, the typical event duration is ≈ 10 days (e.g., Schneider et al. 2006), which is much longer than the solar lensing duration \approx half a day. Third, if radio observations are available for the stellar lensing, double signals (with different amplitudes and arrival times) from the single pulsar may be detected.

7. CONCLUSIONS

We have studied the detectability of solar density profiling using GW lensing with known pulsars. After selecting suitable pulsars with high frequencies and small impact parameters to the Sun from the ATNF catalog, we calculated the measurement accuracy of the overall amplitude of the solar gravitational potential using a Fisher analysis. The lensing signature can be detected with 3σ confidence when SNR $\approx 100 (f/300 \text{ Hz})^{-1}$ during one year of observation (Table 2 and Eq. (21)). The detection is therefore improved with high-frequency pulsars. The signature can be detected even if the pulsar trajectory does not pass behind the Sun (in such cases, the impact parameter to the Sun does not significantly affect the detectability). We found that the detectability is degraded by parameter degeneracy with the constant

=

¹¹ https://chime-experiment.ca/en

¹² https://www.sarao.ac.za/gallery/meerkat/

¹³ https://fast.bao.ac.cn

¹⁴ https://www3.mpifr-bonn.mpg.de/staff/pfreire/GCpsr. html

¹⁵ Data may be downloaded from https://physics.mcmaster. ca/Fac_Harris/mwgc.dat

¹⁶ https://www3.mpifr-bonn.mpg.de/staff/pfreire/GCpsr. html

phase (ϕ_0 in Eq. (16)) of the waveform. Next, we divided the projected density profile into N annuli ($N = 2, 3, \cdots$) and calculated the measurement accuracy of each annulus mass. If three known pulsars move behind the Sun with f > 100 Hz, a high SNR ($\gtrsim 10^4$) in a year-long observation is required for measuring the density profile in the two-annulus case (Fig. 11). The SNR must be raised to $\gtrsim 10^5$ in the three-annulus case (Fig. 12) and $\gtrsim 10^6$ in the six-annulus case (Fig. 13).

Currently, only two known MSPs move behind the Sun, but 200–400 can potentially exist within our Galaxy (Section 5). Ongoing and future radio surveys such as SKA, CHIME, and FAST will find tens of these MSPs in the coming decades.

The solar density profile has already been measured to percent-level precision based on global helioseismology (see the recent review by Basu 2016). GW lensing will likely become a complementary and independent method for probing the Sun in the future.

ACKNOWLEDGMENTS

We thank Hideyuki Hotta for his useful comment on helioseismology. We thank the CASS (CSIRO (Commonwealth Scientific and Industrial Research Organisation) Astronomy and Space Science) pulsar group for developing and maintaining the ATNF pulsar catalog. This work is supported in part by MEXT Japan and JSPS KAKENHI Grant Numbers of JP22H00130 (RT), JP20H05855 (RT), JP17H06359 (TS), JP21H05453 (TS), and JP19K03864 (TS).

APPENDIX

VALIDITY OF THE THIN-LENS APPROXIMATION

Suyama et al. (2005) investigated the validity of the thin-lens approximation in wave optics. They considered plane waves entering a lens and numerically solved the wave equation to obtain the exact lensed waveform (without the thinlens approximation). They considered three spherical lens models (a uniform density sphere, a singular isothermal sphere truncated at a finite radius, and the Hernquist profile). They found that the relative error of F is smaller than the lens radius over $D_{\rm L}$ and the maximum error is achieved when the wavelength is comparable to the Schwarzschild radius of the lens. This suggests that the error of F is $\leq R_{\odot}/(1 \, {\rm au}) \simeq 0.5 \%$ in our case.

REFERENCES

- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2016,
- Phys. Rev. Lett., 116, 061102
- —. 2017, ApJ, 839, 12
- —. 2019, ApJ, 879, 10
- Abbott, R., Abbott, T. D., Abraham, S., et al. 2021,
- Phys. Rev. D, 103, 064017
- Abbott, R., Abe, H., Acernese, F., et al. 2022a, ApJ, 935, 1
- —. 2022b, Phys. Rev. D, 106, 042003
- Abdo, A. A., Ackermann, M., Ajello, M., et al. 2009, Science, 325, 840
- Backer, D. C., & Hellings, R. W. 1986, ARA&A, 24, 537
- Bahcall, J. N., Serenelli, A. M., & Basu, S. 2005, ApJ, 621, L85
- Bartelmann, M., & Schneider, P. 2001, Phys. Rep., 340, 291
- Basu, S. 2016, Living Reviews in Solar Physics, 13, 2
- Baumgardt, H., & Hilker, M. 2018, MNRAS, 478, 1520
- Bhattacharyya, B., & Roy, J. 2022, Astrophysics and Space Science Library, 465, 1
- Bliokh, P. V., & Minakov, A. A. 1975, Ap&SS, 34, L7
- Bontz, R. J., & Haugan, M. P. 1981, Ap&SS, 78, 199
- Cameron, A. D., Champion, D. J., Bailes, M., et al. 2020, MNRAS, 493, 1063
- Camilo, F., Nice, D. J., Shrauner, J. A., & Taylor, J. H. 1996, ApJ, 469, 819
- Choi, H. G., Park, C., & Jung, S. 2021, Phys. Rev. D, 104, 063001
- Cutler, C. 1998, Phys. Rev. D, 57, 7089
- Cutler, C., & Flanagan, É. E. 1994, Phys. Rev. D, 49, 2658
- Cyranski, J. F., & Lubkin, E. 1974, Annals of Physics, 87, 205
- Dai, L., Li, S.-S., Zackay, B., Mao, S., & Lu, Y. 2018,
- Phys. Rev. D, 98, 104029 Dalang, C., Cusin, G., & Lagos, M. 2022, Phys. Rev. D, 105,
- 024005
- Edwards, R. T., Hobbs, G. B., & Manchester, R. N. 2006, MNRAS, 372, 1549
- Ezquiaga, J. M., Holz, D. E., Hu, W., Lagos, M., & Wald, R. M. 2021, Phys. Rev. D, 103, 064047
- Faulkner, A. J., Stairs, I. H., Kramer, M., et al. 2004, MNRAS, 355, 147
- Ferdman, R. D., Stairs, I. H., Kramer, M., et al. 2014, MNRAS, 443, 2183
- Freire, P. C. C. 2013, Neutron Stars and Pulsars: Challenges and Opportunities after 80 years, 291, 243
- Glampedakis, K., & Gualtieri, L. 2018, Astrophysics and Space Science Library, 457, 673

- Guo, X., & Lu, Y. 2020, Phys. Rev. D, 102, 124076
- Han, C. 2008, ApJ, 681, 806
- Harris, W. E. 1996, AJ, 112, 1487
- —. 2010, arXiv:1012.3224
- Hobbs, G. B., Edwards, R. T., & Manchester, R. N. 2006, MNRAS, 369, 655
- Hou, S., Fan, X.-L., & Zhu, Z.-H. 2019, Phys. Rev. D, 100, 064028 Jung, S., & Kim, S. 2022, arXiv:2210.02649
- Keane, E., Bhattacharyya, B., Kramer, M., et al. 2015, Advancing Astrophysics with the Square Kilometre Array (AASKA14), 40 Keith, M. J., Jameson, A., van Straten, W., et al. 2010, MNRAS,
- 409, 619 Kıroğlu, F., Weatherford, N. C., Kremer, K., et al. 2022, ApJ,
- 928, 181 Liao, K., Biesiada, M., & Zhu, Z.-H. 2022, Chinese Physics
- Liao, K., Biesiada, M., & Zhu, Z.-H. 2022, Chinese Physics Letters, 39, 119801
- Lorimer, D., Pol, N., Rajwade, K., et al. 2019, BAAS, 51, 261
- Lorimer, D. R., Nicastro, L., Lyne, A. G., et al. 1995, ApJ, 439, 933
- Lynch, R. S., Ransom, S. M., Freire, P. C. C., & Stairs, I. H. 2011, ApJ, 734, 89
- Macquart, J. P. 2004, A&A, 422, 761
- Manchester, R. N., Hobbs, G. B., Teoh, A., & Hobbs, M. 2005, AJ, 129, 1993
- Marchant, P., Breivik, K., Berry, C. P. L., Mandel, I., & Larson, S. L. 2020, Phys. Rev. D, 101, 024039
- Nakamura, T. T. 1998, Phys. Rev. Lett., 80, 1138
- Nakamura, T. T., & Deguchi, S. 1999, Progress of Theoretical Physics Supplement, 133, 137
- Ng, C., Guillemot, L., Freire, P. C. C., et al. 2020, MNRAS, 493, 1261
- Nice, D., Demorest, P., Stairs, I., et al. 2015, Tempo: Pulsar
- timing data analysis, Astrophysics Source Code Library, record ascl:1509.002
- Oguri, M. 2019, Reports on Progress in Physics, 82, 126901
- Oguri, M., & Takahashi, R. 2020, ApJ, 901, 58
- Ohanian, H. C. 1974, International Journal of Theoretical Physics, 9, 425
- Park, R. S., Folkner, W. M., Konopliv, A. S., et al. 2017, AJ, 153, 121
- Patla, B., & Nemiroff, R. J. 2008, ApJ, 685, 1297
- Ray, P. S., Kerr, M., Parent, D., et al. 2011, ApJS, 194, 17
- Ray, P. S., Abdo, A. A., Parent, D., et al. 2012, arXiv:1205.3089

- Reardon, D. J., Shannon, R. M., Cameron, A. D., et al. 2021, MNRAS, 507, 2137
- Riles, K. 2022, arXiv:2206.06447
- Roy, J., & Bhattacharyya, B. 2013, ApJ, 765, L45
- Sanpa-Arsa, S. 2016, PhD thesis, University of Virginia
- Schneider, P., Ehlers, J., & Falco, E. E. 1992, Gravitational Lenses (Springer-Verlag, Berlin)
 Schneider, P., Kochanek, C. S., & Wambsganss, J. 2006,
- Schneider, P., Kochanek, C. S., & Wambsganss, J. 2006 Gravitational Lensing: Strong, Weak and Micro (Springer-Verlag, Berlin)
- Sonnabend, D. 1979, PhD thesis, Stanford University, California Stix, M. 2004, The Sun : an Introduction (Springer-Verlag, Berlin)
- Suvorov, A. G. 2022, ApJ, 930, 13

- Suyama, T., Takahashi, R., & Michikoshi, S. 2005, Phys. Rev. D, 72, 043001
- Takahashi, R. 2004, A&A, 423, 787
- —. 2006, ApJ, 644, 80
- Takahashi, R., & Nakamura, T. 2003, ApJ, 595, 1039
- The LIGO Scientific Collaboration, the Virgo Collaboration, the KAGRA Collaboration, et al. 2021, arXiv:2111.03606
- —. 2022, arXiv:2201.00697
 Vinyoles, N., Serenelli, A. M., Villante, F. L., et al. 2017, ApJ, 835, 202
- Yao, J. M., Manchester, R. N., & Wang, N. 2017, ApJ, 835, 29