

First Observation of a Three-Resonance Structure in $e^+e^- \rightarrow$ Nonopen Charm Hadrons

M. Ablikim¹, M. N. Achasov^{5,b}, P. Adlarson⁷⁵, X. C. Ai⁸¹, R. Aliberti³⁶, A. Amoroso^{74A,74C}, M. R. An⁴⁰, Q. An^{71,58}, Y. Bai⁵⁷, O. Bakina³⁷, I. Balossino^{30A}, Y. Ban^{47,g}, V. Batozskaya^{1,45}, K. Begzsuren³³, N. Berger³⁶, M. Berlowski⁴⁵, M. Bertani^{29A}, D. Bettini^{30A}, F. Bianchi^{74A,74C}, E. Bianco^{74A,74C}, A. Bortone^{74A,74C}, I. Boyko³⁷, R. A. Briere⁶, A. Brueggemann⁶⁸, H. Cai⁷⁶, X. Cai^{1,58}, A. Calcaterra^{29A}, G. F. Cao^{1,63}, N. Cao^{1,63}, S. A. Cetin^{62A}, J. F. Chang^{1,58}, T. T. Chang⁷⁷, W. L. Chang^{1,63}, G. R. Che⁴⁴, G. Chelkov^{37,a}, C. Chen⁴⁴, Chao Chen⁵⁵, G. Chen¹, H. S. Chen^{1,63}, M. L. Chen^{1,58,63}, S. J. Chen⁴³, S. M. Chen⁶¹, T. Chen^{1,63}, X. R. Chen^{32,63}, X. T. Chen^{1,63}, Y. B. Chen^{1,58}, Y. Q. Chen³⁵, Z. J. Chen^{26,h}, W. S. Cheng^{74C}, S. K. Choi^{11A}, X. Chu⁴⁴, G. Cibinetto^{30A}, S. C. Coen⁴, F. Cossio^{74C}, J. J. Cui⁵⁰, H. L. Dai^{1,58}, J. P. Dai⁷⁹, A. Dbeyssi¹⁹, R. E. de Boer⁴, D. Dedovich³⁷, Z. Y. Deng¹, A. Denig³⁶, I. Denysenko³⁷, M. Destefanis^{74A,74C}, F. De Mori^{74A,74C}, B. Ding^{66,1}, X. X. Ding^{47,g}, Y. Ding⁴¹, Y. Ding³⁵, J. Dong^{1,58}, L. Y. Dong^{1,63}, M. Y. Dong^{1,58,63}, X. Dong⁷⁶, M. C. Du¹, S. X. Du⁸¹, Z. H. Duan⁴³, P. Egorov^{37,a}, Y. L. Fan⁷⁶, J. Fang^{1,58}, S. S. Fang^{1,63}, W. X. Fang¹, Y. Fang¹, R. Farinelli^{30A}, L. Fava^{74B,74C}, F. Feldbauer⁴, G. Felici^{29A}, C. Q. Feng^{71,58}, J. H. Feng⁵⁹, K. Fischer⁶⁹, M. Fritsch⁴, C. Fritschsch⁶⁸, C. D. Fu¹, J. L. Fu⁶³, Y. W. Fu¹, H. Gao⁶³, Y. N. Gao^{47,g}, Yang Gao^{71,58}, S. Garbolino^{74C}, I. Garzia^{30A,30B}, P. T. Ge⁷⁶, Z. W. Ge⁴³, C. Geng⁵⁹, E. M. Gersabeck⁶⁷, A. Gilman⁶⁹, K. Goetzen¹⁴, L. Gong⁴¹, W. X. Gong^{1,58}, W. Gradl³⁶, S. Gramigna^{30A,30B}, M. Greco^{74A,74C}, M. H. Gu^{1,58}, Y. T. Gu¹⁶, C. Y. Guan^{1,63}, Z. L. Guan²³, A. Q. Guo^{32,63}, L. B. Guo⁴², M. J. Guo⁵⁰, R. P. Guo⁴⁹, Y. P. Guo^{13,f}, A. Guskov^{37,a}, T. T. Han⁵⁰, W. Y. Han⁴⁰, X. Q. Hao²⁰, F. A. Harris⁶⁵, K. K. He⁵⁵, K. L. He^{1,63}, F. H. He⁵⁵, Z. L. Hou¹, H. H. Heinsius⁴, C. H. Heinz³⁶, Y. K. Heng^{1,58,63}, C. Herold⁶⁰, T. Holtmann⁴, P. C. Hong^{13,f}, G. Y. Hou^{1,63}, X. T. Hou^{1,63}, Y. R. Hou⁶³, Z. L. Hou¹, H. M. Hu^{1,63}, J. F. Hu^{56,i}, T. Hu^{1,58,63}, Y. Hu¹, G. S. Huang^{71,58}, K. X. Huang⁵⁹, L. Q. Huang^{32,63}, X. T. Huang⁵⁰, Y. P. Huang¹, T. Hussain⁷³, N. Hüskens^{28,36}, W. Imoehl²⁸, M. Irshad^{71,58}, J. Jackson²⁸, S. Jaeger⁴, S. Janchiv³³, J. H. Jeong^{11A}, Q. Ji¹, Q. P. Ji²⁰, X. B. Ji^{1,63}, X. L. Ji^{1,58}, Y. Y. Ji⁵⁰, X. Q. Jia⁵⁰, Z. K. Jia^{71,58}, H. J. Jiang⁷⁶, L. L. Jiang¹, P. C. Jiang^{47,g}, S. S. Jiang⁴⁰, T. J. Jiang¹⁷, X. S. Jiang^{1,58,63}, Y. Jiang⁶³, J. B. Jiao⁵⁰, Z. Jiao²⁴, S. Jin⁴³, Y. Jin⁶⁶, M. Q. Jing^{1,63}, T. Johansson⁷⁵, X. Kui¹, S. Kabana³⁴, N. Kalantar-Nayestanaki⁶⁴, X. L. Kang¹⁰, X. S. Kang⁴¹, R. Kappert⁶⁴, M. Kavatsyuk⁶⁴, B. C. Ke⁸¹, A. Khokkaz⁶⁸, R. Kiuchi¹, R. Kliemt¹⁴, O. B. Kolcu^{62A}, B. Kopf⁴, M. K. Kuessner⁴, A. Kupsc^{45,75}, W. Kühn³⁸, J. J. Lane⁶⁷, P. Larin¹⁹, A. Lavanina²⁷, L. Lavezzi^{74A,74C}, T. T. Lei^{71,k}, Z. H. Lei^{71,58}, H. Leithoff³⁶, M. Lellmann³⁶, T. Lenz³⁶, C. Li⁴⁴, C. Li⁴⁸, C. H. Li⁴⁰, Cheng Li^{71,58}, D. M. Li⁸¹, F. Li^{1,58}, G. Li¹, H. Li^{71,58}, H. B. Li^{1,63}, H. J. Li²⁰, H. N. Li^{56,i}, Hui Li⁴⁴, J. R. Li⁶¹, J. S. Li⁵⁹, J. W. Li⁵⁰, K. L. Li²⁰, Ke Li¹, L. J. Li^{1,63}, L. K. Li¹, Lei Li³, M. H. Li⁴⁴, P. R. Li^{39,j,k}, Q. X. Li⁵⁰, S. X. Li¹³, T. Li⁵⁰, W. D. Li^{1,63}, W. G. Li¹, X. H. Li^{71,58}, X. L. Li⁵⁰, Xiaoyu Li^{1,63}, Y. G. Li^{47,g}, Z. J. Li⁵⁹, Z. X. Li¹⁶, C. Liang⁴³, H. Liang³⁵, H. Liang^{1,63}, H. Liang^{71,58}, Y. F. Liang⁵⁴, Y. T. Liang^{32,63}, G. R. Liao¹⁵, L. Z. Liao⁵⁰, Y. P. Liao^{1,63}, J. Libby²⁷, A. Limphirat⁶⁰, D. X. Lin^{32,63}, T. Lin¹, B. J. Liu¹, B. X. Liu⁷⁶, C. Liu³⁵, C. X. Liu¹, F. H. Liu⁵³, Fang Liu¹, Feng Liu⁷, G. M. Liu^{56,i}, H. Liu^{39,j,k}, H. B. Liu¹⁶, H. M. Liu^{1,63}, Huanhuan Liu¹, Huihui Liu²², J. B. Liu^{71,58}, J. L. Liu⁷², J. Y. Liu^{1,63}, K. Liu¹, K. Y. Liu⁴¹, Ke Liu²³, L. Liu^{71,58}, L. C. Liu⁴⁴, Lu Liu⁴⁴, M. H. Liu^{13,f}, P. L. Liu¹, Q. Liu⁶³, S. B. Liu^{71,58}, T. Liu^{13,f}, W. H. Liu⁴⁴, W. M. Liu^{71,58}, X. Liu^{39,j,k}, Y. Liu^{39,j,k}, Y. Liu⁸¹, Y. B. Liu⁴⁴, Z. A. Liu^{1,58,63}, Z. Q. Liu⁵⁰, X. C. Lou^{1,58,63}, F. X. Lu⁵⁹, H. J. Lu²⁴, J. G. Lu^{1,58}, X. L. Lu¹, Y. Lu⁸, Y. P. Lu^{1,58}, Z. H. Lu^{1,63}, C. L. Luo⁴², M. X. Luo⁸⁰, T. Luo^{13,f}, X. L. Luo^{1,58}, X. R. Lyu⁶³, Y. F. Lyu⁴⁴, F. C. Ma⁴¹, H. L. Ma¹, J. L. Ma^{1,63}, L. L. Ma⁵⁰, M. M. Ma^{1,63}, Q. M. Ma¹, R. Q. Ma^{1,63}, R. T. Ma⁶³, X. Y. Ma^{1,58}, Y. Ma^{47,g}, Y. M. Ma³², F. E. Maas¹⁹, M. Maggiora^{74A,74C}, S. Malde⁶⁹, Q. A. Malik⁷³, A. Mangoni^{29B}, Y. J. Mao^{47,g}, Z. P. Mao¹, S. Marcello^{74A,74C}, Z. X. Meng⁶⁶, J. G. Messchendorp^{14,64}, G. Mezzadri^{30A}, H. Miao^{1,63}, T. J. Min⁴³, R. E. Mitchell²⁸, X. H. Mo^{1,58,63}, N. Yu. Muchnoi^{5,b}, Y. Nefedov³⁷, F. Nerling^{19,d}, I. B. Nikolaev^{5,b}, Z. Ning^{1,58}, S. Nisar^{12,l}, Y. Niu⁵⁰, S. L. Olsen⁶³, Q. Ouyang^{1,58,63}, S. Pacetti^{29B,29C}, X. Pan⁵⁵, Y. Pan⁵⁷, A. Pathak³⁵, P. Patteri^{29A}, Y. P. Pei^{71,58}, M. Pelizaeus⁴, H. P. Peng^{71,58}, K. Peters^{14,d}, J. L. Ping⁴², R. G. Ping^{1,63}, S. Plura³⁶, S. Pogodin³⁷, V. Prasad³⁴, F. Z. Qi¹, H. Qi^{71,58}, H. R. Qi⁶¹, M. Qi⁴³, T. Y. Qi^{13,f}, S. Qian^{1,58}, W. B. Qian⁶³, C. F. Qiao⁶³, J. J. Qin⁷², L. Q. Qin¹⁵, X. P. Qin^{13,f}, X. S. Qin⁵⁰, Z. H. Qin^{1,58}, J. F. Qiu¹, S. Q. Qu⁶¹, C. F. Redmer³⁶, K. J. Ren⁴⁰, A. Rivetti^{74C}, V. Rodin⁶⁴, M. Rolo^{74C}, G. Rong^{1,63}, Ch. Rosner¹⁹, S. N. Ruan⁴⁴, N. Salone⁴⁵, A. Sarantsev^{37,c}, Y. Schelhaas³⁶, K. Schoenning⁷⁵, M. Scodreggio^{30A,30B}, K. Y. Shan^{13,f}, W. Shan²⁵, X. Y. Shan^{71,58}, J. F. Shangguan⁵⁵, L. G. Shao^{1,63}, M. Shao^{71,58}, C. P. Shen^{13,f}, H. F. Shen^{1,63}, W. H. Shen⁶³, X. Y. Shen^{1,63}, B. A. Shi⁶³, H. C. Shi^{71,58}, J. L. Shi¹³, J. Y. Shi¹, Q. Q. Shi⁵⁵, R. S. Shi^{1,63}, X. Shi^{1,58}, J. J. Song²⁰, T. Z. Song⁵⁹, W. M. Song^{35,1}, Y. J. Song¹³, Y. X. Song^{47,g}, S. Sosio^{74A,74C}, S. Spataro^{74A,74C}, F. Stieler³⁶, Y. J. Su⁶³, G. B. Sun⁷⁶, G. X. Sun¹, H. Sun⁶³, H. K. Sun¹, J. F. Sun²⁰, K. Sun⁶¹, L. Sun⁷⁶, S. S. Sun^{1,63}, T. Sun^{1,63}, W. Y. Sun³⁵, Y. Sun¹⁰, Y. J. Sun^{71,58}, Y. Z. Sun¹, Z. T. Sun⁵⁰, Y. X. Tan^{71,58}, C. J. Tang⁵⁴, G. Y. Tang¹, J. Tang⁵⁹, Y. A. Tang⁷⁶, L. Y. Tao⁷², Q. T. Tao^{26,h}, M. Tat⁶⁹, J. X. Teng^{71,58}, V. Thoren⁷⁵, W. H. Tian⁵⁹, W. H. Tian⁵², Y. Tian^{32,63}, Z. F. Tian⁷⁶, I. Uman^{62B}, S. J. Wang⁵⁰, B. Wang¹, B. L. Wang⁶³, Bo Wang^{71,58}, C. W. Wang⁴³, D. Y. Wang^{47,g}, F. Wang⁷², H. J. Wang^{39,j,k}, H. P. Wang^{1,63}, J. P. Wang⁵⁰, K. Wang^{1,58}, L. L. Wang¹, M. Wang⁵⁰, Meng Wang^{1,63}, S. Wang^{39,j,k}, S. Wang^{13,f}, T. Wang^{13,f}, T. J. Wang⁴⁴, W. Wang⁷², W. Wang⁵⁹, W. P. Wang^{71,58}, X. Wang^{47,g}, X. F. Wang^{39,j,k}, X. J. Wang⁴⁰, X. L. Wang^{13,f}, Y. Wang⁶¹, Y. D. Wang⁴⁶, Y. F. Wang^{1,58,63}, Y. H. Wang⁴⁸, Y. N. Wang⁴⁶, Y. Q. Wang¹, Yaqian Wang^{18,1}, Yi Wang⁶¹, Z. Wang^{1,58}, Z. L. Wang⁷², Z. Y. Wang^{1,63}, Ziyi Wang⁶³, D. Wei⁷⁰, D. H. Wei¹⁵, F. Weidner⁶⁸, S. P. Wen¹, C. W. Wenzel⁴, U. W. Wiedner⁴, G. Wilkinson⁶⁹, M. Wolke⁷⁵, L. Wollenberg⁴, C. Wu⁴⁰, J. F. Wu^{1,63}, L. H. Wu¹, L. J. Wu^{1,63}, X. Wu^{13,f}, X. H. Wu³⁵, Y. Wu⁷¹, Y. J. Wu³², Z. Wu^{1,58}, L. Xia^{71,58}, X. M. Xian⁴⁰, T. Xiang^{47,g}, D. Xiao^{39,j,k}, G. Y. Xiao⁴³, H. Xiao^{13,f}, S. Y. Xiao¹, Y. L. Xiao^{13,f}, Z. J. Xiao⁴², C. Xie⁴³, X. H. Xie^{47,g}, Y. Xie⁵⁰, Y. G. Xie^{1,58}, Y. H. Xie⁷, Z. P. Xie^{71,58}, T. Y. Xing^{1,63}, C. F. Xu^{1,63}, C. J. Xu⁵⁹, G. F. Xu¹, H. Y. Xu⁶⁶, Q. J. Xu¹⁷, Q. N. Xu³¹, W. Xu^{1,63}, W. L. Xu⁶⁶, X. P. Xu⁵⁵, Y. C. Xu⁷⁸, Z. P. Xu⁴³, Z. S. Xu⁶³, F. Yan^{13,f}, L. Yan^{13,f}, W. B. Yan^{71,58}, W. C. Yan⁸¹, X. Q. Yan¹, H. J. Yang^{51,e}, H. L. Yang³⁵, H. X. Yang¹, Tao Yang¹, Yang^{13,f}, Y. F. Yang⁴⁴, Y. X. Yang^{1,63}, Yifan Yang^{1,63}, Z. W. Yang^{39,j,k}, Z. P. Yao⁵⁰, M. Ye^{1,58}, M. H. Ye⁹, J. H. Yin¹, Z. Y. You⁵⁹, B. X. Yu^{1,58,63}, C. X. Yu⁴⁴, G. Yu^{1,63}, J. S. Yu^{26,h}, T. Yu⁷², X. D. Yu^{47,g}, C. Z. Yuan^{1,63}, L. Yuan², S. C. Yuan¹, X. Q. Yuan¹, Y. Yuan^{1,63}, Z. Y. Yuan⁵⁹, C. X. Yue⁴⁰, A. A. Zafar⁷³, F. R. Zeng⁵⁰, X. Zeng^{13,f}, Y. Zeng^{26,h}, Y. J. Zeng^{1,63}, X. Y. Zhai³⁵, Y. C. Zhai⁵⁰, Y. H. Zhan⁵⁹, A. Q. Zhang^{1,63}, B. L. Zhang^{1,63}, B. X. Zhang¹, D. H. Zhang⁴⁴, G. Y. Zhang²⁰, H. Zhang⁷¹, H. H. Zhang⁵⁹, H. H. Zhang³⁵, H. Q. Zhang^{1,58,63}, H. Y. Zhang^{1,58}, J. J. Zhang⁵², J. L. Zhang²¹, J. Q. Zhang⁴², J. W. Zhang^{1,58,63}, J. X. Zhang^{39,j,k}, J. Y. Zhang¹, J. Z. Zhang^{1,63}, Jianyu Zhang⁶³, Jiawei Zhang^{1,63}, L. M. Zhang⁶¹, L. Q. Zhang⁵⁹, Lei Zhang⁴³, P. Zhang¹, Q. Y. Zhang^{40,81}, Shuihan Zhang^{1,63}, Shulei Zhang^{26,h}, X. D. Zhang⁴⁶, X. M. Zhang¹, X. Y. Zhang⁵⁰, Xuyan Zhang⁵⁵, Y. Zhang⁷², Y. Zhang⁶⁹, Y. T. Zhang⁸¹, Y. H. Zhang^{1,58}, Yan Zhang^{71,58}, Yao Zhang¹, Z. H. Zhang¹, Z. L. Zhang³⁵, Z. Y. Zhang⁴⁴, Z. Y. Zhang⁷⁶

G. Zhao¹, J. Zhao⁴⁰, J. Y. Zhao^{1,63}, J. Z. Zhao^{1,58}, Lei Zhao^{71,58}, Ling Zhao¹, M. G. Zhao⁴⁴, S. J. Zhao⁸¹, Y. B. Zhao^{1,58}, Y. X. Zhao^{32,63}, Z. G. Zhao^{71,58}, A. Zhemchugov^{37,a}, B. Zheng⁷², J. P. Zheng^{1,58}, W. J. Zheng^{1,63}, Y. H. Zheng⁶³, B. Zhong⁴², X. Zhong⁵⁹, H. Zhou⁵⁰, L. P. Zhou^{1,63}, X. Zhou⁷⁶, X. K. Zhou⁷, X. R. Zhou^{71,58}, X. Y. Zhou⁴⁰, Y. Z. Zhou^{13,f}, J. Zhu⁴⁴, K. Zhu¹, K. J. Zhu^{1,58,63}, L. Zhu³⁵, L. X. Zhu⁶³, S. H. Zhu⁷⁰, S. Q. Zhu⁴³, T. J. Zhu^{13,f}, W. J. Zhu^{13,f}, Y. C. Zhu^{71,58}, Z. A. Zhu^{1,63}, J. H. Zou¹, J. Zu^{71,58}

(BESIII Collaboration)

- ¹ Institute of High Energy Physics, Beijing 100049, People's Republic of China
² Beihang University, Beijing 100191, People's Republic of China
³ Beijing Institute of Petrochemical Technology, Beijing 102617, People's Republic of China
⁴ Bochum Ruhr-University, D-44780 Bochum, Germany
⁵ Budker Institute of Nuclear Physics SB RAS (BINP), Novosibirsk 630090, Russia
⁶ Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA
⁷ Central China Normal University, Wuhan 430079, People's Republic of China
⁸ Central South University, Changsha 410083, People's Republic of China
⁹ China Center of Advanced Science and Technology, Beijing 100190, People's Republic of China
¹⁰ China University of Geosciences, Wuhan 430074, People's Republic of China
¹¹ Chung-Ang University, Seoul, 06974, Republic of Korea
¹² COMSATS University Islamabad, Lahore Campus, Defence Road, Off Raiwind Road, 54000 Lahore, Pakistan
¹³ Fudan University, Shanghai 200433, People's Republic of China
¹⁴ GSI Helmholtzcentre for Heavy Ion Research GmbH, D-64291 Darmstadt, Germany
¹⁵ Guangxi Normal University, Guilin 541004, People's Republic of China
¹⁶ Guangxi University, Nanning 530004, People's Republic of China
¹⁷ Hangzhou Normal University, Hangzhou 310036, People's Republic of China
¹⁸ Hebei University, Baoding 071002, People's Republic of China
¹⁹ Helmholtz Institute Mainz, Staudinger Weg 18, D-55099 Mainz, Germany
²⁰ Henan Normal University, Xinxiang 453007, People's Republic of China
²¹ Henan University, Kaifeng 475004, People's Republic of China
²² Henan University of Science and Technology, Luoyang 471003, People's Republic of China
²³ Henan University of Technology, Zhengzhou 450001, People's Republic of China
²⁴ Huangshan College, Huangshan 245000, People's Republic of China
²⁵ Hunan Normal University, Changsha 410081, People's Republic of China
²⁶ Hunan University, Changsha 410082, People's Republic of China
²⁷ Indian Institute of Technology Madras, Chennai 600036, India
²⁸ Indiana University, Bloomington, Indiana 47405, USA
²⁹ INFN Laboratori Nazionali di Frascati, (a)INFN Laboratori Nazionali di Frascati, I-00044, Frascati, Italy; (b)INFN Sezione di Perugia, I-06100, Perugia, Italy; (C)University of Perugia, I-06100, Perugia, Italy
³⁰ INFN Sezione di Ferrara, (a)INFN Sezione di Ferrara, I-44122, Ferrara, Italy; (b)University of Ferrara, I-44122, Ferrara, Italy
³¹ Inner Mongolia University, Hohhot 010021, People's Republic of China
³² Institute of Modern Physics, Lanzhou 730000, People's Republic of China
³³ Institute of Physics and Technology, Peace Avenue 54B, Ulaanbaatar 13330, Mongolia
³⁴ Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica 1000000, Chile
³⁵ Jilin University, Changchun 130012, People's Republic of China
³⁶ Johannes Gutenberg University of Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany
³⁷ Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia
³⁸ Justus-Liebig-Universitaet Giessen, II. Physikalisches Institut, Heinrich-Buff-Ring 16, D-35392 Giessen, Germany
³⁹ Lanzhou University, Lanzhou 730000, People's Republic of China
⁴⁰ Liaoning Normal University, Dalian 116029, People's Republic of China
⁴¹ Liaoning University, Shenyang 110036, People's Republic of China
⁴² Nanjing Normal University, Nanjing 210023, People's Republic of China
⁴³ Nanjing University, Nanjing 210093, People's Republic of China
⁴⁴ Nankai University, Tianjin 300071, People's Republic of China
⁴⁵ National Centre for Nuclear Research, Warsaw 02-093, Poland
⁴⁶ North China Electric Power University, Beijing 102206, People's Republic of China
⁴⁷ Peking University, Beijing 100871, People's Republic of China
⁴⁸ Qufu Normal University, Qufu 273165, People's Republic of China
⁴⁹ Shandong Normal University, Jinan 250014, People's Republic of China
⁵⁰ Shandong University, Jinan 250100, People's Republic of China
⁵¹ Shanghai Jiao Tong University, Shanghai 200240, People's Republic of China
⁵² Shanxi Normal University, Linfen 041004, People's Republic of China
⁵³ Shanxi University, Taiyuan 030006, People's Republic of China
⁵⁴ Sichuan University, Chengdu 610064, People's Republic of China
⁵⁵ Soochow University, Suzhou 215006, People's Republic of China
⁵⁶ South China Normal University, Guangzhou 510006, People's Republic of China

- ⁵⁷ Southeast University, Nanjing 211100, People's Republic of China
- ⁵⁸ State Key Laboratory of Particle Detection and Electronics, Beijing 100049, Hefei 230026, People's Republic of China
- ⁵⁹ Sun Yat-Sen University, Guangzhou 510275, People's Republic of China
- ⁶⁰ Suranaree University of Technology, University Avenue 111, Nakhon Ratchasima 30000, Thailand
- ⁶¹ Tsinghua University, Beijing 100084, People's Republic of China
- ⁶² Turkish Accelerator Center Particle Factory Group, (a)Istinye University, 34010, Istanbul, Turkey; (b)Near East University, Nicosia, North Cyprus, 99138, Mersin 10, Turkey
- ⁶³ University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China
- ⁶⁴ University of Groningen, NL-9747 AA Groningen, The Netherlands
- ⁶⁵ University of Hawaii, Honolulu, Hawaii 96822, USA
- ⁶⁶ University of Jinan, Jinan 250022, People's Republic of China
- ⁶⁷ University of Manchester, Oxford Road, Manchester, M13 9PL, United Kingdom
- ⁶⁸ University of Muenster, Wilhelm-Klemm-Strasse 9, 48149 Muenster, Germany
- ⁶⁹ University of Oxford, Keble Road, Oxford OX13RH, United Kingdom
- ⁷⁰ University of Science and Technology Liaoning, Anshan 114051, People's Republic of China
- ⁷¹ University of Science and Technology of China, Hefei 230026, People's Republic of China
- ⁷² University of South China, Hengyang 421001, People's Republic of China
- ⁷³ University of the Punjab, Lahore-54590, Pakistan
- ⁷⁴ University of Turin and INFN, (a)University of Turin, I-10125, Turin, Italy; (b)University of Eastern Piedmont, I-15121, Alessandria, Italy; (c)INFN, I-10125, Turin, Italy
- ⁷⁵ Uppsala University, Box 516, SE-75120 Uppsala, Sweden
- ⁷⁶ Wuhan University, Wuhan 430072, People's Republic of China
- ⁷⁷ Xinyang Normal University, Xinyang 464000, People's Republic of China
- ⁷⁸ Yantai University, Yantai 264005, People's Republic of China
- ⁷⁹ Yunnan University, Kunming 650500, People's Republic of China
- ⁸⁰ Zhejiang University, Hangzhou 310027, People's Republic of China
- ⁸¹ Zhengzhou University, Zhengzhou 450001, People's Republic of China
- ^a Also at the Moscow Institute of Physics and Technology, Moscow 141700, Russia
- ^b Also at the Novosibirsk State University, Novosibirsk, 630090, Russia
- ^c Also at the NRC "Kurchatov Institute", PNPI, 188300, Gatchina, Russia
- ^d Also at Goethe University Frankfurt, 60323 Frankfurt am Main, Germany
- ^e Also at Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education; Shanghai Key Laboratory for Particle Physics and Cosmology; Institute of Nuclear and Particle Physics, Shanghai 200240, People's Republic of China
- ^f Also at Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) and Institute of Modern Physics, Fudan University, Shanghai 200443, People's Republic of China
- ^g Also at State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, People's Republic of China
- ^h Also at School of Physics and Electronics, Hunan University, Changsha 410082, China
- ⁱ Also at Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter, South China Normal University, Guangzhou 510006, China
- ^j Also at Frontiers Science Center for Rare Isotopes, Lanzhou University, Lanzhou 730000, People's Republic of China
- ^k Also at Lanzhou Center for Theoretical Physics, Lanzhou University, Lanzhou 730000, People's Republic of China
- ^l Also at the Department of Mathematical Sciences, IBA, Karachi 75270, Pakistan

We report the measurement of the inclusive cross sections for $e^+e^- \rightarrow \text{nOCH}$ (where nOCH denotes non-open charm hadrons) with improved precision at center-of-mass (c.m.) energies from 3.645 to 3.871 GeV. We observe three resonances: $\mathcal{R}(3760)$, $\mathcal{R}(3780)$, and $\mathcal{R}(3810)$ with significances of 8.1σ , 13.7σ , and 8.8σ , respectively. The $\mathcal{R}(3810)$ state is observed for the first time, while the $\mathcal{R}(3760)$ and $\mathcal{R}(3780)$ states are observed for the first time in the nOCH cross sections. Two sets of resonance parameters describe the energy-dependent line shape of the cross sections well. In set I [set II], the $\mathcal{R}(3810)$ state has mass $(3805.7 \pm 1.1 \pm 2.7)$ [(3805.7 \pm 1.1 \pm 2.7)] MeV/ c^2 , total width $(11.6 \pm 2.9 \pm 1.9)$ [(11.5 \pm 2.8 \pm 1.9)] MeV, and an electronic width multiplied by the nOCH decay branching fraction of $(10.9 \pm 3.8 \pm 2.5)$ [(11.0 \pm 3.4 \pm 2.5)] eV. In addition, we measure the branching fractions $\mathcal{B}[\mathcal{R}(3760) \rightarrow \text{nOCH}] = (25.2 \pm 16.1 \pm 30.4)\%[(6.4 \pm 4.8 \pm 7.7)\%]$ and $\mathcal{B}[\mathcal{R}(3780) \rightarrow \text{nOCH}] = (12.3 \pm 6.6 \pm 8.3)\%[(10.4 \pm 4.8 \pm 7.0)\%]$ for the first time. The $\mathcal{R}(3760)$ state can be interpreted as an open-charm (OC) molecular state, but containing a simple four-quark state component. The $\mathcal{R}(3810)$ state can be interpreted as a hadrocharmonium state.

Until two decades ago, it was widely believed that hadron resonances with masses higher than the open-charm (OC) pair thresholds decay entirely to OC final states via the strong interaction. However, in July, 2003, BES reported the obser-

vation of seven events of hadron resonance(s) in this mass regime decaying to nOCH [1–4]. This discovery overturned the understanding of resonance decays and opened up a new era in hadron spectroscopy. In this Letter, we denote these

resonances as X_{aboveOC} [3], which encompasses both supposed pure quark-antiquark $c\bar{c}$ states, i.e. $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$, and nonpure quark-antiquark $c\bar{c}$ states (hereafter referred to as ‘non- $c\bar{c}$ ’), such as four-quark states, OC-pair molecular states, hadrocharmonium states, and hybrid charmonium states [5–8]. Quantum chromodynamics (QCD) expects that the non- $c\bar{c}$ states exist in nature, and thus a discovery of these systems would be an important validation of the QCD predictions. The first nOCH final-state decay of a X_{aboveOC} resonance to be observed after the 2002 discoveries was $J/\psi\pi^+\pi^-$ [1, 2, 9], seen by BES-II. This final state can originate from a $c\bar{c}$ state, a non- $c\bar{c}$ state, or a combination of both [4]. This discovery stimulated strong interest in using $J/\psi\pi^+\pi^-$ or similar final states as golden channels to probe other nOC decays from the X_{aboveOC} , and led to the discovery of several exotic states [10–20].

Subsequent studies of the $\psi(3770)$ resonance showed that it has a branching fraction of about 15% into nOCH final states [21–24]. This fact indicates the contribution of some undiscovered states [25] with masses around $3.773 \text{ GeV}/c^2$, which predominantly decay into nOCH. In 2008, the BES-II experiment observed for the first time a double-peaked structure named the $\mathcal{R}_s(3770)$ in $e^+e^- \rightarrow \text{hadrons}$ at c.m. energies around 3.76 GeV [26], which is composed of two states labeled the $\mathcal{R}(3760)$ and $\mathcal{R}(3780)$. The existence of the $\mathcal{R}(3760)$ state was confirmed in $e^+e^- \rightarrow J/\psi X$ by BES-II [4]. It is seen that the study of the inclusive nOCH decays of X_{aboveOC} both helps the understanding of known states, and is a sensitive probe for undiscovered resonances, particularly non- $c\bar{c}$ states.

In this Letter, we report a measurement of the cross sections for $e^+e^- \rightarrow \text{nOCH}$ at c.m. energies from 3.645 to 3.871 GeV, studies of $\mathcal{R}(3760)$ and $\mathcal{R}(3780)$ production and decays, and a search for an additional resonant state in this energy region. The data samples used in this analysis were collected at 42 c.m. energies in 2010 and correspond to a total integrated luminosity of 75.5 pb^{-1} .

The BESIII detector [27] response is studied using Monte Carlo (MC) samples. The simulations are performed using a GEANT4-based [28] software package. Simulated samples for $q\bar{q}$ vector states (i.e. $u\bar{u}$, $d\bar{d}$, $s\bar{s}$, and $c\bar{c}$) and their decays to hadrons are generated using the MC event generators KKMC [29], EVTGEN [30], and LUNDCHARM [31]. Background sources are estimated with MC samples generated using KKMC, and the MC event generators BABAYAGA [32] and TWOGAM [33].

We select the inclusive nOCH events from the events with charged particles or charged and neutral particles. In order to reject background contributions from $e^+e^- \rightarrow (\gamma)e^+e^-$ and $e^+e^- \rightarrow (\gamma)\mu^+\mu^-$, we require the events to have more than two charged tracks (N_{CTrk}), and impose the following selection criteria for each track: (i) the distance (R_{xy}) of the point of the closest approach to the beam pipe must satisfy the condition $R_{xy} \leq 1.0 \text{ cm}$; (ii) the polar angle θ must satisfy




FIG. 1. Z_{AVRG} distribution of the vertices of the selected events from the data sample collected at $\sqrt{s} = 3.779 \text{ GeV}$, where the dots with error bar represent data, the solid line in blue is the best fit, and the dashed line in red is the fitted background shape.

fy the condition $|\cos\theta| < 0.93$; (iii) the momentum p must be less than $E_b + 0.02E_b\sqrt{1 + E_b^2}$, where E_b is the beam energy; (iv) the time-of-flight t_{TOF} must satisfy $2.0 < t_{\text{TOF}} < 20.0 \text{ ns}$ and $|t_{\text{TOF}} - t_p| < 2.0 \text{ ns}$, where t_p is the expected time-of-flight of protons; (v) the energy E_{EMC} deposited in the electromagnetic calorimeter (EMC) must be less than 1 GeV; (vi) the penetration depth in the muon-chamber system must be less than 30 cm.

For the selection of photons, we require the deposited energy of a neutral cluster in the EMC to be greater than 25 MeV in the barrel and 50 MeV in the end caps. We do not apply any requirements on the number of photons in the event. To suppress electronic noise and showers unrelated to the event, we impose the condition that the difference between the EMC time and the event start time be within $[0, 700] \text{ ns}$. To reduce the beam-associated events (beam interactions with gas or material), we demand that at least one charged track or photon must point into each hemisphere of $\cos\theta < 0$ and $\cos\theta > 0$. In addition, for each event, we require the total energy ($E_{\text{EMC}}^{\text{tot}}$) deposited in the EMC by the charged and neutral particles to be greater than $0.28E_b$.

Some beam-associated background sources still survive this selection. These background sources are produced at random z positions [21], while genuine nOCH events are produced around $z = 0$, where z is the distance to the interaction point along the BESIII axis. To distinguish the nOCH events from the background sources, we calculate the averaged z (Z_{AVRG}) of the charged tracks in each event. Figure 1 shows the distribution of the averaged Z_{AVRG} of the accepted events from the data sample collected at $\sqrt{s} = 3.779 \text{ GeV}$. Using a double-Gaussian function to describe the signal shape plus a second-order Chebychev function to parameterize the background shape, we fit the Z_{AVRG} distribution of event vertices to extract the number of nOCH candidates, $N_{\text{had}}^{\text{fit}}$ at each energy point.

The background to the $N_{\text{had}}^{\text{fit}}$ distribution comes from var-

ious sources, e.g. (i) $e^+e^- \rightarrow (\gamma)e^+e^-$, (ii) $e^+e^- \rightarrow (\gamma)\mu^+\mu^-$, (iii) $e^+e^- \rightarrow \gamma\gamma$, (iv) $e^+e^- \rightarrow (\gamma)\tau^+\tau^-$, (v) $e^+e^- \rightarrow (\gamma)e^+e^-\ell^+\ell^-$ ($\ell = e, \mu$ or τ), (vi) $e^+e^- \rightarrow (\gamma)e^+e^-X_{\text{had}}$ (where X_{had} denotes hadrons), and (vii) $e^+e^- \rightarrow (\gamma)D\bar{D}$. The total amount of background N_b at each c.m. energy \sqrt{s} is determined by $N_b = \sum_{i=1}^{i=7} N_{b,i}$ with $N_{b,i} = \mathcal{L} \times \sigma_{b,i} \times \eta_i$, where \mathcal{L} is the integrated luminosity of the data sample, $\sigma_{b,i}$ is the cross section for the i th background source and η_i is the probability of misidentifying a candidate from the i th background source as a nOCH event, which is determined by analyzing the large background MC samples. The cross sections σ_b for sources (i), (ii) and (iii) are taken from BABAYAGA. For source (iv), σ_b is calculated using the formulae given in Refs. [34, 35]. The cross sections for sources (v) and (vi) are taken from TWOGAM, while that for source (vii) is taken from the observed cross sections $\sigma_{D\bar{D}}^o(s)$ for $e^+e^- \rightarrow D\bar{D}$ determined using the same data samples. For example, at $\sqrt{s} = 3.7731$ GeV, $N_{\text{had}}^{\text{fit}} = (35235 \pm 225)$, $N_{b,\ell^+\ell^-, \gamma\gamma} = (1206 \pm 7)$, $N_{b,e^+e^-\ell^+\ell^-, e^+e^-X_{\text{had}}} = (341 \pm 6)$, and $N_{b,D\bar{D}} = (9458 \pm 186)$, where the uncertainties on the first two backgrounds arise from the statistical uncertainties on \mathcal{L} , $\sigma_{b,i}$, and η_i , and that on the third is due to the statistical uncertainty on $\sigma_{D\bar{D}}^o(s)$. To provide the most conservative signal significance for the resonance searches that are discussed below, we directly subtract N_b from $N_{\text{had}}^{\text{fit}}$ and allow the statistical uncertainty of $N_{b,D\bar{D}}$ to fully contribute to the uncertainty in the difference between $N_{\text{had}}^{\text{fit}}$ and N_b . This procedure yields $N_{\text{nOCH}}^o = (24230 \pm 292)$, where the uncertainty is statistical and includes the contribution from the background estimates.

We determine the detection efficiency using MC simulated events for the four components of the process $e^+e^- \rightarrow \text{nOCH}$: (i) $e^+e^- \rightarrow$ light-hadron (LH) continuum processes including lower-mass resonances (LMRs) with masses below 2 GeV (CPLMRs) (ii) $J/\psi \rightarrow$ hadrons, (iii) $\psi(3686) \rightarrow$ hadrons, and (iv) $\mathcal{R}_s(3770) \rightarrow \text{nOCH}$ [26]. We generate simulated samples for these events with the KKMC package. These events include initial-state radiation (ISR) and final-state radiation (FSR) processes. For the subsequent decays of the J/ψ , $\psi(3686)$ and $\mathcal{R}_s(3770)$, we use EVTGEN to generate the known final states, given in Ref. [36], and use LUNDCHARM to generate the remaining unknown final states. The resulting selection efficiency is determined using $\epsilon = \sum_1^4 w_i \epsilon_i$, where w_i is the number of nOCH events simulated for the i th component over the total number of MC simulated nOCH events for all four components, and ϵ_i is the corresponding efficiency for selection of the nOCH events from the i th component. Figure 2 (top) shows the efficiencies determined at the 42 energy points.

At $\sqrt{s} = 3.7731$ GeV, the efficiency is $\epsilon = 62.11\%$. The integrated luminosity corresponding to the data collected at this energy is $\mathcal{L} = (1831.63 \pm 4.49) \text{ nb}^{-1}$. Dividing the number $N_{\text{nOCH}}^o = (24230 \pm 292)$ by both the integrated luminosity and the efficiency yields the observed nOCH cross section $\sigma_{\text{nOCH}}^o = (21.299 \pm 0.262) \text{ nb}$, where the uncertainty aris-




FIG. 2. The efficiency ϵ (top) versus the c.m energy; the ISR correction factor $[1 + \delta(s)]$ (bottom) versus the c.m energy.

es from the statistical uncertainty on the N_{OCH}^o , the size of the MC samples, and the statistical uncertainty of the luminosity measurement. Similarly, we measure $\sigma_{\text{nOCH}}^o(s)$ at the remaining 41 energy points.

Table I summarizes the systematic uncertainties assigned to the $\sigma_{\text{nOCH}}^o(s)$ measurements. To determine the systematic uncertainties due to the choice of the selection criteria, we vary each criterion from its baseline value to an alternative setting, as given in Table I, and assign the resulting change in $\sigma_{\text{nOCH}}^o(s)$ as the systematic uncertainty. Removing the requirement that tracks must point into different hemispheres in z changes $\sigma_{\text{nOCH}}^o(s)$ by 0.80%, which is taken as the corresponding uncertainty associated with this possible source of bias. To estimate the uncertainty associated with the fit to the Z_{AVRG} distribution, we reperform the fit changing the background shape from a second-order to third-order, and then from a third-order to fourth-order Chebychev function. These result in 0.83% change in the cross sections, which is assigned as the corresponding systematic uncertainty. The uncertainty on the number of background sources is dominated by the estimation of the two-photon contribution, which induces a 0.75% uncertainty of the $\sigma_{\text{nOCH}}^o(s)$.

The choice of the MC generator for $e^+e^- \rightarrow \text{nOCH}$ impacts the selection efficiency. We recalculate the efficiency with different MC packages and take the 1.70% variation observed as the associated uncertainty. There is an uncertainty on $\sigma_{\text{nOCH}}^o(s)$ of 1.00% arising from the corresponding un-

certainty on the luminosity. Adding these contributions in quadrature yields a total systematic uncertainty of 2.89% on $\sigma_{\text{nOCH}}^{\text{o}}(s)$. This total does not include an energy-dependent uncertainty on $\sigma_{\text{nOCH}}^{\text{o}}(s)$, caused by the 1.43% uncertainty on the $\sigma_{D\bar{D}}^{\text{o}}(s)$ shape, which is accounted for when considering the systematic uncertainties on the fitted parameter values, as discussed below.

TABLE I. Systematic uncertainties on $\sigma_{\text{nOCH}}^{\text{o}}(s)$.

Source	Variation range	Uncertainty (%)
R_{xy} cut	[0.9, 1.1]	0.20
$\cos\theta$ cut	[0.875, 0.930]	0.40
$p < E_b + 0.02E_b\sqrt{1 + E_b}$	[nominal value, ∞]	0.01
t_{TOF} cut and $t_{\text{TOF}} - t_p$	[0, ∞] and [2, ∞]	0.09
E_{EMC} cut	[1, ∞]	0.39
Penetration depth	[30, ∞]	0.04
$E_{\text{EMC}}^{\text{tot}}$ cut	[0, 0.28]	0.34
N_{CTrk} cut	[2, 3]	1.45
Different hemisphere in z		0.80
Fitting Z_{AVRG} distribution		0.83
N_b		0.75
MC signal model		1.70
Integrated luminosity		1.00
Total		2.89

To investigate whether the $\mathcal{R}(3760)$ and $\mathcal{R}(3780)$ states decay to nOCH and if a new resonance \mathcal{R} exists in this energy region, we perform a least- χ^2 fit to the nOCH cross sections. The dressed nOCH cross section is modeled by:

$$\sigma_{\text{nOCH}}^{\text{D}}(s') = \sigma_{\text{LH}}^{\text{D}}(s') + \sigma_{J/\psi}^{\text{D}}(s') + \sigma_{\mathcal{R}_{\text{Sup}3680}}^{\text{D}}(s'), \quad (1)$$

where $s' = s(1 - x)$, x is the radiative-photon energy fraction, $\sigma_{\text{LH}}^{\text{D}}(s')$, $\sigma_{J/\psi}^{\text{D}}(s')$ and $\sigma_{\mathcal{R}_{\text{Sup}3680}}^{\text{D}}(s')$ are the cross sections of $e^+e^- \rightarrow \text{LH}$, $J/\psi \rightarrow \text{hadrons}$, and $\mathcal{R}_{\text{Sup}3680} \rightarrow \text{nOCH}$, respectively. Here $\mathcal{R}_{\text{Sup}3680}$ indicates the states with masses above 3.680 GeV/ c^2 . The cross sections for CPLMRs are taken to be $\sigma_{\text{LH}}^{\text{D}}(s') = f\sigma_{\mu^+\mu^-}^{\text{B}}(s') + \sigma_{\text{LMRs}}^{\text{D}}(s')$, where $\sigma_{\mu^+\mu^-}^{\text{B}}(s')$ is the Born cross section for continuum $e^+e^- \rightarrow \mu^+\mu^-$ production, f is a free parameter, $f\sigma_{\mu^+\mu^-}^{\text{B}}(s')$ gives the cross section for continuum $e^+e^- \rightarrow \text{hadrons}$ production [24], and $\sigma_{\text{LMRs}}^{\text{D}}(s')$ is the cross section for the production of LMRs decaying into LH, which is determined using the zeroth-order cross sections [36] multiplied by the vacuum-polarization correction factor $\frac{1}{|1 - \Pi(s)|^2}$ [37, 38] at energies below 2 GeV. The cross sections for J/ψ and $\mathcal{R}_{\text{Sup}3680}$ decaying into nOCH are, respectively, taken as $\sigma_{J/\psi}^{\text{D}}(s') = |A_{J/\psi}(s')|^2$ and $\sigma_{\mathcal{R}_{\text{Sup}3680}}^{\text{D}}(s') = |A_{\psi(3686)}(s') + \sum_1^3 A_k e^{i\phi_k}(s')|^2$, where $k = 1, 2, 3$ are for the $\mathcal{R}(3760)$ state, $\mathcal{R}(3780)$ state, and \mathcal{R} , respectively, and ϕ_k are relative phases.

In the above formulations, A_S are the generic decay amplitudes for these states, which are parameterized by $A_S(s') =$

$\sqrt{12\pi\Gamma_S^{ee}\Gamma_S^{\text{tot}}\mathcal{B}(\mathcal{S} \rightarrow \text{nOCH})}/[(s' - M_S^2) + i(\Gamma_S^{\text{tot}}M_S)]$, in which \mathcal{S} stands for the J/ψ , $\psi(3686)$, $\mathcal{R}(3760)$, and $\mathcal{R}(3780)$ states, as well as resonance \mathcal{R} ; M_S , Γ_S^{tot} and Γ_S^{ee} are, respectively, the mass, total and electronic widths of the \mathcal{S} , and $\mathcal{B}(\mathcal{S} \rightarrow \text{nOCH})$ is the decay branching fraction of the \mathcal{S} . For the $\mathcal{R}(3780)$ state, the total width is set to be energy dependent, as in Ref. [24].

In the fit, the masses, the total widths, the electronic widths, and the hadronic decay branching fractions of J/ψ and $\psi(3686)$ resonances are fixed to those given in Ref. [36]. The $\mathcal{R}(3780)$ mass and total width are fixed to $(3781.0 \pm 1.3 \pm 0.5)$ MeV/ c^2 and $(19.3 \pm 3.1 \pm 0.1)$ MeV [26], respectively. The remaining parameters are left as free parameters in the fit.

The observed cross section is described by $\sigma_{\text{nOCH}}^{\text{O}}(s) = \int_0^{1-(4m_\pi^2/s)} dx \sigma_{\text{nOCH}}^{\text{D}}[s(1-x)] \mathcal{F}(x, s)$, where $\mathcal{F}(x, s)$ is the sampling function [24, 37], and m_π is the pion mass. The cross section as a function of energy is determined by a fit to $\sigma_{\text{nOCH}}^{\text{O}}(s)$ at the 42 energy points. Using $\sigma_{\text{nOCH}}^{\text{D}}(s)$ obtained from the parameter values, we determine the ISR correction factors $[1 + \delta(s)] = \sigma_{\text{nOCH}}^{\text{O}}(s)/\sigma_{\text{nOCH}}^{\text{D}}(s)$ at the 42 energy points, which are shown in Fig. 2 (bottom). Dividing the observed nOCH cross section $\sigma_{\text{nOCH}}^{\text{O}}(s)$ by $(1 + \delta(s))$ yields the observed dressed nOCH cross section $\sigma_{\text{nOCH}}^{\text{d}}(s)$, where the lower case superscripts are used to distinguish the measured from the predicted (upper case superscripts) quantities.

The circles with error bars in Fig. 3 show the $\sigma_{\text{nOCH}}^{\text{d}}(s)$ measurements, where the uncertainties are statistical. Using the same fit procedure described above, we fit $\sigma_{\text{nOCH}}^{\text{d}}(s)$ with the function $\sigma_{\text{nOCH}}^{\text{D}}(s)$ given in Eq. (1), with x fixed to zero. The fit converges at two sets of acceptable parameter values. We denote these as Result I and Result II. Table II summarizes the parameter values, where the first uncertainties are from the fit to the $\sigma_{\text{nOCH}}^{\text{d}}(s)$, and the second are systematic. The two sets of results have the fit quality of χ^2/ndof of 22.1/31, and 22.0/31, respectively. As the mass of the fitted resonance is close to 3810 MeV, we denote it as $\mathcal{R}(3810)$. The cross section described by Result I is superimposed on the fit results in Fig. 3. Also shown are the fit results including one contribution to $\sigma_{\mathcal{R}_{\text{Sup}3680}}^{\text{D}}(s)$ included at a time. The measured mass and total width of the $\mathcal{R}(3760)$ state are consistent within 1.8σ and 0.9σ , respectively, with those measured by the BES-II experiment [26].

To estimate the systematic uncertainties of the fitted parameter in Table II, we vary the values of the $\sigma_{\text{nOCH}}^{\text{d}}(s)$, the values of the $\sigma_{D\bar{D}}^{\text{o}}(s)$, and the fixed parameters by $\pm 1\sigma$, refit the $\sigma_{\text{nOCH}}^{\text{d}}(s)$, and take the difference between the refitted parameter value and the baseline fit result as the corresponding systematic uncertainty. The estimation of the systematic uncertainty due to the uncertainty of the c.m. energies is similar to that described in Ref. [4]. Adding these uncertainties in quadrature yields the total systematic uncertainty.

Dividing the measured values for $\Gamma_{\mathcal{R}(3760)}^{ee}\mathcal{B}_{\mathcal{R}(3760)}$ and $\Gamma_{\mathcal{R}(3780)}^{ee}\mathcal{B}_{\mathcal{R}(3780)}$ in Table II by $\Gamma_{\mathcal{R}(3760)}^{ee} = (186 \pm 201 \pm$




FIG. 3. The dressed cross sections for $e^+e^- \rightarrow n\text{OCH}$, showing also the Result-I fit (line 1) and the contributions to this fit including separate contributions to $\sigma_{\mathcal{R}_{\text{Sup}3680}}^{\text{D}}(s)$ (lines 2–6).

TABLE II. Results of the fit to the cross sections for $e^+e^- \rightarrow n\text{OCH}$ showing the values of the mass M_k [MeV/c^2], total width Γ_k^{tot} [MeV], the product of electronic width (Γ_k^{ee}) and nOCH branching fraction (\mathcal{B}_k) $\Gamma_k^{ee} \mathcal{B}_k$ [eV], and relative phase ϕ_k [degree], where k represents $\mathcal{R}(3760)$, $\mathcal{R}(3780)$, and $\mathcal{R}(3810)$. \mathcal{B}_k is the branching fractions [%] for $\mathcal{R}(3760)$ and $\mathcal{R}(3780)$ decays into nOCH and OC.

Parameters	Result I	Result II
$M_{\mathcal{R}(3760)}$	$3739.9 \pm 4.2 \pm 2.6$	$3739.7 \pm 3.9 \pm 2.6$
$\Gamma_{\mathcal{R}(3760)}^{\text{tot}}$	$23.9 \pm 8.2 \pm 4.8$	$22.5 \pm 8.3 \pm 4.5$
$\Gamma_{\mathcal{R}(3760)}^{ee} \mathcal{B}_{\mathcal{R}(3760)}$	$46.8 \pm 29.9 \pm 25.1$	$11.9 \pm 9.0 \pm 6.4$
$\phi_{\mathcal{R}(3760)}$	$228 \pm 52 \pm 58$	$113 \pm 51 \pm 29$
$\Gamma_{\mathcal{R}(3780)}^{ee} \mathcal{B}_{\mathcal{R}(3780)}$	$29.9 \pm 16.1 \pm 3.7$	$25.3 \pm 11.6 \pm 3.1$
$\phi_{\mathcal{R}(3780)}$	$82 \pm 126 \pm 17$	$250 \pm 119 \pm 52$
$M_{\mathcal{R}(3810)}$	$3805.7 \pm 1.1 \pm 2.7$	$3805.7 \pm 1.1 \pm 2.7$
$\Gamma_{\mathcal{R}(3810)}^{\text{tot}}$	$11.6 \pm 2.9 \pm 1.9$	$11.5 \pm 2.8 \pm 1.9$
$\Gamma_{\mathcal{R}(3810)}^{ee} \mathcal{B}_{\mathcal{R}(3810)}$	$10.9 \pm 3.8 \pm 2.5$	$11.0 \pm 3.4 \pm 2.5$
$\phi_{\mathcal{R}(3810)}$	$52 \pm 149 \pm 25$	$215 \pm 148 \pm 103$
f	$2.28 \pm 0.01 \pm 0.07$	$2.28 \pm 0.01 \pm 0.07$
$\mathcal{B}[\mathcal{R}(3760) \rightarrow n\text{OCH}]$	$25.2 \pm 16.1 \pm 30.4$	$6.4 \pm 4.8 \pm 7.7$
$\mathcal{B}[\mathcal{R}(3780) \rightarrow n\text{OCH}]$	$12.3 \pm 6.6 \pm 8.3$	$10.4 \pm 4.8 \pm 7.0$
$\mathcal{B}[\mathcal{R}(3760) \rightarrow \text{OC}]$	$74.8 \pm 16.1 \pm 30.4$	$93.6 \pm 4.8 \pm 7.7$
$\mathcal{B}[\mathcal{R}(3780) \rightarrow \text{OC}]$	$87.7 \pm 6.6 \pm 8.3$	$89.6 \pm 4.8 \pm 7.0$

8) eV [26] and $\Gamma_{\mathcal{R}(3780)}^{ee} = (243 \pm 160 \pm 9)$ eV [26], respectively, yields the nOCH branching fractions for the decays $\mathcal{R}(3760) \rightarrow n\text{OCH}$ and $\mathcal{R}(3780) \rightarrow n\text{OCH}$, which are shown in Table II. Their corresponding OC branching fractions are also presented in Tab. II. These nOCH branching fractions of $\mathcal{R}(3780)$ decays are in good agreement with $\mathcal{B}[\psi(3770) \rightarrow \text{non-}D\bar{D}] = (15.1 \pm 5.6 \pm 1.8)\%$ [21–24] measured by the BES-II experiment.

By removing the $\mathcal{R}(3810)$ from the $\sigma_{\mathcal{R}_{\text{Sup}3680}}^{\text{D}}(s)$ as discussed above, the χ^2/ndof of the fit changes from 22.1/31 to 116.1/36, indicating that the significance of the $\mathcal{R}(3810)$

signal is 8.8σ . Similarly, the significance of the $\mathcal{R}(3760)$ signal is determined to be 8.1σ by comparing the difference of χ^2/ndof relative to the number degrees of freedom with and without including the $\mathcal{R}(3760)$ component in the fit. A similar procedure is applied to determine the $\mathcal{R}(3780)$ significance, which is 13.7σ . These significances include the systematic uncertainties.

The charmonium model [39] predicts only the 1^3D_1 state existing in the c.m. energy range from 3.733 to 3.870 GeV, which is generally assumed to be the $\psi(3770)$, and so the $\mathcal{R}(3760)$ and $\mathcal{R}(3810)$ state are presumably two non- $c\bar{c}$ states. The $\mathcal{R}(3760)$ state can be explained as a p -wave resonance of a four-quark ($c\bar{c}q\bar{q}$) state [40, 41]. It can be thought of either as an OC molecular state, or as a four-quark bound state [41]. Reference [42] interprets $\mathcal{R}(3760)$ as a possible molecular OC threshold resonance. The most salient features of the $\mathcal{R}(3760)$ state is that its mass, which is $(3739.9 \pm 4.2 \pm 2.6)$ MeV/c^2 for Result I [$3739.7 \pm 3.9 \pm 2.6$ MeV/c^2 for Result II], is just at the D^+D^- threshold (3739.3 ± 0.1 MeV/c^2 , and its branching fraction is $\mathcal{B}[\mathcal{R}(3760) \rightarrow \text{OC}] = (74.8 \pm 16.1 \pm 30.4)\%$ [$\mathcal{B}[\mathcal{R}(3760) \rightarrow \text{OC}] = (93.6 \pm 4.8 \pm 7.7)\%$]. These experimental facts can lead one naturally to interpret the $\mathcal{R}(3760)$ state as an OC pair molecular state, but containing a simple four-quark state component. As no signal for the decay $\mathcal{R}(3810) \rightarrow D\bar{D}$ is observed in the cross sections for $e^+e^- \rightarrow D\bar{D}$ at c.m. energies around 3.810 GeV [43, 44], and the $\mathcal{R}(3810)$ mass ($3805.7 \pm 1.1 \pm 2.7$ MeV/c^2 for Result I [$3805.7 \pm 1.1 \pm 2.7$ MeV/c^2 for Result II]) is exactly at the $h_c\pi^+\pi^-$ threshold (3804.5 ± 0.1 MeV/c^2 , the $\mathcal{R}(3810)$ state can be interpreted as a hadrocharmonium resonance [8].

In summary, we have measured the inclusive nOCH cross sections of $e^+e^- \rightarrow n\text{OCH}$ with improved precision at c.m. energies from 3.645 to 3.871 GeV. We observe three resonances: $\mathcal{R}(3760)$, $\mathcal{R}(3780)$, and $\mathcal{R}(3810)$ in the energy-dependent line shape of the nOCH cross sections with significances of 8.1σ , 13.7σ , and 8.8σ , respectively. The $\mathcal{R}(3760)$ and $\mathcal{R}(3780)$ states are observed for the first time in the nOCH cross sections, while the $\mathcal{R}(3810)$ state is observed for the first time with mass ($3805.7 \pm 1.1 \pm 2.7$) [$3805.7 \pm 1.1 \pm 2.7$] MeV/c^2 , total width ($11.6 \pm 2.9 \pm 1.9$) [($11.5 \pm 2.8 \pm 1.9$)] MeV , and the product of the electronic width and the nOCH decay branching fraction ($10.9 \pm 3.8 \pm 2.5$) [($11.0 \pm 3.4 \pm 2.5$)] eV for Result I [Result II]. In addition, for the first time, we measure $\mathcal{B}[\mathcal{R}(3760) \rightarrow n\text{OCH}] = (25.2 \pm 16.1 \pm 30.4)\%$ [($6.4 \pm 4.8 \pm 7.7$)%] and $\mathcal{B}[\mathcal{R}(3780) \rightarrow n\text{OCH}] = (12.3 \pm 6.6 \pm 8.3)\%$ [($10.4 \pm 4.8 \pm 7.0$)%]. The $\mathcal{R}(3760)$ state can be interpreted as an OC pair molecular state, but containing a simple four-quark state component. The $\mathcal{R}(3810)$ state can be interpreted as a hadrocharmonium state [8].

The BESIII Collaboration thanks the staff of BEPCII and the IHEP computing center for their strong support. This work is supported in part by National Key R&D Program of China under Contracts No. 2009CB825204, No. 2020YFA0406300, No. 2020YFA0406400; National Natural Science Foundation of China (NSFC) under Contracts No. 10935007, No.

11635010, No. 11735014, No. 11835012, No. 11935015, No. 11935016, No. 11935018, No. 11961141012, No. 12022510, No. 12025502, No. 12035009, No. 12035013, No. 12061131003, No. 12192260, No. 12192261, No. 12192262, No. 12192263, No. 12192264, No. 12192265, No. 12221005, No. 12225509, No. 12235017; the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; the CAS Center for Excellence in Particle Physics (CCEPP); CAS Key Research Program of Frontier Sciences under Contracts No. QYZDJ-SSW-SLH003, No. QYZDJ-SSW-SLH040; 100 Talents Program of CAS; The CAS Research Program under Code No. Y41G1010Y1; The CAS Other Research Program under Code No. Y129360; The Institute of Nuclear and Particle Physics (INPAC) and Shanghai Key Laboratory for Particle Physics and Cosmology; ERC under Contract No. 758462; European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement under Contract No. 894790; German Research Foundation DFG under Contracts No. 443159800, No. 455635585, Collaborative Research Center CRC 1044, FOR5327, GRK 2149; Istituto Nazionale di Fisica Nucleare, Italy; Ministry of Development of Turkey under Contract No. DPT2006K-120470; National Research Foundation of Korea under Contract No. NRF-2022R1A2C1092335; National Science and Technology fund of Mongolia; National Science Research and Innovation Fund (NSRF) via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation of Thailand under Contract No. B16F640076; Polish National Science Centre under Contract No. 2019/35/O/ST2/02907; The Swedish Research Council; U. S. Department of Energy under Contract No. DE-FG02-05ER41374.

-
- [1] J. Z. Bai *et al.* (BES Collaboration), arXiv:hep-ex/0307028v1; W. G. Li, G. Rong, D. G. Cassel, in Proceedings of Tenth International Conference on Hadron Spectroscopy, Aschaffenburg, Germany, 31 August – 6 September 2003, p. 495, p. 592, p. 937; J. Z. Bai *et al.* (BES Collaboration), High Energy Physics and Nuclear Physics **28**, 325 (2004).
- [2] G. Rong, in Proceedings of the Symposium of 30 Years of BES Physics, Beijing, China, 2019 (World Scientific, Singapore, 2019), p. 48.
- [3] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. D **102**, 112009 (2020).
- [4] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. Lett. **127**, 082002 (2021).
- [5] F. E. Close and P. R. Page, Phys. Lett. B **578**, 119 (2004).
- [6] M. B. Voloshin and L. B. Okun, JETP Lett. **23**, 369 (1976).
- [7] N. Brambilla *et al.*, Eur. Phys. J C **71**, 1534 (2011).
- [8] S. Dubynskiy and M. B. Voloshin, Phys. Letts. B **666**, 344 (2008).
- [9] J. Z. Bai *et al.* (BES Collaboration), Phys. Lett. B **605**, 63 (2005).
- [10] S. K. Choi *et al.* (Belle Collaboration), Phys. Rev. Lett. **91**, 262001 (2003).
- [11] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. Lett. **95**, 142001 (2005).
- [12] Q. He *et al.* (CELO Collaboration), Phys. Rev. D **74**, 091104 (2006).
- [13] T. E. Coan *et al.* (CLEO Collaboration), Phys. Rev. Lett. **96**, 162003 (2006).
- [14] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. Lett. **110**, 252001 (2013).
- [15] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. Lett. **112**, 092001 (2014).
- [16] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. Lett. **111**, 242001 (2013).
- [17] X.L. Wang *et al.* (Belle Collaboration), Phys. Rev. Lett. **99**, 142002 (2007).
- [18] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. Lett. **98**, 212001 (2007).
- [19] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. Lett. **118**, 092001 (2017).
- [20] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. Lett. **118**, 092002 (2017).
- [21] M. Ablikim *et al.* (BES Collaboration), Phys. Letts. B **641**, 145 (2006).
- [22] M. Ablikim *et al.* (BES Collaboration), Phys. Rev. D **76**, 122002 (2007).
- [23] M. Ablikim *et al.* (BES Collaboration), Phys. Letts. B **659**, 74 (2008).
- [24] M. Ablikim *et al.* (BES Collaboration), Phys. Rev. Lett. **97**, 121801 (2006).
- [25] G. Rong, Chin. Phys. C **34**, 788 (2010).
- [26] M. Ablikim *et al.* (BES Collaboration), Phys. Rev. Lett. **101**, 102004 (2008).
- [27] M. Ablikim *et al.* (BESIII Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **614**, 345 (2010).
- [28] S. Agostinelli *et al.* (GEANT4 Collaboration), Nucl. Instrum. Methods Phys. Res. Sect. A **506**, 250 (2003).
- [29] S. Jadach, B. F. L. Ward, and Z. Was, Comput. Phys. Commun. **130**, 260 (2000).
- [30] D. J. Lange, Nucl. Instrum. Methods Phys. Res., Sect. A **462**, 152 (2001); R. G. Ping, Chin. Phys. C **32**, 599 (2008).
- [31] J. C. Chen, G. S. Huang, X. R. Qi, D. H. Zhang, and Y. S. Zhu, Phys. Rev. D **62**, 034003 (2000).
- [32] G. Balossini, C. M. Carloni Calame, G. Montagna, O. Nicrosini and F. Piccinini, Nucl. Phys. **B758**, 227 (2006); G. Balossini, C. Bignamini, C. M. C. Calame, G. Montagna, O. Nicrosini and F. Piccinini, Phys. Lett. B **663**, 209 (2008).
- [33] S. Nova, A. Olchevski and T. Todorov, TWOGAM, a Monte Carlo event generator for two photon physics, DELPHI Note 90-35 PROG 152.
- [34] J. Z. Bai, *et al.* (BES Collaboration), Phys. Rev. D **53**, 20 (1996).
- [35] H. Burkhardt and B. Pietrzyk, Phys. Lett. **B513**, 46 (2001).
- [36] R. L. Workman *et al.* (Particle Data Group), Prog. Theor. Exp. Phys. **2022**, 083C01 (2022).
- [37] E. A. Kuraev and V. S. Fadin, Yad. Fiz. **41**, 733 (1985); Sov. J. Nucl. Phys. **41**, 466 (1985).
- [38] D. Zhang, G. Rong, and J. C. Chen, Phys. Rev. D **74**, 054012 (2006).
- [39] E. Eichten, K. Gottfried, T. Kinoshita, K. D. Lane, and T.-M. Yan, Phys. Rev. D **17**, 3090 (1978); E. Eichten, K. Gottfried, T. Kinoshita, K. D. Lane, and T. M. Yan, Phys. Rev. D **21**, 203 (1980);
- [40] A. De Rujula, Howard Georgi, and S. L. Glashow, Phys. Rev. Lett. **38**, 317 (1977).
- [41] Rafe Hyam Schindler, Stanford Linear Accelerator Center,

- Stanford University, Stanford, California, Report No. 219, p. 98, May 1979.
- [42] S. Dubynskiy and M. B. Voloshin, Phys. Rev. D **78**, 116014 (2008).
- [43] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **76**, 111105(R) (2007).
- [44] G. Pakhlova *et al.* (Belle Collaboration), Phys. Rev. D **77**, 011103(R) (2008).