# NON-ISOTOPIC SPLITTING SPHERES FOR A SPLIT LINK IN $S^4$

MARK HUGHES, SEUNGWON KIM, AND MAGGIE MILLER

ABSTRACT. We show that there exist split, orientable, 2-component surface-links in  $S^4$  with non-isotopic splitting spheres in their complements. In particular, for non-negative integers m, n with  $m \geq 4$ , the unlink  $L_{m,n}$  consisting of one component of genus m and one component of genus n contains in its complement two smooth splitting spheres that are not topologically isotopic in  $S^4 \setminus L_{m,n}$ . This contrasts with link theory in the classical dimension, as any two splitting spheres in the complement of a 2-component split link  $L \subset S^3$  are isotopic in  $S^3 \setminus L$ .

### 1. INTRODUCTION

In this paper, we study *splitting spheres* in surface-link complements in  $S^4$ . Unless otherwise specified, statements are in the topological locally-flat category.

**Definition 1.1.** Let  $L = L_1 \sqcup L_2$  be a compact, codimension-2 submanifold of  $S^n$  with two connected components  $L_1$  and  $L_2$ . A splitting sphere S for L is an embedded  $S^{n-1}$  in  $S^n \setminus L$  such that  $L_1$  and  $L_2$  lie in distinct components of  $S^n \setminus S$ .

It is a consequence of the Schoenflies theorem in dimension three that any two splitting spheres  $S_1, S_2$  for a 2-component link L in  $S^3$  are isotopic in the complement of L. Laudenbach showed that more generally, any two homotopic 2-spheres in a 3-manifold are isotopic [Lau73]. However, the situation is quite different in dimension four. For example, Iida– Konno–Mukherjee-Taniguchi [IKMT] (following work of Donaldson and Wall) gave many examples of pairs of separating 3-spheres in 4-manifolds that are not smoothly isotopic. In this setting, the 3-spheres are topologically isotopic and the smooth obstruction necessitates  $b_2^+$  of the ambient 4-manifold to be positive. However, avoiding any  $b_2$  requirements, Budney–Gabai [BG] recently produced many homotopic yet smoothly non-isotopic essential 3-spheres in  $S^1 \times S^3$ .

As a consequence, Budney–Gabai showed that the unknotted 2-sphere in  $S^4$  bounds 3balls in  $S^4$  that are not smoothly isotopic rel. boundary. Inspired by their work, in [HKM23] we proved (using very different methods) that for  $g \ge 2$ , the unknotted genus-g surface in  $S^4$  bounds smooth handlebodies that are homeomorphic rel. boundary but not topologically isotopic rel. boundary. In this paper, we adapt our previous techniques to prove the following theorem.

**Theorem 1.2.** Let m, n be non-negative integers with  $m \ge 4$ . Let  $L = L_{m,n}$  be the 2component unlink in  $S^4$  with one component of genus m and one component of genus n. There exist two splitting spheres  $S_1, S_2$  for L such that  $S_1$  and  $S_2$  are not isotopic in  $S^4 \setminus L$ .

Here, an *unlink* in  $S^4$  is a split union of *unknots*, while an *unknot* in  $S^4$  is a surface that bounds a 3-dimensional 1-handlebody in  $S^4$ . For each g, the genus-g unknot is unique up

MH was supported by a grant from the NSF (LEAPS-MPS-2213295). MM is supported by a Clay Research Fellowship and a Stanford Science Fellowship.

to isotopy. Note that the conclusion of Theorem 1.2 greatly differs from those of [BG] or [IKMT]: the 3-spheres we obtain are separating, not topologically isotopic, and are contained in a 4-manifold  $(S^4 \setminus \nu(L_{m,n}))$  with  $b_2^+ = 0$ .

**Remark 1.3.** In the proof of Theorem 1.2, the splitting 3-spheres  $S_1, S_2$  can be taken to be smoothly embedded in the complement of the smooth unlink L, and each divide  $S^4$  into two smooth 4-balls. Yet, the splitting spheres  $S_1, S_2$  are not even topologically isotopic.

**Remark 1.4.** Theorem 1.2 can also be extended to the case when the pair (m, n) is in the set  $\{(2, 2), (2, 3), (3, 3)\}$  (see Remark 4.2).

Theorem 1.2 may seem counterintuitive, since each of  $S_1, S_2$  bound a 4-ball in  $S^4$  and hence are isotopic in  $S^4$  [Qui82]. However, to construct an isotopy from  $S_1$  to  $S_2$ , we would generally first shrink the ball bounded by  $S_1$  until  $S_1$  and  $S_2$  were disjoint, then isotope  $S_1$ along the  $S^3 \times I$  cobounded by  $S_1$  and  $S_2$  (which is where we make use of [Qui82]). In the setting of a link complement, if  $S_1, S_2 \subset S^4 \setminus L$  were disjoint then they would still cobound an  $S^3 \times I$  and would hence be isotopic. However, shrinking a 4-ball B bounded by  $S_1$  in order to separate  $S_1$  and  $S_2$  is now problematic, since one component of L lies in B.

Moreover, we point out that while uniqueness of splitting spheres for 2-component links in  $S^3$  follows from the 3-dimensional Schoenflies theorem, this argument does not extend to higher dimensions. In dimension three, we remove intersections between a pair of splitting spheres using a standard innermost circle argument: we surger one of the splitting spheres  $S_1$  along a disk contained in the other sphere  $S_2$  to split  $S_1$  into two 2-spheres, one of which is a splitting sphere isotopic to  $S_1$  that intersects  $S_2$  in fewer circles. In contrast, two splitting 3-spheres in  $S^4$  intersect in a surface, which we expect to include positive-genus components. Surgering one splitting sphere using such an intersection will split it into two 3-manifolds, but in general neither needs be a 3-sphere.

Using the proof of Theorem 1.2, we also prove the following link version of the main theorem of [HKM23].

**Theorem 1.5.** Let m, n be non-negative integers with  $m \ge 4$ . There exist two smooth, properly embedded links of handlebodies  $H_1 \sqcup H_2$  and  $\tilde{H}_1 \sqcup \tilde{H}_2$  in  $B^5$ , with  $g(H_1) = g(\tilde{H}_1) = m$ and  $g(H_2) = g(\tilde{H}_2) = n$ , such that the following are true:

- The 2-component links H<sub>1</sub> ⊔ H<sub>2</sub> and H̃<sub>1</sub> ⊔ H̃<sub>2</sub> are each smoothly boundary-parallel in B<sup>5</sup>.
- The boundaries agree componentwise, i.e.,  $\partial H_1 = \partial \tilde{H}_1, \partial H_2 = \partial \tilde{H}_2$ .
- For each i = 1, 2, the handlebodies  $H_i, \tilde{H}_i$  are smoothly isotopic rel. boundary.
- The 2-component links H<sub>1</sub> ⊔ H<sub>2</sub> and H̃<sub>1</sub> ⊔ H̃<sub>2</sub> are not topologically isotopic rel. boundary.

Given that the methods of this paper were established in our previous paper [HKM23] and were inspired by the results of Budney–Gabai [BG], it would be reasonable to hope that the techniques of [BG] could be adapted to prove the existence of non-isotopic splitting spheres in the complement of a link of 2-spheres in  $S^4$ .

**Question 1.6.** Can one adapt the methods of Budney–Gabai to prove that the unlink of two spheres in  $S^4$  has multiple splitting spheres?

One approach to Question 1.6 would be to take one 3-sphere  $S_1$  to be standard, and obtain other 3-spheres from  $S_1$  by ambient 2-handle surgery on either side of  $S_1$ , analogous to the barbells of [BG]. However, it seems to the present authors technically difficult to extend the isotopy obstruction used by Budney–Gabai from the original setting of the homotopy class of pt  $\times B^3$  in  $S^1 \times B^3$  to a homotopy class of separating 3-spheres in  $S^4 \setminus \nu(L_{0,0}) \cong$  $S^1 \times B^3 \# S^1 \times B^3.$ 

**Organization.** In Section 2 we discuss Bing doubling of codimension-2 submanifolds, particularly in ambient dimensions four and five. In Section 3, we discuss simple 3-knots, which are 3-spheres embedded in  $S^5$  whose complements have cyclic fundamental group. Finally in Section 4, we make use of the constructions in the previous two sections to produce two distinct splitting spheres in the complement of a trivial surface link, proving Theorems 1.2 and 1.5.

Acknowledgement. The work in this paper took place while the first and third authors were visiting the second author at Sungkyunkwan University (Suwon campus), Republic of Korea during February 2023.

#### 2. Bing doubles

In order to produce interesting codimension-2 links, we make use of a well-known satellite operation, Bing doubling. Since this well-known construction in classical knot theory is less well-known in higher dimensions, we review its definition here.

**Definition 2.1.** Let  $K^{n-2}$  be a closed, orientable, codimension-2 submanifold of  $S^n$  for some  $n \geq 3$ . Let  $\Sigma \subset K$  be a separating (n-3)-sphere in K. The Bing double of K along  $\Sigma$ , which we refer to as  $B_{\Sigma}(K)$ , is a 2-component codimension-2 link in  $S^n$  obtained as follows.

- (1) Start with two parallel disjoint copies K', K'' of K. In order to obtain these parallel copies, we choose a framing of K. Since the normal bundle of K in  $S^n$  is trivial, framings are in 1-to-1 correspondence with elements of  $H^1(K;\mathbb{Z})$ . We specifically choose the zero framing, which has the property that the parallel copies K', K'' are nullhomologous in  $S^n \setminus K$ . (In other words, we first find an (n+1)-manifold M in  $S^n$  bounded by K, and we take K', K'' to be parallel copies of K that bound disjoint, parallel copies of M.)
- (2) Let V be a copy of  $B^3 \times S^{n-3}$  in  $S^n$  intersecting  $K' \sqcup K''$  in tubular neighborhoods P' and P'' of pushed-off copies of  $\Sigma$  in each of K' and K'' respectively.
- (3) Parameterize  $V = B^3 \times S^{n-3}$  so that

$$(V, V \cap (P' \sqcup P'')) = (B^3 \times S^{n-3}, T_\infty \times S^{n-3}),$$

where  $(B^3, T_{\infty})$  is the  $\infty$ -tangle (see Figure 1). (4) Delete  $V = B^3 \times S^{n-3}$  and replace it with  $(B^3, T_1) \times S^{n-3}$ , where  $T_1$  is the +1-tangle (again, see Figure 1).

Let A, B denote the closures of the two components of  $K \setminus \Sigma$ . By construction, each component of  $B_{\Sigma}(K)$  bounds a copy of  $A \times I$  or  $B \times I$ . In particular, if K is a sphere then each component of  $B_{\Sigma}(K)$  bounds a ball (and hence is unknotted), though these balls are not disjoint and in general we do not expect the components of  $B_{\Sigma}(K)$  to form an unlink (see Lemma 3.2).

**Remark 2.2.** Let P be a 3-sphere in  $S^5$  intersecting an equatorial  $S^4$  in a connected surface S, and let  $\Sigma$  be a 2-sphere in P intersecting S in a circle  $\gamma$ . Then  $B_{\Sigma}(P)$  intersects  $S^4$  in a Bing double  $B_{\gamma}(S)$ .




FIGURE 1. Top left: a schematic of a codimension-2 manifold  $K^{n-2}$  of  $S^n$ and a separating (n-3)-sphere  $\Sigma$  in K. Top right: two parallel copies K', K'' of K and a tubular neighborhood  $V = B^3 \times S^{n-3}$  of  $\Sigma$  intersecting  $K' \cup K''$  in  $T_{\infty} \times S^{n-3}$  where  $T_{\infty}$  is the  $\infty$ -tangle. Bottom left: we replace the intersection of  $(K' \cup K'') \cap V$  with  $T_1 \times S^{n-3}$  where  $T_1$  is the +1-tangle. Bottom right: the Bing double  $B_{\Sigma}(K)$ .

**Proposition 2.3.** Let S be a smoothly unknotted genus-g surface in  $S^4$ . Then for any separating curve  $\gamma$  in S,  $B_{\gamma}(S)$  is a smooth 2-component unlink.

Note that it is obvious from Definition 2.1 that each component of  $B_{\gamma}(S)$  is unknotted for any surface S in  $S^4$ , as  $\gamma$  cuts S into two 2-dimensional 1-handlebodies and hence each component of  $B_{\gamma}(S)$  bounds a 3-dimensional 1-handlebody.

Proof of Proposition 2.3. Let  $F_1, F_2$  be the closures of the two components of  $S \setminus \gamma$ . Let  $\gamma'$  be any other separating curve on S so that the closures  $F'_1, F'_2$  of the components of  $S \setminus \gamma'$  satisfy  $g(F'_i) = g(F_i)$  for each i.

**Claim 2.4.** There is a self-diffeomorphism  $\phi$  of  $S^4$  fixing S setwise but with  $\phi(\gamma) = \gamma'$ .

Proof of Claim. Let  $(\alpha_1, \beta_1)$  be a symplectic basis of  $H_1(F_1; \mathbb{Z})$  and  $(\alpha_2, \beta_2)$  be a symplectic bases of  $H_1(F_2; \mathbb{Z})$ . Take all  $\alpha_i, \beta_i$  curves to have trivial Rokhlin invariant (as in [Hir02]). Similarly, choose vanishing-Rokhlin symplectic bases  $(\alpha'_1, \alpha'_2), (\beta'_1, \beta'_2)$  for  $H_1(F'_1; \mathbb{Z})$  and  $H_1(F'_2; \mathbb{Z})$  respectively. Let  $\psi : S \to S$  be a surface automorphism with  $\psi(\alpha_i) = \alpha'_i$  and  $\psi(\beta_i) = \beta'_i$  for each *i*. Then up to isotopy,  $\psi(\gamma) = \gamma'$ . By Hirose [Hir02], the map  $\psi$  extends as a diffeomorphism over  $S^4$ .

In particular, there is a diffeomorphism  $\phi : S^4 \to S^4$  taking  $(S, \gamma)$  to the pair  $(S, \gamma')$  illustrated in Figure 2. In Figure 2 we construct  $B_{\gamma'}(S)$  and explicitly observe that this 2-component link is an unlink. Since  $(S^4, B_{\gamma'}(S)) \cong (S^4, B_{\gamma}(S))$ , we conclude that  $B_{\gamma}(S)$  is also an unlink.

When  $K^{n-2}$  is a codimension-2 sphere, then we can omit the separating sphere  $\Sigma$  from the notation  $B_{\Sigma}(K)$ , since any two codimension-1 subspheres of K are isotopic in K (remember that we are working in the topological locally-flat category). In this setting, it is also a simple exercise to check that B(K) is well-defined up to isotopy in a neighborhood of K.




FIGURE 2. Top row: For S an unknotted surface in  $S^4$ , there is a selfdiffeomorphism  $\phi$  of  $S^4$  fixing S setwise and taking a separating curve  $\gamma$  to a standard separating curve  $\gamma'$ . Taking S to be contained in an equatorial 3-sphere,  $\gamma'$  is the intersection of S with an equatorial  $S^2$  of the 3-sphere. We illustrate here the case that  $\gamma$  cuts S into a genus-1 and a genus-2 piece. Bottom row: we draw a banded unlink diagram of the Bing double  $B_{\gamma'}(S)$ and observe that  $B_{\gamma'}(S)$  is an unlink.




FIGURE 3. First: A schematic of a codimension-2 sphere K in  $S^n$ . Second: Four parallel copies of K with indicated orientations, and two framed arcs connecting different pairs of copies of K. Third: We surger the four parallel copies of K along the two arcs to obtain a link of two spheres, each of which is unknotted. Fourth: The link is isotopic to the Bing double B(K).

**Remark 2.5.** Let  $K^{n-2}$  be a codimension-2 sphere in  $S^n$  for  $n \ge 3$ . Then the Bing double of K can also be obtained by the following procedure.

Let  $K_1, K_2, K_3, K_4$  be disjoint parallel copies of K. (If n = 3 then ensure these copies are parallel with the 0-framing; if n > 3 then  $H^1(K; \mathbb{Z}) = 0$  so there is no ambiguity.) Give  $K_1, K_2$  parallel orientations and  $K_3, K_4$  the opposite orientation. Surger  $K_1 \sqcup K_2 \sqcup K_3 \sqcup K_4$ along two arcs as specified in Figure 3, taking framing on the arcs that are compatible with the relative orientations of  $K_1, K_3$  and  $K_2, K_4$ .




FIGURE 4. When  $K^{n-2}$  is a codimension-2 sphere in  $S^n$ , the Bing double of K is obtained by the illustrated satellite operation. Left:  $U_1, U_2$  form a codimension-2 unlink in  $S^n$  and C is a curve representing  $\mu_1 \mu_2 \mu_1^{-1} \mu_2^{-1}$ in the complement of  $U_1 \sqcup U_2$ , where  $\mu_i$  is a meridian of  $U_i$ . Right: the pattern  $(X, B) = (S^n \setminus \nu(C), U_1 \sqcup U_2)$ .

Framings of an arc in dimension  $n \ge 4$  are in bijection with  $\pi_1(SO(n-1)) = \mathbb{Z}/2\mathbb{Z}$ ; requiring that the surgeries respect the orientations on  $K_1, K_2, K_3, K_4$  determines the framings on the two arcs, and thus the Bing double B(K) is well-defined up to isotopy when  $K^{n-2}$  is a codimension-2 sphere with  $n \ge 4$ .

Additionally, it is useful to note that B(K) can be obtained by a certain satellite operation.

**Remark 2.6.** Let  $K^{n-2}$  be a codimension-2 sphere in  $S^n$  for  $n \ge 3$ . Then the Bing double of K can be obtained by the following satellite operation (see Figure 4).

Let  $U_1 \sqcup U_2$  be a 2-component unlink of (n-2)-spheres in  $S^n$ . Then  $\pi_1(S^n \setminus (U_1 \sqcup U_2))$ is a free group generated by meridians  $\mu_1$ , and  $\mu_2$  for  $U_1, U_2$  respectively. Let C be a curve in the complement of  $U_1 \sqcup U_2$  freely representing  $\mu_1 \mu_2 \mu_1^{-1} \mu_2^{-1}$ .

Now set  $(X, B) = (S^n \setminus \nu(C), U_1 \sqcup U_2)$ . The Bing double B(K) of K is then obtained by

$$(S^n, B(K)) = ((S^n \setminus \nu(K)) \cup X, B).$$

If n = 3, then the union above must be chosen so that a meridian of C is glued to a 0-framed longitude of K. If  $n \ge 4$  the resulting link does not depend on the gluing map ([Sta63, Corollary 9.3] for  $n \ge 5$ , [LP72] for n = 4).

Both Remarks 2.5 and 2.6 contain standard phrasings of the definition of the Bing double of a classical knot in  $S^3$ , extended to hold in general dimension.

## 3. SIMPLE 3-KNOTS

In this section, we study (smooth/topological) 3-knots, which are (smooth/locally-flat) submanifolds of  $S^5$  that are homeomorphic to  $S^3$ . We say that a 3-knot P is simple if  $\pi_1(S^5 \setminus \nu(P)) \cong \mathbb{Z}$ . A 3-knot is unknotted if it bounds a 4-ball in  $S^5$  (with no mention of category necessary, as any smooth 3-knot that bounds a locally flat 4-ball also bounds a smooth 4-ball [Sha68, Theorem 1.1]).

While the unknotted 3-knot is simple, there are also many nontrivial simple 3-knots. Consider the following lemma, which appears in [HKM23] and is mostly a consequence of Hirose's work on extending surface automorphisms from an unknotted surface to all of  $S^4$  [Hir02].

**Lemma 3.1.** Every smooth 2-knot K arises as the cross-section of a smooth simple 3-knot. If K can be turned into a smoothly unknotted genus-g surface via surgery along g arcs, then the height function h on  $S^5$  can be taken to restrict to this simple 3-knot as a Morse function with one index-0 point, 2g index-1 points, 2g index-2 points, and one index-3 point.

Lemma 3.1 is particularly interesting in the case that K is smooth and not doubly slice (e.g., when K is the 2-twist spun trefoil [Sto79]), as then we automatically know that K is the cross-section of some nontrivial simple 3-knot.

The existence of nontrivial simple 3-knots makes the following lemma surprisingly nontrivial to prove.

**Lemma 3.2.** Let P be a nontrivial 3-knot. Then the Bing double B(P) of P is not split.

*Proof.* The components of B(P) are each unknotted, so if B(P) is split then B(P) is an unlink. Thus if B(P) is split,  $S^5 \setminus \nu(B(P)) \simeq S^1 \vee S^1 \vee S^4$ . In particular, if B(P) is split then  $\pi_1(S^5 \setminus \nu(B(P)))$  is free and  $\pi_2(S^5 \setminus \nu(B(P)))$  is trivial.

As observed in Remark 2.6, the complement  $S^5 \setminus \nu(B(P))$  decomposes as a union of a manifold  $X_1 \cong S^5 \setminus \nu(P)$  and a manifold  $X_2$  homeomorphic to the complement in  $S^5$ of a generalized Borromean rings of two 3-spheres and a circle. The manifolds  $X_1$  and  $X_2$  are glued along an  $S^1 \times S^3$  boundary component (which in  $X_2$  is the boundary of the deleted tubular neighborhood of a circle). Let  $U_1, U_2$  be small tubular open neighborhoods of  $X_1, X_2$  respectively. The Seifert-van Kampen theorem applied to  $\{U_1, U_2\}$  yields the following pushout.

$$\mathbb{Z} = \pi_1(S^1 \times S^3, *) \xrightarrow{1 \mapsto \mu} \pi_1(S^5 \setminus \nu(P)) = \langle \langle \mu \rangle \rangle$$

$$\begin{array}{c}1\\\downarrow\\aba^{-1}b^{-1}\\\langle a, b \rangle = \pi_1(X_2) \longrightarrow \pi_1(S^5 \setminus \nu(B(P)))\end{array}$$

If  $\pi_1(S^5 \setminus \nu(P)) \cong \mathbb{Z}$ , then  $\pi_1(S^5 \setminus \nu(B(P))$  is not free and we conclude that B(P) is not split.

We now restrict to the case that  $\pi_1(S^5 \setminus \nu(P)) \cong \mathbb{Z}$ , i.e., P is simple. We will show that  $\pi_2(S^5 \setminus B(P)) \cong 0$ .

By Kreck–Su [KS17], if a 3-knot  $Q \subset S^5$  is such that  $\pi_1(S^5 \setminus Q) \cong \mathbb{Z}$  and  $\pi_2(S^5 \setminus Q) = 0$ , then  $S^5 \setminus \nu(Q)$  is homeomorphic to  $S^1 \times B^4$ , so by e.g. [Sta63, Corollary 9.3] the 3-knot Qbounds a ball in  $S^5$ . Since  $\pi_1(S^5 \setminus \nu(P)) \cong \mathbb{Z}$  and P does not bound a ball, we conclude that  $\pi_2(S^5 \setminus \nu(P)) \neq 0$ .

We will make use of Althoen's analogue of the Seifert van Kampen theorem for  $\pi_2$  [Alt78]. This theorem applies to an open cover  $\{V_1, V_2\}$  of a path-connected topological space when  $V_1, V_2$ , and  $V_1 \cap V_2$  are all path-connected and the maps induced by inclusion from  $\pi_1(V_1 \cap V_2) \rightarrow \pi_1(V_i)$  are injective for both i = 1, 2. We have already seen that this is the case for the open cover  $\{U_1, U_2\}$  of  $S^5 \setminus \nu(B(P))$ .

We thus obtain the following pushout diagram, with \* a basepoint in  $U_3 := U_1 \cap U_2 \simeq S^1 \times S^3$  and indexing sets  $I_j = \pi_1(S^5 \setminus \nu(B(P)), *)/\pi_1(U_j, *)$ .

Because  $q \equiv 0$ , the map f is an injection. Since  $\pi_2(S^5 \setminus \nu(P)) \not\cong 0$  we conclude that  $\pi_2(S^5 \setminus \nu(B(P))) \not\cong 0$  and hence B(P) is not split. 

**Proposition 3.3.** Let K be a smooth 2-knot with the property that K can be turned into a smoothly unknotted torus via surgery along some arc. There is a smooth link  $P_1 \cup P_2$  of unknotted 3-spheres in  $S^5$  so that  $(P_1 \cup P_2) \cap S^4 = B(K) = B_1 \sqcup B_2$  (with  $P_i \cap S^4 = B_i$ ), and the height function h on  $S^5$  restricts to each of  $P_1, P_2$  as follows.

- The function  $h|_{P_1}$  is Morse with one local minimum, four index-1 critical points, four index-2 critical points, and one local maximum.
- The function  $h|_{P_2}$  is Morse with one local minimum, one local maximum, and no other critical points.

Moreover,  $P_1 \cup P_2$  is isotopic to the Bing double B(P) of P for P some simple 3-knot with cross-section K.

*Proof.* Choose a curve  $\gamma$  on K and isotope B(K) to agree with  $B_{\gamma}(K)$ . The curve  $\gamma$  separates K into disks  $D_1$  and  $D_2$  so that  $B_1, B_2$  are respectively doubles of  $D_1, D_2$ . Let H be a 3dimensional 1-handle with ends on K so that surgering K along H yields an unknotted torus T. Isotope H so that the ends of H lie in  $D_1$ . By Hirose [Hir02], T bounds a solid torus V so that there is a curve on T bounding a disk in V that is geometrically dual to the core of H. As in Lemma 3.1, we produce a simple 3-knot P with cross-section K by specifying the intersections of P with various subsets of  $S^4 \times [-1,1]$  as follows (see left of Figure 5).

- $(S^4 \times [-1,0], S^4 \times [-1,0] \cap P)$  is the mirror of  $(S^4 \times [0,1], S^4 \times [0,1] \cap P)$ .
- $P \cap (S^4 \times [0, 1/3)) = K \times [0, 1/3),$
- $P \cap (S^4 \times \{1/3\}) = (K \cup H) \times \{1/3\},$   $P \cap (S^4 \times (1/3, 1)) = T \times (1/3, 1),$
- $P \cap (S^4 \times \{1\}) = V \times \{1\}.$

Note that we could smooth P so that  $h|_P$  is Morse with one local minimum, two index-1 critical points, two index-2 critical points, and one local maximum.<sup>1</sup>

Now let  $\Sigma$  be a separating 2-sphere in P as in Figure 5. That is,  $\Sigma \cap h^{-1}(0) = C$ , both index-1 points of  $h|_P$  lie in the same component of  $P \setminus \Sigma$  as does  $D_1$ , and  $h|_{\Sigma}$  has one local minimum and one local maximum, which coincide with the local minimum and maximum of  $h|_P$ . We can then take  $\Sigma \cap (V \times \{1\})$  to be a disk. Let  $V_1, V_2$  be the genus-2 handlebody and 3-ball (respectively) each obtained by taking one component of  $(V \times \{1\}) \setminus \Sigma$  and doubling along the intersection with  $\Sigma$ . Let  $H_1, H_2$  denote parallel copies of the 3-dimensional 1handle H, with ends on distinct sheets of  $B_1$ . Let  $B'_1$  denote the result of surgering  $B_1$ along  $H_1 \sqcup H_2$  (note that  $B'_1$  bounds a copy of  $V_1$  and  $B_2$  bounds a copy of  $V_2$ ).

Now  $B_{\Sigma}(P) = P_1 \sqcup P_2$  is (up to isotopy) as follows (see right of Figure 5).

<sup>&</sup>lt;sup>1</sup>The resulting 3-manifold P thus admits a handle decomposition with one 0-handle, two cancelling 1/2handle pairs, and one 3-handle, hence is a 3-sphere. Since  $h|_P$  has a unique local minimum,  $\pi_1(S^5 \setminus \nu(P))$ is cyclic and hence P is simple.




FIGURE 5. The construction of Proposition 3.3. Left: the portion of the simple 3-knot P contained in  $h^{-1}[0,1]$ . Right: the portion of the Bing double  $B(P) = P_1 \sqcup P_2$  contained in  $h^{-1}[0, 1]$ .

- $(S^4 \times [-1,0], S^4 \times [-1,0] \cap P_1 \cup P_2)$  is the mirror of  $(S^4 \times [0,1], S^4 \times [0,1] \cap P_1 \cup P_2)$ .
- $P_1 \cap (S^4 \times [0, 1/3)) = B_1 \times [0, 1/3),$
- $P_1 \cap (S^4 \times \{0, 1/3\}) = D_1 \times [0, 1/3),$   $P_1 \cap (S^4 \times \{1/3\}) = (P_1 \cup H_1 \cup H_2) \times \{1/3\},$   $P_1 \cap (S^4 \times (1/3, 1)) = B'_1 \times (1/3, 1),$   $P_1 \cap (S^4 \times \{1\})) = V_1 \times \{1\},$   $P_2 \cap (S^4 \times \{0, 1\}) = B_2 \times [0, 1],$   $P_2 \cap (S^4 \times \{1\}) = V_2 \times \{1\}.$

We smooth  $P_1 \sqcup P_2$  and obtain the desired restricted height function.

Remark 3.4. By Satoh [Sat04], Proposition 3.3 applies to any twist-spun knot of a classical knot of unknotting number one. In particular, we may take K to be the 2-twist spun trefoil.

**Remark 3.5.** In Proposition 3.3, if we take the 2-sphere  $\Sigma$  to separate the two index-1 critical points, then we can arrange for  $h|_{P_i}$  to have one local minimum, two index-1 critical points, two index-2 critical points, and one local maximum for each i = 1, 2.

### 4. Constructing splitting spheres

In this section we show how to construct distinct splitting spheres in the setting of Theorem 1.2. We first present a lemma that is analogous to [HKM23, Lemma 4.1].

**Lemma 4.1.** Let  $H_1 \sqcup H_2$  be a disjoint union of two 3-dimensional handlebodies properly embedded in  $B^5$ . Suppose that with respect to the radial height function  $h: B^5 \to \mathbb{R}$ , both  $h|_{H_1}$  and  $h|_{H_2}$  have exactly one local minimum and no index-2 or index-3 critical points. Suppose also that  $\partial H_1 \sqcup \partial H_2$  is an unlink in  $\partial B^5 = S^4$ . Then  $H_1 \sqcup H_2$  is smoothly boundary parallel.

Proof. Parameterize  $h: B^5 \to \mathbb{R}$  so that  $h(\partial B^5) = 1$  and h maps the center of  $B^5$  to 0. Isotope  $H_1 \sqcup H_2$  so that both index-0 critical points of  $h|_{(H_1 \sqcup H_2)}$  lie in  $h^{-1}(1/4)$ , and all index-1 critical points lie in disjoint heights  $t_1 < \ldots < t_{m+n}$  in  $h^{-1}(2/3, 3/4)$ .

Now  $H_1 \sqcup H_2$  intersects the 4-sphere  $h^{-1}(1/2)$  in a 2-component unlink of 2-spheres  $S_1 \sqcup S_2$ . As the value of h increases, the surfaces  $S_1 \sqcup S_2$  are successively surgered along 3-dimensional 1-handles. Fix a basepoint of  $B^5$  in  $h^{-1}(1/2)$ , so  $\pi_1(h^{-1}(1/2) \setminus (S_1 \sqcup S_2)) = \langle a, b \rangle$ , where a is a meridian of  $S_1$  and b is a meridian of  $S_2$ .

Consider the effect of surgering  $S_1$  along a 3-dimensional 1-handle at height  $t_1$  whose core arc traces out an arc representing a word w in the letters a, b. (Such an arc is completed to a loop by choosing a fixed whisker from  $S_1$  to the basepoint; w is thus defined only up to conjugation and multiplication by a at either end.) We obtain the relation  $waw^{-1} = a$  in the presentation of  $\pi_1(h^{-1}(t_1 + \epsilon) \setminus (H_1 \sqcup H_2))$ . Similarly if the handle instead has ends on  $S_2$ , then we obtain a relation of the form  $wbw^{-1} = b$ .

Let  $w_1, \ldots, w_m$  be words represented by the cores of the 3-dimensional 1-handles along which  $S_1$  is surgered as h increases. Let  $W_1, \ldots, W_n$  be words represented by the cores of the 3-dimensional 1-handles along which  $S_2$  is surgered. Then we obtain a presentation of  $\pi_1(h^{-1}(1) \setminus (H_1 \sqcup H_2)) = \pi_1(S^4 \setminus \partial(H_1 \sqcup H_2))$  as

$$\langle a, b \mid w_1 a w_1^{-1} a^{-1} = \dots = w_m a w_m a^{-1} = W_1 b W_1^{-1} b^{-1} = \dots = W_n b W_n^{-1} b^{-1} = 1. \rangle$$

On the other hand, by assumption  $\pi_1(S^4 \setminus \partial(H_1 \sqcup H_2))$  is free on two generators. We conclude that  $w_i, W_j = 1$  for all i, j. Since the cores of the 3-dimensional 1-handles attached to  $S_1, S_2$  are 1-dimensional and each  $h^{-1}(t)$  is 4-dimensional, up to smooth isotopy preserving h we can trivialize the 3-dimensional 1-handles. Thus,  $H_1 \sqcup H_2$  is smoothly boundary-parallel.

We are now ready to construct splitting spheres. We restate the main theorem again here.

**Theorem 1.2.** Let  $L = L_1 \sqcup L_2$  be a smooth 2-component unlink in  $S^4$  with  $g(L_1) \ge 4$ . There exist smooth splitting spheres  $S_1, S_2$  for L so that  $S_1, S_2$  are not topologically isotopic in  $S^4 \setminus L$ .

We include a schematic of the proof of Theorem 1.2 in the case  $g(L_1) = 4$ ,  $g(L_2) = 0$  in Figure 6.

Proof. Let  $U_1 \sqcup U_2$  be the 2-component unlink in  $S^4$  with  $U_1$  a genus-4 surface and  $U_2$  a 2-sphere. Let K be the 2-twist spun trefoil. By Proposition 3.3 and Remark 3.4, there is a standard height function  $h: S^5 \to \mathbb{R}$  and a smooth simple 3-knot  $P \subset S^5$  so that P intersects  $h^{-1}(0) \cong S^4$  in a copy of K and  $B(P) = P_1 \sqcup P_2$  intersects  $h^{-1}(0)$  in  $B(K) = K_1 \sqcup K_2$  (with  $P_i \cap S^4 = K_i$ ). Moreover, the height function h on  $S^5$  restricts to P in a Morse function with one local minimum, two index-1 points, two index-2 points, and one local maximum; and restricts to B(P) in a Morse function with one local minimum and local maximum on each  $P_1, P_2$  in addition to four index-1 and four index-2 critical points on  $P_1$ . By Stoltzfus [Sto79], P is a nontrivial 3-knot.

Smoothly isotope P and B(P) to reorder the critical points of  $h|_P$  and  $h|_{B(P)}$  so that index-0 and index-1 points are in  $h^{-1}(-\infty, 0)$  and index-2 and index-3 points are in  $h^{-1}(0, \infty)$ 

Now P intersects  $h^{-1}(0)$  in a trivial genus-2 surface  $\Sigma$  and B(P) intersects  $h^{-1}(0)$  in a Bing double  $B_{\gamma}(\Sigma)$ , where  $\gamma$  is a curve bounding a disk in  $\Sigma$ . By Proposition 2.3,  $B_{\gamma}(\Sigma)$  is smoothly isotopic to the unlink  $U_1 \sqcup U_2$ . Thus, B(P) intersects  $h^{-1}(0)$  in a copy of  $U_1 \sqcup U_2$ , with  $P_i \cap h^{-1}(0) = U_i$ .




FIGURE 6. The proof of Theorem 1.2. We first construct a nontrivial, simple 3-knot P and its Bing double B(P) with the property that  $h|_{B(P)}$  is as in Proposition 3.3 (here we indicate a critical point with \* and write the index). Then by Lemma 4.1, the top and bottom halves (that is, on either side of  $S^4 \times 0$ ) of B(P) are smoothly boundary-parallel, and hence splittable by properly embedded smooth 4-balls in  $h^{-1}(-\infty, 0]$  and  $h^{-1}[0, \infty)$ . These 4-balls intersect  $h^{-1}(0)$  in smooth splitting spheres for an unlink L of a genus-0 and a genus-4 surface. These splitting spheres cannot be topologically isotopic in  $S^4 \setminus L$ , or else the link B(P) would be split, contradicting Lemma 3.2.

Consider the 2-component 3-manifold  $V_1 = B(P) \cap h^{-1}(\infty, 0]$  properly embedded in  $h^{-1}(-\infty, 0] \cong B^5$ . By Lemma 4.1,  $V_1$  is boundary-parallel. Let  $S_1$  be a splitting sphere for  $U_1 \sqcup U_2$  in  $S^4 = h^{-1}(0)$  so that  $V_1$  can be pushed into  $S^4 = h^{-1}(0)$  in the complement of  $S_1$ . Then there exists a trivial 4-ball  $B_1$  properly embedded in  $h^{-1}(\infty, 0]$  with boundary  $S_1$  that is disjoint from  $V_1$ . Similarly, apply Lemma 4.1 to the 2-component 3-manifold  $V_2 = B(P) \cap h^{-1}[0,\infty)$  properly embedded in  $h^{-1}[0,\infty) \cong B^5$  to obtain another splitting sphere  $S_2$  for  $U_1 \sqcup U_2$  in  $h^{-1}(0)$  that bounds a 4-ball  $B_2$  in  $h^{-1}[0,\infty)$  in the complement of  $V_2$ .

Now suppose that there is a topological isotopy of  $S^4 \setminus (U_1 \sqcup U_2)$  that takes  $S_1$  to  $S_2$ . Then we can extend this isotopy over  $h^{-1}(-\infty, 0] \setminus P$ . This isotopy takes the ball  $B_1$  to some other 4-ball  $B'_1$  whose boundary is  $S_2$ . Then  $B'_1 \cup B_2$  forms a splitting sphere for B(P). But since P is nontrivial, by Lemma 3.2 the Bing double B(P) is not split and hence we obtain a contradiction. We thus conclude that the splitting spheres  $S_1, S_2$  for  $U_1 \sqcup U_2$  are not topologically isotopic in  $S^4 \setminus (U_1 \sqcup U_2)$ , completing the proof in the case that  $g(L_1) = 4, g(L_2) = 0$  (here setting  $L_i = U_i$ ).

Now consider perturbing either component  $P_1, P_2$  of B(P) to introduce a new index-1 critical point of  $h|_{P_i}$  in  $h^{-1}(-\infty, 0]$  and a new index-2 critical point of  $h|_{P_i}$  in  $h^{-1}[0, \infty)$ . Now  $S_1, S_2$  are splitting spheres for an unlink obtained by adding a tube to either component of  $U_1 \sqcup U_2$ , and again cannot be topologically isotopic in the unlink complement or else B(P) would be split. By repeating this argument, we obtain non-isotopic splitting spheres in the complement of  $L = L_1 \sqcup L_2$ . **Remark 4.2.** We can adapt the proof of Theorem 1.2 to the case when  $(g(L_1), g(L_2))$  lies in the set  $\{(2, 2), (2, 3), (3, 3)\}$  using the observation of Remark 3.5: the Morse function on B(P) as constructed in Proposition 3.3 can be taken to have two index-1 critical points on each component, so that the unlink cross-section of B(P) is a split union of two genus-2 surfaces, rather than a genus-4 surface and a 2-sphere.

**Corollary 4.3.** Let  $K_1 \sqcup K_2$  be a split, doubly slice link of 2-spheres in  $S^4$ . Let  $\Sigma_1$  be a genus-4 surface obtained by trivially stabilizing  $K_1$  four times. Then the link  $\Sigma_1 \sqcup K_2$  has two smooth splitting spheres that are not topologically isotopic in the complement of  $\Sigma_1 \sqcup K_2$ .

*Proof.* We repeat the proof of Theorem 1.2 in the case  $g(L_1) = 4, g(L_2) = 0$ . Let  $Q_1, Q_2$  be small unknotted 3-knots meeting  $S^4$  in  $K_1, K_2$ , respectively. Then we can connect-sum on  $Q_1, Q_2$  to  $P_1, P_2$  respectively and continue the argument of Theorem 1.2 to obtain two non-isotopic splitting spheres for  $\Sigma_1 \sqcup K_2$ .

As a consequence of the proof of Theorem 1.2, we are now ready to prove Theorem 1.5.

**Theorem 1.5.** Let m, n be non-negative integers with  $m \ge 4$ . There exist two smooth, properly embedded links  $H_1 \sqcup H_2$ ,  $\tilde{H}_1 \sqcup \tilde{H}_2$  of handlebodies in  $B^5$  with  $g(H_1) = g(\tilde{H}_1) = m$ ,  $g(H_2) = g(\tilde{H}_2) = n$  so that the following are true.

- The 2-component links H<sub>1</sub> ⊔ H<sub>2</sub> and H̃<sub>1</sub> ⊔ H̃<sub>2</sub> are each smoothly boundary-parallel in B<sup>5</sup>.
- The boundaries agree componentwise, i.e.,  $\partial H_1 = \partial \tilde{H}_1, \partial H_2 = \partial \tilde{H}_2$ .
- For each i = 1, 2, the handlebodies  $H_i, \tilde{H}_i$  are smoothly isotopic rel. boundary.
- The 2-component links H<sub>1</sub> ⊔ H<sub>2</sub> and H̃<sub>1</sub> ⊔ H̃<sub>2</sub> are not topologically isotopic rel. boundary.

*Proof.* As in the proof of Theorem 1.2 (and indeed, the proof of the main theorem of [HKM23]), we obtain a smooth, nontrivial 3-knot P so that the standard height function  $h: S^5 \to \mathbb{R}$  when restricted to  $B(P) = P_1 \sqcup P_2$  has one local minimum on each component, four index-1 critical points on  $P_1$  four index-2 critical points on  $P_1$ , one local maximum on each component and no other critical points.

We obtain a new 3-link as follows (see Figure 7 for a schematic). Smoothly translate  $P_2$ while preserving  $h|_{P_2}$ , until the resulting 3-knot  $P'_2$  is split from  $P_1$ . Let  $\Sigma_1, \Sigma_2, \Sigma'_2$  denote the intersections of  $P_1, P_2, P'_2$  (respectively) with  $h^{-1}(0)$ . Then  $\Sigma_2, \Sigma'_2$  are each 2-spheres that bound balls in the complement of  $\Sigma_1$ , and hence there is a smooth isotopy of  $S^4$  fixing  $\Sigma_1$  pointwise and taking  $\Sigma'_2$  to  $\Sigma_2$ . Smoothly extend this isotopy over  $S^5$ , and let  $\tilde{P}_1, \tilde{P}_2$  be the images of  $P_1, P'_2$  of this isotopy. Then  $\tilde{P}_1 \sqcup \tilde{P}_2$  is a smooth 3-link intersecting  $h^{-1}(0)$ with  $\tilde{P}_1 \cap h^{-1}(0) = \Sigma_1, \tilde{P}_2 \cap h^{-1}(0) = \Sigma_2$ .

Let  $W^T, W^B$  denote the 5-balls  $h^{-1}[0, \infty)$  and  $h^{-1}(-\infty, 0]$ , respectively. For i = 1, 2 and  $* \in \{T, B\}$ , let  $H_i^* := P_i \cap W^*$  and  $\tilde{H}_i^* = \tilde{P}_i \cap W^*$ . By Lemma 4.1, the links  $H_1^* \sqcup H_2^*$  and  $\tilde{H}_1^* \sqcup \tilde{H}_2^*$  are each boundary-parallel in  $W^*$ . By construction, the genus-4 handlebodies  $H_1^*, \tilde{H}_1^*$  are smoothly isotopic rel. boundary. The 3-balls  $H_2^*, \tilde{H}_2^*$  are boundary-parallel with the same boundary and hence are isotopic rel. boundary in  $W^*$  by [Har]. However, since  $P_1 \sqcup P_2$  is not split and  $\tilde{P}_1 \sqcup \tilde{P}_2$  is split, we conclude that either the pair  $(H_1^T \sqcup H_2^T, \tilde{H}_1^T \sqcup \tilde{H}_2^T)$  are not topologically isotopic rel. boundary in  $W^T$  or the pair  $(H_1^B \sqcup H_2^B, \tilde{H}_1^B \sqcup \tilde{H}_2^B)$  are not topologically isotopic rel. boundary in  $W^B$ .

Perturbing B(P) to add additional (1,2)-pairs to  $h|_{B(P)}$  allows us to increase the genus of the resulting pair of non-isotopic handlebody links, as before.




FIGURE 7. A schematic of the construction of Theorem 1.5. First and second: the Bing double  $B(P) = P_1 \sqcup P_2$  of a nontrivial 3-knot P. We include the second frame to emphasize that B(P) intersects  $h^{-1}(0)$  in an unlink. Third: we translate  $P_2$  (while preserving  $h|_{P_2}$ ) to obtain a 3-knot  $P'_2$  split from  $P_1$ . Fourth: we smoothly isotope  $h^{-1}(0)$  (while keeping the intersection of  $P_1$  with  $h^{-1}(0)$  fixed) and extend over all of  $S^5$  so as to isotope  $P_1, P'_2$  to 3-knots  $\tilde{P}_1, \tilde{P}_2$  whose intersections with  $h^{-1}(0)$  agree with those of  $P_1, P_2$ . For  $i \in \{1, 2\}$  and  $* \in \{B, T\}$ , the handlebodies  $H_i^*$  and  $\tilde{H}_i^*$  are smoothly isotopic rel. boundary in the 5-ball  $W^*$ .

**Remark 4.4.** One can remove the ambiguity from the final conclusion in the proof of Theorem 1.5 via repeated applications of [Hir02] (see discussion in [HKM23]), but this is tedious and not particularly enlightening in practice – though [Hir02] is in principle constructive, in practice it is very difficult to implement the construction.

## References

- [Alt78] Steven C. Althoen. A van Kampen theorem for  $\pi_2$ . J. Pure Appl. Algebra, 10(3):257–269, 1977/78.
- $[BG] Ryan Budney and David Gabai. Knotted 3-balls in <math>S^4$ . ArXiv:1912.09029 [math.GT], Apr. 2021.
- [Har] Daniel Hartman. Unknotting 3-Balls in the 5–Ball. ArXiv:2206.11243 [math.GT], June 2022.
- [Hir02] Susumu Hirose. On diffeomorphisms over surfaces trivially embedded in the 4-sphere. Algebr. Geom. Topol., 2:791–824, 2002.
- [HKM23] Mark Hughes, Seungwon Kim, and Maggie Miller. Knotted handlebodies in the 4-sphere and 5-ball. to appear in J. Eur. Math. Soc., 2023.
- [IKMT] Nobuo Iida, Hokuto Konno, Anubhav Mukherjee, and Masaki Taniguchi. Diffeomorphisms of 4-manifolds with boundary and exotic embeddings. ArXiv:2203.14878 [math.GT], Mar. 2022.
- [KS17] Matthias Kreck and Yang Su. On 5-manifolds with free fundamental group and simple boundary links in S<sup>5</sup>. Geom. Topol., 21(5):2989–3008, 2017.
- [Lau73] F. Laudenbach. Sur les 2-sphères d'une variété de dimension 3. Ann. of Math. (2), 97:57–81, 1973.
- [LP72] François Laudenbach and Valentin Poénaru. A note on 4-dimensional handlebodies. Bull. Soc. Math. France, 100:337–344, 1972.
- [Qui82] Frank Quinn. Ends of maps. III. Dimensions 4 and 5. J. Differential Geometry, 17(3):503-521, 1982.
- [Sat04] Shin Satoh. A note on unknotting numbers of twist-spun knots. Kobe J. Math., 21(1-2):71-82, 2004.
- [Sha68] Julius L. Shaneson. Embeddings with codimension two of spheres in spheres and *H*-cobordisms of  $S^1 \times S^3$ . Bull. Amer. Math. Soc., 74:972–974, 1968.
- [Sta63] John Stallings. On topologically unknotted spheres. Ann. of Math. (2), 77:490–503, 1963.
- [Sto79] Neal W. Stoltzfus. Isometries of inner product spaces and their geometric applications. In Geometric topology (Proc. Georgia Topology Conf., Athens, Ga., 1977), pages 527–541. Academic Press, New York-London, 1979.

BRIGHAM YOUNG UNIVERSITY, PROVO, UT, 84602 USA  $\mathit{Email}\ address:\ \texttt{hughes@mathematics.byu.edu}$ 

SUNGKYUNKWAN UNIVERSITY, SUWON, GYEONGGI, 16419 REPUBLIC OF KOREA Email address: seungwon.kim@skku.edu

STANFORD UNIVERSITY, STANFORD, CA, 94305 USA UNIVERSITY OF TEXAS AT AUSTIN, AUSTIN, TX, 78712 USA Email address: maggie.miller.math@gmail.com