

Density streams in the disc winds of Classical T Tauri stars

P. P. Petrov,¹★ K. N. Grankin,¹ E. V. Babina,¹ S. A. Artemenko,¹ M.M. Romanova,² S. Yu. Gorda,³
A. A. Djupvik,^{4,5} J. F. Gameiro,^{6,7}

¹*Crimean Astrophysical Observatory, p/o Nauchny, 298409, Republic of Crimea*

²*Cornell University, Ithaca, NY 14853, USA*

³*Ural Federal University, 51, Lenin av., Ekaterinburg, Russia, 620000*

⁴*Nordic Optical Telescope, Rambla José Ana Fernández Pérez, 7, 38711 Breña Baja, Spain*

⁵*Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, 8000 Aarhus C, Denmark*

⁶*Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, CAUP, Rua das Estrelas, PT4150-762 Porto, Portugal*

⁷*Departamento de Física e Astronomia, Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre 687, PT4169-007 Porto, Portugal*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

Spectral and photometric variability of the Classical T Tauri stars RY Tau and SU Aur from 2013 to 2022 is analyzed. We find that in SU Aur the $H\alpha$ line's flux at radial velocity $RV = -50 \pm 7 \text{ km s}^{-1}$ varies with a period $P = 255 \pm 5$ days. A similar effect previously discovered in RY Tau is confirmed with these new data: $P = 21.6$ days at $RV = -95 \pm 5 \text{ km s}^{-1}$. In both stars, the radial velocity of these variations, the period, and the mass of the star turn out to be related by Kepler's law, suggesting structural features on the disc plane orbiting at radii of 0.2 AU in RY Tau and 0.9 AU in SU Aur, respectively. Both stars have a large inclination of the accretion disc to the line of sight - so that the line of sight passes through the region of the disc wind. We propose there is an azimuthal asymmetry in the disc wind, presumably in the form of 'density streams', caused by substructures of the accretion disc surface. These streams cannot dissipate until they go beyond the Alfvén surface in the disc's magnetic field. These findings open up the possibility to learn about the structure of the inner accretion disc of CTTS on scales less than 1 AU and to reveal the orbital distances related to the planet's formation.

Key words: Stars: variables: T Tauri, Herbig Ae/Be – Stars: winds, outflows – Line: – Stars: individuals: RY Tau, SU Aur

1 INTRODUCTION

The Classical T Tauri Stars (CTTSs) are commonly recognized as the counterparts of the young Sun, allowing us to study the solar system's past evolution. The formation of a planetary system around a young star occurs simultaneously with the formation of the star itself - when the central star and the surrounding protoplanetary disc are still hidden from the observer by the gas-dust shell (Nixon et al. 2018). By the time we see a star as a CTTS, there are already possible indirect signatures of planets in the surrounding disc. However, detecting these planets is quite difficult. The classical method of precision measurements of radial velocities is hardly applicable for CTTSs: as a result of magnetospheric accretion, hot spots form on the stellar surface, distorting the photospheric spectrum (e.g. Petrov et al. 2011). Unlike long-lived cool spots, these hot spots can change on a time scale of a few days, resulting in jittering the measured radial velocity. Although thousands of exoplanets are currently discovered and their properties and migrations are being investigated, we still have very limited information about planets around CTTSs (see Grandjean et al. 2021 and references therein).

In this article, we consider the possibility of detecting emerging planets in the inner accretion disc by their influence on the disc wind.

The wind properties in CTTS are usually inferred from the kinematics of spatially unresolved, blueshifted forbidden emission lines including [O I], [S II], [N II], [Fe II], [Ne II], as well as P Cygni profiles in strong permitted emission lines, such as He I 10830 Å (see review by Pascucci et al. 2022 and references therein). Analysis of the FUV spectra of CTTS showed that the mass loss is concentrated in the innermost region of the disc, < 1-2 AU (Xu et al. 2021).

In the visible spectral region the most obvious signature of the wind is the variable blue-shifted depression in the $H\alpha$ emission profile. The radial velocity of the blue-shifted depression depends on the wind velocity and the inclination angle of the disc axis to the line of sight.

However, when observing the variability of the $H\alpha$ line formed in the CTTS wind, it is difficult to identify the type of wind responsible for the changes in the line profile. Different types of outflow can start at the boundary between the stellar magnetosphere and the inner accretion disc. The disc wind, accelerated by the magneto-centrifugal force, emanates from the disc surface (Blandford & Payne 1982). Besides, the X-wind (Shu et al. 2016), conical wind (Romanova et al. 2009; Kurosawa & Romanova 2012), magnetospheric ejections (Zanni & Ferreira 2013), and magnetic propeller were discussed (see Romanova & Owocki 2015; Romanova et al. 2018 for review).

Emission line profiles provide information about the dynamics of gas flows around a CTTS. The formation of various types of the $H\alpha$ emission profiles in spectra of CTTS was considered by Kuro-

★ E-mail: petrov@craocrimea.ru

Table 1. The basic parameters of SU Aur and RY Tau

Star	Sp. type	T_{eff} K	R/R_{\odot}	L/L_{\odot}	M/M_{\odot}	\dot{M} $10^{-8} M_{\odot} \text{ yr}^{-1}$	inner disc inclination degrees
SU Aur	G1	5945	2.6 ± 0.4	7.8 ± 1.2	1.7 ± 0.2	$0.5 - 0.6 (\pm 0.4)$	50.9 ± 1
RY Tau	G1	5945	2.9 ± 0.4	9.6 ± 0.5	2.0 ± 0.3	$6.4 - 9.1 (\pm 4.9)$	60.0 ± 1

Stellar parameters: Calvet et al. (2004).

Inner disc inclinations: Labdon et al. (2019) (SU Aur); Perraut et al. (2021) (RY Tau).

sawa et al. (2006) within models of the magnetosphere and the disc wind. They have shown that at a sufficiently high inclination angle of the line of sight, the main feature in the spectral line is blue-shifted absorption due to the disc wind. A similar result has been obtained while modeling the disc wind in the He I (10830) spectral line (Kurosawa et al. 2011).

Models of H α line profiles formed in the magnetosphere and the disk wind of CTTSs were calculated and compared with the observed ones (Kurosawa et al. 2011; Tambovtseva et al. 2014).

Variability of spectral line profiles caused by the gas flows, including the outflows from the boundary between the magnetosphere and the accretion disk, is well documented for some CTTS, e.g. by Sousa et al. 2021 (V2129 Oph), Alencar et al. 2018 (LkCa15), Bouvier et al. 2007b (AA Tau), Bouvier et al. (2023) (GM Aur), while less is known about variations in the disc wind.

There is a class of young stars - FUors - where the disc wind dominates over other types of outflow. The prototype of the class, FU Ori, is a low-mass CTTS with the mass-loss rate in the disc wind $2 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$ (Zhu et al. 2007), while in a CTTS the average value is $10^{-9} M_{\odot} \text{ yr}^{-1}$ (Bouvier et al. 2007a). Herbig et al. (2003) analyzed a series of spectra of FU Ori, obtained in 1997 - 1999. They found that the intensity of the P Cyg absorption in the H α line varied with a period of 14.8 days. A decade later, Powell et al. (2012) analyzed their series of spectral observations of FU Ori of 2007. They confirmed the existence of the periods and refined its values: 13.5 days. The case of FU Ori shows that periodic variations in the disc wind are real and worth to be investigated.

A long time series of spectroscopic observations of a CTTS may potentially reveal rotationally modulated variations in the wind density at certain radial velocities. Such a Doppler probe may provide information about the inner disc structure. In this article we focus on time variability of H α line fluxes at different radial velocities across the broad line profile.

We present the results of spectral and photometric monitoring of RY Tau and SU Aur, both with large inclinations, so that line of sight intersects the region of the disc winds. The stars are bright, and each has been explored in detail before. The stellar parameters, according to Calvet et al. (2004), are presented in Table 1. The stars are relatively massive and may be classified as the intermediate objects between the CTTS and the HAeBe stars.

Our program of spectroscopic and photometric monitoring of RY Tau were started in 2013, then later in 2015 another object, SU Aur, was included. The first results of this program were published in Petrov et al. (2019), hereafter referred as Paper I: from analysis of the variable outflow velocity and the circumstellar extinction of RY Tau we found that the obscuring dust is near the star, at the dust sublimation radius of about 0.2 AU. During events of enhanced outflow, the circumstellar extinction becomes lower. As more data on RY Tau accumulated, we found another effect: the spectral indicators of the accretion and outflows (in H α and Na I lines) vary with a period of about 22 d. The infall and outflow vary in antiphase: an increase

of infall is accompanied by a decrease of outflow, and vice versa. We considered two possible interpretations: 1) quasi-periodic magnetohydrodynamics processes at the disc-magnetosphere boundary in the propeller mode, or 2) a massive planet on inclined orbit in the accretion disc (Petrov et al. 2021).

By now we have accumulated 9 seasons of observation of RY Tau and 7 seasons of SU Aur, including high resolution spectroscopy and photometry. In this paper we investigate variability of the H α profile in spectra of RY Tau and SU Aur, with the aim to confirm the periodic variations and find their possible relationship with the structure of the accretion disc and the disc wind.

The paper is organized as follows. In Section 2 we overview the basic properties of RY Tau and SU Aur. In Section 3 we address briefly the instruments and observation sites. Then, in Section 4 we analyze the variability of brightness and H α line fluxes in RY Tau and SU Aur. The major part is devoted to the frequency analysis of variability of the H α line flux at different radial velocities. Finally, in Section 5 we discuss the results in terms of the disc wind axial asymmetry and the accretion disc structures.

2 BASIC PROPERTIES OF RY TAU AND SU AUR

RY Tau and SU Aur are young stars with emission line spectra, showing signatures of outflows and accretion. Both stars possess accretion discs. Excess continuous radiation is present in the far UV spectrum of both stars, larger in RY Tau than in SU Aur. The mass accretion rates (Table 1) were estimated from the accretion luminosities in the UV spectrum (Calvet et al. 2004).

Inclinations of the *inner* disc were derived from *K*-band interferometry: $60^{\circ} \pm 1^{\circ}$ in RY Tau (Perraut et al. 2021) and $50.9^{\circ} \pm 1^{\circ}$ in SU Aur (Labdon et al. 2019). The inclination of the *outer* disc of RY Tau, as derived from ALMA observations, is $65^{\circ} \pm 0.1^{\circ}$ (Long et al. 2019). *K*-band interferometry of RY Tau by Davies et al. (2020) indicated that the central star is occulted by the disc surface layers close to the sublimation rim. They suggested that the aperiodic photometric variability of RY Tau is likely related to temporal and/or azimuthal variations in the structure of the disc surface layers.

Interferometric and polarimetric observations of SU Aur revealed more complicated structures with several dust tails connected to the Keplerian disk, indicating that late accretion events can still occur (Ginski et al. 2021; Akiyama et al. 2019).

Both stars are rapid rotators. In RY Tau, $v \sin i = 52 \pm 2 \text{ km s}^{-1}$ (Bouvier 1990; Petrov et al. 1999). The period of axial rotation was not detected directly either in photometric data or in variations of spectral line profiles. Assuming the inclination of the stellar rotational axis $i = 60^{\circ}$ (the same as in the accretion disc), stellar radius $2.9 \pm 0.4 R_{\odot}$ and $v \sin i = 52 \text{ km s}^{-1}$, the expected period of the axial rotation may be about 3 days.

In SU Aur, $v \sin i = 60 - 66 \text{ km s}^{-1}$ (Nguyen et al. 2012). The period of axial rotation within 2.7 - 3.0 days was estimated from variations in emission line profiles (Giampapa et al. 1993; Johns &

Table 2. Reported periods in photometric and spectroscopic observations of SU Aur and RY Tau

Star	Period, days	Method	Year of obs.	Reference
SU Aur	1.55; 2.73	phot	1985 - 1986	1
	~ 1.7	phot	1995 - 1999	2
	2.661	phot	2009 - 2010	3
	2.98	spec	1986 - 1990	4
	3.0	spec	1986 - 1992	5
	3.0	spec	1990 - 1994	6
	2.742	spec	1993 - 1994	7
	18.7	phot	1981 - 1991	8
RY Tau	20; 29.4	phot	1993, 1996	9
	23	spec	1975 - 1985	10
	21.6	spec	2013 - 2021	11

References: 1 - [Herbst et al. \(1987\)](#), 2 - [DeWarf et al. \(2003\)](#), 3 - [Cody et al. \(2013\)](#), 4 - [Giampapa et al. \(1993\)](#), 5 - [Johns & Basri \(1995\)](#), 6 - [Petrov et al. \(1996\)](#), 7 - [Strassmeier et al. \(1997\)](#), 8 - [Gahm et al. \(1993\)](#), 9 - [Zajtseva \(2010\)](#), 10 - [Ismailov et al. \(2015\)](#), 11 - [Petrov et al. \(2019, 2021\)](#).

[Basri 1995](#)) and in the red-shifted absorptions of He I and Balmer lines ([Petrov et al. 1996](#)).

RY Tau has an extended jet with a few knots of young dynamical ages ([St-Onge & Bastien 2008](#); [Agra-Amboage et al. 2009](#); [Skinner et al. 2011](#)). The jet is wiggled, which may indicate the presence of an unseen planetary or sub-stellar companion to the star ([Garufi et al. 2019](#)). Interferometric images of the protoplanetary disc around RY Tau at millimeter wavelengths did not reveal planets more massive than $5 M_J$ at distances between 10 and 60 AU ([Isella et al. 2010](#)).

Both RY Tau and SU Aur are irregular variables with long photometric records ([Herbst & Stine 1984](#); [Herbst et al. 1994](#); [Zajtseva et al. 1996](#)). No stable periods in the light variations were recorded, although quasi-periodic variations on different time scales were reported by observers. The most extended series of photometric observations of RY Tau from 1965 to 2000 were analyzed by [Zajtseva \(2010\)](#): periods of 20.0 and 29.4 days were revealed in the data of 1993 and 1996 respectively and interpreted as probably caused by inhomogeneities in the circumstellar disc. A period of about 23 days in variations of emission line intensities in UV and optical spectrum of RY Tau was reported by [Ismailov et al. \(2015\)](#) and interpreted as due to protostellar objects orbiting in the inner accretion disc.

The light curve of SU Aur is characterized by a nearly constant maximum brightness level with a usually small amplitude of variability but interrupted at times by deep fading episodes ([Grankin et al. 2007](#)). Searches for periodic photometric variations gave conflicting results. Various authors reported finding different periods: 1.76 d ([DeWarf et al. 2003](#)), 1.55 d or 2.73 d ([Herbst et al. 1987](#)), 2.7 d ([Strassmeier et al. 1997](#)) or their absence ([Percy et al. 2006](#); [Artemenko et al. 2012](#)). Only analysis of intense space observations performed on the MOST satellite made it possible to reliably detect a low-amplitude periodic process (2.661 days) against the background of a longer and higher-amplitude brightness change ([Cody et al. 2013](#)).

All the reported periods in photometric and spectroscopic observations of SU Aur and RY Tau are summarized in Table 2. More detailed review of the previously reported photometric periods in SU Aur was presented in [Cody et al. \(2013\)](#). In summary: the short periods (< 3 days) in SU Aur indicate the axial rotation of the star, while the longer periods observed in RY Tau are probably related to rotations of the inner region of the accretion disk. A more detailed overview of the basic properties of RY Tau and SU Aur was presented in [Paper I](#).

3 OBSERVATIONS

A major part of our spectral data was obtained with an echelle spectrograph at the 2.6 meters Shajn reflector of the Crimean Astrophysical Observatory (CrAO) and with a fiber-fed echelle spectrograph at the 1.21-meter telescope of Kourouka Observatory of the Ural Federal University. The spectral resolutions $\lambda/\Delta\lambda$ were 27000 and 15000 correspondingly. Two spectral orders, including H α and D Na I lines, were extracted from the echelle frame for analysis. The Crimean observations were carried out in short time series by 4-6 consequent nights every month since September to March, in each season. The spectral regions including H α and D Na I lines were registered.

In one season, 2015-2016, multi-site monitoring of RY Tau and SU Aur was performed, including observations with the ALFOSC grism spectrograph at the 2.56-m Nordic Optical Telescope ($\lambda/\Delta\lambda = 15000$), and the CAFE echelle spectrograph at the 2.2-m telescope at Centro Astronomico Hispano-Aleman (CAHA), $\lambda/\Delta\lambda = 30000$. More information about the observations and data reductions was presented in our previous publication ([Paper I](#)).

As the data from our monitoring of these stars accumulated, the results obtained were published by [Petrov et al. \(2019, 2021\)](#). At present, our dataset contains 193 spectra (193 nights) of RY Tau and 176 spectra of SU Aur, suitable for analysis of H α line variability. There are fewer spectra of the absorption D Na I lines, suitable for analysis, since these lines are noisy on nights with poor image quality, and often contaminated with atmospheric water lines. On most of the nights of spectral observations, BVR photometry of both targets was carried out at a 1.25 m telescope at CrAO. The log of spectral observations and the V-magnitudes for the dates of observations are presented in Tables A1 and A2 in the same format as in the [Paper I](#).

In the next section, we analyze the H α line profile variability in the whole set of data obtained in 2013-2022: 9 seasons of RY Tau and 7 seasons of SU Aur.

4 RESULTS

In both stars, RY Tau and SU Aur, the equivalent width of the H α emission increases as the star gets weaker (Fig. 1). This effect is caused by the circumstellar dust which obscures the star, but not the whole region radiating in H α line. The straight line in Fig. 1 indicates the slope expected for the case of constant H α flux at variable stellar continuum. The observed scatter of H α equivalent widths is due to the intrinsic variability of the H α flux.

Figs. 2 and 3 show stellar brightness and H α flux variability in SU Aur and RY Tau according to our observations, presented in Tables A1 and A2. The H α flux is calculated as $F = EW \times 10^{-0.4(V-10)}$, where the line flux is expressed in units of the continuum flux density of a star with $V = 10$ mag, which is 3.67×10^{-13} erg cm $^{-2}$ s $^{-1}$. The magnitudes in the R band would be more appropriate for this conversion, but only V magnitudes are available to cover the full set of spectroscopic data. However we have checked that for RY Tau the color $V - R$ does not change considerably with brightness: on average, $V - R = 1.1 \pm 0.1$. In the case of SU Aur the probable error in the H α flux, caused by variations in the $V - R$ color, is about 5%, while the H α flux varies by a factor of 3 and more (see Fig. 2).

For analysis of spectral line profiles, the spectra obtained at different instruments were re-binned to get a common sampling of the wavelength scale. In the following analysis of H α line profile variations, we use the astrometric radial velocity scale, assuming that the radial velocity of RY Tau and SU Aur is +18 km s $^{-1}$ ([Petrov et al. 1996, 1999](#)). Radial velocities variations due to possible photospheric inhomogeneities (spots) are expected to be less than a few




Figure 1. Equivalent width of the $H\alpha$ emission as a function of stellar brightness in SU Aur (blue triangles) and RY Tau (red circles).




Figure 2. Light curve (above) and $H\alpha$ flux variations (below) of SU Aur in 2015-2022. The $H\alpha$ flux is expressed in units $3.67 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$.

km s^{-1} . A spot would also result in rotational modulation of stellar brightness. This effect was not found in RY Tau but may have been observed in SU Aur (see Table 2).

4.1 $H\alpha$ profile variability in SU Aur

A sample of 30 $H\alpha$ profiles in SU Aur, observed in different seasons of 2015-2022, is presented in Fig. 4 to show the pattern and amplitude of the line profile variability, typical for this star. All the 176 $H\alpha$ spectra were used to quantify the line profile variability.

A power spectrum of the flux variations was calculated in each radial velocity bin across the $H\alpha$ line profile. This results in a two-dimensional (2D) periodogram in the space "Frequency vs Radial Velocity", which reveals the areas of probable periodic signals. The power spectra were calculated following the procedure CORRELOGRAM described in S. Lawrence (1987). For better visibility of the area of a periodic signal, the power spectra were smoothed to a lower frequency resolution, about 10^{-3} day^{-1} . The 2D periodogram of the $H\alpha$ line flux is shown in Fig. 5. The power of the periodic oscillations is color-coded from blue to red as it increases. Most of the power is




Figure 3. Light curve (above) and $H\alpha$ flux variations (below) of RY Tau in 2013-2022. The $H\alpha$ flux units: $3.67 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$.




Figure 4. A sample of $H\alpha$ line profile variability in SU Aur.

concentrated in the blueward of $H\alpha$, at low frequencies. This region of the periodogram is expanded in the lower part of Fig. 5. The most significant period $P = 255 \pm 5$ days ($1/P = 0.004$) is at a radial velocity $RV = -45 \pm 7 \text{ km s}^{-1}$. The same result was obtained when only high-resolution ($R \geq 27000$) spectra were used in the analysis.

The period of 255 days is seen also in the ratio of fluxes measured at -50 and $+25 \text{ km s}^{-1}$, corresponding to the deepest absorption and the highest emission in the profile. This ratio may be considered as a parameter of the $H\alpha$ profile asymmetry. The power spectrum and convolution of this parameter with a period of 255 days are presented in Fig. 6. The first and the second halves of the time series (2015-2018 and 2019-2022 respectively) are displayed with different symbols, to show that periodic variations persisted throughout the whole data set. The same period is also present in the intensity variations of the blue-shifted absorptions in the D Na I lines.

4.2 $H\alpha$ profile variability in RY Tau

A sample of 30 $H\alpha$ profiles in RY Tau, observed in different seasons between 2018 and 2021, is presented in Fig. 7 showing the pattern and amplitude of the profile variability, typical for this star. All the 193 $H\alpha$ spectra were used to quantify the line profile variability.




Figure 5. Upper: two-dimensional periodogram of the $H\alpha$ line flux in SU Aur. Lower: enlarged fragment of the periodogram. The red spot indicates the area of the most powerful oscillations with a period of 255 ± 5 days.




Figure 6. Top panel: power spectrum of the $H\alpha$ profile asymmetry in SU Aur. Bottom panel: $H\alpha$ profile asymmetry, convolved with the period $P = 255$ days. Dots and circles correspond to the first and the second halves of the time series, correspondingly.




Figure 7. A sample of $H\alpha$ line profile variability in RY Tau.

Periodic variations in the $H\alpha$ profile of RY Tau were previously reported in our publication, based on observations of 2013-2018: $P=21.6$ days at $RV=-100 \text{ km s}^{-1}$ (Petrov et al. 2021). A similar analysis has been done using the whole set of data (9 seasons), corresponding to the years 2013-2022.

Fig. 8 shows a 2D periodogram of the $H\alpha$ flux variations in RY Tau. The most significant period is present at radial velocity $-95 \pm 5 \text{ km s}^{-1}$: the flux in this section of the profile varies with a period $P=21.6$ days ($1/P=0.0463$). The lower part of the Fig. 8 shows this region with better resolution.

The period is also present in the variations of the line profile asymmetry. In the case of RY Tau, the $H\alpha$ line profile is less regular, as compared to SU Aur: the position of the blue-shifted depression is not fixed at the same radial velocity, as well as the position of the red-shifted emission peak (see Fig. 7). Therefore, to measure the line asymmetry we use a more integrated parameter: $(b-r)/(b+r)$, where 'b' and 'r' are fluxes in the 'blue' and 'red' halves of $H\alpha$ profile, relative to zero radial velocity. The most negative values of the asymmetry correspond to the P Cyg type profile.

The power spectrum of $H\alpha$ line asymmetry and convolution of the asymmetry with the period of 21.6 days, are shown in Fig. 9. About the same sinusoidal signal with the period of 21.6 days appears when the asymmetry of $H\alpha$ is expressed as the ratio of flux at radial velocity -95 km s^{-1} to the total flux in the line: F_{-95}/F_{total} . The period of 21.6 days is also detected in the variability of the blue-shifted absorption of the D Na I lines, at a radial velocity of about -100 km s^{-1} (Petrov et al. 2021).

The level of false alarm probability $FAP=0.001$, indicated in Figs. 6 and 9, was estimated via the Fisher's Method of Randomization (Nemec & Nemec 1985). The highest peaks in the power spectra of SU Aur (Fig. 6) and RY Tau (Fig. 9) correspond to the "red spots" in the 2D periodograms in Figs. 5 and 8. In the case of RY Tau, the power spectrum shows two peaks above the $FAP=0.001$. The less significant peak indicates variations with a period of about 34 days in the blue wing of $H\alpha$ emission. In the 2D periodogram (Fig. 8) it looks like a yellow strip at frequency $1/P \sim 0.03$, starting from $RV=-110 \text{ km s}^{-1}$.

5 DISCUSSION

The light variations of SU Aur and RY Tau, observed during our spectroscopic monitoring (Figs. 2 and 3), are typical for these stars. Taking into account the large inclination of the accretion discs, the




Figure 8. Upper: two-dimensional periodogram of the $H\alpha$ line flux in RY Tau. Lower: enlarged fragment of the periodogram. The red spot at $RV = -95 \pm 5$ km s^{-1} indicates the area of the most powerful oscillations with a period of 21.6 days.




Figure 9. Top panel: Power spectrum of $H\alpha$ profile asymmetry in RY Tau. Bottom panel: $H\alpha$ profile asymmetry convolved with the period of 21.6 days. Dots and circles correspond to the first and the second halves of the time series, correspondingly.

short-term dimmings of SU Aur may be explained as a variable extinction in the dusty disc wind (Labdon et al. 2019). In RY Tau, the brightness variations over one magnitude can also be explained as due to circumstellar dust, because the veiling in the visible region of the photospheric spectrum remains low, < 0.1 (Petrov et al. 1999).

In both stars, RY Tau and SU Aur, the periodic changes appear in the blue wings of $H\alpha$ and the D Na I lines, in the radial velocities range that correspond to absorption in the wind. In the case of RY Tau, the period (21.6 days) could be assigned to the quasi-periodic outflows in the regime of the magnetic propeller. Therefore, in our previous publication (Petrov et al. 2021) we discussed two scenarios: 1) a stable magnetic propeller, or 2) a Keplerian motion of a planet at 0.2 AU. In the case of SU Aur, the period is much longer and cannot be identified with the magnetic propeller regime. Therefore, we have to consider whether the periods are related to rotation in the accretion disk.

Accretion discs around CTTSs are expected to have magnetic fields, driving the disc wind out of the disc surface. The poloidal magnetic field lines diverge outward due to the inward flow of matter in the disc plane and are twisted by the disc rotation. Thus, the ionized gas from the disc surface is accelerated into the wind by the magneto-centrifugal forces (Blandford & Payne 1982). Azimuthal asymmetries in the density distribution of the accretion disc lead to asymmetries in the wind.

The masses of RY Tau and SU Aur are known, so we can check whether the periods found are related to the Keplerian motion. In Fig. 10 the straight lines show the Keplerian relationship between the orbital velocity and the orbital period for the masses of RY Tau and SU Aur. The dots with error bars indicate the observed values of the periodic variations of the $H\alpha$ flux, and the radial velocities, at which these changes occur.

The spatial velocity of a gas particle in a disk wind may be considered as the sum of two vectors: toroidal and poloidal (e.g. Kurosawa et al. 2006). A gas particle starts from the surface of the disk and moves along an expanding spiral, rising above the disk plane. The blue-shifted absorption in the $H\alpha$ emission line profile is formed in the wind along the line of sight to the star. The observed radial velocity of this absorption is defined mainly by the wind acceleration in poloidal direction. The projection of the poloidal component to the line of sight depends on two angles: the inclination of the disc axis i , and the inclination of the wind stream to the disc axis (the semi-open angle), which must be $\geq 30^\circ$ in order to start the centrifugal acceleration (Blandford & Payne 1982). Hence, the deprojected poloidal velocity: $V_p \leq RV/\cos(i - 30)$. In Fig. 10, the estimated poloidal velocities of wind are shown with triangles.

In the magneto-centrifugal wind model, the terminal velocity of wind is proportional to the Keplerian velocity at the launching point on the accretion disc (Ferreira et al. 2006). In the case of high inclination i , the line of sight to the star crosses the wind streams launched from different radii in the disc plane. Assuming there is an axial asymmetry in the wind, caused by an azimuthal asymmetry in the structure of the accretion disc at a certain radius R_w , the column density of the disc wind on the line of sight to the star varies with a Keplerian period, corresponding to the orbital radius R_w .

The observed periodic changes in the disc wind do not yet mean the presence of a planet in the disc. Moreover, a planet by itself does not produce a density enhancement in the disc wind. There must be some feature in the accretion disc that rotates at Keplerian velocity and can create a non-axisymmetric dense structure in the wind.

What causes this asymmetry? Images of accretion discs of young stars obtained with the ALMA interferometer show that the surface brightness of the discs is significantly inhomogeneous, with different




Figure 10. Two straight lines show the Keplerian relationship between the orbital velocity and the orbital period for the masses of RY Tau (red dashed line) and SU Aur (blue solid line). Dots with error bars show the observed radial velocities and periods in the $H\alpha$ flux variations in RY Tau and SU Aur. Open triangles indicate the deprojected radial velocities.

substructures, presumably reflecting different instabilities in the disc leading to maxima of gas pressure, or indicating the locations of planet formation (Andrews 2020).

Possible candidates for azimuthal asymmetry in the disc are vortices formed due to the Rossby wave instability (Lovelace et al. 1999; Li et al. 2000). They typically form in parts of the disc with a sharp density gradient, such as at the dead zone boundaries, dust sublimation radii, magnetospheric cavity boundary, or the edges of gaps opened by giant planets (e.g. Armitage & Kley 2019). A vortex represents the region of high pressure and density, which attracts dust particles. The vortex moves around a star approximately with a Keplerian velocity and rotates around the center of the vortex.

The derived orbital radius of 0.2 AU in RY Tau is very close to the inner dusty disc radius of 0.2 – 0.3 AU measured from near IR interferometry (e.g. Perraut et al. 2021) where some density jump is expected and vortices may form. Even higher density jump is expected at the inner dead zone transition (Gammie 1996), where the inner parts of the disc are strongly ionized and matter accretes more rapidly due to the MRI-driven turbulence (e.g. Balbus & Hawley 1991), while accretion rate is low in the external, weakly-ionized parts of the disc. This density jump is shown to be a region where vortices form (e.g. Varniere & Tagger 2006; Lyra & Mac Low 2012). The derived radii of 0.2 – 0.9 AU for the location of the density enhancements in the disc are in the range of dead zones expected in considered stars. Simulations show that typically, the vortex is stable during many rotations (e.g. Li et al. 2001), however detailed radiative simulations by Faure et al. (2015) show that at the dead zone boundaries vortices form episodically and migrate inward in a cyclic manner. On average, the vortex position coincides with the location of the dead zone, and the period of azimuthal rotation will change if the dead zone will change its position.

If the vortex forms at the edge of the gap formed by a giant planet, then its period will be associated with a period of the planet and is expected to be more stable for a long time because a planet migrates slowly (see review by Armitage & Kley 2019). Periods obtained in our work can be connected with one of the above mechanisms.

Outflows from Rossby vortices are expected to have a higher density than those from surrounding parts of the disc and may produce

a local stream of denser matter in the wind. The poloidal magnetic field threading the disc does not prevent the formation of Rossby vortices and driving matter into the wind (Yu & Lai 2013). Moreover, the rotation of the vortex leads to the generation of an additional, azimuthal magnetic field around the vortex which may contribute to driving denser matter into the wind (Lovelace & Romanova 2014; Matilsky et al. 2018).

The key condition for the stability of a stream of dense wind is the prevalence of magnetic pressure in the disc wind. There is the Alfvén surface below which the magnetic pressure exceeds the gas pressure and the kinetic pressure: $B^2/8\pi > P_g + \rho \cdot v^2$. Therefore, the streams of higher density in the wind cannot dissipate until they go above the Alfvén surface. Below the Alfvén surface, the disc wind ‘remembers’ the structure of the disc surface. This is a kind of magnetic ‘memory’ of the disc wind: it is permanently written at the base of the wind and erased above the Alfvén surface.

These results open up new perspectives in studying the structure of the inner regions of the accretion discs. A method for mapping structural features on a moving surface, using changes in spectral line profiles, is known as the Doppler Imaging technique. It is widely applied in mapping spots on stars, where the inclination of the rotational axis and the period of rotation are well known. A similar technique, Doppler tomography, is applied for the analysis of the accretion disc emission line regions in cataclysmic variables (Marsh & Horne 1988).

In the case of Classical T Tauri stars, by studying the density streams in the disc wind, we learn about the inner accretion disc structure and the orbital distances, related to planets formation. The method may be applied for the discs with high inclination, where the line of sight crosses the disc wind.

6 CONCLUSIONS

Spectroscopic monitoring of CTTS provides information about the gas flows around the stars. In the case of high inclination of the accretion disc, when the line of sight crosses the region of the disc wind below Alfvén’s surface, observations may reveal azimuthal asymmetry in the disc wind’s density caused by the structural features of the disc surface. Our time series of RY Tau and SU Aur revealed stable periodic variations in the strength of the blue-shifted absorptions in $H\alpha$ and $D\text{N}\text{I}$ lines. The radial velocity of these variable absorptions, the periods of variations, and the masses of the stars are found to be related by Kepler’s law. The structural details in the discs, presumably associated with Rossby waves, appeared at a distance of 0.2 AU in RY Tau and 0.9 AU in SU Aur. These findings open a possibility to research the structure of the inner accretion disc of a CTTS on scales of the order of one AU or less and to reveal the orbital distances, related to the formation of planets.

ACKNOWLEDGEMENTS

Partly based on observations made with the Nordic Optical Telescope, owned in collaboration by the University of Turku and Aarhus University, and operated jointly by Aarhus University, the University of Turku, and the University of Oslo, representing Denmark, Finland, and Norway, the University of Iceland and Stockholm University at the Observatorio del Roque de los Muchachos, La Palma, Spain, of the Instituto de Astrofísica de Canarias, using ALFOSC, which is provided by the Instituto de Astrofísica de Andalucía (IAA) under a joint agreement with the University of Copenhagen and NOT.

The Crimean observations in 2019-2020 by S. Artemenko, E. Babina, and P. Petrov were supported by a grant from the Russian Science Foundation 19-72-10063. The work of S. Yu. Gorda was partially supported by the Ministry of Science and Higher Education of Russia, topic No. FEUZ-2023-0019. M. M. Romanova was supported in part by NSF grant AST-2009820. J. F. Gameiro was supported by fundação para a Ciência e Tecnologia (FCT) through the research grants UIDB/04434/2020 and UIDP/04434/2020.

We thank R. V. E. Lovelace for useful discussions and an anonymous referee for a thorough review that improved the paper.

DATA AVAILABILITY STATEMENT

The data underlying this article will be shared on reasonable request to the corresponding author [PP].

REFERENCES

- Agra-Amboage V., Dougados C., Cabrit S., Garcia P. J. V., Ferruit P., 2009, *A&A*, 493, 1029
- Akiyama E., Vorobyov E. I., Liu H. B., Dong R., de Leon J., Liu S.-Y., Tamura M., 2019, *AJ*, 157, 165
- Alencar S. H. P., et al., 2018, *A&A*, 620, 195
- Andrews S. M., 2020, *ARA&A*, 58, 483
- Armitage P. J., Kley W., 2019, *From Protoplanetary Disks to Planet Formation*, Saas-Fee Advanced Course, Volume 45. Swiss Society for Astrophysics and Astronomy. Berlin: Springer Berlin Heidelberg
- Artemenko S. A., Grankin K. N., Petrov P. P., 2012, *AstL*, 38, 783
- Balbus S. A., Hawley J. F., 1991, *ApJ*, 376, 214
- Blandford R. D., Payne D. G., 1982, *MNRAS*, 199, 883
- Bouvier J., 1990, *AJ*, 99, 946
- Bouvier J., Alencar S., Harries T., Johns-Krull C. M., Romanova M. M., 2007a, in *Protostars and Planets V*. University of Arizona Press, pp 479–494
- Bouvier J., Alencar S. H. P., Boutelier T., Dougados C., Balog Z., Grankin K., Hodgkin S. T., et al. 2007b, *A&A*, 463, 1017
- Bouvier J., et al., 2023, *A&A*, 672, 5
- Calvet N., Muzerolle J., Briceno C., Hernandez J., Hartmann L., Saucedo J. L., Gordon K. D., 2004, *AJ*, 128, 1294
- Cody A. M., Tayar J., Hillenbrand L. A., Matthews J. M., Kallinger T., 2013, *AJ*, 145, 79
- Davies C. L., et al., 2020, *ApJ*, 897, 31
- DeWarf L. E., Sepinsky J. F., Guinan E. F., Ribas I., Nadalin I., 2003, *ApJ*, 590, 357
- Faure J., Fromang S., Latter H., Meheut H., 2015, *A&A*, 573, 132
- Ferreira J., Dougados C., Cabrit S., 2006, *A&A*, 453, 785
- Gahm G. F., Gullbring E., Fischerstrom C., Lindroos K. P., Loden K., 1993, *A&AS*, 100, 371
- Gammie C., 1996, *ApJ*, 457, 355
- Garufi A., Podio L., Bacciotti F., Antonucci S., et al. 2019, *A&A*, 628, 68
- Giampapa M. S., Basri G. S., Johns C. M., Imhoff C., 1993, *ApJS*, 89, 321
- Ginski C., et al., 2021, *ApJ*, 908, 25
- Grandjean A., et al., 2021, *A&A*, 650, 39
- Grankin K. N., Melnikov S. Y., Bouvier J., Herbst W., Shevchenko V. S., 2007, *A&A*, 461, 183
- Herbig G. H., Petrov P. P., Duemmler R., 2003, *ApJ*, 595, 384
- Herbst W., Stine P. C., 1984, *AJ*, 89, 1716
- Herbst W., et al., 1987, *AJ*, 94, 137
- Herbst W., Herbst D. K., Grossman E. J., Weinstein D., 1994, *AJ*, 108, 1906
- Isella A., Carpenter J. M., Sargent A. I., 2010, *ApJ*, 714, 1746
- Ismailov N. Z., Adigezalzade A. N., Bahaddinova G. R., 2015, *PKAS*, 30, 229
- Johns C., Basri G., 1995, *AJ*, 109, 2800
- Kurosawa R., Romanova M. M., 2012, *MNRAS*, 426, 2901
- Kurosawa R., Harries T. J., Symington N. H., 2006, *MNRAS*, 370, 580
- Kurosawa R., Romanova M. M., Harries T. J., 2011, *MNRAS*, 416, 2623
- Labdon A., et al., 2019, *A&A*, 561, 23
- Li H., Finn J. M., Lovelace R. V. E., A. C. S., 2000, *ApJ*, 533, 1023
- Li H., Colgate S. A., Wendroff B., Liska R., 2001, *ApJ*, 551, 874
- Long F., et al., 2019, *ApJ*, 882, 49
- Lovelace R. V. E., Romanova M. M., 2014, *FIDyR*, 46, 1401
- Lovelace R. V. E., Li H., Colgate S. A., Nelson A. F., 1999, *ApJ*, 513, 805
- Lyra W., Mac Low M.-M., 2012, *ApJ*, 756, 62
- Marsh T., Horne K., 1988, *MNRAS*, 235, 269
- Matilsky L., Dyda S., Lovelace R. V. E., Lii P. S., 2018, *MNRAS*, 480, 3671
- Nemec A. F. L., Nemec J. M. A., 1985, *AJ*, 90, 2317
- Nguyen D. C., Brandeker A., van Kerkwijk M. H., Jayawardhana R., 2012, *ApJ*, 745, 119
- Nixon C. J., King A. R., Pringle J. E., 2018, *MNRAS*, 477, 3273
- Pascucci I., Cabrit S., Edwards S., Gorti U., Gressel O., Suzuki T., 2022, arXiv:2203.10068
- Percy J. R., Gryc W. K., Wong J. C. Y., Herbst W., 2006, *PASP*, 118, 1390
- Perraut K., Gravity Collaboration 2021, [The 20.5th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun](#)
- Petrov P. P., Gullbring E., Ilyin I., Gahm G. F., Tuominen I., Hackman T., Loden K., 1996, *A&A*, 314, 821
- Petrov P., Zajtseva G., Efimov Y., Duemmler R., Ilyin I. V., Tuominen I., Shcherbakov V. A., 1999, *A&A*, 341, 553
- Petrov P., Gahm G. F., Stempels H. C., Walter F. M., Artemenko S. A., 2011, *A&A*, 535, 6
- Petrov P. P., et al., 2019, *MNRAS*, 483, 132
- Petrov P. P., Romanova M. M., Grankin K. N., Artemenko S. A., Babina E. V., Gorda S. Y., 2021, *MNRAS*, 504, 871
- Powell S. L., Irwin M., Bouvier J., Clarke C. J., 2012, *MNRAS*, 426, 3315
- Romanova M. M., Owocki S. P., 2015, *SSRv*, 191, 339
- Romanova M. M., Ustyugova G. V., Koldoba A. V., Lovelace R. V. E., 2009, *MNRAS*, 399, 1802
- Romanova M. M., Blinova A. A., Ustyugova G. V., Koldoba A. V., Lovelace R. V. E., 2018, *NewA*, 62, 94
- S. Lawrence M. J., 1987, *Digital spectral analysis with applications*. Prentice-Hall
- Shu F., Najita J., Ostriker E., Wilkin F., Ruden S., Lizano S., 2016, *ApJ*, 820, 139
- Skinner S., Audard M., Guedel M., 2011, *ApJ*, 737, 19
- Sousa A. P., et al., 2021, *A&A*, 649, 68
- St-Onge G., Bastien P., 2008, *ApJ*, 674, 1032
- Strassmeier K. G., Bartus J., Cutispoto G., Rodono M., 1997, *A&AS*, 125, 11
- Tambovtseva L. V., Grinin V. P., Weigelt G., 2014, *A&A*, 562, 104
- Varniere P., Tagger M., 2006, *A&A*, 446, 13
- Xu Z., Herczeg G., Johns-Krull C., France C., 2021, *ApJ*, 921, 181
- Yu C., Lai D., 2013, *MNRAS*, 429, 2748
- Zajtseva G. V., 2010, *Ap*, 53, 212
- Zajtseva G., Petrov P., Ilyin I., Duemmler R., Tuominen I., 1996, *IBVS*, 4408, 1
- Zanni C., Ferreira J., 2013, *A&A*, 550, 99
- Zhu Z., Hartmann L., Calvet N., Hernandez J., Muzerolle J., Tannirkulam A.-K., 2007, *ApJ*, 669, 483

APPENDIX A: OBSERVATION DATA

This paper has been typeset from a \TeX/L\AA\TeX file prepared by the author.

Table A1. RY Tau V-magnitudes for the dates of spectral observations in 2018-2022. The Heliocentric Julian Date (HJD) in the first column is followed by the site of the observation (CrAO - Crimean Astrophysical Observatory, UrFU - Kourvka astronomical observatory of Ural Federal University), magnitude measured in the V-band, corresponding error and source (CrAO, AAVSO - American Association of Variable Star Observers, Int - value interpolated from AAVSO and CrAO photometry).

HJD-2400000	Site	V	V error	Source of V	HJD-2400000	Site	V	V error	Source of V
58385.4814	CrAO	9.95	0.01	CrAO	59221.1563	CrAO	9.72	0.01	CrAO
58387.4837	CrAO	9.93	0.10	Int	59251.1820	CrAO	10.01	0.01	CrAO
58388.3744	CrAO	9.92	0.01	CrAO	59278.2196	CrAO	9.89	0.01	CrAO
58389.3881	CrAO	9.80	0.01	CrAO	59279.2029	CrAO	9.94	0.01	CrAO
58390.4398	CrAO	9.87	0.01	CrAO	59280.2914	CrAO	9.95	0.10	Int
58410.4249	CrAO	9.46	0.01	CrAO	59288.2045	CrAO	9.94	0.01	CrAO
58413.3276	CrAO	9.58	0.01	CrAO	59475.4156	CrAO	10.00	0.10	Int
58414.4158	CrAO	9.59	0.01	CrAO	59477.4852	CrAO	9.96	0.01	CrAO
58415.4108	CrAO	9.64	0.01	CrAO	59505.4315	CrAO	10.28	0.01	CrAO
58508.1821	CrAO	9.84	0.01	CrAO	59508.3752	CrAO	10.28	0.01	CrAO
58509.1849	CrAO	9.71	0.01	CrAO	59531.2225	CrAO	10.44	0.01	CrAO
58511.3288	CrAO	9.80	0.10	Int	59532.2283	CrAO	10.65	0.01	AAVSO
58539.3164	CrAO	10.78	0.01	CrAO	59533.2331	CrAO	10.67	0.10	Int
58567.2115	CrAO	10.78	0.10	Int	59534.2230	CrAO	10.69	0.01	CrAO
58733.4181	CrAO	10.01	0.01	CrAO	59544.2192	CrAO	10.37	0.01	CrAO
58734.4325	CrAO	9.86	0.01	CrAO	59545.2085	CrAO	10.30	0.10	Int
58735.4190	CrAO	10.03	0.01	CrAO	59586.3711	CrAO	10.10	0.10	Int
58736.4114	CrAO	10.02	0.01	CrAO	59599.3919	CrAO	10.30	0.10	Int
58737.4201	CrAO	9.98	0.01	CrAO	59624.1807	CrAO	10.38	0.01	CrAO
58765.4148	CrAO	9.74	0.01	CrAO	59632.2225	CrAO	9.94	0.01	CrAO
58766.3949	CrAO	9.65	0.01	CrAO	59652.2014	CrAO	10.03	0.01	CrAO
58767.4075	CrAO	9.60	0.01	CrAO	59654.2259	CrAO	10.23	0.01	CrAO
58768.4256	CrAO	9.56	0.01	CrAO	59663.2065	CrAO	9.76	0.01	CrAO
58793.4984	CrAO	9.80	0.01	CrAO	59665.2201	CrAO	9.72	0.01	CrAO
58794.4122	CrAO	9.81	0.10	Int	59666.2159	CrAO	9.77	0.01	CrAO
58795.3102	CrAO	9.82	0.01	CrAO					
58796.4116	CrAO	9.78	0.01	CrAO					
58822.1974	CrAO	10.07	0.01	CrAO					
58823.1794	CrAO	9.95	0.01	CrAO					
58824.1764	CrAO	9.83	0.01	CrAO					
58825.1786	CrAO	9.87	0.01	CrAO					
58826.3344	CrAO	9.96	0.01	CrAO					
58832.1747	CrAO	9.72	0.01	CrAO					
58834.1948	CrAO	9.74	0.01	CrAO					
58835.1767	CrAO	9.72	0.01	CrAO					
58836.1848	CrAO	9.72	0.01	CrAO					
58837.1782	CrAO	9.67	0.01	CrAO					
59094.4484	CrAO	10.21	0.01	CrAO					
59095.4479	CrAO	10.15	0.10	Int					
59096.4027	CrAO	10.10	0.01	AAVSO					
59118.4230	CrAO	10.09	0.10	Int					
59119.4668	CrAO	10.09	0.01	CrAO					
59120.3924	CrAO	10.10	0.01	CrAO					
59121.4180	CrAO	10.10	0.01	CrAO					
59123.4526	CrAO	10.24	0.01	AAVSO					
59150.4246	CrAO	9.47	0.01	CrAO					
59151.4660	CrAO	9.43	0.01	CrAO					
59151.5010	UrFU	9.43	0.01	CrAO					
59157.4042	UrFU	9.50	0.10	Int					
59174.3277	CrAO	9.70	0.10	Int					
59176.4235	CrAO	9.73	0.01	CrAO					
59178.4348	CrAO	9.77	0.01	CrAO					
59189.1893	CrAO	10.27	0.01	CrAO					
59190.2879	CrAO	10.03	0.01	CrAO					
59194.4090	UrFU	9.75	0.10	Int					
59195.4826	UrFU	9.73	0.10	Int					
59203.5020	UrFU	9.54	0.10	Int					
59217.1873	CrAO	9.66	0.01	CrAO					

Table A2. SU Aur V-magnitudes for the dates of spectral observations in 2018-2022. The Heliocentric Julian Date (HJD) in the first column is followed by the site of the observation (CrAO - Crimean Astrophysical Observatory, UrFU - Kourvka astronomical observatory of Ural Federal University), magnitude measured in the V-band, corresponding error and source (CrAO, AAVSO - American Association of Variable Star Observers, Int - value interpolated from AAVSO and CrAO photometry).

HJD-2400000	Site	V	V error	Source of V	HJD-2400000	Site	V	V error	Source of V
58385.5254	CrAO	9.35	0.01	CrAO	59095.5174	CrAO	9.47	0.10	Int
58387.5306	CrAO	9.55	0.01	CrAO	59096.4822	CrAO	9.50	0.10	Int
58388.4419	CrAO	9.33	0.01	CrAO	59119.5152	CrAO	9.86	0.01	CrAO
58389.4523	CrAO	9.36	0.01	CrAO	59120.4622	CrAO	10.00	0.01	CrAO
58390.5045	CrAO	9.40	0.01	CrAO	59121.4901	CrAO	10.03	0.01	CrAO
58410.4894	CrAO	9.34	0.01	CrAO	59150.5046	CrAO	9.50	0.01	CrAO
58413.3925	CrAO	9.32	0.01	CrAO	59151.4993	UrFU	9.56	0.01	CrAO
58414.4808	CrAO	9.35	0.01	CrAO	59151.5310	CrAO	9.56	0.01	CrAO
58415.4753	CrAO	9.34	0.01	CrAO	59174.3962	CrAO	9.64	0.10	Int
58508.2357	CrAO	9.53	0.01	CrAO	59176.4926	CrAO	9.65	0.01	CrAO
58509.2595	CrAO	9.50	0.01	CrAO	59189.2371	CrAO	9.58	0.01	CrAO
58511.3531	CrAO	9.40	0.10	Int	59190.3823	CrAO	9.64	0.01	CrAO
58521.3400	UrFU	9.35	0.10	Int	59217.2541	CrAO	9.68	0.01	CrAO
58535.2368	UrFU	9.51	0.01	CrAO	59218.2252	CrAO	9.68	0.01	CrAO
58539.2519	CrAO	9.44	0.01	CrAO	59219.2124	CrAO	9.70	0.01	CrAO
58559.2333	UrFU	9.62	0.02	AAVSO	59221.2569	CrAO	9.83	0.01	CrAO
58567.2619	CrAO	9.60	0.10	Int	59249.2494	CrAO	9.40	0.01	CrAO
58568.2508	CrAO	9.52	0.01	CrAO	59257.3505	UrFU	9.33	0.10	AAVSO
58733.4841	CrAO	9.40	0.01	CrAO	59275.3060	UrFU	9.36	0.10	Int
58734.4970	CrAO	9.44	0.01	CrAO	59279.2716	CrAO	9.34	0.01	CrAO
58735.4834	CrAO	9.34	0.01	CrAO	59287.2796	UrFU	9.34	0.10	Int
58736.4770	CrAO	9.26	0.01	CrAO	59288.2700	CrAO	9.35	0.01	CrAO
58737.4841	CrAO	9.27	0.01	CrAO	59288.2886	UrFU	9.35	0.01	CrAO
58765.4834	CrAO	9.32	0.01	CrAO	59296.2526	UrFU	10.03	0.10	AAVSO
58766.4600	CrAO	9.33	0.01	CrAO	59318.2247	UrFU	9.90	0.10	Int
58767.4518	CrAO	9.29	0.01	CrAO	59477.5286	CrAO	9.72	0.01	CrAO
58768.4906	CrAO	9.34	0.01	CrAO	59505.4855	CrAO	9.55	0.01	CrAO
58793.4281	CrAO	9.27	0.10	Int	59533.2996	CrAO	9.50	0.10	Int
58794.4564	CrAO	9.28	0.01	CrAO	59544.3023	CrAO	9.73	0.01	CrAO
58795.3754	CrAO	9.29	0.01	CrAO	59586.4188	CrAO	9.60	0.10	Int
58796.4773	CrAO	9.24	0.01	CrAO	59598.2719	CrAO	9.58	0.01	CrAO
58822.2870	CrAO	9.34	0.01	CrAO	59599.4372	CrAO	9.58	0.10	Int
58823.2465	CrAO	9.25	0.01	CrAO	59624.2683	CrAO	9.53	0.01	CrAO
58824.2414	CrAO	9.29	0.01	CrAO	59628.2886	CrAO	9.49	0.01	CrAO
58825.2342	CrAO	9.28	0.01	CrAO	59630.2814	CrAO	9.45	0.10	Int
58826.3785	CrAO	9.27	0.01	CrAO	59632.2727	CrAO	9.38	0.01	CrAO
58832.2412	CrAO	9.27	0.01	CrAO	59652.2676	CrAO	9.44	0.01	CrAO
58834.2621	CrAO	9.26	0.01	CrAO	59663.2538	CrAO	9.43	0.01	CrAO
58835.2633	CrAO	9.24	0.01	CrAO	59665.2656	CrAO	9.43	0.01	CrAO
58836.2718	CrAO	9.23	0.01	CrAO	59666.2610	CrAO	9.55	0.01	CrAO
58837.2858	CrAO	9.21	0.01	CrAO	59667.2560	CrAO	9.53	0.01	CrAO
58847.3812	UrFU	9.32	0.10	Int					
58851.2413	CrAO	9.36	0.01	CrAO					
58862.3010	CrAO	9.26	0.01	CrAO					
58864.2685	CrAO	9.24	0.01	CrAO					
58865.2706	CrAO	9.25	0.01	CrAO					
58877.2560	UrFU	9.27	0.01	Int					
58889.3070	CrAO	9.41	0.01	CrAO					
58890.2680	CrAO	9.45	0.01	CrAO					
58914.2987	UrFU	9.55	0.10	Int					
58916.2893	CrAO	9.51	0.01	CrAO					
58918.2852	CrAO	9.37	0.01	CrAO					
58923.2899	CrAO	9.35	0.01	CrAO					
58925.2692	CrAO	9.55	0.01	CrAO					
58926.2454	CrAO	9.63	0.01	CrAO					
58932.2576	UrFU	9.60	0.10	Int					
58936.2847	UrFU	9.60	0.10	Int					
59094.5135	CrAO	9.45	0.01	CrAO					