ON THE DIRECT IMAGE OF THE ADJOINT LINE BUNDLE

INDRANIL BISWAS, FATIMA LAYTIMI, D. S. NAGARAJ, AND WERNER NAHM

ABSTRACT. We give an algebro-geometric proof of the fact that for a smooth fibration $\pi : X \longrightarrow Y$ of projective varieties, the direct image $\pi_*(L \otimes K_{X/Y})$ of the adjoint line bundle of an ample (respectively, nef and π -strongly big) line bundle L is ample (respectively, nef and big).

1. INTRODUCTION

Throughout we work over the field \mathbb{C} of complex numbers. For the standard notation used here the reader is referred to [1], [2].

We first give an algebraic proof of the following:

Theorem 1.1. Let $\pi : X \longrightarrow Y$ be a smooth fibration of smooth projective varieties and $K_{X/Y}$ the relative canonical line bundle for it. For any ample line bundle L on X the direct image $\pi_*(L \otimes K_{X/Y})$ is either zero or an ample vector bundle.

Mourougane in [3] has proved by analytic methods that the vector bundle $\pi_*(L \otimes K_{X/Y})$ is Griffith positive (hence ample).

We define below the notion of π -strongly big line bundle (see, Definition 3.1) and prove the following:

Theorem 1.2. Let $\pi : X \longrightarrow Y$ be a smooth fibration of projective varieties. If L is a π -strongly big line bundle on X, then the vector bundle $\pi_*(L \otimes K_{X/Y})$ is big. Moreover, if L is nef and π -strongly big, then $\pi_*(L \otimes K_{X/Y})$ is also nef and big.

2. Proof of Theorem 1.1

We need to recall following known results.

²⁰¹⁰ Mathematics Subject Classification. 14F17, 14J60.

Key words and phrases. Nef bundle; ample bundle; adjoint system; relative ample bundle.

Theorem 2.1 ([1, Theorem 4.1.10]). Let ξ be a line bundle on a projective variety Y. Fix a positive integer d. Then there exists a projective variety \widetilde{Y} , a finite surjective morphism $f : \widetilde{Y} \longrightarrow Y$ and a line bundle M on \widetilde{Y} , such that $f^*\xi \simeq M^{\otimes d}$.

Theorem 2.2 ([4, Theorem 2.43]). Let $\pi : X \longrightarrow Y$ be a smooth fibration of projective varieties. If L is semi-ample, then $\pi_*(L \otimes K_{X/Y})$ is nef.

Take π as above. If L is an ample bundle on X, then the restriction $L|_{X_y}$ of L to the fiber $X_y := \pi^{-1}(y)$ over $y \in Y$ is ample, and hence from Kodaira vanishing theorem it follows that $\pi_*(L \otimes K_{X/Y})$ is either 0 or a vector bundle on Y.

Now we start the proof Theorem 1.1. Let L be an ample line bundle on X and ξ a line bundle on Y. Then there is an integer d > 0 such that $L^d \otimes \pi^* \xi^*$ is ample, where ξ^* is the dual of the line bundle ξ . We assume that ξ is ample. By Theorem 2.1 there exists a projective variety \widetilde{Y} , a finite surjective morphism $f : \widetilde{Y} \longrightarrow Y$ and a line bundle M on \widetilde{Y} , such that

$$(2.1) f^*\xi = M^{\otimes d}.$$

Note that $f^*\xi$ is ample because f is a finite map and ξ is ample. Hence from (2.1) it follows that M is ample (recall that d > 0). We have a commutative diagram:

(2.2)
$$\begin{array}{ccc} \widetilde{X} & \xrightarrow{f} & X \\ & & \downarrow_{\widetilde{\pi}} & & \downarrow_{\pi} \\ & & \widetilde{Y} & \xrightarrow{f} & Y \end{array}$$

The line bundle $L^d \otimes \pi^* \xi^*$ is ample if and only if $\tilde{f}^*(L^d \otimes \pi^* \xi^*)$ is ample, because \tilde{f} is finite. We have

$$\begin{split} \widetilde{f}^*(L^d \otimes \pi^*\xi^*) &= (\widetilde{f}^*L^d) \otimes (\widetilde{f}^*\pi^*\xi^*) \\ &= (\widetilde{f}^*L^d) \otimes (\widetilde{\pi}^*f^*\xi^*) \\ &= (\widetilde{f}^*L^d) \otimes (\widetilde{\pi}^*(M^*)^d) \\ &= ((\widetilde{f}^*L) \otimes (\widetilde{\pi}^*M^*))^d. \end{split}$$

Set $\mathcal{L} = \widetilde{f}^* L \otimes \widetilde{\pi}^* M^*$; then \mathcal{L} is ample.

From Theorem 2.2 we know that $\tilde{\pi}_*(\mathcal{L} \otimes K_{\widetilde{X}/\widetilde{Y}})$ is nef. Using projection formula,

$$\widetilde{\pi}_*(\mathcal{L} \otimes K_{\widetilde{X}/\widetilde{Y}}) = \widetilde{\pi}_*(\widetilde{f}^*L \otimes K_{\widetilde{X}/\widetilde{Y}}) \otimes M^*.$$

The vector bundle $\tilde{\pi}_*((\tilde{f}^*L) \otimes K_{\tilde{X}/\tilde{Y}}) = \tilde{\pi}_*((\tilde{f}^*L) \otimes K_{\tilde{X}/\tilde{Y}}) \otimes M^* \otimes M$ is ample, since the tensor product of a nef vector bundle and an ample vector bundle is ample (see [5, Lemma 1.3]).

But $\tilde{\pi}_*((\tilde{f}^*L) \otimes K_{\tilde{X}/\tilde{Y}}) = f^*(\pi_*(L \otimes K_{X/Y}))$ by the commutative diagram in (2.2). Hence $\pi_*(L \otimes K_{X/Y})$ is ample. This completes the proof of Theorem 1.1.

Remark 2.3. For any ample line bundle L on a smooth projective variety Z, the vanishing theorem of Kodaira says that $H^i(Z, L \otimes K_Z) =$ 0 for all $i \geq 1$, where K_Z is the canonical line bundle of Z [1]. Let $\pi : X \longrightarrow Y$ be a smooth fibration of smooth projective varieties. Let L be an ample line bundle on X. From Kodaira vanishing theorem it now follows that $R^i \pi_*(L \otimes K_{X/Y}) = 0$ for all $i \geq 1$.

Remark 2.4. In Theorem 1.1 set $X = \mathbb{P}(E)$, where E is a vector bundle on Y. Let L be an ample line bundle on $\mathbb{P}(E)$. Assume that $\pi_*(L \otimes K_{X/Y}) \neq 0$. Then there is a unique positive integer d and a unique line bundle L_0 on Y such that

$$L \otimes K_{X/Y} = \mathcal{O}_{\mathbb{P}(E)}(d) \otimes \pi^* L_0.$$

Hence using the projection formula, we have

$$\pi_*(L \otimes K_{X/Y}) = \pi_*(\mathcal{O}_{\mathbb{P}(E)}(d) \otimes \pi^*L_0) = \operatorname{Sym}^d(E) \otimes L_0,$$

where $\operatorname{Sym}^{d}(E)$ is the *d*-th symmetric product of *E*. Now Theorem 1.1 says that $\operatorname{Sym}^{d}(E) \otimes L_{0}$ is ample.

When Y is a smooth projective curve, $\operatorname{Sym}^{d}(E) \otimes L_{0}$ is ample if and only if $\mu_{\min}(\operatorname{Sym}^{d}(E) \otimes L_{0}) > 0$ [Ha, p. 84, Theorem 2.4].

3. Proof of Theorem 1.2

Definition 3.1. Let $\pi : X \longrightarrow Y$ be a smooth fibration of smooth projective varieties. A line bundle L on X is said to be π -strongly big if there is an effective divisor D on Y with simple normal crossing support (see [1, Definition 4.1.1]) such that $L^d \otimes \pi^*(\mathcal{O}_Y(D))^*$ is ample for some integer d > 0, where $\pi^*(\mathcal{O}_Y(D))^* = \pi^*(\mathcal{O}_Y(-D))$.

The following proposition states some properties of π -strongly big line bundles.

Proposition 3.2. Take π and L as in Definition 3.1. If L is π -strongly big, then it is big and π -ample. In addition, if L is nef, then $\pi_*(L \otimes K_{X/Y})$ is also nef.

Proof. Since $\pi^*(\mathcal{O}_Y(D)) = \mathcal{O}_X(\pi^{-1}(D))$, and $\pi^{-1}(D)$ is effective, it follows from a characterization of big divisors (see [1, p. 141, Corollary

2.2.7]) that L is big. On the other hand, by definition, and the fact that $L^d(-\pi^*\mathcal{O}_Y(D))|_{X_y} \simeq L^d|_{X_y}$ for all $y \in Y$, we obtain that L is π -ample; here X_y denotes the fiber of π over y. With these observations, the second statement of the proposition follows from [3, Theorem 1]. \Box

Before proving Theorem 1.2 we need the following lemma which is a consequence Kawamata coverings result (see [1, Proposition 4.1.12]).

Lemma 3.3. In Theorem 2.1 assume Y is non-singular and $\xi = \mathcal{O}_Y(D)$, where D is an effective divisor on Y with simple normal crossing support (i.e., the reduced divisor D_{red} is normal crossing divisor). Then given any integer d > 0, there is a finite surjective morphism $f : \tilde{Y} \longrightarrow Y$, and a line bundle $M = \mathcal{O}_{\tilde{Y}}(\tilde{D})$ on \tilde{Y} , such that $f^*\xi \simeq M^d$, and \tilde{D} is effective.

For the proof of Theorem 1.2 we proceed as in the proof of Theorem 1.1. Let L be a π -strongly big line bundle on X. Then there exists an effective divisor D on Y with simple normal crossing support and an integer d > 0, such that $L^d \otimes \pi^* \mathcal{O}_Y(D)^*$ is an ample line bundle. By Lemma 3.3 there exists a projective variety \widetilde{Y} , a finite surjective morphism $f : \widetilde{Y} \longrightarrow Y$ and a line bundle $M = \mathcal{O}_{\widetilde{Y}}(\widetilde{D})$ with \widetilde{D} effective, such that $f^*(\mathcal{O}_Y(D)) = M^d$. We have the commutative diagram:

$$(3.1) \qquad \begin{array}{c} \widetilde{X} & \stackrel{f}{\longrightarrow} X \\ \downarrow_{\widetilde{\pi}} & \qquad \downarrow_{\pi} \\ \widetilde{Y} & \stackrel{f}{\longrightarrow} Y \end{array}$$

The line bundle $L^d \otimes \pi^*(\mathcal{O}_Y(D))^*$ is ample if and only if $\widetilde{f}^*(L^d \otimes \pi^*(\mathcal{O}_Y(D))^*)$ is ample. We have

$$\widetilde{f}^{*}(L^{d} \otimes \pi^{*}(\mathcal{O}_{Y}(D))^{*}) = (\widetilde{f}^{*}L^{d}) \otimes (\widetilde{f}^{*}\pi^{*}(\mathcal{O}_{Y}(D))^{*})$$

$$= (\widetilde{f}^{*}L^{d}) \otimes (\widetilde{\pi}^{*}(f^{*}\mathcal{O}_{Y}(D))^{*})$$

$$= (\widetilde{f}^{*}L^{d}) \otimes (\widetilde{\pi}^{*}M^{*})^{d}$$

$$= ((\widetilde{f}^{*}L) \otimes (\widetilde{\pi}^{*}M^{*}))^{d}.$$

Set $\mathcal{L} = (\tilde{f}^*L) \otimes \tilde{\pi}^*M^*$. Then \mathcal{L} is ample, and hence from Theorem 1.1 it follows that $\tilde{\pi}_*(\mathcal{L} \otimes K_{\tilde{X}/\tilde{Y}})$ is ample. Now

$$\widetilde{\pi}_*(\mathcal{L} \otimes K_{\widetilde{X}/\widetilde{Y}}) = \widetilde{\pi}_*((\widetilde{f}^*L) \otimes K_{\widetilde{X}/\widetilde{Y}}) \otimes M^*,$$

where $M = \mathcal{O}_{\widetilde{Y}}(\widetilde{D})$ is effective. Then $\widetilde{\pi}_*((\widetilde{f}^*L) \otimes K_{\widetilde{X}/\widetilde{Y}})$ is big, because an ample vector bundle tensored with an effective line bundle is big (see [2, Example 6.1.23]).

But $\widetilde{\pi}_*(\mathcal{L} \otimes K_{\widetilde{X}/\widetilde{Y}}) = f^*(\pi_*(L \otimes K_{X/Y}))$ by (3.1). Since f is finite we conclude that $\pi_*(L \otimes K_{X/Y})$ is big.

Moreover, if L is π -strongly big and nef, by Proposition 3.2 the direct image $\pi_*(\mathcal{L} \otimes K_{\widetilde{X}/\widetilde{Y}})$ is nef. This completes the proof.

4. Some observations

Proposition 4.1. Let L be a line bundle on a non-singular projective variety X and E a vector bundle on X. Assume that the vector bundle $E^{\otimes s} \otimes L$ is generated by its global sections for every integer $s \gg 0$. Then E is nef.

Proof. Assume that E is not nef. Then a criterion for nef bundles (see [2, Proposition 6.1.18]) gives that there is a non-constant morphism $f : C \longrightarrow X$ from a non-singular projective curve C, and a line bundle M on C of degree n < 0, such that there is a surjective map

$$f^*E \longrightarrow M.$$

So for all s > 0 there is a surjection

$$(f^*E)^{\otimes s} \otimes f^*L = f^*(E^{\otimes s} \otimes L) \longrightarrow M^{\otimes s} \otimes (f^*L).$$

Note that the degree of the line bundle $M^{\otimes s} \otimes (f^*L)$ is n.s + d, where d is the degree of the line bundle f^*L . Since n < 0, we have n.s+d < 0 for all large s. The bundle $(f^*E)^{\otimes s} \otimes (f^*L)$ is generated by its global sections, by hypothesis, for all large s. Hence it cannot admit a surjective map to a negative line bundle. Thus we get a contradiction to the assumption that E is not nef. This proves the proposition. \Box

We give an example to show that the assumptions do hold.

Example 4.2. Let Z and Y be two smooth projective varieties, and let $X = Y \times Z$. Let $\pi_1 : X \longrightarrow Y$ and $\pi_2 : X \longrightarrow Z$ be the natural projections. If L_1 is a big line bundle on Y, and L_2 is an ample line bundle on Z, then the line bundle $M = (\pi_1^*L_1) \otimes (\pi_2^*L_2)$ is big on X and $(\pi_1)_*(M \otimes K_{X/Y}) = \operatorname{H}^0(Z, L_2 \otimes K_Z) \otimes L_1$ is big provided $\operatorname{H}^0(Z, L_2 \otimes K_Z) \neq 0$.

Remark 4.3. Note that in Example 4.2 the line bundle M is π_1 strongly big provided the big bundle L_1 satisfies the following property: There is an integer d > 0, and a divisor D on Y with a simple normal
crossing support, such that $L^d(-D)$ is ample on Y. Let $\pi : X \longrightarrow Y$ be a smooth fibration of smooth projective varieties. Next example shows that when L is nef and π -big, the direct image $\pi_*(L \otimes K_{X/Y})$ is not necessarily big.

Example 4.4. Let Z and Y be two smooth projective varieties, and let $X = Y \times Z$. Let $\pi_1 : X \longrightarrow Y$ and $\pi_2 : X \longrightarrow Z$ be the natural projections. Let M be an ample line bundle on Z. Then the line bundle π_2^*M is nef and π_1 -ample (and hence π_1 -big) on X and $(\pi_1)_*((\pi_2^*M) \otimes K_{X/Y}) = \operatorname{H}^0(Z, M \otimes K_Z) \otimes \mathcal{O}_Y$ is a trivial bundle, if $\operatorname{H}^0(Z, L_2 \otimes K_Z) \neq 0$, which is only nef but not big.

Definition 4.5. Let $\pi : X \longrightarrow Y$ be a smooth fibration of smooth projective varieties. A line bundle L on X is said to be π -weakly big if there is an effective divisor D on Y such that $L^d \otimes f^*(\mathcal{O}_Y(D))^*$ is ample for some integer d > 0, where $f^*(\mathcal{O}_Y(D))^* = f^*(\mathcal{O}_Y(-D))$.

Remark 4.6. For a $\pi : X \longrightarrow Y$ smooth fibration of smooth projective varieties a π -strongly big line bundle is π -weakly big but is π -weakly big then it may not be π -strongly big.

Proposition 4.7. Let $\pi : X \longrightarrow Y$ as in Theorem 1.1 and L be a π -weakly big line bundle on X. Then there is an integer r > 0 such that the bundles $\pi_*(L^{rm} \otimes K_{X/Y})$ is big for all m > 0.

Proof: By definition of π -weakly big line bundle there is a effective divisor D on Y and a positive integer r such that $L^r \otimes f^*(\mathcal{O}_Y(D))^*$ is an ample and hence $L^{rm} \otimes f^*(\mathcal{O}_Y(mD))^*$ is ample for all m > 0. By Theorem 1.1 $\pi_*(L^{rm} \otimes f^*(\mathcal{O}_Y(mD))^* \otimes K_{X/Y})$ is ample for all m > 0. But by projection formula

$$\pi_*(L^{rm} \otimes \otimes K_{X/Y}) = \pi_*(L^{rm} \otimes f^*(\mathcal{O}_Y(mD))^* \otimes K_{X/Y}) \otimes \mathcal{O}_Y(mD)$$

for all m > 0. Thus $\pi_*(L^{rm} \otimes \otimes K_{X/Y})$ is big for m > 0.

Conjecture: Let $\pi : X \longrightarrow Y$ as in Theorem 1.1 and L be a big and nef line bundle on X. Then $\pi_*(L \otimes K_{X/Y})$ is nef and big.

ACKNOWLEDGEMENTS

DSN thanks CEMPI of Lille university for the financial support.

References

- [Ha] R. Hartshorne, Ample vector bundles on curves, Nagoya Math. J. 43 (1971), 73–89.
 - R. Lazarsfeld, *Positivity in algebraic geometry*. I, Ergeb. Math. Grenzgeb. 48, Springer-Verlag, Berlin, 2004.

- [2] R. Lazarsfeld, *Positivity in algebraic geometry. II*, Ergeb. Math. Grenzgeb. 49, Springer-Verlag, Berlin, 2004.
- [3] C. Mourougane, Image direct de fibré en droites adjoints, Proc. Res. Ins. Math. Sci. 33 (1997), 893–916.
- [4] E. Viehweg, Quasi-projective Moduli for Polarised Manifolds, Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge, A Series of Modern Surveys in Mathematics, Springer, 1995.
- [5] F. Laytimi, On Degeneracy Loci, Int. Jour. of Math. 7 (1996), 745–754.

School of Mathematics, Tata Institute of Fundamental Research, 1 Homi Bhabha Road, Colaba, Mumbai 400005

 $Email \ address: \verb"indranil@math.tifr.res.in"$

Mathématiques - bât. M2, Université Lille 1, F-59655 Villeneuve d'Ascq Cedex, France

Email address: fatima.laytimi@univ-lille.fr

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH, TIRUPATI *Email address*: dsn@iisertirupati.ac.in

DUBLIN INSTITUTE FOR ADVANCED STUDIES, 10 BURLINGTON ROAD, DUBLIN 4, IRELAND

Email address: wnahm@stp.dias.ie