Free electron laser in the magnetically dominated regime: simulations with the ONEDFEL code

Maxim Lyutikov ¹ and Henry Freund ²

¹Department of Physics and Astronomy, Purdue University, 525 Northwestern Avenue, West Lafayette, IN, 47907-2036, USA

²Institute for Research in Electronics and Applied Physics, University of Maryland, College Park, MD 20742 and Department of Electrical and Computer Engineering, University of New Mexico, Albuquerque, NM 87131

(Received xx; revised xx; accepted xx)

Using the ONEDFEL code we perform Free Electron Laser simulations in the astrophysically important guide-field dominated regime. For wigglers' (Alfvén waves) wavelengths of tens of kilometers and beam Lorentz factor $\sim 10^3$, the resulting coherently emitted waves are in the centimeter range. Our simulations show a growth of the wave intensity over fourteen orders of magnitude, over the astrophysically relevant scale of \sim few kilometers. The signal grows from noise (unseeded). The resulting spectrum shows fine spectral substructures, reminiscent of the ones observed in Fast Radio Bursts (FRBs).

Key words:

1. Introduction: Free Electron Laser in astrophysical setting

Pulsars' radio emission mechanism(s) eluded identification for nearly half a century (e.g. Melrose 2000; Lyubarsky 2008; Eilek and Hankins 2016). Most likely, several types of coherent processes operate in different sources (e.g. magnetars versus pulsars), and in different parts of pulsar magnetospheres (see e.g. discussion in Lyutikov et al. 2016).

The problem of pulsar coherent emission generation has been brought back to the research forefront by the meteoritic developments over the last years in the field of mysterious Fast Radio Bursts (FRBs). Especially important was the detection of a radio burst from a Galactic magnetar by CHIME and STARE2 collaborations in coincidence with high energy bursts (CHIME/FRB Collaboration et al. 2020; Ridnaia et al. 2021; Bochenek et al. 2020; Mereghetti et al. 2020; Li et al. 2021). The similarity of properties of magnetars' bursts to the Fast Radio Bursts gives credence to the magnetar origin of FRBs (even though the radio powers are quite different - there is a broad distribution).

The phenomenon of Fast Radio Bursts challenges our understating of relativistic plasma coherent processes to the extreme. In this case radio waves can indeed carry an astrophysically important amount of the energy. For example, radio luminosity in FRBs can match, for a short period of time, the macroscopic Eddington luminosity and exceed total Solar luminosity by many orders of magnitude. Still, the fraction emitted in radio remains small - this relatively small fraction of total energy that pulsars and FRBs emit in radio ($\sim 10^{-5}$ is typical) is theoretically challenging: simple order-of-magnitude estimates

cannot be used. Emission production and saturation levels of instabilities depend on the kinetic details of the plasma distribution function.

Lyutikov (2021) developed a model of the generation of coherent radio emission in the Crab pulsar, magnetars and Fast Radio Bursts (FRBs) due to a variant of the free-electron laser (FEL) mechanism, operating in a weakly-turbulent, guide-field dominated plasma. This presents a new previously unexplored way (in astrophysical settings) of producing coherent emission via *parametric* instability.

A particular regime of the FEL (SASE - Self-Amplified Spontaneous Emission) microbunching is initiated by the spontaneous radiation. In the beam frame the wiggler and the electromagnetic wave have the same frequency/wave number, but propagate in the opposite direction. The addition of two counter-propagating waves creates a standing wave in the beam frame. The radiation energy density is smaller at the nodes of the standing wave: this creates a ponderomotive force that pushes the particles towards the nodes - bunches are created. These bunches are still shaken by the electromagnetic wiggler: they emit in phase, coherently.

Somewhat surprisingly, the FEL model in magnetically dominated regimes (Lyutikov 2021) is both robust to the underlying plasma parameters and succeeded in reproducing a number of subtle observed features: (i) emission frequencies depend mostly on the scale of turbulent fluctuations and the Lorentz factor of the reconnection generated beam, Eq. (2.5); it is independent of the absolute value of the underlying magnetic field. (ii) The model explained both broadband emission and the presence of emission stripes, including multiple stripes observed in the High Frequency Interpulse of the Crab pulsar. (iii) The model reproduced correlated spectrum-polarization properties: the presence of narrow emission bands in the spectrum favors linear polarization, while broadband emission can have arbitrary polarization. The model is applicable to a very broad range of neutron star parameters: the model is mostly independent of the value of the magnetic field. It is thus applicable to a broad variety of NSs, from fast spin/weak magnetic field millisecond pulsars to slow spin/super-critical magnetic field in magnetars, and from regions near the surface up to (and a bit beyond of) the light cylinder.

The guide field dominance plays a tricky role in the operation of an FEL. On the one hand it suppresses the growth rate. But what turns out to be more important in astrophysical applications is that the guide field dominance helps to maintain beam coherence. Without the guide field, particles with different energies follow different trajectories in the magnetic field of the wiggler, and quickly lose coherence even for small initial velocity spread. In contrast, in the guide-field dominated regime all particles follow, basically, the same trajectory. Hence coherence is maintained as long as the velocity spread in the beam frame is $\Delta \beta \leqslant 1$. Such tolerance to velocity spread is an unusual property of guide-field dominaed FEL.

2. FEL in guide-field dominated regime: theoretical summary

2.1. Model parameters

Let us next discuss the basic model parameters. (Unfortunately, there is some confusion in standard definitions used in literature.)

The model starts with an assumption that guiding magnetic field lines are perturbed by a packet of linearly polarized Alfvén waves of intensity B_w and frequency ω . In highly magnetized force-free plasma Alfvén waves propagating along the magnetic field are nearly luminal, $v_A \approx c$.

The first parameter is dimensionless wave intensity. We chose notation a_0 , which is

standard in the laser community and sometimes used in the FEL community as well,

$$a_0 = \frac{eB_w}{m_e c\omega} \gg 1 \tag{2.1}$$

It should be remarked that in the FEL literature parameters K and a_0 are often used interchangeably. In this paper, we follow the nomenclature used for K in Eq. (2.4) below in Jackson (1975), parag. 14.7

For guide-field dominated regime another parameter is the relative intensity

$$\delta = \frac{B_w}{B_0} \ll 1 \tag{2.2}$$

where B_0 is a guide field. We assume that the wave is relatively weak, $\delta \ll 1$.

A (reconnection-generated) beam of particles with Lorentz factor γ_b propagates along the rippled magnetic field in a direction opposite to the direction of Alfvén waves. In the frame of the beam the waves are seen with $k'_w = 2\gamma_b k_w$. In the guide-field dominated regime the cyclotron frequency associated with the guide field is much larger than the frequency of the wave in the beam frame, and the cyclotron frequency associated with the fluctuating field; hence

$$\Omega_0 \gg k_w' c, \Omega_w$$
(2.3)

where $\Omega_0 = eB_0/(m_ec)$ is the cyclotron frequency (non-relativistic) of the guide field, $\Omega_w = eB_w/(m_ec)$ is the cyclotron frequency associated with the wiggler field.

Another important parameter, defined by (Jackson 1975, parag. 14.7) is wiggler-undulator parameter

$$K = \delta \gamma_b \tag{2.4}$$

This parameter is related to the magnitude of the wiggler-induced oscillations in the beam trajectory, which is also related to the opening angle of the cone of the generated radiation. When $K\gg 1$ this oscillation is large and the pump field is sometimes referred to as a "wiggler". In the opposite regime where $K\ll 1$ the pump field sometimes referred to as an "undulator". In this paper, we will refer to the pump field as a wiggler throughout.

The K parameter (2.4) is a product of two quantities, relative amplitude $\delta \ll 1$ and Lorentz factor $\gamma_b \gg 1$, so generally its values can be either large and small. A relativistically moving electron emits in a cone with opening angle $\Delta\theta \sim 1/\gamma$. In the $K \ll 1$ regime regime that opening angle is much larger than the variation in the bulk direction of emission at different points in the trajectory, Fig. 1, left panel. The radiation detected by an observer is an almost coherent superposition of the contributions from all the oscillations in the trajectory at a frequency

$$\omega_C = \gamma_b^2(ck_w) \times \begin{cases} 4, & \text{EM wiggler} \\ 2, & \text{static wiggler} \end{cases}$$
(2.5)

We note that the axial velocity in the strong axial guide field regime is nearly constant and close to the speed of light for high energy electrons, see Fig. 7 in §4.3. Hence, the resonant frequency is also nearly constant.

In the $K \gg 1$ regime the variations in the direction of emission is much larger than the angular width of the emission at any point in the trajectory, Fig. 1, right panel. As a result, an observer located within angle $\leq \delta$ with respect to the overall guide field see periodic bursts of emission with typical frequency

$$\omega_c \approx \gamma_b^2 K \times (ck_w) \tag{2.6}$$




Figure 1: Different regimes of wave-particle interaction depending on the parameter K (2.4). Gray cones illustrate instantaneous emisison cone.

Since $K \gg 1$ in this regime, the resulting frequency is higher than that for the case where $K \ll 1$. In this paper, we work in the $K \ll 1$ regime which is the regime more typical of FELs.

Another terminology issue: here we contrast the term "Compton" with "curvature", not with "Raman" regime, which is Compton-like scattering, but on collective plasma fluctuations.

In this paper we work in the regime $K \ll 1$ (this is the usual regime of FELs). The scattered frequency is then given by (2.5); below we drop the subscript C.

2.2. Overview of main results of Lyutikov (2021)

Lyutikov (2021) developed a model of the generation of coherent radio emission in the Crab pulsar, magnetars and Fast Radio Bursts (FRBs) whereby the emission is produced by a reconnection-generated beam of particles via a variant of Free Electron Laser (FEL) mechanism, operating in a weakly-turbulent, guide-field dominated plasma. The guide-field dominance is a key new feature that distiguishes this regime from the conventional laboratory FELs: a (reconnection-generated) beam of particles with Lorentz factor γ_b propagates along the wiggled magnetic field in a direction opposite to the direction of Alfvén waves. In the frame of the beam the waves are seen with $k_{w,b} = 2\gamma_b k_w$. Guide field dominance requires $2\gamma_b k_w c \ll \Omega_0$.

The key results of Lyutikov (2021) are as follows: †

•The interaction Hamiltonian (this is particularly important for the present work, as it describes evolution of the instability and its saturation). Particle motion in the combined fields of the wiggler B_w , the EM wave E_{EM} (both with wave vector k'_w in the beam frame) and the guide-field $B_0 \gg E_w, E_{EM}$ can be described by a simple ponderomotive Hamiltonian (Eq. (39) Lyutikov 2021)

$$\mathcal{H} = \frac{\beta_z^2}{2} + \delta \left(\frac{E_{EM}}{B_0}\right) \left(\frac{\Omega_0}{k_w c}\right) \sin^2(k_w' z) \tag{2.7}$$

where β_z is axial velocity, see Fig. 2. The second term is the ponderomotive potential. The Hamiltonian formulation allows powerful analytical methods to be applied to the system (adiabatic invariant, phase space separatrix etc). This is especially important for the estimates of the non-linear saturation, one of the main goals of the present work.

† Notations used in Lyutikov (2021) are somewhat different: $a_H \to \delta$, $a_A \to a_0$, $a_{H,b} \to K = \gamma_b \delta$, $\omega_B \to \Omega_0$, $\delta \to a_0 \delta \gamma_b \Omega_0$.




Figure 2: Particle dynamics for FEL in the guide-field dominated plasma with the ponderomotive potential (2.7) (arbitrary unites chosen to illustrate the trajectory Lyutikov 2021). Left panel: 3D rendering of particles trajectories in the beam frame (a particle experiences a saddle-like trajectory). Right panel: trajectory of trapped particles in phase $\beta_z - z$ plane; velocity is normalized to the separatrix β_S , Eq. (2.9).

•The growth rate of the parametric instability is (Lyutikov 2021, Eq. (62))

$$\Gamma = \left(\left(\frac{E_w}{B_0} \right) \left(\frac{E_{EM}}{B_0} \right) \Omega_0 k_w c \right)^{1/2} \propto B_0^{-1/2}$$
(2.8)

Importantly, it is only mildly suppressed by the strong guide field.

•Saturation level. The ponderomotive potential increases linearly with EM wave intensity, while energy density of EM waves increases quadratically. The corresponding saturated velocity jitter (Lyutikov 2021, Eq. (73))

$$\beta_S = \frac{\delta}{\sqrt{2\gamma}} \frac{\omega_p}{k_w c} \tag{2.9}$$

- •The guide field dominance plays a dual role. First, it reduced the growth rate, but only mildly, $\propto B_0^{-1/2}$, Eq. (2.8). What is more important, the strong guide field helps to maintain beam coherence. Without the guide field, particles with different energies follow different trajectories, and quickly lose coherence even for small initial velocity spread. In contrast, in the guide-field dominated regime all particles follow, basically, the same trajectory. Hence coherence is maintained as long as the velocity spread in the beam frame is $\Delta\beta \leqslant 1$ (Fig. 13 of Lyutikov 2021). As a result, the model require only a mildly narrow distribution of the beam's particles, $\Delta p/p_0 \leqslant 1$ and the spectrum of turbulence $\Delta k_{w,b}/k_{w,b} \leqslant 1$
- •The model operates in a very broad range of neutron star's parameters: the model is independent of the value of the magnetic field. It is thus applicable to a broad variety of NSs, from fast spin/weak magnetic field millisecond pulsars to slow spin/super-critical magnetic field in magnetars, and from regions near the surface up to (and a bit beyond of) the light cylinder.

3. Simulations with ONEDFEL codes

3.1. The codes

In this work we performed simulations of the interaction of a single-charged relativistic beam with the wiggler using FEL code ONEDFEL (Freund and Antonsen 2024). ONEDFEL is time-dependent code that simulates the FEL interaction in one-dimension.

The radiation fields are tracked by integration of the wave equation under the slowlyvarying envelope approximation. As such, the wave equation is averaged over the fast time scale under the assumption that the wave amplitudes vary slowly over a wave period. The dynamical equations are a system of ordinary differential equations for the mode amplitudes of the field and the Lorentz force equations for the electrons which are integrated simultaneously using a 4th order Runge-Kutta algorithm. Time dependence is treated by including multiple temporal "slices" in the simulation which are separated by an integer number of wavelengths. The numerical procedure is that each slice is advanced from $z \to z + \Delta z$ separately by means of the Runge-Kutta algorithm. Time dependence is imposed as an additional operation by using the forward time derivative as an additional source term to treat the slippage of the radiation field with respect to the electrons. Slippage occurs at the rate of one wavelength per undulator period in the low gain regime and after saturation of the high gain regime but at the rate of one third of a wavelength per undulator period in the exponential gain regime (Bonifacio et al. 1990; Saldin et al. 1995; Freund and Antonsen 2024). ONEDFEL self-consistently describes slippage in each of these regimes. The simplest way to accomplish this is to use a linear interpolation algorithm to advance the field from the (i-1)th slice to the *i*th slice. Using this procedure ONEDFEL can treat electron beams and radiation fields with arbitrary temporal profiles and it is possible to simulate complex spectral properties.

Simulations, effectively, work in the lab frame, The general set-up consists of

- •guide magnetic field B_0 (as strong as numerically possible);
- •a wiggler with wave number k_w and relative amplitude $\delta = B_w/B_0 \ll 1$ is as EM wave (with adiabatic turning-on); wiggler's frequency in the beam frame is below the cyclotron frequency associated with the guide field, $\omega_w \sim \gamma_b \ll \omega_{B_0}$ (but can be comparable to the cyclotron frequency of the wiggler, ω_{B_w});
- •charged beam with "solid" (dead) neutralizing background; the corresponding Alfvén wave is relativistic, $v_A \sim c$. The beam is initially propagating along the magnetic field (no gyration)
- •pulse duration is much longer than the wiggler wavelength.

By using a pure EM wave, and not as a self-consistent Alfvén wave, eliminates complications related to setting the correct particle currents. In the highly magnetized regime $\sigma \gg 1$ Alfvén waves are nearly luminal (here $\sigma = B^2/(4\pi\rho c^2)$ is the magnetization parameter Kennel and Coroniti 1984).

Let us next comment on the applicability of the 1D regime. The post-eruption magnetic field lines are mostly radial. Development of a Coronal Mass Ejection (CME), accompanying magnetospheric FRBs, leads to opening of the magnetosphere from radii R_0 , much smaller than the light cylinder (Sharma et al. 2023), Fig. 3. After the generation of a CME the magnetosphere becomes open, with nearly radial magnetic field lines for $r \geq R_O$

4. Results

In simulating the magnetar magnetosphere environment we consider an electron beam propagating along the magnetic field in the presence of a plane-polarized electromagnetic wave. The basic parameters are shown in Table 1. We consider a mono-energetic 50 MeV beam over a bunch length/charge of 1.8 μ s/5.9 mC with a peak current of 5 kA. The beam plasma frequency corresponding to a 5 kA beam with a radius of 100 cm is about 1.6 kHz. The electromagnetic undulator is taken to have a period of 100 m and an amplitude of 0.01 kG. The excited radiation, therefore is also plane-polarized. We study





Figure 3: Post-flare opening of the magnetosphere (Sharma et al. 2023). Color is values $r \sin \theta B_{\phi}$, lines are poloidal field lines. The spin parameter is $\Omega = 0.2$, so that before the ejection of a CME the light cylinder is at $R_{LC} = 5$ (last closed magnetic field line is dashed white). Post-flare magnetosphere is open starting $R_O \ll R_{LC}$. At $r \geqslant R_O$ the magnetosphere has monopolar-like magnetic field structure. Appearance of plasmoids ("ejected plasmoid") is not important for present work.

the interaction for various values of the axial field so that the resonant wavelength will vary with the axial field.

The parameters of simulations nearly match the real physical condition (except the value of the guide field): for a beam Lorentz factor $\gamma_b \sim 100$ and a wiggler length $\lambda_w \sim 100$ meters the resonant wavelength is a few centimeters. These values are close to the real scales we expect in neutron star magnetospheres. As mentioned previously, the guide field is below that expected but the numerical simulation becomes more and more computationally challenging as the resonant linewidth becomes narrower for high guide fields. However, the wavelength becomes independent of the guide field (Fig. 4). In this particular example, over a few kilometers (also a realistic physical value) the intensity grows by fourteen orders of magnitude and reaches saturation.

| Beam Energy | $50~{\rm MeV}$ |
|------------------------------|--------------------|
| Peak Current | 5000 A |
| Bunch Duration | $1.8~\mu s$ |
| Beam Radius | 100 cm |
| Pitch Angle Spread | 0 |
| Period | 100 m |
| Amplitude | $0.01~\mathrm{kG}$ |
| Polarization | Planar |
| Axial Magnetic Field (B_0) | Variable |
| Amplitude | Variable |

Table 1: Parameters of simulations






Figure 4: Steady-state runs. Left panel: Variation in the FEL resonant wavelength with magnetic field in the Group II regime. Right panel: The saturated power and saturation distance versus wavelength for $B_0 = 0.2$ kG.

4.1. Steady-state runs

Using steady-state (i.e., time independent) ONEDFEL simulations, we have studied the variation in the resonant wavelength with increases in the magnetic field. As shown in Fig. 4, the resonant wavelength for the FEL interaction decreases from about 0.25 m for a magnetic field of 0.15 kG to 0.053 m when the magnetic field increases to 0.20 kG. We observe that the curve is approaching an asymptote as B_0 increases past 0.20 kG. This means that the resonant wavelength will remain relatively constant as the magnetic field increases above this value and we expect that the interaction properties will not change significantly for still higher field levels. This is important because simulations become increasing challenging as the field increases beyond this point. Independence of the resonant wavelength on the value of the guide magnetic field is expected, see Eq. (2.5).

The variation in the saturated power and saturation distance (when starting from noise) are shown in Fig. 4 for $B_0 = 0.20$ kG. Here we observe that the full width of the gain band extends from about 0.051 m to 0.056 m and the optimal wavelength, corresponding to the shortest saturation distance is 0.053 m (as indicated in Fig. 4) and that the decreases rapidly as the wavelength increases within this gain band.

4.2. Time-dependent simulations

Next, we ran time-dependent simulations using ONEDFEL. Simulations were conducted to determine the resonant wavelengths for different values of the axial magnetic




Figure 5: FEL power as a function of distance. We observe a growth of the EM signal over \sim fourteen orders of magnitude, from noise (no seeding)! (over a scale of \sim few kilometers - realistic neutron star setting).

field. Note that while these are 1D simulations, we need to specify the beam radius in order to determine the current density and beam plasma frequency.

Our results are plotted in Fig. 5. Importantly, the signal evolves from noise - there is no seeding. The simulations nearly match the real physical condition: for beam Lorentz factor $\gamma \sim 100$ and wiggler length $\lambda_w = 100$ meters the resonant wave length is few centimeters! These values are close to the real scales we expect in neutron star magnetospheres!). The guide field is below that of the expected, though. For high guide fields the procedure becomes more and more computationally challenging as the resonant line becomes narrower. Over a scale of few kilometers (also a realistic physical value) the intensity grows by fourteen orders of magnitude, reaching saturation.

The resulting spectral structure is most revealing, Fig. 6. We find, first, that there is a typical wavelength of the produced emission (top left panel) - this is a natural consequence of our assumptsion. We also find that the pulse has a complicated internal spectral structure, (top right panel in Fig. 6) - which is a natural property of a SASE FEL. The spectral width of the central spike is less than 1% (FWHM). This complicated internal structure resembles what is indeed seen in Fast Radio Bursts, Fig. 6, bottom right panel. FRBs display a wide variety of complex time-frequency structures (Ravi et al. 2016; Michilli et al. 2018), including strong modulations in both frequency and time (with characteristic bandwidth of $\sim 100~\rm kHz$.)

We also attempted a statistical description of the peaks, bottom left panel in Fig. 6. Since we are not aware of any theoretical prediction for the distribution of the peak, we cannot do a proper statistical analysis. Plus, naturally, any particular realization of the peaks is subject to numerous uncertainties, both physical, numerical, and statistical.




Figure 6: Top row: Time-dependent simulations of the FEL with ONEDFEL showing a line-like feature (left) with zoomed-in (right) complicated internal spectral structure of FEL in SASE regime. Red line is averaged spectrum. Bottom row, left panel: distribution of brightest peaks normalized to maximal; Bottow right: Dynamic spectrum of FRB150807 (Ravi et al. 2016) showing strong modulations in both frequency and time. (Time and frequency structures are inter-dependent in the de-dispersion procedures.)

4.3. Connection to CARM regime

The particle-wave interaction may lead to the excitation of the cyclotron motion and ensuing azimuthal bunching of emitting electrons. This will take the us into the Cyclotron Auto Resonance Maser (CARM regime). In this case, the resonant wavelength is governed by the axial velocity of the electron beam and, for fixed beam energy and undulator parameters, this will vary with the axial field. The combination of the wiggler and axial magnetic fields results in particle trajectories that exhibit a magneto-resonance in which the transverse velocity increases as the difference between the wiggler and Larmor periods decreases. The energy corresponding to the transverse velocity cannot exceed the total energy; hence, there are two distinct classes of trajectories corresponding to cases where the Larmor period is longer than the wiggler period (termed Group I) and shorter than the wiggler period (termed Group II). The variation in the axial velocity, β_{\parallel} , as the axial magnetic field increases (for given values if the wiggler period and field strength and electron energy) is illustrated in Fig. 7 (taken from Freund and Antonsen 2024) where the dashed line indicates unstable trajectories. The figure is meant to show the generic variation in the average axial velocity versus the axial field strength for a magneto-static




Figure 7: Illustration of the variation in the axial velocity with the axial magnetic field for Group I and Group II trajectories (Freund and Antonsen 2024).

wiggler, hence, it is not meant to correspond to the parameters used in the simulation. Note that the average transverse velocity $\beta_{\perp}^2 = 1 - 1/\gamma_b^2 - \beta_{\parallel}^2$. Group I trajectories are generally found in the weak axial field regime below the magneto-resonance $\Omega_0 \leq \gamma_b k_w c$, where the Larmor period is longer than the wiggler period. Group II trajectories occur in the strong axial field regime where $\Omega_0 \gg \gamma_b k_w c$. We are most concerned with Group II trajectories in which $\Omega_0 \gg \gamma_b k_w c$, which we expect to be relevant to the conditions in magnetar magnetospheres. As shown in Fig. 7, the axial velocity increases but asymptotes with increasing magnetic field in the Group II regime which implies that the FEL resonant wavelength decreases with increasing magnetic field but reaches a value which is relatively independent of the axial field strength.

We remark that there are two possible interactions of an electron beam streaming along the field lines corresponding to the FEL resonance and that of a cyclotron autoresonance maser (CARM). The ratio of the resonant frequencies of these two interaction mechanisms is given by

$$\frac{\omega_{CARM}}{\omega_{FEL}} = \frac{\Omega_0}{\gamma_b k_w c},\tag{4.1}$$

so that the FEL resonance is found at a lower frequency than that for the CARM for strong axial guide fields in the Group II regime. For the parameters of interest here, the magneto-resonance is found for an axial field of about 0.10 kG as shown in Fig. 8. We are primarily concerned here with the strong axial field regime. This is a separate, and possibly astrophysically important emission mechanism. We leave the analysis of CARM regime to a separate future investigation.

5. Conclusion

In this work we numerically study operation of Free Electron Laser in the astrophysically important guide-field dominated regime. In this regime particles mostly slide along the dominant guiding magnetic field and experience $\mathbf{E} \times \mathbf{B}$ drift in the field of the wiggler.




Figure 8: Plot showing the variation in $\Omega_0/\gamma_b k_w c$ versus the axial field showing that Group I orbits are found when $B_0 \leq 0.10$ kG and Group II orbits when $B_0 \geq 0.10$ kG.

Our parameters (energy of the beam, wavelength of the wiggler) closely match what is expected in the magnetospheres of neutron stars. The value of the guide field is, though, much smaller than expected. However, we verified that the wavelength becomes independent of the guide field, Fig. 4.

Our results are encouraging. First, we see unseeded growth over 14 orders of magnitude over the real physical scale of \sim few kilometers. It is expected that the real magnetospheres are much "noisier", with mild level of intrinsically present turbulence. The presence of such turbulent Alfvén waves will provide seeds to jump-start the operation of the FEL.

The most intriguing result is, perhaps, the fine spectral structure, Fig. 6, that qualitatively matches observations. Such fine structure is an inherent property of SASE FEL, as different narrow modes are amplified parametrically.

Limitations of our approach include:

- •One-dimensional approximation. In this case we neglect curvatures of the magnetic field lines, and corresponding particles' trajectories. We plan to address this in a separate work, using MINERVA code.
- •The saturation level will be affected by the higher guiding field. For a single quasimonochromatic wave the ponderomotive potential (2.7) increases linearly with EM wave intensity E_w , while energy density of EM waves increases quadratically $\propto E_w^2$. The balance is achieved at

$$\frac{E_{EM}}{E_w} = \frac{\omega_{p,b}^2}{k_w c \Omega_0},\tag{5.1}$$

where $\omega_{p,b}$ is beam plasma density. This is an estimate of the saturation level of the EM waves in the beam frame.

•We have not addressed the energy spread in the beam. It is expected that in the guide-field dominated regime the operation of FEL is much more tolerant to the beam spread Lyutikov (2021) since in this regime the particle trajectory is *independent of energy*. In the broad-band case the saturation will be determined \sim by a (random phase)

quasilinear diffusion. In this regime the growth rate of the EM energy of the wave due to the development of the parametric instability, Eq. (2.8), will be balanced by the particle diffusion (random phases!) in the turbulent EM field (the diffusion coefficient $\propto E_n^2$).

•Coherence of the wiggler. We assumed purely monochromatic wiggler. Spectral spread of the wiggler will tend to reduce the FEL efficiency.

We plan to address theses issues in a future publication.

This work had been supported by NASA grants 80 NSSC 17 K 0757 and NSF grants 1903332 and 1908590.

REFERENCES

- D. B. Melrose, in IAU Colloq. 177: Pulsar Astronomy 2000 and Beyond, edited by M. Kramer, N. Wex, and R. Wielebinski (2000), vol. 202 of Astronomical Society of the Pacific Conference Series, pp. 721-+.
- Y. Lyubarsky, in 40 Years of Pulsars: Millisecond Pulsars, Magnetars and More, edited by C. Bassa, Z. Wang, A. Cumming, and V. M. Kaspi (2008), vol. 983 of American Institute of Physics Conference Series, pp. 29–37.
- J. A. Eilek and T. H. Hankins, Journal of Plasma Physics 82, 635820302 (2016), 1604.02472.
- M. Lyutikov, L. Burzawa, and S. B. Popov, Mon. Not. Roy. Astron. Soc. 462, 941 (2016), 1603.02891.
- CHIME/FRB Collaboration, B. C. Andersen, K. M. Bandura, M. Bhardwaj, A. Bij, M. M. Boyce, P. J. Boyle, C. Brar, T. Cassanelli, P. Chawla, et al., *Nature* 587, 54 (2020), 2005.10324.
- A. Ridnaia, D. Svinkin, D. Frederiks, A. Bykov, S. Popov, R. Aptekar, S. Golenetskii, A. Lysenko, A. Tsvetkova, M. Ulanov, et al., Nature Astronomy 5, 372 (2021), 2005.11178.
- C. D. Bochenek, V. Ravi, K. V. Belov, G. Hallinan, J. Kocz, S. R. Kulkarni, and D. L. McKenna, Nature 587, 59 (2020), 2005.10828.
- S. Mereghetti, V. Savchenko, C. Ferrigno, D. Götz, M. Rigoselli, A. Tiengo, A. Bazzano, E. Bozzo, A. Coleiro, T. J. L. Courvoisier, et al., Astrophys. J. Lett. 898, L29 (2020), 2005.06335.
- C. K. Li, L. Lin, S. L. Xiong, M. Y. Ge, X. B. Li, T. P. Li, F. J. Lu, S. N. Zhang, Y. L. Tuo, Y. Nang, et al., Nature Astronomy (2021), 2005.11071.
- M. Lyutikov, Astrophys. J. 922, 166 (2021), 2102.07010.
- J. D. Jackson, Classical electrodynamics (1975).
- H. P. Freund and T. M. Antonsen, *Principles of Free-electron Lasers* (Springer Nature Switzerland, 4rd Edition, 2024).
- R. Bonifacio, F. Casagrande, G. Cerchioni, L. Salvo Souza, P. Pierini, and N. Piovella, Nuovo Cimento Rivista Serie 13, 1 (1990).
- E. L. Saldin, E. A. Schneidmiller, and M. V. Yurkov, Phys. Rep. 260, 187 (1995).
- C. F. Kennel and F. V. Coroniti, Astrophys. J. 283, 694 (1984).
- P. Sharma, M. V. Barkov, and M. Lyutikov, Mon. Not. Roy. Astron. Soc. 524, 6024 (2023), 2302.08848.
- V. Ravi, R. M. Shannon, M. Bailes, K. Bannister, S. Bhandari, N. D. R. Bhat, S. Burke-Spolaor, M. Caleb, C. Flynn, A. Jameson, et al., Science 354, 1249 (2016), 1611.05758.
- D. Michilli, A. Seymour, J. W. T. Hessels, L. G. Spitler, V. Gajjar, A. M. Archibald, G. C. Bower, S. Chatterjee, J. M. Cordes, K. Gourdji, et al., Nature 553, 182 (2018), 1801.03965.