

K-inflation: the legitimacy of classical treatment

Y. Ageeva^{a,b,1}, P. Petrov^{a,2}

^a *Institute for Nuclear Research of the Russian Academy of Sciences,
60th October Anniversary Prospect, 7a, 117312 Moscow, Russia*

^b *Institute for Theoretical and Mathematical Physics, M.V. Lomonosov Moscow State University,
Leninskie Gory 1, 119991 Moscow, Russia*

Abstract

In this paper we consider general theory of k-inflation and find out, that it may be in strong coupling regime. We derive accurate conditions of classical description validity using unitarity bounds for this model. Next, we choose simple toy model of k-inflation and obtain the explicit condition, which guarantees that the generation of perturbations is performed in a controllable way, i.e the exit from effective horizon occurs in the weak coupling regime. However, for the same toy model the corresponding experimental bounds on a non-linear parameter $f_{\text{NL}}^{\text{equil}}$ associated with non-Gaussianities of the curvature perturbation provide much stronger constraint than strong coupling absence condition. Nevertheless, for other known models of inflation this may not be the case. Generally, one should always check if classical description is legitimate for chosen models of inflation.

1 Introduction

Nowadays, inflation [1, 2, 3, 4] is a very successful paradigm for understanding the properties of the early Universe. Among many models of inflation, we choose k-inflation model [5, 6] for our purposes. In such models the lagrangian involves non-canonical kinetic term, which drives the cosmological evolution. Although k-inflation theories are known as free from obvious pathologies, we address to the examination of *strong coupling problem* in k-inflation. The energy scale of strong coupling is an important parameter in an effective QFT. In other words, it is the maximum energy below which the effective QFT description is valid. The strong coupling energy scale can often be qualitatively estimated by naive dimensional analysis, see for example [7, 8, 9, 10]. However, in [11, 12] it was shown, that more accurate

¹**email:** ageeva@inr.ac.ru

²**email:** petrov@inr.ac.ru

estimates using *unitarity bounds* that follow from general unitarity relations must be used in order to proceed to the correct analysis of mentioned problem.

We show - firstly by the preliminary analysis of cubic order action for scalar perturbation - that strong coupling problem indeed arises in the most general setup for k-inflation. The simple estimation of s -channel matrix element for the $2 \rightarrow 2$ scattering processes and applied unitarity bound, provide some non-trivial conditions on model functions and parameters. Next, in order to improve our estimation, we turn to the explicit calculation of all channels: s -, t -, and u -matrix elements. The structure of k-inflation model lagrangian and cubic order action for scalar perturbation in this model lead to non-trivial cancellations in the final answer for matrix element. It also turned out that t - and u -elements are suppressed compared to s -channel and the factor of suppression is the slow roll parameter ϵ , which is usually a small quantity during inflation $\epsilon \ll 1$. Using more accurate result for s -channel element, unitarity bound provides final constraint on the parameters of the model.

Our next step is to choose a simple toy model of k-inflation in order to show how to apply the unitarity constraints. It turns out that the latter gives the lower bound on the slow-roll parameter ϵ .

As we mention above, the analysis of strong coupling involves cubic order action for scalar perturbation. The same expansion is used in the calculations of *non-Gaussianity of the curvature perturbation*. Thus, it is interesting to compare the conditions on the parameters of the model that comes from observational bound on non-Gaussianity [13] and from the validity of classical description. Note that these bounds have different nature: conditions from non-Gaussianity are experimental constraints, while the strong coupling absence guarantees that our classical description is legitimate during considered times.

Again, working with a specific toy model of k-inflation, the non-Gaussianities also lead to the lower bound on the slow roll parameter ϵ . However, the condition from non-Gaussianities turns out to be much stronger than the bound from strong coupling analysis for the chosen toy model of k-inflation. We emphasize that this result is obtained for the specific model of k-inflation: the situation may differ in other models of inflation. In other words, one should check if the classical description is legitimate for the chosen theory. For example, other models of inflation may not lead to the cancellations in cubic order action between the leading terms, so this can make the conditions of strong coupling absence more restrictive.

This paper is organized as follows: a brief review of general k-inflation model is given in Sec.2. Then the analysis of the strong coupling problem is addressed in Sec.3: simple estimations of s -channel matrix element and naive condition from unitarity bound are given in subsection 3.1. This allows us to highlight the terms which provide the strongest constraints. More accurate calculations of s -, t -, u - channels matrix elements and the final condition from unitarity bound are given in subsection 3.2. The Sec. 4 dedicated to short discussion of the formulas for a non-linear parameter $f_{\text{NL}}^{\text{equil}}$ associated with non-Gaussianities of the curvature perturbation. Finally, in Sec.5 we stick to the specific simple model of k-inflation, find corresponding constraints on model parameter from strong coupling analysis as well as from

bounds for non-Gaussianities. The paper ends with the conclusion in Sec.6. There are two Appendices: Appendix A collects full expressions of couplings from cubic order action for scalar perturbation, while Appendix B contains the discussion of one interesting subtlety which arises in the calculation of s -channel matrix element from subsection 3.2.

2 Generalities

In this paper we consider a class of k-inflation models in the framework of the following action:

$$\mathcal{S} = \int d^3x dt \sqrt{-g} \mathcal{L}, \quad (1)$$

where $\sqrt{-g} \equiv \sqrt{\gamma}$ with three dimensional metric tensor with the determinant $\gamma \equiv \det^{(3)}\gamma_{ij}$ and

$$\begin{aligned} \mathcal{L} &= G_2(\phi, X) + \frac{M_{Pl}^2}{2} R, \\ X &= -\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi, \end{aligned} \quad (2)$$

and $G_2(\phi, X)$ is an arbitrary function of scalar field and its kinetic term; R is the Ricci scalar. Here we also note that we work in the Einstein frame through the whole paper. Further we will use the metric signature as $(-, +, +, +)$.

We consider the flat FLRW space-time with a scale factor $a(t)$, where t is the cosmic time, so the background equations read [14]

$$3M_{Pl}^2 H^2 + G_2 - 2XG_{2X} = 0, \quad (3a)$$

$$3M_{Pl}^2 H^2 + 2M_{Pl}^2 \dot{H} + G_2 = 0, \quad (3b)$$

where $H = \dot{a}/a$ is the Hubble parameter. As it was pointed out in [14, 15, 16], one can obtain k-inflation cosmology solving these equations for the specific form of G_2 function. The inflation occurs in the slow roll regime, i.e. $\epsilon \ll 1$ [17], where ϵ is a standard slow-roll parameter which is given by:

$$\epsilon \equiv -\frac{\dot{H}}{H^2} = \frac{XG_{2X}}{M_{Pl}^2 H^2}. \quad (4)$$

The condition $\epsilon \ll 1$ can be satisfied with some specific choice of G_2 form. For instance, one can choose G_2 as [5, 18]

$$G_2(\phi, X) = K(\phi)X + L(\phi)X^2, \quad (5)$$

where the dimensions of the functions $K(\phi)$, $L(\phi)$, and X are as follows $[K] = 2$, $[L] = 0$, and $[X] = 2$; moreover, we note that in our setup we have $[\phi] = 0$. This form of G_2 indeed admits the slow-roll inflation solution, and a necessary condition for the accelerated expansion in this case reads [18]:

$$\frac{X(K + 2XL)}{M_{Pl}^2 H^2} \ll 1.$$

To obtain the latter we also use an expression for the Hubble parameter during inflation (up to the leading order by ϵ)[18]:

$$H^2 \approx -\frac{G_2}{3M_{Pl}^2}.$$

In order to explore the stability of the model, the strong coupling problem as well as to calculate the primordial scalar non-Gaussianities we need to expand the action (1) up to the second and the third order in the perturbations. In this paper we concentrate on the scalar sector of perturbations only, since this sector usually provides the strongest conditions; for instance, see Ref. [19]. Later, when we turn to the concrete model of k-inflation, we prove that scalar sector indeed gives the strongest constraints. To this end, considering the perturbations about some background solution, we choose the following form of the metric [14]:

$$ds^2 = -[(1 + \alpha)^2 - a^{-2}e^{-2\mathcal{R}}(\partial\beta)^2]dt^2 + 2\partial_i\beta dt dx^i + a^2 e^{2\mathcal{R}} d\mathbf{x}^2,$$

where α and β are non-dynamical scalar perturbations, while \mathcal{R} is a physical one. We also note, that we work with the *unitary* gauge, i.e. $\delta\phi = 0$, which fixes the time-component of a gauge-transformation vector, see [14, 19, 20] for details. Solving the constraints for α and β , we write the unconstrained action for scalar perturbation \mathcal{R} [14, 21]

$$\mathcal{S}_{\mathcal{R}\mathcal{R}}^{(2)} = \int dt a^3 d^3x \mathcal{G}_S \left(\dot{\mathcal{R}}^2 - \frac{c_S^2}{a^2} (\vec{\nabla}\mathcal{R})^2 \right), \quad (6)$$

where

$$\mathcal{G}_S = \frac{XG_{2X} + 2X^2G_{2XX}}{H^2} = \frac{\Sigma}{H^2}, \quad (7)$$

where $\Sigma \equiv XG_{2X} + 2X^2G_{2XX}$; next

$$c_S^2 = \frac{M_{Pl}^2 H^2 \epsilon}{XG_{2X} + 2X^2G_{2XX}} = \frac{M_{Pl}^2 H^2 \epsilon}{\Sigma}. \quad (8)$$

Using the expression (8), we can rewrite formula (7) as

$$\mathcal{G}_S = M_{Pl}^2 \frac{\epsilon}{c_S^2},$$

where the ratio $\frac{\epsilon}{c_S^2}$ generally is not small.

Briefly turning to the stability analysis, we require that

$$\mathcal{G}_S > 0, \quad c_S^2 > 0, \quad (9)$$

to avoid ghost and gradient instabilities as well as we require that the speed of perturbations does not exceed the speed of light,

$$c_S^2 \leq 1.$$

The latter condition is necessary for the existence of the UV completion, see [22, 23] for the details.

3 Strong coupling regime in k-inflation model

This Section is dedicated to the computation of the unitarity bounds and corresponding constraints on the parameters of the model. We remind, that we consider pure scalar sector and we take into account only cubic order expansion of the action (1) by the scalar perturbation \mathcal{R} . This Section consists of two parts: in the first part we estimate which terms from cubic order action provide the leading contributions to unitarity bound, while in the second part we use these leading terms in order to proceed to the accurate calculation of the corresponding matrix elements and final conditions for the validity of the classical description.

3.1 Preliminary analysis

In order to show, that we indeed face the strong coupling regime in the considered class of k-inflation model (1), let us firstly carry out the simple dimensional analysis of noted problem. To this end we write the full unconstrained cubic order action for scalar perturbation \mathcal{R} [8]:

$$\begin{aligned}
\mathcal{S}_{\mathcal{R}\mathcal{R}\mathcal{R}}^{(3)} = & \int dt a^3 d^3x \left\{ \Lambda_1 \dot{\mathcal{R}}^3 + \Lambda_2 \dot{\mathcal{R}}^2 \mathcal{R} + \Lambda_3 \dot{\mathcal{R}}^2 \frac{\partial^2 \mathcal{R}}{a^2} + \Lambda_4 \dot{\mathcal{R}} \mathcal{R} \frac{\partial^2 \mathcal{R}}{a^2} \right. \\
& + \Lambda_5 \dot{\mathcal{R}} \frac{(\partial_i \mathcal{R})^2}{a^2} + \Lambda_6 \mathcal{R} \frac{(\partial_i \mathcal{R})^2}{a^2} + \Lambda_7 \dot{\mathcal{R}} \frac{(\partial^2 \mathcal{R})^2}{a^4} + \Lambda_8 \mathcal{R} \frac{(\partial^2 \mathcal{R})^2}{a^4} + \Lambda_9 \frac{\partial^2 \mathcal{R} (\partial_i \mathcal{R})^2}{a^4} \\
& + \Lambda_{10} \dot{\mathcal{R}} \frac{(\partial_i \partial_j \mathcal{R})^2}{a^4} + \Lambda_{11} \mathcal{R} \frac{(\partial_i \partial_j \mathcal{R})^2}{a^4} + \Lambda_{12} \dot{\mathcal{R}} \partial_i \mathcal{R} \partial_i \psi + \Lambda_{13} \frac{\partial^2 \mathcal{R} \partial_i \mathcal{R} \partial_i \psi}{a^2} + \Lambda_{14} \dot{\mathcal{R}} (\partial_i \partial_j \psi)^2 \\
& \left. + \Lambda_{15} \mathcal{R} (\partial_i \partial_j \psi)^2 + \Lambda_{16} \dot{\mathcal{R}} \frac{\partial_i \partial_j \mathcal{R} \partial_i \partial_j \psi}{a^2} + \Lambda_{17} \mathcal{R} \frac{\partial_i \partial_j \mathcal{R} \partial_i \partial_j \psi}{a^2} \right\}, \tag{10}
\end{aligned}$$

where $\partial^2 = \partial_i \partial_i$ and

$$\psi = \partial^{-2} \dot{\mathcal{R}}. \tag{11}$$

Actually, there are non-trivial cancellations in the models with the lagrangian (2) among $\Lambda_7, \dots, \Lambda_{11}$ terms from the action (10). Indeed, substituting the lagrangian (2), as well as expressions (7) and (8) into the general formulas for these coefficients, which are listed in Appendix A¹, we arrive to

$$\Lambda_7 = \frac{M_{Pl}^2}{2H^3}, \quad \Lambda_8 = -\frac{3M_{Pl}^2}{2H^2}, \quad \Lambda_9 = -\frac{2M_{Pl}^2}{H^2}, \quad \Lambda_{10} = -\frac{M_{Pl}^2}{2H^3}, \quad \Lambda_{11} = \frac{3M_{Pl}^2}{2H^2}, \tag{12}$$

and after quite simple integration by parts [14] this part of cubic action significantly simplifies as follows

$$\begin{aligned}
\mathcal{S}_{7,8,9,10,11}^{(3)} = & \int dt d^3x \frac{1}{a} \left\{ \Lambda_9 \partial^2 \mathcal{R} (\partial_i \mathcal{R})^2 + (\Lambda_{10} \dot{\mathcal{R}} + \Lambda_{11} \mathcal{R}) \left((\partial_i \partial_j \mathcal{R})^2 - (\partial^2 \mathcal{R})^2 \right) \right\} \\
= & \int dt d^3x \left\{ \frac{d}{dt} \left(\frac{\Lambda_{10}}{3a} \right) - \frac{\Lambda_{11}}{a} - \frac{2}{3a} \Lambda_9 \right\} \mathcal{R} \left((\partial^2 \mathcal{R})^2 - (\partial_i \partial_j \mathcal{R})^2 \right),
\end{aligned}$$

¹All other couplings Λ_1 - Λ_6 , Λ_{12} - Λ_{17} expressions are listed in Appendix A as well.

where curly brackets read

$$-\frac{1}{a} \left(H \left(\frac{\Lambda_{10}}{3} \right) - \frac{d}{dt} \left(\frac{\Lambda_{10}}{3} \right) + \Lambda_{11} + \frac{2}{3} \Lambda_9 \right) = -\frac{M_{\tilde{P}l}^2 \epsilon}{2aH^2}, \quad (13)$$

where non-zero contribution comes from the second term with the time derivative, i.e. from $\frac{d}{adt} \left(\frac{\Lambda_{10}}{3} \right)$, while the combination of other three terms with $\Lambda_9, \Lambda_{10}, \Lambda_{11}$ give zero. After that, we will denote this contribution from formula (13) as

$$\Lambda_* \equiv -\frac{M_{\tilde{P}l}^2 \epsilon}{2H^2}. \quad (14)$$

To find the conditions of the validity of the classical description, we turn to the generalized unitarity bound and use the method which was described in [12]. According to this method, we need to rewrite the quadratic action (6) as:

$$\mathcal{S}_{\mathcal{R}\mathcal{R}}^{(2)} = \frac{1}{2} \int d^3x d\eta \left[\tilde{\mathcal{R}}'^2 - c_S^2 (\vec{\nabla} \tilde{\mathcal{R}})^2 \right], \quad (15)$$

so for that one introduces new field $\tilde{\mathcal{R}} = z\mathcal{R}$ with $z = a\sqrt{2\mathcal{G}_S}$. Here also $d\eta = adt$ is a conformal time, which we will use in the calculations below; the prime means the derivative with respect to conformal time $' \equiv \frac{d}{d\eta}$. In Appendix A we write down the cubic order action (10) in terms of $\tilde{\mathcal{R}}$, formula (A.1). Having the latter we can proceed to the analysis of the potential strong coupling problem. To this end, making use of all terms in the cubic action (A.1), with Λ_i replaced by $\Lambda_{i,(j)} \propto \frac{\Lambda_i}{\mathcal{G}_S^{3/2}}$,² it is straightforward to estimate $2 \rightarrow 2$ scattering amplitude, while in the subsection 3.2 we calculate this amplitude accurately. Firstly, the dimensional analysis leads us to the schematic formula for the tree $2 \rightarrow 2$ matrix element³ [19]

$$M_{i,(j)} \sim \frac{1}{E^2} \cdot \left\{ \Lambda_{i,(j)} \cdot E^a \cdot \left(\frac{E}{c_S} \right)^b \right\}^2, \quad (16)$$

where a and b are the number of time and spatial derivatives for each term in (A.1). We consider the center-of-mass frame for our purposes. The conservation laws for the latter are as follows

$$\vec{p}_1 + \vec{p}_2 = \vec{p}_3 + \vec{p}_4 = 0, \quad (17a)$$

$$E_1 + E_2 = E_3 + E_4 = E, \quad (17b)$$

$$|\vec{p}_1| = |\vec{p}_2|, \quad |\vec{p}_3| = |\vec{p}_4|, \quad (17c)$$

where $\vec{p}_{1,2}, E_{1,2}$ and $\vec{p}_{3,4}, E_{3,4}$ are the incoming and outgoing particles momenta and energies, respectively. Next, we find

$$E_{1,2,3,4} = \frac{E}{2}, \quad (17d)$$

²New index (j) can be explained by the replacement of \mathcal{R} to $\tilde{\mathcal{R}}/z$, where z depends on conformal time, so taking the derivative with respect to conformal time provide several terms with different $\Lambda_{i,(j)}$, see Appendix A for details.

³For this kind of estimations we consider the s-channel matrix element only.

where E is the center-of-mass energy. Due to (15), the dispersion relation reads

$$E_{1,2,3,4} = c_S p_{1,2,3,4}, \quad (17e)$$

thus

$$p_{1,2,3,4} = \frac{E}{2c_S}. \quad (17f)$$

Coming back to the formula (16), the factor $\frac{1}{E^2}$ presents the s-channel propagator. Next, since the energy and momentum of the scalar are related by $\omega = c_S p$ (note, that we reserve the notation E for the center-of-mass energy), spatial momentum of an incoming or outgoing scalar is of order $p \sim E/c_S$. This clarifies the factor $\left(\frac{E}{c_S}\right)^b$, coming from the Fourier of spatial derivative. Moreover, in the case of center-of-mass frame the energies of incoming (noted as $\omega_{1,2}$) and outgoing (noted as $\omega_{3,4}$) scalars are $\omega_{1,2,3,4} \sim E$, thus we count the possible factor E^a from the Fourier of time derivative. We square the expression in curve brackets in eq. (16) since for our naive estimations we consider the easier case when both vertices are the same. The corresponding partial wave amplitude (PWA) is given by [12, 19, 24]

$$\tilde{a}_l = \frac{1}{2c_S^3} \frac{1}{32\pi} \int d(\cos x) P_l(\cos x) M, \quad (18)$$

so, omitting all numerical coefficients we can write for $l = 0$ and for each $M_{i,(j)}$

$$(\tilde{a}_0)_{i,(j)} \sim \frac{M_{i,(j)}}{c_S^3}. \quad (19)$$

It is known from Refs. [12, 19, 24] that the amplitudes at classical energy scales saturate the unitarity bound $|\tilde{a}_0| \leq 1/2$. The classical energy scale is given by Hubble parameter H , and the latter was obtained in cosmic time t , see eqs. (3). However, the amplitudes (19) are given in conformal time η , so one should substitute conformal energies E at which unitarity bound saturates as $E = E_{class} = aH$. Finally, bound $|\tilde{a}_0| \leq 1/2$ provides the set of constraints from each matrix element $M_{i,(j)}$ ⁴:

$$\frac{1}{\epsilon^{3/2}} \leq \frac{H^3 M_{Pl}^7}{\Sigma^{5/2}}, \quad (20a)$$

$$\epsilon \leq \frac{\Sigma^{3/2}}{H^4 M_{Pl}^2}, \quad (20b)$$

$$\frac{1}{\epsilon^{3/2}} \leq \frac{M_{Pl}^3 \Sigma^{3/2}}{H \lambda_1^2}, \quad (20c)$$

$$\frac{1}{\epsilon^{3/2}} \leq \frac{M_{Pl}^3}{H \Sigma^{1/2}}, \quad (20d)$$

$$\frac{1}{\epsilon^{7/2}} \leq \frac{H^3 M_{Pl}^7}{\Sigma^{5/2}}, \quad (20e)$$

$$\frac{1}{\epsilon^{7/2}} \leq \frac{M_{Pl}^3}{H \Sigma^{1/2}}, \quad (20f)$$

⁴Some of the amplitudes provide the same constraint.

where $\lambda_1 \equiv X^2 G_{2XX} + X^3 G_{2XXX}/3$ and we put all related calculations in Appendix A. Since ϵ^{-1} is an enhancement factor, the strongest conditions are (20e) and (20f), coming from $\Lambda_3 - \Lambda_6, \Lambda_*, \Lambda_{13}, \Lambda_{16}, \Lambda_{17}$ terms. We will not consider the terms with other couplings in our more accurate analysis of the amplitudes since they provide suppressed contribution.

3.2 Strong coupling absence: accurate analysis

In this subsection we go ahead to precisely calculate tree matrix elements - s -, t -, and u -channels - and find more accurate constraints on model parameters from the strong coupling problem analysis. We mention once again, that we work with the center-of-mass frame. We start with s -channel, corresponding diagram is shown in Fig. 1 (left one) and corresponding conservation laws are given by eqs. (17). Thus, the s -channel matrix element is

$$iM_s = -\frac{i(E^6 \Sigma^2 + E^4 \Sigma(-8H^2 M_{Pl}^2 + 5\Sigma)a^2 H^2 + 4E^2(-2H^2 M_{Pl}^2 + \Sigma)^2 a^4 H^4)}{128\epsilon^2 \Sigma M_{Pl}^4 a^6 H^6}. \quad (21)$$

Next, the expression for the t -channel matrix element (corresponding diagram is given on Fig. 1, central one) reads

$$iM_t = -\frac{i\left\{E^3 \epsilon(x^2 - 1) + 8Ea^2 H^2(3 + 2x - 4x^2 + \frac{2H^2 M_{Pl}^2(x-2)}{\Sigma})\right\}^2}{1024\epsilon M_{Pl}^2 a^6 H^4(x-1)},$$

where $x \equiv \cos \theta$. Changing $x \rightarrow -x$, one obtains the u -channel amplitude:

$$iM_u = \frac{i\left\{E^3 \epsilon(x^2 - 1) + 8Ea^2 H^2(3 - 2x - 4x^2 - \frac{2H^2 M_{Pl}^2(x+2)}{\Sigma})\right\}^2}{1024\epsilon M_{Pl}^2 a^6 H^4(1-x)},$$

and the diagram for this process is the right one on Fig. 1.




Figure 1: Tree level diagrams for $2 \rightarrow 2$ process: s -, t -, and u -channels, respectively.

The matrix elements for t - and u -channels can be obtained straightforwardly (though the calculations are quite cumbersome), while the s -channel element calculation involves some subtlety, which is related to the terms with ψ factor (11) in the cubic order action (10). We discuss how to deal with such a subtlety in Appendix B. Before turning

to the partial wave amplitude, we note that M_t and M_u are suppressed by ϵ as compared to M_s , so we will use $M \approx M_s$, where initially M is the full matrix element, given by the sum of all channels amplitudes. Finally, we find the PWA (18) with $l = 0$, which provides the lowest bound on the amplitude

$$\tilde{a}_0 = \frac{(-8H^4 M_{Pl}^4 + 12H^2 M_{Pl}^2 \Sigma - 5\Sigma^2)\sqrt{\Sigma}}{4096\pi\epsilon^{7/2} M_{Pl}^7 H^3}, \quad (22)$$

where we substitute classical $E = aH$ which saturates the unitarity bound

$$|\tilde{a}_0| \leq \frac{1}{2}. \quad (23)$$

In Section 5 we will choose a specific model of k-inflation and obtain the concrete constraint on model parameters. If the parameters of the model satisfy these constraint then the classical description is valid.

4 Primordial non-Gaussianities

Another conditions on the parameters of the model of k-inflation with the lagrangian (2) comes from the observational constraints on primordial scalar non-Gaussianities. The extent of non-Gaussianity can be quantified by evaluating the bispectrum of curvature perturbations \mathcal{R} , as

$$\langle \mathcal{R}(\vec{k}_1)\mathcal{R}(\vec{k}_2)\mathcal{R}(\vec{k}_3) \rangle = (2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) B_{\mathcal{R}}(k_1, k_2, k_3),$$

where $\mathcal{R}(\vec{k})$ is a Fourier component of \mathcal{R} with a wave number \vec{k} and the bispectrum is

$$B_{\mathcal{R}}(k_1, k_2, k_3) = \frac{(2\pi)^4 (\mathcal{P}_{\mathcal{R}})^2}{\prod_{i=1}^3 k_i^3} \mathcal{A}_{\mathcal{R}}(k_1, k_2, k_3),$$

which translates into a non-linear parameter f_{NL} as

$$f_{NL} = \frac{10}{3} \frac{\mathcal{A}_{\mathcal{R}}}{\sum_{i=1}^3 k_i^3},$$

where $\mathcal{P}_{\mathcal{R}}$ is a power spectrum and $\mathcal{A}_{\mathcal{R}}$ being its amplitude. The bispectrum can be of different forms depending on the relation between the $\vec{k}_1, \vec{k}_2, \vec{k}_3$. In this paper, we stick to the well-known equilateral configuration f_{NL}^{equil} with $k_1 = k_2 = k_3$. The corresponding calculations of the scalar non-Gaussianities for the k-inflation with the lagrangian (2) are given in Ref. [14]. The non-linear parameter f_{NL}^{equil} for the equilateral form is given by [14]

$$f_{NL}^{\text{equil}} = \frac{85}{324} \left(1 - \frac{1}{c_S^2}\right) - \frac{10}{81} \frac{\lambda}{\Sigma} + \frac{55}{36} \frac{\epsilon}{c_S^2} + \frac{5}{12} \frac{\eta}{c_S^2} - \frac{85}{54} \frac{s}{c_S^2}, \quad (24)$$

where the following notations were used:

$$\eta \equiv \dot{\epsilon}/(H\epsilon),$$

$$s \equiv \dot{c}_S/(Hc_S),$$

$$\lambda \equiv X^2 G_{2XX} + 2X^3 G_{2XXX}/3,$$

and $\eta \ll 1$, $s \ll 1$, while λ is generally not small. In the next Section we choose a concrete model of k-inflation and find the specific form of conditions of model parameters coming from primordial scalar non-Gaussianities.

5 Constraints on the model parameters from strong coupling problem and scalar non-Gaussianities

In this Section we choose a concrete model of k-inflation and show that some non-trivial condition on the parameter of the model indeed arises from the requirement of the classical description validity. To this end, we take the lagrangian (2) with

$$G_2(\phi, X) = -\frac{16M_{Pl}^2}{9\gamma^2\phi^2}X + \frac{16M_{Pl}^2}{9\gamma^2\phi^2M^2}X^2, \quad (25)$$

where γ is a parameter with $[\gamma] = 0$, and M is another dimensional parameter, $[M] = 1$; the setup with eq. (25) is similar to the one from Ref. [16]. For this model equations of motion (3) provide

$$H = \frac{2M}{3\sqrt{3}\gamma\phi}, \quad X = \frac{M^2}{2}, \quad (26)$$

and for the scalar field we obtain

$$\phi = Mt + c, \quad (27)$$

choosing $\phi > 0$ during $0 < t < +\infty$, without the loss of generality. Here c is a dimensionless constant. After that we find all other functions and they read:

$$\begin{aligned} \Sigma &= \frac{16M^2M_{Pl}^2}{9\gamma^2\phi^2}, \\ \mathcal{G}_S &= 12M_{Pl}^2, \quad c_S^2 = \frac{\sqrt{3}}{8}\gamma, \end{aligned} \quad (28)$$

so $\gamma > 0$ due to the stability requirement (9). Also, the slow roll parameter (4) for the model (25) is

$$\epsilon = \frac{3 \cdot \sqrt{3} \cdot \gamma}{2} \ll 1, \quad (29)$$

which provides that $\gamma \ll 1$ as well as $c_S^2 \ll 1$. This situation is similar to Ref.[19], so this justifies that scalar sector provides the strongest conditions of classical description validity.

In the considered model of k-inflation the cosmological perturbations with a slightly red-tilted power spectrum may be generated [16]. The power spectrum of \mathcal{R} perturbations is given by [14]:

$$\mathcal{P}_{\mathcal{R}} = \mathcal{A}_{\mathcal{R}} \left(\frac{k}{k_*} \right)^{n_S-1} = \frac{H^2}{8\pi^2 \mathcal{G}_S c_S^3}, \quad (30)$$

where $\mathcal{A}_{\mathcal{R}}$ is an amplitude, k_* is a pivot momentum, n_S is a spectral tilt. Surely, we require that the exit beyond effective horizon must occur in the weak coupling regime. To this end we turn to unitarity bound (23) to see whether

this condition can be satisfied at the times when the relevant modes of perturbations exit the effective horizon. The corresponding PWA (22) in the chosen model (25) reads

$$\tilde{a}_0 = -\frac{73M^2}{8748 \cdot 3^{3/4} \cdot \sqrt{2} \cdot \pi \cdot \gamma^{11/2} M_{Pl}^2 (Mt + c)^2}, \quad (31)$$

where we substitute eqs. (26), (27), and (28). To obtain a rough estimate, we find the exit time t_f at $k = k_*$, keeping in mind the smallness of $|n_S - 1|$:

$$(Mt_f + c)^2 = \frac{h_0^2}{8\pi^2 \mathcal{G}_S c_S^3 \mathcal{A}_{\mathcal{R}}}, \quad h_0 = \frac{2M}{3\sqrt{3}\gamma},$$

where we use eqs. (26), (27), (28), and (30).

At t_f eq. (31) takes quite simple form

$$\tilde{a}_0 = -\frac{73\pi \mathcal{A}_{\mathcal{R}}}{432\gamma^2},$$

and counting all numerical factors together as well as substituting observational value of $\mathcal{A}_{\mathcal{R}} = 2 \cdot 10^{-9}$ [13] we arrive to

$$\tilde{a}_0 = -\frac{1.1 \cdot 10^{-9}}{\gamma^2},$$

and, finally, the unitarity bound (23) provides

$$\gamma \geq 4.6 \cdot 10^{-5}. \quad (32)$$

Next, one can obtain an additional condition on the k-inflation model parameter based on the current experimental bounds for scalar non-Gaussianities, i.e. $f_{\text{NL}}^{\text{equil}} = -26 \pm 47$ [13]. The leading term from eq. (24) is

$$f_{\text{NL}}^{\text{equil}} \approx -\frac{85}{324c_S^2},$$

and in the model with (25) and c_S^2 given by eq. (28) we obtain

$$f_{\text{NL}}^{\text{equil}} \approx -\frac{1.2}{\gamma},$$

where the behaviour $\sim 1/\gamma$ coincides with Ref. [16]. The observational value of $f_{\text{NL}}^{\text{equil}}$ has an error larger than the value itself, i.e. $f_{\text{NL}}^{\text{equil}} = -26 \pm 47$, for 68 % CL [13]. Thus, let us choose the biggest confidence region, for example, 99.7 % CL from Ref. [13] (see Fig. 19 therein). Roughly, this confidence region provides the constraint $|f_{\text{NL}}^{\text{equil}}| < 180$, so:

$$\gamma > 0.0067. \quad (33)$$

The result is as follows: in the considered model of k-inflation the absence of strong coupling problem (32) is guaranteed in the presence of the observational bound for scalar non-Gaussianities (33).

Finally, let us find other constraints coming from the calculations of the spectral tilt n_S and r -ratio. We start with the spectral tilt, which reads [14]

$$n_S - 1 \approx -2\epsilon = -3\sqrt{3}\gamma,$$

where for the second equality we have substituted ϵ from eq. (29). For the observational values $n_S = 0.9649 \pm 0.0042$ [25] the corresponding γ satisfies

$$0.0060 < \gamma < 0.0075. \quad (34)$$

Next and final constraint comes from the observational upper bound on r -ratio [14, 25, 26]:

$$r \equiv \frac{\mathcal{P}_T}{\mathcal{P}_R} = \frac{4\mathcal{G}_S c_S^3}{\mathcal{G}_T c_T^3}, \quad (35)$$

where \mathcal{G}_T is a coupling from second order action for tensor perturbations

$$\mathcal{S}_T = \sum_{\lambda} \int dt d^3x a^3 \mathcal{G}_T \left[\dot{h}_{\lambda} - \frac{c_T^2}{a^2} (\partial h_{\lambda})^2 \right],$$

and c_T is a tensor perturbation sound speed; λ means two polarization of tensor perturbation. The corresponding experimental upper bound is [25, 26, 27]

$$r < 0.032. \quad (36)$$

For the model (25) we have

$$\mathcal{G}_T = \frac{1}{4} M_{Pl}^2,$$

and so r -ratio (35) is

$$r = 6 \cdot 3^{3/4} \cdot \sqrt{2} \cdot \gamma^{3/2},$$

where we also substitute eq. (28). Finally, applying eq. (36) we arrive to

$$\gamma < 0.014. \quad (37)$$

Thus we conclude, that the strongest conditions coming both from the observational bounds on n_S and f_{NL}^{equil} are

$$0.0067 < \gamma < 0.0075. \quad (38)$$

However, if one takes another confidence region when calculating f_{NL}^{equil} , for example 68 % CL, see Ref. [13], then the model of k-inflation with (25) will be ruled out due to the inconsistency among the conditions from n_S , r -ratio and non-Gaussianities.

6 Conclusion

This paper demonstrates that the specific model of k-inflation (2) with G_2 given by eq. (25) meets strong coupling problem. However, it is possible to find such parameters of the model that the approach of classic field theory is legitimate during considered k-inflation. We prove this statement with the proceeding to the accurate analysis of $2 \rightarrow 2$ processes and corresponding matrix elements, and then apply unitarity bound in order to obtain a non-trivial

condition of the model parameter γ . Another constraint comes from the recent observational data for scalar non-Gaussianities from Planck [13]. We find out that the latter is much stronger than the condition from strong coupling absence. Thus, we conclude that the model of k-inflation with (25) is healthy: choosing γ parameter from the permitted area, one obtains a stable theory, where the classical description is valid, and corresponding f_{NL}^{equil} is allowed by the current experimental bounds. We should also note, that for simplicity we make our calculations for the scalar sector of primordial perturbations only, however, there are mixed and tensor sectors as well. We expect that, as usual (see, for instance Ref. [19]), these sectors give even weaker constraints than the scalar one. There remains another important question: does the same result hold for each known (and phenomenologically interesting) model of inflation or maybe it is not the case? For example, models of G -inflation contain higher order partial derivatives in the cubic action for scalars, thus, it potentially can strengthen strong coupling absence condition.

Acknowledgments

The authors are thankful to Valery Rubakov for valuable discussions and very useful comments at early stages of this work back in the days. The authors are grateful to Mikhail Shaposhnikov, Eugeny Babichev, Pavel Demidov, Sergei Demidov, Sergei Mironov, Petr Satunin, Vladislav Barinov, Andrei Kataev and Victoria Volkova for fruitful discussions and careful reading of this manuscript. This work has been supported by Russian Science Foundation Grant No. 19-12-00393.

A Expressions for $\Lambda_{i,(j)}$

The purpose of this Appendix is to list the cubic order action coefficients from eq. (10). The general expressions for Λ_i , with $i = 1, \dots, 17$ are given in Ref. [8], and the formulas for the specific model (2) read:

$$\begin{aligned}
\Lambda_1[\dot{\mathcal{R}}^3] &= \frac{3\Sigma^2 - 2M_{Pl}^2 H^2 X (3G_{2X} + 4X(3G_{2XX} + XG_{2XXX}))}{6M_{Pl}^2 H^5}, \\
\Lambda_2[\dot{\mathcal{R}}^2 \mathcal{R}] &= -\frac{3\Sigma(-2M_{Pl}^2 H^2 + \Sigma)}{2M_{Pl}^2 H^4}, \\
\Lambda_3[(\dot{\mathcal{R}}^2/a^2)\partial^2 \mathcal{R}] &= -\frac{\Sigma}{H^4}, \\
\Lambda_4[(\dot{\mathcal{R}}/a^2)\mathcal{R}\partial^2 \mathcal{R}] &= \frac{-2M_{Pl}^2 H^2 + 3\Sigma}{H^3}, \\
\Lambda_5[(\dot{\mathcal{R}}/a^2)(\partial_i \mathcal{R})^2] &= \frac{-M_{Pl}^2 H^2 + 2\Sigma}{H^3}, \\
\Lambda_6[(\mathcal{R}/a^2)(\partial_i \mathcal{R})^2] &= M_{Pl}^2, & \Lambda_7[(\dot{\mathcal{R}}/a^4)(\partial^2 \mathcal{R})^2] &= \frac{M_{Pl}^2}{2H^3}, \\
\Lambda_8[(\mathcal{R}/a^4)(\partial^2 \mathcal{R})^2] &= -\frac{3M_{Pl}^2}{2H^2}, & \Lambda_9[(\partial^2 \mathcal{R}/a^4)(\partial_i \mathcal{R})^2] &= -\frac{2M_{Pl}^2}{H^2}, \\
\Lambda_{10}[(\dot{\mathcal{R}}/a^4)(\partial_i \partial_j \mathcal{R})^2] &= -\frac{M_{Pl}^2}{2H^3}, & \Lambda_{11}[(\mathcal{R}/a^4)(\partial_i \partial_j \mathcal{R})^2] &= \frac{3M_{Pl}^2}{2H^2}, \\
\Lambda_{12}[\dot{\mathcal{R}}\partial_i \mathcal{R}\partial^i \psi] &= -\frac{2\Sigma^2}{M_{Pl}^2 H^4}, & \Lambda_{13}[(\partial^2 \mathcal{R}/a^2)\partial_i \mathcal{R}\partial^i \psi] &= \frac{2\Sigma}{H^3}, \\
\Lambda_{14}[\dot{\mathcal{R}}(\partial_i \partial_j \psi)^2] &= -\frac{\Sigma^2}{2M_{Pl}^2 H^5}, & \Lambda_{15}[\mathcal{R}(\partial_i \partial_j \psi)^2] &= \frac{3\Sigma^2}{2M_{Pl}^2 H^4}, \\
\Lambda_{16}[(\dot{\mathcal{R}}/a^2)\partial_i \partial_j \mathcal{R}\partial^i \partial^j \psi] &= \frac{\Sigma}{H^4}, & \Lambda_{17}[(\mathcal{R}/a^2)\partial_i \partial_j \mathcal{R}\partial^i \partial^j \psi] &= -\frac{3\Sigma}{H^3}.
\end{aligned}$$

One can rewrite the expressions above, using eq. (8) and introducing

$$\lambda_1 \equiv X^2 G_{2XX} + X^3 G_{2XXX}/3,$$

thus, we arrive to

$$\begin{aligned}
\Lambda_1[\dot{\mathcal{R}}^3] &= \frac{\left(3M_{Pl}^2\left(\frac{H^2\epsilon}{c_S^2}\right)^2 - 2H^2[3\epsilon M_{Pl}^2 H^2 + 12\lambda_1]\right)}{6H^5}, \\
\Lambda_2[\dot{\mathcal{R}}^2\mathcal{R}] &= -\frac{3M_{Pl}^2\epsilon}{2c_S^2}\left(-2 + \frac{\epsilon}{c_S^2}\right), \\
\Lambda_3[(\dot{\mathcal{R}}^2/a^2)\partial^2\mathcal{R}] &= -\frac{M_{Pl}^2\epsilon}{c_S^2 H^2}, \\
\Lambda_4[(\dot{\mathcal{R}}/a^2)\mathcal{R}\partial^2\mathcal{R}] &= \frac{-2M_{Pl}^2 + \frac{3M_{Pl}^2\epsilon}{c_S^2}}{H}, \\
\Lambda_5[(\dot{\mathcal{R}}/a^2)(\partial_i\mathcal{R})^2] &= \frac{-M_{Pl}^2 + \frac{2M_{Pl}^2\epsilon}{c_S^2}}{H}, \\
\Lambda_6[(\mathcal{R}/a^2)(\partial_i\mathcal{R})^2] &= M_{Pl}^2, & \Lambda_7[(\dot{\mathcal{R}}/a^4)(\partial^2\mathcal{R})^2] &= \frac{M_{Pl}^2}{2H^3}, \\
\Lambda_8[(\mathcal{R}/a^4)(\partial^2\mathcal{R})^2] &= -\frac{3M_{Pl}^2}{2H^2}, & \Lambda_9[(\partial^2\mathcal{R}/a^4)(\partial_i\mathcal{R})^2] &= -\frac{2M_{Pl}^2}{H^2}, \\
\Lambda_{10}[(\dot{\mathcal{R}}/a^4)(\partial_i\partial_j\mathcal{R})^2] &= -\frac{M_{Pl}^2}{2H^3}, & \Lambda_{11}[(\mathcal{R}/a^4)(\partial_i\partial_j\mathcal{R})^2] &= \frac{3M_{Pl}^2}{2H^2}, \\
\Lambda_{12}[\dot{\mathcal{R}}\partial_i\mathcal{R}\partial^i\psi] &= -2M_{Pl}^2\left(\frac{\epsilon}{c_S^2}\right)^2, & \Lambda_{13}[(\partial^2\mathcal{R}/a^2)\partial_i\mathcal{R}\partial^i\psi] &= \frac{2M_{Pl}^2\epsilon}{c_S^2 H}, \\
\Lambda_{14}[\dot{\mathcal{R}}(\partial_i\partial_j\psi)^2] &= -\frac{M_{Pl}^2\left(\frac{\epsilon}{c_S^2}\right)^2}{2H}, & \Lambda_{15}[\mathcal{R}(\partial_i\partial_j\psi)^2] &= \frac{3M_{Pl}^2\left(\frac{\epsilon}{c_S^2}\right)^2}{2}, \\
\Lambda_{16}[(\dot{\mathcal{R}}/a^2)\partial_i\partial_j\mathcal{R}\partial^i\partial^j\psi] &= \frac{M_{Pl}^2\left(\frac{\epsilon}{c_S^2}\right)}{H^2}, & \Lambda_{17}[(\mathcal{R}/a^2)\partial_i\partial_j\mathcal{R}\partial^i\partial^j\psi] &= -\frac{3M_{Pl}^2\left(\frac{\epsilon}{c_S^2}\right)}{H}.
\end{aligned}$$

Using these expressions, as well as keeping in mind the discussion about $\Lambda_7 - \Lambda_{11}$, see eqs. (12)-(14), we substitute field $\mathcal{R} = \tilde{\mathcal{R}}/z$ into eq. (10) and obtain:

$$\begin{aligned}
S_{\mathcal{R}\mathcal{R}\mathcal{R}}^{(3)} = & \int d\eta d^3x \left\{ \Lambda_{1(1)} \tilde{\mathcal{R}}'^3 + \Lambda_{1(2)} \tilde{\mathcal{R}} \tilde{\mathcal{R}}'^2 + \Lambda_{1(3)} \tilde{\mathcal{R}}^2 \tilde{\mathcal{R}}' + \Lambda_{1(4)} \tilde{\mathcal{R}}^3 \right. \\
& + \Lambda_{2(1)} \tilde{\mathcal{R}} \tilde{\mathcal{R}}'^2 + \Lambda_{2(2)} \tilde{\mathcal{R}}^2 \tilde{\mathcal{R}}' + \Lambda_{2(3)} \tilde{\mathcal{R}}^3 \\
& + \Lambda_{3(1)} \tilde{\mathcal{R}}^2 \partial^2 \tilde{\mathcal{R}} + \Lambda_{3(2)} \tilde{\mathcal{R}} \tilde{\mathcal{R}}' \partial^2 \tilde{\mathcal{R}} + \Lambda_{3(3)} \tilde{\mathcal{R}}'^2 \partial^2 \tilde{\mathcal{R}} \\
& + \Lambda_{4(1)} \tilde{\mathcal{R}}^2 \partial^2 \tilde{\mathcal{R}} + \Lambda_{4(2)} \tilde{\mathcal{R}} \tilde{\mathcal{R}}' \partial^2 \tilde{\mathcal{R}} \\
& + \Lambda_{5(1)} \tilde{\mathcal{R}} (\partial_i \tilde{\mathcal{R}})^2 + \Lambda_{5(2)} \tilde{\mathcal{R}}' (\partial_i \tilde{\mathcal{R}})^2 \\
& + \Lambda_{6(1)} \tilde{\mathcal{R}} (\partial_i \tilde{\mathcal{R}})^2 \\
& + \Lambda_{*(1)} \tilde{\mathcal{R}} \left((\partial^2 \tilde{\mathcal{R}})^2 - (\partial_i \partial_j \tilde{\mathcal{R}})^2 \right) \\
& + \Lambda_{12(1)} \partial_i \tilde{\mathcal{R}} \tilde{\mathcal{R}}' \partial_i \partial^{-2} \tilde{\mathcal{R}}' + \Lambda_{12(2)} \tilde{\mathcal{R}}' \partial_i \tilde{\mathcal{R}} \partial_i \partial^{-2} \tilde{\mathcal{R}} + \Lambda_{12(3)} \tilde{\mathcal{R}} \partial_i \tilde{\mathcal{R}} \partial_i \partial^{-2} \tilde{\mathcal{R}}' + \Lambda_{12(4)} \tilde{\mathcal{R}} \partial_i \tilde{\mathcal{R}} \partial_i \partial^{-2} \tilde{\mathcal{R}} \\
& + \Lambda_{13(1)} \partial^2 \tilde{\mathcal{R}} \partial_i \tilde{\mathcal{R}} \partial_i \partial^{-2} \tilde{\mathcal{R}} + \Lambda_{13(2)} \partial^2 \tilde{\mathcal{R}} \partial_i \tilde{\mathcal{R}} \partial_i \tilde{\psi} \\
& + \Lambda_{14(1)} \tilde{\mathcal{R}}' (\partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}}')^2 + \Lambda_{14(2)} \tilde{\mathcal{R}}' \partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}} \partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}}' + \Lambda_{14(3)} \tilde{\mathcal{R}} (\partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}}')^2 + \Lambda_{14(4)} \tilde{\mathcal{R}}' (\partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}})^2 \\
& + \Lambda_{14(5)} \tilde{\mathcal{R}} \partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}} \partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}}' + \Lambda_{14(6)} \tilde{\mathcal{R}} (\partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}})^2 \\
& + \Lambda_{15(1)} \tilde{\mathcal{R}} (\partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}}')^2 + \Lambda_{15(2)} \tilde{\mathcal{R}} \partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}} \partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}}' + \Lambda_{15(3)} \tilde{\mathcal{R}} (\partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}})^2 \\
& + \Lambda_{16(1)} \tilde{\mathcal{R}} \partial_i \partial_j \tilde{\mathcal{R}} \partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}} + \Lambda_{16(2)} \tilde{\mathcal{R}} \partial_i \partial_j \tilde{\mathcal{R}} \partial_i \partial_j \tilde{\psi} + \Lambda_{16(3)} \tilde{\mathcal{R}}' \partial_i \partial_j \tilde{\mathcal{R}} \partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}} + \Lambda_{16(4)} \tilde{\mathcal{R}}' \partial_i \partial_j \tilde{\mathcal{R}} \partial_i \partial_j \tilde{\psi} \\
& \left. + \Lambda_{17(1)} \tilde{\mathcal{R}} \partial_i \partial_j \tilde{\mathcal{R}} \partial_i \partial_j \partial^{-2} \tilde{\mathcal{R}} + \Lambda_{17(2)} \tilde{\mathcal{R}} \partial_i \partial_j \tilde{\mathcal{R}} \partial_i \partial_j \tilde{\psi} \right\}, \tag{A.1}
\end{aligned}$$

where

$$\begin{aligned}
\Lambda_{1(1)} &= \frac{\Lambda_1}{2\sqrt{2}\mathcal{G}_S^{3/2}a^2}, & \Lambda_{1(2)} &= -\frac{3\Lambda_1 H}{2\sqrt{2}\mathcal{G}_S^{3/2}a}, & \Lambda_{1(3)} &= \frac{3\Lambda_1 H^2}{2\sqrt{2}\mathcal{G}_S^{3/2}}, & \Lambda_{1(4)} &= -\frac{\Lambda_1 a H^3}{2\sqrt{2}\mathcal{G}_S^{3/2}}, \\
\Lambda_{2(1)} &= \frac{\Lambda_2}{2\sqrt{2}\mathcal{G}_S^{3/2}a}, & \Lambda_{2(2)} &= -\frac{\Lambda_2 H}{\sqrt{2}\mathcal{G}_S^{3/2}}, & \Lambda_{2(3)} &= \frac{\Lambda_2 a H^2}{2\sqrt{2}\mathcal{G}_S^{3/2}}, \\
\Lambda_{3(1)} &= \frac{\Lambda_3 H^2}{2\sqrt{2}\mathcal{G}_S^{3/2}a}, & \Lambda_{3(2)} &= -\frac{\Lambda_3 H}{\sqrt{2}\mathcal{G}_S^{3/2}a^2}, & \Lambda_{3(3)} &= \frac{\Lambda_3}{2\sqrt{2}\mathcal{G}_S^{3/2}a^3}, \\
\Lambda_{4(1)} &= -\frac{\Lambda_4 H}{2\sqrt{2}\mathcal{G}_S^{3/2}a}, & \Lambda_{4(2)} &= \frac{\Lambda_4}{2\sqrt{2}\mathcal{G}_S^{3/2}a^2}, \\
\Lambda_{5(1)} &= -\frac{\Lambda_5 H}{2\sqrt{2}\mathcal{G}_S^{3/2}a}, & \Lambda_{5(2)} &= \frac{\Lambda_5}{2\sqrt{2}\mathcal{G}_S^{3/2}a^2}, \\
\Lambda_{6(1)} &= \frac{\Lambda_6}{2\sqrt{2}\mathcal{G}_S^{3/2}a}, \\
\Lambda_{*(1)} &= \frac{\Lambda_*}{2\sqrt{2}\mathcal{G}_S^{3/2}a^3}.
\end{aligned}$$

$$\begin{aligned}
\Lambda_{12(1)} &= \frac{\Lambda_{12}}{2\sqrt{2}\mathcal{G}_S^{3/2}a}, & \Lambda_{12(2)} &= -\frac{\Lambda_{12}H}{2\sqrt{2}\mathcal{G}_S^{3/2}}, & \Lambda_{12(3)} &= -\frac{\Lambda_{12}H}{2\sqrt{2}\mathcal{G}_S^{3/2}}, & \Lambda_{12(4)} &= \frac{\Lambda_{12}aH^2}{2\sqrt{2}\mathcal{G}_S^{3/2}}, \\
\Lambda_{13(1)} &= -\frac{\Lambda_{13}H}{2\sqrt{2}\mathcal{G}_S^{3/2}a}, & \Lambda_{13(2)} &= \frac{\Lambda_{13}}{2\sqrt{2}\mathcal{G}_S^{3/2}a^2}, \\
\Lambda_{14(1)} &= \frac{\Lambda_{14}}{2\sqrt{2}\mathcal{G}_S^{3/2}a^2}, & \Lambda_{14(2)} &= -\frac{\Lambda_{14}H}{\sqrt{2}\mathcal{G}_S^{3/2}a}, & \Lambda_{14(3)} &= -\frac{\Lambda_{14}H}{2\sqrt{2}\mathcal{G}_S^{3/2}a}, & \Lambda_{14(4)} &= \frac{\Lambda_{14}H^2}{2\sqrt{2}\mathcal{G}_S^{3/2}}, \\
\Lambda_{14(5)} &= \frac{\Lambda_{14}H^2}{\sqrt{2}\mathcal{G}_S^{3/2}}, & \Lambda_{14(6)} &= -\frac{\Lambda_{14}aH^3}{2\sqrt{2}\mathcal{G}_S^{3/2}}, \\
\Lambda_{15(1)} &= \frac{\Lambda_{15}}{2\sqrt{2}\mathcal{G}_S^{3/2}a}, & \Lambda_{15(2)} &= -\frac{\Lambda_{15}H}{\sqrt{2}\mathcal{G}_S^{3/2}}, & \Lambda_{15(3)} &= \frac{\Lambda_{15}aH^2}{2\sqrt{2}\mathcal{G}_S^{3/2}}, \\
\Lambda_{16(1)} &= \frac{\Lambda_{16}H^2}{2\sqrt{2}\mathcal{G}_S^{3/2}a}, & \Lambda_{16(2)} &= -\frac{\Lambda_{16}H}{2\sqrt{2}\mathcal{G}_S^{3/2}a^2}, & \Lambda_{16(3)} &= -\frac{\Lambda_{16}H}{2\sqrt{2}\mathcal{G}_S^{3/2}a^2}, & \Lambda_{16(4)} &= \frac{\Lambda_{16}}{2\sqrt{2}\mathcal{G}_S^{3/2}a^3}, \\
\Lambda_{17(1)} &= -\frac{\Lambda_{17}H}{2\sqrt{2}\mathcal{G}_S^{3/2}a}, & \Lambda_{17(2)} &= \frac{\Lambda_{17}}{2\sqrt{2}\mathcal{G}_S^{3/2}a^2}.
\end{aligned}$$

We use these $\Lambda_{i,(j)}$ to naively estimate the matrix elements $M_{i,(j)}$ (16). Finally, unitarity bound $|(\tilde{a}_0)_{i,(j)}| \leq \frac{1}{2}$ for each $(\tilde{a}_0)_{i,(j)}$ (19) provide the conditions on the slow roll parameter ϵ (20).

B Subtlety in the calculation of s -matrix element

In this Appendix we discuss a subtlety, which arises in calculations for s -channel matrix element (21). Formulas for t - and u -channels can be obtained in a quite straightforward way. We remind, that we consider only such vertices in the matrix element which involve only $\Lambda_3 - \Lambda_6, \Lambda_*, \Lambda_{13}, \Lambda_{16}, \Lambda_{17}$ couplings, since these terms provide the strongest naive constraints (20e) and (20f), i.e. contributions from other terms are suppressed with ϵ .

The mentioned subtlety in s -channel is related to the terms with $\Lambda_{13}, \Lambda_{16}$, and Λ_{17} couplings which involve $\psi = \partial^{-2}\tilde{\mathcal{R}}$ in the cubic action (10). Recalling that the momentum of propagator equals to zero for the s -channel (see conservation law for momenta (17a)), we consider the following terms firstly

$$\Lambda_{13(1)}\partial^2\tilde{\mathcal{R}}\partial_i\tilde{\mathcal{R}}\partial_i\partial^{-2}\tilde{\mathcal{R}} + \Lambda_{13(2)}\partial^2\tilde{\mathcal{R}}\partial_i\tilde{\mathcal{R}}\partial_i\tilde{\psi}. \quad (\text{B.1})$$

One can easily see, that we get a $\frac{1}{0^2}$ factor as ∂^{-2} acting on the propagator. To deal with such contributions, let us introduce a new parameter $\vec{\eta} \rightarrow 0$, which satisfies $\vec{p}_{1,2} \perp \vec{\eta}$, so we change the center-of-mass frame to a new frame with

$$\vec{p}'_1 + \vec{p}'_2 = \vec{\eta},$$

where

$$\vec{p}'_{1,2} \rightarrow \vec{p}_{1,2} + \frac{\vec{\eta}}{2}. \quad (\text{B.2a})$$

Next, we find

$$(p'_{1,2})^2 = p_{1,2}^2 + \frac{(\vec{\eta})^2}{4}, \quad (\vec{p}'_1, \vec{p}'_2) = (\vec{p}_1, \vec{p}_2) + \frac{(\vec{\eta})^2}{4}, \quad (\text{B.2b})$$

as well as

$$E'_1 = E_1 \sqrt{1 + \frac{c_S^2(\vec{\eta})^2}{4E_1^2}} \approx E_1 \left(1 + \frac{c_S^2(\vec{\eta})^2}{8E_1^2}\right). \quad (\text{B.2c})$$




Figure 2: Tree level diagram for $2 \rightarrow 2$ process: s -channel.

Thus, considering the left vertex from the diagram in Fig.2 we write a related expression for the vertex connected with (B.1) terms

$$\begin{aligned} & i\Lambda_{13(1)}(i\vec{p}'_1)^2 \frac{(i\vec{p}'_2, -i\vec{\eta})}{(-i\vec{\eta})^2} + i\Lambda_{13(1)}(i\vec{p}'_2)^2 \frac{(i\vec{p}'_1, -i\vec{\eta})}{(-i\vec{\eta})^2} \\ & + i\Lambda_{13(2)}(i\vec{p}'_1)^2 \frac{(i\vec{p}'_2, -i\vec{\eta})}{(-i\vec{\eta})^2} (iE') + i\Lambda_{13(2)}(i\vec{p}'_2)^2 \frac{(i\vec{p}'_1, -i\vec{\eta})}{(-i\vec{\eta})^2} (iE'). \end{aligned}$$

The same “trick” should be done for Λ_{16} and Λ_{17} terms. The final result with all contributions from Λ_{13} , Λ_{16} , and Λ_{17} for the left vertex in Fig.2 reads

$$\begin{aligned} V_1 &= \frac{1}{2} \left(i\Lambda_{13(1)}\vec{p}_1^2 - \Lambda_{13(2)}\vec{p}_1^2 E + i\Lambda_{13(1)}\vec{p}_2^2 - \Lambda_{13(2)}\vec{p}_2^2 E \right) \\ &+ \frac{|\vec{\eta}|^2}{2} \left(-i\Lambda_{16(1)} + \Lambda_{16(2)}E - \frac{1}{2}\Lambda_{16(3)}E_1 - \frac{1}{2}\Lambda_{16(3)}E_2 - \frac{i}{2}\Lambda_{16(4)}E_1E - \frac{i}{2}\Lambda_{16(4)}E_2E - i\Lambda_{17(1)} + \Lambda_{17(2)}E \right) \\ &= \frac{1}{2} \left(i\Lambda_{13(1)}\vec{p}_1^2 - \Lambda_{13(2)}\vec{p}_1^2 E + i\Lambda_{13(1)}\vec{p}_2^2 - \Lambda_{13(2)}\vec{p}_2^2 E \right), \end{aligned} \quad (\text{B.3})$$

where in the last equality we take $\vec{\eta} = 0$ and also we substitute (B.2). Here we use $(i\vec{p}'_1, -i\vec{\eta}) = (\vec{p}_1 + \frac{\vec{\eta}}{2}, \vec{\eta}) = \frac{|\vec{\eta}|^2}{2}$ as well. Using the same logic one can consider the second right vertex with outgoing particles in Fig. 2 and find out that similar contribution proportional to $\vec{\eta}$ vanishes in the same way as in (B.3).

This concludes our discussion related to a subtlety coming from ψ factor. We note once again, that t - and u -channels do not suffer from this problem, since the propagator’s momentum is not zero in these cases.

References

- [1] A. A. Starobinsky, Phys. Lett. B **91**, 99-102 (1980) doi:10.1016/0370-2693(80)90670-X
- [2] A. H. Guth, Phys. Rev. D **23**, 347-356 (1981) doi:10.1103/PhysRevD.23.347
- [3] K. Sato, Mon. Not. Roy. Astron. Soc. **195**, 467-479 (1981) NORDITA-80-29.
- [4] A. D. Linde, Phys. Lett. B **108**, 389-393 (1982) doi:10.1016/0370-2693(82)91219-9
- [5] C. Armendariz-Picon, T. Damour and V. F. Mukhanov, Phys. Lett. B **458**, 209-218 (1999) doi:10.1016/S0370-2693(99)00603-6 [arXiv:hep-th/9904075 [hep-th]].
- [6] J. Garriga and V. F. Mukhanov, Phys. Lett. B **458**, 219-225 (1999) doi:10.1016/S0370-2693(99)00602-4 [arXiv:hep-th/9904176 [hep-th]].
- [7] Y. A. Ageeva, O. A. Evseev, O. I. Melichev and V. A. Rubakov, EPJ Web Conf. **191**, 07010 (2018) doi:10.1051/epjconf/201819107010 [arXiv:1810.00465 [hep-th]].
- [8] Y. Ageeva, O. Evseev, O. Melichev and V. Rubakov, Phys. Rev. D **102**, no.2, 023519 (2020) doi:10.1103/PhysRevD.102.023519 [arXiv:2003.01202 [hep-th]].
- [9] Y. Ageeva, P. Petrov and V. Rubakov, JHEP **12**, 107 (2020) doi:10.1007/JHEP12(2020)107 [arXiv:2009.05071 [hep-th]].
- [10] Y. Ageeva, P. Petrov and V. Rubakov, Phys. Rev. D **104**, no.6, 063530 (2021) doi:10.1103/PhysRevD.104.063530 [arXiv:2104.13412 [hep-th]].
- [11] Y. Ageeva and P. Petrov, Mod. Phys. Lett. A **37**, no.26, 2250171 (2022) doi:10.1142/S0217732322501711 [arXiv:2206.10646 [gr-qc]].
- [12] Y. Ageeva and P. Petrov, [arXiv:2206.03516 [hep-th]].
- [13] Y. Akrami *et al.* [Planck], Astron. Astrophys. **641**, A9 (2020) doi:10.1051/0004-6361/201935891 [arXiv:1905.05697 [astro-ph.CO]].
- [14] A. De Felice and S. Tsujikawa, JCAP **04**, 029 (2011) doi:10.1088/1475-7516/2011/04/029 [arXiv:1103.1172 [astro-ph.CO]].
- [15] D. Seery and J. E. Lidsey, JCAP **06**, 003 (2005) doi:10.1088/1475-7516/2005/06/003 [arXiv:astro-ph/0503692 [astro-ph]].

- [16] X. Chen, M. x. Huang, S. Kachru and G. Shiu, JCAP **01**, 002 (2007) doi:10.1088/1475-7516/2007/01/002 [arXiv:hep-th/0605045 [hep-th]].
- [17] S. Weinberg, “Cosmology”, Oxford, UK: Oxford Univ. Pr. (2008) 593 p.
- [18] Z. P. Peng, J. N. Yu, X. M. Zhang and J. Y. Zhu, Phys. Rev. D **94**, no.10, 103531 (2016) doi:10.1103/PhysRevD.94.103531 [arXiv:1611.02789 [gr-qc]].
- [19] Y. Ageeva, P. Petrov and V. Rubakov, JHEP **01**, 026 (2023) doi:10.1007/JHEP01(2023)026 [arXiv:2207.04071 [hep-th]].
- [20] T. Kobayashi, Rept. Prog. Phys. **82**, no.8, 086901 (2019) doi:10.1088/1361-6633/ab2429 [arXiv:1901.07183 [gr-qc]].
- [21] T. Kobayashi, M. Yamaguchi and J. Yokoyama, JCAP **07**, 017 (2015) doi:10.1088/1475-7516/2015/07/017 [arXiv:1504.05710 [hep-th]].
- [22] A. Adams, N. Arkani-Hamed, S. Dubovsky, A. Nicolis and R. Rattazzi, JHEP **10**, 014 (2006) doi:10.1088/1126-6708/2006/10/014 [arXiv:hep-th/0602178 [hep-th]].
- [23] C. de Rham, M. Fasiello and A. J. Tolley, Phys. Lett. B **733**, 46-51 (2014) doi:10.1016/j.physletb.2014.03.061 [arXiv:1308.2702 [hep-th]].
- [24] C. Grojean, Phys. Usp. **50**, 1-35 (2007) doi:10.1070/PU2007v050n01ABEH006157
- [25] N. Aghanim *et al.* [Planck], Astron. Astrophys. **641**, A6 (2020) [erratum: Astron. Astrophys. **652**, C4 (2021)] doi:10.1051/0004-6361/201833910 [arXiv:1807.06209 [astro-ph.CO]].
- [26] P. A. R. Ade *et al.* [BICEP and Keck], Phys. Rev. Lett. **127**, no.15, 151301 (2021) doi:10.1103/PhysRevLett.127.151301 [arXiv:2110.00483 [astro-ph.CO]].
- [27] M. Tristram, A. J. Banday, K. M. Górski, R. Keskitalo, C. R. Lawrence, K. J. Andersen, R. B. Barreiro, J. Borrill, L. P. L. Colombo and H. K. Eriksen, *et al.* Phys. Rev. D **105**, no.8, 083524 (2022) doi:10.1103/PhysRevD.105.083524 [arXiv:2112.07961 [astro-ph.CO]].