

A STUDY ON VARIOUS GENERALIZATIONS OF GENERALIZED CENTERS (GC) IN BANACH SPACES

SYAMANTAK DAS AND TANMOY PAUL

ABSTRACT. In [*Generalized centers of finite sets in Banach spaces*, Acta Math. Univ. Comenian. (N.S.) **66**(1) (1997), 83–115], Veselý developed the idea of generalized centers for finite sets in Banach spaces. In this work, we explore the concept of *restricted \mathcal{F} -center property* for a triplet $(X, Y, \mathcal{F}(X))$, where Y is a subspace of a Banach space X and $\mathcal{F}(X)$ is the family of finite subsets of X . In addition, we generalize the analysis to include all closed, bounded subsets of X . Similar to how Lindenstrauss characterized *n.2.I.P.*, we characterize *n.X.I.P.*. So, it is possible to figure out that Y has *n.X.I.P.* in X for all natural numbers n if and only if $\text{rad}_Y(F) = \text{rad}_X(F)$ for all finite subsets F of Y . It then turns out that, for all continuous, monotone functions f , the f -radii viz. $\text{rad}_Y^f(F), \text{rad}_X^f(F)$ are same whenever the generalized radii viz. $\text{rad}_Y(F), \text{rad}_X(F)$ are also same, for all finite subsets F of Y . We establish a variety of characterizations of central subspaces of Banach spaces. With reference to an appropriate subfamily of closed and bounded subsets, it appears that a number of function spaces and subspaces exhibit the restricted weighted Chebyshev center property.

1. INTRODUCTION

1.1. **Objectives:** In this study, we focus on the minimization problem $\inf_{y \in Y} f(\|z_1 - y\|, \dots, \|z_n - y\|)$ within the framework of Banach spaces X and its closed subspaces Y . The problem involves finding the minimum value of a given function $f : \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}_{\geq 0}$, which is continuous, monotone (point wise), coercive, corresponding to a finite subset $\{z_1, z_2, \dots, z_n\}$ of X . Additionally, we explore variations of this problem in Banach spaces. The outcomes in [16] make it clear that this phenomenon is connected to the intersection properties of balls in Banach spaces. We extend the concept

2000 *Mathematics Subject Classification.* Primary 41A28, 41A65 Secondary 46B20
41A50 May 7, 2024 .

Key words and phrases. Chebyshev center, generalized center, weighted Chebyshev center, central subspace, *n.X.I.P.*

of f -centers, as introduced in the work by Veselý in [16], to encompass all closed and bounded subsets of the space X .

In [16], the author observed that the generalized radius for a finite set in a Banach space remains same when it is considered in the bidual of the space. In this paper, we investigate the following problem.

Problem 1.1. *Let Y be a subspace of X . For every finite subset F of Y , under what necessary and sufficient conditions do the generalized radii of F remain the same when it is viewed as a subset of both X and Y ?*

Additionally, we aim to study various properties viz. \mathcal{F} -rcp, wrcp, \mathcal{A} - C -subspace, (GC) , \mathcal{A} -IP (see Definition 1.3, 1.6) in various function spaces and their subspaces.

1.2. Prerequisites: We list some standard notations used in this study: X denotes a Banach space, and by subspace, we indicate a closed linear subspace. B_X and S_X represent the closed unit ball and the unit sphere of X , respectively. For $x \in X$ and $r > 0$, $B_X(x, r)$ denotes the closed ball in X , centered at x and radius r . When there is no chance of confusion, we simply write $B(x, r)$ for the closed ball in X . We consider an element $x \in X$ to be canonically embedded in X^{**} . $\mathcal{F}(X)$, $\mathcal{C}(X)$, $\mathcal{K}(X)$, $\mathcal{WC}(X)$, $\mathcal{B}(X)$, and $\mathcal{P}(X)$ represent the set of all nonempty finite, closed and convex, compact, weakly compact, closed and bounded, and power set of X , respectively. It is assumed that the real line corresponds to the scalar field for the spaces. For a nonempty subset B of X , we consider an ordered tuple $(t_b)_{b \in B}$, indexed by the set B itself. We allow the repetitions, and by the well-ordering principle, such an ordering exists always. We consider coordinate-wise ordering on a subclass of $\Pi_{b \in B} \mathbb{R}_{\geq 0}$, viz. $\ell_\infty(B) = \{\varphi : B \rightarrow \mathbb{R}_{\geq 0} : \sup_{b \in B} \varphi(b) < \infty\}$, defined as: for $\varphi_1, \varphi_2 \in \ell_\infty(B)$, $\varphi_1 \leq \varphi_2$ if and only if $\varphi_1(b) \leq \varphi_2(b)$ for all $b \in B$. For a subset B of X , we consider $\ell_\infty(B)$ to be endowed with the supremum norm.

Definition 1.2. For a subset B of X , and a function $f : \ell_\infty(B) \rightarrow \mathbb{R}_{\geq 0}$, we call:

- (a) f is *monotone* if for $\varphi_1, \varphi_2 \in \ell_\infty(B)$, $\varphi_1 \leq \varphi_2$ implies $f(\varphi_1) \leq f(\varphi_2)$.
- (b) f is *strictly monotone* if for $\varphi_1, \varphi_2 \in \ell_\infty(B)$, $\varphi_1 \leq \varphi_2$ and $\varphi_1 \neq \varphi_2$ imply $f(\varphi_1) < f(\varphi_2)$.

(c) f is *coercive* if $f(\varphi) \rightarrow \infty$ as $\|\varphi\|_\infty \rightarrow \infty$.

- Definition 1.3.** (a) Let Y be a subspace of X , $F \in \mathcal{B}(X)$, and a function $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$. We define $r_f(x, F) = f(\|x - a\|_{a \in F})$ and $\text{rad}_Y^f(F) = \inf_{y \in Y} r_f(y, F)$ for $x \in X$. The collection of all points in Y where the infimum that defines $\text{rad}_Y^f(F)$ is achieving is referred to as the *restricted f -centers of F in Y* and is denoted by $\text{Cent}_Y^f(F)$. That is, $\text{Cent}_Y^f(F) = \{y \in Y : r_f(y, F) = \text{rad}_Y^f(F)\}$. When $Y = X$, then the restricted f -centers are called the f -centers of F in X . If the set of f -centers (restricted f -centers) is nonempty, we say that $X(Y)$ admits f -centers (restricted f -centers) for F .
- (b) When f is of the form $f(a) = \sup_{t \in F} \rho(t)a(t)$, where $\rho = (\rho(t))_{t \in F} \in \ell_\infty(F)$, then $r_f(x, F)$, $\text{rad}_Y^f(F)$, and $\text{Cent}_Y^f(F)$ are rewritten as $r_\rho(x, F)$, $\text{rad}_Y^\rho(F)$, and $\text{Cent}_Y^\rho(F)$, respectively. In this case, we refer to the f -centers as the *restricted weighted Chebyshev centers*. In this case, we call Y admits restricted weighted Chebyshev center for F for weights $(\rho(t))_{t \in F}$. When $\rho(t) = 1$ for all t , we denote the previous quantities by $r(x, F)$, $\text{rad}_Y(F)$ and $\text{Cent}_Y(F)$ respectively.
- (c) For a subspace Y of X , $\mathfrak{F} \subseteq \mathcal{B}(X)$ and a family of functions \mathcal{F} consisting of $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$ for all $F \in \mathfrak{F}$, the triplet (X, Y, \mathfrak{F}) is said to have the *restricted \mathcal{F} -center property* (\mathcal{F} -**rcp** in short) if for all $F \in \mathfrak{F}$ and $f \in \mathcal{F}$, we have $\text{Cent}_Y^f(F) \neq \emptyset$.
- (d) For a subspace Y of X and $\mathfrak{F} \subseteq \mathcal{B}(X)$, the triplet (X, Y, \mathfrak{F}) is said to have *restricted weighted Chebyshev center property* (**wrcp** in short) if for all $F \in \mathfrak{F}$ and bounded weights $\rho : F \rightarrow \mathbb{R}_{\geq 0}$, $\text{Cent}_Y^\rho(F) \neq \emptyset$.

We state the following result based on [16].

Theorem 1.4. *For a Banach space X and any n -tuple $(a_1, \dots, a_n) \in \mathcal{F}(X)$, the following are equivalent:*

- (a) *If $r_1, \dots, r_n > 0$ then $\bigcap_{i=1}^n B_{X^{**}}(a_i, r_i) \neq \emptyset$ implies $\bigcap_{i=1}^n B_X(a_i, r_i) \neq \emptyset$.*
- (b) *X admits weighted Chebyshev centers for (a_1, \dots, a_n) for all weights $\rho_1, \dots, \rho_n > 0$.*
- (c) *X admits f -centers for (a_1, \dots, a_n) for each continuous monotone coercive function f on $\mathbb{R}_{\geq 0}^n$.*

A Banach space X belongs to the class (GC) if for every ordered n -tuple $(a_1, \dots, a_n) \in \mathcal{F}(X)$ it satisfies any one of the equivalent conditions stated in Theorem 1.4. It is denoted by $X \in (GC)$.

Bandyopadhyay and Rao in [3] introduced the notion of *central subspace*, which is a generalization of class (GC) given in [16].

Definition 1.5. A subspace Y of X is said to be a *central subspace* of X if for any finite family of balls in X with centers in Y which intersect in X , also intersect in Y .

Bandyopadhyay and Dutta in [2] further generalized the notion of central subspace to $\mathcal{A} - C$ -subspace, for a subfamily \mathcal{A} of $\mathcal{P}(X)$.

Definition 1.6. [2] Let Y be a subspace of Banach space X and \mathcal{A} be a family of subsets of Y .

- (a) Y is said to be an *almost $\mathcal{A} - C$ -subspace* of X if for all $x \in X$, $A \in \mathcal{A}$ and $\varepsilon > 0$, there exists $y \in Y$ such that $\|y - a\| \leq \|x - a\| + \varepsilon$ for all $a \in A$.
- (b) Y is said to be a *$\mathcal{A} - C$ -subspace* of X if we can take $\varepsilon = 0$ in (a).
- (c) If \mathcal{A} is a family of subsets of X , then it is referred to that X has (almost) \mathcal{A} -IP, if X is an (almost) $\mathcal{A} - C$ -subspace of X^{**} .

Note, $X \in (GC)$ if it has $\mathcal{F}(X)$ -IP. Moreover, $\mathcal{F}(Y) - C$ -subspaces are called central subspaces in [3].

Definition 1.7. (a) [9] A subspace Y of X is said to have *n -X.I.P.* in X if for any n closed balls $\{B_X(a_i, r_i)\}_{i=1}^n$ having centers in Y and $\bigcap_{i=1}^n B_X(a_i, r_i) \neq \emptyset$, then $\bigcap_{i=1}^n B_Y(a_i, r_i + \varepsilon) \neq \emptyset$ for all $\varepsilon > 0$.

- (b) [9] A subspace Y of X is said to be an *ideal* of X if there exists a projection $P : X^* \rightarrow X^*$ such that $\|P\| = 1$ and $\ker(P) = Y^\perp$.

Remark 1.8. (a) *The subspaces Y which have n -X.I.P. for all n are precisely those which are almost $\mathcal{F}(Y) - C$ -subspaces of X . In general, n -X.I.P. does not imply k -X.I.P. for $k > n$, although under certain assumptions, n -X.I.P. for all n is equivalent to 3-X.I.P.*

- (b) *A subspace Y of X which is known to be an ideal in X also satisfies n -X.I.P. (see [9, Proposition 3.2]).*
- (c) *A subspace that is a range of a norm-1 projection is clearly a central subspace and also an ideal.*

For a subspace Y of X , we introduce the following notions here.

Definition 1.9. A subspace Y of X is said to have *restricted n .X.I.P. in X* (be a *restricted central subspace of X*) if for any n closed balls $\{B_X(a_i, r)\}_{i=1}^n$ having centers in Y and $\bigcap_{i=1}^n B_X(a_i, r) \neq \emptyset$, then $\bigcap_{i=1}^n B_Y(a_i, r + \varepsilon) \neq \emptyset$ for all $\varepsilon > 0$ ($\bigcap_{i=1}^n B_Y(a_i, r) \neq \emptyset$).

For a Banach space X , and a finite measure space (Ω, Σ, μ) and $1 \leq p < \infty$, $L_p(\Sigma, X)$ represents the space of all *p -Bochner integrable functions*, which are precisely $f : \Omega \rightarrow X$ strongly measurable and $\int_{\Omega} \|f(t)\|^p d\mu(t) < \infty$. $\|f\|_p := \left(\int_{\Omega} \|f(t)\|^p d\mu(t)\right)^{1/p}$ defines a norm on $L_p(\Sigma, X)$, which makes $L_p(\Sigma, X)$ a Banach space. $L_{\infty}(\Sigma, X)$ represents the set of all *essentially bounded functions* $f : \Omega \rightarrow X$, which are strongly measurable. [5, Ch. 2] is a standard reference for these spaces and all the properties used in this article.

Let us recall from [5, Ch. 5] that for a sub σ -algebra $\Sigma' \subseteq \Sigma$, by considering $L_p(\Sigma', X)$ as a subspace of $L_p(\Sigma, X)$, $1 \leq p < \infty$, the *conditional expectation operator* is a mapping $E : L_p(\Sigma, X) \rightarrow L_p(\Sigma', X)$ such that $E(f) = g$, where $\int_B f d\mu = \int_B g d\mu$ for all $B \in \Sigma'$ indicates a linear projection of norm-1.

Definition 1.10. [6]

- (a) A bounded linear projection $P : X \rightarrow X$ is said to be an *L -projection* if $\|x\| = \|Px\| + \|x - Px\|$ for all $x \in X$.
- (b) A Banach space X is said to be *L -embedded* if X , under its canonical image in X^{**} , is the range of an L -projection on X^{**} .
- (c) A closed subspace $J \subseteq X$ is an *M -ideal* in X if J^{\perp} is the range of an L -projection on X^* ([6]).

Reference [6, Ch. 4] provides examples and other properties of L -embedded spaces. We call a subspace Y of X is *1-complemented* if Y is a range of a norm-1 projection from X . If Y is a 1-complemented subspace of an L -embedded space X , then Y is also L -embedded (see [6, Theorem IV.1.5.]). For a sub σ -algebra $\Sigma' \subseteq \Sigma$, due to the conditional expectation $E : L_1(\Sigma, X) \rightarrow L_1(\Sigma', X)$, $L_1(\Sigma', X)$ is L -embedded if $L_1(\Sigma, X)$ is so.

A Banach space X is said to be a *Lindenstrauss space* (or *L_1 -predual*) if $X^* \cong L_1(\mu)$ for some measure μ . In [10], Lindenstrauss characterizes these spaces as Banach spaces, where any collection of pairwise intersecting closed balls whose centers form a compact set has a nonempty intersection.

An ideal of a Lindenstrauss space is also a Lindenstrauss space and, hence, a central subspace (see [8, Lemma 10, Theorem 15]).

In [16], Veselý derived that for spaces X with the Radon-Nikodým-property (RNP in short) and 1-complemented in its bidual, $L_p(\mu, X) \in (GC)$, for $1 \leq p < \infty$. Additionally, if X is a dual space that is strictly convex and has (w^*K) , then $C_b(T, X) \in (GC)$ for a Hausdorff space T .

We refer to the articles [2, 16] for various examples of the spaces that are discussed in this paper.

1.3. Observations: In this subsection, we give an overview of our observations. Suppose that, Y has $n.X.I.P.$ in X and $F \in \mathcal{F}(Y)$, where $\text{card}(F) = n$ and X admits weighted Chebyshev centers for F . In section 2, it is demonstrated that Y admits restricted f -centers for F , where $\text{card}(F) = n$, for all continuous, monotone functions $f : \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}_{\geq 0}$, if and only if for any collection of n closed balls with centers in F having a nonempty intersection in X also intersect in Y . We obtain a similar characterization to that [16, Theorem 2.7] for the family of closed and bounded subsets of Y .

Problem 1.1 is answered in Theorem 2.6, which extends a characterization to the notion $n.X.I.P.$ in Banach spaces. Consequently, we obtain various characterizations for central subspaces in Banach spaces, as stated in Theorem 2.7 and Theorem 2.8. It is easy to observe that if Y is a subspace of X and if there exists $P : X \rightarrow Y$ onto, where $P(\lambda x + y) = \lambda P(x) + y$, for $x \in X, y \in Y$ and scalar λ , $\|P(x)\| \leq \|x\|$ for all x , a *quasi-linear projection* $P : X \rightarrow X$, then for any $F \in \mathcal{B}(Y)$, $\text{rad}_Y(F) = \text{rad}_X(F)$. Veselý's example in [16, Pg.9] ensures there does not exist any quasi-linear projection $P : \ell_\infty \rightarrow c_0$. We do not know the solution to the question posed in Problem 1.1 for closed bounded subsets, with the exception of some trivial circumstances, such as when Y is 1-complemented in X or, more generally, a range of a quasi-linear projection. The requirement for Y being a central subspace can be more simply expressed when $Y \in (GC)$ is observed.

Let us give an example of a space X such that $L_1(\mu, X) \notin (GC)$. Let $f \in \ell_1$ be such that support of f is infinite and $2\|f\|_\infty > \|f\|_1$. Now identify f as a linear functional on c_0 and suppose that $X = \ker(f)$. Thus, $X \notin (GC)$ (see [15]). Since X is 1-complemented in $L_1(\mu, X)$, $L_1(\mu, X) \notin (GC)$.

In section 3, we investigate the restricted \mathcal{F} -center property for various triplets (X, Y, \mathfrak{F}) , where Y represents a subspace of X and $\mathfrak{F} \subseteq \mathcal{B}(X)$. It is

evident that when a given subspace $Y \subseteq X$, $(X, Y, \mathcal{F}(X))$ possesses **wrcp**, then $Y \in (GC)$. On the other hand, from Theorem 2.3 one can conclude in the above instances that Y is a central subspace of X whenever X admits f -centers for $F \in \mathcal{F}(Y)$, for all continuous, monotone functions $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$.

Methods for dealing with the aforementioned ideas, including (GC) , central subspace, $n.X.I.P.$, $\mathcal{A} - C$ -subspace are adapted from [2, 3, 14, 15, 16]. In section 3, we use certain measure-theoretic tools from [5] to derive our observations.

2. $\mathcal{A} - C$ -SUBSPACES OF BANACH SPACES

The following is obtained by employing justifications similar to those given in [16, Theorem 2.6]. For an $F \in \mathcal{F}(X)$, by $card(F)$, we mean the cardinality of the ordered tuple.

Theorem 2.1. *Let Y be a subspace of X and $n \in \mathbb{N}$. Suppose Y has $n.X.I.P.$ in X , and $f : \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}_{\geq 0}$ be a continuous, monotone function. Then, for all $F \in \mathcal{F}(Y)$ with $card(F) = n$, we have $rad_Y^f(F) = rad_X^f(F)$.*

Proof. Suppose that $F = (z_1, \dots, z_n)$ be an ordered n -tuple. Let $\mathbf{t}_0 = (\|x - z_1\|, \dots, \|x - z_n\|)$. Choose $\varepsilon > 0$ and let $\delta > 0$ be such that $|f(\mathbf{t}) - f(\mathbf{t}_0)| \leq \varepsilon$ whenever $d(\mathbf{t}, \mathbf{t}_0) \leq \delta$. Without loss of generality, we may assume $\delta < \varepsilon$.

Let $x \in X$ be such that, $f(\|x - z_1\|, \dots, \|x - z_n\|) < rad_X^f(F) + \delta$. Since Y has $n.X.I.P.$ in X , there exists $y \in \cap_{i=1}^n B_Y(z_i, \|x - z_i\| + \delta)$.

$$\begin{aligned} \text{Now } rad_Y^f(F) &\leq f(\|y - z_1\|, \dots, \|y - z_n\|) \\ &\leq f(\|x - z_1\| + \delta, \dots, \|x - z_n\| + \delta) \\ &\leq f(\|x - z_1\|, \dots, \|x - z_n\|) + \varepsilon \\ &< rad_X^f(F) + \delta + \varepsilon < rad_X^f(F) + 2\varepsilon. \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, the result follows. \square

The proof of the following theorem follows in the same manner as the proof of [16, Theorem 2.7].

Theorem 2.2. *Let \mathcal{A} be one of the families \mathcal{F}, \mathcal{K} and \mathcal{B} . Assume that $F \in \mathcal{A}(Y)$, where $F = (y_i)_{i \in F}$. Suppose that for each continuous, monotone function $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$, $rad_Y^f(F) = rad_X^f(F)$ and X admits f -centers for F . Then the following are equivalent.*

- (a) If for $i \in F$, $r_i \in \mathbb{R}_{>0}$, $\cap_{i \in F} B_X(y_i, r_i) \neq \emptyset$, then $\cap_{i \in F} B_Y(y_i, r_i) \neq \emptyset$.
- (b) Y admits restricted weighted Chebyshev centers for F for all weights $\rho : F \rightarrow \mathbb{R}_{>0}$.
- (c) Y admits restricted f -centers for F for all continuous, monotone functions $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$.

Proof. (a) \Rightarrow (c). Let $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$ be a continuous, monotone function and $x \in \text{Cent}_X^f(F)$. As $x \in \cap_{i \in F} B_X(y_i, \|x - y_i\|)$, there exists $y \in \cap_{i \in F} B_Y(y_i, \|x - y_i\|)$. Since $\text{rad}_Y^f(F) = \text{rad}_X^f(F)$, it can be clearly understood that $y \in \text{Cent}_Y^f(F)$.

(c) \Rightarrow (b). This is obvious.

(b) \Rightarrow (a). Let $x \in \cap_{i \in F} B_X(y_i, r_i)$. Define $\rho_i = \frac{1}{r_i}$ for all $i \in F$. Accordingly, $r_\rho(x, F) \leq 1$. Based on our assumption, we obtain $\text{rad}_Y^\rho(F) = \text{rad}_X^\rho(F) \leq r_\rho(x, F) \leq 1$. Moreover, there exists $y \in Y$ such that $y \in \text{Cent}_Y^\rho(F)$. Thus $\frac{1}{r_i} \|y - y_i\| \leq 1$ for all $i \in F$. Thus $y \in \cap_{i \in F} B_Y(y_i, r_i)$. \square

By varying $F \in \mathcal{A}(Y)$ in Theorem 2.2 we obtain the following.

Theorem 2.3. *Let \mathcal{A} be one of the families \mathcal{F}, \mathcal{K} and \mathcal{B} . Let Y be a subspace of X and for all $F \in \mathcal{A}(Y)$, $\text{rad}_Y^f(F) = \text{rad}_X^f(F)$ and X admit f -centers for F , for all continuous monotone $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$. Then the following are equivalent.*

- (a) Y is a \mathcal{A} - C -subspace of X .
- (b) The triplet $(X, Y, \mathcal{A}(Y))$ has **wrcp**.
- (c) $\text{Cent}_Y^f(F) = \text{Cent}_X^f(F) \cap Y \neq \emptyset$, for all $F \in \mathcal{A}(Y)$ and each continuous, monotone $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$.

We now focus on the family $\mathcal{F}(Y)$ for a subspace Y of X . Theorem 2.3 reduces to the characterizations of central subspaces by taking $\mathcal{A} = \mathcal{F}$. Some other characterizations of central subspaces, (GC) are also obtained in the subsequent part of this section.

Lemma 2.4. *Let Y be a subspace of X . If Y has restricted n -X.I.P. in X , then Y has n -X.I.P. in X .*

Proof. Let $\{B_X(y_i, r_i)\}_{i=1}^n$ be a collection of n closed balls in X , where $y_1, \dots, y_n \in Y$ and $r_1, \dots, r_n > 0$ such that $\cap_{i=1}^n B_X(y_i, r_i) \neq \emptyset$.

Suppose there exists an $\varepsilon > 0$ for which $\cap_{i=1}^n B_Y(y_i, r_i + \varepsilon) = \emptyset$. Let $r \in \mathbb{R}$ be such that $r - \frac{\varepsilon}{2} > r_i$ for all $i = 1, \dots, n$. We now construct n closed balls

$\{B_X(w_i, r - \frac{\varepsilon}{2})\}_{i=1}^n$ with $w_1 \cdots, w_n \in Y$ such that $B_X(w_i, r - \frac{\varepsilon}{2}) \supseteq B_X(y_i, r_i)$ and $\bigcap_{i=1}^n B_Y(w_i, r) = \emptyset$, which contradict our assumption. We now choose w_i inductively.

We define $L_1 = \bigcap_{i=2}^n B_Y(y_i, r_i + \varepsilon)$ and $L_2 = B_Y(y_1, r_1 + \varepsilon)$.

CASE 1: Suppose $L_1 = \emptyset$. Then we chose $w_1 = y_1$ and obtain $B_Y(w_1, r) \cap (\bigcap_{i=2}^n B_Y(y_i, r_i + \varepsilon)) = \emptyset$ and proceed further to obtain w_2 .

CASE 2: Suppose that $L_1 \neq \emptyset$. Thus, $L_1 \cap L_2 = \emptyset$ and $(L_1 - y_1) \cap B_Y(0, r_1 + \varepsilon) = \emptyset$. Thus, we get $g \in Y^*$ such that $g(y - y_1) \geq 1 \geq g(x)$ for all $y \in L_1$ and $x \in B_Y(0, r_1 + \varepsilon)$.

Now, $\|g\| = \sup_{y \in B_Y} g(y) = \frac{1}{r_1 + \varepsilon} \sup_{y \in B_Y(0, r_1 + \varepsilon)} g(y) \leq \frac{1}{r_1 + \varepsilon}$.

Choose $\delta > 0$ such that $\delta < \frac{1}{r - \frac{\varepsilon}{2} - r_1} (1 - \frac{r_1 + \frac{\varepsilon}{2}}{r_1 + \varepsilon})$. Let $z \in S_Y$ be such that $g(z) \leq -\|g\| + \delta$. Let $w_1 = y_1 + (r - \frac{\varepsilon}{2} - r_1)z \in Y$.

Let $x \in B_X(y_1, r_1)$. Thus,

$$\begin{aligned} \|x - w_1\| &\leq \|x - y_1\| + (r - \frac{\varepsilon}{2} - r_1)\|z\| \\ &\leq r_1 + (r - \frac{\varepsilon}{2} - r_1) \\ &= r - \frac{\varepsilon}{2}. \end{aligned}$$

Thus, $x \in B_X(w_1, r - \frac{\varepsilon}{2})$ and hence $B_X(y_1, r_1) \subseteq B_X(w_1, r - \frac{\varepsilon}{2})$.

Let $y \in B_Y(w_1, r)$. Thus,

$$\begin{aligned} g(y - y_1) &= g(y - w_1) + g(w_1 - y_1) \\ &\leq \|g\|r + (r - \frac{\varepsilon}{2} - r_1)g(z) \\ &\leq \|g\|r + (r - \frac{\varepsilon}{2} - r_1)(-\|g\| + \delta) \\ &\leq \frac{r_1 + \frac{\varepsilon}{2}}{r_1 + \varepsilon} + \delta(r - \frac{\varepsilon}{2} - r_1) < 1. \end{aligned}$$

Thus, for all $y \in B_Y(w_1, r)$, $y \notin L_1$. Hence, $L_1 \cap B_Y(w_1, r) = \emptyset$.

Now, suppose that we have w_i for $i \leq j (< n)$ such that,

$$\begin{aligned} B_X(w_i, r - \frac{\varepsilon}{2}) &\supseteq B_X(y_i, r_i) \text{ for } i \leq j \text{ and} \\ \bigcap_{i=1}^j B_Y(w_i, r) \cap (\bigcap_{i=j+1}^n B_Y(y_i, r_i + \varepsilon)) &= \emptyset. \end{aligned}$$

Let us define $K_1 = \bigcap_{i=1}^j B_Y(w_i, r) \cap (\bigcap_{i=j+2}^n B_Y(y_i, r_i + \varepsilon))$ and $K_2 = B_Y(y_{j+1}, r_{j+1} + \varepsilon)$.

We follow the similar techniques that are used in order to obtain w_1 . The two sets K_1 and K_2 play similar roles as the sets L_1 and L_2 like before, and hence we get w_{j+1} . This completes the induction, and the proof follows. \square

Corollary 2.5. *Let Y be a subspace of X such that $\text{rad}_Y(F) = \text{rad}_X(F)$ for all $F \in \mathcal{F}(Y)$. Then for all $n \in \mathbb{N}$, Y possesses n .X.I.P. in X .*

Proof. Let $y_1, \dots, y_n \in Y$ and $\{B_X(y_i, r)\}_{i=1}^n$ be a finite family of closed balls in X , where $\cap_{i=1}^n B_X(y_i, r) \neq \emptyset$ in X . Hence, $\text{rad}_Y(F) = \text{rad}_X(F) \leq r$. As $\cap_{i=1}^n B_Y(y_i, \text{rad}_Y(F) + \varepsilon) \neq \emptyset$ holds for all $\varepsilon > 0$, we have that the balls $\{B_X(y_i, r)\}_{i=1}^n$ almost intersect in Y . The result now follows from Lemma 2.4. \square

Combining Theorem 2.1, Lemma 2.4, and Corollary 2.5, we obtain the following.

Theorem 2.6. *Let Y be a subspace of X . Then the following are equivalent.*

- (a) Y has n .X.I.P. in X , for all n .
- (b) Y has restricted n .X.I.P. in X , for all n .
- (c) $\text{rad}_Y(F) = \text{rad}_X(F)$, for all $F \in \mathcal{F}(Y)$.
- (d) $\text{rad}_Y^f(F) = \text{rad}_X^f(F)$, for all $F \in \mathcal{F}(Y)$, for all continuous, monotone functions $f : \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}_{\geq 0}$, $n \geq 1$.

We now derive a few characterizations of central subspaces of Banach spaces.

Theorem 2.7. *Let Y be a subspace of X and $Y \in (GC)$. Thus, Y is a restricted central subspace of X if and only if Y is a central subspace of X .*

Proof. Let $\{B_X(y_i, r_i)\}_{i=1}^n$ be a collection of n closed balls in X , where $y_1, \dots, y_n \in Y$ and $r_1, \dots, r_n > 0$ such that $\cap_{i=1}^n B_X(y_i, r_i) \neq \emptyset$.

CLAIM: $\cap_{i=1}^n B_Y(y_i, r_i) \neq \emptyset$.

By [3, Proposition 2.9], it can be sufficiently shown that $\cap_{i=1}^n B_Y(y_i, r_i + \varepsilon) \neq \emptyset$ for all $\varepsilon > 0$. Accordingly, from Lemma 2.4, we conclude the proof. \square

Theorem 2.8. *Let Y be a subspace of X and X admit weighted Chebyshev centers for all finite subsets of Y . Suppose that Y has n .X.I.P. in X for all n . Then, Y is a central subspace of X if and only if $Y \in (GC)$.*

Proof. Suppose that Y is a central subspace of X . Thus, from Theorems 2.1 and 2.2, we obtain $(X, Y, \mathcal{F}(Y))$ has **wrcp**.

Conversely, assume that $Y \in (GC)$. Hence, by Theorem 2.7, it is sufficient to consider balls with centers in Y of equal radii. Let $\{B_X(y_i, r)\}_{i=1}^n$ be a collection of n balls with $y_1, \dots, y_n \in Y$, $r > 0$ such that $\bigcap_{i=1}^n B_X(y_i, r) \neq \emptyset$. Let $F = (y_1, \dots, y_n)$. Thus, $\text{rad}_X(F) \leq r$ and by our assumption, $\text{rad}_Y(F) \leq r$. Owing to the fact that $(X, Y, \mathcal{F}(Y))$ has **wrcp**, $\bigcap_{i=1}^n B_Y(y_i, \text{rad}_Y(F)) \neq \emptyset$ and consequently $\bigcap_{i=1}^n B_Y(y_i, r) \neq \emptyset$. \square

Theorem 2.9. *Let \mathcal{A} be any of the families $\mathcal{F}, \mathcal{K}, \mathcal{B}$, or \mathcal{P} , P be a projection on X of norm-1, $Y = \ker(P)$, and $Z \subseteq P(X)$ be a subspace. If $Y + Z$ is an (almost) $\mathcal{A}(Y + Z) - C$ -subspace of X , then Z is an (almost) $\mathcal{A}(Z) - C$ -subspace of $P(X)$.*

Proof. Let $A \in \mathcal{A}(Z)$. Suppose $\bigcap_{a \in A} B_X(a, r_a) \cap P(X) \neq \emptyset$. Thus, $\bigcap_{a \in A} B_X(a, r_a) \cap X \neq \emptyset$. Based on our assumption, $\bigcap_{a \in A} B_X(a, r_a) \cap (Y + Z) \neq \emptyset$. Let $y_0 + z_0 \in \bigcap_{a \in A} B_X(a, r_a) \cap (Y + Z) \neq \emptyset$. Accordingly, for all $a \in A$,

$$\|z_0 - a\| = \|P(z_0 - y_0 - a)\| \leq \|z_0 - y_0 - a\| \leq r_a.$$

This proves Z is an $\mathcal{A}(Z) - C$ -subspace of $P(X)$. The remaining follows in a similar way as stated above. \square

Proposition 2.10. *Let \mathcal{A} be any of the families $\mathcal{F}, \mathcal{K}, \mathcal{B}$, or \mathcal{P} . Let $Z \subseteq Y \subseteq X$ represent subspaces of X such that Y is an M -ideal in X . If Z is an (almost) $\mathcal{A}(Z) - C$ -subspace of Y and Y has (almost) $\mathcal{A}(Y)$ -IP, then Z is an (almost) $\mathcal{A}(Z) - C$ -subspace of X .*

Proof. Let $A \in \mathcal{A}(Z)$ and $\{B_X(a, r_a)\}_{a \in A}$ be a collection of closed balls, where $r_a > 0$ for all $a \in A$, such that $\bigcap_{a \in A} B_X(a, r_a) \neq \emptyset$. Since Y is an M -ideal in X , we obtain $X^{**} = Y^{\perp\perp} \oplus_{\infty} N^{\perp}$ for a subspace N of X^* . If $x \in \bigcap_{a \in A} B_X(a, r_a)$, then it can be assumed that $x = y + z$, where $y \in Y^{\perp\perp}$ and $z \in N^{\perp}$. Now $\max\{\|y - a\|, \|z\|\} \leq r_a$ for all $a \in A$. Since Y has $\mathcal{A}(Y)$ -IP, there exists $y_0 \in Y$ such that $\|y_0 - a\| \leq r_a$ for all $a \in A$. Thus, based on our assumption, $\bigcap_{a \in A} B_Z(a, r_a) \neq \emptyset$. \square

Theorem 2.11 and Corollary 2.12 indicate the consequences of our previous observations and are derived from the techniques employed in [17, Theorem 2.1, Corollary 2.2]. We now use the well-known Michael selection theorem ([11, Theorem 3.2'']) several times to obtain our observations.

Theorem 2.11. *Let X be a Lindenstrauss space, K and S compact Hausdorff spaces, and $\psi : K \rightarrow S$ a continuous onto map. Let $\psi^* : C(S, X) \rightarrow C(K, X)$ be a continuous isometric embedding expressed as $\psi^*f = f\psi$. Accordingly, $\psi^*C(S, X)$ is a $\mathcal{K}(\psi^*C(S, X)) - C$ -subspace of $C(K, X)$.*

Proof. Let $A \in \mathcal{K}(\psi^*C(S, X))$ and $B = \{f \in C(S, X) : \psi^*f \in A\}$. Suppose $\cap_{f \in B} B(\psi^*f, r_f) \neq \emptyset$, where $r_f > 0$ for all $f \in B$. We define a multivalued map $F : S \rightarrow \mathcal{C}(X)$ by $F(y) = \cap_{f \in B} B(f(y), r_f)$ for all $y \in S$. Evidently, $F(y)$ is closed, convex, and $F(y) \neq \emptyset$ for all $y \in S$.

CLAIM : F is lower semicontinuous.

Let $G \subseteq X$ be open and $y_0 \in \{y \in S : F(y) \cap G \neq \emptyset\}$. Let $a \in F(y_0) \cap G$. Thus, we have $a \in \cap_{f \in B} B(f(y_0), r_f)$ and $B(a, \varepsilon) \subseteq G$ for some $\varepsilon > 0$. Accordingly, there exists an open neighbourhood N of y_0 such that $\|f(y) - f(y_0)\| < \varepsilon$ for all $y \in N$ and $f \in B$. Now for all $y \in N$, we have $B(f(y), r_f) \cap B(a, \varepsilon) \neq \emptyset$ for all $f \in B$ and since X is a Lindenstrauss space, $F(y) \cap G \neq \emptyset$. Thus, $N \subseteq \{y \in S : F(y) \cap G \neq \emptyset\}$ and F is lower semicontinuous. Thus, in accordance with Michael's selection theorem, there exists a continuous $h : S \rightarrow X$ such that $h(y) \in F(y)$ for all $y \in S$. It is evident that $\psi^*h \in \cap_{f \in B} B(\psi^*f, r_f) \cap \psi^*C(S, X)$. \square

Corollary 2.12. *Let S, K, ψ and X be as in Theorem 2.11. Furthermore, fix $y_0 \in S$ and let $M = \{\psi^*f : f \in C(S, X) \text{ and } f(y_0) = 0\}$. Then, M is a $\mathcal{K}(M) - C$ -subspace of $C(K, X)$.*

Proof. Let $A \in \mathcal{K}(M)$ and $B = \{f \in C(S, X) : \psi^*f \in A\}$. Suppose $\cap_{f \in B} B(\psi^*f, r_f) \neq \emptyset$, where $r_f > 0$ for all $f \in B$. Define $F : S \rightarrow \mathcal{C}(X)$ by $F(y) = \cap_{f \in B} B(f(y), r_f)$ for all $y \in S$. As in the proof of Theorem 2.11, we obtain $F(y) \neq \emptyset$ for all $y \in S$. Now we define $F_0 : S \rightarrow \mathcal{C}(X)$ by:

$$F_0(y) = \begin{cases} F(y) & \text{if } y \neq y_0 \\ \{0\} & \text{if } y = y_0 \end{cases}$$

Clearly, $F_0(y)$ is closed, convex and $F_0(y) \neq \emptyset$ for all $y \in S$. Similarly, as done in the proof of Theorem 2.11, it can be shown that F_0 is lower semicontinuous. Thus, by Michael's selection theorem, we obtain $h : S \rightarrow X$ such that $h(y) \in F_0(y)$ for all $y \in S$. Thus, $\psi^*h \in \cap_{f \in B} B(\psi^*f, r_f) \cap M$. \square

Proceeding similarly to the proof of Theorem 2.11, the following can be obtained.

Corollary 2.13. (a) Let X be a Lindenstrauss space, T compact Hausdorff space and S be a closed subset of T . If $M = \{f \in C(T, X) : f|_S = 0\}$, then M is a $\mathcal{K}(M) - C$ -subspace of $C(T, X)$.
 (b) Let X be a Lindenstrauss space and T be a compact Hausdorff space. Then, $C(T, X)$ is a $\mathcal{K}(C(T, X)) - C$ -subspace of $\ell_\infty(T, X)$.

Our following observation shows that the characteristic of being a central subspace is stable in the spaces of continuous functions. Let us recall the manner in which the Lindenstrauss spaces are characterized in the introduction.

Theorem 2.14. Let \mathcal{A} be any of the families \mathcal{F}, \mathcal{K} . Let X be a Lindenstrauss space, Y a $\mathcal{A}(Y) - C$ -subspace of X , and T a compact Hausdorff space. Then, $C(T, Y)$ is a $\mathcal{A}(C(T, Y)) - C$ -subspace of $\ell_\infty(T, X)$.

Proof. We follow the similar techniques that are used in Theorem 2.11.

Let $A \in \mathcal{A}(C(T, Y))$ and $\cap_{f \in A} B(f, r_f) \cap \ell_\infty(T, X) \neq \emptyset$, where $r_f > 0$ for all $f \in A$. Define $F : T \rightarrow \mathcal{C}(Y)$ by $F(t) = \cap_{f \in A} B(f(t), r_f) \cap Y$. Now since for all $t \in T$, $\cap_{f \in A} B(f(t), r_f) \cap X \neq \emptyset$, based on our assumption, $\cap_{f \in A} B(f(t), r_f) \cap Y \neq \emptyset$ for all $t \in T$. Clearly, $F(t)$ is closed and convex for all $t \in T$.

By using similar arguments stated in Theorem 2.11 it follows that F is lower semicontinuous.

By Michael's selection theorem, we obtain $h : T \rightarrow Y$ such that $h(t) \in F(t)$ for all $t \in T$. This completes the proof. \square

We end this section by providing a straightforward application of the conditional expectation projection $E : L_p(\Sigma, X) \rightarrow L_p(\Sigma', X)$, for $1 \leq p < \infty$. In the following (Ω, Σ, μ) denotes a finite measure space and Σ' is a sub σ -algebra of Σ .

Proposition 2.15. Let Y be a subspace of X and \mathcal{A} be any of the families $\mathcal{F}, \mathcal{K}, \mathcal{B}$, or \mathcal{P} . Let $L_1(\Sigma, Y)$ be a $\mathcal{A}(L_1(\Sigma, Y)) - C$ -subspace of $L_1(\Sigma, X)$. Consequently, $L_1(\Sigma', Y)$ is an $\mathcal{A}(L_1(\Sigma', Y)) - C$ -subspace of $L_1(\Sigma', X)$.

Remark 2.16. In section 3, we encounter cases in Theorem 3.9 and 3.11 when $L_1(\Sigma, Y)$ is a $\mathcal{A}(L_1(\Sigma, Y)) - C$ -subspace of $L_1(\Sigma, X)$, for subspaces Y of X .

3. WEIGHTED RESTRICTED CHEBYSHEV CENTERS IN SPACES OF VECTOR-VALUED FUNCTIONS

First, we note that the following may be established by the arguments given in [1, Theorem 2].

Theorem 3.1. *Let X be a Banach space which is uniformly convex and let T be a topological space. The Banach space $C_b(T, X)$ admits weighted Chebyshev centers for all closed bounded subsets of $C_b(T, X)$ and bounded weights ρ .*

The following result demonstrates that in the spaces of Bochner integrable functions $L_p(\Sigma, X)$, where (Ω, Σ, μ) is a finite measure space, the property **\mathcal{F} -rcp** is stable for a particular class of functions \mathcal{F} . For $n \in \mathbb{N}$ and $1 \leq p < \infty$, \mathcal{F}_p denotes a countable family of functions viz. $(f_p^n)_{n=1}^\infty$, where $f_p^n : \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}_{\geq 0}$ is defined by $f_p^n(\alpha_1, \dots, \alpha_n) = (\sum_{i=1}^n |\alpha_i|^p)^{\frac{1}{p}}$.

Theorem 3.2. *Consider the measure space (Ω, Σ, μ) as stated above. Let Y be a separable subspace of X and consider the corresponding function spaces $L_p(\Sigma, Y) \subseteq L_p(\Sigma, X)$. Let \mathcal{F}_p be the family of functions as stated above, then:*

- (a) $(X, Y, \mathcal{F}(X))$ has **wrcp** if and only if $(L_\infty(\Sigma, X), L_\infty(\Sigma, Y), \mathcal{F}(L_\infty(\Sigma, X)))$ has **wrcp**.
- (b) for $1 \leq p < \infty$, $(X, Y, \mathcal{F}(X))$ has **\mathcal{F}_p -rcp** if and only if $(L_p(\Sigma, X), L_p(\Sigma, Y), \mathcal{F}(L_p(\Sigma, X)))$ has **\mathcal{F}_p -rcp**.

Proof. (a). Let $F \in \mathcal{F}(L_\infty(\Sigma, X))$. We prove the result when $\text{card}(F) = 2$ because new ideas are not involved for higher values of $\text{card}(F)$. Let $F = \{f_1, f_2\}$ and $\rho = (\rho_1, \rho_2)$ be the corresponding weights. Now, there exists a measurable subset $E \subseteq \Omega$ such that $\mu(E) = 0$ and f_1, f_2 are bounded on $\Omega \setminus E$. Define for all $t \in \Omega \setminus E$, $F_t = \{f_1(t), f_2(t)\}$. Let $G = \{(t, y) \in (\Omega \setminus E) \times Y : r_\rho(y, F_t) = \text{rad}_Y^\rho(F_t)\}$. Since $(X, Y, \mathcal{F}(X))$ has **wrcp**, the projection of G on $\Omega \setminus E$ is $\Omega \setminus E$.

Suppose (y_n) is dense in Y . Then, $G = \cap_{n \in \mathbb{N}} \{(t, y) \in (\Omega \setminus E) \times Y : r_\rho(y, F_t) \leq r_\rho(y_n, F_t)\}$. Since all the involved functions are measurable, G is a measurable set. Thus, as a consequence of the von Neumann selection theorem [13, Theorem 7.2], we obtain a measurable function $g_0 : (\Omega \setminus E) \rightarrow Y$ such that $(t, g_0(t)) \in G$ for μ -a.e t . Then, $g_0(t) \in \text{Cent}_Y^\rho(F_t)$ for μ -a.e.

$t \in \Omega \setminus E$. Clearly, $g_0 \in L_\infty(\Sigma, Y)$. For $g \in L_\infty(\Sigma, Y)$, we obtain

$$\begin{aligned}
r_\rho(g, F) &= \max_{i=1,2} \operatorname{ess\,sup}_{t \in \Omega} \rho_i \|g(t) - f_i(t)\| \\
&= \operatorname{ess\,sup}_{t \in \Omega} \max_{i=1,2} \rho_i \|g(t) - f_i(t)\| \\
&\geq \operatorname{ess\,sup}_{t \in \Omega} \max_{i=1,2} \rho_i \|g_0(t) - f_i(t)\| \\
&= \max_{i=1,2} \operatorname{ess\,sup}_{t \in \Omega} \rho_i \|g_0(t) - f_i(t)\| \\
&= r_\rho(g_0, F).
\end{aligned}$$

Hence, $g_0 \in \operatorname{Cent}_{L_\infty(\Sigma, Y)}^\rho(F)$.

(b). Let $F \in \mathcal{F}(L_p(\Sigma, X))$. We prove the result when $\operatorname{card}(F) = 2$, as no new ideas are involved for higher values of $\operatorname{card}(F)$. Let $F = \{f_1, f_2\}$ and $f_p^2 : \mathbb{R}_{\geq 0}^2 \rightarrow \mathbb{R}_{\geq 0}$ represent the function in \mathcal{F}_p as stated above. Accordingly, there exists a measurable subset $E \subseteq \Omega$ such that $\mu(E) = 0$ and f_1, f_2 are finite valued on $\Omega \setminus E$. Define for all $t \in \Omega \setminus E$, $F_t = \{f_1(t), f_2(t)\}$. Let $G = \{(t, y) \in (\Omega \setminus E) \times Y : r_{f_p^2}(y, F_t) = \operatorname{rad}_Y^{f_p^2}(F_t)\}$. Since $(X, Y, \mathcal{F}(X))$ has \mathcal{F}_p -**r**cp, the projection of G on $\Omega \setminus E$ is $\Omega \setminus E$.

Thus, in a similar manner as the proof of (a), we obtain a measurable function $g_0 : (\Omega \setminus E) \rightarrow Y$ such that $(t, g_0(t)) \in G$ μ -a.e. t . Clearly, $g_0 \in L_p(\Sigma, Y)$. It is easy to see that $g_0 \in \operatorname{Cent}_{L_p(\Sigma, Y)}^{f_p^2}(F)$. \square

Our next few results are derived under an assumption that the function $r_f(\cdot, F)$ is lower semicontinuous w.r.t. a suitable topology on X , for a given bounded set $F \in \mathcal{B}(X)$, which is now stated in the following remark. For an arbitrary set Γ , we define $\ell_1(\Gamma) := \{\vartheta : \Gamma \rightarrow \mathbb{R}_{\geq 0} : \sum_\gamma \vartheta(\gamma) < \infty\}$.

Remark 3.3. (a) Clearly $r_f(\cdot, F) : X \rightarrow \mathbb{R}_{\geq 0}$ is a continuous and coercive function, where $r_f(x, F) = f(\|x - a\|_{a \in F})$, when $F \in \mathcal{B}(X)$ and $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$ is a continuous, monotone, and coercive function.

(b) In (a), if we choose $\rho : F \rightarrow \mathbb{R}_{\geq 0}$ a bounded weight then $r_\rho(\cdot, F) : (X, w) \rightarrow \mathbb{R}_{\geq 0}$ is lower semicontinuous. Moreover $r_\rho(\cdot, F) : (X, w^*) \rightarrow \mathbb{R}_{\geq 0}$ is also lower semicontinuous, when X is a dual space.

(c) For $F \in \mathcal{B}(X)$ and for any $\vartheta \in \ell_1(F)$, since ϑ induces a continuous, monotone functional on $\ell_\infty(F)$, it is easy to observe that $r_\vartheta(\cdot, F) : (X, w) \rightarrow \mathbb{R}_{\geq 0}$ is lower semicontinuous and so is when X is a dual space endowed with the w^* -topology.

Proposition 3.4. *Suppose that X is a dual space, Y is a w^* -closed subspace of X , $F \in \mathcal{B}(X)$ and $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$ be a monotone, coercive function such that $r_f(\cdot, F) : X \rightarrow \mathbb{R}_{\geq 0}$ is w^* -lower semicontinuous. Then, $\text{Cent}_Y^f(F) \neq \emptyset$.*

Proof. Let $(y_n) \subseteq Y$ be such that $r_f(y_n, F) \rightarrow \text{rad}_Y^f(F)$. As f is coercive, the sequence (y_n) is bounded. Hence there exists $y_0 \in Y$ such that $y_\alpha \rightarrow y_0$ in w^* -topology, for some subnet (y_α) of (y_n) . It is clear that $r_f(y_0, F) = \text{rad}_Y^f(F)$ and hence $y_0 \in \text{Cent}_Y^f(F)$. \square

The following is a variation of [6, Proposition IV.1.2]. For a $B \in \mathcal{B}(X)$ and $f : \ell_\infty(B) \rightarrow \mathbb{R}_{\geq 0}$, f is said to be *weakly strictly monotone* if $f(\varphi_1) < f(\varphi_2)$, whenever $\varphi_1(t) < \varphi_2(t)$, for all $t \in B$.

Theorem 3.5. *Let X be an L -embedded space and Y be a subspace of X , which is also L -embedded and $F \in \mathcal{B}(X)$. Then,*

- (a) *for all monotone, coercive functions $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$, such that $r_f(\cdot, F) : X^{**} \rightarrow \mathbb{R}_{\geq 0}$ is w^* -lower semicontinuous, $\text{Cent}_Y^f(F) \neq \emptyset$.*
- (b) *for all weakly strictly monotone, coercive functions $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$, such that $r_f(\cdot, F) : X^{**} \rightarrow \mathbb{R}_{\geq 0}$ is w^* -lower semicontinuous, $\text{Cent}_Y^f(F)$ is weakly compact.*

Proof. (a). Let $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$ be a monotone, coercive function such that $r_f(\cdot, F)$ is w^* -lower semicontinuous. Let $P : X^{**} \rightarrow X$ be the L -projection. Thus, from [6, Theorem IV.1.2], we have $P(\overline{Y}^{w^*}) \subseteq Y$. By Proposition 3.4 there exists $y^{**} \in \overline{Y}^{w^*}$ such that $r_f(y^{**}, F) = \text{rad}_{\overline{Y}^{w^*}}^f(F)$. Based on the monotonicity of f , we obtain,

$$(1) \quad \text{rad}_Y^f(F) \geq \text{rad}_{\overline{Y}^{w^*}}^f(F) = r_f(y^{**}, F) \geq r_f(Py^{**}, F) \geq \text{rad}_Y^f(F).$$

Since $Py^{**} \in Y$, our first assertion follows.

(b). Let $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$ be a weakly strictly monotone, coercive function such that $r_f(\cdot, F)$ is w^* -lower semicontinuous. By (1), we obtain a $y^{**} \in \overline{Y}^{w^*}$ such that, $r_f(Py^{**}, F) = r_f(y^{**}, F)$. Now for all $x \in F$,

$$\|y^{**} - x\| = \|Py^{**} - x + (Id - P)y^{**}\| = \|Py^{**} - x\| + \|y^{**} - Py^{**}\|.$$

This leads to $r_f(Py^{**}, F) = f((\|Py^{**} - x\| + \|y^{**} - Py^{**}\|)_{x \in F})$.

Since f is weakly strictly monotone, we have $y^{**} = Py^{**}$ and consequently, $\text{Cent}_Y^f(F) = \text{Cent}_{\overline{Y}^{w^*}}^f(F)$. Hence, the w^* -closure of $\text{Cent}_Y^f(F)$ is contained in X , so it is weakly compact. \square

As a corollary of the Theorem 3.5, we have,

Corollary 3.6. (a) *Let X be an L -embedded space and Y be a subspace of X , which is also L -embedded. Suppose that Y has $n.X.I.P.$ in X for all $n \in \mathbb{N}$. Then, Y is a central subspace of X .*

(b) *Let $L_1(\Sigma, X)$ be an L -embedded space. Let $F \in \mathcal{B}(L_1(\Sigma, X))$ and $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$ be a monotone, coercive function such that $r_f(\cdot, F) : L_1(\Sigma, X)^{**} \rightarrow \mathbb{R}_{\geq 0}$ is w^* -lower semicontinuous. Then, for all sub σ -algebra $\Sigma' \subseteq \Sigma$, $\text{Cent}_{L_1(\Sigma', X)}^f(F) \neq \emptyset$.*

Proof. (a). From our assumption, it follows that both X and Y admit weighted Chebyshev center for finite subsets of Y . The result now follows from Theorem 2.8, and Theorem 3.5.

(b). Since $L_1(\Sigma', X)$ is one complemented in $L_1(\Sigma, X)$, it follows from [6, Proposition IV.1.5], that $L_1(\Sigma', X)$ is also L -embedded. Thus, our conclusion follows from Theorem 3.5. \square

Similar arguments stated in the proof of Theorem 3.5(b) also lead to the following.

Lemma 3.7. *Let X be an L -embedded space and Y be a subspace of X , which is also L -embedded. Let $F \in \mathcal{B}(X)$ and $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$ be a strictly monotone, coercive function such that $r_f(\cdot, F) : X^{**} \rightarrow \mathbb{R}_{\geq 0}$ is w^* -lower semicontinuous. Suppose that (y_n) is a sequence in Y such that $\lim_n r_f(y_n, F) = \text{rad}_Y^f(F)$. Then, (y_n) is relatively weakly compact.*

As a consequence of Lemma 3.7, we have the following.

Theorem 3.8. *Let X be a Banach space such that $L_1(\Sigma, X)$ is L -embedded. Let $\{\Sigma_n\}_{i=1}^\infty$ be an increasing sequence of sub σ -algebras. Let Σ_∞ be the σ -algebra generated by $\cup_{n \in \mathbb{N}} \Sigma_n$. Let $F \in \mathcal{B}(L_1(\Sigma, X))$ and $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$ be a strictly monotone, coercive function such that $r_f(\cdot, F) : L_1(\Sigma, X)^{**} \rightarrow \mathbb{R}_{\geq 0}$ is w^* -lower semicontinuous. Suppose that $\text{rad}_{L_1(\Sigma_n, X)}^f(F) = r_f(f_n, F)$ for some $f_n \in L_1(\Sigma_n, X)$. Then, (f_n) is relatively weakly compact, and any weak limit point of this sequence belongs to $\text{Cent}_{L_1(\Sigma_\infty, X)}^f(F)$.*

Proof. We follow the techniques similar to those used in [14, Theorem 12]. Let E_n be the conditional expectation projection of $L_1(\Sigma_\infty, X)$ onto $L_1(\Sigma_n, X)$ for all $n \in \mathbb{N}$. It is known that $E_n(g) \rightarrow g$ for all $g \in L_1(\Sigma_\infty, X)$.

We have for any $n \in \mathbb{N}$, $\text{rad}_{L_1(\Sigma_\infty, X)}^f(F) \leq \text{rad}_{L_1(\Sigma_n, X)}^f(F) \leq r_f(f_n, F)$ and hence $\text{rad}_{L_1(\Sigma_\infty, X)}^f(F) \leq \liminf_n r_f(f_n, F)$.

Now for any $g \in L_1(\Sigma_\infty, X)$, we have,

$$\limsup_n \text{rad}_{L_1(\Sigma_n, X)}^f(F) \leq \lim_n r_f(E_n(g), F) \leq r_f(g, F).$$

Hence, $\limsup_n r_f(f_n, F) = \limsup_n \text{rad}_{L_1(\Sigma_n, X)}^f(F) \leq \text{rad}_{L_1(\Sigma_\infty, X)}^f(F)$.

Thus, $\text{rad}_{L_1(\Sigma_\infty, X)}^f(F) = \lim_n r_f(f_n, F) = \lim_n \text{rad}_{L_1(\Sigma_n, X)}^f(F)$. Hence, by Lemma 3.7, (f_n) is relatively weakly compact. Using the w^* -lower semicontinuity of $r_f(\cdot, F)$, we have the weak limit point of (f_n) belonging to $\text{Cent}_{L_1(\Sigma_\infty, X)}^f(F)$. \square

Our next observation follows from [16, Theorem 5.1] and the existence of the conditional expectation operator.

Theorem 3.9. *Let X denote a Banach space such that X^* has RNP. Let (Ω, Σ, μ) be a finite measure space. Let $F \in \mathcal{B}(L_1(\Sigma, X^*))$ and $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$ be a monotone, coercive function such that $r_f(\cdot, F) : L_1(\Sigma, X^{**}) \rightarrow \mathbb{R}_{\geq 0}$ is w^* -lower semicontinuous. Then, for all sub σ -algebra $\Sigma' \subseteq \Sigma$, $\text{Cent}_{L_1(\Sigma', X^*)}(F) \neq \emptyset$.*

Let us recall that $\mathcal{WC}(X)$ denotes the set of all weakly compact subsets of X . Additionally, let us also recall the following theorem of Dunford (see [5, pg. 101]).

Theorem 3.10. *Let X be a reflexive Banach space. Then, $K \subseteq L_1(\mu, X)$ is relatively weakly compact if and only if K is bounded and uniformly integrable.*

It is apparent from the proof of the above Theorem that for any Banach space X , a relatively weakly compact subset of $L_1(\mu, X)$ is bounded and uniformly integrable.

Theorem 3.11. *Let Y be a reflexive subspace of X and (Ω, Σ, μ) be a finite measure space.*

- (a) *Then for $F \in \mathcal{WC}(L_1(\Sigma, X))$, for all sub σ -algebra $\Sigma' \subseteq \Sigma$ and for all monotone, coercive function $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$ such that $r_f(\cdot, F) : L_1(\Sigma, X) \rightarrow \mathbb{R}_{\geq 0}$ is w -lower semicontinuous, $\text{Cent}_{L_1(\Sigma', Y)}^f(F) \neq \emptyset$.*
- (b) *Then for all $F \in \mathcal{B}(L_\infty(\Sigma, X))$, monotone, coercive function $f : \ell_\infty(F) \rightarrow \mathbb{R}_{\geq 0}$ such that $r_f(\cdot, F) : L_1(\Sigma, X^*) \rightarrow \mathbb{R}_{\geq 0}$ is w^* -lower semicontinuous, $\text{Cent}_{L_\infty(\Sigma, Y)}^f(F) \neq \emptyset$.*

Proof. (a). Let $(g_n) \subseteq L_1(\Sigma', Y)$ be a such that $r_f(g_n, F) \rightarrow \text{rad}_{L_1(\Sigma', Y)}^f(F)$. Clearly, (g_n) is bounded. Hence by a vector-valued version of Kadec-Pełczyński-Rosenthal theorem ([4, Lemma 2.1.3]), there exists a subsequence (g_{n_k}) of (g_n) and a sequence of pairwise disjoint sets $(A_k) \subseteq \Sigma'$ such that $(g_{n_k} \chi_{A_k^c})$ is uniformly integrable in $L_1(\Sigma', Y)$. Since, $\sum_{k=1}^{\infty} \mu(A_k) = \mu(\cup_{k=1}^{\infty} A_k) \leq \mu(\Omega) < \infty$, we have $\mu(A_k) \rightarrow 0$.

Let us choose $h \in F$ then,

$$\|g_{n_k} \chi_{A_k^c} - h\|_1 = \int_{A_k} \|h\| d\mu + \int_{A_k^c} \|g_{n_k} - h\| d\mu \leq \int_{A_k} \|h\| d\mu + \|g_{n_k} - h\|_1.$$

As F is uniformly integrable, we get $\lim_k \int_{A_k} \|h\| d\mu = 0$ uniformly for all $h \in F$. Now, it is easy to see that $r_f(g_{n_k} \chi_{A_k^c}, F) \rightarrow \text{rad}_{L_1(\Sigma', Y)}^f(F)$. Also since $(g_{n_k} \chi_{A_k^c})$ is a bounded sequence in $L_1(\Sigma', Y)$, by Theorem 3.10, $(g_{n_k} \chi_{A_k^c})$ is relatively weakly compact in $L_1(\Sigma', Y)$. Let us denote by $(g_{n_{k(j)}})$, a weakly convergent subsequence of $(g_{n_k} \chi_{A_k^c})$ converging weakly to $g \in L_1(\Sigma', Y)$. Then $r_f(g, F) \leq \liminf_j r_f(g_{n_{k(j)}}, F) = \text{rad}_{L_1(\Sigma', Y)}^f(F)$. Hence $r_f(g, F) = \text{rad}_{L_1(\Sigma', Y)}^f(F)$.

(b). We consider $L_\infty(\Sigma, X)$ to be canonically embedded in $L_1(\Sigma, X^*)^*$. It follows from Proposition 3.1 of [12] that $L_\infty(\Sigma, Y)$ is w^* -closed in $L_1(\Sigma, X^*)^*$. Hence by the w^* -lower semicontinuity of $r_f(\cdot, F)$, we have an $g \in L_\infty(\Sigma, Y)$ such that $r_f(g, F) = \text{rad}_{L_\infty(\Sigma, Y)}^f(F)$. \square

Remark 3.12. *Using similar arguments, it can be shown that for a reflexive subspace Y of X and for a sub- σ -algebra $\Sigma' \subseteq \Sigma$, $(L_1(\Sigma, X), B_{L_1(\Sigma', Y)}, \mathcal{WC}(L_1(\Sigma, X)))$ has **rcp**.*

Let \mathcal{F} be the family of all monotone functions $f : (\mathbb{R}_{\geq 0}^n, \|\cdot\|_\infty) \rightarrow \mathbb{R}_{\geq 0}$, where $n \geq 1$.

Proposition 3.13. *Let P be a projection on X of norm-1 and $Y = \ker(P)$. Let $Z \subseteq P(X)$ be a subspace. If $(X, Y + Z, \mathcal{F}(X))$ has **\mathcal{F} -rcp**, then $(P(X), Z, \mathcal{F}(P(X)))$ has **\mathcal{F} -rcp**.*

Proof. Let $F \in \mathcal{F}(P(X))$ and $\text{card}(F) = n$. Suppose that, $f \in \mathcal{F}$, where $f : \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}_{\geq 0}$ be monotone. Based on our assumption, there exists $y_0 + z_0 \in Y + Z$ such that $r_f(y_0 + z_0, F) = \text{rad}_{Y+Z}^f(F) \leq \text{rad}_Z^f(F)$. Then, $r_f(z_0, F) = r_f(P(y_0 + z_0), F) \leq r_f(y_0 + z_0, F) \leq \text{rad}_Z^f(F)$. \square

- Remark 3.14.** (a) *An identical arguments stated in the proof of Proposition 3.13 also work to conclude that F has \mathcal{F} -**rcp** in Z whenever $F \in \mathcal{B}(P(X))$ and F has \mathcal{F} -**rcp** in $Y + Z$. Here we consider \mathcal{F} to be the family $\ell_\infty(F)$.*
- (b) *Similar conditions as that in Proposition 3.13, it can be used to prove that if $(X, B_{Y+Z}, \mathcal{B}(X))$ has **rcp**, then $(P(X), B_Z, \mathcal{B}(P(X)))$ has **rcp**.*

REFERENCES

- [1] Amir Dan, *Chebyshev centers and uniform convexity*, Pacific J. Math. **77**(1), (1978) 1–6.
- [2] Bandyopadhyay Pradipta, Dutta S., *Weighted Chebyshev centers and intersection properties of balls in Banach spaces*, Function spaces (Edwardsville, IL, 2002), 43–58, Contemp. Math., **328**, Amer. Math. Soc., Providence, RI, 2003.
- [3] Bandyopadhyay Pradipta, Rao T. S. S. R. K., *Central subspaces of Banach spaces*, J. Approx. Theory **103**(2) (2000), 206–222.
- [4] Cembranos Pilar, Mendoza José, *Banach spaces of vector-valued functions*, Lecture Notes in Mathematics, **1676**, Springer, Berlin, 1997.
- [5] Diestel J., Uhl J.J., *Vector measures*, Mathematical Surveys, No. 15, American Mathematical Society, Providence, RI, 1977.
- [6] Harmand P., Werner D., Werner W., *M-ideals in Banach spaces and Banach algebras*, Lecture Notes in Mathematics **1547**, Springer-Verlag, Heidelberg, 1993.
- [7] Indumathi V., Lalithambigai S., *Ball proximal spaces*, J. Convex Anal. **18**(2), (2011) 353–366.
- [8] Jayanarayanan C.R., *Intersection properties of balls in Banach spaces*, J. Funct. Spaces Appl. (2013), Article ID 902694.
- [9] Lima Á., *Uniqueness of Hahn-Banach extensions and liftings of linear dependences*, Math. Scand. **53**(1) (1983), 97–113.
- [10] Lindenstruass J., *Extension of compact operators*. Mem. Amer. Math. Soc. **48** (1964).
- [11] Michael E., *Continuous selections. I*, Ann. of Math. **63**(2) (1956), 361–382.
- [12] Pakhrou Tijani, *Best simultaneous approximation in $L_\infty(\mu, X)$* . Math. Nachr. **281**(3) (2008), 396–401.
- [13] Parthasarathy T., *Selection theorems and their applications*, Lecture Notes in Mathematics, **263**. Springer-Verlag, Berlin-New York, 1972.
- [14] Rao T.S.S.R.K., *Approximation properties for spaces of Bochner integrable functions*, J. Math. Anal. Appl. **423**(2) (2015) 1540–1545.
- [15] Veselý Libor, *Chebyshev centers in hyperplanes of c_0* , Czechoslovak Math. J. **52**(127) (2002), 721–729.
- [16] Veselý Libor, *Generalized centers of finite sets in Banach spaces* Acta Math. Univ. Comenian. (N.S.) **66**(1) (1997), 83–115.

- [17] Yost D.T., *Best approximation and intersections of balls in Banach spaces*. Bull. Austral. Math. Soc. **20**(2), (1979) 285–300.

DEPT. OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY HYDERABAD, INDIA
Email address: `ma20resch11006@iith.ac.in` & `tanmoy@math.iith.ac.in`