Recent $B^+ \rightarrow K^+ \nu \bar{\nu}$ Excess and Muon g - 2 Illuminating Light Dark Sector with Higgs Portal

Shu-Yu Ho,* Jongkuk Kim,[†] and Pyungwon Ko[‡]

Korea Institute for Advanced Study, Seoul 02455, Republic of Korea

The Belle II collaboration recently announced that they observed the $B^+ \to K^+ \nu \bar{\nu}$ decay process for the first time. This dineutrino mode of $B^+ \to K^+ \nu \bar{\nu}$ has been theoretically identified as a very clean channel. However, their result encounters a 2.7 σ deviation from the Standard Model (SM) calculation. On the other hand, last year, Fermilab released new data on muon g - 2 away from the SM expectation with 5σ . In this letter, we study the simplest UV-complete U(1)_{Lµ-L₇}-charged complex scalar Dark Matter (DM) model. Thanks to the existence of light dark Higgs boson and light dark photon, we can explain the observed relic density of DM and resolve the results reported by both Belle II and Fermilab experiments simultaneously. As a byproduct, the Hubble tension is alleviated by taking $\Delta N_{\rm eff} \simeq 0.3$ induced by the light dark photon.

INTRODUCTION

Recently, the Belle II collaboration has found the first evidence of the $B^+ \rightarrow K^+ \nu \bar{\nu}$ rare decay predicted by the SM [1]. The decay branching fraction was measured with two different techniques: Hadronic-tagged and inclusivetagged analyses. The combined result is given by

$$\mathcal{B}(B^+ \to K^+ \nu \bar{\nu})_{\text{exp}} = (2.3 \pm 0.7) \times 10^{-5} .$$
 (1)

On the other hand, it is known that the decay branching fraction of $B^+ \to K^+ \nu \bar{\nu}$ is calculated with high accuracy in the SM [2]:

$$\mathcal{B}(B^+ \to K^+ \nu \bar{\nu})_{\mathsf{SM}} = (4.97 \pm 0.37) \times 10^{-6}$$
. (2)

Comparing the difference between these two results, the Belle II measurement has a 2.7σ deviation from the SM prediction. Although this deviation is not yet statistically significant enough to be interpreted as Beyond Standard Model (BSM) signals. However, it would be worthwhile to entertain the possibility of accommodating it in some well-motivated BSM models with light dark sectors. If we adopt this line of thought, this deviation implies the presence of a BSM effect on the $b \rightarrow s\nu\bar{\nu}$ transition, which is treated as the missing energy $\not E$ in the final states of the B^+ to K^+ decay with a branching fraction of

$$\mathcal{B}(B^+ \to K^+ \not\!\!\!E)_{\rm NP} = (1.8 \pm 0.7) \times 10^{-5} .$$
 (3)

This discrepancy has been interpreted by effective theories [3–6], three-body $B \to K\chi\bar{\chi}$ decay with light DM or dark photon [7–12], and the other approaches [13, 14].

In Ref. [9], the authors stressed that the Belle II analysis provides information on the q^2 spectrum, indicating that there is a peak localized around $q^2 = 4 \text{ GeV}^2$. This can be described by a dark boson with a mass of 2 GeVprovided that its couplings to the SM fermions are sufficiently tiny. In comparison to Ref. [9], Ref. [10] studied a three-body decay involving a DM pair in a Higgs portal DM model. However, they predicted superabundant DM relic density. To obtain the correct relic abundance, they have to either introduce a new DM annihilation channel or modify the standard cosmology. Nonetheless, Ref. [11] performed likelihood analyses and concluded that the B three-body decay is most favored. Combining all of these interpretations, the observed enhancement of the signals might indicate that new particles in the dark sector leave their imprints.

In 2023, the Fermilab Muon g-2 Collaboration released a new measurement of the muon magnetic moment [15]

$$a_{\mu}^{\exp} = 116\ 592\ 059(22) \times 10^{-11}$$
, (4)

and the theoretical value computed by the SM is [16]

$$a_{\mu}^{\text{SM}} = 116\ 591\ 810(43) \times 10^{-11}$$
 (5)

This tension has increased from 4.2σ released in 2021 to 5.1σ . However, it seems to reduce to 1.6σ by taking into account the new lattice QCD result from BMW Collaboration [17]. Ongoing efforts are made to understand the discrepancy in the results obtained by two different theoretical approaches. In this work, we will consider 5.1σ is real to be explained by new physics BSM.

A simple $U(1)_{L_{\mu}-L_{\tau}}$ -charged DM model can explain both $\Delta a_{\mu} = a_{\mu}^{\exp} - a_{\mu}^{\text{SM}}$ and thermal DM if the vector boson mass in the Proca Lagrangian is near the resonance region, where $m_V \simeq 2m_{\text{DM}}$ [18–20]. However, it was emphasized in Ref. [21] that this fine-tuning mass condition can be relaxed if we promote these models with the Abelian Higgs mechanism.

In this letter, we investigate the simplest UV-complete $U(1)_{L_{\mu}-L_{\tau}}$ -charged complex scalar DM model. We will demonstrate that this DM model can produce the correct relic abundance of DM, and resolve the tensions of Δa_{μ} at Fermilab and $B^+ \to K^+ \nu \bar{\nu}$ at Belle II at the same time by fully taking into account of dark Higgs boson in the whole analyses (See Ref. [22] for reviews on the roles of dark Higgs boson in (astro) particle physics and cosmology). Remarkably, we can explain the $B^+ \to K^+ \nu \bar{\nu}$ excess through either two- or three-body decay accom-

panied by the dark Higgs boson with a nonzero vacuum expectation value (VEV) or DM particle, respectively.

MODEL OVERVIEW

A simple extension of the SM to account for the Δa_{μ} and $B^+ \to K^+ \nu \bar{\nu}$ excess is the gauged U(1)_{L_µ-L_τ} model (known to be anomaly-free without adding extra chiral fermions [23, 24]) including a dark Higgs boson. In this scenario, the $B^+ \to K^+ \nu \bar{\nu}$ excess can be addressed by the dark Higgs boson (or its dark decay product) that is interpreted as the missing energy akin to neutrinos. On the other hand, the Δa_{μ} can be explained by the dark $U(1)_{L_{\mu}-L_{\tau}}$ gauge boson with a mass generated by the Higgs mechanism. To account for DM, we further introduce a complex scalar DM charged under the $U(1)_{L_{\mu}-L_{\tau}}$ symmetry which, as we will discuss soon, would mainly annihilate into the dark gauge bosons via the Higgs portal. Then these dark gauge bosons will eventually decay into neutrinos if they are lighter than the muon, which is the case we shall consider here. This is in contrast to the real singlet scalar DM via the Higgs portal where the final states of the DM annihilation are electrically charged SM particles [25–27].

With the above model setup, the $U(1)_{L_{\mu}-L_{\tau}}$ charge assignments for the relevant SM particles and new particles are given as follows

$$\widehat{\mathcal{Q}}_{\mathsf{L}_{\mu}-\mathsf{L}_{\tau}}\left(\nu_{\mu},\nu_{\tau},\mu,\tau,X,\Phi\right) = \left(1,-1,1,-1,\mathcal{Q}_{X},\mathcal{Q}_{\Phi}\right),\quad(6)$$

(

where X is the complex singlet scalar DM, and $\Phi = \frac{1}{\sqrt{2}} (v_{\Phi} + \phi)$ is the dark Higgs with a nonvanishing VEV v_{Φ} that breaks the U(1)_{Lµ-L_τ} gauge symmetry spontaneously. To ensure the DM to be stable or very long-lived $(\tau_X \gtrsim 10^{26} \text{ sec})$, here we assume that $Q_X = 1$ and Q_{Φ} in such a way that there are no gauge invariant operators up to dim-5 that would make the DM decay into the SM particles [28, 29].

Given this particle charge assignments, the renormalizable and gauge invariant Lagrangian density associated with the present study is given by

$$\mathcal{L} = |\mathcal{D}_{\rho}\Phi|^{2} + |\mathcal{D}_{\rho}X|^{2} - \frac{1}{4} \left(\partial_{\rho}Z'_{\omega} - \partial_{\omega}Z'_{\rho}\right)^{2} - m_{X}^{2}|X|^{2} - g_{X} \left(\bar{\mu}\gamma^{\rho}\mu - \bar{\tau}\gamma^{\rho}\tau + \bar{\nu}_{L\mu}\gamma^{\rho}\nu_{L\mu} - \bar{\nu}_{L\tau}\gamma^{\rho}\nu_{L\tau}\right)Z'_{\rho} - \lambda_{\Phi X}|X|^{2} \left(|\Phi|^{2} - \frac{1}{2}v_{\Phi}^{2}\right) - \lambda_{HX}|X|^{2} \left(|\mathcal{H}|^{2} - \frac{1}{2}v_{H}^{2}\right) - \lambda_{\Phi H} \left(|\Phi|^{2} - \frac{1}{2}v_{\Phi}^{2}\right) \left(|\mathcal{H}|^{2} - \frac{1}{2}v_{H}^{2}\right) + \cdots,$$
(7)

where $\mathcal{D}_{\rho} = \partial_{\rho} + ig_X \widehat{\mathcal{Q}}_{{\sf L}_{\mu}-{\sf L}_{\tau}} Z'_{\rho}$ is the covariant derivative with g_X denoting the dark gauge coupling and Z'being the dark gauge boson with mass $m_{Z'} = g_X |\mathcal{Q}_{\Phi}| v_{\Phi}$, m_X is the mass of DM, and $\mathcal{H} = \frac{1}{\sqrt{2}} (0, v_H + h)^{\mathsf{T}}$ is the SM Higgs doublet (in the unitary gauge) with the VEV $v_H \simeq 246.22 \,\text{GeV}$. From now on, we call Z' as dark photon because it couples to the DM. Note that the kinetic mixing between Z' and the usual photon arises through $\ell = \mu, \tau \text{ loops}; -\epsilon e \bar{\ell} \gamma^{\rho} \ell Z'_{\rho} \text{ with } \epsilon \simeq -g_X/70 \text{ [30]. In}$ this Lagrangian density, the DM relic abundance is determined by the $\lambda_{\Phi X}, \lambda_{HX}$, and g_X couplings. However, we will take $\lambda_{HX} = 0$ for simplicity. The $\lambda_{\Phi H}$ coupling allows the CP-even neutral components of Φ and \mathcal{H} , ϕ and h, respectively, to mix after electroweak and $U(1)_{L_{\mu}-L_{\tau}}$ symmetry breakings. The dark Higgs (SM-like Higgs) boson in the mass eigenstate is denoted as $H_1(H_2)$, where $H_1 = \phi \cos \theta - h \sin \theta$ and $H_2 = \phi \sin \theta + h \cos \theta$ with θ being the mixing angle. In this work, we will assume that the mass of H_1 is smaller than that of H_2 , $m_{H_1} <$ $m_{H_2} \simeq 125 \,\text{GeV}$. The parameter set for our numerical calculations is then

$$\left\{m_{Z'}, m_X, m_{H_1}, \lambda_{\Phi X}, g_X, \mathcal{Q}_{\Phi}, \theta\right\}$$

Note that the Δa_{μ} is only determined by g_X and $m_{Z'}$, whereas the $B^+ \rightarrow K^+ \not\!\!\! E$ depends on all these parameters. We will discuss in more details in the following sections.

MUON g - 2 & HUBBLE TENSION

In the U(1)_{$L_{\mu}-L_{\tau}$} model, the massive dark photon Z' provides an additional contribution to the muon magnetic moment via the vertex correction [31, 32]:

$$\Delta a_{\mu} = \frac{g_X^2}{4\pi^2} \int_0^1 \mathrm{d}x \, \frac{x^2(1-x)}{x^2 + (1-x)m_{Z'}^2/m_{\mu}^2} \,. \tag{8}$$

Taking $g_X \sim 10^{-4}$ and $m_{Z'} < m_{\mu}$, one can alleviate the discrepancy in Δa_{μ} . The MeV-scale Z' can produce substantial entropy and energy via the $Z' \rightarrow \nu \bar{\nu}$ decay process, which would spoil the successful predictions of the Big Bang Nucleosynthesis. We impose $\Delta N_{\rm eff} < 3.5$ as an exclusion bound. On the other hand, there exists an inconsistency between the Hubble constant observed today from the Cosmic Microwave Background (CMB) and the value measured from the celestial sources [30, 33]. To relieve this tension, $\Delta N_{\rm eff} > 0.2$ is preferred [30]. Taking into account the constraints from BOREXINO [34–36], CCFR [37, 38], and NA64 [39], we identify a benchmark point $(m_{Z'}, g_X) = (11.5 \,{\rm MeV}, 5 \times 10^{-4})$ as our numerical inputs, from which $\Delta N_{\rm eff} \simeq 0.3$, thereby relaxing the Hubble tension [30]. Also, $v_{\Phi} \sim m_{Z'}/g_X \sim \mathcal{O}(10) \,{\rm GeV}$.

HIGGS INVISIBLE DECAY

With the dark sector, the SM-like Higgs boson H_2 has additional decay processes : $H_2 \rightarrow H_1H_1, Z'Z'$, and XX^{\dagger} if kinematically open. These decay channels lead to invisible or non-standard decays of the SM-like Higgs boson,




FIG. 1. Feynman graphs for complex scalar DM annihilating into the Z' bosons, H_1 bosons, and SM leptons, where $\ell = \mu, \tau$ and the arrow represents the $L_{\mu} - L_{\tau}$ charge flow.

which is strongly bounded by the LHC data [40]:

$$\mathcal{B}(H_2 \to \text{Inv.}) < 0.13 . \tag{9}$$

It is easy to check that the H_2 mainly decays into H_1 and Z' pairs if $\lambda_{\Phi X} \lesssim \mathcal{O}(1)$, where $\Gamma_{H_2 \to H_1 H_1} \simeq \Gamma_{H_2 \to Z'Z'} \propto \sin^2 \theta \, m_{H_2}^3 / v_{\Phi}^2 \gg \Gamma_{H_2 \to XX^{\dagger}} \propto \sin^2 \theta \, \lambda_{\Phi X}^2 v_{\Phi}^2 / m_{H_2}$. Hence, the constraint from the Higgs invisible decay on the $\sin \theta$ can be insensitive to $\lambda_{\Phi X}$ as long as $\lambda_{\Phi X}$ is small enough. Typically, the $\sin \theta$ should be less than $\sim 10^{-2}$ to satisfy the Higgs invisible decay constraint. Note that the produced dark Higgs from the SM Higgs decay can further decays into the SM fermions, f. However, these decay channels are highly suppressed thanks to small $\sin \theta$ and yukawa couplings, where $\Gamma_{H_1 \to XX^{\dagger}} \propto \lambda_{\Phi X}^2 v_{\Phi}^2 / m_{H_1} \gg \Gamma_{H_1 \to Z'Z'} \propto m_{H_1}^3 / v_{\Phi}^2 \gg \Gamma_{H_1 \to f\bar{f}} \propto \sin^2 \theta \, m_f^2 m_{H_1} / v_H^2$ in our interesting parameter space. Thus, all of the additional decay channels of the SM Higgs are invisible. It follows that the decay process $B^0 \to K^{*0} H_1 \to K^{*0} \mu^+ \mu^-$ is also suppressed [41].

DARK MATTER

Let us now discuss the production of thermal freezeout DM. The evolution of the DM density is governed by the Boltzmann equation. To get the right amount of DM relic density, it is required that the total DM annihilation cross section $\langle \sigma v \rangle$ for symmetric DM, $\Omega_X = \Omega_{X^{\dagger}}$, satisfying [42]

$$\Omega_{\rm DM} \hat{h}^2 = 2\Omega_X \simeq \frac{1.75 \times 10^{-10} {\rm GeV}^{-2} x_f}{\sqrt{g_*} \langle \sigma v \rangle} , \qquad (10)$$

where $g_* \simeq 10$ and $x_f = m_X/T_f \sim 12 - 19$ for DM in the MeV-GeV range. In U(1)_{Lµ-Lτ}-charged DM models without a dark Higgs boson, the $\langle \sigma v \rangle$ only depends on $m_{Z'}, g_X$, and m_X . To explain the $\Delta a_{\mu}, m_{Z'}$ and g_X are fixed. Also, since the dominant DM annihilation process in these models is $XX^{\dagger} \rightarrow \nu \bar{\nu}$ via s-channel Z' exchange (see the bottom-right diagram in Fig. 1), the correct DM relic abundance can only be produced when $m_{Z'} \sim 2m_X$ [19]. However, the authors in Ref. [21] pointed out that this tight correlation between $m_{Z'}$ and m_X can be completely bypassed if we include a dark Higgs boson, which plays an important role not only in the generation of dark photon mass but also opening new DM annihilation channels and related processes.

We depict in Fig. 1 all of the Feynman diagrams of the DM annihilation processes in this model. The dominant annihilation channels determining the relic abundance of DM are $XX^{\dagger} \rightarrow Z'Z', H_1Z', H_1H_1$, where the latter two processes are important only when $m_X \gtrsim m_{H_1}$. The processes $XX^{\dagger} \rightarrow \ell^+\ell^-, \nu_\ell\bar{\nu}_\ell$ are suppressed by $\sin\theta$ and g_X for $\ell = \mu, \tau$, respectively, and by an additional kinetic mixing for $\ell = e$. Also, the $XX^{\dagger} \rightarrow Z'Z'$ process is governed by the s-channel H_1 exchange diagram (see the top-left diagram in Fig. 1) because of the smallness of g_X^4 in the other $XX^{\dagger} \rightarrow Z'Z'$ diagrams. Thus, the total DM annihilation cross section in the $\sin\theta \ll 1$ and $\cos\theta \simeq 1$ limit (adopting hereafter) is approximately given by

$$\langle \sigma v \rangle \simeq \frac{\lambda_{\Phi X}^2}{16\pi m_X^2} \frac{4m_X^4 - 4m_X^2 m_{Z'}^2 + 3m_{Z'}^4}{(m_{H_1}^2 - 4m_X^2)^2 + \Gamma_{H_1}^2 m_{H_1}^2} \sqrt{1 - \frac{m_{Z'}^2}{m_X^2}} ,$$
(11)

where the decay width of H_1 is described as

$$\Gamma_{H_1} \simeq \frac{\lambda_{\Phi X}^2 v_{\Phi}^2}{16\pi m_{H_1}} \sqrt{1 - \frac{4m_X^2}{m_{H_1}^2}} \ . \tag{12}$$

Notice that the dark Higgs can also decay into a pair of either Z' or the SM charged particle. However, again due to the small values of $\sin \theta$ and g_X , these decay channels are negligible.

In U(1)_{Lµ-Lτ} DM models without a dark Higgs boson, DM has to be light, $m_X \sim m_{Z'}/2 \sim \mathcal{O}(10)$ MeV. The elastic scattering cross section of such light DM with the proton (electron) through the Z' mediator is $\sigma_{\rm el}^{X-p} \sim$ $10^{-46} {\rm cm}^2 (\sigma_{\rm el}^{X-e} \sim 10^{-51} {\rm cm}^2)$. Hence, the ongoing direct detection experiments are not sensitive to theoretical expectations [19]. However, in our model, we have a DMnucleon scattering process induced by the SM and dark Higgs boson exchanges. The corresponding elastic scattering cross section is given by

$$\sigma_{\rm el} \simeq \frac{4\mu_n^2 f_n^2 \lambda_{\Phi X}^2 \sin^2 \theta}{\pi} \left(\frac{m_n}{m_X}\right)^2 \left(\frac{\upsilon_{\Phi}}{\upsilon_H}\right)^2 \left(\frac{1}{m_{H_1}^2} - \frac{1}{m_{H_2}^2}\right)^2,\tag{13}$$

where $f_n = 0.326$ [43, 44], m_n is nucleon mass and $\mu_n = m_X m_n / (m_X + m_n)$ is the reduced mass of nucleon.

When $m_X \lesssim \mathcal{O}(10)$ GeV, the CMB provides the most stringent bounds [45]. To avoid the CMB constraints on the low DM mass region, asymmetric DM [46], *p*-wave annihilation [42], or forbidden channel [47] have been proposed. In our DM model, the dominant DM annihilation channels are $XX^{\dagger} \rightarrow Z'Z', H_1H_1$. These channels are s-wave annihilation. However, the dark Higgs boson eventually decays into a DM pair, $H_1 \rightarrow XX^{\dagger}$, and the dark photon would decay into the active neutrinos, $Z' \rightarrow \nu \bar{\nu}$ since $m_{Z'} < m_{\mu}$. Hence, we can naturally escape the CMB bound. Considering the modification of $N_{\rm eff}$ via light DM s-wave annihilation to neutrinos, complex scalar DM mass below 8.2 MeV is disfavored [48]. In our study, we concentrate on $m_X \geq 20$ MeV.

TWO- OR THREE-BODY DECAYS AT BELLE II

In Ref. [49], they first proposed a bound on DM mass and its coupling using the $B^+ \to K^+ \nu \bar{\nu}$ decay channel in BaBar. They considered a real singlet scalar DM with the SM Higgs portal. However, this simple DM model is already ruled out by the CMB constraints since the relic density of DM is determined by the *s*-wave annihilation $SS \to H_2^{(*)} \to f\bar{f}$, where *S* is real singlet scalar DM.

In comparison with Ref. [49], owing to the dark Higgs boson, the B^+ meson can go through a two-body decay, $B^+ \to K^+ H_1$ when $m_{B^+} - m_{K^+} > m_{H_1}$ in our model. The corresponding two-body decay rate is

$$\Gamma_{B^+ \to K^+ H_1} \simeq \frac{|\kappa_{cb}|^2 \sin^2 \theta}{64\pi m_{B^+}^3} \left[f_0(m_{H_1}^2) \right]^2 \left(\frac{m_{B^+}^2 - m_{K^+}^2}{m_b - m_s} \right)^2 \\ \times \sqrt{\mathcal{K}(m_{B^+}^2, m_{K^+}^2, m_{H_1}^2)} , \qquad (14)$$

where $\kappa_{cb} \simeq 6.7 \times 10^{-6}$ is the one-loop vertex correction after integrating out the top quark and W boson, $f_0(q^2)$ with $q^2 = (p_B - p_K)^2$ is the $B^- \to K^-$ transition form factor which can be found in Ref. [50], and $\mathcal{K}(a, b, c) \equiv a^2 + b^2 + c^2 - 2(ab + bc + ac)$.

Similar to Ref. [49], when the dark Higgs becomes offshell, where $m_{H_1} > m_{B^+} - m_{K^+} > 2m_X$, the B^+ meson can have a three-body decay, $B^+ \to K^+ H_1^{(*)} \to K^+ X X^{\dagger}$. The corresponding three-body decay rate is

$$\Gamma_{B^+ \to K^+ X X^{\dagger}} \simeq \frac{\lambda_{\Phi X}^2 v_{\Phi}^2 |\kappa_{cb}|^2 \sin^2 \theta}{1024 \pi^3 m_{B^+}^3} \left(\frac{m_{B^+}^2 - m_{K^+}^2}{m_b - m_s}\right)^2 \\ \times \int dq^2 \frac{\mathcal{I}(q^2) [f_0(q^2)]^2 (m_{H_1}^2 - m_{H_2}^2)^2}{(q^2 - m_{H_1}^2)^2 (q^2 - m_{H_2}^2)^2} ,$$
(15)

where $4m_X^2 \le q^2 \le (m_{B^+} - m_{K^+})^2$, and

$$\mathcal{I}(q^2) = \sqrt{1 - \frac{4m_X^2}{q^2}} \sqrt{\mathcal{K}(m_{B^+}^2, m_{K^+}^2, q^2)} .$$
(16)

Note that the $B^+ \to K^+ Z' Z'$ decay is also open, but its contribution is negligible as $\Gamma_{B^+ \to K^+ Z' Z'} / \Gamma_{B^+ \to K^+ X X^{\dagger}} \propto m_B^4 / (\lambda_{\Phi X}^2 v_{\Phi}^4) \lesssim 7 \times 10^{-3}$, where $\lambda_{\Phi X} \gtrsim \mathcal{O}(0.1)$ to explain the observed DM relic density and $B^+ \to K^+ \nu \bar{\nu}$ excess (see the bottom panel of Fig. 2).




FIG. 2. (Top) Allowed parameter space of $(\sin \theta, m_{H_1})$ plane. The region inside green (yellow) is allowed at $1\sigma(2\sigma)$ C.L. by Belle II $B^+ \to K^+H_1$, where H_1 decays to either DM or Z'. The gray area is excluded by various experiments from Belle $B^0 \to K^{*0}\nu\bar{\nu}$, and $K^+ \to \pi^+$ inv.. (Bottom) The preferred parameter space in the $(m_{H_1}, \lambda_{\Phi X})$ plane. The region inside green (yellow) is allowed at $1\sigma(2\sigma)$ C.L. by Belle II $B^+ \to K^+XX^{\dagger}$. The orange area is ruled out by the SM Higgs invisible decay at LHC [40]. The solid lines correspond to the thermal freeze-out DM relic density which is consistent with the Planck observation, $\Omega_{\rm DM}\hat{h}^2 = 0.12$ [51]. The dashed lines are disfavored by the DM direct detection bound from XENONnT [52]. The gray region is excluded by $B^0 \to K^{*0}\nu\bar{\nu}$ bound.

NUMERICAL RESULTS

We present in the top panel of Fig. 2 the mixing angle versus the dark Higgs mass with our benchmark point, where the gray region is ruled out by Belle II $B^0 \to K^{*0}\nu\bar{\nu}$ channel [53], KOTO [54], and $K^+ \to \pi^+ +$ invisibles from NA62 [55] and the red-shaded region is excluded by the Higgs invisible decay constraint [40]. The $B^+ \to K^+\nu\bar{\nu}$ excess at Belle II is explained by the twobody decay channel $B^+ \to K^+H_1$ in the green and yellow bands. According to Ref. [9], 2 GeV new light particle is preferred because of q^2 spectrum information provided by Belle II. The allowed $\sin \theta$ lies between 2×10^{-3} and 7×10^{-3} . We will take $\sin \theta = 6 \times 10^{-3}$ as our input below.

In the bottom panel of Fig. 2, we show the parameter region of $\lambda_{\Phi X}$ varying with m_{H_1} , where the color curves with different DM masses satisfy the relic abundance of DM. The orange- and cyan-shaded regions are excluded by the Higgs invisible decay constraint and perturbativity, respectively. On the left side of the dash-dotted line, where $m_{H_1} = m_{B^+} - m_{K^+}$, the $B^+ \to K^+ \nu \bar{\nu}$ excess can be explained by the $B^+ \to K^+ H_1$ when $m_{H_1} \gtrsim 0.4 \,\text{GeV}$ based on the top panel of Fig. 2. Notice that $m_X \gtrsim 10$ GeV is not allowed due to the DM direct detections. On the right side of the dash-dotted line, the $B^+ \to K^+ \nu \bar{\nu}$ excess is explained within the green and yellow regions by the three-body decay channel $B^+ \to K^+ X X^{\dagger}$. The bounds from $\mathcal{B}(B^0 \to K^{*0} \nu \bar{\nu}) < 1.8 \times 10^{-5}$ (gray region) [53] partially excludes the preferred region for Belle II excess. Note that these regions are only valid for the light DM masses, where $m_X \ll m_{B^+}$.

The behavior of the color curves in the low- and high- H_1 mass regions is simple to understand. This is because $\langle \sigma v \rangle \propto \lambda_{\Phi X}^2/m_X^2 \left(\lambda_{\Phi X}^2 m_X^2/m_{H_1}^4 \right)$ for $m_{H_1} \ll m_X \left(m_{H_1} \gg m_X \right)$ and $\langle \sigma v \rangle$ is nearly a constant. However, these color curves have an interesting behavior around $\lambda_{\Phi X} \sim \mathcal{O}(1)$ and $m_X \sim \mathcal{O}(10)$ MeV. This behavior originates from the H_1 decay width in the denominator of the $XX^{\dagger} \rightarrow Z'Z'$ cross section, where $\Gamma_{H_1}^2 m_{H_1}^2 \propto \lambda_{\Phi X}^4 v_{\Phi}^4$ is dominant over $m_{H_1}^4$ when $\lambda_{\Phi X} v_{\Phi}$ is larger. Thus, there are two positive solutions to get the correct relic abundance of DM. The inflection point is around $m_{H_1} \sim m_X/(\langle \sigma v \rangle^{1/2} v_{\Phi})$. After this inflection point, the H_1 decay width enters the phase space suppression when $m_{H_1} \gtrsim 2m_X$, then the required $\lambda_{\Phi X}$ to fit the DM relic density becomes much larger. Due to this unexpected behavior, the light DM mass between 20 MeV $\lesssim m_X \lesssim 60$ MeV is allowed.

CONCLUSIONS

After the Belle II excess announcement, a number of literature tried without success to explain both the observed DM relic density and the enhancement of Belle II at the same time. That is, fitting the Belle II excess with light DM provides an extremely large relic abundance of DM. To reproduce the correct relic density, either introducing new DM annihilation channels or allowing DM to decay were necessary.

In this letter, we have considered the simplest UVcomplete gauged U(1)_{L_µ-L_τ}-charged complex scalar DM model. We have found that the dark Higgs boson mass $0.4 \text{ GeV} \lesssim m_{H_1} \lesssim 10 \text{ GeV}$ with the upper (lower) bound coming from the $B^+ \to K^+ \nu \bar{\nu}$ excess (KOTO if $\sin \theta \sim 6 \times 10^{-3}$), the complex scalar DM mass $10 \text{ MeV} \lesssim m_X \lesssim$ 10 GeV with the upper (lower) bound coming from the direct detection experiments $(\Delta N_{\rm eff})$, and the dark photon mass $m_{Z'} \sim 10 \,{\rm MeV}$ with $g_X \sim 5 \times 10^{-4}$ to explain the muon g - 2. Attributing to these light dark particles in this model, we can achieve the integrated solution of $B^+ \to K^+ \nu \bar{\nu}$ at Belle II, Δa_{μ} , thermal DM relic density, and Hubble tension. Our analysis makes another case where the dark Higgs boson plays a crucial role in DM phenomenology.

ACKNOWLEDGMENTS

We are grateful to Josef Pradler, Michael Schmidt, Satoshi Shirai, and Fanrong Xu for useful discussions. This work is supported by KIAS Individual Grants under Grants No. PG081202 (SYH), PG021403 (PK), and by National Research Foundation of Korea (NRF) Research Grant NRF-2019R1A2C3005009 (JK, PK).

- * phyhunter@kias.re.kr
- [†] jkkim@kias.re.kr
- [‡] pko@kias.re.kr
- [1] I. Adachi *et al.* [Belle-II], "Evidence for $B^+ \to K^+ \nu \bar{\nu}$ Decays," [arXiv:2311.14647 [hep-ex]].
- W. G. Parrott *et al.* [HPQCD], Phys. Rev. D 107, no.1, 014511 (2023) [erratum: Phys. Rev. D 107, no.11, 119903 (2023)] doi:10.1103/PhysRevD.107.014511 [arXiv:2207.13371 [hep-ph]].
- [3] P. Athron, R. Martinez and C. Sierra, "B meson anomalies and large $B^+ \to K^+ \nu \bar{\nu}$ in non-universal U(1)' models," [arXiv:2308.13426 [hep-ph]].
- [4] R. Bause, H. Gisbert and G. Hiller, Phys. Rev. D 109, no.1, 015006 (2024) [arXiv:2309.00075 [hep-ph]].
- [5] L. Allwicher, D. Becirevic, G. Piazza, S. Rosauro-Alcaraz and O. Sumensari, Phys. Lett. B 848, 138411 (2024) [arXiv:2309.02246 [hep-ph]].
- [6] X. G. He, X. D. Ma and G. Valencia, "Revisiting models that enhance $B^+ \to K^+ \nu \bar{\nu}$ in light of the new Belle II measurement," [arXiv:2309.12741 [hep-ph]].
- [7] A. Berezhnoy and D. Melikhov, " $B \to K^* M_X$ vs $B \to K M_X$ as a probe of a scalar-mediator dark matter scenario," [arXiv:2309.17191 [hep-ph]].
- [8] A. Datta, D. Marfatia and L. Mukherjee, "B → Kνν̄, MiniBooNE and muon g − 2 anomalies from a dark sector," [arXiv:2310.15136 [hep-ph]].
- [9] W. Altmannshofer, A. Crivellin, H. Haigh, G. Inguglia and J. Martin Camalich, "Light New Physics in $B \rightarrow K^{(*)}\nu\bar{\nu}$," [arXiv:2311.14629 [hep-ph]].
- [10] D. McKeen, J. N. Ng and D. Tuckler, "Higgs Portal Interpretation of the Belle II $B^+ \to K^+ \nu \nu$ Measurement," [arXiv:2312.00982 [hep-ph]].
- [11] K. Fridell, M. Ghosh, T. Okui and K. Tobioka, "Decoding the $B \to K\nu\nu$ excess at Belle II: kinematics, operators, and masses," [arXiv:2312.12507 [hep-ph]].
- [12] K. Cheung, Y. Kim, Y. Kwon, C. J. Ouseph, A. Soffer and Z. S. Wang, "Probing dark photons from a light scalar at Belle II," [arXiv:2401.03168 [hep-ph]].

- [13] T. Felkl, A. Giri, R. Mohanta and M. A. Schmidt, "When energy goes missing: new physics in $b \rightarrow s\nu\nu$ with sterile neutrinos," Eur. Phys. J. C 83, no.12, 1135 (2023) [arXiv:2309.02940 [hep-ph]].
- [14] H. K. Dreiner, J. Y. Günther and Z. S. Wang, "The Decay $B \to K \nu \bar{\nu}$ at Belle II and a Massless Bino in R-parity-violating Supersymmetry," [arXiv:2309.03727 [hep-ph]].
- [15] D. P. Aguillard *et al.* [Muon g-2], "Measurement of the Positive Muon Anomalous Magnetic Moment to 0.20 ppm," Phys. Rev. Lett. **131**, no.16, 161802 (2023) [arXiv:2308.06230 [hep-ex]].
- [16] T. Aoyama, N. Asmussen, M. Benayoun, J. Bijnens, T. Blum, M. Bruno, I. Caprini, C. M. Carloni Calame, M. Cè and G. Colangelo, *et al.* "The anomalous magnetic moment of the muon in the Standard Model," Phys. Rept. 887, 1-166 (2020) [arXiv:2006.04822 [hep-ph]].
- [17] S. Borsanyi, Z. Fodor, J. N. Guenther, C. Hoelbling, S. D. Katz, L. Lellouch, T. Lippert, K. Miura, L. Parato and K. K. Szabo, *et al.* "Leading hadronic contribution to the muon magnetic moment from lattice QCD," Nature 593, no.7857, 51-55 (2021) [arXiv:2002.12347 [hep-lat]].
- [18] P. Foldenauer, "Light dark matter in a gauged $U(1)_{L_{\mu}-L_{\tau}}$ model," Phys. Rev. D **99**, no.3, 035007 (2019) [arXiv:1808.03647 [hep-ph]].
- [19] I. Holst, D. Hooper and G. Krnjaic, Phys. Rev. Lett. 128, no.14, 141802 (2022) [arXiv:2107.09067 [hep-ph]].
- [20] M. Drees and W. Zhao, Phys. Lett. B 827, 136948 (2022) [arXiv:2107.14528 [hep-ph]].
- [21] S. Baek, J. Kim and P. Ko, "Muon (g-2) and Thermal WIMP DM in U(1)_{$L_{\mu}-L_{\tau}$} Models," [arXiv:2204.04889 [hep-ph]].
- [22] P. Ko, J. Korean Phys. Soc. 73, no.4, 449-465 (2018) doi:10.3938/jkps.73.449
- [23] X. G. He, G. C. Joshi, H. Lew and R. R. Volkas, "NEW Z-prime PHENOMENOLOGY," Phys. Rev. D 43, 22-24 (1991)
- [24] X. G. He, G. C. Joshi, H. Lew and R. R. Volkas, "Simplest Z-prime model," Phys. Rev. D 44, 2118-2132 (1991)
- [25] V. Silveira and A. Zee, "SCALAR PHANTOMS," Phys. Lett. B 161, 136-140 (1985)
- [26] X. G. He, T. Li, X. Q. Li, J. Tandean and H. C. Tsai, "The Simplest Dark-Matter Model, CDMS II Results, and Higgs Detection at LHC," Phys. Lett. B 688, 332-336 (2010) [arXiv:0912.4722 [hep-ph]].
- [27] X. G. He, S. Y. Ho, J. Tandean and H. C. Tsai, "Scalar Dark Matter and Standard Model with Four Generations," Phys. Rev. D 82, 035016 (2010) [arXiv:1004.3464 [hep-ph]].
- [28] S. Baek, P. Ko and W. I. Park, "Singlet Portal Extensions of the Standard Seesaw Models to a Dark Sector with Local Dark Symmetry," JHEP 07, 013 (2013) [arXiv:1303.4280 [hep-ph]].
- [29] P. Ko, C. T. Lu and U. Min, "Crossing two-component dark matter models and implications for 511 keV γ-ray and XENON1T excesses," [arXiv:2202.12648 [hep-ph]].
- [30] M. Escudero, D. Hooper, G. Krnjaic and M. Pierre, "Cosmology with A Very Light L_μ – L_τ Gauge Boson," JHEP 03, 071 (2019) [arXiv:1901.02010 [hep-ph]].
- [31] S. Baek, N. G. Deshpande, X. G. He and P. Ko, "Muon anomalous g-2 and gauged L(muon) - L(tau) models," Phys. Rev. D 64, 055006 (2001) [arXiv:hep-ph/0104141 [hep-ph]].

- [32] S. Baek and P. Ko, "Phenomenology of U(1)(L(mu)-L(tau)) charged dark matter at PAMELA and colliders," JCAP 10, 011 (2009) [arXiv:0811.1646 [hep-ph]].
- [33] E. Di Valentino, O. Mena, S. Pan, L. Visinelli, W. Yang, A. Melchiorri, D. F. Mota, A. G. Riess and J. Silk, "In the realm of the Hubble tension—a review of solutions," Class. Quant. Grav. 38, no.15, 153001 (2021) [arXiv:2103.01183 [astro-ph.CO]].
- [34] A. Kamada and H. B. Yu, "Coherent Propagation of PeV Neutrinos and the Dip in the Neutrino Spectrum at IceCube," Phys. Rev. D 92, no.11, 113004 (2015) [arXiv:1504.00711 [hep-ph]].
- [35] Y. Kaneta and T. Shimomura, "On the possibility of a search for the $L_{\mu} - L_{\tau}$ gauge boson at Belle-II and neutrino beam experiments," PTEP **2017**, no.5, 053B04 (2017) [arXiv:1701.00156 [hep-ph]].
- [36] S. Gninenko and D. Gorbunov, "Refining constraints from Borexino measurements on a light Z'-boson coupled to $L\mu$ - $L\tau$ current," Phys. Lett. B **823**, 136739 (2021) [arXiv:2007.16098 [hep-ph]].
- [37] S. R. Mishra *et al.* [CCFR], "Neutrino Tridents and W Z Interference," Phys. Rev. Lett. **66**, 3117-3120 (1991)
- [38] W. Altmannshofer, S. Gori, M. Pospelov and I. Yavin, "Neutrino Trident Production: A Powerful Probe of New Physics with Neutrino Beams," Phys. Rev. Lett. 113, 091801 (2014) [arXiv:1406.2332 [hep-ph]].
- [39] Y. M. Andreev, D. Banerjee, B. B. Oberhauser, J. Bernhard, P. Bisio, N. Charitonidis, P. Crivelli, E. Depero, A. V. Dermenev and S. V. Donskov, *et al.* "Exploration of the Muon g 2 and Light Dark Matter explanations in NA64 with the CERN SPS high energy muon beam," [arXiv:2401.01708 [hep-ex]].
- [40] R. L. Workman *et al.* [Particle Data Group], "Review of Particle Physics," PTEP **2022**, 083C01 (2022)
- [41] M. Ovchynnikov, M. A. Schmidt and T. Schwetz, "Complementarity of $B \to K^{(*)} \mu \bar{\mu}$ and $B \to K^{(*)} + \text{inv}$ for searches of GeV-scale Higgs-like scalars," Eur. Phys. J. C 83, no.9, 791 (2023) [arXiv:2306.09508 [hep-ph]].
- [42] K. Griest and D. Seckel, "Three exceptions in the calculation of relic abundances," Phys. Rev. D 43, 3191-3203 (1991)
- [43] R. D. Young and A. W. Thomas, "Octet baryon masses and sigma terms from an SU(3) chiral extrapolation," Phys. Rev. D 81, 014503 (2010) [arXiv:0901.3310 [heplat]].
- [44] A. Crivellin, M. Hoferichter and M. Procura, "Accurate evaluation of hadronic uncertainties in spinindependent WIMP-nucleon scattering: Disentangling two- and three-flavor effects," Phys. Rev. D 89, 054021 (2014) [arXiv:1312.4951 [hep-ph]].
- [45] T. R. Slatyer, "Indirect dark matter signatures in the cosmic dark ages. I. Generalizing the bound on s-wave dark matter annihilation from Planck results," Phys. Rev. D 93, no.2, 023527 (2016) [arXiv:1506.03811 [hep-ph]].
- [46] D. E. Kaplan, M. A. Luty and K. M. Zurek, "Asymmetric Dark Matter," Phys. Rev. D 79, 115016 (2009) [arXiv:0901.4117 [hep-ph]].
- [47] R. T. D'Agnolo and J. T. Ruderman, "Light Dark Matter from Forbidden Channels," Phys. Rev. Lett. 115, no.6, 061301 (2015) [arXiv:1505.07107 [hep-ph]].
- [48] X. Chu and J. Pradler, "On the minimal mass of thermal dark matter and the viability of millicharged particles affecting 21cm cosmology," [arXiv:2310.06611 [hep-ph]].

- [49] C. Bird, P. Jackson, R. V. Kowalewski and M. Pospelov, "Search for dark matter in b —> s transitions with missing energy," Phys. Rev. Lett. 93, 201803 (2004) [arXiv:hep-ph/0401195 [hep-ph]].
- [50] W. G. Parrott *et al.* [(HPQCD collaboration)§ and HPQCD], "B→K and D→K form factors from fully relativistic lattice QCD," Phys. Rev. D **107**, no.1, 014510 (2023) [arXiv:2207.12468 [hep-lat]].
- [51] N. Aghanim *et al.* [Planck], "Planck 2018 results. VI. Cosmological parameters," Astron. Astrophys. **641**, A6 (2020) [erratum: Astron. Astrophys. **652**, C4 (2021)] [arXiv:1807.06209 [astro-ph.CO]].
- [52] E. Aprile *et al.* [XENON], "First Dark Matter Search with Nuclear Recoils from the XENONnT Experiment," Phys. Rev. Lett. **131**, no.4, 041003 (2023) [arXiv:2303.14729 [hep-ex]].
- [53] J. Grygier *et al.* [Belle], "Search for $B \to h\nu\bar{\nu}$ decays with semileptonic tagging at Belle," Phys. Rev. D **96**, no.9, 091101 (2017) [arXiv:1702.03224 [hep-ex]].
- [54] J. K. Ahn *et al.* [KOTO], "Study of the $K_L \rightarrow \pi^0 \nu \bar{\nu}$ Decay at the J-PARC KOTO Experiment," Phys. Rev. Lett. **126**, no.12, 121801 (2021) [arXiv:2012.07571 [hepex]].
- [55] E. Cortina Gil *et al.* [NA62], JHEP **06**, 093 (2021) [arXiv:2103.15389 [hep-ex]].