Comments on "Can quantum statistics help distinguish Dirac from Majorana neutrinos?" (arXiv 2402.05172 [hep-ph])

C. S. Kim*1, M. V. N. Murthy^{†2}, and Dibyakrupa Sahoo^{‡3}

¹Department of Physics and IPAP, Yonsei University, Seoul 03722, Korea
²The Institute of Mathematical Sciences (Retired), Taramani, Chennai 600113, India
³Institute of Theoretical Physics, Faculty of Physics, University of Warsaw,
ul. Pasteura 5, 02-093 Warsaw, Poland

March 8, 2024

Abstract

In a recent article [1], the authors discuss the question "whether quantum statistics can help distinguish between Dirac and Majorana neutrinos." The paper contains, among other things, an unsubstantiated critique of the results derived in our papers [2] and [3]. One of the criticisms is related to results obtained using back-to-back kinematics in the B^0 meson differential decay rate. We show that the claim is wrong and point out how the correct result was obtained. The second criticism is related to the implementation of the anti-symmetrization as dictated by quantum statistics for Majorana neutrinos and antineutrinos (which are identical, by definition). Any direct observation of the neutrinos, as done in Ref. [1], would project the neutrinos into distinguishable helicity states, thus nullifying all observable effects of quantum statistics. They have missed the point that our procedure holds when the neutrino-antineutrino remain undetected by the detector. In the back-to-back kinematic configuration, one can infer the neutrino energies without directly detecting their identities. This smartly ensures that the quantum statistical effects are not erased. Their overriding assertion that our papers [2] and [3] are incorrect fails to recognize that in both Ref. [2] and [3] we also point out generic conditions under which the practical Dirac-Majorana confusion theorem holds. "Clearly there is no confusion over confusion theorem."

^{*}cskim@yonsei.ac.kr

[†]murthymundur@gmail.com

[‡]Dibyakrupa.Sahoo@fuw.edu.pl

1 Rebuttal of comments made in Sec. 3.1 of Ref. [1]

Below we present our arguments highlighting our disagreement with the incorrect statements in Ref. [1] in their Sec.3.1. Their criticism is primarily centered around Eqs. (48a) and (48b) of our paper [2] which give the differential decay rate for the decay $B^0 \to \mu^- \mu^+ \nu_\mu \overline{\nu}_\mu$ when the neutrino and antineutrino are back-to-back in the rest frame of the parent meson B^0 . Our work basically aims at using the quantum statistics of identical particles to distinguish between Dirac or Majorana nature of neutrinos in general and using the above decay process in particular.

1.1 The misunderstanding!

The authors of Ref. [1] have clearly not followed our step-by-step analysis of the back-to-back case which is correctly and unambiguously described in Sec.IV.H of our paper [2]. They simply state in Sec. 3.1 of their paper [1] that "the denominators of the left-hand sides of Eqs. (48a) and (48b)" in our paper [2] "erroneously contain $d \sin \theta$ instead of $d \cos \theta$ " without giving details but in a hand waving manner. As explained in the next subsection, the phase space is fixed by the final states of the process chosen. However, the variables chosen to describe the differential decay rate are a matter of choice once the kinematic frame is defined.¹

The authors of Ref. [1] correctly state that the back-to-back kinematics is "fully characterized by just two physical variables – the muon energy E_{μ} and the angle θ between the muon and neutrino directions." However, the article suggests that "since the neutrinos are not observed, one must integrate over the solid angle of the neutrino direction, which involves the integration over $d\cos\theta$, not over $d\sin\theta$." If we start from the general kinematics and correctly follow the change of variables required to describe the back-to-back kinematics, we do find that one gets $d\sin\theta$ instead of $d\cos\theta$. This important result is explained in detail in our paper [2]. We refer the reader to see Sec.IV.F–H of our Ref. [2] in order to get the correct understanding of the origin of $d\sin\theta$ as deduced from general kinematics to the back-to-back kinematics.

1.2 Why a simple approach does not yield the correct decay rate

The authors of [1] suggest that "One can consider the problem in the back-to-back kinematics from the outset, without any limiting procedures." However, such a simple approach of imposing the kinematics constraints artificially does not lead to the decay rate but to some unphysical quantity. The correct approach is to start with general formulation of the phase space and reduce it to the desired kinematic situation as shown in in Sec.IV.F–H of our paper [2].

For the decay considered in Ref. [2] $B^0(p_B) \to \mu^-(p_-) \mu^+(p_+) \overline{\nu}_\mu(p_1) \nu_\mu(p_2)$, the decay

¹Changing the variable of integration from $\sin \theta$ to $\cos \theta$ requires change of the integrand as well so that the result after integration remains the same. One can not simply change the variable of integration while keeping the integrand unchanged.

rate (considering all allowed kinematic configurations) is given by,

$$\Gamma^{D/M} = \frac{1}{2 m_B} \int \left| \mathcal{M}^{D/M}(p_+, p_-, p_1, p_2) \right|^2 dLIPS (2\pi)^4 \delta^{(4)} \Big(p_B - p_+ - p_- - p_1 - p_2 \Big), (1)$$

where $\mathcal{M}^{D/M}(p_+, p_-, p_1, p_2)$ is the general amplitude in Dirac (D) or Majorana (M) neutrino case, and the Lorentz invariant phase space element is given by,

$$dLIPS = \left(\frac{d^3\vec{p}_+}{(2\pi)^3 2E_+}\right) \left(\frac{d^3\vec{p}_-}{(2\pi)^3 2E_-}\right) \left(\frac{d^3\vec{p}_1}{(2\pi)^3 2E_1}\right) \left(\frac{d^3\vec{p}_2}{(2\pi)^3 2E_2}\right),\tag{2}$$

where $p_i = (E_i, \vec{p_i})$ with $i \in \{+, -, 1, 2\}$. Since the mass dimension of the amplitude square in our case is -2, Eq. (1) yields the correct mass dimension for the decay rate $\Gamma^{D/M}$,

$$\operatorname{mass dim} \left[\Gamma^{D/M} \right] = 1. \tag{3}$$

Following the suggestion of Ref. [1] to impose the back-to-back kinematic condition (i.e. $\vec{p}_+ + \vec{p}_- = \vec{0} = \vec{p}_1 + \vec{p}_2$) "without any limiting procedures" amounts to multiplying the right hand side of Eq. (1) by the Dirac delta function $\delta^{(3)}$ ($\vec{p}_1 + \vec{p}_2$). Thus, we get some quantity, say $X^{D/M}$, given by

$$X^{D/M} = \frac{1}{2 m_B} \int \left| \mathcal{M}^{D/M}(p_+, p_-, p_1, p_2) \right|^2 dLIPS (2\pi)^4 \delta^{(4)} \left(p_B - p_+ - p_- - p_1 - p_2 \right) \delta^{(3)} (\vec{p}_1 + \vec{p}_2) . \tag{4}$$

It is indeed true that the integration on right hand side is now much simpler and straightforward and it leads to

$$X^{D/M} = \frac{1}{32 (2\pi)^8 m_B} \int \left| \mathcal{M}^{D/M} (\underline{p_+, p_-, p_1, p_2}) \right|^2 \left(\frac{d^3 \vec{p}_+}{\left| \vec{p}_+ \right|^2 + m_\mu^2} \right) \left(\frac{d^3 \vec{p}_1}{\left| \vec{p}_1 \right|^2 + m_\nu^2} \right) \times \delta \left(m_B - 2\sqrt{\left| \vec{p}_+ \right|^2 + m_\mu^2} - 2\sqrt{\left| \vec{p}_1 \right|^2 + m_\nu^2} \right). \tag{5}$$

From usual analysis of solid angles we have, $d^3\vec{p}_+ = (2\pi) E_\mu \sqrt{E_\mu^2 - m_\mu^2} dE_\mu d\cos\theta$, as well as, $d^3\vec{p}_1 = (4\pi) E_\nu \sqrt{E_\nu^2 - m_\nu^2} dE_\nu$. Neglecting m_μ and m_ν dependence it is easy to show that,

$$X^{D/M} \approx \frac{1}{32 (2\pi)^6 m_B} \int \left| \mathcal{M}^{D/M} (\underline{p_+, p_-, p_1, p_2}) \right|^2 dE_\mu d\cos\theta, \tag{6}$$

which indeed has d $\cos \theta$ instead of d $\sin \theta$ but the mass dimension of the quantity $X^{D/M}$ is

$$\operatorname{mass\,dim}\left[X^{D/M}\right] = -2,\tag{7}$$

due to the fact that $\delta^{(3)}$ ($\vec{p}_1 + \vec{p}_2$) is a dimensionful quantity. Therefore $X^{D/M}$ is not the decay rate for the back-to-back kinematic configuration. On the other hand, our expressions Eq. (48a) and (48b) in Ref. [2] have the correct mass dimension 1 for the corresponding decay rate in the back-to-back case. It should be obvious from the outset, since the Dirac delta-function is not a projection operator.

Therefore, any specific choice of kinematics has to follow from the general analysis by a suitable change of variables which appropriately defines the differential decay rate. We follow this step-by-step approach in [2] to get the differential decay rate for back-to-back kinematic configuration. And this automatically leads us to $d \sin \theta$ and not $d \cos \theta$ in Eqs. (48a) and (48b) in Ref. [2]. Further details on the analysis of back-to-back kinematics can also be found in Appendix A of Ref. [3].

1.3 Main conclusion of our papers [2,3]

The authors of Ref. [1] add in their conclusion that the main conclusion of our papers is dependent on the angular distribution with respect to $\sin \theta$. We would like to emphasise that the process $B^0 \to \mu^- \mu^+ \nu_\mu \overline{\nu}_\mu$ and the choice of back-to-back kinematics considered in [2, 3] are meant to serve as illustrative examples of the many possibilities where the effect of quantum statistics may be realised directly. Thus the main conclusion of our papers is that quantum statistical effects can be used to pin down the nature of the neutrino – whether it is Majorana or Dirac type.

In the context of processes allowed in the Standard Model, this follows when

- 1. the direct and exchange terms are non-trivially different² (e.g. see Eqs. (31) and (32) in Ref. [2]), and
- 2. the observable defined in terms of neutrino momenta is accessible even when the neutrinos are not directly observed in the final state.

This is more clearly mentioned in Sec.2.2 and Fig. 1 of [3], as well as in beginning of Sec.III of [2]. We stress that direct observation of the state of neutrinos destroys the effect of quantum statistics since the neutrinos in the final state are projected on to opposite helicity states (and thus no longer remain indistinguishable) by observation. Hence, any inference from quantum statistical probe necessarily requires no direct detection of the final neutrinos, as stated in point 2 above. This is precisely achieved in the back-to-back kinematic configuration where the energy-momentum information is inferred purely by observing the muons leaving the antisymmetrization effects of quantum statistics intact.

The differential decay rate in terms of $\sin \theta$ is the final expression for the back-to-back configuration of the decay $B^0 \to \mu^- \mu^+ \nu_\mu \overline{\nu}_\mu$. This just comes from phase-space analysis alone

²Due to this non-trivial difference between direct and exchange terms in amplitude square, the phase space integration for certain specific kinematic configuration, such as our back-to-back case, need not vanish as well. Finding such kinematic configuration and doing the relevant phase space integration depend on the process under consideration. One can not make a process independent observation at this level contrary to the goal of Ref. [1].

and not from quantum statistical considerations at all. We have clearly mentioned all the steps that lead to the specific θ dependence of our result in Ref. [2].

2 Comments on Sec. 4 and 5 of Ref. [1]

In Sec. 4 of Ref. [1] the authors study the effect of observing the active neutrinos via weak charged current (CC) and neutral current (NC) processes in some neutrino detector (see their Fig. 1b, 1c and Fig. 2). Their study suggests that when neutrinos get detected, the difference between Dirac and Majorana neutrinos vanishes. This is a well-known quantum measurement effect: if neutrino and antineutrino can be distinguished by their detection, it destroys the quantum indistinguishability of Majorana neutrino and antineutrino. To study the effects of quantum statistics, one needs to consider only Fig. 1a, without the quantum measurement effects that come in Fig. 1b, 1c and 2. In Figs. 1b, 1c and 2 of Ref. [1], each of which is not a single Feynman diagram but a combination of independent Feynman diagrams for production and detection processes, the propagating neutrinos are not virtual but real on-shell neutrino mass eigenstates which lead to neutrino flavor oscillations. Such neutrinos even if they have Majorana nature can not be considered as identical as one observes their identity (ν , $\bar{\nu}$ or chirality) and position (left detector and/or right detector which could be a few tens of meters apart). It is very important that one takes into consideration whether effects of measurement would adversely affect the quantum statistical effects one is trying to probe. As is shown in Ref. [3] there is no difference between Dirac and Majorana neutrino possibilities in $Z^{(*)} \rightarrow \nu \overline{\nu}$ decay mode, if one sums over the final spins of neutrinos. If one wants to study spin dependent effects, one would need to detect the neutrinos and that would have the measurement effect leading to no difference between Dirac and Majorana possibilities, as the authors of Ref. [1] have shown.³

This quantum measurement effect is analogous to the well-known fact that the interference pattern in a double slit experiment is destroyed if one could somehow know through which slit the photon has passed. This is clearly highlighted in the last paragraph of Sec. 2.1 of Ref. [3], which says "It should be noted that in this work we discuss processes where the effect of measurements does not destroy the identical nature of Majorana neutrino and antineutrino. This is akin to putting the constraint that in a double-slit experiment, meant to observe the interference of light, no measurement should identify the slit through which the photon has passed." In summary, to correctly study signatures of quantum statistics it is important that no measurements should affect the amplitude for the process, or alter the predictions for the appropriate observable, *i.e.* quantum statistics requires absolutely identical indistinguishable particles.

Therefore, the authors of Ref. [1] in their Sec.4 are mistaken in wrongly suggesting that the quantum statistical differences between Dirac and Majorana natures of active neutrinos could be

³If the neutrino and antineutrino interact with the detector by NC interaction, then their spin states are still undetected and therefore the final spins must be summed over. This, as we know, results in no difference between Dirac and Majorana cases in $Z^{(*)} \rightarrow \nu \overline{\nu}$ decay in the SM, see Ref. [3] and Sec. 4.2 of Ref. [1].

probed by detecting the active neutrinos and antineutrinos. As explained above and in Ref. [3] such measurement effects would alter the predictions for the very observable itself which one is trying to study. Therefore, we disagree with the view of the authors of Ref. [1] that quantum statistics is not a useful tool to distinguish between Dirac and Majorana neutrinos. In our works [2, 3], we smartly used the back-to-back kinematics to deduce the energies of neutrinos, without detecting their identities so as to avoid quantum measurement effect, and correctly probe the quantum statistics.

3 Comment on anti-symmerization requirement for Majoarana neutrino antineutrino pair and the massless limit

When the neutrinos are not detected in the detector and their polarisations are not directly deducible from the process itself, the "chiral" nature of neutrino or antineutrino is irrelevant. In any case massive neutrinos are not chiral eigenstates. The full calculation then involves sum over the spins of the unobserved neutrinos and antineutrinos. This automatically take into account any m_{ν} dependence. This is preciesly what is done in other processes involving identical fermions, Moller scattering for example.

Besides, if one considers neutrino and antineutrino as chiral fermions, then in the massless limit the Majorana condition is mathematically impossible to satisfy (see Appendix B of [3]). Strictly massless neutrino and antineutrino, are Weyl fermions and are distinct from one another for all purposes. Thus a smooth massless limit for Majorana neutrino is a red herring. The effect of quantum statistics, an intrinsic property, can not depend on a dimensional parameter! This is the reason why we consider only non-zero mass for the neutrinos in Refs. [2] and [3].

For more clarifications on the issues (i) one-to-one correspondence between Dirac and Majorana neutrinos in the massless limit and (ii) amplitude anti-symmetrization for the pair of Majorana neutrinos of the same flavor, we urge the reader to see Sec. 5 of Ref. [3].

4 Summary

To summarise, we have pointed out how the step-by-step approach [2] automatically leads us to $d \sin \theta$ and not $d \cos \theta$ for differential decay rate in Eqs. (48a) and (48b) in Ref. [2].

Furthermore, we again reiterate that when the final state neutrino and antineutrino are directly detected there will be no difference between Dirac and Majorana scenarios due to the well known quantum measurement effect. In our case, we are considering a scenario where there is no direct detection of the neutrino and antineutrino. The momenta are gleaned indirectly by a clever use of the back-to-back geometry which does not destroy the identical nature of Majorana neutrino and antineutrino. Hence the effect of quantum statistics remains intact and this is the focus of our study. In the process we have clearly shown the conditions under which the "confusion theorem" may be evaded and bring out the effect of quantum statistics. For

a detailed explanation Sec. 2.2 and Fig. 1 of [3] especially for (A) the requirement that there should be non-trivial difference between the squares of direct and exchange terms, (B) the need to infer the energies or 3-momenta of the neutrinos instead of direct detection of the neutrinos, and (C) when one considers beyond SM interactions.

We conclude by emphasizing that the detection and identification of final neutrinos would erase all quantum statistical differences between Dirac and Majorana neutrinos. However, this does not preclude existence of quantum statistical differences when one is not directly detecting and identifying the neutrinos and antineutrinos. It only emphasizes that study of quantum statistical effects requires careful consideration of only those signatures which can be measured or probed without directly detecting the neutrinos. Therefore, we can not agree with the conclusion of authors of Ref. [1] that "quantum statistics does not lead to any exceptions to the Practical Dirac-Majorana Confusion Theorem." Their "general proof" of the practical Dirac Majorana confusion theorem fails to consider such possibilities and takes into account only the measurement effects. We would like to encourage that quantum statistical probes be further explored carefully avoiding the nuisance of measurement effects.

Note added: The authors of Ref. [1] also cite Ref. [4] to erroneously claim that integration over the angle θ between the muon and neutrino directions in our back-to-back kinematic configuration for $B^0 \to \mu^- \mu^+ \nu_\mu \overline{\nu}_\mu$ would give no difference between Dirac and Majorana neutrinos, whereas in [4] they integrate over the angle ϕ between the two decay planes (planes made by the di-neutrino and the di-muon sub-systems) which leads to no difference between Dirac and Majorana cases. In this context we would like to point out our comments in [5] where we highlight how [4] is considering a different process than ours and how their analysis of back-to-back kinematics is erroneous as they integrate over the angle ϕ between the two planes instead of fixing it to $\phi = 0$.

Acknowledgments

We thank Prof. Evgeny Akhmedov for sharing their manuscript with us before submission and for discussions which brought more clarity to some issues. The work of CSK is supported by NRF of Korea (NRF-2022R1A5A1030700 and NRF-2022R1I1A1A01055643). The work of DS is supported by the Polish National Science Centre under the grant number DEC-2019/35/B/ST2/02008.

References

[1] E. Akhmedov and A. Trautner, "Can quantum statistics help distinguish Dirac from Majorana neutrinos?," arXiv:2402.05172 [hep-ph]

- [2] C. S. Kim, M. V. N. Murthy and D. Sahoo, "Inferring the nature of active neutrinos: Dirac or Majorana?," Phys. Rev. D **105**, no.11, 113006 (2022) doi:10.1103/PhysRevD.105.113006 [arXiv:2106.11785 [hep-ph]].
- [3] C. S. Kim, "Practical Dirac Majorana confusion theorem: issues and applicability," Eur. Phys. J. C **83**, no.10, 972 (2023) doi:10.1140/epjc/s10052-023-12156-9 [arXiv:2307.05654 [hep-ph]].
- [4] J. M. Márquez, D. Portillo-Sánchez and P. Roig, "Dirac-Majorana neutrinos distinction in four-body decays," Phys. Rev. D **109**, no.3, 033005 (2024) doi:10.1103/PhysRevD.109.033005 [arXiv:2305.14140 [hep-ph]].
- [5] C. S. Kim, M. V. N. Murthy and D. Sahoo, "Comments on "On the Dirac-Majorana neutrinos distinction in four-body decays" (arXiv:2305.14140 [hep-ph])," [arXiv:2308.08464 [hep-ph]].