SN 2023zaw: an ultra-stripped, nickel-poor supernova from a low-mass progenitor

KAUSTAV K. DAS,^{1,*} CHRISTOFFER FREMLING,² MANSI M. KASLIWAL,¹ STEVE SCHULZE,³ JESPER SOLLERMAN,⁴ VIRAJ KARAMBELKAR,¹ SAM ROSE,¹ SHREYA ANAND,¹ IGOR ANDREONI,^{5,6,7} MARIE AUBERT,⁸ SEAN J. BRENNAN,⁴ S. BRADLEY CENKO,^{5,9} MICHAEL W. COUGHLIN,¹⁰ B. O'CONNOR,¹¹ KISHALAY DE,¹² JIM FULLER,¹ MATTHEW GRAHAM,¹ ERICA HAMMERSTEIN,⁶ ANNASTASIA HAYNIE,^{13,14} K-RYAN HINDS,¹⁵ IO KLEISER,¹ S. R. KULKARNI,¹ ZEREN LIN,¹ CHANG LIU,^{16,3} ASHISH A. MAHABAL,^{17,18} CHRISTOPHER MARTIN,¹ ADAM A. MILLER,^{16,3} JAMES D. NEILL,¹ DANIEL A. PERLEY,¹⁵ PRISCILA J. PESSI,⁴ NIKOLAUS Z. PRUSINSKI,¹ JOSIAH PURDUM,² VIKRAM RAVI,¹ BEN RUSHOLME,¹ SAMANTHA WU,¹ AVERY WOLD,¹ AND LIN YAN²

¹Cahill Center for Astrophysics, California Institute of Technology, MC 249-17, 1200 E California Boulevard, Pasadena, CA, 91125, USA

²Caltech Optical Observatories, California Institute of Technology, Pasadena, CA 91125, USA

³Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA), Northwestern University, 1800 Sherman Ave,

Evanston, IL 60201, USA

⁴ The Oskar Klein Centre, Department of Astronomy, Stockholm University, AlbaNova, SE-10691 Stockholm, Sweden

⁵ Joint Space-Science Institute, University of Maryland, College Park, MD 20742, USA

⁶Department of Astronomy, University of Maryland, College Park, MD 20742, USA

⁷Astrophysics Science Division, NASA Goddard Space Flight Center, Mail Code 661, Greenbelt, MD 20771, USA

⁸ Université Clermont-Auvergne, CNRS, LPCA, 63000 Clermont-Ferrand, France

⁹Astrophysics Science Division, NASA Goddard Space Flight Center, MC 661, Greenbelt, MD 20771, USA

¹⁰School of Physics and Astronomy, University of Minnesota, Minneapolis, MN 55455, USA

¹¹McWilliams Center for Cosmology and Astrophysics, Department of Physics, Carnegie Mellon University, Pittsburgh, PA 15213, USA

¹²MIT-Kavli Institute for Astrophysics and Space Research 77 Massachusetts Ave. Cambridge, MA 02139, USA

¹³Department of Physics and Astronomy, University of Southern California, Los Angeles, CA 90089, USA,

¹⁴ The Observatories of the Carnegie Institution for Science, 813 Santa Barbara St., Pasadena, CA 91101, USA

¹⁵Astrophysics Research Institute, Liverpool John Moores University, IC2, Liverpool L3 5RF, UK

¹⁶Department of Physics and Astronomy, Northwestern University, 2145 Sheridan Rd, Evanston, IL 60208, USA

¹⁷Division of Physics, Mathematics and Astronomy, California Institute of Technology, Pasadena, CA 91125, USA

¹⁸Center for Data Driven Discovery, California Institute of Technology, Pasadena, CA 91125, USA

ABSTRACT

We present SN 2023zaw - a sub-luminous (M_r = -16.7 mag) and rapidly-evolving supernova $(t_{1/2,r} = 4.9 \text{ days})$, with the lowest nickel mass ($\approx 0.002 \text{ M}_{\odot}$) measured among all stripped-envelope supernovae discovered to date. The photospheric spectra are dominated by broad He I and Ca NIR emission lines with velocities of $\sim 10\ 000 - 12\ 000\ \mathrm{km\ s^{-1}}$. The late-time spectra show prominent narrow He I emission lines at ~ 1000 km s⁻¹, indicative of interaction with He-rich circumstellar material. SN 2023zaw is located in the spiral arm of a star-forming galaxy. We perform radiation-hydrodynamical and analytical modeling of the lightcurve by fitting with a combination of shock-cooling emission and nickel decay. The progenitor has a best-fit envelope mass of $\approx 0.2 \, M_{\odot}$ and an envelope radius of $\approx 50 \text{ R}_{\odot}$. The extremely low nickel mass and low ejecta mass ($\approx 0.5 \text{ M}_{\odot}$) suggest an ultra-stripped SN, which originates from a mass-losing low mass He-star (ZAMS mass < 10 M_{\odot}) in a close binary system. This is a channel to form double neutron star systems, whose merger is detectable with LIGO. SN 2023zaw underscores the existence of a previously undiscovered population of extremely low nickel mass ($< 0.005 \text{ M}_{\odot}$) stripped-envelope supernovae, which can be explored with deep and high-cadence transient surveys.

Keywords: supernovae: general – supernovae: individual (SN 2023zaw, ZTF23absbqun) – stars: massive - stars: mass-loss - stars: neutron

1. INTRODUCTION

Modern wide-field and high-cadence transient surveys such as the Zwicky Transient Facility (ZTF; Bellm et al. 2019; Graham et al. 2019) and the Asteroid Terrestrialimpact Last Alert System (ATLAS; Tonry et al. 2018; Smith et al. 2020) have expanded the discovery space of unusual supernovae (SNe). The class of rapidlyevolving, faint SNe constitutes such a population of peculiar SNe that evolve on the shortest timescales of a few days and have a peak absolute magnitude fainter than -17 mag. There are only a handful of well-studied examples in the literature – SN 2005ek (Drout et al. 2013), SN 2010X (Kasliwal et al. 2010), iPTF14gqr (De et al. 2018a), SN 2018kzr (McBrien et al. 2019), SN 2019dge (Yao et al. 2020), SN 2019wxt (Shivkumar et al. 2023; Agudo et al. 2023), SN 2019bkc (Prentice et al. 2020; Chen et al. 2020), and SN 2022agco (Yan et al. 2023).

Despite advances in understanding the photometric and spectroscopic diversity, their progenitors and powering mechanisms remain unknown. The various theoretical scenarios include ultra-stripped core-collapse SNe of massive stars that lead to binary neutron star systems (Tauris et al. 2013, 2015), non-terminal thermonuclear detonations of a helium shell on the surface of a white dwarf called a '.Ia' SN (Bildsten et al. 2007; Shen et al. 2010), shock-cooling of H-poor stars with an extended envelope (Kleiser & Kasen 2014; Kleiser et al. 2018), fallback in a core-collapse SN (Zhang et al. 2008; Moriya et al. 2010), accretion-induced collapse of a white dwarf (Dessart et al. 2007), electron-capture SN (Moriya & Eldridge 2016), and neutron-star white-dwarf mergers (Margalit & Metzger 2016).

In this paper, we present SN 2023zaw – a Type Ib SN with the lowest nickel mass measured among all stripped-envelope SNe discovered to date. The late-time spectra are dominated by narrow He I emission lines, suggestive of interaction with He-rich circumstellar material (CSM). SN 2023zaw exhibits the most rapid rise and fade time among all stripped-envelope SNe in the literature. Additionally, it is the nearest SN among all fast and faint SNe in the literature. The photometric and spectroscopic properties suggest that it is an ultrastripped SN, originating from a progenitor with an initial mass < 10 M_{\odot}, which likely formed a neutron star.

2. DISCOVERY AND FOLLOW-UP OBSERVATIONS 2.1. Discovery

* E-mail: kdas@astro.caltech.edu

SN 2023zaw (ZTF23absbgun) was discovered with the ZTF camera (Dekany et al. 2020), which is mounted on the 48-inch Samuel Oschin Telescope at the Palomar Observatory. It was first detected at $\alpha = 04^{h}29^{m}20.235^{s}$, $\delta = +70^{\circ}25'37.52''$ (J2000) on December 7, 2023 at 05:34:06 UTC and reported to the Transient Name Server¹ (Sollerman 2023). At the time of discovery, the AB apparent magnitude in the g-band was 19.34 \pm 0.15 mag. The transient stood out due to its faintness and extremely fast evolution (see Figure A1). It was also flagged as a fast transient candidate by the ZT-FReST framework for kilonova and fast transient discovery (Andreoni et al. 2021) and was saved as part of the magnitude-limited (Bright Transient Survey; Fremling et al. 2020) and volume-limited (Census of the Local Universe; De et al. 2020) surveys of ZTF.

SN 2023zaw is located in the UGC 03048 galaxy at a redshift of z = 0.0101 (Springob et al. 2005). It is located on the edge of one of its spiral arms (see Figure 1). The angular separation from the nucleus of the galaxy is 21.44", which corresponds to a physical separation of 4.36 kpc. We correct for the Virgo, Great Attractor, and Shapley supercluster infall (Mould et al. 2000) based on the NASA Extragalactic Database object page (NED)² for UGC 03048. We adopt a Hubble-flow distance of 43.9 \pm 3.1 Mpc, which corresponds to a distance modulus of 33.21 \pm 0.15 mag.

2.2. Optical photometry

We obtained multiple epochs of q, r and i-band photometry with the ZTF camera. The images were reduced using the ZTF image analysis pipeline (Masci et al. 2019). We also perform forced photometry at the location of the transient for the ZTF images in the gribands. We obtained c and o-band photometry from the forced-photometry service of the Asteroid Terrestrialimpact Last Alert System (ATLAS; Tonry et al. 2018; Smith et al. 2020; Shingles et al. 2021). We took regular cadence multiband photometry in the q, r, i and z bands with the Optical Imager (IO:O) at the 2.0 m robotic Liverpool Telescope (Steele et al. 2004). In addition, photometric data was obtained with the Rainbow Camera on the automated Palomar 60-inch telescope (P60; Cenko et al. 2006). LT and P60 images were processed using the FPipe (Fremling et al. 2016) image subtraction pipeline with PanSTARRS (PS1; Chambers et al. 2016) reference images. The Alhambra Faint Object Spectrograph and Camera (ALFOSC) at the 2.5m Nordic Op-

¹ https://www.wis-tns.org/

² https://ned.ipac.caltech.edu/




Figure 1. Left: Pan-STARRS image of the host-galaxy of SN 2023zaw, UGC 03048 in the grizy filters. The location of the transient is indicated by the white cross at the center. Right: Star-formation rate (SFR) map of the environment around SN 2023zaw (see Section 3.7). SN 2023zaw (position marked by the red circle) exploded in a star-forming region of the host galaxy close to regions of more vigorously star-forming regions. The image has a size of 4.1×6.7 kpc. The SFR scale is not corrected for attenuation.




Figure 2. Left: The multiband lightcurve collage of SN 2023zaw. The x-axis shows rest-frame days since the r-band peak (MJD = 60287.7). The photometry data has been corrected for Milky Way ($A_{V,MW} = 0.77$ mag) and host extinction ($A_{V,host} = 1.12$ mag) as described in Section 3.1. The vertical magenta lines represent epochs of spectral observations. Right: The g-band lightcurve of SN 2023zaw compared with other fast-evolving transients in the literature – fast-declining Type I SNe – SN 2010X (Kasliwal et al. 2010), SN 2019bkc (Prentice et al. 2020; Chen et al. 2020), the kilonova AT 2017gfo (Abbott et al. 2017), ultra-stripped SN candidates – SN 2019dge (Yao et al. 2020), iPTF14gqr (De et al. 2018b), SN 2019wxt (Shivkumar et al. 2023; Agudo et al. 2023), SN 2022agco (Yan et al. 2023). We also compare with Type Ibc and Type Ibn SN lightcurve templates from Taddia et al. (2015) and Hosseinzadeh et al. (2017), respectively.

tical Telescope was used to obtain g, r, and i-band photometry. The ALFOSC data was reduced using the PyNOT³ pipeline. The collage of all the multi-band lightcurves is shown in Figure 2. The photometry data is listed in Table C1.

2.3. Optical spectroscopy

We acquired four epochs of spectroscopy with the Spectral Energy Distribution Machine (SEDM; Blagorodnova et al. 2018) on the Palomar 60-inch telescope. The SEDM data was reduced using the pipeline described in Rigault et al. (2019) and Kim et al. (2022). We obtained three epochs of spectroscopy with the

³ https://github.com/jkrogager/PyNOT

Low-Resolution Imaging Spectrometer (LRIS; Oke et al. 1995) on the Keck I telescope, with data reduced using the automated LPIPE (Perley 2019) pipeline. The Alhambra Faint Object Spectrograph and Camera (AL-FOSC) instrument at the 2.5m Nordic Optical Telescope was used to obtain one epoch of low-resolution spectrum, which was reduced using the $PyNOT^4$ pipeline. We took one spectrum with the Binospec spectrograph (Fabricant et al. 2019) on the Multiple Mirror Telescope, which was reduced using the PypeIt package (Prochaska et al. 2020; Prochaska et al. 2020). We used the Keck Cosmic Web Imager (KCWI; Martin et al. 2010; Morrissey et al. 2018a) to obtain spectra of the transient and its host environment. This was reduced using the KCWI data reduction pipeline⁵. We also show one spectrum observed with the Gemini-GMOS instrument from the Transient Name Server⁶. The collage of all the spectra is shown in Figure D1 and the spectroscopy log can be found in Table E1.

3. METHODS AND ANALYSIS

3.1. Extinction Correction

We correct for Milky Way extinction using the dust maps from Schlafly & Finkbeiner (2011). Along the line of sight of SN 2023zaw, E(B-V) = 0.25 mag. For reddening corrections, we use the extinction law in Cardelli et al. (1989) with $R_V = 3.1$.

For host extinction correction, we use the Na I D absorption lines of the host-galaxy (Poznanski et al. 2012; Stritzinger et al. 2018). We measure an equivalent width of 1.5 ± 0.2 Å, from the KCWI spectrum obtained at a phase of +2.5 days past *r*-band peak. To compute $A_{\rm V}$, we use $A_{\rm V}^{\rm host}[{\rm mag}] = 0.78(\pm 0.15) \times {\rm EW}_{\rm Na I D}[{\rm \AA}]$ (Stritzinger et al. 2018). We note that there are caveats in using the Na I D EW measurement to estimate host extinction, especially for low-resolution spectra (Poznanski et al. 2011) and in the presence of circumstellar material (Phillips et al. 2013).

We plot the g - r and r - i colors of SN 2023zaw before and after host extinction correction in Figure F1 and compare with other rapidly evolving transients and the SN Ib color template from Stritzinger et al. (2018).

3.2. Explosion epoch estimation

We fit for the r-band maximum epoch with a polynomial fit to the r-band photometry and obtain a peak magnitude of -16.7 mag and peak MJD of 60287.7. All phases mentioned in the paper will be in rest frame days

measured from this r-band peak epoch. The last nondetection before the first detection was on 60283 MJD (-4.7 days), with an upper limit of -14.8 mag in the o-band. We perform a power-law fit to the r-band data prior to the maximum epoch to estimate the explosion epoch. We find that the explosion epoch is 60284.4 \pm 0.5 MJD (-3.3 \pm 0.5 days).

3.3. Lightcurve properties

In Figures 2 and G1, we compare the q- and r-band evolution of SN 2023zaw to other fast-evolving and subluminous transients in the literature – fast-declining Type I SNe - SN 2005ek (Drout et al. 2013), SN 2010X (Kasliwal et al. 2010), SN 2018kzr (McBrien et al. 2019), SN 2019bkc (Prentice et al. 2020; Chen et al. 2020), the kilonova AT 2017gfo (Abbott et al. 2017), ultrastripped SN candidates - SN 2019dge (Yao et al. 2020), iPTF14gqr (De et al. 2018b), SN 2019wxt (Shivkumar et al. 2023; Agudo et al. 2023), SN 2022agco (Yan et al. 2023). We also compare with lightcurve templates for Type Ibn (Hosseinzadeh et al. 2017) and Type Ibc SNe (Taddia et al. 2015). SN 2023zaw is much fainter and faster evolving than the typical Type Ibn or Ibc SN. However, we note that some Type Ibn/Icn SNe also show rapid evolution: LSQ13ccw (Hosseinzadeh et al. 2017), SN 2023emg (Pursiainen et al. 2023). We fit a polynomial to the lightcurves and measure the rise time and fade time as the duration above the half-maximum of the peak. SN 2023zaw has a rise time $(t_{rise}^{1/2})$ of 1.8 days and a decline time $(t_{dec}^{1/2})$ of 3.1 days. The time above half maximum of the peak is shorter than for all other fast-evolving faint transients in the literature (see Figures 2, A1, and G1). The lightcurve declines by 2.7 mag in 10 days in the r-band and by 2.7 mag in 6 days in the q-band. SN 2023zaw is only slower than kilonova AT 2017 gfo and has a comparable decline timescale to SN 2019bkc and SN 2018kzr. In the r-band, SN 2023zaw has a very sharp initial decline of $\sim 0.28 \text{ mag day}^{-1}$ followed by a transition to a relatively slower decline rate of ~ 0.07 mag day⁻¹ after ~10 days since peak. In the g band, SN 2023zaw shows a very sharp initial decline of $\sim 0.44 \text{ mag day}^{-1}$ till 10 days post-peak. The peak absolute magnitude in the r band before and after host-extinction correction is -15.6 mag and -16.7 mag respectively. In the g-band, the peak absolute magnitude before and after host-extinction correction is -15.2mag and -16.6 mag respectively. We compare the time above half maximum and the absolute luminosity of SN 2023zaw to a sample of SESNe detected by ZTF and other fast and faint SNe in the literature in Figure A1. SN 2023zaw shows the fastest evolution among all transients at similar luminosities.

⁴ https://github.com/jkrogager/PyNOT

⁵ https://kcwi-drp.readthedocs.io

⁶ https://www.wis-tns.org/




Figure 3. The spectra of SN 2023zaw (in black) compared with other fast-evolving transients in the literature – fast-declining Type I SNe – SN 2010X (Kasliwal et al. 2010), ultra-stripped SNe candidates – SN 2019dge (Yao et al. 2020), iPTF14gqr (De et al. 2018b), SN 2019wxt (Shivkumar et al. 2023; Agudo et al. 2023), SN 2022agco (Yan et al. 2023), canonical Type Ib SN – iPTF14bvn (Fremling et al. 2018), Type Ibn – SN 2006jc (Foley et al. 2007). We also compare with theoretical predictions for '.Ia' SNe (Shen et al. 2010) and nickel-free SN models (Kleiser & Kasen 2014). The spectra are corrected for Milky Way and host extinction. The x-axis represents the rest wavelength. The He I and Ca II emission lines are shown in dashed vertical lines. The telluric absorption features are indicated by the \oplus symbol.

3.4. Spectral Properties

In the early SEDM spectra taken before r-band peak, we see a blue, featureless continuum. We see multiple broad helium lines – He I $\lambda\lambda$ 5876, 6678, 7065, in the subsequent spectra. The expansion velocity measured from the absorption in the P-Cygni profile of the He I lines evolved from 12,000 to 9700 km s⁻¹ from 0 days to 8 days post-peak. These He I absorption lines resemble those seen in Type Ib SNe and we obtain a decent fit to Type Ib SNe at similar or later phases using the SuperNova Identification (SNID; Blondin & Tonry 2007) code. This is consistent with the initial classification made by Gillanders et al. (2023). There is also a redshifted component to the He I λ 5876 line profile in the spectrum taken at +8 and +29 days, probably due to asymmetrical ejecta. He I line velocities drop to ~ 3000 $\rm km~s^{-1}$ at a phase of 29 days. Prominent narrow He I emission lines can be seen in the LRIS spectra obtained at 29 and 61 days after peak. The FWHM velocities of the He I emission lines in the latest LRIS spectrum are $\sim 1050 \pm 100 \text{ km s}^{-1}$.

In the photospheric phase, we do not see any broad O I λ 7774 emission line, which is common in strippedenvelope SNe. We also see broad, high-velocity lines of the Ca II NIR triplet at ~ 12,000 km s⁻¹ as early as 2 days after peak. Such high-velocity early Ca NIR absorption lines are not seen in other fast-evolving SNe (see Figure 3). Also, unlike other rapid-declining SNe, SN 2023zaw does not reach nebular phase even at 29 days after peak. Broad nebular lines such as [O I] $\lambda\lambda$ 6300, 6364, Ca II NIR triplet are also not seen in the latest LRIS spectrum taken at +61 days.

3.5. Bolometric Lightcurve

We bin the data in intervals of one day and fit with the Planck blackbody function where there are detections in at least three filters. We perform the fit using the Python EMCEE package (Foreman-Mackey et al. 2013). The blackbody temperature, radius and luminosity thus obtained are shown in Table H1. The errors in the model parameters are determined by extracting the 16th and 84th percentiles of the posterior probability distribution. For the later epochs, where we have photometry in only one band, we use the extrapolated blackbody temperatures obtained by fitting an exponential decay function to the temperature data from previous epochs. The temperature stays roughly constant at ≈ 4000 K. For these later epochs, we also estimate the bolometric luminosity as νL_{ν} . These estimates agree with each other within 3σ , with the latter method slightly underestimating the bolometric luminosity as expected.

We plot the best-fit blackbody parameters and compare these with other ultra-stripped SN candidates in Figure 4. We can see that the bolometric light curve of SN 2023zaw is similar in peak brightness to the other SNe but has a faster decline rate within the first 10 days after *r*-band peak. The decline of the bolometric luminosity slows down after around 6 days post-peak. The photospheric radius increases up to 4 days post-peak, reaching a maximum of ~10 000 R_{\odot}. The temperature decreases over time, from a maximum of ~17 000 K on day 0 to ~4000 K on day 8.

3.6. Modeling the lightcurve

3.6.1. Radiation-Hydrodynamics Modeling

We use the pre-supernova model grid presented in Wu & Fuller (2022). This grid was produced through binary evolution of He-stars with an initial mass of 2.5 M_{\odot} – 3 M_{\odot} evolved up to silicon burning using the Modules for Experiments in Stellar Astrophysics code (MESA; Paxton et al. 2010). The binary systems evolved in Wu & Fuller (2022) include orbital periods of 1, 10, and 100 days. Predictions of the properties of the unbound CSM in the vicinity of each pre-supernova model were also produced. In this work we compare our data to both models with and without unbound CSM.

We use the SuperNova Explosion code (SNEC; Morozova et al. 2015) to explode the grid of pre-supernova stellar models described above. First, we exploded all models in the grid using a thermal bomb with $E_{\rm kin} =$ 0.4×10^{51} erg, and $M_{\rm Ni} = 0.0035$ M_{\odot}, with ⁵⁶Ni mixed all the way through the remaining star after mass excision of 1.4 M_{\odot} in the center.

The resulting bolometric lightcurves and expansion velocities were then compared to our data on SN 2023zaw. We find that the model Wu & Fuller (2022) produced from a He-star with an initial mass of $2.651 \,\mathrm{M}_{\odot}$ and an orbit period of 100 days reproduces the overall behavior seen in our data well, including the expansion velocities. Other models fail to reproduce both the early cooling-phase of the bolometric lightcurve (governed by the radius and mass in the bound envelope) or the late-time decline rate (governed by the total ejecta mass). To produce our best-fitting MESA+SNEC model shown in Figure 4, we modified the bound envelope radius of the raw MESA model so that it extends to $\approx 50 \ R_{\odot}$, by cutting the model grid appropriately (compared to $\approx 110 \text{ R}_{\odot}$ in the raw model). This would correspond to an orbital period between 10 and 100 days. We also modified the mass excision in the center to be $1.3 M_{\odot}$. These modifications result in the explosion parameters $E_{\rm kin}=0.4\times 10^{51}~{\rm erg},\,M_{\rm Ni}=0.0035~M_{\odot},$ with complete mixing of ⁵⁶Ni through the ejecta, which has SN 2023zaw




Figure 4. Top: Comparison of the bolometric luminosity, best-fit radius and temperature evolution of SN 2023zaw from the blackbody fits described in Section 3.5 with ultra-stripped SN candidates – SN 2019dge (Yao et al. 2020), iPTF14gqr (De et al. 2018b), SN 2019wxt (Shivkumar et al. 2023; Agudo et al. 2023), SN 2022agco (Yan et al. 2023). Bottom left: Shock cooling emission model fit to the early-phase multiband lightcurve as described in Section 3.6.2. Bottom right: Best-fits for bolometric luminosity with radiation-hydrodymanics and analytical models as described in Section 3.6.

a total mass of $M_{\rm ej}=0.56~M_{\odot}$, including the bound envelope. Models including the unbound CSM predicted by Wu & Fuller (2022) are largely inconsistent with our data. The luminosity during the first week can be similar to what we observe if the explosion energy is adjusted to be low ($<0.1\times10^{51}$ erg). However, such models are inconsistent (too bright) at later times and inconsistent with the overall decay rate.

3.6.2. Analytical Modeling

The bolometric luminosity declines at a rapid rate for the first ~ 6 days after peak (see Figure 4). Radioactivity alone is unable to power this early-time luminosity under the constraint that the ejecta mass must be greater than the nickel mass. The peak luminosity of 2.5×10^{42} erg s⁻¹ and timescale of evolution (\approx 4.9 days) falls in the forbidden region of nickel-powered lightcurves (see Figure 1 in Kasen 2017). Instead, we model this early lightcurve as being powered by shockcooling emission from the shock-heated bound or unbound extended stellar material. We use the shockcooling emission model presented in Piro et al. (2021) to fit the early-time multiband photometry. We obtain a good fit for all the filters for photometry up to 6 days past peak (Figure 4). The explosion time is constrained to be -2.7 ± 0.2 days before the peak. Based on the fit, we find that the best-fit envelope mass is 0.20 ± 0.01 M_{\odot} and envelope radius is $55 \pm 11 \text{ R}_{\odot}$. This is comparable to the envelope properties measured from shockcooling emission modeling in other Type Ibc SNe (see Figure I1). The envelope properties of SN 2023zaw are also consistent with theoretical models for bound and unbound stellar material (Figure I2).

For estimating the Ni mass, we first subtract the contribution of the shock cooling emission estimated above from the bolometric luminosity, which we then assume to be entirely powered by radioactive decay. We use the analytical expression for a Ni-powered bolometric lightcurve provided in Arnett et al. (1989), Valenti et al. (2008), and Wheeler et al. (2015). Further de-




Figure 5. We compare the estimated ejecta mass and nickel mass with Type Ibc SNe from Taddia et al. (2018), double-peaked Type Ibc SNe from Das et al. (2023a) and other ultra-stripped SN candidates – SN 2019dge (Yao et al. 2020), iPTF14gqr (De et al. 2018b), SN 2019wxt (Shivkumar et al. 2023; Agudo et al. 2023), SN 2022agco (Yan et al. 2023). The red plus sign denotes the best-fit parameters by fitting shock cooling emission + nickel decay to the lightcurve after host-extinction correction.

tails on the model fitting can be found in Yao et al. (2020, their Appendix B). Based on the fit, we find that the required Ni-mass to power the supernova is $0.24 \pm 0.14 \times 10^{-2} \,\mathrm{M_{\odot}}$. We measure a photon diffusion timescale (τ_m) of $6.21^{+0.11}_{-0.11}$ days and characteristic γ -ray diffusion timescale (t_o) of $36.04^{+3.41}_{-2.74}$ days.

We assume that the photospheric velocity is ~ 12000 km s⁻¹, as assumed from the He I absorption lines in the ALFOSC spectrum taken at r-band peak. We use an opacity (κ) value of 0.07 cm² g⁻¹, appropriate for H-poor SNe (Taddia et al. 2018). We estimate the ejecta mass using the following relation: $\tau_m = \left(\frac{2\kappa M_{ej}}{13.8 \text{cv}_{\text{phot}}}\right)^{1/2}$, which gives $M_{ej} \approx 0.52^{+0.02}_{-0.02} \times \left(\frac{v_{\text{phot}}}{12000 \text{ km/s}}\right) M_{\odot}$. The calculated explosion kinetic energy is $\approx 4.82^{+0.18}_{-0.18} \times 10^{50} \text{ erg s}^{-1}$.

These explosion parameters are an order of magnitude lower than for regular Type Ibc SNe. The estimated ejecta mass is comparable to other ultra-stripped SNe in the literature while the Ni-mass is smaller by a factor of ~5. The rapid light curve decline is explained by the low ejecta mass, which allows γ -rays to escape already a few days after the peak. Overall, SN 2023zaw occupies a unique location in the phase space plot of nickel mass versus ejecta mass (see Figure 5). For comparison, we also plot the nickel and ejecta masses estimated for a sample of stripped-envelope SNe from Taddia et al. (2018) and double-peaked Type Ibc SNe from Das et al. (2023a). We note that the double-peaked SN 2021inl (Jacobson-Galán et al. 2022; Das et al. 2023a) is the other SN with Ni-mass < 0.01 M_{\odot}.

We also use the analytical models given in Khatami & Kasen (2019) to obtain a nickel mass estimate of $\approx 2.5 \times 10^{-2} M_{\odot}$ and an ejecta mass estimate of $\approx 0.45 M_{\odot}$.

3.7. Host galaxy environment

We observed SN 2023zaw and its immediate environment with the integral-field unit KCWI (Section 2.3). These data allow us to address two outstanding questions: i) did SN 2023zaw explode in a star-forming region, and ii) how high is the host attenuation along the line of sight towards SN 2023zaw?

We cleanly detect emission lines from a starforming region at the SN position in the unbinned KCWI data. To measure the properties of the star-forming region, we extract the SN spectrum using a 4 × 4-px aperture, tie the flux scale to our g- and *i*-band photometry, and extract the flux of H α -H γ , [O II] $\lambda\lambda$ 3729,3729, [O III] $\lambda\lambda$ 4959,5007, [N II] $\lambda\lambda$ 6548,6584, and [S II] $\lambda\lambda$ 6717,6731 by fitting each

Table 1. Comparison of explosion properties of SN 2023zaw with ultra-stripped SN candidates – SN 2019dge (Yao et al. 2020), iPTF14gqr (De et al. 2018b), iPTF16hgs (De et al. 2018c), SN 2019wxt (Shivkumar et al. 2023; Agudo et al. 2023), SN 2022agco (Yan et al. 2023).

Source	Redshift	Host Type	r-band Peak	$M_{\rm ej}$	$M_{\rm Ni}$	$R_{\rm ext}$	$M_{\rm ext}$
			(mag)	$({ m M}_{\odot})$	$(10^{-2} {\rm M}_{\odot})$	(10^{13} cm)	$(10^{-2} {\rm M}_{\odot})$
SN 2023zaw	0.0101	Spiral	-16.7 ± 0.1	$0.52\substack{+0.02\\-0.02}$	$0.24\substack{+0.02\\-0.02}$	$0.4^{+0.1}_{-0.1}$	$19.53\substack{+0.99\\-0.98}$
SN 2021 agco	0.01056	Spiral	-16.2 ± 0.24	$0.26\substack{+0.04\\-0.02}$	$2.2^{+0.2}_{-0.3}$	$0.55_{-0.14}^{+0.18}$	$10.04^{+1.87}_{-1.05}$
AT 2019 wxt	0.036	Compact	-16.6 ± 0.4	$0.20_{-0.11}^{+0.12}$	$2.7^{+0.33}_{-0.18}$	$35.8^{+4.06}_{-3.68}$	$3.55_{-0.11}^{+0.12}$
SN 2019dge	0.021	Compact	-16.3 ± 0.2	$0.30\substack{+0.02 \\ -0.02}$	$1.6\substack{+0.04\\-0.03}$	$1.2^{+0.06}_{-0.05}$	$9.71^{+0.28}_{-0.27}$
iPTF14gqr	0.063	Spiral	-17.5 ± 0.2	$0.20_{\pm 0.10}^{-0.10}$	$5.0\substack{+0.14 \\ -0.15}$	$6.1^{+8.73}_{-3.18}$	$2.59^{+0.46}_{-0.34}$
iPTF16hgs	0.017	Spiral	-15.5 ± 0.2	$1.68^{+0.28}_{-0.25}$	$2.5^{+0.20}_{-0.22}$	$2.6^{+14.08}_{-1.80}$	$9.27^{+3.40}_{-2.48}$

line with a Gaussian. All measurements are summarised in Table B1.

We infer the attenuation by ionised gas from the flux ratio between H β and H γ (Momcheva et al. 2013). Using the Calzetti et al. (2000) attenuation model, we measure $E(B-V)_{\rm gas} = 0.58 \pm 0.20$ mag, which translates to a stellar continuum colour excess of 0.26 ± 0.09 mag. This value is comparable to the reddening inferred from the Na I D absorption lines (Section 3.1) and corroborates that the line of sight is suffering from significant host attenuation.

We note that the stellar populations near SN 2023zaw are evolved, and their spectra show conspicuous Balmer absorption lines. This could introduce a bias into the observed Balmer decrement. To quantify the impact, we centre a box annulus on the SN position and extract a spectrum from the blue KCWI data between 2 to 4 px from the SN position. We fit the stellar continuum with the stellar population fitting code FIREFLY version 1.0.3 (Wilkinson et al. 2017; Neumann et al. 2022), utilising the stellar population models from Maraston et al. (2020) and assuming the Kroupa initial mass function (IMF), a spectral resolution of 1000–1250 appropriate for our spectroscopic observation (Morrissey et al. 2018b) and a velocity dispersion of $130 \pm 20 \text{ km s}^{-1}$. The best-fit yields an $E(B-V)_{\text{star}}$ of 0.36 mag, comparable to the value at the explosion site. Henceforth, we use $E(B-V)_{gas} = 0.58 \pm 0.20$ mag to characterise the properties of the star-forming region further.

The attenuation-corrected H α flux of $1.2 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ translates to a star-formation rate of $1.4^{+1.2}_{-0.6} \times 10^{-2} M_{\odot} \text{ yr}^{-1}$ utilising the relationship between the H α luminosity and the star-formation rate from Kennicutt (1998) and the scaling factor from Madau & Dickinson (2014) to convert from the Salpeter to the Chabrier IMF in the Kennicutt (1998) relation. The error estimate includes the uncertainty of the flux measurements, the gas-phase attenuation and the lumi-

nosity distance of SN 2023zaw's host galaxy. We stress that the large SFR error reflects the uncertainty in the attenuation correction, not a low precision of the H α flux measurement (Table B1).

Detecting emission lines from hydrogen, nitrogen and oxygen also allows us to measure the metallicity of the star-forming region. Using the O3N2 metallicity indicator and the calibration from Curti et al. (2017), we infer a metallicity of 1.18 ± 0.02 solar (statistical error), adopting a solar oxygen abundance of 8.67 (Asplund et al. 2009).

Next, we draw our attention to SN 2023zaw's neighbourhood. Figure 1 shows a map of the star-formation activity out to a distance of 6.7 kpc from SN 2023zaw's location. This image was generated from the KCWI data. First, we extracted a 20-Å-wide image centred on the wavelength of H α . Then, we extracted two additional 20-Å-wide images from emission-line free regions adjacent to H α and interpolated between them to remove the galaxy/SN flux at the wavelength of H α . Finally, the intensity scale was corrected for MW extinction, scaled to the SN photometry and converted to star-formation rate as we did for the explosion site. The SFRs in Figure 1 are strict lower limits. They account for neither host attenuation nor stellar Balmer absorption lines.

The explosion site, marked by the blue circle in Figure 1, is located near the outskirts of a larger region with ongoing star-formation activity. Although the level of star formation activity at the explosion site is in the lower half of the intensity distribution, this star-forming region extends much farther. Inspecting the neighbouring spaxels reveals that the star-formation activity extends to regions of even less intensity. Thus, the location of SN 2023zaw favors a massive star progenitor, but it does not rule out a thermonuclear origin.

4. DISCUSSION: PROGENITOR AND EVOLUTIONARY PATHWAY

In this section, we discuss the possible progenitors and evolutionary pathways for SN 2023zaw.

4.1. Thermonuclear '.Ia' SN

In the '.Ia' SN scenario, the progenitor is a white dwarf in a close binary ($P_{orb} < 1$ hour) system, accreting helium from its companion. They have a tenth of the ejecta mass and explosion energy of a normal Type Ia supernova (Bildsten et al. 2007). We compare the r-band lightcurve and bolometric luminosity with '.Ia' SN models from Shen et al. (2010) in Figure 6. The peak luminosities range from $0.5 - 5 \times 10^{42}$ erg s⁻¹ or peak bolometric magnitudes from -15.5 to -18 mag. consistent with SN 2023zaw. We highlight the best-fit '.Ia' SN bolometric lightcurve which is an excellent fit to the early-time data before host-extinction correction. Prominent He I features from unburnt He are predicted, which is consistent with the spectra of SN 2023zaw. We note that the absence of broad O I λ 7774 and the presence of Ca II NIR lines at early phases resemble the spectra of SN 2010X (Kasliwal et al. 2010) which is a '.Ia' SN candidate (see Figure 3). However, the fact that the decline of the r-band lightcurve slows down after \sim 10 days post-peak, is not consistent with the '.Ia' SN models (Shen et al. 2010).

4.2. Accretion-induced collapse (AIC) of white dwarf

O-Ne white dwarfs collapse to form a neutron star when it approaches Chandrasekhar mass. Numerical simulations predict very low explosion energy (~ 0.5×10^{49} erg). We compare the bolometric lightcurve prediction from Darbha et al. (2010) with SN 2023zaw in Figure 6. The predicted peak luminosity is ~ 10 times fainter than that observed for SN 2023zaw. The models do not predict the late-time tail seen in SN 2023zaw. The nickel mass predicted in AIC models ($\leq 0.001 \text{ M}_{\odot}$) is slightly smaller than that estimated for SN 2023zaw. The ejecta velocities predicted for AIC explosions are very high (~ 0.1c), which is not consistent with SN 2023zaw.

4.3. White $dwarf - neutron \ star/black$ hole merger

Spectroscopic models of WD-NS/BH mergers (Gillanders et al. 2020) predict that the ejecta should be rich in intermediate elements such as O, Mg, S. This is not consistent with the observed spectra for SN 2023zaw.

4.4. Nickel free core-collapse supernova

A potential powering mechanism for a fast-evolving, low-Ni mass SN is the 'Ni-free' SN scenario (Kleiser & Kasen 2014; Kleiser et al. 2018). In this case, the supernova is not powered by decay of radionuclides like ⁵⁶Ni but is instead powered by shock-interactions with an extended hydrogen-poor circumstellar medium (CSM). We compare the *r*-band lightcurve and bolometric luminosity of SN 2023zaw with Ni-free SN models from Kleiser et al. (2018) in Figure 6. If there is a significant amount of Ni present (> 0.05 M_☉), it shows up as a second peak in the model lightcurves, which is not seen for SN 2023zaw. The models with 0 or 0.01 M_☉ of nickel produce bright and short-lived peaks. However, none of the lightcurves can explain the rapid rise and the slower-evolving late-time lightcurve tail of SN 2023zaw.

4.5. Electron-capture SN

The low nickel mass is consistent with super Asymptotic Giant Branch (sAGB) progenitor stars with an O-Ne-Mg core that explodes as an electron-capture supernova (ECSN). However, a sAGB star has a hydrogenrich envelope. Since, the spectra do not show hydrogen, a binary system is necessary, where the hydrogen has been stripped off through Roche-lobe overflow or common envelope ejection. Stripped-envelope ECSN models in Moriya & Eldridge (2016) predict ejecta mass of $\sim 0.3 - 0.6 \ M_{\odot}$, nickel mass of $\sim 0.003 \ M_{\odot}$, and explosion energy of 10^{50} erg, which is consistent with SN 2023zaw. Enhanced production of calcium is predicted for these SN, which is also consistent with the broad Ca emission lines seen in SN 2023zaw, as early as 2 days after peak. We compared the theoretical lightcurves from Moriya & Eldridge (2016) with SN 2023zaw in Figure 6. The predicted peak bolometric luminosities $(\sim 10^{41} \text{ erg s}^{-1})$ are ~ 10 times fainter than the peak luminosity measured for SN 2023zaw.

4.6. Ultra-stripped core-collapse supernova from a low-mass progenitor embedded in a He-rich CSM

An ultra-stripped SN (Tauris et al. 2015) is a possible channel for the formation of double neutron-star systems. They arise from a binary star system where a Hestar undergoes a high degree of stripping by its smaller companion, such that the remaining ejecta mass is < 1 M_{\odot} . The nickel mass is also expected to be quite low (~ 0.01 M_{\odot}). Due to the low nickel and ejecta mass, the lightcurves are expected to be faint and show fast evolution. The multiband lightcurve and bolometric properties of SN 2023zaw are consistent with an ultra-stripped SN scenario (Figures 2, 4 and G1). We compare the measured ejecta and nickel mass for SN 2023zaw with other ultra-stripped SN candidates in the literature in Table 1 and Figure 5. The estimated ejecta mass is comparable with other ultra-stripped SN candidates, while the nickel mass is a factor of ~ 5 lower. We could also reproduce the observed lightcurve by using ultra-stripped SN progenitor models evolved with MESA and exploded using SNEC (see Section 3.6.1).

4.6.1. Implication of narrow He I lines in the late-time spectra

The presence of narrow He I emission lines in the latetime spectra at 30 and 60 days post-peak implies the presence of a He-rich dense CSM shell, which is consistent with the predictions of ultra-stripped SN models. We note that we do not see prominent narrow He I emission lines in the spectra taken prior to 9 days after peak. It is likely the broad He I lines from the ejecta dominate over the narrow CSM emission lines in the initial days. But if there is actually no CSM interaction till +8 days, we can constrain the CSM shell to be at $\sim 10^{15}$ cm, which is consistent with the short-period binary models from Wu & Fuller (2022). While the mass transfer model in Wu & Fuller (2022) assumes that the stellar material lost through the L2 point is distributed spherically, a torus-like distribution of the CSM is also likely (Pejcha et al. 2016; Lu et al. 2023). The viewing angle will strongly influence the observed spectral line widths. In this scenario, the early spectra do not show narrow lines as the CSM is engulfed in the SN ejecta, similar to the model described in Andrews & Smith (2018) for iPTF14hls. Once the photosphere recedes, the CSM is accelerated by the collision of the ejecta to intermediate velocities of $\approx 1000 \text{ km s}^{-1}$. In this scenario, the absence of narrow lines in the ALFOSC spectrum taken at +0 days can be explained if the CSM is confined within 10^{14} cm, which is consistent with the majority of binary mass transfer models in Wu & Fuller (2022). We note that the CSM interaction does not show up as an additional bump in the optical lightcurve. This could be explained by a very low mass He-rich CSM, which is consistent with the estimated low progenitor and ejecta masses. If CSM interaction does contribute significantly to the luminosity, the nickel mass we estimate should be considered as an upper limit.

4.6.2. Implication of very low nickel mass

Observations and parametrized 1D models of massive star explosions suggest that typically ~ 0.05 M_{\odot} Ni should be ejected from a core-collapse SN. Anderson (2019) and Meza & Anderson (2020) measured the Ni-mass for a sample of Type II and Type Ibc SNe in the literature and found that the Ni-mass synthesized in SESNe was significantly higher than for Type II SNe. The lowest Ni-mass in their Type Ibc sample was 0.015 M_{\odot}, and 25% of all Type II SNe has a Ni-mass below this limit. However, this might be due to an observational bias as low Ni-mass SESNe are harder to detect as they are likely to be faint as well as fast-evolving. The extremely low amount of Ni-mass in SN 2023zaw ($\sim 0.002 M_{\odot}$) could imply a fallback event, where most of the Ni produced in the core-collapse immediately fall into the newly formed black hole (Turatto et al. 1998; Sollerman 2002; Balberg et al. 2000). However, the observed high velocities do not favor this scenario.

The other explanation is that the progenitor's initial mass lies in the low-mass end of core-collapse SN progenitors, as stars with low-mass iron cores produce a low amount of Ni. Stockinger et al. (2020) and Sandoval et al. (2021) obtained ~ 0.002-0.005 M_{\odot} Ni mass for a 9.6 M_{\odot} progenitor. More recent 3D simulations (Burrows et al. 2024) estimate $0.002 - 0.006 \,\mathrm{M}_{\odot}$ Nimass for a 9 M_{\odot} ZAMS mass progenitor and > 0.01 M_{\odot} for progenitors with initial mass greater than 9.25 M_{\odot} . We note that these models are not for strippedenvelope SNe. However, if they undergo mass-loss after He-core burning, the core evolution of a star is decoupled from the envelope evolution and the nickel mass is expected to be the same. The amount of Ni synthesized in ultra-stripped models for low-mass CO stars is ≤ 0.01 M_{\odot} (Moriya et al. 2017; Sawada et al. 2022), which is consistent with our estimate for SN 2023zaw.

5. CONCLUSION

In this paper, we have presented the observations and analysis of an unusual stellar explosion - SN 2023zaw. It has the lowest nickel mass among all stripped-envelope SNe discovered so far. SN 2023zaw also shows the fastest evolution among all stripped-envelope SNe, with a time above half maximum of 4.9 days in the r-band. The photospheric spectra show broad He I emission lines. It also shows broad Ca NIR emission lines as early as 2 days after peak.

We performed radiation-hydrodynamical and analytical modeling of the lightcurve by fitting with a combination of shock-cooling emission and nickel decay. The early-time lightcurve fits well with shock-cooling emission, with an envelope mass of $\approx 0.2 \, M_{\odot}$ and an envelope radius of $\approx 50 \, R_{\odot}$. The estimated ejecta and nickel masses are $\approx 0.5 \, M_{\odot}$ and $\approx 0.002 \, M_{\odot}$ respectively.

We interpret SN 2023zaw as an ultra-stripped SN from a low-mass He-star – compact object binary. The late-time evolution of the He-star results in an extended He-rich envelope. There is also spectral evidence for thick He-rich CSM, likely from pre-SN mass loss episodes from late-time Roche lobe overflow. The very low nickel mass, ejecta mass, and explosion energy can only be explained by a progenitor with an initial mass less than around 10 M_{\odot} .

Figure 6. Comparison of the r-band and bolometric luminosity of SN 2023zaw with theoretical models for '.Ia' SNe (Shen et al. 2010), nickel-free SNe (Kleiser et al. 2018), electron-capture SN (Moriya & Eldridge 2016), and accretion-induced collapse of white dwarf (Darbha et al. 2010). The bold blue and green lines indicate the best-fit '.Ia' and nickel-free SN models, respectively.

The late-time spectra show prominent narrow He I at a FWHM velocity of $\sim 1000 \text{ km s}^{-1}$, indicating the presence of He-rich CSM. It is unclear if CSM interaction also provides a powering mechanism for SN 2023zaw. A number of other models for faint and rapidly evolving transients such as the nickel-free SN, accretion induced collapse, electron capture supernova were ruled out based on the peak luminosity and the presence of a late-time tail in the lightcurve of SN 2023zaw. The excess luminosity might be explained if the luminosity is boosted by CSM interaction. In this case, the estimated nickel mass should be treated as an upper limit. Including CSM interaction modeling in the future will help conclusively favor or rule out these scenarios. Also, late-time multi-wavelength follow-up will provide useful diagnostics in determining the CSM properties, the pre-SN mass-loss mechanism, and the evolutionary channel (Matsuoka & Maeda 2020; Kashiyama et al. 2022).


Spectroscopic modeling is currently missing for most of the theoretical scenarios for producing faint and rapid transients. Detailed modeling is required to explain some unusual spectroscopic features of SN 2023zaw such as – the presence of strong Ca NIR lines close to peak, redshifted component of He I λ 5876 emission line, not reaching nebular phase till 30 days in spite of the low ejecta mass, the emergence of narrow He I emission lines and the absence of [Ca II], [O I] and Ca NIR emission lines in the nebular phase.

The extremely low nickel mass of SN 2023zaw gives rise to the question – what is the lowest nickel mass that is synthesized in a stripped-envelope SN? Current literature shows that the nickel mass measured in strippedenvelope SNe is significantly higher than in Type II SNe (Anderson 2019). It is likely that this is due to an observational bias, as SESNe with low nickel masses are likely to be faint and fast evolving and hence harder to detect, classify, and follow-up. SN 2023zaw underscores the existence of this undiscovered population of low nickel mass SESNe ($< 0.005 M_{\odot}$) with progenitor masses in the low mass end of core-collapse SNe ($< 10 M_{\odot}$). A systematic study of the lowest-nickel mass stripped-envelope SNe, including their rates and the nickel mass distribution of low-luminosity SESNe will be explored in future work. Candidates for such low nickel mass strippedenvelope SNe include double peaked Type Ibc SNe such as SN 2021inl (Jacobson-Galán et al. 2022; Das et al. 2023a), rapidly-evolving ($t_{1/2,r} < 10$ days) Type IIb SNe (e.g., Ho et al. 2023; Das et al. 2023b, Fremling et al. in prep), rapidly-declining H-poor SNe such as SN 2020ghq (Wang et al. in prep). For SNe with extremely low Nimass, the shock cooling emission is likely to dominate over nickel decay. The ultra-stripped SNe candidates in the literature show an early luminosity excess that is powered by shock-cooling. Future wide-field UV surveys such as ULTRASAT (Sagiv et al. 2014; Shvartzvald et al. 2023), UVEX (Kulkarni et al. 2021), and deep ground-based surveys such as LSST (Ivezić et al. 2008) in synergy with high-cadenced surveys such as ZTF will provide an exciting opportunity to explore this phase space.

6. DATA AVAILABILITY

All the photometric and spectroscopic data used in this work will be made available on GitHub after publication.

The optical photometry and spectroscopy will also be made public through WISeREP, the Weizmann Interactive Supernova Data Repository (Yaron & Gal-Yam 2012).



7. ACKNOWLDEGEMENT

KKD acknowledges support from the Schwartz Reisman Collaborative Science Program, which is supported by the Gerald Schwartz and Heather Reisman Foundation.

M.W.C acknowledges support from the National Science Foundation with grant numbers PHY-2308862 and PHY-2117997.

S. Schulze is partially supported by LBNL Subcontract NO. 7707915.

Based on observations obtained with the Samuel Oschin Telescope 48-inch and the 60-inch Telescope at the Palomar Observatory as part of the Zwicky Transient Facility project. ZTF is supported by the National Science Foundation under Grant No. AST-2034437 and a collaboration including Caltech, IPAC, the Oskar Klein Center at Stockholm University, the University of Maryland, University of California, Berkeley, the University of Wisconsin at Milwaukee, University of Warwick, Ruhr University, Cornell University, Northwestern University and Drexel University. Operations are conducted by COO, IPAC, and UW.

SED Machine is based upon work supported by the National Science Foundation under Grant No. 1106171.

The ZTF forced-photometry service was funded under the Heising-Simons Foundation grant #12540303 (PI: Graham).

The Gordon and Betty Moore Foundation, through both the Data-Driven Investigator Program and a dedicated grant, provided critical funding for SkyPortal.

This research has made use of the NASA/IPAC Extragalactic Database (NED), which is funded by the National Aeronautics and Space Administration and operated by the California Institute of Technology.

The data presented here were obtained in part with ALFOSC, which is provided by the Instituto de Astrofisica de Andalucia (IAA) under a joint agreement with the University of Copenhagen and NOT.

The Liverpool Telescope is operated on the island of La Palma by Liverpool John Moores University in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofisica de Canarias with financial support from the UK Science and Technology Facilities Council. Based on observations made with the Italian Telescopio Nazionale Galileo (TNG) operated on the island of La Palma by the Fundación Galileo Galilei of the INAF (Istituto Nazionale di Astrofisica) at the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofisica de Canarias.

The W. M. Keck Observatory is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation. The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Maunakea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.

MMT Observatory access was supported by Northwestern University and the Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA).

REFERENCES

Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017, PhRvL, 119, 161101,

doi: 10.1103/PhysRevLett.119.161101

Agudo, I., Amati, L., An, T., et al. 2023, A&A, 675, A201, doi: 10.1051/0004-6361/202244751

Anderson, J. P. 2019, A&A, 628, A7, doi: 10.1051/0004-6361/201935027

- Andreoni, I., Coughlin, M. W., Kool, E. C., et al. 2021, ApJ, 918, 63, doi: 10.3847/1538-4357/ac0bc7
- Andrews, J. E., & Smith, N. 2018, MNRAS, 477, 74, doi: 10.1093/mnras/sty584
- Anupama, G. C., Sahu, D. K., Gurugubelli, U. K., et al. 2009, MNRAS, 392, 894,

doi: 10.1111/j.1365-2966.2008.14129.x

Arnett, W. D., Bahcall, J. N., Kirshner, R. P., & Woosley, S. E. 1989, ARA&A, 27, 629,

doi: 10.1146/annurev.aa.27.090189.003213

Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, ARA&A, 47, 481,

doi: 10.1146/annurev.astro.46.060407.145222

- Balberg, S., Zampieri, L., & Shapiro, S. L. 2000, ApJ, 541, 860, doi: 10.1086/309472
- Bellm, E. C., Kulkarni, S. R., Graham, M. J., et al. 2019, PASP, 131, 018002, doi: 10.1088/1538-3873/aaecbe
- Bildsten, L., Shen, K. J., Weinberg, N. N., & Nelemans, G. 2007, ApJL, 662, L95, doi: 10.1086/519489
- Blagorodnova, N., Neill, J. D., Walters, R., et al. 2018, PASP, 130, 035003, doi: 10.1088/1538-3873/aaa53f
- Blondin, S., & Tonry, J. L. 2007, ApJ, 666, 1024, doi: 10.1086/520494

- Burrows, A., Wang, T., & Vartanyan, D. 2024, arXiv e-prints, arXiv:2401.06840, doi: 10.48550/arXiv.2401.06840
- Calzetti, D., Armus, L., Bohlin, R. C., et al. 2000, ApJ, 533, 682, doi: 10.1086/308692
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, ApJ, 345, 245, doi: 10.1086/167900
- Cenko, S. B., Fox, D. B., Moon, D.-S., et al. 2006, PASP, 118, 1396, doi: 10.1086/508366
- Chambers, K. C., Magnier, E. A., Metcalfe, N., et al. 2016, arXiv e-prints. https://arxiv.org/abs/1612.05560
- Chen, P., Dong, S., Stritzinger, M. D., et al. 2020, ApJL, 889, L6, doi: 10.3847/2041-8213/ab62a4
- Clark, P., Maguire, K., Inserra, C., et al. 2020, MNRAS, 492, 2208, doi: 10.1093/mnras/stz3598
- Curti, M., Cresci, G., Mannucci, F., et al. 2017, MNRAS, 465, 1384, doi: 10.1093/mnras/stw2766
- Darbha, S., Metzger, B. D., Quataert, E., et al. 2010, MNRAS, 409, 846, doi: 10.1111/j.1365-2966.2010.17353.x
- Das, K. K., Kasliwal, M. M., Sollerman, J., et al. 2023a, arXiv e-prints, arXiv:2306.04698,
 - doi: 10.48550/arXiv.2306.04698
- Das, K. K., Kasliwal, M. M., Fremling, C., et al. 2023b, ApJ, 959, 12, doi: 10.3847/1538-4357/acfeeb
- De, K., Kasliwal, M. M., Cantwell, T., et al. 2018a, ApJ, 866, 72, doi: 10.3847/1538-4357/aadf8e
- De, K., Kasliwal, M. M., Ofek, E. O., et al. 2018b, Science, 362, 201, doi: 10.1126/science.aas8693
- De, K., Kasliwal, M. M., Cantwell, T., et al. 2018c, ApJ, 866, 72, doi: 10.3847/1538-4357/aadf8e
- De, K., Kasliwal, M. M., Tzanidakis, A., et al. 2020, ApJ, 905, 58, doi: 10.3847/1538-4357/abb45c
- Dekany, R., Smith, R. M., Riddle, R., et al. 2020, PASP, 132, 038001, doi: 10.1088/1538-3873/ab4ca2
- Dessart, L., Burrows, A., Livne, E., & Ott, C. D. 2007, ApJ, 669, 585, doi: 10.1086/521701
- Drout, M. R., Soderberg, A. M., Mazzali, P. A., et al. 2013, ApJ, 774, 58, doi: 10.1088/0004-637X/774/1/58
- Fabricant, D., Fata, R., Epps, H., et al. 2019, PASP, 131, 075004, doi: 10.1088/1538-3873/ab1d78
- Foley, R. J., Smith, N., Ganeshalingam, M., et al. 2007, ApJL, 657, L105, doi: 10.1086/513145
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, PASP, 125, 306, doi: 10.1086/670067
- Fremling, C., Sollerman, J., Taddia, F., et al. 2016, A&A, 593, A68, doi: 10.1051/0004-6361/201628275
- Fremling, C., Sollerman, J., Kasliwal, M. M., et al. 2018, A&A, 618, A37, doi: 10.1051/0004-6361/201731701
- Fremling, C., Miller, A. A., Sharma, Y., et al. 2020, ApJ, 895, 32, doi: 10.3847/1538-4357/ab8943

- Gillanders, J. H., Sim, S. A., & Smartt, S. J. 2020, MNRAS, 497, 246, doi: 10.1093/mnras/staa1822
- Gillanders, J. H., Huber, M., Chambers, K., et al. 2023, Transient Name Server AstroNote, 341, 1
- Graham, M. J., Kulkarni, S. R., Bellm, E. C., et al. 2019, PASP, 131, 078001, doi: 10.1088/1538-3873/ab006c
- Ho, A. Y. Q., Perley, D. A., Kulkarni, S. R., et al. 2020, ApJ, 895, 49, doi: 10.3847/1538-4357/ab8bcf
- Ho, A. Y. Q., Perley, D. A., Gal-Yam, A., et al. 2023, ApJ, 949, 120, doi: 10.3847/1538-4357/acc533
- Hosseinzadeh, G., Arcavi, I., Valenti, S., et al. 2017, ApJ, 836, 158, doi: 10.3847/1538-4357/836/2/158
- Ivezić, Ž., Tyson, J. A., Acosta, E., et al. 2008, ArXiv e-prints. https://arxiv.org/abs/0805.2366
- Jacobson-Galán, W. V., Venkatraman, P., Margutti, R., et al. 2022, ApJ, 932, 58, doi: 10.3847/1538-4357/ac67dc
- Kasen, D. 2017, in Handbook of Supernovae, ed. A. W. Alsabti & P. Murdin, 939,
- doi: 10.1007/978-3-319-21846-5_32 Kashiyama, K., Sawada, R., & Suwa, Y. 2022, ApJ, 935,
- 86, doi: 10.3847/1538-4357/ac7ff7
- Kasliwal, M. M., Kulkarni, S. R., Gal-Yam, A., et al. 2010, ApJL, 723, L98, doi: 10.1088/2041-8205/723/1/L98
- Kennicutt, Robert C., J. 1998, ARA&A, 36, 189, doi: 10.1146/annurev.astro.36.1.189
- Khatami, D. K., & Kasen, D. N. 2019, ApJ, 878, 56, doi: 10.3847/1538-4357/ab1f09
- Kim, Y. L., Rigault, M., Neill, J. D., et al. 2022, PASP, 134, 024505, doi: 10.1088/1538-3873/ac50a0
- Kleiser, I. K. W., & Kasen, D. 2014, MNRAS, 438, 318, doi: 10.1093/mnras/stt2191
- Kleiser, I. K. W., Kasen, D., & Duffell, P. C. 2018, MNRAS, 475, 3152, doi: 10.1093/mnras/stx3321
- Kulkarni, S. R., Harrison, F. A., Grefenstette, B. W., et al. 2021, arXiv e-prints, arXiv:2111.15608. https://arxiv.org/abs/2111.15608
- Laplace, E., Götberg, Y., de Mink, S. E., Justham, S., & Farmer, R. 2020, A&A, 637, A6, doi: 10.1051/0004-6361/201937300
- Leung, S.-C., Fuller, J., & Nomoto, K. 2021, ApJ, 915, 80, doi: 10.3847/1538-4357/abfcbe
- Leung, S.-C., Nomoto, K., & Blinnikov, S. 2019, ApJ, 887, 72, doi: 10.3847/1538-4357/ab4fe5
- Lu, W., Fuller, J., Quataert, E., & Bonnerot, C. 2023, MNRAS, 519, 1409, doi: 10.1093/mnras/stac3621
- Madau, P., & Dickinson, M. 2014, ARA&A, 52, 415, doi: 10.1146/annurev-astro-081811-125615
- Maraston, C., Hill, L., Thomas, D., et al. 2020, MNRAS, 496, 2962, doi: 10.1093/mnras/staa1489

- Margalit, B., & Metzger, B. D. 2016, MNRAS, 461, 1154, doi: 10.1093/mnras/stw1410
- Martin, C., Moore, A., Morrissey, P., et al. 2010, in Society of Photo-Optical Instrumentation Engineers (SPIE)
 Conference Series, Vol. 7735, Ground-based and Airborne Instrumentation for Astronomy III, ed. I. S. McLean,
 S. K. Ramsay, & H. Takami, 77350M,
 doi: 10.1117/12.858227
- Masci, F. J., Laher, R. R., Rusholme, B., et al. 2019, PASP, 131, 018003, doi: 10.1088/1538-3873/aae8ac
- Matsuoka, T., & Maeda, K. 2020, ApJ, 898, 158, doi: 10.3847/1538-4357/ab9c1b
- McBrien, O. R., Smartt, S. J., Chen, T.-W., et al. 2019, ApJL, 885, L23, doi: 10.3847/2041-8213/ab4dae
- Meza, N., & Anderson, J. P. 2020, A&A, 641, A177, doi: 10.1051/0004-6361/201937113
- Momcheva, I. G., Lee, J. C., Ly, C., et al. 2013, AJ, 145, 47, doi: 10.1088/0004-6256/145/2/47
- Moriya, T., Tominaga, N., Tanaka, M., et al. 2010, ApJ, 719, 1445, doi: 10.1088/0004-637X/719/2/1445
- Moriya, T. J., & Eldridge, J. J. 2016, MNRAS, 461, 2155, doi: 10.1093/mnras/stw1471
- Moriya, T. J., Mazzali, P. A., Tominaga, N., et al. 2017, MNRAS, 466, 2085, doi: 10.1093/mnras/stw3225
- Morozova, V., Piro, A. L., Renzo, M., et al. 2015, ApJ, 814, 63, doi: 10.1088/0004-637X/814/1/63
- Morrissey, P., Matuszewski, M., Martin, D. C., et al. 2018a, ApJ, 864, 93, doi: 10.3847/1538-4357/aad597
- —. 2018b, ApJ, 864, 93, doi: 10.3847/1538-4357/aad597
- Mould, J. R., Huchra, J. P., Freedman, W. L., et al. 2000, ApJ, 529, 786, doi: 10.1086/308304
- Neumann, J., Thomas, D., Maraston, C., et al. 2022, MNRAS, 513, 5988, doi: 10.1093/mnras/stac1260
- Oke, J. B., Cohen, J. G., Carr, M., et al. 1995, PASP, 107, 375, doi: 10.1086/133562
- Paxton, B., Bildsten, L., Dotter, A., et al. 2010, MESA: Modules for Experiments in Stellar Astrophysics, Astrophysics Source Code Library. http://ascl.net/1010.083
- Pejcha, O., Metzger, B. D., & Tomida, K. 2016, MNRAS, 461, 2527, doi: 10.1093/mnras/stw1481
- Pellegrino, C., Howell, D. A., Terreran, G., et al. 2022, arXiv e-prints, arXiv:2205.07894. https://arxiv.org/abs/2205.07894
- Perley, D. A. 2019, PASP, 131, 084503, doi: 10.1088/1538-3873/ab215d
- Perley, D. A., Mazzali, P. A., Yan, L., et al. 2019, MNRAS, 484, 1031, doi: 10.1093/mnras/sty3420
- Perley, D. A., Fremling, C., Sollerman, J., et al. 2020, ApJ, 904, 35, doi: 10.3847/1538-4357/abbd98

- Phillips, M. M., Simon, J. D., Morrell, N., et al. 2013, ApJ, 779, 38, doi: 10.1088/0004-637X/779/1/38
- Piro, A. L., Haynie, A., & Yao, Y. 2021, ApJ, 909, 209, doi: 10.3847/1538-4357/abe2b1
- Poznanski, D., Ganeshalingam, M., Silverman, J. M., & Filippenko, A. V. 2011, MNRAS, 415, L81, doi: 10.1111/j.1745-3933.2011.01084.x
- Poznanski, D., Prochaska, J. X., & Bloom, J. S. 2012, MNRAS, 426, 1465, doi: 10.1111/j.1365-2966.2012.21796.x
- Prentice, S. J., Maguire, K., Flörs, A., et al. 2020, A&A, 635, A186, doi: 10.1051/0004-6361/201936515
- Prochaska, J. X., Hennawi, J. F., Westfall, K. B., et al. 2020, Journal of Open Source Software, 5, 2308, doi: 10.21105/joss.02308
- Prochaska, J. X., Hennawi, J., Cooke, R., et al. 2020, pypeit/PypeIt: Release 1.0.0, v1.0.0, Zenodo, doi: 10.5281/zenodo.3743493
- Pursiainen, M., Leloudas, G., Schulze, S., et al. 2023, ApJL, 959, L10, doi: 10.3847/2041-8213/ad103d
- Renzo, M., Farmer, R., Justham, S., et al. 2020, A&A, 640, A56, doi: 10.1051/0004-6361/202037710
- Rigault, M., Neill, J. D., Blagorodnova, N., et al. 2019, A&A, 627, A115, doi: 10.1051/0004-6361/201935344
- Sagiv, I., Gal-Yam, A., Ofek, E. O., et al. 2014, AJ, 147, 79, doi: 10.1088/0004-6256/147/4/79
- Sandoval, M. A., Hix, W. R., Messer, O. E. B., Lentz, E. J., & Harris, J. A. 2021, ApJ, 921, 113, doi: 10.3847/1538-4357/ac1d49
- Sawada, R., Kashiyama, K., & Suwa, Y. 2022, ApJ, 927, 223, doi: 10.3847/1538-4357/ac53ae
- Schlafly, E. F., & Finkbeiner, D. P. 2011, ApJ, 737, 103, doi: 10.1088/0004-637X/737/2/103
- Shen, K. J., Kasen, D., Weinberg, N. N., Bildsten, L., & Scannapieco, E. 2010, ApJ, 715, 767, doi: 10.1088/0004-637X/715/2/767
- Shingles, L., Smith, K. W., Young, D. R., et al. 2021, Transient Name Server AstroNote, 7, 1
- Shiode, J. H., & Quataert, E. 2014, ApJ, 780, 96, doi: 10.1088/0004-637X/780/1/96
- Shivkumar, H., Jaodand, A. D., Balasubramanian, A., et al. 2023, ApJ, 952, 86, doi: 10.3847/1538-4357/acd5d5
- Shvartzvald, Y., Waxman, E., Gal-Yam, A., et al. 2023, arXiv e-prints, arXiv:2304.14482, doi: 10.48550/arXiv.2304.14482
- Smith, K. W., Smartt, S. J., Young, D. R., et al. 2020, PASP, 132, 085002, doi: 10.1088/1538-3873/ab936e
- Sollerman, J. 2002, NewAR, 46, 493, doi: 10.1016/S1387-6473(02)00189-6

—. 2023, Transient Name Server Discovery Report, 2023-3158, 1

Springob, C. M., Haynes, M. P., Giovanelli, R., & Kent, B. R. 2005, ApJS, 160, 149, doi: 10.1086/431550

Steele, I. A., Smith, R. J., Rees, P. C., et al. 2004, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 5489, Ground-based Telescopes, ed. J. Oschmann, Jacobus M., 679–692, doi: 10.1117/12.551456

Stockinger, G., Janka, H. T., Kresse, D., et al. 2020, MNRAS, 496, 2039, doi: 10.1093/mnras/staa1691

Stritzinger, M. D., Taddia, F., Burns, C. R., et al. 2018, A&A, 609, A135, doi: 10.1051/0004-6361/201730843

Taddia, F., Sollerman, J., Leloudas, G., et al. 2015, A&A, 574, A60, doi: 10.1051/0004-6361/201423915

Taddia, F., Stritzinger, M. D., Bersten, M., et al. 2018, A&A, 609, A136, doi: 10.1051/0004-6361/201730844

Tauris, T. M., Langer, N., Moriya, T. J., et al. 2013, ApJL, 778, L23, doi: 10.1088/2041-8205/778/2/L23

Tauris, T. M., Langer, N., & Podsiadlowski, P. 2015, MNRAS, 451, 2123, doi: 10.1093/mnras/stv990 Tonry, J. L., Denneau, L., Heinze, A. N., et al. 2018, PASP, 130, 064505, doi: 10.1088/1538-3873/aabadf

Turatto, M., Mazzali, P. A., Young, T. R., et al. 1998, ApJL, 498, L129, doi: 10.1086/311324

Valenti, S., Benetti, S., Cappellaro, E., et al. 2008, MNRAS, 383, 1485, doi: 10.1111/j.1365-2966.2007.12647.x

Wheeler, J. C., Johnson, V., & Clocchiatti, A. 2015, MNRAS, 450, 1295, doi: 10.1093/mnras/stv650

Wilkinson, D. M., Maraston, C., Goddard, D., Thomas, D., & Parikh, T. 2017, MNRAS, 472, 4297, doi: 10.1093/mnras/stx2215

- Wu, S. C., & Fuller, J. 2022, ApJL, 940, L27, doi: 10.3847/2041-8213/ac9b3d
- Yan, S., Wang, X., Gao, X., et al. 2023, ApJL, 959, L32, doi: 10.3847/2041-8213/ad0cc3
- Yao, Y., De, K., Kasliwal, M. M., et al. 2020, ApJ, 900, 46, doi: 10.3847/1538-4357/abaa3d

Yaron, O., & Gal-Yam, A. 2012, PASP, 124, 668, doi: 10.1086/666656

Zhang, W., Woosley, S. E., & Heger, A. 2008, ApJ, 679, 639, doi: 10.1086/526404

APPENDIX

A. TIMESCALE - LUMINOSITY PHASE SPACE

In Figure A1, we compare the time above half maximum and peak luminosity of SN 2023zaw with stripped-envelope supernovae and other fast-evolving transients in the literature with rise and decline time constraints.




Figure A1. We compare the time above half maximum and peak luminosity of SN 2023zaw with stripped-envelope supernovae from the BTS experiment (Perley et al. 2020). For sources with two peaks in the r-band lightcurve, the second peak is used. We also compare with other fast-evolving transients – SN 2019dge (Yao et al. 2020), SN 2019bkc (Prentice et al. 2020; Chen et al. 2020), iPTF14gqr (De et al. 2018b), iPTF16hgs (De et al. 2018c), AT 2018cow (Perley et al. 2019), ZTF18abvkwla or AT2023abbd (Ho et al. 2020).

B. HOST GALAXY EMISSION LINES

Table B1 shows the emission line flux measurements of the star-forming region at the SN site.

Table B1. Emission line flux measurements of the star-forming region at the SN site.

Transition	Flux		
	$(10^{-17} \mathrm{erg} \mathrm{cm}^{-2} \mathrm{s}^{-1})$		
$[O II] \lambda\lambda 3726,3729$	61.10 ± 5.23		
$[{\rm O~III}]\lambda4959$	4.63 ± 0.15		
$[{\rm O~III}]\lambda5007$	13.90 ± 0.46		
${ m H}\gamma$	14.83 ± 1.20		
${ m H}eta$	42.48 ± 2.51		
$[{\rm N~II}]\lambda6549$	26.04 ± 1.27		
$\mathrm{H}lpha$	203.98 ± 6.74		
$[{\rm N~II}]\lambda6584$	78.12 ± 3.81		
$[{\rm S~II}]\lambda6718$	40.84 ± 1.61		
$[{\rm S~II}]\lambda6732$	28.69 ± 1.43		

NOTE— No measurement is corrected for reddening.

Das et al.

C. PHOTOMETRY DATA

A truncated version of the photometry table is shown in Table C1. All the photometric and spectroscopic data used in this work will be made available on GitHub after publication.

Table C1. Truncated photometry data for SN 2023zaw. The full machine-readable version will be made available on GitHub after publication. The photometry data has been corrected for Milky Way extinction.

MJD	Phase ^a	Instrument	Filter	AB Magnitude
60285	-2	ATLAS	0	19.33 ± 0.26
60285	-2	P48	g	18.34 ± 0.06
60285	-2	ATLAS	0	18.16 ± 0.07
60285	-2	SEDM	r	18.09 ± 0.07
60285	-2	SEDM	g	18.23 ± 0.05
60285	-2	SEDM	r	18.08 ± 0.04
60285	-2	SEDM	i	17.91 ± 0.06
60286	-1	SEDM	r	17.72 ± 0.06
60286	-1	SEDM	g	17.98 ± 0.05
60286	-1	SEDM	r	17.78 ± 0.04
60286	-1	SEDM	i	17.65 ± 0.02
60286	-1	ATLAS	c	17.88 ± 0.05
60287	0	P48	g	17.96 ± 0.05
60287	0	P48	g	17.97 ± 0.04

^{*a*} Days since *r*-band peak (MJD = 60287.7).

D. SPECTROSCOPY COLLAGE

A collage all the spectra of SN 2023zaw is shown in Figure D1.




Figure D1. Spectral sequence for SN 2023zaw. The phases are in rest-frame days since the *r*-band peak (MJD = 60287.7). See Section 2.3 for details on the obtained spectra.

E. SPECTROSCOPY LOG

The spectroscopy log is shown in Table E1.

DAS ET AL.

Date	Phase ^a	Instrument
Dec 12 2023	-1.5	SEDM
Dec 08 2023	-0.6	SEDM
Dec 09 2023	0.0	ALFOSC
Dec 11 2023	+2.4	SEDM
Dec 12 2023	+2.5	KCWI
Dec 12 2023	+3.1	SEDM
Dec 13 2023	+4.0	GMOS
Dec 14 2023	+5.1	Binospec
Dec 17 2023	+8.0	LRIS
Jan 07 2024	+29.0	LRIS
Feb 07 2024	+61.0	LRIS
Dec 17 2023 Jan 07 2024	+8.0 +29.0	LRIS LRIS

Table E1. Spectroscopy log.

 a Days since r-band peak (MJD = 60287.7).

F. COLOR EVOLUTION PLOT

The g-r and r-i color evolution of SN 2023zaw compared to other SNe in the literature is shown in Figure F1.




Figure F1. The g - r (*left*) and r - i (*right*) color evolution of SN 2023zaw compared to other fast transients – SN 2005ek (Drout et al. 2013), SN 2019dge (Yao et al. 2020), iPTF14gqr (De et al. 2018b), iPTF14hgs (De et al. 2018c). The gray line shows the intrinsic color template for Type Ib SNe from Stritzinger et al. (2018).

G. LIGHTCURVE COMPARISON

In Figure G1, we compare the r-band lightcurve of SN 2023zaw with other fast-evolving transients in the literature.




Figure G1. The r-band lightcurve of SN 2023zaw compared with other fast-evolving transients in the literature – fast-declining Type I SNe – SN 2010X (Kasliwal et al. 2010), SN 2019bkc (Prentice et al. 2020; Chen et al. 2020), the kilonova AT 2017gfo (Abbott et al. 2017), ultra-stripped SN candidates – SN 2019dge (Yao et al. 2020), iPTF14gqr (De et al. 2018b), SN 2019wxt (Shivkumar et al. 2023; Agudo et al. 2023), SN 2022agco (Yan et al. 2023).

H. BLACKBODY BEST-FIT PARAMETERS

The summary of the best-fit blackbody properties can be found in Table H1.

blication.				
	Phase	Log Luminosity	Temperature	Radius
	(days)	(erg s^{-1})	(K)	(R_{\odot})
	-2	$42.40_{-0.08}^{+0.10}$	15620^{+1850}_{-1440}	3500^{+360}_{-360}
	$^{-1}$	$42.29_{-0.03}^{+0.03}$	11050^{+510}_{-490}	6180^{+340}_{-320}
	0	$42.24_{-0.02}^{+0.02}$	8960^{+500}_{-460}	8820^{+760}_{-700}
	2	$41.92^{+0.02}_{-0.02}$	6900^{+340}_{-330}	10270^{+860}_{-750}
	3	$41.80^{+0.02}_{-0.02}$	6710^{+300}_{-280}	9530^{+710}_{-670}
	4	$41.60^{+0.01}_{-0.01}$	5520^{+230}_{-210}	11190^{+880}_{-830}
	5	$41.42_{-0.02}^{+0.02}$	6100^{+380}_{-340}	7440^{+790}_{-720}
	6	$41.24_{-0.03}^{+0.02}$	5060^{+470}_{-430}	8780^{+1710}_{-1360}
	7	$41.17\substack{+0.02 \\ -0.02}$	5370^{+320}_{-290}	7140^{+870}_{-760}
	8	$41.16\substack{+0.07 \\ -0.08}$	5160^{+10}_{-10}	7640^{+590}_{-660}
	10	$41.00\substack{+0.04\\-0.05}$	5070^{+10}_{-10}	6630^{+310}_{-330}
	17	$40.70\substack{+0.04 \\ -0.04}$	5000^{+10}_{-10}	4830^{+220}_{-240}

Table H1. Summary of the blackbody properties for SN 2023zaw. A machine-readable version of this table will be made available on GitHub after publication.

I. COMPARISON OF ENVELOPE PROPERTIES

We compare the envelope properties of SN 2023zaw with USSNe candidates and Type Ibn SNe in Figure I1. We compare the envelope properties of SN 2023zaw with theoretical models for bound and unbound stellar material in Figure I2.




Figure I1. The envelope properties of SN 2023zaw (red cross) compared with double-peaked Type Ibc SNe (black squares) from Das et al. (2023a), USSN candidates (circles) – iPTF14gqr, SN 2019dge, SN 2019wxt, SN 2022agco (De et al. 2018b; Yao et al. 2020; Shivkumar et al. 2023; Agudo et al. 2023; Yan et al. 2023), and Type Ibn SNe (purple diamonds) – SN 2006jc, SN 2019wep, SN 2019up, SN 2012jpk, SN 2019deh, LSQ13ddu (Anupama et al. 2009; Pellegrino et al. 2022; Clark et al. 2020).




Figure I2. Left: Comparison of the envelope properties of SN 2023zaw with theoretical models for *bound* stellar material – binary and single star models from Wu & Fuller (2022) (WF22), Laplace et al. (2020) (L20) and Tauris et al. (2015) (T15). *Right:* Comparison of the the envelope properties of SN 2023zaw with theoretical models for *unbound* stellar material – late-time binary mass transfer (BMT; Wu & Fuller 2022), wave-driven mass loss (Leung et al. 2021; Shiode & Quataert 2014), pulsation-pair instability driven mass loss (PPI; Renzo et al. 2020; Leung et al. 2019).

J. PRIORS AND CORNER PLOTS

The priors and corner plots for shock-cooling emission and radioactive decay model fitting are shown in Tables J1, J2 and Figure J1.

Table J1. Priors used for shock-cooling emission model fitting described in Section 3.6.2.

Parameter	Prior
$\log R_{\rm ext}$	$\mathcal{U}(-5,25)$
$\log M_{\rm ext}$	$\mathcal{U}(-4,1)$
$E_{\rm ext,49}$	$\mathcal{U}(0.1, 100)$

Table J2. Priors used for nickel decay model fitting described in Section 3.6.2.

Parameter	Prior
$ au_{ m m}$	$\mathcal{U}(1,20)$
$\log M_{\rm Ni}$	$\mathcal{U}(-4,0)$
t_0	$\mathcal{U}(2,100)$




Figure J1. Corner plots showing the posterior constraints of the model parameters in the shock-cooling emission (left) and radioactive decay (right) models described in Section 3.6.2.