

# Quantized polarization in a generalized Rice-Mele model at arbitrary filling

Yasuhiro Tada<sup>1,2,\*</sup>

<sup>1</sup>*Quantum Matter Program, Graduate School of Advanced Science and Engineering, Hiroshima University, Higashihiroshima, Hiroshima 739-8530, Japan*

<sup>2</sup>*Institute for Solid State Physics, University of Tokyo, Kashiwa 277-8581, Japan*

We discuss the charge polarization in a generalized Rice-Mele model at arbitrary particle filling per site as a model of charge ordered systems in one dimension. The model possesses neither the conventional bond-centered inversion symmetry nor the one site translation symmetry alone, but has combinations of these symmetries. We show that the charge polarization in the ground state is quantized by the combined symmetry and is characterized solely by the filling. Especially, the polarization can be  $1/2 \pmod{1}$  in the zero filling limit. Under the open boundary condition, there exist excess charges accumulated near edges of the system irrespective of existence or absence of edge modes. Correspondingly, we decompose the polarization into a bulk contribution and an edge contribution, and numerically demonstrate that the polarization is dominated by the former (latter) when the energy gap is small (large). We also discuss a simple generalization of our model and examine absence/existence of a gapless edge mode protected by the inversion symmetry by introducing intra unit cell and inter unit cell contributions of the charge polarization.

## I. INTRODUCTION

An insulator can have polarization which characterizes asymmetry of a charge configuration. A classical example is the ionic insulator where positive point charges and negative point charges are alternatively distributed. Careful discussions are necessary for an appropriate definition of the polarization even for such a simple system under the periodic boundary condition. It is now widely accepted that the polarization can well be described by the Berry-Zak formula, which clarifies an intimate relationship of the polarization to topology and symmetry<sup>1-6</sup>. The celebrated prototypical model of dipole topological insulators is the Su-Schrieffer-Heeger (SSH) model, where the hopping integral has a staggered modulation<sup>7-9</sup>. The polarization of the SSH model is quantized by an inversion symmetry, and there have been extensive theoretical studies of the generalized SSH models and many interesting properties have been clarified<sup>10-19</sup>. Topological nature of the charge polarization is clearly seen in charge pumping where a change of the polarization in an adiabatic cycle in a parameter space is quantized by the Chern number<sup>20,21</sup>. Indeed, the Rice-Mele (RM) model<sup>22</sup> was experimentally realized in cold atoms and the Thouless pumping was observed<sup>23-25</sup>. In this way, the SSH and RM models are regarded as one of the most important models in the context of topological insulators.

The very basic physics behind the SSH and RM models is charge ordering due to the Peierls instability<sup>26,27</sup> as they were originally proposed for polyacetylene and diatomic polymers<sup>7,8,22</sup>. A trimerized variant of the SSH model was also studied with an application to the Peierls system TTF-TCNQ<sup>8</sup>, although related models have been discussed in relation to various systems such as engineered materials and cold atoms<sup>14-19</sup>. In a general one-dimensional metal with the Fermi wavenumber  $k_F$ , the charge susceptibility shows singularity at the wavenumber  $Q = 2k_F$  and the system has strong instability to-

ward a charge order. The Peierls phase transition and charge orders have been discussed in both organic materials and inorganic materials, and there are a variety of charge orders with different modulations  $Q$ <sup>26,27</sup>. In this context, the standard SSH and RM models correspond to the simplest charge order with  $Q = \pi$ . In spite of the extensive studies on the charge polarization, however, focus has been made mainly on this particular case and polarizations for other charge orders with general  $Q$ -vectors have not been well explored. Since there are a lot of charge ordered one-dimensional materials<sup>26,27</sup>, it is natural to consider generalization of the  $Q = \pi$  order to  $Q = 2k_F \neq \pi$  orders. Such generalization may be interesting also in the context of symmetry protected topological phases beyond the non-interacting fermions. For example, an  $S = 1/2$  trimer spin chain exhibits a  $1/3$  magnetization plateau implying a uniquely gapped ground state under a magnetic field, which can be regarded as a generalization of more familiar spin dimer orders<sup>28-30</sup>. Modulated spin systems could have “charge polarization” associated with the  $U(1)$  spin rotation symmetry similarly to charge ordered fermion systems<sup>31-35</sup>.

In this study, we investigate the charge polarization in a generalized RM model with a wavenumber  $Q = 2k_F$  at arbitrary filling per site  $\nu$ . The model has neither the conventional bond-centered inversion symmetry nor the one site translation symmetry, but has a combination of these symmetries. The combined symmetry leads to quantization of the charge polarization, and it is independent of the parameters and determined solely by the filling  $\nu$  in presence of the symmetry. Especially, the polarization can remain non-zero in the low density limit  $\nu \rightarrow 0$ . We also discuss excess charges accumulated near edges of the system under the open boundary condition. Correspondingly, the polarization can be decomposed into a bulk contribution and an edge contribution. It is found that these two contributions depend on parameters, while their sum remains constant under the symmetry. Finally, we investigate effects of a phase

shift in the modulation as a simple generalization of the model, where we introduce the intra unit cell and inter unit cell contributions of the charge polarization and examine absence/existence of gapless edge modes protected by the inversion symmetry.

## II. QUANTIZED CHARGE POLARIZATION UNDER SYMMETRY

It is well known that a one dimensional fermion system has the Peierls instability where the modulation period is determined by the Fermi wavevector or equivalently the particle density<sup>26,27</sup>. In this study, we consider fermion systems with a modulated hopping integral and a modulated on-site potential as a toy model for charge ordered insulators. Such a system can be described by a generalized Rice-Mele model for spinless fermions<sup>7-15,17,18,22</sup>,

$$H_0 = - \sum_j t_j (c_j^\dagger c_{j+1} + c_{j+1}^\dagger c_j) + \sum_j \Delta_j c_j^\dagger c_j, \quad (1)$$

$$t_j = t + \delta \cos(Qj), \quad \Delta_j = \Delta \cos(Qj), \quad (2)$$

where the sites are labeled as  $j = 0, 1, 2, \dots, L-1$ . See Fig. 1. The modulation wavenumber is given by  $Q = 2\pi\nu$ , where  $\nu = N/L = p/q$  is the particle filling per site with  $N$  being the total number of fermions and  $p, q$  being coprime integers ( $p$  fermions per  $q$ -site unit cell on average). The system size is assumed to be an integer multiple of the unit cell for both the periodic and open boundary conditions,  $L = L_q q$ , and is taken to be  $L = 600$  for numerical calculations in this study. The modulation could be induced by interactions or arise from different atom species in real materials, but here  $(\delta, \Delta)$  are model parameters of the system. The modulation factor  $\cos(Qj)$  reduces to the conventional factor  $(-1)^j$  at the half-filling  $\nu = 1/2$ . One can introduce a phase shift such as  $\cos(Q(j - j_0))$  with  $j_0 = 1, 2, \dots, q-1$  for which the corresponding Hamiltonian is related to the above  $H_0$  by the  $j_0$ -site translation (see Sec. IV). It is straightforward to numerically calculate single-particle spectra of Eq. (1) and there exists an energy gap  $\Delta E$  between the  $N$ -th and  $(N+1)$ -th single-particle energy eigenvalues under the periodic boundary condition as shown in Fig. 2. One can add interactions such as  $H_{\text{int}} = \sum_{ij} V_{ij} n_i n_j$  where  $n_j = c_j^\dagger c_j$  and  $V_{ij}$  keeps the symmetry of the system. A large part of our discussions including symmetry-based arguments presented below is applicable also to interacting systems, which will be important when one considers bosonic particles and spin systems. Note that our ground state is a uniquely gapped insulator with an integer filling per unit cell, which is distinct from interaction-driven degenerate ground states<sup>36</sup>.

Let us discuss symmetry of the Hamiltonian (1) and the resulting quantization of the charge polarization under the periodic boundary condition. Obviously, neither the one site translation symmetry  $T_x : c_j \rightarrow c_{j+1}$  nor the bond-centered inversion symmetry  $I_0 : c_j \rightarrow c_{L-j-1}$




FIG. 1. Schematic pictures of the system for (a) the periodic boundary condition and (b) the open boundary condition.  $I_0$  is the inversion about the bond  $\langle L-1, 0 \rangle$  and  $T_x$  is the one site translation. The open boundary condition is realized by cutting the hopping integral on the bond  $\langle L-1, 0 \rangle$  in the periodic boundary condition.




FIG. 2. Numerical results of the energy gap at (a)  $\delta = 0, \Delta = 0.5t$  and (b)  $\delta = 0.5t, \Delta = 0$  for the system under the periodic boundary condition.

about the bond  $\langle L-1, 0 \rangle$  alone is preserved for general  $Q$  when  $\delta$  and  $\Delta$  are non-zero. ( $\langle j, j+1 \rangle$  denotes the bond between the sites  $j$  and  $j+1$ .) However, one can see that appropriate combinations of these symmetries are preserved. Indeed, the Hamiltonian is invariant under the combined symmetry when either  $\delta$  or  $\Delta$  is zero,

$$I_k = (T_x)^k I_0 = \begin{cases} T_x I_0 & (\delta = 0, \Delta \neq 0), \\ T_x^2 I_0 & (\delta \neq 0, \Delta = 0). \end{cases} \quad (3)$$

Namely, the system has the  $I_1$ -symmetry (which is the site-centered inversion symmetry about the site  $j = 0$ ) when  $\delta = 0$  and  $I_2$ -symmetry (which is the bond-centered inversion symmetry about the bond  $\langle 0, 1 \rangle$ ) when  $\Delta = 0$ . This can easily be shown as follows. For the potential term,  $I_0 n_j I_0^{-1} = n_{L-j-1}$  and  $T_x^k n_j T_x^{-k} = n_{j+k}$ , which leads to  $T_x^k I_0 \sum \cos(Qj) n_j (T_x^k I_0)^{-1} = \sum \cos(Q(-j+k-1)) n_j$ . This implies  $Qj = Q(j-k+1)$  in modulo  $2\pi$  and hence  $k = 1 \pmod{q}$ . For the hopping term,  $I_0 c_j^\dagger c_{j+1} I_0^{-1} = c_{L-j-1}^\dagger c_{L-j-2}$  and therefore  $Qj = Q(j-k+2)$  in modulo  $2\pi$ , implying  $k = 2 \pmod{q}$ . Note that the  $I_k$ -symmetry is just an inversion symmetry about an appropriate site or bond translated by  $T_x^k$  from the original inversion center  $\langle L-1, 0 \rangle$ , but it turns out that clearly regarding  $I_k$ -symmetry as a combined symmetry is highly helpful. The inversion center dependence will be essentially important for a system under the open boundary condition. A generalization for shifted modulations and inversion center dependence will be discussed in Sec. IV.

The combined symmetries can put strong constraints

on the charge polarization as in the standard SSH model at half filling  $\nu = 1/2$  where it is quantized to either 0 or  $1/2$  by the bond-centered inversion symmetry  $I_0$ . Here we consider the charge polarization in the ground state  $|\Psi\rangle$  based on the Resta formula which is equivalent to the Berry-Zak formula in one dimension<sup>1,2,37,38</sup>,

$$P_x = \frac{1}{2\pi} \text{Im} \log \langle \Psi | U_x | \Psi \rangle, \quad (4)$$

$$U_x = \exp \left( \frac{2\pi}{L} \sum_j j(n_j - \nu) \right). \quad (5)$$

The summand  $(n_j - \nu)$  corresponds to the total electric charge at the site  $j$  including the uniform ion contribution  $-\nu$ . We note that the Resta formula itself is well-defined and can be used for both the periodic boundary condition and the open boundary condition. The operator  $U_x$  is often called Lieb-Schultz-Mattis twist operator<sup>31–33,39–41</sup> and its expectation value in a uniquely gapped ground state is known to be  $|\langle U_x \rangle| \simeq 1$  in one dimension. This is because the energy of the variational state  $|\Phi\rangle = U_x |\Psi\rangle$  converges to the ground state energy in the thermodynamic limit and therefore  $|\Phi\rangle$  must be essentially proportional to the ground state with a phase factor,  $|\Phi\rangle \simeq e^{i2\pi P_x} |\Psi\rangle$ , in a uniquely gapped system. See also Appendix A. Thus the Resta formula for the polarization is well-defined in one dimensional uniquely gapped ground state, although the expectation value is vanishing in the thermodynamic limit for general dimensions and an appropriate modification is necessary to resolve such a problem<sup>42,43</sup>.

The Lieb-Schultz-Mattis operator behaves under the symmetries as

$$I_0 U_x I_0^{-1} = U_x^{-1}, \quad (6)$$

$$T_x U_x T_x^{-1} = U_x e^{-i2\pi\nu}. \quad (7)$$

Note that these properties themselves are true for both the periodic boundary condition and the open boundary condition, since they are just algebraic relations of the operators and independent of a quantum state within the fixed filling Hilbert space. The above equations imply that, for the uniquely gapped ground state  $|\Psi\rangle$  which is an eigenstate of the combined operator  $I_k$  under the periodic boundary condition,

$$e^{2\pi i P_x} = \langle \Psi | U_x | \Psi \rangle = e^{2\pi i k\nu} \langle \Psi | U_x^\dagger | \Psi \rangle = e^{2\pi i(k\nu - P_x)}. \quad (8)$$

Therefore, the charge polarization is quantized as

$$2P_x = k\nu \pmod{1}. \quad (9)$$

$P_x$  has two kinds of branches, namely,  $P_x = k\nu/2 + n$  and  $P_x = -1/2 + k\nu/2 + n$  with  $n = 0, \pm 1, \pm 2, \dots$  as shown in Fig. 3. For example,  $P_x = +1/4$  and  $P_x = -1/4$  at  $\nu = 1/2$  for  $\delta = 0$  are inequivalent. Similarly,  $P_x = 0$  and  $P_x = 1/2$  at  $\nu = 1/2$  for  $\Delta = 0$  correspond to distinct

gapped states of the standard SSH model. We stress that Eq. (9) is valid in presence of an interaction under the  $I_k$ -symmetry as long as the total Hamiltonian also preserves the symmetry and the many-body energy gap does not close.




FIG. 3. The charge polarization quantized by (a)  $I_1$ -symmetry for  $\delta = 0, \Delta \neq 0$  and (b)  $I_2$ -symmetry for  $\delta \neq 0, \Delta = 0$ . The branches in the same color are equivalent, while those with different colors are inequivalent. Only the region  $-1 \leq P_x \leq 1$  is shown for simplicity.

Interestingly, the symmetry arguments claim that the polarization is  $P_x \rightarrow 1/2 \pmod{1}$  in the limit of zero-filling per site,  $\nu \rightarrow 0$ , for the blue-colored branches in Fig. 3. This may be counter-intuitive at first sight, but it does not contradict thermodynamics because charge polarization is *not* a thermodynamic quantity. The non-thermodynamic nature of the charge polarization is evident in a gedankenexperiment where the polarization is easily changed by an additional tiny amount of charges put on a surface of a system. Formally, there is no healthy thermodynamic limit under a uniform electric field as an intensive quantity which is conjugate to the polarization, because the energy cost due to the corresponding scalar potential is super-extensive similarly to the orbital angular momentum in superfluids<sup>44,45</sup>. Nevertheless, it is widely considered that a fractional part of the polarization is an intrinsic or “bulk” quantity, and therefore one might naively expect that it vanishes in the low density limit. This seemingly antagonistic problem can be resolved once one realizes that the charge polarization is not a standard thermodynamic observable but is a kind of a center of mass of the particles. The symmetry arguments simply mean that the center of mass in the present system can be  $O(1)$  at  $\nu = p/q \rightarrow 0$ , where there are  $p$  fermions in each  $q$ -site unit cell on average. Indeed, a single particle can have an  $O(1)$  center of mass in the unit cell of large  $q \gg 1$ . This can be confirmed by a straightforward calculation in the strong potential limit  $|\Delta| \gg t$  at  $\delta = 0$ , where the Hamiltonian is reduced to  $H = \sum \Delta \cos(Qj)n_j$ . The ground state of this Hamiltonian is simply given by a classical configuration of fermions which minimizes the potential  $\Delta \cos(2\pi p j/q)$ . We suppose  $p = 1$  and  $q \in 2\mathbb{Z}$  for simplicity, and then one fermion is localized at every  $j = 0 \pmod{q}$  in the ground

state when  $\Delta < 0$ . The polarization for this state is

$$\begin{aligned} P_x &= \frac{1}{2\pi} \text{Im} \log \exp \left( i \frac{2\pi}{L} \sum_{k=0}^{L_q-1} kq - i \frac{2\pi}{L} \sum_{j=0}^{L-1} j\nu \right) \\ &= \frac{1}{2} \left( \frac{L}{q} - 1 \right) - \frac{1}{2q} (L-1) \\ &= -\frac{1}{2} + \frac{\nu}{2} \pmod{1}. \end{aligned} \quad (10)$$

Note that the  $L$ -linear terms cancel out in the present charge neutral system, and the final result is fully consistent with Eq. (9). Similarly, one fermion is located at every  $j = q/2 \pmod{q}$  when  $\Delta > 0$ , and thus

$$P_x = \frac{\nu}{2} \pmod{1}. \quad (11)$$

We stress that these results obtained in the limit  $|\Delta| \rightarrow \infty$  are valid for general  $\Delta$  as long as the energy gap does not close as  $|\Delta|$  is decreased, because  $P_x$  is quantized by the  $I_k$ -symmetry.

The above symmetry arguments are easily confirmed by numerical calculations. We exemplify numerical results of the Resta formula in Fig. 4 (see Appendix B for technical details). They are fully quantized and characterized solely by the particle filling  $\nu$  under the  $I_k$ -symmetry in exact agreement with the symmetry arguments. As was shown above, the two inequivalent branches correspond to different signs of  $\delta$  and  $\Delta$ . The red-colored branches correspond to  $\Delta > 0$  in Fig. 4 (a) and  $\delta < 0$  in (b), while blue-colored branches to  $\Delta < 0$  in (a) and  $\delta > 0$  in (b).




FIG. 4. Numerical results of the charge polarization at (a)  $\delta = 0, |\Delta| = 0.5t$  and (b)  $|\delta| = 0.5t, \Delta = 0$ . The branches in the same color are equivalent, while those with different colors are inequivalent. Only the region  $-1 \leq P_x \leq 1$  is shown for simplicity.

In general, both of  $\delta$  and  $\Delta$  can be non-zero simultaneously. One can parameterize them as

$$\delta(\tau) = \delta_0 \cos \tau, \quad (12)$$

$$\Delta(\tau) = \Delta_0 \sin \tau. \quad (13)$$

We consider charge pumping during an adiabatic cycle  $\tau = 0 \rightarrow 2\pi$ . An example of the calculated polarization in this process is shown in Fig. 5. The pumped charge  $\Delta Q$  is quantized as

$$\Delta Q = \int_0^{2\pi} \frac{dP_x}{d\tau} d\tau = 1, \quad (14)$$

as expected from the general theory of the Thouless pumping<sup>20,21</sup>. This is a straightforward generalization of the Thouless pumping in the standard RM model at  $\nu = 1/2$  to that at arbitrary filling.




FIG. 5. Numerical demonstration of the charge pumping at  $\delta_0 = \Delta_0 = 0.5t$  calculated with the Resta formula for the system under the periodic boundary condition. The filling is  $\nu = 3/10$ .

### III. CHARGE POLARIZATION UNDER OPEN BOUNDARY CONDITION

In this section, we consider the model Eq. (1) with the sites  $j = 0, 1, \dots, L-1$  under the open boundary condition where there is no hopping between the right edge site  $j = L-1$  and the left edge site  $j = 0$  (Fig. 1). The inversion acts as  $I_0 : c_j \rightarrow c_{L-j-1}$ , as in the periodic boundary condition. In the standard SSH model ( $\delta \neq 0, \Delta = 0$ ) at  $\nu = 1/2$ , there can be gapless edge modes thanks to the chiral symmetry, but existence of such single-particle edge modes are not generally guaranteed in a system only with point group symmetry<sup>46,47</sup>. Similarly, there are no edge modes in the present system at almost all filling except for some special cases which will be discussed in Sec. IV. Nevertheless, one may naively expect that there are extra charges accumulated near edges of the system for a given filling. Here we introduce excess charges over one unit cell (length  $q$ ) compared to the uniform background charges,

$$\delta n_k = \sum_{j=kq}^{(k+1)q-1} (\langle n_j \rangle - \nu). \quad (15)$$

The excess charge  $\delta n_k$  corresponds to the total charge of electrons and ions contained in the  $k$ -th unit cell. Figure 6 shows examples of the fermion density profiles  $\langle n_j \rangle$  and the corresponding excess charges  $\delta n_k$ . Clearly, the excess charges vanish in the bulk region of the system, although the fermion density itself oscillates with the period  $q$ . This is a general property for a gapped periodic system which has unit cell translation symmetry. There are excess edge charges localized around the ends of the




FIG. 6. The fermion particle density  $\langle n_j \rangle$  and the excess charge  $\delta n_k$  for (a), (b)  $\delta = 0, \Delta = 0.5t$  and (c), (d)  $\delta = 0.5t, \Delta = 0$  for the system under the open boundary condition. The filling is  $\nu = 3/10$ . Note that  $\langle n_j \rangle$  and  $\delta n_k$  are not inversion symmetric since there is no inversion symmetry in the system.

system and they are quantified as

$$Q_{\text{left}} = \sum_{k=0}^{k_e} \delta n_k, \quad Q_{\text{right}} = \sum_{k=L_q-1-k_e}^{L_q-1} \delta n_k. \quad (16)$$

$k_e$  specifies an edge region of length  $O(1)$  and is supposed to be sufficiently larger than the correlation length of the system. In the numerical calculations, however, it is chosen as  $k_e = \lfloor L_q/4 \rfloor$  for numerical stability when the energy gap is small or equivalently the correlation length is large. We can numerically confirm  $Q_{\text{left}} + Q_{\text{right}} = 0$  holds because of the charge neutrality in the entire system, although  $\delta n_k$  itself is not inversion symmetric in general. (Typically,  $|Q_{\text{left}} + Q_{\text{right}}| = O(10^{-8 \sim -7})$  for the system size  $L = 600$ . The equation  $Q_{\text{left}} + Q_{\text{right}} = 0$  is satisfied in the thermodynamic limit,  $\lim_{k_e \rightarrow \infty} \lim_{L \rightarrow \infty} (Q_{\text{left}} + Q_{\text{right}}) = 0$ , because  $Q_k \simeq 0$  for the  $k$ -th unit cell in the bulk region in a gapped system.) We emphasize that the present excess edge charges are different from the (smeared or weighted) edge charges widely discussed in the literatures<sup>1,2,5,34,35</sup>. The smeared edge charge is given by a convolution of the charge density with a broadening function and it exactly equals the polarization. On the other hand, our excess charge only partly contributes to the polarization as will be discussed in the following. Our excess charge in the present lattice model would correspond to the extra charge in Bloch electron systems discussed in the previous study<sup>48</sup>. It may be regarded as a many-body variant of the previously introduced extra edge charge based on the trivial decomposition of the Resta formula.

Let us now discuss the polarization under the open boundary condition. There will be contributions to the polarization from the edge region with width  $O(1)$ , when the excess edge charges are non-zero. At the same time, microscopic charge configuration within a unit cell will also contribute to the polarization and such a contribu-

tion comes uniformly from a bulk region of the system. Correspondingly, we introduce a bulk contribution and an edge contribution under the open boundary condition by decomposing the polarization operator. The polarization operator is given by

$$\hat{P}_x = \frac{1}{L} \sum_{j=0}^{L-1} j(n_j - \nu) \quad (17)$$

and it is well-defined under the open boundary condition. For a state with the above mentioned charge configuration, the expectation value of this operator is simplified as

$$\begin{aligned} \langle \hat{P}_x \rangle &= \frac{1}{L} \sum_{k=0}^{L_q-1} \sum_{j=0}^{q-1} (kq + j) (\langle n_{kq+j} \rangle - \nu) \\ &\simeq \frac{1}{q} \sum_{j=0}^{q-1} j (\langle n_{kq+j} \rangle - \nu) \Big|_{k \gg 1} \\ &\quad + \frac{1}{L} \sum_{k \simeq 0, L_q} kq \sum_{j=0}^{q-1} (\langle n_{kq+j} \rangle - \nu). \end{aligned} \quad (18)$$

Therefore, the expectation value of the polarization operator is simply evaluated as

$$\langle \hat{P}_x \rangle \simeq P_{\text{bulk}} + P_{\text{edge}}, \quad (19)$$

$$P_{\text{bulk}} = \frac{1}{q} \sum_{j=j_b}^{j_b+q-1} j (\langle n_j \rangle - \nu), \quad (20)$$

$$P_{\text{edge}} = (Q_{\text{right}} - Q_{\text{left}})/2. \quad (21)$$

$j_b = nq (n \gg 1)$  defines a bulk region and is chosen as  $n = \lfloor L_q/4 \rfloor$  in the numerical calculations. This is an almost trivial decomposition and is valid for general systems with gapped bulk excitations, for which effects of a boundary condition are exponentially localized only around an edge. In such a system, the edge contribution can be evaluated under the periodic boundary condition as  $P_{\text{edge}} = P_x - P_{\text{bulk}}$  in modulo 1, where  $P_x$  and  $P_{\text{bulk}}$  are calculated under the periodic boundary condition (see also Sec. IV). Therefore, the excess edge charge is regarded as an intrinsic quantity of the system, which holds true not only in non-interacting models but also in interacting models.

In Fig. 7, we exemplify numerical results of the bulk and edge contributions when one of  $\delta$  and  $\Delta$  is zero. Each of the contributions depends on the parameters, but their sum remains constant and is quantized under the  $I_k$ -symmetry. We numerically see that

$$\langle \hat{P}_x \rangle \simeq P_{\text{bulk}} + P_{\text{edge}} \simeq P_x \pmod{1}, \quad (22)$$

where  $P_x$  is the polarization evaluated with use of the Resta formula under the open boundary condition. (As mentioned in the previous section, the Resta formula is well-defined both for the periodic boundary condition



and the open boundary condition.) Note that  $P_x$  under the open boundary condition numerically coincides with the one under the periodic boundary condition, which would be a general property as discussed in Appendix C. We confirm that typical difference between  $\langle \hat{P}_x \rangle$ ,  $(P_{\text{bulk}} + P_{\text{edge}})$ , and  $P_x$  is small ( $O(10^{-4} \sim -3)$  when  $L = 600$ ), and it decreases as the system size increases in the numerical calculations. Although the total polarization is a quantized constant under the symmetry, each of  $P_{\text{bulk}}$  and  $P_{\text{edge}}$  changes continuously with the parameters. The bulk contribution decreases as the modulation strength is reduced and approaches zero,  $P_{\text{bulk}} \rightarrow 0$ , when  $\delta \rightarrow 0$  or  $\Delta \rightarrow 0$ , because the system becomes a simple metal with a uniform bulk charge density in that limit. At the same time, the edge contribution increases and approaches the quantized value, which means that the excess edge charges  $Q_{\text{left/right}}$  become quantized in the weak modulation limit under the  $I_k$ -symmetry. Similar quantization was discussed previously, but the fundamental difference is that the edge charges in the previous studies are defined so that they are equal to the total polarization for any parameters<sup>1,2,5,34,35</sup>. In contrast to the weak modulation limit, the total polarization is dominated by the bulk contribution while the edge contribution is decreased when the modulation is large, because the system is deep in an insulating regime. In the strong modulation limit ( $|\delta| \rightarrow \infty$  or  $|\Delta| \rightarrow \infty$ ), the edge contribution vanishes,  $P_{\text{edge}} \rightarrow 0$ . These behaviors are generally seen except for special cases which will be discussed in the next section.




FIG. 7. The bulk (blue) and edge (red) contributions of the charge polarization for (a)  $\delta = 0$  and (b)  $\Delta = 0$  for the system under the open boundary condition. The filling is  $\nu = 3/10$ . The total polarization (green) is the numerical sum of these two contributions.

#### IV. GENERALIZATION AND BULK-BOUNDARY CORRESPONDENCE

The discussions on the model Eq. (1) in the previous sections can be generalized to some extent. In this section, we discuss a simple generalization of the model when either  $\delta$  or  $\Delta$  is zero and examine existence/absence of gapless edge modes protected by the inversion symmetry. We introduce a phase shift in the cosine term and focus on a “commensurate” phase shift  $\cos(Q(j - j_0))$  where  $j_0 = 1, 2, \dots, q - 1$  for the filling  $\nu = p/q$ . This modulation is also reduced to the standard staggered

factor  $(-1)^{j-j_0}$  at the half filling  $\nu = 1/2$ . Under the periodic boundary condition, the Hamiltonian  $H_{j_0}$  with  $\cos(Q(j - j_0))$  is related to  $H_0$  with  $\cos(Qj)$  by the  $j_0$ -site translation,

$$H_{j_0} = T_x^{j_0} H_0 T_x^{-j_0}. \quad (23)$$

This means that the Hamiltonian  $H_{j_0}$  has the  $I_{k+2j_0}$ -symmetry when  $H_0$  has the  $I_k$ -symmetry. Besides, the energy spectra for these Hamiltonians are unitary equivalent and especially the ground states are uniquely gapped under the periodic boundary condition. Thanks to the  $I_{k+2j_0}$ -symmetry, the charge polarization is quantized as

$$2P_x = (k + 2j_0)\nu \pmod{1}. \quad (24)$$

This is a natural result for the potential  $\Delta \cos(Q(j - j_0))$ , because the phase shift will change the positions of the potential minima by  $j_0$  sites and correspondingly the center of mass of the particles will move by  $j_0\nu$  to minimize the ground state energy. Note that the Hamiltonian has the  $q$ -site (one unit cell) translation symmetry and the center of mass comes to the position which is equivalent to the original position under the  $q$ -site translation. A similar thing will take place for the system with the phase shifted hopping modulation, although it is not easy to provide an intuitive picture.

Under the open boundary condition where there is no hopping integral on the bond  $(L - 1, 0)$ , Eq. (23) does not hold and the Hamiltonians  $H_{j_0}$  and  $H_0$  are not unitary equivalent. In this case, we numerically find that the energy spectrum of  $H_{j_0}$  becomes gapless at particular filling  $\nu = \nu^*$  implying existence of gapless edge modes as shown in Fig. 8. Alternatively, existence or absence of an gapless edge mode depends on the position of cutting a bond of the system (setting the hopping integral to zero) to realize an open boundary condition. The gap closing takes place when the filling satisfies the condition

$$P_x = \frac{1}{2} \pmod{1}, \quad (25)$$

or equivalently  $(k/2 + j_0)\nu^* = 0, 1/2 \pmod{1}$  depending on the parameters as shown in Fig. 9. This includes the well-known gapless edge mode in the topologically non-trivial state of the SSH model at  $\nu = 1/2$  ( $\delta < 0, \Delta = 0, j_0 = 0$  in our model). It is easy to see that, under this condition, the Hamiltonian  $H_{j_0}$  has the conventional inversion symmetry  $I_0$  about the bond  $(L - 1, 0)$ ,  $I_0 H_{j_0} I_0^{-1} = H_{j_0}$ . In presence of this inversion symmetry, the bulk contribution of the polarization is zero,  $P_x^{\text{bulk}} = (1/q) \sum_{j=L-j_b-1}^{L-j_b-1} (L - 1 - j) (\langle n_j \rangle - \nu) \simeq -P_x^{\text{bulk}} \simeq 0$ , because the summation of  $(\langle n_j \rangle - \nu)$  over one unit cell in the bulk should be exponentially small. Therefore, Eq. (25) simply means  $P_x = P_x^{\text{edge}} = \pm 1/2$ , which can be confirmed numerically as seen in Fig. 10. Strictly speaking, the inversion symmetric ground states under the open boundary condition are nearly degenerate and  $Q_{\text{left}} = Q_{\text{right}} = 0$  for each state due to the inversion




FIG. 8. Numerical results of the energy gap at (a)  $j_0 = 1, \delta = 0, \Delta = 0.5t$  and (b)  $j_0 = 2, \delta = 0.5t, \Delta = 0$  for the system under the open boundary condition. The gap closing takes place at (a)  $\nu^* = 1/3$  and (b)  $\nu^* = 1/4, 3/4$ .




FIG. 9. Numerical results of the charge polarization at (a)  $j_0 = 1, \delta = 0, |\Delta| = 0.5t$  and (b)  $j_0 = 2, |\delta| = 0.5t, \Delta = 0$  for the system under the open boundary condition. The solid lines are drawn with an emphasis on the gap closing for the eyes.

symmetry and charge neutrality.  $P_x^{\text{edge}} = \pm 1/2$  should be understood as either the limit  $\nu \nearrow \nu^*$  or  $\nu \searrow \nu^*$ .

The above discussions are consistent with the general argument for Bloch band systems in the previous study<sup>48</sup>. Here, we provide a brief explanation of the bulk-boundary correspondence based on the Resta formula for the many-body (interacting) ground state. Similarly to Eq. (18), we decompose the Lieb-Schultz-Mattis opera-




FIG. 10. The bulk (blue) and edge (red) contributions of the charge polarization for (a)  $j_0 = 1, \delta = 0, \Delta = 0.5t$  and (b)  $j_0 = 2, \delta = 0.5t, \Delta = 0$  for the system under the open boundary condition. The total polarization (green) is the numerical sum of these two contributions. The bulk contribution vanishes and  $P_x = P_x^{\text{edge}} = \pm 1/2$  holds (a) at  $\nu^* = 1/3$  and (b)  $\nu^* = 1/4, 3/4$ .

tor as  $U_x = U_{\text{intra}}U_{\text{inter}}$ , where

$$U_{\text{intra}} = e^{i2\pi\hat{P}_x^{\text{intra}}}, \quad \hat{P}_x^{\text{intra}} = \frac{1}{L_q} \sum_{k=0}^{L_q-1} \frac{1}{q} \sum_{l=0}^{q-1} l(n_{kq+l} - \nu), \quad (26)$$

$$U_{\text{inter}} = e^{i2\pi\hat{P}_x^{\text{inter}}}, \quad \hat{P}_x^{\text{inter}} = \frac{1}{L_q} \sum_{k=0}^{L_q-1} k \sum_{l=0}^{q-1} (n_{kq+l} - \nu). \quad (27)$$

We call them the intra unit cell contribution and inter unit cell contribution, respectively. Note that the bond  $\langle L-1, 0 \rangle$  is an inter unit cell bond under the periodic boundary condition and the sites  $j=0$  and  $j=L-1$  are the edges under the open boundary condition in the present study. The intra (inter) unit cell contribution under the periodic boundary condition corresponds to the bulk (edge) contribution  $P_{\text{bulk}}(P_{\text{edge}})$  under the open boundary condition. Indeed, it turns out that

$$P_{\text{intra}} = P_{\text{bulk}}, \quad P_{\text{inter}} = P_{\text{edge}}, \quad (28)$$

where  $P_a = \arg \langle \Psi | U_a | \Psi \rangle / 2\pi$  ( $a = \text{intra}, \text{inter}$ ). This is because the intra unit cell contribution has no fluctuations in a uniquely gapped ground state (see the discussion in the next paragraph and Appendix B) and thus  $P_{\text{intra}} = P_{\text{bulk}}$  just follows. Besides, the total polarization is independent of the boundary conditions (in modulo 1) as mentioned in the previous section and Appendix C, which means that  $P_{\text{inter}} = P_x - P_{\text{intra}} = \langle \hat{P}_x \rangle - P_{\text{bulk}} = P_{\text{edge}}$ .

The variational energy is  $\langle \Phi_a | H_{j_0} | \Phi_a \rangle = O(1/L)$  for each of the states  $|\Phi_a\rangle = \hat{U}_a |\Psi\rangle$  similarly to the original Lieb-Schultz-Mattis operator<sup>39,40</sup>. Correspondingly, deviations of the variational states from the uniquely gapped ground state under the periodic boundary condition are  $\|\delta\Psi_a\|^2 = O(1/L)$  when written as  $|\Phi_a\rangle = e^{i2\pi P_a} |\Psi\rangle + |\delta\Psi_a\rangle$  (see Appendix A). The ground state is almost an eigenstate of  $U_a$  in this sense, and thus the polarization is given by  $\langle U_x \rangle = \langle U_{\text{intra}} \rangle \langle U_{\text{inter}} \rangle + O(1/L)$ , where  $\langle \dots \rangle = \langle \Psi | \dots | \Psi \rangle$ . An important point is that  $\hat{P}_x^{\text{intra}}$  is a normalized extensive operator whose norm is  $O(1)$ , although  $\hat{P}_x^{\text{inter}}$  is super-extensive with the norm  $O(L)$ . As a result, there are no fluctuations of the intra unit cell contribution and  $\langle U_{\text{intra}} \rangle = e^{i2\pi P_{\text{intra}}} + O(1/L) \simeq 1$  holds, because  $(\partial/\partial\theta) \langle U_{\text{intra}}(\theta) \rangle = i \langle \hat{P}_x^{\text{intra}} \rangle \langle U_{\text{intra}}(\theta) \rangle + i \langle \Psi | \hat{P}_x^{\text{intra}} | \delta\Psi_{\text{intra}} \rangle = 0 + O(1/\sqrt{L})$  for the inversion symmetric ground state with  $\langle \hat{P}_x^{\text{intra}} \rangle = 0$ , where  $U_{\text{intra}}(\theta) = e^{i\theta\hat{P}_x^{\text{intra}}}$  (see also Appendix B). On the other hand, there can be non-negligible fluctuations for the inter unit cell contribution, since  $\hat{P}_x^{\text{inter}}$  is super-extensive. Therefore, the charge polarization is dominated by the inter unit cell contribution for the inversion symmetric uniquely gapped ground state in the thermodynamic limit,

$$\langle U_x \rangle = \langle U_{\text{inter}} \rangle, \quad P_x = P_{\text{inter}}. \quad (29)$$

The inter unit cell Lieb-Schultz-Mattis operator has been used in the previous study, where it was shown that a non-trivial expectation value  $\langle U_{\text{inter}} \rangle = -1$  in the inversion symmetric ground state (for an infinite system) implies existence of a gapless edge state in presence of an edge (for a semi-infinite system)<sup>33,41</sup>. Roughly speaking, a local variant of the inter unit cell Lieb-Schultz-Mattis operator  $U_{\text{loc}}(x)$  was introduced to characterize nature of a finite large region around a point  $x \in \mathbb{R}$ , where its expectation value is  $\pm 1$  under the inversion symmetry. The condition  $\langle U_{\text{loc}}(x) \rangle = -1$  for  $x$  in the bulk of the system together with the trivial condition  $\langle U_{\text{loc}}(x) \rangle = +1$  for  $x$  outside of the system implies that there must be gap closing around the edge of the system. This index theorem was rigorously proved as bulk-boundary correspondence for an infinite system without an edge and a semi-infinite system with an edge<sup>33,41</sup>, but we assume that it is applicable also to a finite large system with the periodic boundary condition and the same system with the open boundary condition. Thus, we can summarize the above argument on the bulk-boundary correspondence for a (interacting) uniquely gapped state as follows.

**Claim.** *In an inversion symmetric insulator, there exists a gapless edge state under the open boundary condition if the charge polarization is  $P_x = 1/2 \pmod{1}$  under the periodic boundary condition.*

Finally, we remark that the existence of a gapless (single-particle) edge mode or more generally degenerate ground states under the open boundary condition could also be understood as the filling anomaly<sup>43,49</sup>. We discuss the ground state wavefunctions when  $\delta = 0, \Delta \gg t$  in view of the filling anomaly to have an intuitive understanding. Let us consider a simple case with  $j_0 = 1$  for which  $P_x = 1/2$  is achieved when  $\nu = 1/3 (p = 1, q = 3)$ . There are degenerate potential minima at  $j = 0$  and  $j = 2 \pmod{3}$  in a unit cell and the middle site  $j = 1 \pmod{3}$  corresponds to the potential maximum. Here, we define a unit cell as the sites  $\{3k, 3k + 1, 3k + 2\}, k = 0, 1, 2, \dots, L_3 - 1$  for which the bonds  $\langle 3k - 1, 3k \rangle$  including  $\langle L - 1, 0 \rangle$  connect neighboring unit cells, where  $L_3 = L/3 \in \mathbb{Z}$ . The ground state under the periodic boundary condition is

$$|\Psi\rangle \simeq \frac{1}{\sqrt{2}}(c_{L-1}^\dagger + c_0^\dagger) \prod_{k=0}^{L_3-2} \frac{1}{\sqrt{2}}(c_{3k+2}^\dagger + c_{3k+3}^\dagger) |0\rangle. \quad (30)$$

A dimer  $(c_j^\dagger + c_{j+1}^\dagger) |0\rangle$  is formed by the hopping  $t > 0$  between neighboring unit cells, which is analogous to the standard SSH model with  $P_x = 1/2$  where there are dimers on inter unit cell bonds. (See Appendix D for direct calculations of the polarization in the ground state (30).) The local excited state on the dimer bond  $\langle j, j + 1 \rangle$  is  $(c_j^\dagger - c_{j+1}^\dagger) |0\rangle$  with an energy gap  $2t$  which has a different inversion eigenvalue from that of the ground state. Under the open boundary condition without a hopping integral on the bond  $\langle L - 1, 0 \rangle$ , the energy gap

between the two states  $(c_{L-1}^\dagger + c_0^\dagger) |0\rangle$  and  $(c_{L-1}^\dagger - c_0^\dagger) |0\rangle$  vanishes and the state

$$|\Psi'\rangle \simeq \frac{1}{\sqrt{2}}(c_{L-1}^\dagger - c_0^\dagger) \prod_{k=0}^{L_3-2} \frac{1}{\sqrt{2}}(c_{3k+2}^\dagger + c_{3k+3}^\dagger) |0\rangle \quad (31)$$

becomes another ground state. (To be precise, there is an exponentially small energy gap between these two ground states in a finite size system, but they are exactly degenerate in the thermodynamic limit.) We can also consider linear combinations of these states which explicitly break the inversion symmetry,  $|\Psi_{\text{left}}\rangle = (|\Psi\rangle - |\Psi'\rangle)/\sqrt{2}$  and  $|\Psi_{\text{right}}\rangle = (|\Psi\rangle + |\Psi'\rangle)/\sqrt{2}$ . The state  $|\Psi_{\text{left}}\rangle$  ( $|\Psi_{\text{right}}\rangle$ ) has a localized fermion at the left (right) edge  $j = 0 (j = L - 1)$ , corresponding to the excess edge charges  $Q_{\text{left/right}} = \pm 1/2$ . We stress that it is impossible to realize a uniquely gapped ground state under the open boundary condition with keeping the inversion symmetry and the fermion particle number. Essentially the same argument applies to other filling as well. This is a filling anomaly<sup>49</sup> and it can be discussed more rigorously with use of a polarization index<sup>43</sup>.

## V. DISCUSSION AND SUMMARY

We have studied the charge polarization in the generalized RM model in one dimension. Motivated by the various charge ordered materials, we introduced modulation wavenumber  $Q = 2\pi\nu$  with filling  $\nu$  per site corresponding to a charge order driven by the Peierls instability. The model has the combination of the bond-centered inversion and one site translation symmetries when either the hopping modulation or the potential modulation is zero. This symmetry leads to the quantization of the charge polarization. Especially, the charge polarization can be  $1/2 \pmod{1}$  at the low density limit  $\nu \rightarrow 0$ . These results are confirmed by the numerical calculations of the non-interacting model, and Thouless pumping at arbitrary filling was also demonstrated. Under the open boundary condition, there exist excess charges accumulated near the edges of the system. Correspondingly, the charge polarization can be decomposed into the bulk and edge contributions. The sum of these two contributions is a quantized constant under the combined symmetry, while each of them depends on the modulation strength. The polarization is dominated by the edge (bulk) contribution in the weak (strong) modulation limit. We also discussed the effects of a phase shift and examined the absence/existence of gapless edge modes or more generally degenerate ground states. Especially, we introduced the intra and inter unit cell contributions of the charge polarization in the Resta formula and clarified the role of the inversion symmetry.

A large part of our discussions including the symmetry argument is applicable to general uniquely gapped systems beyond the non-interacting fermions such as interacting fermions, bosons and spin systems. For exam-



ple, it would be interesting to study spin systems with large unit cells such as spin Peierls systems and trimer spin chains in view of the  $I_k$ -symmetry protected topological state. Our work could provide an insight for a further development in the understanding of the charge polarization.

### ACKNOWLEDGMENTS

We thank Ken Shiozaki for valuable comments and discussions. This work is supported by JSPS KAKENHI Grant No. 22K03513.

### Appendix A: The variational argument with the Lieb-Schultz-Mattis operator

We quickly review the Lieb-Schultz-Mattis argument in one dimension<sup>31–33,39–41</sup>. We consider the variational state  $|\Phi\rangle = U_x |\Psi\rangle$  for the uniquely gapped ground state  $|\Psi\rangle$ . The fermion creation operator behaves as  $U_x c_j^\dagger U_x^{-1} = e^{i2\pi j/L} c_j^\dagger$  and a straightforward calculation gives an estimate for the energy difference between these two states,

$$\langle \Psi | U_x^\dagger H_0 U_x | \Psi \rangle - \langle \Psi | H_0 | \Psi \rangle \leq \frac{C}{L}, \quad (\text{A1})$$

where  $C$  is a positive constant which is independent of  $L$ . From this inequality, one can obtain an inequality for the variational wavefunction as follows. We expand the variational state in terms of the normalized energy eigenstates  $|\Psi_n\rangle$  as  $|\Phi\rangle = z |\Psi\rangle + \sum_{n \geq 1} z_n |\Psi_n\rangle$ , where  $|z|^2 + \sum_{n \geq 1} |z_n|^2 = 1$  because  $U_x$  is a unitary operator. Then, Eq. (A1) is rewritten as

$$|z|^2 E_0 + \sum_{n \geq 1} |z_n|^2 E_n - E_0 \leq \frac{C}{L}, \quad (\text{A2})$$

where  $E_n$  is the  $n$ -th eigenenergy. We denote the energy gap between the ground state and the first excited state as  $E_1 - E_0 = \Delta E > 0$ . Clearly,  $E_n - E_0 \geq \Delta E$  for higher excited states. The above inequality immediately implies that the deviation of the variational state from the ground state,  $|\delta\Psi\rangle = \sum_{n \geq 1} z_n |\Psi_n\rangle$ , is vanishing in the thermodynamic limit as

$$\langle \delta\Psi | \delta\Psi \rangle = \sum_{n \geq 1} |z_n|^2 \leq \frac{C}{L \cdot \Delta E}. \quad (\text{A3})$$

Equivalently,  $|z|^2 = 1 - C/(L\Delta E)$  and the variational state is nearly proportional to the ground state,  $|\Phi\rangle = U_x |\Psi\rangle = e^{i2\pi P_x} |\Psi\rangle + |\delta\Psi'\rangle$  with  $|\delta\Psi'\rangle = (z - e^{i2\pi P_x}) |\Psi\rangle + |\delta\Psi\rangle$  and  $2\pi P_x = \arg z$ . This property is expressed as  $\langle \delta\Psi' | \delta\Psi' \rangle = O(1/L)$  for simplicity and is used in the main text and Appendix C.

### Appendix B: Details of numerical calculations of Resta formula

The Resta formula for the charge polarization can be easily evaluated for a non-interacting fermion system with use of the single-particle wavefunctions. Let  $\{u_{jn}\}_{n=0}^{L-1}$  be the eigenvectors of the single-particle Hamiltonian, where the eigenvalues are in the ascending-order. Then the many-body ground state is given by

$$|\Psi\rangle = \prod_{n=0}^{N-1} \gamma_n^\dagger |0\rangle, \quad (\text{B1})$$

where  $\gamma_n = \sum_j u_{jn}^* c_j$  are the single-particle modes diagonalizing the Hamiltonian. Using  $U_x c_j^\dagger U_x^{-1} = e^{i2\pi j/L} c_j^\dagger$ , we obtain

$$\begin{aligned} \langle \Psi | U_x | \Psi \rangle &= \langle \Psi | \prod_n (U_x \gamma_n^\dagger U_x^{-1}) U_x | 0 \rangle \\ &= \det \mathcal{U}_N \cdot e^{-2\pi i/L \sum_j j\nu}, \end{aligned} \quad (\text{B2})$$

where  $(\mathcal{U}_N)_{mn} = \sum_{j=0}^{L-1} u_{jm}^* u_{jn} e^{i2\pi j/L}$ ,  $(m, n = 0, 1, \dots, N-1)$ .

We remark on fluctuations of the polarization within the Resta formula. One can introduce a parameter  $\theta$  in the Resta formula as

$$P_x(\theta) = \frac{1}{2\pi} \text{Im} \log \langle \Psi | U_x(\theta) | \Psi \rangle, \quad (\text{B3})$$

$$U_x(\theta) = \exp \left( \frac{i\theta}{L} \sum_j j(n_j - \nu) \right). \quad (\text{B4})$$

It is straightforward to calculate  $\langle U_x(\theta) \rangle$  for general  $\theta$  similarly to the above. Then, higher order powers of the charge polarization can be obtained as

$$P_x^{(n)} = \left| \left\langle \frac{\partial^n U_x(\theta)}{\partial (i\theta)^n} \Big|_{\theta=0} \right\rangle \right|^{\frac{1}{n}}. \quad (\text{B5})$$

We can numerically evaluate the higher order powers by polynomial fitting of  $\langle U_x(\theta) \rangle$  as a function of  $\theta$ . Under the periodic boundary condition, we find that  $P_x^{(2n+1)} < P_x^{(2n+3)}$  and  $P_x^{(2n)} < P_x^{(2n+2)}$  respectively, and they all take different values. Therefore, fluctuations of the charge polarization seem to exist in the present insulator, which may be simply due to the fact that the operator  $\hat{P}_x$  is not a good operator under the periodic boundary condition. On the other hand, under the open boundary condition, we see that  $P_x^{(n)} \simeq |P_x(2\pi)|$  numerically holds for all  $n = 1, 2, \dots$ . Therefore, there are no fluctuations of the charge polarization in the present system similar to standard thermodynamic quantities, although the polarization itself is not a thermodynamic quantity. We note that, in the numerical calculations, the fluctuations under the periodic boundary condition arise only from the inter unit cell contribution  $\langle U_{\text{inter}} \rangle$  and the

intra unit cell contribution  $\langle U_{\text{intra}} \rangle$  does not have fluctuations, which is consistent with the general argument in the main text. Under the open boundary condition, there are no fluctuations in both contributions.

### Appendix C: Robustness of the charge polarization to boundary conditions

One would naively expect that the charge polarization  $P_x = \arg \langle \Psi | U_x | \Psi \rangle / 2\pi$  is robust to boundary conditions in an insulator. Indeed, we have numerically checked that  $P_x$  of the present non-interacting model does not change under the periodic boundary condition and the open boundary condition. It was shown previously that the expectation value of a local operator whose support is far away from the boundary is almost unchanged by boundary conditions in a gapped ground state based on the exponential decay of correlation functions<sup>50</sup>. However, the operator  $U_x$  is non-local and some discussions may be necessary for the robustness of  $\langle U_x \rangle$ .

Here, we provide an argument based on the locality of the boundary condition and gapped nature of the ground state by following the previous studies<sup>50-52</sup>, although it is not rigorous. We introduce a parameter  $\lambda \in [0, 1]$  in the hopping integral on the bond  $\langle L-1, 0 \rangle$  as  $t_{L-1} \rightarrow \lambda t_{L-1}$ , where  $\lambda = 0$  corresponds to the open boundary condition and  $\lambda = 1$  to the periodic boundary condition. It is assumed that there are no gapless edge states under the open boundary condition and the system is uniquely gapped for all  $\lambda \in [0, 1]$ . (If there exists a gapless edge state at  $\lambda = 0$ , one may introduce a small perturbation to gap out it and turn off the perturbation at the end.) Then, the ground state is almost an eigenstate of  $U_x$  and the variational state can be written as  $U_x |\Psi(\lambda)\rangle = e^{i2\pi P_x(\lambda)} |\Psi(\lambda)\rangle + |\delta\Psi(\lambda)\rangle$  with  $\| |\delta\Psi(\lambda)\rangle \|^2 = O(1/L)$  as mentioned in Appendix A. We have

$$\begin{aligned} \frac{d}{d\lambda} \langle \Psi | U_x | \Psi \rangle &= \left\langle \frac{d\Psi}{d\lambda} \left| U_x \right| \Psi \right\rangle + \left\langle \Psi \left| U_x \right| \frac{d\Psi}{d\lambda} \right\rangle \\ &\simeq e^{i2\pi P_x} \frac{d}{d\lambda} \langle \Psi | \Psi \rangle \\ &\quad + \left\langle \frac{d\Psi}{d\lambda} \left| \delta\Psi \right\rangle + \left\langle \delta\Psi \left| \frac{d\Psi}{d\lambda} \right\rangle, \quad (\text{C1}) \end{aligned}$$

where we have neglected the small error from  $|z - e^{i2\pi P_x}|$ . The first term in the above equation is zero. On the other hand, the inner product  $\langle \frac{d\Psi}{d\lambda} | \delta\Psi \rangle$  can be  $O(1)$  for a general change of the Hamiltonian, although  $\| |\delta\Psi \rangle \|^2 = O(1/L)$ . For example,  $\langle U_x \rangle$  can change when we vary the model parameters in the bulk region of the system such as the Thouless pumping as discussed in the main text. However, since the change of the Hamiltonian is local in the present setup, we can claim that the inner product is exponentially small under an assumption. Here, we suppose that there exist local operators such that  $|\langle \Psi_n | O_n^{(0)} | \Psi \rangle| \geq f_n > 0$ , where the distance between their supports  $\Omega_n^{(0)}$  and the boundary bond  $\langle L-1, 0 \rangle$

is  $O(L)$ . Namely, the excited states above the gap can be derived from local perturbations on the ground state. Then, it was shown that one can construct local operators  $O_n$  whose supports  $\Omega_n$  are finite and include those of  $O_n^{(0)}$ <sup>50-52</sup>. The excited states can be well approximated by these operators as  $|\Psi_n\rangle = O_n |\Psi\rangle / \|O_n |\Psi\rangle\|$ , where errors are exponentially small with respect to  $R = \text{dist}(\partial\Omega_n, \Omega_n^{(0)})$ .

Now, we write the Hamiltonian as  $H(\lambda) = H(0) + \lambda W$  with  $W = -t_{L-1}(c_{L-1}^\dagger c_0 + c_0^\dagger c_{L-1})$ . By differentiating  $H(\lambda) |\Psi(\lambda)\rangle = E_0(\lambda) |\Psi(\lambda)\rangle$ , we obtain

$$\left\langle \Psi_n \left| \frac{d\Psi}{d\lambda} \right\rangle = \frac{1}{E_n - E_0} \langle \Psi_n | \delta W | \Psi \rangle, \quad (\text{C2})$$

where  $\delta W = W - \langle \Psi | W | \Psi \rangle$ . Note that the energy difference is greater than the excitation gap,  $E_n - E_0 \geq \Delta E$ . Then the inner product is rewritten as

$$\left\langle \delta\Psi \left| \frac{d\Psi}{d\lambda} \right\rangle \simeq \sum_{n \geq 1} \frac{z_n}{E_n - E_0} \frac{\langle \Psi | O_n^\dagger \delta W | \Psi \rangle}{\|O_n |\Psi\rangle\|}. \quad (\text{C3})$$

It is known that, for local operators  $A, B$  with the distance  $d$  between their supports, the correlation function is exponentially small,  $|\langle \Psi | \delta A \delta B | \Psi \rangle| \leq \text{const} \cdot e^{-\text{const} \cdot d}$  in a gapped system<sup>53</sup>. From this exponential decay of correlation functions and  $0 = \langle \Psi_n | \Psi \rangle \simeq \langle \Psi | O_n^\dagger | \Psi \rangle / \|O_n |\Psi\rangle\|$ , we conclude that the numerator in the above equation is exponentially small in the system size  $L$  with an appropriately chosen  $\Omega_n$  such that  $\text{dist}(\Omega_n, \langle L-1, 0 \rangle) = O(L)$ . This implies that  $\frac{d}{d\lambda} \langle \Psi | U_x | \Psi \rangle \simeq 0$  and the charge polarization within the Resta formula is independent of the parameter  $\lambda$ . Especially, it does not change for the open boundary condition and periodic boundary condition,  $P_x(0) = P_x(1)$ . Further, the above argument is applicable also to general perturbations localized around the edges and hence the charge polarization is robust to them.

Finally, one might think that the robustness of the charge polarization will immediately follow from the previous study for local operators<sup>50</sup>, if one uses a local variant of the Lieb-Schultz-Mattis operator whose support is far away from the boundary bond<sup>41</sup>. We naively expect that expectation values of the global Lieb-Schultz-Mattis operator and the local one are equal under the periodic boundary condition. However, we are interested in the physical quantity  $P_{\text{edge}}$  where the edge sites are included in the operator support. In this case, the supports of the local Lieb-Schultz-Mattis operator and the boundary operator  $W$  are not separated and the robustness does not follow immediately.

### Appendix D: Direct calculation of polarization for Eq. (30)

We can directly calculate the polarization in the ground state Eq. (30) under the periodic boundary con-

dition. Since  $U_x c_j^\dagger U_x^{-1} = e^{i2\pi j/L} c_j^\dagger$ , the dimers behave as  $U_x (c_j^\dagger + c_{j+1}^\dagger) |0\rangle = e^{i2\pi(j+1/2)/L} (e^{-i\pi/L} c_j^\dagger + e^{i\pi/L} c_{j+1}^\dagger) U_x |0\rangle$ . Note that  $|\langle U_x \rangle| \rightarrow 1$  as  $L \rightarrow \infty$ , because  $\prod_{k=0}^{L_3-1} \cos(\pi/L) \rightarrow 1$ . The argument of the expectation value is

$$\frac{1}{2\pi} \arg \langle U_x \rangle = \frac{1}{L} \sum_{k=0}^{L_3-1} (3k + 5/2) - \frac{1}{L} \sum_{j=0}^{L-1} j\nu = \frac{1}{2}. \quad (\text{D1})$$

It is also straightforward to calculate  $P_{\text{bulk}}$  for the  $k$ -th unit cell (the sites  $j = 3k, 3k + 1, 3k + 2$ ),

$$\begin{aligned} 3P_{\text{bulk}} &= \left\langle \sum_{l=0}^2 (3k + l)(n_{3k+l} - 1/3) \right\rangle \\ &= \frac{1}{4} [3k \cdot 2 + (3k + 1) \cdot 0 + (3k + 2) \cdot 2] - (3k + 1) \\ &= 0. \end{aligned} \quad (\text{D2})$$

These results are fully consistent with the symmetry argument in the main text.

\* ytada@hiroshima-u.ac.jp

- <sup>1</sup> Raffaele Resta and David Vanderbilt, “Theory of polarization: A modern approach,” in *Physics of Ferroelectrics: A Modern Perspective* (Springer Berlin Heidelberg, Berlin, Heidelberg, 2007) pp. 31–68.
- <sup>2</sup> David Vanderbilt, *Berry Phases in Electronic Structure Theory* (Cambridge University Press, 2018).
- <sup>3</sup> Nicola A. Spaldin, “A beginner’s guide to the modern theory of polarization,” *Journal of Solid State Chemistry* **195**, 2–10 (2012).
- <sup>4</sup> R. D. King-Smith and David Vanderbilt, “Theory of polarization of crystalline solids,” *Phys. Rev. B* **47**, 1651–1654 (1993).
- <sup>5</sup> David Vanderbilt and R. D. King-Smith, “Electric polarization as a bulk quantity and its relation to surface charge,” *Phys. Rev. B* **48**, 4442–4455 (1993).
- <sup>6</sup> J. Zak, “Berry’s phase for energy bands in solids,” *Phys. Rev. Lett.* **62**, 2747–2750 (1989).
- <sup>7</sup> W. P. Su, J. R. Schrieffer, and A. J. Heeger, “Solitons in polyacetylene,” *Phys. Rev. Lett.* **42**, 1698–1701 (1979).
- <sup>8</sup> W. P. Su and J. R. Schrieffer, “Fractionally charged excitations in charge-density-wave systems with commensurability 3,” *Phys. Rev. Lett.* **46**, 738–741 (1981).
- <sup>9</sup> András Pályi János K. Asbóth, László Oroszlány, *A Short Course on Topological Insulators: Band-structure topology and edge states in one and two dimensions*, Lecture Notes in Physics, Vol. 919 (Springer, 2016).
- <sup>10</sup> Linhu Li, Zhihao Xu, and Shu Chen, “Topological phases of generalized su-schrieffer-heeger models,” *Phys. Rev. B* **89**, 085111 (2014).
- <sup>11</sup> Yuan Yao, Masahiro Sato, Tetsuya Nakamura, Nobuo Furukawa, and Masaki Oshikawa, “Theory of electron spin resonance in one-dimensional topological insulators with spin-orbit couplings: Detection of edge states,” *Phys. Rev. B* **96**, 205424 (2017).
- <sup>12</sup> Beatriz Pérez-González, Miguel Bello, Álvaro Gómez-León, and Gloria Platero, “Interplay between long-range hopping and disorder in topological systems,” *Phys. Rev. B* **99**, 035146 (2019).
- <sup>13</sup> Myles Scollon and Malcolm P. Kennett, “Persistence of chirality in the su-schrieffer-heeger model in the presence of on-site disorder,” *Phys. Rev. B* **101**, 144204 (2020).
- <sup>14</sup> Huaiming Guo and Shu Chen, “Kaleidoscope of symmetry-protected topological phases in one-dimensional periodically modulated lattices,” *Phys. Rev. B* **91**, 041402 (2015).
- <sup>15</sup> V. M. Martinez Alvarez and M. D. Coutinho-Filho, “Edge states in trimer lattices,” *Phys. Rev. A* **99**, 013833 (2019).
- <sup>16</sup> Md Nurul Huda, Shawulienu Kezilebieke, Teemu Ojanen, Robert Drost, and Peter Liljeroth, “Tuneable topological domain wall states in engineered atomic chains,” *npj Quantum Information* **5**, 17 (2020).
- <sup>17</sup> Maria Maffei, Alexandre Dauphin, Filippo Cardano, Maciej Lewenstein, and Pietro Massignan, “Topological characterization of chiral models through their long time dynamics,” *New Journal of Physics* **20**, 013023 (2018).
- <sup>18</sup> Dizhou Xie, Wei Gou, Teng Xiao, Bryce Gadway, and Bo Yan, “Topological characterizations of an extended su-schrieffer-heeger model,” *npj Quantum Information* **5**, 55 (2019).
- <sup>19</sup> Balázs Hetényi, Yetkin Pulcu, and Serkan Doğan, “Calculating the polarization in bipartite lattice models: Application to an extended su-schrieffer-heeger model,” *Phys. Rev. B* **103**, 075117 (2021).
- <sup>20</sup> D. J. Thouless, “Quantization of particle transport,” *Phys. Rev. B* **27**, 6083–6087 (1983).
- <sup>21</sup> Roberta Citro and Monika Aidelsburger, “Thouless pumping and topology,” *Nature Reviews Physics* **5**, 87–101 (2023).
- <sup>22</sup> M. J. Rice and E. J. Mele, “Elementary excitations of a linearly conjugated diatomic polymer,” *Phys. Rev. Lett.* **49**, 1455–1459 (1982).
- <sup>23</sup> M. Lohse, C. Schweizer, O. Zilberberg, M. Aidelsburger, and Immanuel Bloch, “A thouless quantum pump with ultracold bosonic atoms in an optical superlattice,” *Nature Physics* **12**, 350–354 (2016).
- <sup>24</sup> Shuta Nakajima, Takafumi Tomita, Shintaro Taie, Tomohiro Ichinose, Hideki Ozawa, Lei Wang, Matthias Troyer, and Yoshiro Takahashi, “Topological thouless pumping of ultracold fermions,” *Nature Physics* **12**, 296–300 (2016).
- <sup>25</sup> Shuta Nakajima, Nobuyuki Takei, Keita Sakuma, Yoshihito Kuno, Pasquale Marra, and Yoshiro Takahashi, “Competition and interplay between topology and quasi-periodic disorder in thouless pumping of ultracold atoms,” *Nature Physics* **17**, 844–849 (2021).
- <sup>26</sup> Seiichi Kagoshima, “Peierls phase transition,” *Japanese Journal of Applied Physics* **20**, 1617 (1981).
- <sup>27</sup> Sander van Smaalen, “The peierls transition in low-dimensional electronic crystals,” *Acta Cryst.* **A61**, 51–61 (2004).
- <sup>28</sup> Kazuo Hida, “Magnetic properties of the spin-1/2 ferromagnetic-ferromagnetic-antiferromagnetic trimerized heisenberg chain,” *Journal of the Phys-*

- ical Society of Japan **63**, 2359–2364 (1994), <https://doi.org/10.1143/JPSJ.63.2359>.
- <sup>29</sup> Haiping Hu, Chen Cheng, Zhihao Xu, Hong-Gang Luo, and Shu Chen, “Topological nature of magnetization plateaus in periodically modulated quantum spin chains,” *Phys. Rev. B* **90**, 035150 (2014).
- <sup>30</sup> Anup Kumar Bera, S. M. Yusuf, Sudip Kumar Saha, Manoranjan Kumar, David Voneshen, Yurii Skourski, and Sergei A. Zvyagin, “Emergent many-body composite excitations of interacting spin-1/2 trimers,” *Nature Communications* **13**, 6888 (2022).
- <sup>31</sup> Masaki Oshikawa, Masanori Yamanaka, and Ian Affleck, “Magnetization plateaus in spin chains: “haldane gap” for half-integer spins,” *Phys. Rev. Lett.* **78**, 1984–1987 (1997).
- <sup>32</sup> Masaaki Nakamura and Synge Todo, “Order parameter to characterize valence-bond-solid states in quantum spin chains,” *Phys. Rev. Lett.* **89**, 077204 (2002).
- <sup>33</sup> Hal Tasaki, “Topological phase transition and  $F_2$  index for  $s = 1$  quantum spin chains,” *Phys. Rev. Lett.* **121**, 140604 (2018).
- <sup>34</sup> Haruki Watanabe, Yasuyuki Kato, Hoi Chun Po, and Yukiotoshi Motome, “Fractional corner magnetization of collinear antiferromagnets,” *Phys. Rev. B* **103**, 134430 (2021).
- <sup>35</sup> Shunsuke C. Furuya and Masahiro Sato, “Quantized edge magnetizations and their symmetry protection in one-dimensional quantum spin systems,” *Phys. Rev. B* **104**, 184401 (2021).
- <sup>36</sup> A. A. Aligia and G. Ortiz, “Quantum mechanical position operator and localization in extended systems,” *Phys. Rev. Lett.* **82**, 2560–2563 (1999).
- <sup>37</sup> Raffaele Resta, “Quantum-mechanical position operator in extended systems,” *Phys. Rev. Lett.* **80**, 1800–1803 (1998).
- <sup>38</sup> Raffaele Resta and Sandro Sorella, “Electron localization in the insulating state,” *Phys. Rev. Lett.* **82**, 370–373 (1999).
- <sup>39</sup> D. Bohm, “Note on a theorem of Bloch concerning possible causes of superconductivity,” *Phys. Rev.* **75**, 502–504 (1949).
- <sup>40</sup> Elliott Lieb, Theodore Schultz, and Daniel Mattis, “Two soluble models of an antiferromagnetic chain,” *Annals of Physics* **16**, 407–466 (1961).
- <sup>41</sup> Hal Tasaki, “Rigorous index theory for one-dimensional interacting topological insulators,” *ArXiv:2111.07335*.
- <sup>42</sup> Haruki Watanabe and Masaki Oshikawa, “Inequivalent Berry phases for the bulk polarization,” *Phys. Rev. X* **8**, 021065 (2018).
- <sup>43</sup> Yasuhiro Tada and Masaki Oshikawa, “Many-body multipole index and bulk-boundary correspondence,” *Phys. Rev. B* **108**, 235150 (2023).
- <sup>44</sup> Yasuhiro Tada, Wenxing Nie, and Masaki Oshikawa, “Orbital angular momentum and spectral flow in two-dimensional chiral superfluids,” *Phys. Rev. Lett.* **114**, 195301 (2015).
- <sup>45</sup> Yasuhiro Tada, “Nonthermodynamic nature of the orbital angular momentum in neutral fermionic superfluids,” *Phys. Rev. B* **97**, 214523 (2018).
- <sup>46</sup> Hao Song, Sheng-Jie Huang, Liang Fu, and Michael Hermele, “Topological phases protected by point group symmetry,” *Phys. Rev. X* **7**, 011020 (2017).
- <sup>47</sup> Sheng-Jie Huang, Hao Song, Yi-Ping Huang, and Michael Hermele, “Building crystalline topological phases from lower-dimensional states,” *Phys. Rev. B* **96**, 205106 (2017).
- <sup>48</sup> Jun-Won Rhim, Jan Behrends, and Jens H. Bardarson, “Bulk-boundary correspondence from the intercellular Zak phase,” *Phys. Rev. B* **95**, 035421 (2017).
- <sup>49</sup> Wladimir A. Benalcazar, Tianhe Li, and Taylor L. Hughes, “Quantization of fractional corner charge in  $C_n$ -symmetric higher-order topological crystalline insulators,” *Phys. Rev. B* **99**, 245151 (2019).
- <sup>50</sup> Haruki Watanabe, “Insensitivity of bulk properties to the twisted boundary condition,” *Phys. Rev. B* **98**, 155137 (2018).
- <sup>51</sup> Jutho Haegeman, Spyridon Michalakis, Bruno Nachtergaele, Tobias J. Osborne, Norbert Schuch, and Frank Verstraete, “Elementary excitations in gapped quantum spin systems,” *Phys. Rev. Lett.* **111**, 080401 (2013).
- <sup>52</sup> Tohru Koma, “Topological current in fractional Chern insulators,” *ArXiv:1504.01243*.
- <sup>53</sup> Matthew B. Hastings and Tohru Koma, “Spectral gap and exponential decay of correlations,” *Communications in Mathematical Physics* **265**, 781–804 (2006).