CLASSIFICATION OF POSITIVE SOLUTIONS OF CRITICAL ANISOTROPIC SOBOLEV EQUATION WITHOUT THE FINITE VOLUME CONSTRAINT

LU CHEN AND YABO YANG

Abstract. In this paper, we classify all positive solutions of the critical anisotropic Sobolev equation

$$
-\Delta_p^H u = u^{p^*-1}, \quad x \in \mathbb{R}^n
$$

without the finite volume constraint for $n \geq 2$ and $\frac{(n+1)}{3} \leq p \lt n$, where $p^* = \frac{np}{n-p}$ denotes the critical Sobolev exponent and $-\Delta_p^H = -div(H^{p-1}(\cdot)\nabla H(\cdot))$ denotes the anisotropic p -Laplace operator. This result removes the finite volume assumption on the classification of critical anisotropic p-Laplace equation which was obtained by Ciraolo-Figalli-Roncoroni in the literature [\[8\]](#page-25-0). The method is based on constructing suitable vector fields integral inequality and using Newton's type inequality.

Keywords: Critical anisotropic Sobolev equation, Classification, Without finite volume constraint, Integral identity, Regularity

2020 MSC. 35J92, 35B33, 35B06.

1. INTRODUCTION

Given $n \geq 2$ and $1 < p < n$, the classical Sobolev inequality [\[29\]](#page-26-0) in \mathbb{R}^n states that for any $u \in W^{1,p}(\mathbb{R}^n)$, there holds

(1.1)
$$
\int_{\mathbb{R}^n} |u|^{p^*} dx \leq C(n, p, s) \int_{\mathbb{R}^n} |\nabla u|^p dx,
$$

where $p^* = \frac{np}{n-p}$ $\frac{np}{n-p}$ denotes the critical Sobolev exponent. Aubin [\[1\]](#page-25-1) and Talenti [\[33\]](#page-26-1) applied the technique of symmetry and rearrangement combining the Bliss Lemma to show that all radial extremals of Sobolev inequality must take the form as

$$
U = \left(1 + |x|^{\frac{p}{p-1}}\right)^{-\frac{n-p}{p}},
$$

up to some dilations and calculated the sharp constant of Sobolev inequality. However, they didn't classify all extremals of Sobolev inequality. Later, Erausquin, Nazaret and Villani [\[7\]](#page-25-2) show that all extremals must take the form as

$$
U = \left(1 + |x|^{\frac{p}{p-1}}\right)^{-\frac{n-p}{p}},
$$

The first author was partly supported by the National Key Research and Development Program (No. 2022YFA1006900) and National Natural Science Foundation of China (No. 12271027), the second author was supported by a grant from Beijing Institute of Technology (No. 2022CX01002).

2 LU CHEN AND YABO YANG

up to some dilation and translation by using the optimal transportation method. Obviously, the extremals of Sobolev inequality satisfy the critical Sobolev equation:

(1.2)
$$
\begin{cases}\n-\Delta_p u = u^{\frac{np}{n-p}-1}, & x \in \mathbb{R}^n \\
u \ge 0, & x \in \mathbb{R}^n \\
u \in W^{1,p}(\mathbb{R}^n).\n\end{cases}
$$

The classification of positive solutions to (1.2) in \mathbb{R}^n started in the crucial papers [\[17\]](#page-25-3) and [\[18\]](#page-25-4) and it has been the object of several studies. Damascelli-Merchán-Montoro-Sciunzi [\[11\]](#page-25-5) and Sciunzi [\[32\]](#page-26-2) established the symmetry of positive solutions of equation [\(1.2\)](#page-1-0), which together with Aubin and Talenti's results deduces the uniqueness of extremals of Sobolev inequality.

A natural problem is whether we can classify the positive solutions of critical Sobolev equations without the finite volume assumption. In fact, this is proved to be true by Caffralli, Giddas and Spruck [\[5\]](#page-25-6) applying Kelvin transform and moving plane method to classify all the positive solutions of the Yamabe equation [\[21\]](#page-25-7) when $p = 2$. We also note that Dai and Qin [\[12,](#page-25-8) [13,](#page-25-9) [14\]](#page-25-10) applied the method of scaling sphere to establish Liouvilletype theorem for Laplace operator or high-order Laplace operator involving polynomial nonlinearity in bounded or unbounded domain.

However, the Kelvin transform is not available for the general p -Laplace equation, hence the classification problem of critical Sobolev equation for $p \neq 2$ without the finite volume assumption is a challenging problem. Recently, Ou [\[28\]](#page-26-3) solved the classification problem under the assumption $\frac{n+1}{3} \leq p < n$. The same method has been also used successfully in the analogous problems on critical Sobolev and Hardy-Sobolev equation on Euclidean space and Heisenberg group (see [\[10\]](#page-25-11), [\[20\]](#page-25-12), [\[26\]](#page-26-4) and [\[36\]](#page-26-5)). However, the classification result for critical Sobolev equation without the finite volume assumption in the remaining index still keeps open.

Now, let us turn to the introduction of anisotropic Sobolev inequality. Anisotropic Sobolev inequality can be stated as follows: for any $u \in W^{1,p}(\mathbb{R}^n)$, there holds

(1.3)
$$
\left(\int_{\mathbb{R}^n} |u|^{p^*} dx\right)^{\frac{1}{p^*}} \leq C(n,p) \left(\int_{\mathbb{R}^n} H^p(\nabla v) dx\right)^{\frac{1}{p}},
$$

where H is a 1-homogenous convex function (see details in Section 2.1) and $C_{n,p}$ denotes the best possible constant which make the anisotropic Sobolev inequality hold. This sharp inequality was first obtained by Alvino, Ferone, Trombetti and Lions [\[2\]](#page-25-13) using the convex symmetrization technique. However, they did not solve the uniqueness problem of extremals of anisotropic Sobolev inequality. Ciraolo, Figalli and Roncoroni[[\[8\]](#page-25-0),Appendix A] solved the uniqueness problem by adapting the optimal transportation method. Furthermore, they proved that all positive solutions of anisotropic Sobolev equation with finite volume constraint

(1.4)
$$
\begin{cases}\n-\text{div } (a(\nabla u)) = u^{p^* - 1}, & x \in \mathbb{R}^n \\
u \ge 0, & x \in \mathbb{R}^n \\
\int_{\mathbb{R}^n} |u|^{\frac{np}{n-p}} dx < +\infty.\n\end{cases}
$$

must take the form as

$$
U_{\lambda}(x) = \left(\frac{(\lambda^{\frac{1}{p-1}}(n^{\frac{1}{p}}(\frac{n-p}{p-1})^{\frac{p-1}{p}})}{\lambda^{\frac{p}{p-1}} + H_0(x)^{\frac{p}{p-1}}}\right)^{\frac{n-p}{p}},
$$

where $a(\nabla u) = H^{p-1}(\nabla u) \nabla H(\nabla u)$, up to some translation and calculated the sharp constant of critical anisotropic Sobolev inequality. They provided a new approach based on an integral identity to classify all positive solutions and furthermore extended the classification result to the case of critical anisotropic Sobolev equation in convex cone. It should be noted that in the above research of the classification problem, the finite volume assumption plays an important role. However, the classification problem of critical anisotropic Sobolev equation without the finite volume constraint is still unknown.

In this paper, we are devoted to classified positive solutions of critical anisotropic Sobolev equation

(1.5)
$$
\begin{cases} -\text{div } (a(\nabla u)) = u^{p^* - 1} & x \in \mathbb{R}^n, \\ u \ge 0 & x \in \mathbb{R}^n. \end{cases}
$$

We are motivated by recent progress in Ma-Ou's work in [\[26\]](#page-26-4), Ma-Ou-Wu's work [\[27\]](#page-26-6) and Ou's work in [\[28\]](#page-26-3). Our main result states

Theorem 1.1. For $\frac{n+1}{3} \leq p < n$, assume that $u \in W^{1,p}_{loc}(\mathbb{R}^n)$ is a positive weak solution *of* [\(1.5\)](#page-2-0)*. Then* u *must take the form as*

(1.6)
$$
U_{\lambda}(x) = \left(\frac{(\lambda^{\frac{1}{p-1}}(n^{\frac{1}{p}}(\frac{n-p}{p-1})^{\frac{p-1}{p}})}{\lambda^{\frac{p}{p-1}} + H_0(x)^{\frac{p}{p-1}}}\right)^{\frac{n-p}{p}},
$$

up to some translation.

Remark 1.2. *To be precise, the proof of this Theorem [1.1](#page-2-1) is based on researching integral inequality involving the suitable vector fields. By constructing the correlation between the matrix* $W = \{W_{ij}\}\$ and vector fields and applying the integral inequality, we obtain that *the matrix is trace free, which could help us to classify all positive solutions of critical anisotropic Sobolev equation.*

Remark 1.3. *A function* $u \in W^{1,p}_{loc}(\mathbb{R}^n) \cap L^{\infty}_{loc}(\mathbb{R}^n)$ is said to be a weak solution of [\(1.5\)](#page-2-0) *if*

(1.7)
$$
\int_{\mathbb{R}^n} H^{p-1}(\nabla v) \nabla H(\nabla v) \cdot \nabla \psi dx - \int_{\mathbb{R}^n} u^{p^*-1} \psi = 0,
$$

for any $\psi \in W_0^{1,p}$ $i_{0}^{1,p}(\mathbb{R}^{n}).$

Here we mention some well-known facts about solutions of [\(1.5\)](#page-2-0), which are instrumental in the proof of Theorem [1.1.](#page-2-1) By the strong maximum principle, all nonnegative nontrivial solutions of [\(1.5\)](#page-2-0) must be strictly positive. In what follows, for any positive weak solution u of (1.5) , we have

(1.8)
$$
u \ge C(n, p, \min_{|x|=1} u)|x|^{-\frac{n-p}{p-1}} \quad \text{for } |x| > 1,
$$

4 LU CHEN AND YABO YANG

where C is denote as a general positive constant. In fact, the estimate (1.8) has been derived for positive weak sup-p-harmonic functions (see [\[8\]](#page-25-0)).

Organization of the paper: This paper is organized as follows. In section 2 we introduce some notations involving anisotropic norms and give a brief proof of regularity of solutions of critical anisotropic Sobolev equation. In Section 3, we construct suitable vector fields and solve the vital integral inequality which plays a very important role on classification of critical anisotropic Sobolev equation. In Section 4, we classify the positive solutions of critical anisotropic Sobolev equation without the finite volume constraint.

2. Preliminaries

In this section, we introduce some basic notations and properties about anisotropic norms and present the regularity of weak solutions of anisotropic equation.

2.1 Some basic properties of anisotropic norms: Let $H : \mathbb{R}^n \to \mathbb{R}$ is a norm such that H^2 is of class $C^2(\mathbb{R}\setminus\{0\})$ and it is uniformly convex. This fact is easily seen to be equivalent to the following three properties:

H is convex;(2.1)

(2.2)
$$
H(\xi) \ge 0 \text{ for } \xi \in \mathbb{R}^n \text{ and } H(\xi) = 0 \text{ if and only if } \xi = 0;
$$

(2.3)
$$
H(t\xi) = tH(\xi) \text{ for } \xi \in \mathbb{R}^n \text{ and for } t > 0.
$$

All norms in \mathbb{R}^n are equivalent. Hence, there exist positive constants C_1 and C_2 such that

(2.4)
$$
C_1|\xi| \le H(\xi) \le C_2|\xi|, \text{ for } \xi \in \mathbb{R}^n.
$$

Accordingly, H_0 denotes the dual norm to H given by

(2.5)
$$
H_0(\xi) = \sup_{\xi \neq 0} \frac{\xi \cdot \eta}{H(\xi)}, \quad \forall \eta \in \mathbb{R}^n.
$$

The following properties hold provided $H \in C^1(\mathbb{R}^n \setminus \{0\})$ (see [\[3\]](#page-25-14), section 2.2)

(2.6)
$$
H(\nabla_{\eta} H_0(\eta)) = 1, \quad H_0(\nabla_{\xi} H(\xi)) = 1,
$$

for every $\xi, \eta \in \mathbb{R}^n \setminus \{0\}$. Furthermore, the map $H \nabla_{\xi} H$ is invertible with

$$
(2.7) \t\t H\nabla_{\xi}H = (H_0\nabla_{\xi}H_0)^{-1}.
$$

From (2.6) and the homogeneity of H_0 , (2.7) is equivalent to

(2.8)
$$
H(\xi)\nabla_{\eta}H_0(\nabla_{\xi}H(\xi))=\xi.
$$

Sometimes we write

$$
\Delta_p^H u = -\text{div}\, (a(\nabla u)),
$$

in the sense of distributions, where Δ_p^H is called the *Finsler* p-Laplace(or anisotropic p-Laplace) operator and $a(\nabla u)$ is given by [\(1.4\)](#page-1-1). More precisely, [\(1.5\)](#page-2-0) reads as

$$
(2.9) \qquad \qquad -\Delta_p^H u = u^{p^*-1},
$$

where

$$
p^* = \frac{np}{n-p}.
$$

Lemma 2.1. *Assume that* H in $C^2(\mathbb{R}^n\setminus\{0\})$ *, it holds that*

(1)
$$
\sum_{i=1}^{n} H_i(\xi)\xi_i = H(\xi),
$$

\n(2)
$$
\sum_{i=1}^{n} H_{ij}(\xi)\xi_i = 0,
$$

\n(3)
$$
H_{ij}(t\xi) = \frac{1}{t}H_{ij}(\xi).
$$

Proof. Although we believe this result to be well-known to the experts (see [\[3\]](#page-25-14)) and other references, we still present all the details here just for the convenience of the reader. Using the equation [\(2.3\)](#page-3-2), we can derive that

(2.10)
$$
\sum_{i=1}^{n} H_i(\xi) \xi_i = \frac{d}{dt}|_{t=1} H(t\xi)
$$

$$
= H(\nabla u).
$$

Then the proof of statement (1) is accomplished. Furthermore, derivation the left side of (2.10) with respect to ξ_j gives

$$
\sum_{i=1}^n (H_i(\xi)\xi_i)_{\xi_j} = \sum_{i=1}^n (H_{ij}(\xi)\xi_i + H_i(\xi)(\xi_i)_{\xi_j}).
$$

From statement (1) , we obtain

$$
\sum_{i=1}^{n} (H_i(\xi)\xi_i)_{\xi_j} = (H(\xi))_{\xi_j} = H_j(\xi).
$$

Combining the above estimate, we deduce that

(2.11)
$$
\sum_{i=1}^{n} H_{ij}(\xi) \xi_i = 0,
$$

and accomplish the proof of statement (2).

Finally, we can proceed to compute the derivatives of both sides of the equation [\(2.3\)](#page-3-2) with respect to ξ_i , and then obtain that $H_i(t\xi) = H_i(\xi)$ (implies that $H_i(\xi)$ is 0-homogeneous). Thus, by a straightforward computation it is easy to show that

$$
(H_i(t\xi))_{\xi_j} = tH_{ij}(t\xi).
$$

On the other hand, since $H_i(\xi)$ is 0-homogeneous, we also have

$$
(H_i(t\xi))_{\xi_j} = (H_i(\xi))_{\xi_j}
$$

= $H_{ij}(\xi)$.

Then the proof of statement (3) is proved. \square

The regularity theory for Sobolev equation in divergence form, modeled upon the p-Laplacian, has extensively been developed in the past years (see [\[4,](#page-25-15) [6,](#page-25-16) [15,](#page-25-17) [16,](#page-25-18) [19,](#page-25-19) [22,](#page-25-20) [23,](#page-25-21) [24,](#page-25-22) [34,](#page-26-7) [35,](#page-26-8) [36\]](#page-26-5) and the references therein). However, we could not find a convenient literature for the results of regularity of anisotropic Sobolev equation, so for completeness and reader's convenience we present some results regarding regularity of anisotropic equation. Notice that Einstein summation convention of summation is used throughout the paper, we will omit the sum sign below.

2.2 Regularity of solutions of critical anisotropic Sobolev equation.

Proposition 2.2. For $1 < p < n$, let $u \in W^{1,p}_{loc}(\mathbb{R}^n)$ be any weak solution of [\(2.9\)](#page-4-1), then *u belongs to* $C^{1,\tau}_{loc}(\mathbb{R}^n)$ *for some* $\tau > 0$ *.*

Before giving the proof of Proposition [2.2,](#page-5-0) we first introduce the following Lemma [2.3,](#page-5-1) Lemma [2.4,](#page-6-0) Lemma [2.5](#page-6-1) from references $[31], [25]$ $[31], [25]$ and $[8]$.

Lemma 2.3 (See [\[31\]](#page-26-9), Theorem 1). *Let* u *be a weak solution of the equation*

$$
(2.12) \qquad \qquad \text{div}\mathcal{A}(x, u, \nabla u) = \mathcal{B}(x, u)
$$

defined in some bounded domain $\Omega \subseteq \mathbb{R}^n$. If $\mathcal{A}(x, u, \nabla u)$ and $\mathcal{B}(x, u)$ satisfies the following *condition:*

(2.13)
$$
\begin{cases} |\mathcal{A}| \le a |\nabla u|^{p-1}, & (2.13a) \\ |\mathcal{B}| \le b |u|^{p-1} + c, & (2.13b) \\ \nabla u \cdot \mathcal{A} \ge d |\nabla u|^p, & (2.13c) \end{cases}
$$

where $a, d \in \mathbb{R}^+$ and $b, c \in L^r(\Omega)$, with $r > \frac{n}{p}$, then $u \in L^{\infty}_{loc}(\Omega)$.

CRITICAL ANISOTROPIC SOBOLEV EQUATION WITHOUT THE FINITE VOLUME CONSTRAINTS7

Lemma 2.4 (See [\[25\]](#page-25-23), Theorem 1). *Let* u *be a weak solution of the equation*

(2.14)
$$
\operatorname{div} \mathcal{A}(x, u, \nabla u) = \mathcal{B}(x, u),
$$

defined in some bounded domain $\Omega \subseteq \mathbb{R}^n$. If $\mathcal{A}(x, u, \nabla u)$ and $\mathcal{B}(x, u)$ satisfies the following *condition:*

(2.15)
$$
\begin{cases} |\frac{\partial A_i}{\partial u_j} |\zeta_i \zeta_j \ge a |\nabla u|^{p-2} |\zeta|^2, & (2.15a) \\ |\frac{\partial A_i}{\partial u_j}| \le b |\nabla u|^{p-2}, & (2.15b) \\ |\mathcal{B}| \le c, & (2.15c) \end{cases}
$$

where a, b, c *be positive constants with* $b \geq a$ *. If* u *is a bounded weak solution of the* problem [\(2.14\)](#page-6-2)*, then* $u \in C_{loc}^{1,\tau}(\Omega)$ *for some* $\tau > 0$ *.*

Lemma 2.5 (See [\[8\]](#page-25-0), Proposition 2.8). *Let* u *be a local weak solution of equation* [\(2.9\)](#page-4-1) *with* a(·) given by [\(1.4\)](#page-1-1), where H satisfies the assumptions in subsection 2.1. Then $a(\nabla u) \in$ $W^{1,2}_{loc}(\mathbb{R}^n)$.

Proof of Proposition [2.2](#page-5-0)

Proof. We first show that $u \in L^{\infty}_{loc}(\mathbb{R}^n)$. From [\(2.4\)](#page-3-3) and Lemma [2.1,](#page-4-2) there exist constants a, d such that

$$
|H^{p-1}(\nabla u)\nabla H(\nabla u)| \le a|\nabla u|^{p-1},
$$

and

$$
|\nabla u \cdot H^{p-1}(\nabla u) \nabla H(\nabla u)| = |H^p(\nabla u)| \ge d|\nabla u|^p.
$$

Thus (2.13a) and (2.13c) have been proved. Thus, according to Lemma [2.3,](#page-5-1) we only need to verify the condition (2.13b) is applicative to obtain the desired result. Now, we rewrite the equation (2.9) satisfied by u as follows:

$$
-\text{div}(H^{p-1}(\nabla u)\nabla H(\nabla u)) = b|u|^{p-1} + c,
$$

where

(2.16)
$$
b = \begin{cases} 0 & \text{if } u < 1, \\ u^{p^* - p} & \text{if } u \ge 1, \end{cases}
$$

and

(2.17)
$$
c = \begin{cases} 0 & \text{if } u > 1, \\ u^{p^* - 1} & \text{if } u \le 1. \end{cases}
$$

Asin the proof of [[\[8\]](#page-25-0), Lemma 2.1], we know that $b \in L_{loc}^r(\mathbb{R}^n)$, with $r > \frac{n}{p}$ and $c \in$ $L^{\infty}(\mathbb{R}^n)$. This proves that u is bounded in $L^{\infty}_{loc}(\mathbb{R}^n)$.

Next, we start to show that $u \in C^{1,\tau}_{loc}(\mathbb{R}^n)$. According to Lemma [2.4,](#page-6-0) we need to prove that the conditions $(2.15a)-(2.15c)$ are applicative for the equation

$$
-\text{div}(H^{p-1}(\nabla u)\nabla H(\nabla u)) = u^{p^*-1}.
$$

We notice that H is positively homogeneous of degree 1 and H_i is homogeneous of degree 0 for all $i \in (1, n)$ from Lemma [2.1.](#page-4-2) Therefore, there exists a constant M_1 such that

$$
|H_i(\nabla u)| = \left| H_i(|\nabla u| \frac{\nabla u}{|\nabla u|}) \right|
$$

=
$$
\left| H_i(\frac{\nabla u}{|\nabla u|}) \right|
$$

$$
\leq \left| \max_{\xi \in \mathbb{S}^{n-1}} H_i(\xi) \right|
$$

$$
\leq M_1.
$$

Since $H \in C^2(\mathbb{R}^n \setminus \{0\}), H_{ij} \in L^{\infty}(\mathbb{S}^{n-1})$. Thanks to the Lemma [2.1](#page-4-2) (3), there exists a constant M_2 such that

$$
|H_{ij}(\nabla u)| = \left| H_{ij}(|\nabla u| \frac{\nabla u}{|\nabla u|}) \right|
$$

$$
= \left| \frac{1}{|\nabla u|} H_{ij}(\frac{\nabla u}{|\nabla u|}) \right|
$$

$$
\leq \frac{1}{|\nabla u|} \left| \max_{\xi \in \mathbb{S}^{n-1}} H_{ij}(\xi) \right|
$$

$$
\leq \frac{M_2}{|\nabla u|}.
$$

Recalling H^2 is uniformly convex and positive definite, there exists a constant C such that

(2.20)
\n
$$
(H^2)_{\xi_i\xi_j}\zeta_i\zeta_j = (2HH_i)_{\xi_j}\zeta_i\zeta_j
$$
\n
$$
= (H_iH_j + HH_{ij})\zeta_i\zeta_j
$$
\n
$$
\geq C|\zeta|^2.
$$

Inserting [\(2.20\)](#page-7-0) into (2.15a) and choosing $\varepsilon = \frac{p-1}{p-1}$ $\frac{p-1}{n-1}$, $a = \frac{p-1}{n-1}CC_1$, we obtain that for any $1 < p < n$ there holds

$$
\left| \frac{\partial (H^{p-1}(\nabla u)H_i(\nabla u))}{\partial u_j} \right| \zeta_i \zeta_j = \left| (p-1)H^{p-2}(\nabla u)H_i(\nabla u)H_j(\nabla u) \zeta_i \zeta_j + H^{p-2}(\nabla u)H(\nabla u)H_{ij}(\nabla u) \right| \zeta_i \zeta_j
$$
\n
$$
\geq \left| (p-1)H^{p-2}(\nabla u)H_i(\nabla u)H_j(\nabla u) \right| \zeta_i \zeta_j + \left| \varepsilon H^{p-2}(\nabla u)H(\nabla u)H_{ij}(\nabla u) \right| \zeta_i \zeta_j
$$
\n
$$
\geq \left| (p-1)H^{p-2}(\nabla u)H_i(\nabla u)H_j(\nabla u) \right| \zeta_i \zeta_j + \varepsilon \left| H^{p-2}(\nabla u) \right| C|\zeta|^2
$$
\n
$$
- \varepsilon \left| H^{p-2}(\nabla u)H_i(\nabla u)H_j(\nabla u) \right| \zeta_i \zeta_j
$$
\n
$$
= \frac{(n-2)(p-1)}{n-1} \left| H^{p-2}(\nabla u)H_i(\nabla u)H_j(\nabla u) \right| \zeta_i \zeta_j + \frac{p-1}{n-1} C \left| H^{p-2}(\nabla u) \right| |\zeta|^2
$$
\n
$$
\geq \frac{p-1}{n-1} C \left| H^{p-2}(\nabla u) \right| |\zeta|^2
$$
\n
$$
= a|\nabla u|^{p-2}|\zeta|^2,
$$

where we use $H_i H_j \zeta_i \zeta_j = (H_i \zeta_i)^2 \geq 0$ and $H_{ij} \zeta_i \zeta_j \geq 0$. A slight adjustment of constants shows that the statement (2.15a) must hold. Moreover, applying [\(2.18\)](#page-7-1) and [\(2.19\)](#page-7-2), we get

$$
\left| \frac{\partial (H^{p-1}(\nabla u)H_i(\nabla u))}{\partial u_j} \right| = |(p-1)H^{p-2}(\nabla u)H_i(\nabla u)H_j(\nabla u) + H^{p-2}(\nabla u)H(\nabla u)H_{ij}(\nabla u)|
$$

\n
$$
\leq (p-1)M_1^2 C_2 |\nabla u|^{p-2} + M_2 C_2^2 |\nabla u|^{p-2},
$$

\n
$$
\leq b |\nabla u|^{p-2},
$$

where $b = \max\{(p-1)M_1^2C_2, M_2C_2^2\}$, which ends the proof of (2.15b). The proof of the condition (2.15c) is easily modified by Lemma [2.3.](#page-5-1) Hence, applying the Lemma [2.4,](#page-6-0) we conclude that $u \in C^{1,\tau}_{loc}(\mathbb{R}^n)$. Recalling the definition of H, we obtain $H(\nabla u) \in C^{1,\tau}_{loc}(\mathbb{R}^n)$ and $\nabla H(\nabla u) \in C^{\tau}_{loc}(\mathbb{R}^n)$). $\qquad \qquad \Box$

3. Some differential identities and a key integral inequality on vector **FIELDS**

In this section, we need some preliminaries results before proving Theorem [1.1.](#page-2-1) More precisely, the key integral inequality [\(3.16\)](#page-11-0) plays a key role in proving Theorem [1.1.](#page-2-1) Hence our main goal in this section is to prove the key integral inequality [\(3.16\)](#page-11-0). Before presenting of [\(3.16\)](#page-11-0), we first define vector fields and show some lemmas that we need.

3.1 Definition of vector fields. Let $u > 0$ be any weak solution of [\(2.9\)](#page-4-1). By the regularity of solutions of critical anisotropic p-Laplace equation in part two, we take $v = u^{-\frac{p}{n-p}}$ and v satisfies, also in the weak sense

(3.1)
$$
-\Delta_p^H v = \frac{n(p-1)}{p} v^{-1} H^p(\nabla v) + \left(\frac{p}{n-p}\right)^{p-1} v^{-1} \text{ in } \mathbb{R}^n.
$$

Obviously, v inherits some properties from u. Notice that, applying the property (1.8) to u we deduce

(3.2)
$$
v \le C(n, p, \max_{|x|=1} v(x)) |x|^{\frac{p}{p-1}} \quad \text{for } |x| > 1.
$$

Recalling that we have already obtained $u \in C^{1,\tau}_{loc}(\mathbb{R}^n)$ in the previous Proposition [2.2,](#page-5-0) one could immediately deduce that

$$
(3.3) \t v \in C^{1,\tau}_{loc}(\mathbb{R}^n).
$$

Furthermore, the Proposition [2.2](#page-5-0) and Lemma [2.5](#page-6-1) are used to infer that

(3.4)
$$
H^{p-1}(\nabla v)\nabla H(\nabla v) \in W^{1,2}_{loc}(\mathbb{R}^n).
$$

Now we introduce the following vector fields

$$
\overrightarrow{\mathbf{v}}^i = H^{p-1}(\nabla v) H_i(\nabla v),
$$

$$
W_{ij} = \overrightarrow{\mathbf{v}}^i_{\cdot,j} - \frac{1}{n} \overrightarrow{\mathbf{v}}^h_{\cdot,h} \,\delta_{ij},
$$

and

$$
W_j = v^{-1} v_i W_{ij},
$$

where and in the sequel. With the help of [\(3.4\)](#page-8-0) and $C^{1,\alpha}$ regularity of $v, \vec{v}^i \in L^{\infty}_{loc}(\mathbb{R}^n)$ and $W_{i,j} \in L^2_{loc}(\mathbb{R}^n)$. Obviously, the matrix $W = W_{ij}$ is trace free, i.e., $\text{Tr}W = W_{ii} \equiv 0$, but may not be symmetric.

Denote the function

$$
g = \alpha v^{-1} H^p(\nabla v) + \beta v^{-1},
$$

with $\alpha = \frac{p-1}{n}$ $\frac{-1}{p}n, \beta = \left(\frac{p}{n-p}\right)^{p-1}$. Then v satisfies the equation

$$
\Delta_p v = \overrightarrow{\mathbf{v}}_{,k}^k = g \quad \text{in} \quad \mathbb{R}^n
$$

in the weak sense, that is

(3.5)
$$
-\int_{\mathbb{R}^n} \overrightarrow{v} \psi_k = \int_{\mathbb{R}^n} g \psi.
$$

Furthermore, direct calculation also gives

$$
W_{ij} = \overrightarrow{\mathbf{v}}^i_{\ j} - \frac{1}{n} g \delta_{ij}.
$$

3.2 Some fundamental differential identities. Before proving the key integral inequalities Lemma [3.3,](#page-11-1) we first give some fundamental differential identities below.

Lemma 3.1. *With the notations as in above, then we have*

(1)
$$
g_j = nv^{-1}v_iW_{ij} = nW_j
$$

\n(2) $W_{ij,i} = \frac{n-1}{n}g_j = (n-1)W_j$
\n(3) $(\vec{\nabla}^j W_{ij})_{,i} = W_{ij}W_{ji} + (n-1)\vec{\nabla}^j W_j = Tr\{W^2\} + (n-1)\vec{\nabla}^j W_j$

Proof. The proof of the statements go in the same way as the proof of Lemma 2.1 from [\[28\]](#page-26-3), we present statements here just for the convenience of the reader.

We first prove the statement (1). Careful computation gives

$$
v^{-1}v_iW_{ij} = v^{-1}v_i(\overrightarrow{\mathbf{v}}^i, -\frac{1}{n}\overrightarrow{\mathbf{v}}^k, k \,\delta_{ij})
$$

\n
$$
= v^{-1}v_i((H^{p-1}(\nabla v)H_i(\nabla v))_j - \frac{1}{n}g\delta_{ij})
$$

\n
$$
= v^{-1}v_i((p-1)H^{p-2}(\nabla v)H_j(\nabla v)H_i(\nabla v) + H^{p-1}(\nabla v)H_{ij}(\nabla v) - \frac{1}{n}g\delta_{ij})
$$

\n
$$
= (p-1)v^{-1}H^{p-1}(\nabla v)H_j(\nabla v) - \frac{1}{n}gv^{-1}v_j.
$$

Using the definition of g , we have

(3.7)
\n
$$
g_j = (\alpha v^{-1} H^p (\nabla v) + \beta v^{-1})_j
$$
\n
$$
= ((\alpha H^p (\nabla v) + \beta)_j v^{-1} + (\alpha H^p (\nabla v) + \beta)(v^{-1})_j)
$$
\n
$$
= \alpha p H^{p-1} (\nabla v) H_j (\nabla v) v^{-1} + (\alpha H^p (\nabla v) + \beta)(-v^{-2} v_j)
$$
\n
$$
= \frac{n(p-1)}{p} p H^{p-1} (\nabla v) H_j (\nabla v) v^{-1} - (\alpha v^{-1} H^p (\nabla v) + \beta v^{-1}) v^{-1} v_j
$$
\n
$$
= n(p-1) v^{-1} H^{p-1} (\nabla v) H_j (\nabla v) - g v^{-1} v_j.
$$

Hence, we find that

(3.8)
$$
g_j = nv^{-1}v_iW_{ij},
$$

that is the statement (1) holds true.

For the proof of (2), by applying the statement (1) and using the definition of W_j , we conclude that

(3.9)
\n
$$
W_{ij,i} = (\overrightarrow{\mathbf{v}}^i, _j - \frac{1}{n}g\delta_{ij})_i
$$
\n
$$
= \overrightarrow{\mathbf{v}}^i, _{ji} - \frac{1}{n}g_i\delta_{ij}
$$
\n
$$
= \frac{n-1}{n}g_j
$$
\n
$$
= (n-1)W_j.
$$

Hence, statement (2) holds true. By virtue of the statement (1) , (2) we notice that W is trace free and yields

(3.10)
\n
$$
(\overrightarrow{\mathbf{v}}^{j}W_{ij})_{,i} = \overrightarrow{\mathbf{v}}^{j}_{,i}W_{ij} - \overrightarrow{\mathbf{v}}^{j}W_{ij,i}
$$
\n
$$
= W_{ij}W_{ij} + (n-1)\overrightarrow{\mathbf{v}}^{j}W_{j}
$$
\n
$$
= Tr\{W^{2}\} + (n-1)\overrightarrow{\mathbf{v}}^{j}W_{j}.
$$

Thus, the statement (3) holds true. \Box

Lemma 3.2.

(3.11)
$$
(v^q g^m \vec{v}^j)_{,j} = (q+a)v^{q-1} g^m H^p(\nabla v) + bv^{q-1} g^m + nmv^q g^{m-1} W_j \vec{v}^j
$$

and

$$
(3.12) \quad (v^q g^m \overrightarrow{\mathbf{v}}^j E_{ij})_{,i} = v^q g^m Tr\{E^2\} + (q+n-1)v^q g m \overrightarrow{\mathbf{v}}^j E_{ij} + nm E_i g^{m-1} v^q \overrightarrow{\mathbf{v}}^j E_{ij}
$$

Proof. First we prove that

$$
(3.13)
$$

\n
$$
(v^q g^m \vec{\mathbf{v}}^j)_{,j} = q v^{q-1} v_j g^m \vec{\mathbf{v}}^j + m v^q g^{m-1} g_j \vec{\mathbf{v}}^j + v^q g^m \vec{\mathbf{v}}^j_{,j}
$$

\n
$$
= q v^{q-1} g^m v_j H^{p-1} (\nabla v) H_j (\nabla v) + m v^q g^{m-1} (n W_j) \vec{\mathbf{v}}^j + v^q g^m (a v^{-1} H^p (\nabla v) + b v^{-1})
$$

\n
$$
= q v^{q-1} g^m H^p (\nabla v) + v^{q-1} g^m a H^p (\nabla v) + v^q g^m b v^{-1} + n m v^q g^{m-1} W_j \vec{\mathbf{v}}^j
$$

\n
$$
= (q + a) v^{q-1} g^m H^p (\nabla v) + b v^{q-1} g^m + n m v^q g^{m-1} W_j \vec{\mathbf{v}}^j.
$$

Furthermore, using the statement [\(3.11\)](#page-11-2) and the statement (3) of Lemma 2.1, we obtain (3.14)

$$
(v^q g^m \overrightarrow{\mathbf{v}}^j W_{ij})_{,i} = (v^q g^m)_i \overrightarrow{\mathbf{v}}^j W_{ij} + v^q g^m (\overrightarrow{\mathbf{v}}^j W_{ij})_i
$$

\n
$$
= (qv^{q-1} v_i g^m + mv^q g^{m-1} g_i) \overrightarrow{\mathbf{v}}^j W_{ij} + v^q g^m (Tr{W^2} + (n-1) \overrightarrow{\mathbf{v}}^j W_j)
$$

\n
$$
= v^q g^m Tr{E^2} + (q+n-1) v^q g^m \overrightarrow{\mathbf{v}}^j W_j + nm W_i g^{m-1} v^q \overrightarrow{\mathbf{v}}^j W_{ij}.
$$

3.3 The key integral inequality. In this subsection, we will prove the key integral inequality [\(3.16\)](#page-11-0) which plays a vital role in proving Theorem [1.1.](#page-2-1) Let ρ be a smooth cut-off function satisfying:

(3.15)
$$
\begin{cases} \rho \equiv 1 & \text{in } B_R \\ 0 \le \rho \le 1 & \text{in } B_{2R} \\ \rho \equiv 0 & \text{in } \mathbb{R}^n \setminus B_{2R} \\ |\nabla \rho| \lesssim \frac{1}{R} & \text{in } \mathbb{R}^n \end{cases}
$$

where and in the sequel. Moreover we use " \leq "," \leq " to replace " \leq ", " = ", etc., to drop out some positive constants independent of R and v .

Lemma 3.3. *Let* u *be any weak solution of* [\(2.9\)](#page-4-1) *and using the notations as before, then for every* $0 \leq \varphi \in C_c^{\infty}(\mathbb{R}^n)$ *we have*

$$
(3.16)\quad \int_{\mathbb{R}^n} \varphi v^{1-n} g^m Tr\{W^2\} + nm \int_{\mathbb{R}^n} \varphi v^{1-n} g^{m-1} \overrightarrow{\mathbf{v}}^j W_{ij} W_i \leq - \int_{\mathbb{R}^n} v^{1-n} g^m \overrightarrow{\mathbf{v}}^j W_{ij} \varphi_i
$$

In order to prove the Lemma [3.3,](#page-11-1) we first introduce the following Lemma [3.4.](#page-12-0)

Lemma 3.4. *Let* $1 < p < n$, $\gamma \in \mathbb{R}$ *and u be a positive weak solution of [1.1.](#page-2-1) Then for* $every\ 0 \leq \phi \in C_0^{\infty}(\mathbb{R}^n)$ *we have*

(3.17)
$$
\int_{\mathbb{R}^n} (u^{\gamma} I(x) + \psi) \leq - \int_{\mathbb{R}^n} \omega \cdot \nabla \phi.
$$

where

$$
\omega := u^{\gamma}(\overrightarrow{\mathbf{v}}\cdot\nabla\overrightarrow{\mathbf{v}} - \frac{1}{n}\overrightarrow{\mathbf{v}}\mathrm{div}\overrightarrow{\mathbf{v}}),
$$

and

$$
\psi = u^{\gamma + 2\alpha - 1} \left[\left(1 - \frac{np}{n-p} \right) \gamma g(u) - \frac{n-1}{n} u g'(u) \right] H^p(\nabla u) - \left(\frac{p-1}{p} (\gamma + 2\alpha - 1) \gamma + \frac{n-1}{n} \alpha^2 \right) u^{\gamma + 2\alpha - 2} H^{2p}(\nabla u)
$$

$$
+ \left(\frac{n-1}{n} \alpha + \frac{p-1}{p} \gamma \right) \operatorname{div} \left(u^{\gamma + 2\alpha - 1} \vec{u} H^p(\nabla u) \right).
$$

Proof. Put

(3.18)
$$
\vec{\mathbf{u}} = H^{p-1}(\nabla u)\nabla H(\nabla u), \quad \vec{\mathbf{v}} = u^{\alpha}\vec{\mathbf{u}} = u^{\alpha}H^{p-1}(\nabla u)\nabla H(\nabla u).
$$

Then according to Lemma [2.1,](#page-4-2) we have

(3.19)
$$
\overrightarrow{\mathbf{u}} \cdot \nabla u = H^p(\nabla u), \quad \overrightarrow{\mathbf{v}} \cdot \nabla u = u^{\alpha} H^p(\nabla u), \text{ div } \overrightarrow{\mathbf{u}} = -g(u).
$$

Moreover, since $\overrightarrow{\mathbf{v}} \in W_{loc}^{1,2}(\Omega)$ from section 2, we still have

(3.20)
\n
$$
\nabla \vec{\mathbf{v}} = \alpha u^{\alpha-1} H^{p-1}(\nabla u) \nabla H(\nabla u) \bigotimes \nabla u + u^{\alpha} \nabla \vec{\mathbf{u}} \quad \text{div } \vec{\mathbf{v}} = u^{\alpha-1} [\alpha H^p(\nabla u) - ug(u)],
$$

where $\overrightarrow{U} = \nabla \overrightarrow{u}$, $\overrightarrow{V} = \nabla \overrightarrow{v}$ are the Jacobian matrices of \overrightarrow{u} , \overrightarrow{v} . Finally, with standard matrix multiplication notation and Lemma [2.1,](#page-4-2)

(3.21)
\n
$$
\overrightarrow{U}\nabla u = (\nabla \overrightarrow{u})\nabla u
$$
\n
$$
= \nabla (H^{p-1}(\nabla u)\nabla H(\nabla u))\nabla u
$$
\n
$$
= (p-1)H^{p-2}(\nabla u)\nabla H(\nabla u)\nabla^2 u\nabla H(\nabla u)\nabla u + H^{p-1}(\nabla u)\nabla^2 H(\nabla u)\nabla^2 u\nabla u
$$
\n
$$
= (p-1)H^{p-1}(\nabla u)\nabla H(\nabla u)\nabla^2 u
$$
\n
$$
= \frac{p-1}{p}\nabla (H^p(\nabla u)).
$$

Furthermore, we define

$$
I(x) = |\overrightarrow{V} - \frac{1}{n} Tr{\overrightarrow{V} \cdot I_{d_n}}|^2,
$$

14 LU CHEN AND YABO YANG

where I_{d_n} is the identity tensor. Note that $I(x) \in L^1(\Omega)$ since $\overrightarrow{V} \in L^2(\Omega)$.

Now we are in position to prove Lemma [3.4.](#page-12-0) For the right side of the [\(3.17\)](#page-12-1) and the definition of ω , we have

$$
-\int_{\mathbb{R}^n} \omega \cdot \nabla \phi = -\int_{\mathbb{R}^n} u^\gamma (\vec{\mathbf{v}} \cdot \nabla \vec{\mathbf{v}} - \frac{1}{n} \vec{\mathbf{v}} \operatorname{div} \vec{\mathbf{v}}) \nabla \phi
$$

$$
\equiv I_0 + I_1.
$$

For the second term I_1 , combining with $(3.16)-(3.21)$ $(3.16)-(3.21)$, we conclude that

$$
(3.22)
$$
\n
$$
I_{1} = \frac{1}{n} \int_{\mathbb{R}^{n}} u^{\gamma} (\text{div } \vec{\mathbf{v}}) \vec{\mathbf{v}} \nabla \phi
$$
\n
$$
= -\frac{1}{n} \int_{\mathbb{R}^{n}} \text{div} (u^{\gamma} (\text{div } \vec{\mathbf{v}}) \vec{\mathbf{v}}) \phi
$$
\n
$$
= -\frac{1}{n} \gamma \int_{\mathbb{R}^{n}} u^{\gamma-1} \nabla u \, u^{\alpha-1} [\alpha H^{p} (\nabla u) - u g(u)] u^{\alpha} \vec{\mathbf{u}} \phi - \frac{1}{n} \int_{\mathbb{R}^{n}} u^{\gamma} \nabla [u^{\alpha-1} (\alpha H^{p} (\nabla u) - u g(u))] u^{\alpha} \vec{\mathbf{u}} \phi
$$
\n
$$
- \frac{1}{n} \int_{\mathbb{R}^{n}} (\text{div } \vec{\mathbf{v}})^{2} \phi
$$
\n
$$
= -\frac{1}{n} \alpha \gamma \int_{\mathbb{R}^{n}} u^{\gamma+2\alpha-2} H^{2p} (\nabla u) \phi - \frac{1}{n} \alpha \int_{\mathbb{R}^{n}} u^{\gamma+\alpha} \nabla (u^{\alpha-1} H^{p} (\nabla u)) \vec{\mathbf{u}} \phi - \frac{1}{n} \int_{\mathbb{R}^{n}} u^{\gamma} (\text{div } \vec{\mathbf{v}})^{2} \phi
$$
\n
$$
+ \frac{1}{n} \int_{\mathbb{R}^{n}} u^{\gamma+2\alpha-1} [(\alpha + \gamma) g(u) + u g'(u)] H^{p} (\nabla u) \phi.
$$

For the first term I_0 , combining with $(3.16)-(3.21)$ $(3.16)-(3.21)$, we have

(3.23)
\n
$$
I_0 = -\int_{\mathbb{R}^n} u^\gamma \vec{\mathbf{v}} \cdot \nabla \vec{\mathbf{v}} \nabla \phi
$$
\n
$$
= -\alpha \int_{\mathbb{R}^n} u^{\gamma + \alpha - 1} \vec{\mathbf{v}} \cdot \nabla u \, \vec{\mathbf{u}} \cdot \nabla \phi - \int_{\mathbb{R}^n} u^{\gamma + \alpha} \vec{\mathbf{v}} \cdot \nabla \vec{\mathbf{u}} \cdot \nabla \phi
$$
\n
$$
= I_2 + I_3.
$$

With the help of $(3.16)-(3.21)$ $(3.16)-(3.21)$ one finds that

(3.24)
\n
$$
I_2 = -\alpha \int_{\mathbb{R}^n} u^{\gamma + \alpha - 1} \vec{v} \cdot \nabla u \vec{u} \cdot \nabla \phi
$$
\n
$$
= \alpha \int_{\mathbb{R}^n} \text{div}(u^{\gamma + \alpha - 1} \vec{v} \cdot \nabla u \vec{u}) \phi
$$
\n
$$
= \alpha \int_{\mathbb{R}^n} [(\gamma + 2\alpha - 1)u^{\gamma + 2\alpha - 2} H^p(\nabla u) \vec{u} \cdot \nabla u + u^{\gamma + 2\alpha - 1} \nabla (H^p(\nabla u)) \vec{u} - u^{\gamma + 2\alpha - 1} H^p(\nabla u) g(u)] \phi
$$
\n
$$
= \alpha (\gamma + 2\alpha - 1) \int_{\mathbb{R}^n} u^{\gamma + 2\alpha - 2} H^{2p}(\nabla u) \phi + \alpha \int_{\mathbb{R}^n} u^{\gamma + 2\alpha - 1} \nabla (H^p(\nabla u)) \vec{u} \phi - \alpha \int_{\mathbb{R}^n} u^{\gamma + 2\alpha - 1} H^p(\nabla u) g(u) \phi.
$$

The integral I_3 is difficult involving a delicate interchange of order of differentiation. The proof of this step closely follows the proof of[[\[30\]](#page-26-10) , Proposition 6.2]. For convenience, we state the result that we need here. Readers could see the paper for more details.

CRITICAL ANISOTROPIC SOBOLEV EQUATION WITHOUT THE FINITE VOLUME CONSTRAINTS

Let $\overrightarrow{U_h}$ be the matrix with exponents

$$
\overrightarrow{U}_{h_i}^j = \frac{H^{p-1}(\nabla u(x+h^i))H_j(\nabla u(x+h^i)) - H^{p-1}(\nabla u(x))H_j(\nabla u(x))}{h}
$$

.

Since $\overrightarrow{u} \in W_{loc}^{1,2}(\Omega)$, it is standard that $\overrightarrow{U}_h \to \nabla \overrightarrow{u}$ in $L_{loc}^2(\Omega)$ as $h \to 0$. Hence we can write

$$
I_3 = -\lim_{h \to 0} \int_{\mathbb{R}^n} u^{\gamma + \alpha} \overrightarrow{v} \overrightarrow{U_h} \cdot \nabla \phi
$$

(3.25)

$$
= \lim_{h \to 0} \left(\int_{\mathbb{R}^n} \overrightarrow{v} \overrightarrow{U_h} \cdot \nabla (u^{\gamma + \alpha}) \phi + \int_{\mathbb{R}^n} u^{\gamma + \alpha} Tr \{ \overrightarrow{V} \overrightarrow{U_h} \} \phi + \int_{\mathbb{R}^n} u^{\gamma + \alpha} \overrightarrow{v} \cdot \text{div} (\overrightarrow{U_h^T}) \phi \right)
$$

$$
= I_4 + I_5 + I_6.
$$

By [\(3.16\)](#page-11-0)-[\(3.21\)](#page-12-2), with convergence in the sense of $L^2_{loc}(\Omega)$, we have

(3.26)
\n
$$
I_4 = \lim_{h \to 0} \int_{\mathbb{R}^n} \overrightarrow{\mathbf{v}} \overrightarrow{U_h} \cdot \nabla (u^{\gamma + \alpha}) \phi
$$
\n
$$
= (\gamma + \alpha) \int_{\mathbb{R}^n} u^{\gamma + 2\alpha - 1} \overrightarrow{\mathbf{u}} \cdot \nabla \overrightarrow{\mathbf{u}} \cdot \nabla u \phi
$$
\n
$$
= \frac{p - 1}{p} (\gamma + \alpha) \int_{\mathbb{R}^n} u^{\gamma + 2\alpha - 1} \overrightarrow{\mathbf{u}} \cdot \nabla (H^p(\nabla u)) \phi.
$$

and

$$
(3.27)
$$
\n
$$
I_5 = \lim_{h \to 0} \int_{\mathbb{R}^n} u^{\gamma + \alpha} Tr{\{\overrightarrow{V} \overrightarrow{U_h}\}\phi}
$$
\n
$$
= \int_{\mathbb{R}^n} u^{\gamma + \alpha} Tr{\{\overrightarrow{V} \nabla \overrightarrow{u}\}\phi}
$$
\n
$$
= \int_{\mathbb{R}^n} u^{\gamma + \alpha} Tr{\{\overrightarrow{V} \nabla (u^{-\alpha} \overrightarrow{v})\}\phi}
$$
\n
$$
= \int_{\mathbb{R}^n} u^{\gamma + \alpha} (u^{-\alpha} Tr{\{\overrightarrow{V}^2\} - \alpha u^{-1} \overrightarrow{u} \cdot \nabla u \cdot \nabla \overrightarrow{v})\phi}
$$
\n
$$
= \int_{\mathbb{R}^n} u^{\gamma + \alpha} (u^{-\alpha} Tr{\{\overrightarrow{V}^2\} - \alpha u^{-1} \overrightarrow{u} \cdot \nabla u \cdot \nabla (u^{\alpha} \overrightarrow{u}))\phi}
$$
\n
$$
= \int_{\mathbb{R}^n} u^{\gamma} Tr{\{\overrightarrow{V}^2\}\phi - \alpha^2 \int_{\mathbb{R}^n} u^{\gamma + 2\alpha - 2} H^{2p}(\nabla u)\phi - \alpha \frac{p - 1}{p} \int_{\mathbb{R}^n} u^{\gamma + 2\alpha - 1} \overrightarrow{u} \nabla (H^p(\nabla u))\phi}.
$$

Since $\frac{u(x+h^j)-u(x)}{h} \to \nabla u$ and \int 1 0 $g'(u(x)+t(u(x+h^j)-u(x))) dt \to g'(u)$ as $h \to 0$ uniformly on compact subsets of Ω , the distribution sense I_6 on compact subsets of Ω is (3.28)

$$
I_6 = \lim_{h \to 0} \left(\int_{\mathbb{R}^n} u^{\gamma + \alpha} \vec{\mathbf{v}} \cdot \mathrm{div}(\overrightarrow{U_h^T}) \phi \right)
$$

=
$$
\lim_{h \to 0} \left(\int_{\mathbb{R}^n} u^{\gamma + \alpha} \vec{\mathbf{v}} \cdot (- \int_0^1 g'(u(x) + t(u(x + h^j) - u(x))) dt) \frac{u(x + h^j) - u(x)}{h} \phi \right)
$$

=
$$
\int_{\mathbb{R}^n} u^{\gamma + 2\alpha} H^p(\nabla u) g'(u) \phi.
$$

Finally, inserting [\(3.26\)](#page-14-0), [\(3.27\)](#page-14-1) and [\(3.29\)](#page-15-0) into [\(3.25\)](#page-14-2), we get

$$
(3.29)
$$
\n
$$
I_{3} = \lim_{h \to 0} \left(\int_{\mathbb{R}^{n}} \vec{\mathbf{v}} \vec{U}_{h} \cdot \nabla(u^{\gamma+\alpha}) \phi + \int_{\mathbb{R}^{n}} u^{\gamma+\alpha} Tr{\{\vec{V} \vec{U}_{h} \}} \phi + \int_{\mathbb{R}^{n}} u^{\gamma+\alpha} \vec{\mathbf{v}} \cdot \text{div}(\vec{U}_{h}^{\mathcal{F}}) \phi \right)
$$
\n
$$
= \frac{p-1}{p} (\gamma + \alpha) \int_{\mathbb{R}^{n}} u^{\gamma+2\alpha-1} \vec{\mathbf{u}} \cdot \nabla(H^{p}(\nabla u)) \phi + \int_{\mathbb{R}^{n}} u^{\gamma} Tr{\{\vec{V}^{2} \}} \phi - \alpha^{2} \int_{\mathbb{R}^{n}} u^{\gamma+2\alpha-2} H^{2p}(\nabla u) \phi
$$
\n
$$
- \alpha \frac{p-1}{p} \int_{\mathbb{R}^{n}} u^{\gamma+2\alpha-1} \vec{\mathbf{u}} \nabla(H^{p}(\nabla u)) \phi + \int_{\mathbb{R}^{n}} u^{\gamma+2\alpha} H^{p}(\nabla u) g'(u) \phi
$$
\n
$$
= \frac{p-1}{p} \gamma \int_{\mathbb{R}^{n}} u^{\gamma+2\alpha-1} \vec{\mathbf{u}} \cdot \nabla(H^{p}(\nabla u)) \phi + \int_{\mathbb{R}^{n}} u^{\gamma} Tr{\{\vec{V}^{2} \}} \phi - \alpha^{2} \int_{\mathbb{R}^{n}} u^{\gamma+2\alpha-2} H^{2p}(\nabla u) \phi
$$
\n
$$
+ \int_{\mathbb{R}^{n}} u^{\gamma+2\alpha} H^{p}(\nabla u) g'(u) \phi.
$$

It is clear that the right side of [\(3.17\)](#page-12-1) has the form

(3.30)
$$
-\int_{\mathbb{R}^n} \omega \cdot \nabla \phi = I_0 + I_1.
$$

Combing with the calculated above , we have

(3.31)
$$
\int_{\mathbb{R}^n} (u^{\gamma} I(x) + \psi) \phi = - \int_{\mathbb{R}^n} \omega \cdot \nabla \phi.
$$

where

$$
\omega := u^{\gamma}(\overrightarrow{\mathbf{v}} \cdot \nabla \overrightarrow{\mathbf{v}} - \frac{1}{n} \overrightarrow{\mathbf{v}} \operatorname{div} \overrightarrow{\mathbf{v}}),
$$

and

$$
\psi = u^{\gamma + 2\alpha - 1} \left[(1 - \frac{np}{n-p}) \gamma g(u) - \frac{n-1}{n} u g'(u) \right] H^p(\nabla u) - \left(\frac{p-1}{p} (\gamma + 2\alpha - 1) \gamma + \frac{n-1}{n} \alpha^2 \right) u^{\gamma + 2\alpha - 2} H^{2p}(\nabla u)
$$

$$
+ \left(\frac{n-1}{n} \alpha + \frac{p-1}{p} \gamma \right) \operatorname{div} \left(u^{\gamma + 2\alpha - 1} \vec{\mathbf{u}} H^p(\nabla u) \right).
$$

Next we turn to the proof of inequality [\(3.17\)](#page-12-1). To this end we set

(3.32)
$$
\Omega_{\varepsilon} = \{x \in \mathbb{R}^n : 0 < |\nabla u| < \varepsilon\}.
$$

Then

$$
\int_{\mathbb{R}^n} (u^{\gamma} I(x) + \psi) \phi = \int_{\Omega_{\varepsilon}} (u^{\gamma} I(x) + \psi) \phi + \int_{\mathbb{R}^n \setminus \Omega_{\varepsilon}} (u^{\gamma} I(x) + \psi) \phi
$$

By formula (7.22) in [\[30\]](#page-26-10) we have

$$
\liminf_{\varepsilon \to 0} \int_{\Omega_{\varepsilon}} (u^{\gamma} I(x) + \psi) \phi \ge 0.
$$

Thus by monotone convergence theorem and passing to the limit as ε tends to 0 in [\(3.31\)](#page-15-1), we conclude that

(3.33)
$$
\int_{\mathbb{R}^n} u^{\gamma} I(x) \phi \leq - \int_{\mathbb{R}^n} \omega \cdot \nabla \phi,
$$

which finishes the proof. Now we are in position to prove Lemma [3.3.](#page-11-1) \Box

Proof of Lemma [3.3](#page-11-1)

Proof. Applying the definition of $I(x)$ and ω , and combining with lemma [3.4](#page-12-0) (especially with $\gamma = 1-n$, we could choose the suitable test function φ to obtain needed inequality. Using an argument analogous to the (3.33) with $m = 0$, that is,

(3.34)
$$
\int_{\mathbb{R}^n} \varphi v^{1-n} Tr\{W^2\} \leq - \int_{\mathbb{R}^n} v^{1-n} \overrightarrow{\mathbf{v}}^j W_{ij} \varphi_i.
$$

For $m \neq 0$ we can choose the test function as $(g_{\epsilon})^m \varphi$ in [\(3.33\)](#page-16-0) and get

$$
(3.35)
$$

$$
\int_{\mathbb{R}^n} v^{1-n} (g_\epsilon)^m \varphi Tr\{W^2\} \le -\int_{\mathbb{R}^n} v^{1-n} \vec{\nabla}^j W_{ij} [(g_\epsilon)^m \varphi]_i
$$

$$
= -m \int_{\mathbb{R}^n} v^{1-n} \vec{\nabla}^j W_{ij} (g_\epsilon)^{m-1} (g_\epsilon)_i - \int_{\mathbb{R}^n} v^{1-n} \vec{\nabla}^j W_{ij} (g_\epsilon)^m \varphi_i
$$

$$
= -I_1 - I_2.
$$

For the left side of [\(3.35\)](#page-16-1), we want to prove that

(3.36)
$$
\lim_{\epsilon \to 0} \int_{\mathbb{R}^n} v^{1-n} (g_{\epsilon})^m \varphi Tr\{W^2\} = \int_{\mathbb{R}^n} v^{1-n} g^m \varphi Tr\{W^2\}.
$$

From $C^{1,\alpha}$ regularity of v, we have $g = av^{-1}H^p(\nabla v) + bv^{-1} \in L^{\infty}_{loc}(\mathbb{R}^n)$ and $g_{\epsilon} \to g$ in $L^{\infty}_{loc}(\mathbb{R}^n)$ by the convolution approximation. Combining these with $\varphi \in C^{\infty}_c(\mathbb{R}^n)$, $v \in$ $L^{\infty}_{loc}(\mathbb{R}^n)$, $Tr\{W^2\} \in L^2_{loc}(\mathbb{R}^n)$ and Hölder inequality, we derive [\(3.36\)](#page-16-2).

For the integral term I_1 on the right side, since $\vec{\nabla}^j = H^{p-1}(\nabla v)H_j(\nabla v) \in L^{\infty}_{loc}(\mathbb{R}^n)$, $v \in L^{\infty}_{loc}(\mathbb{R}^n)$, $W_{ij} \in L^2_{loc}(\mathbb{R}^n)$, $g_{\epsilon} \to g$ in $L^{\infty}_{loc}(\mathbb{R}^n)$ and $(g_{\epsilon})_i \to g_i$ in $L^2_{loc}(\mathbb{R}^n)$, we obtain

(3.37)
$$
\lim_{\epsilon \to 0} m \int_{\mathbb{R}^n} v^{1-n} \vec{\mathbf{v}}^j W_{ij}(g_{\epsilon})^{m-1}(g_{\epsilon})_i = m \int_{\mathbb{R}^n} v^{1-n} \vec{\mathbf{v}}^j W_{ij} g^{m-1} g_i
$$

through Hölder inequality. Finally, for the integral term I_2 on the right side, we calculate that

(3.38)
$$
\lim_{\epsilon \to 0} \int_{\mathbb{R}^n} v^{1-n} \vec{\mathbf{v}}^j W_{ij}(g_{\epsilon})^m \varphi_i = \int_{\mathbb{R}^n} v^{1-n} \vec{\mathbf{v}}^j W_{ij} g^m \varphi_i.
$$

The combination of [\(3.35\)](#page-16-1), [\(3.37\)](#page-16-3) and [\(3.38\)](#page-17-0) leads to the result of Lemma [3.3.](#page-11-1)

3.4 Asymptotic estimates on bounded region. The main goal of this subsection is to prove Lemma [3.5](#page-17-1) below. In particular we could only use familiar Hölder inequality to prove the inequality [3.39](#page-17-2) and [3.40.](#page-17-3) It will remove the finite volume assumption, which is crucial in [\[8\]](#page-25-0).

 \Box

Lemma 3.5. *For* $p \le q < \alpha + 1$,

(3.39)
$$
\int_{B_R} v^{-q} H^p(\nabla v) \lesssim R^{n-q},
$$

and for $0 < q \leq \alpha + 1$

$$
(3.40)\qquad \qquad \int_{B_R} v^{-q} \lesssim R^{n-q}.
$$

Before giving the proof of Lemma [3.5,](#page-17-1) we first introduce a useful Lemma as follow.

Lemma 3.6.

(3.41)
$$
(\alpha+1-q)\int_{\mathbb{R}^n} v^{-q} H^p(\nabla v)\psi + \beta \int_{\mathbb{R}^n} v^{-q}\psi = -\int_{\mathbb{R}^n} \overrightarrow{\mathbf{v}}^j v^{1-q}\psi_j
$$

Proof. Replacing ψ by $v^{1-q}\psi$ in [\(3.5\)](#page-9-0), then we consider the term on the left side

$$
(3.42) \qquad \qquad -\int_{\mathbb{R}^n} \overrightarrow{\mathbf{v}}^j (v^{1-q} \psi)_j = -\int_{\mathbb{R}^n} (1-q) \overrightarrow{\mathbf{v}}^j v^{-q} v_j \psi - \int_{\mathbb{R}^n} \overrightarrow{\mathbf{v}}^j v^{1-q} \psi_j
$$
\n
$$
= -\int_{\mathbb{R}^n} (1-q) H^{p-1} (\nabla v) H_j (\nabla v) v^{-q} v_j \psi - \int_{\mathbb{R}^n} \overrightarrow{\mathbf{v}}^j v^{1-q} \psi_j.
$$

From the term on the right side of [\(3.5\)](#page-9-0), we obtain

(3.43)
$$
\int_{\mathbb{R}^n} v^{1-q} g \psi = \int_{\mathbb{R}^n} v^{1-q} (\alpha v^{-1} H^p (\nabla v) + \beta v^{-1}) \psi
$$

$$
= \int_{\mathbb{R}^n} \alpha v^{-q} H^p (\nabla v) \psi + \int_{\mathbb{R}^n} \beta v^{-q} \psi.
$$

Combining (3.42) with (3.43) , we get

(3.44)
$$
(\alpha+1-q)\int_{\mathbb{R}^n} v^{-q}H^p(\nabla v)\psi + \beta\int_{\mathbb{R}^n} v^{-q}\psi = -\int_{\mathbb{R}^n} \overrightarrow{\mathbf{v}}^j v^{1-q}\psi_j,
$$

which ends the proof of Lemma. Now we turn to the proof of Lemma [3.5.](#page-17-1) \Box

Proof of Lemma [3.5](#page-17-1)

Proof. Let $\theta > 0$ be a constant big enough and ρ be the cut-off function as [\(3.15\)](#page-11-3). Using (3.44) with $\psi = \rho^{\theta}$ we have

(3.45)
$$
(\alpha+1-q)\int_{\mathbb{R}^n} v^{-q} H^p(\nabla v)\rho^{\theta} + \beta \int_{\mathbb{R}^n} v^{-q} \rho^{\theta} = -\theta \int_{\mathbb{R}^n} v^{1-q} \overrightarrow{\mathbf{v}}^j \rho_j \rho^{\theta-1}.
$$

Since $H \in C^{1,\tau}_{loc}(\mathbb{R}^n)$ what we got from Proposition [\(2.2\)](#page-5-0), there exists a constant $M_1 > 0$ such that $H_j(\nabla v) \leq M_1$. Moreover, since $|\nabla^j \rho_j| \lesssim \frac{M_1}{R} H^{p-1}(\nabla v)$, we derive that

(3.46)
$$
-\theta \int_{\mathbb{R}^n} v^{1-q} \overrightarrow{\mathbf{v}}^j \rho_j \rho^{\theta-1} \lesssim \frac{M_1}{R} \int_{R^n} v^{1-q} \rho^{\theta-1} H^{p-1}(\nabla v) \leq \varepsilon \rho^{\theta} v^{-q} H^p(\nabla v) + \frac{M_1^p}{\varepsilon^{p-1} R^p} \rho^{\theta-p} v^{p-q},
$$

where $\varepsilon > 0$ is a sufficiently small constant by using the Young's inequality with exponent pair $(\frac{p}{p-1}, p)$. Hence, the inequality [\(3.46\)](#page-18-0) is used to get that

(3.47)

$$
(\alpha+1-q)\int_{\mathbb{R}^n} v^{-q}H^p(\nabla v)\rho^\theta+\beta\int_{\mathbb{R}^n} v^{-q}\rho^\theta\lesssim \varepsilon\int_{\mathbb{R}^n} \rho^\theta v^{-q}H^p(\nabla v)+\frac{M_1^p}{\varepsilon^{p-1}R^p}\int_{\mathbb{R}^n} \rho^{\theta-p}v^{p-q}.
$$

Since $p < q$ implies $\frac{q}{q-p} > 1$, similarly, we obtain

(3.48)
$$
\frac{M_1^p}{\varepsilon^{p-1} R^p} \rho^{\theta-p} v^{p-q} \leq \varepsilon v^{-q} \rho^{\theta} + \frac{M_1^q}{\varepsilon^{q-1} R^q} \rho^{\theta-q},
$$

by using the Young's inequality with exponent pair $(\frac{q}{q-p}, \frac{q}{p})$ $\frac{q}{p}$). Inserting [\(3.48\)](#page-18-1) into [\(3.47\)](#page-18-2) yields

$$
(3.49)
$$

$$
(\alpha+1-q)\int_{\mathbb{R}^n} v^{-q}H^p(\nabla v)\rho^\theta+\beta\int_{\mathbb{R}^n} v^{-q}\rho^\theta\lesssim \varepsilon\int_{\mathbb{R}^n} \rho^\theta v^{-q}H^p(\nabla v)+\varepsilon\int_{\mathbb{R}^n} v^{-q}\rho^\theta+\frac{M_1^q}{\varepsilon^{q-1}R^q}\int_{R^n} \rho^{\theta-q}.
$$

If $a + 1 - q > 0$, by recalling the definition of the test function ρ in [\(3.15\)](#page-11-3) and taking $\varepsilon > 0$ small enough with $\theta > q$, we see that

(3.50)
$$
\int_{B_R} v^{-q} H^p(\nabla v) + \int_{B_R} v^{-q} \lesssim R^{n-q}.
$$

This implies [\(3.39\)](#page-17-2) and [\(3.40\)](#page-17-3) for $p \le q < \alpha + 1$. On the other hand, for $0 < s < p \le q$, it follows from Hölder inequality that

$$
(3.51) \qquad \int_{B_R} v^{-s} \le \left(\int_{B_R} (v^{-s})^{\frac{q}{s}} \right)^{\frac{s}{q}} \cdot \left(\int_{B_R} (1)^{\frac{q}{q-s}} \right)^{\frac{q-s}{q}}
$$
\n
$$
\le (R^{n-q})^{\frac{s}{q}} \cdot (R^n)^{\frac{q-s}{q}}
$$
\n
$$
\le R^{n-s},
$$

where in the second step we have used [\(3.40\)](#page-17-3) with $p \leq q < \alpha + 1$. This implies (3.40) also valid for $0 < q < p$ and hence we get that [\(3.40\)](#page-17-3) holds for all $0 < q < a + 1$.

To prove [\(3.40\)](#page-17-3) for $q = \alpha + 1$, we first notice that the first term in [\(3.45\)](#page-18-3) vanishes and now we have

(3.52)
$$
b \int_{\mathbb{R}^n} v^{-\alpha-1} \rho^{\theta} \lesssim \frac{M_1}{R} \int_{\mathbb{R}^n} v^{-\alpha} \rho^{\theta-1} H^{p-1}(\nabla v),
$$

or that

(3.53)
$$
\int_{B_R} v^{-\alpha - 1} \lesssim \frac{M_1}{R} \int_{B_{2R}} v^{-\alpha} H^{p-1}(\nabla v).
$$

Next we estimate the right hand side of (3.53) . By Hölder inequality

$$
(3.54) \qquad \int_{B_{2R}} v^{-\alpha} H^{p-1}(\nabla v) \leq \left(\int_{B_{2R}} (v^{-\alpha+(p-1)(1-\varepsilon)})\right)^{\frac{1}{p}} \cdot \left(\int_{B_{2R}} (v^{\varepsilon-\alpha-1} H^p(\nabla v))\right)^{\frac{p-1}{p}}.
$$

Choosing $0 < \varepsilon < \min\{1, \frac{n-p}{n}\}$ $\frac{-p}{p}(p-1)$ } implies $p < a+1-\varepsilon < \alpha+1$ and $0 < \alpha-(1-\varepsilon)$ ε)(p - 1) < α + 1. Then for the right hand side of [\(3.54\)](#page-19-1), we can apply [\(3.40\)](#page-17-3) and [\(3.39\)](#page-17-2) to get

$$
(3.55) \quad \int_{B_{2R}} v^{-\alpha} H^{p-1}(\nabla v) \le \left(\int_{B_{2R}} (v^{-\alpha+(p-1)(1-\varepsilon)})^p \right)^{\frac{1}{p}} \cdot \left(\int_{B_{2R}} (v^{\varepsilon-\alpha-1} H^p(\nabla v))^{\frac{p}{p-1}} \right)^{\frac{p-1}{p}} \le R^{n-\alpha}.
$$

Inserting the latter into the preceding integral estimate [\(3.53\)](#page-19-0) then yields

$$
\int_{B_R} v^{-q} = \int_{B_R} v^{-(\alpha+1)}
$$
\n
$$
\leq \frac{1}{R} \cdot R^{n-\alpha}
$$
\n
$$
= R^{n-\alpha-1}
$$
\n
$$
= R^{n-q},
$$

which completed the proof of [\(3.40\)](#page-17-3) for $q = \alpha + 1$.

4. Proof of Theorem

In this section, our main effort is to prove the Theorem 1.1 by constructing the correlation between the matrix $W = \{W_{ij}\}\$ and vector fields and applying the key integral inequality [\(3.16\)](#page-11-0) what we got in Lemma [3.3,](#page-11-1) we obtain that the matrix is trace free. Let $u > 0$ be any weak solution of [\(1.3\)](#page-1-2) and take $v = u^{-\frac{p}{n-p}}$. We will use the notations and the results presented in section 2 and 3. For the sake of the proof of the matrix is trace free, we also need the following lemma. More details of the proof of this Lemma [4.1](#page-19-2) can be founded in the reference [\[28\]](#page-26-3).

Lemma 4.1. Let $n \geq 2$, $1 < p < n$ and $u \in W^{1,p}_{loc}(\mathbb{R}^n) \bigcap C^{1,\alpha}_{loc}(\mathbb{R}^n)$ be a positive, weak *solution of* [\(1.3\)](#page-1-2), *v and* g *be the functions defined as* $v = u^{-\frac{p}{n-p}}$ *and* $g = av^{-1}H^p(\nabla v)$ + bv^{-1} respectively. W be the $n \times n$ square matrix whose elements are denoted by $\{W_{ij}\}.$ *Then we have*

(1)For each $A = \{a_{ij}\}\$ be an $n \times n$ *square matrix*

$$
T_r\{AW\} \le c(p)T_r\{AA^T\} + T_r\{W^2\},\
$$

where $c(p)$ *is a positive constant depending only on p. (2)By summing over* i *and* j*, we have*

$$
T_r\{W^2\} = W_{ij}W_{ji}.
$$

Remark 4.2. *We should explain that the importance of the matrix* W *is trace free. More precisely, we first observe that* $\vec{v}^i_{,j} = (H^{p-1}(\nabla v)H_i(\nabla v))_j = H^{p-2}(\nabla v)BC$, with C is *Hessian matrix of* v and $B = (p-1)\nabla H(\nabla v) \bigotimes \nabla H(\nabla v) + H(\nabla v)H_{ij}(\nabla v)$. Since H^2 is *uniformly convex, Hessian matrix of* H² *is positive definite and we obtain that the matrix* B is positive definite. Hence, there exists an orthogonal matrix T such that $T^{-1}BT = D$ *where* D *is a diagonal matrix and is written as* $D = \{d_{ij}\}\$. Then we can rewrite $W =$ $H^{p-2}(\nabla v)BC - \frac{1}{n}$ $\frac{1}{n}gI = BF$ with $F = H^{p-2}(\nabla v)C - \frac{1}{n}$ $\frac{1}{n}$ g B^{-1} *. Obviously, the matrix* F *is symmetric implies that* $T^{-1}FT = Q$ *where* Q *is symmetric and is written as* $\{q_{ij}\}$ *. Thus, we get*

(4.1)
\n
$$
T_r\{W^2\} = T_r\{BFBF\}
$$
\n
$$
= T_r\{T^{-1}BFBFT\}
$$
\n
$$
= T_r\{T^{-1}BTT^{-1}FTT^{-1}BTT^{-1}FT\}
$$
\n
$$
= T_r\{DQDQ\}.
$$

If $T_r\{W^2\} = 0$, applying D is diagonal and positive definite, and Q is symmetric, then *we have*

$$
0 = T_r \{W^2\}
$$

\n
$$
= T_r \{DQDQ\}
$$

\n
$$
= \sum_{i,j,k,l=1}^n d_{ij} q_{jk} d_{kl} q_{li}
$$

\n
$$
= \sum_{i,k=1}^n d_{ii} d_{kk} q_{ki}^2,
$$

which implies that $DQ = 0$ *. It follows that* $W = BF = T DQT^{-1} = 0$ *. Now, we will prove that the matrix* W *is trace free.*

Proof of the matrix is trace free

Proof. Let $\theta > 0$ be a constant big enough and ρ be the smooth cut off functions as before. Replacing m by $-m$ and φ by ρ^{θ} in [\(3.16\)](#page-11-0), we have (4.3)

$$
\int_{\mathbb{R}^n} \rho^{\theta} v^{1-n} g^{-m} Tr\{W^2\} - nm \int_{\mathbb{R}^n} \rho^{\theta} v^{1-n} g^{-m-1} \overrightarrow{\mathbf{v}}^j W_{ij} W_i \le -\theta \int_{\mathbb{R}^n} v^{1-n} g^{-m} \overrightarrow{\mathbf{v}}^j W_{ij} \rho^{\theta-1} \rho_i
$$

Because $g = \alpha v^{-1} H^p(\nabla v) + \beta v^{-1}$, we can rewrite the left side of [\(4.3\)](#page-20-0) to conclude that

$$
\int_{\mathbb{R}^n} \rho^{\theta} v^{1-n} g^{-m} Tr{W^2} - nm \int_{\mathbb{R}^n} \rho^{\theta} v^{1-n} g^{-m-1} \vec{\mathbf{v}}^{j} W_{ij} W_{i}
$$
\n
$$
= \int_{\mathbb{R}^n} \rho^{\theta} v^{1-n} g^{-m-1} (\alpha v^{-1} H^p(\nabla v) + \beta v^{-1}) Tr{W^2} - nm \int_{\mathbb{R}^n} \rho^{\theta} v^{1-n} g^{-m-1} \vec{\mathbf{v}}^{j} W_{ij} W_{i}
$$
\n
$$
= \beta \int_{\mathbb{R}^n} \rho^{\theta} v^{-n} g^{-m-1} v^{-1} Tr{W^2} + (\alpha - nm) \int_{\mathbb{R}^n} \rho^{\theta} v^{-n} g^{-m-1} H^p(\nabla v) Tr{W^2}
$$
\n
$$
+ nm \int_{\mathbb{R}^n} \rho^{\theta} v^{-n} g^{-m-1} H^p(\nabla v) Tr{W^2} - nm \int_{\mathbb{R}^n} \rho^{\theta} v^{1-n} g^{-m-1} \vec{\mathbf{v}}^{j} W_{ij} W_{i}
$$
\n
$$
\geq \beta \int_{\mathbb{R}^n} \rho^{\theta} v^{-n} g^{-m-1} Tr{W^2} + (\alpha - nm) \int_{\mathbb{R}^n} \rho^{\theta} v^{-n} g^{-m-1} H^p(\nabla v) Tr{W^2}
$$
\n
$$
+ nm \int_{\mathbb{R}^n} \rho^{\theta} v^{-n} g^{-m-1} H^p(\nabla v) (Tr{W^2} - W_{ij} W_{ji}).
$$

By Lemma [4.1](#page-19-2) the bracket pair in above last integral is equal to zero. Now if we take $m = \frac{p-1}{p} - \varepsilon_0$ with $0 < \varepsilon_0 < \frac{p-1}{p}$ $\frac{-1}{p}$, then

$$
\int_{\mathbb{R}^n} \rho^{\theta} v^{1-n} g^{-m} Tr{W^2} - nm \int_{\mathbb{R}^n} \rho^{\theta} v^{1-n} g^{-m-1} \vec{\nabla}^{j} W_{ij} W_{i}
$$
\n
$$
\geq \beta \int_{\mathbb{R}^n} \rho^{\theta} v^{-n} g^{-m-1} Tr{W^2} + n\varepsilon_0 \int_{\mathbb{R}^n} \rho^{\theta} v^{-n} g^{-m-1} H^p(\nabla v) Tr{W^2}
$$
\n
$$
= \frac{p\varepsilon_0}{p-1} \left[\frac{p-1}{p\varepsilon_0} \beta \int_{\mathbb{R}^n} \rho^{\theta} v^{-n} g^{-m-1} Tr{W^2} + \alpha \int_{\mathbb{R}^n} \rho^{\theta} v^{-n} g^{-m-1} H^p(\nabla v) Tr{W^2} \right]
$$
\n
$$
\geq \frac{p\varepsilon_0}{p-1} \int_{\mathbb{R}^n} \rho^{\theta} v^{1-n} g^{-m} Tr{W^2}.
$$

Inserting this inequality into the preceding integral estimate [\(4.3\)](#page-20-0) then yields

(4.5)
$$
\frac{p\varepsilon_0}{p-1} \int_{\mathbb{R}^n} \rho^{\theta} v^{1-n} g^{-m} Tr\{W^2\} \leq \int_{\mathbb{R}^n} v^{1-n} g^{-m} \overrightarrow{\mathbf{v}}^j W_{ij} \rho^{\theta-1} \rho_i.
$$

Furthermore, if we take the matrix B as the form $B_{ij} = \epsilon \rho^{-1} \rho_i \vec{v}^j$ for $\epsilon > 0$ small enough, then we get

(4.6)
$$
\rho^{-1} \rho_i \overrightarrow{\mathbf{v}}^j W_{ij} \leq \frac{c(p)}{\epsilon} \rho^{-2} \rho_i \rho_j \overrightarrow{\mathbf{v}}^i \overrightarrow{\mathbf{v}}^j + \epsilon Tr\{W^2\},
$$

by using Lemma [4.1.](#page-19-2) Plugging this into [\(4.5\)](#page-21-0) we obtain

$$
\left(\frac{p\varepsilon_0}{p-1} - \epsilon\right) \int_{\mathbb{R}^n} \rho^{\theta} v^{1-n} g^{-m} Tr\{W^2\} \le \frac{c(p)}{\epsilon} \int_{\mathbb{R}^n} \rho^{\theta-2} v^{1-n} g^{-m} \rho_i \rho_j \overrightarrow{\mathbf{v}}^i \overrightarrow{\mathbf{v}}^j.
$$

Since $|\rho_i \rho_j| \leq \frac{1}{R^2}$ and $|\vec{\nabla}^i \vec{\nabla}^j| \leq M_1^2 H^{2p-2}(\nabla v)$, taking $\epsilon = \frac{p\varepsilon_0}{2(p-1)}$ we arrive at

(4.7)
$$
\int_{\mathbb{R}^n} \rho^{\theta} v^{1-n} g^{-m} Tr\{W^2\} \le \frac{M_1^2}{R^2} \int_{\mathbb{R}^n} \rho^{\theta-2} v^{1-n} g^{-m} H^{2p-2}(\nabla v).
$$

For the term on the right side of [\(4.7\)](#page-21-1), by

$$
g^{-m} \le (\alpha v^{-1} H^p(\nabla v))^{-\left(\frac{p-1}{p} - \varepsilon_0\right)},
$$

we can compute

$$
(4.8)
$$
\n
$$
\int_{\mathbb{R}^n} \rho^{\theta-2} v^{1-n} g^{-m} H^{2p-2}(\nabla v) \le \int_{\mathbb{R}^n} \rho^{\theta-2} v^{1-n} (v^{-1} H^p(\nabla v))^{-\left(\frac{p-1}{p}-\varepsilon_0\right)} H^{2p-2}(\nabla v)
$$
\n
$$
\lesssim \int_{B_{2R}} v^{1-n+\frac{p-1}{p}-\varepsilon_0} H^{p-1+p\varepsilon_0}(\nabla v)
$$
\n
$$
= \int_{B_{2R}} v^{(\varepsilon_0-\alpha-1)\frac{p-1+p\varepsilon_0}{p}} H^{p-1+p\varepsilon_0}(\nabla v) \cdot v^{1-n+\frac{p-1}{p}-\varepsilon_0-(\varepsilon_0-\alpha-1)\frac{p-1+p\varepsilon_0}{p}},
$$

Recalling $0 < \varepsilon_0 < \frac{p-1}{p}$ $\frac{-1}{p}$, furthermore, we assume $0 < \varepsilon_0 < \min\{\frac{p-1}{p}\}$ $\frac{-1}{p},\frac{1}{p}$ $\frac{1}{p}$. Then we deduce that

$$
(4.9)
$$
\n
$$
\int_{\mathbb{R}^n} \rho^{\theta - 2} v^{1 - n} g^{-m} H^{2p - 2}(\nabla v) \lesssim \int_{B_{2R}} v^{(\varepsilon_0 - \alpha - 1) \frac{p - 1 + p\varepsilon_0}{p}} H^{p - 1 + p\varepsilon_0}(\nabla v) \cdot v^{1 - n + \frac{p - 1}{p} - \varepsilon_0 - (\varepsilon_0 - \alpha - 1) \frac{p - 1 + p\varepsilon_0}{p}} \lesssim \left(\int_{B_{2R}} v^{(\varepsilon_0 - \alpha - 1)} H^p(\nabla v) \right)^{\frac{p - 1 + p\varepsilon_0}{p}} \left(\int_{B_{2R}} v^{(1 - n + \frac{p - 1}{p} - \varepsilon_0 - (\varepsilon_0 - \alpha - 1) \frac{p - 1 + p\varepsilon_0}{p}}) \frac{1 - p\varepsilon_0}{p} \right)^{\frac{1 - p\varepsilon_0}{p}}
$$
\n
$$
= I_1 \cdot I_2,
$$

by using Hölder inequality with exponent pair $(\frac{p}{p-1+p\varepsilon_0},\frac{p}{1-p}\)$ $\frac{p}{1-p\varepsilon_0}$) to (4.8) .

Next, we are going to estimate the two terms I_1 and I_2 of (4.9) by applying lemma [3.5.](#page-17-1) To this ends, we assume furthermore $0 < \varepsilon_0 < \min\{\frac{p-1}{n}\}$ $\frac{-1}{p},\frac{1}{p}$ $\frac{1}{p}$, $\frac{np-n-p^2+p}{p}$ $\frac{-p^2+p}{p}$. So, for the first one I_1 , by $0 < \varepsilon_0 < \frac{np-n-p^2+p}{p}$ $\frac{-p^2+p}{p}$ we have $p < \alpha + 1 - \varepsilon_0 < \alpha + 1$. Analogously to what we did in the proof of Lemma [3.5,](#page-17-1) we see that

(4.10)
$$
I_1 = \left(\int_{B_{2R}} v^{(\varepsilon_0 - \alpha - 1)} H^p(\nabla v) \right)^{\frac{p-1+pe_0}{p}} \lesssim R^{(n+\varepsilon_0 - \alpha - 1)\frac{p-1+pe_0}{p}}.
$$

For the second one I_2 , we observe that the exponent of v could read as

$$
(1 - n + \frac{p-1}{p} - \varepsilon_0 - (\varepsilon_0 - \alpha - 1)\frac{p-1 + p\varepsilon_0}{p})\frac{p}{1 - p\varepsilon_0} = \frac{1}{p}[3p^2 - 2(n+1)p + n] + \left[\frac{3p-1 - n}{\frac{1}{p} - \varepsilon_0} + 1\right]\varepsilon_0.
$$

So we consider the mater in the following two cases:

(i)
$$
\frac{n+1}{3} \le p < \frac{(n+1) + \sqrt{(n+1)^2 - 3n}}{3},
$$

(ii)
$$
\frac{(n+1)+\sqrt{(n+1)^2-3n}}{3} \le p < n.
$$

Case (i). In this case, we see $-\alpha-1 \leq \frac{1}{n}$ $\frac{1}{p}[3p^2-2(n+1)p+n] < 0$ and $\left[\frac{3p-1-n}{\frac{1}{p}-\varepsilon_0}+1\right]\varepsilon_0 > 0$. So if we choose $\varepsilon_0 > 0$ small enough, say $0 < \varepsilon_0 < \min\{\frac{p-1}{n}\}$ $\frac{-1}{p},\frac{1}{2p}$ $rac{1}{2p}, \frac{np-n-p^2+p}{p}$ $\frac{-p^2+p}{p}, \frac{-[3p^2-2(n+1)p+n]}{p[\frac{3p-1-n}{1}+1]}$ $\frac{p^{2}-2(n+1)p+n}{\frac{1}{2p}}\},$ then $-\alpha - 1 < (1 - n + \frac{p-1}{p} - \varepsilon_0 - (\varepsilon_0 - \alpha - 1) \frac{p-1+p\varepsilon_0}{p}) \frac{p}{1-p}$ $\frac{p}{1-p\varepsilon_0} < 0$. Therefore, we can use Lemma [3.5](#page-17-1) to obtain

$$
(4.11)
$$
\n
$$
I_2 = \left(\int_{B_{2R}} v^{(1-n+\frac{p-1}{p}-\varepsilon_0-(\varepsilon_0-\alpha-1)\frac{p-1+p\varepsilon_0}{p})\frac{p}{1-p\varepsilon_0}} \right)^{\frac{1-p\varepsilon_0}{p}} \lesssim (R^{n+(1-n+\frac{p-1}{p}-\varepsilon_0-(\varepsilon_0-\alpha-1)\frac{p-1+p\varepsilon_0}{p})\frac{p}{1-p\varepsilon_0}})^{\frac{1-p\varepsilon_0}{p}} = R^{n \cdot \frac{1-p\varepsilon_0}{p} + (1-n+\frac{p-1}{p}-\varepsilon_0-(\varepsilon_0-\alpha-1)\frac{p-1+p\varepsilon_0}{p})}.
$$

The combination of (4.10) , (4.11) and (4.9) leads to

$$
(4.12)
$$
\n
$$
\int_{\mathbb{R}^n} \rho^{\theta-2} v^{1-n} g^{-m} H^{2p-2}(\nabla v) \lesssim \int_{B_{2R}} v^{(\varepsilon_0 - \alpha - 1) \frac{p-1+p\varepsilon_0}{p}} H^{p-1+p\varepsilon_0}(\nabla v) v^{1-n+\frac{p-1}{p}-\varepsilon_0-(\varepsilon_0-\alpha-1) \frac{p-1+p\varepsilon_0}{p}} \leq \left(\int_{B_{2R}} v^{(\varepsilon_0 - \alpha - 1)} H^p(\nabla v) \right)^{\frac{p-1+p\varepsilon_0}{p}} \left(\int_{B_{2R}} v^{(1-n+\frac{p-1}{p}-\varepsilon_0-(\varepsilon_0-\alpha-1) \frac{p-1+p\varepsilon_0}{p}) \frac{p}{1-p\varepsilon_0}} \right)^{\frac{1-p\varepsilon_0}{p}} \leq R^{(n+\varepsilon_0-\alpha-1) \frac{p-1+p\varepsilon_0}{p}} \cdot R^{n \cdot \frac{1-p\varepsilon_0}{p} + (1-n+\frac{p-1}{p}-\varepsilon_0-(\varepsilon_0-\alpha-1) \frac{p-1+p\varepsilon_0}{p})} = R^{2-\frac{1}{p}-\varepsilon_0}.
$$

Case (ii). In this case, we still have $(1 - n + \frac{p-1}{p} - \varepsilon_0 - (\varepsilon_0 - \alpha - 1) \frac{p-1+p\varepsilon_0}{p}) \frac{p}{1-p}$ $\frac{p}{1-p\varepsilon_0}>0$ easily. Thus, by virtue of [\(3.2\)](#page-8-1) we get

$$
(4.13)
$$

$$
\left(\int_{B_{2R}} v^{(1-n+\frac{p-1}{p}-\varepsilon_0-(\varepsilon_0-\alpha-1)\frac{p-1+ p\varepsilon_0}{p})\frac{p}{1-p\varepsilon_0}}\right)^{\frac{1-p\varepsilon_0}{p}}\lesssim \left(R^{n+(1-n+\frac{p-1}{p}-\varepsilon_0-(\varepsilon_0-\alpha-1)\frac{p-1+ p\varepsilon_0}{p})\frac{p}{1-p\varepsilon_0}\cdot \frac{p}{p-1}}\right)^{\frac{1-p\varepsilon_0}{p}}
$$

.

From [\(4.9\)](#page-22-1), [\(4.10\)](#page-22-2) and [\(4.13\)](#page-23-1) it holds that

$$
(4.14)
$$
\n
$$
\int_{\mathbb{R}^n} \rho^{\theta-2} v^{1-n} g^{-m} H^{2p-2}(\nabla v) \leq R^{(n+\varepsilon_0-\alpha-1)\frac{p-1+ps_0}{p}} \cdot \left(R^{n+(1-n+\frac{p-1}{p}-\varepsilon_0-(\varepsilon_0-\alpha-1)\frac{p-1+ps_0}{p})\frac{p}{1-p\varepsilon_0}\cdot\frac{p}{p-1}}\right)^{\frac{1-p\varepsilon_0}{p}} \leq R^{2-\frac{(2p-1)(n-p)}{p^2(p-1)}+\frac{\varepsilon_0}{p-1}(\alpha+\frac{1}{p}+\varepsilon_0-p)}.
$$

Hence, for any $\frac{n+1}{3} \leq p < n$, combining [\(4.7\)](#page-21-1) with [\(4.12\)](#page-23-2) and combining (4.7) with [\(4.14\)](#page-23-3) separately, we conclude that

(4.15)
$$
\int_{\mathbb{R}^n} \rho^{\theta - 2} v^{1 - n} g^{-m} Tr\{W^2\} \lesssim R^{-\mu(\varepsilon_0)},
$$

with $\mu(\varepsilon_0) = \frac{1}{p} + \varepsilon_0$ in case (i) or $\mu(\varepsilon_0) = \frac{(2p-1)(n-p)}{p^2(p-1)} - \frac{\varepsilon_0}{p-1}$ $\frac{\varepsilon_0}{p-1}(\alpha+\frac{1}{p}+\varepsilon_0-p)$ in case (ii). Finally, if we choose $\varepsilon_0 > 0$ small enough such that $0 < \varepsilon_0 < \min\{\frac{p-1}{p}\}$ $\frac{-1}{p},\frac{1}{2p}$ $\frac{1}{2p}, \frac{np-n-p^2+p}{p}$ $\frac{-p^2+p}{p}, \frac{-[3p^2-2(n+1)p+n]}{p[\frac{3p-1-n}{p+1}]}$ $\frac{p^2-2(n+1)p+n}{p[\frac{3p-1-n}{2p}+1]}, \frac{(2p-1)(n-p)}{p^2(\alpha+2)}$ $\frac{p-1)(n-p)}{p^2(\alpha+2)}\}$ then it holds that

(4.16)
$$
\int_{\mathbb{R}^n} \rho^{\theta - 2} v^{1 - n} g^{-m} Tr\{W^2\} \lesssim R^{-\mu},
$$

for some constant $\mu > 0$ depending only on n, p in both cases. Letting $R \to \infty$ in [\(4.16\)](#page-23-4) we deduce

(4.17)
$$
\int_{\mathbb{R}^n} v^{1-n} g^{-m} Tr\{W^2\} \leq 0,
$$

which implies $W = 0$ a.e. in \mathbb{R}^n , since $Tr{W^2} \ge 0$ and the "=" happens if and only if $W = 0$ (see Remark [4.2\)](#page-20-1). Now we are in position to prove Theorem [1.1.](#page-2-1)

Proof of the Theorem [1.1](#page-2-1)

Proof. By the definition of W_{ij} , $W = 0$ is equivalent to

(4.18)
$$
H^{p-1}(\nabla v)\nabla H(\nabla v) = \lambda(x - x_0),
$$

which implies that

(4.19)
$$
x - x_0 = \frac{1}{\lambda} H^{p-1}(\nabla v) \nabla H(\nabla v).
$$

We notice that, acting H_0 on both sides of (4.18) and applying (2.6) , one could obtain that

(4.20)
$$
H^{p-1}(\nabla v) = \lambda H_0(x - x_0).
$$

Furthermore, according [\(4.18\)](#page-24-0) and [\(4.20\)](#page-24-1) we have

(4.21)

$$
\nabla H(\nabla v) = \frac{\lambda(x - x_0)}{H^{p-1}(\nabla v)}
$$

$$
= \frac{x - x_0}{H_0(x - x_0)}.
$$

Submitting [\(4.20\)](#page-24-1) and [\(4.21\)](#page-24-2) into [\(2.8\)](#page-4-3), and applying the property of 0-homogeneous of ∇H_0 (the proof is same as H_i in Lemma [\(2.1\)](#page-4-2)), then we compute

$$
\nabla v = H(\nabla v) \nabla H_0 (\nabla H(\nabla v))
$$

= $H(\nabla v) \nabla H_0 \left(\frac{x - x_0}{H_0 (x - x_0)} \right)$
= $\lambda^{\frac{1}{p-1}} H_0^{\frac{1}{p-1}} (x - x_0) \nabla H_0 (x - x_0)$
= $\frac{p - 1}{p} \lambda^{\frac{1}{p-1}} \nabla \left(H_0^{\frac{p}{p-1}} (x - x_0) \right),$

which implies that

$$
v = C_1 + C_2 H_0^{\frac{p}{p-1}} (x - x_0),
$$

for some $C_1, C_2 > 0$. Thus we have $u = U_\lambda$ and the proof of Theorem [1.1](#page-2-1) has been \Box

Data availability: Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

26 LU CHEN AND YABO YANG

REFERENCES

- [1] T. Aubin. Equations differentielles non linéaires et problème de Yamabe concernant la courbure scalaire. J. Math. Pures Appl., 55 (1976), 269-296.
- [2] A. Alvino, V. Ferone, G. Trombetti, and P. L. Lions. Convex symmetrization and applications. Annales De Linstitut Henri Poincare Non Linear Analysis, (2)14 (1997), 275-293.
- [3] C. Bianchini and G. Ciraolo. Wulff shape characterizations in overdetermined anisotropic elliptic problems. Comm. Partial Differential Equations, 43 (2018), 790-820.
- [4] Y. Z. Chen and E Dibenedetto. Boundary estimates for solutions of nonlinear degenerate parabolic systems. J. Reine Angew. Math., 395 (1989), 102-131.
- [5] L. A. Caffarelli, B. Gidas, and J. Spruck. Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth. Comm. Pure Appl. Math., (3)42 (1989), 271-297.
- [6] L. A. Caffarelli. Regularity theorems for weak solutions of some nonlinear systems. Communications on Pure and Applied Mathematics, (6)35 (1982), 833-838.
- [7] D. C. Erausquin, B. Nazaret, and C. Villani. A mass transportation approach to sharp Sobolev and Gagliardo-Nirenberg inequalities. Advances in Mathematics, (2)182 (2004), 307-332.
- [8] G. Ciraolo, A. Figalli, and A. Roncoroni. Symmetry results for critical anisotropic p-Laplacian equations in convex cones. Geom. Funct. Anal., 30(2020) 770-803.
- [9] G. Catine, D. D. Monticelli, and A. Roncoroni. On the critical p-Laplace equation. [arXiv:2204.06940v](http://arxiv.org/abs/2204.06940)3, 2022.
- [10] L. Chen and Y. B. Yang. Classification of positive solutions of Hardy-Sobolev equation without the finite volume constraints. [arXiv:2204.06940v](http://arxiv.org/abs/2204.06940)3, 2023.
- [11] L. Damascelli, S. Merch´an, L. Montoro and B. Sciunzi. Radial symmetry and applications for a problem involving the $-\Delta_p(\cdot)$ operator and critical nonlinearity in \mathbb{R}^n . Adv. Math., (10) 265 (2014), 313-335.
- [12] W. Dai and G. L. Qin. Classification of nonnegative classical solutions to third-order equations. Adv. Math., 328 (2018), 822-857.
- [13] W. Dai and G. L. Qin. Method of scaling spheres, Liouville theorems in inner or outer (unbounded or bounded) generalized radially convex domains, blowing-up analysis on domains with not necessarily $C¹$ boundary and other applications. [arXiv:2302.13988.](http://arxiv.org/abs/2302.13988)
- [14] W. Dai and G. Qin. Liouville type theorems for fractional and higher order Hénon-Hardy type equations viathe method of scaling spheres. Int. Math. Res. Not., 11 (2023), 9001-9070.
- [15] E. Dibenedetto and J. Manfredi. On the higher integrability of the gradient of weak solutions of certain degenerate elliptic systems. Am. J. Math., 115 (1993), 1107-1134.
- [16] E. Dibenedetto. $C^{1+\alpha}$ local regularity of weak solutions of degenerate elliptic equations. Nonlinear Anal., 7 (1983), 827-850.
- [17] B. Gidas, W. M. Ni and L, Nirenberg. Symmetry of positive solutions of nonlinear elliptic equations in \mathbb{R}^n . Mathematical analysis and applications. A (1981), 369-402.
- [18] B. Gidas and J. Spruck. Global and local behavior of positive solutions of nonlinear elliptic equations. Comm. Pure Appl. Math., 34 (1981) 525-598.
- [19] T. Iwaniec. Projections onto gradient fields and L^p -estimates for degenerated elliptic operators. Studia Math., 75 (1983), 293-312.
- [20] D. Jerison and J. M. Lee. Extremals for the Sobolev inequality on the Heisenberg group and the C-R Yamabe problem. J. Amer. Math. Soc., (1)1 (1988), 1-13.
- [21] J. M. Lee and T. H. Parker. The Yamabe problem. Bull. Amer. Math. Soc., 17 (1987), 37-91.
- [22] S. C. Lee and H. Yun. $C^{1,\alpha}$ -regularity for functions in solution classes and its application to parabolic normalized p-Laplace equations. J. Differ. Equ., 378 (2024), 539-558.
- [23] J. L. Lewis. Regularity of the derivatives of solutions to certain degenerate elliptic equations. Indiana Univ. Math. J., 32 (1983), 849-858.
- [24] Y. Lian and K. Zhang. Boundary pointwise $C^{1,\alpha}$ and $C^{1,\alpha}$ regularity for fully nonlinear elliptic equations. J. Differ. Equ., (2)269 (2020), 1172-1191.
- [25] G. M. Lieberman. Boundary regularity for solutions of degenerate elliptic equations. Nonlinear Analysis Theory Methods and Applications, (11)12 (1988), 1203-1219.

CRITICAL ANISOTROPIC SOBOLEV EQUATION WITHOUT THE FINITE VOLUME CONSTRAINTS7

- [26] X. N. Ma and Q. Ou. A Liouville theorem for a class semilinear elliptic equations on the Heisenberg group. Adv. Math. 413 (2023), 108851.
- [27] X. N. Ma, Q. Ou and T. Wu. Jerison-Lee identity and Semi-linear subelliptic equation on C-R manifold. [arXiv:2311.16428,](http://arxiv.org/abs/2311.16428) 2023.
- [28] Q. Ou. On the classification of entire solutions to the critical p-Laplace equation. [arXiv:2210.05141v](http://arxiv.org/abs/2210.05141)1, 2022.
- [29] S. L. Sobolev. On a theorem of functional (in russian). Mat. Sb., 4 (1938).
- [30] J. Serrin and H. Zou. Cauchy-Liouville and universal boundedness theorems for quasilinear elliptic equations and inequalities. Acta Math., 189(2002), 79-142.
- [31] J. Serrin. Local behaviour of solutions of quasilinear equations. Acta. Math., 113 (1965), 219-240.
- [32] B. Sciunzi. Classification of positive $D^{1,p}(\mathbb{R}^n)$ -solutions to the critical p-Laplace equation in \mathbb{R}^n . Advance in Mathematics, 291 (2016), 12-23.
- [33] G. Talenti. A weighted version of a rearrangement inequality. Ann. Univ. Ferrara, 43 (1997), 121-133.
- [34] P. Tolksdorf. Regularity for a more general class of quasilinear elliptic equations. J. Differ. Equ., 51 (1983), 126-150.
- [35] K. Uhlenbeck. Regularity for a class of non-linear elliptic systems. Acta Math. 138 (1977), 219-240.
- [36] J. Vétois. A priori estimates and application to the symmetry of solutions for critical p-Laplace equations. J. Differ. Equ., (1)260 (2016) , 149-161.

Key Laboratory of Algebraic Lie Theory and Analysis of Ministry of Education, SCHOOL OF MATHEMATICS AND STATISTICS, BEIJING INSTITUTE OF TECHNOLOGY, BEIJING 100081,

P. R. China

Email address: chenlu5818804@163.com

Key Laboratory of Algebraic Lie Theory and Analysis of Ministry of Education, SCHOOL OF MATHEMATICS AND STATISTICS, BEIJING INSTITUTE OF TECHNOLOGY, BEIJING 100081, P. R. China

Email address: 18048320592@163.com