

UNCOMMON LINEAR SYSTEMS OF TWO EQUATIONS

DINGDING DONG, ANQI LI, AND YUFEI ZHAO

ABSTRACT. A system of linear equations L is common over \mathbb{F}_p if, as $n \rightarrow \infty$, any 2-coloring of \mathbb{F}_p^n gives asymptotically at least as many monochromatic solutions to L as a random 2-coloring.

The notion of common linear systems is analogous to that of common graphs, i.e., graphs whose monochromatic density in 2-edge-coloring of cliques is asymptotically minimized by the random coloring. Saad and Wolf initiated a systematic study on identifying common linear systems, built upon the earlier work of Cameron–Cilleruelo–Serra. When L is a single equation, Fox–Pham–Zhao gave a complete characterization of common linear equations. When L consists of two equations, Kamčev–Liebenau–Morrison showed that irredundant 2×4 linear systems are always uncommon. In this work, (1) we determine commonness of all 2×5 linear systems up to a small number of cases, and (2) we show that all $2 \times k$ linear systems with k even and girth (minimum number of nonzero coefficients of a nonzero equation spanned by the system) $k - 1$ are uncommon, answering a question of Kamčev–Liebenau–Morrison.

1. INTRODUCTION

One of the major open problems in extremal graph theory is *Sidorenko’s conjecture* [24], which roughly states that for bipartite graphs H , a random graph G on n vertices asymptotically minimizes the density of H in G among all G with n vertices and fixed edge density. Sidorenko’s conjecture has been verified for various classes of bipartite graphs H (e.g. [6, 8, 9, 10, 11, 15, 19, 20, 23, 25] and the references therein), although a complete proof of Sidorenko’s conjecture still eludes us. A closely related graph property is the following coloring variant known as *commonness*: a graph H is *common* if the random coloring minimizes the density of monochromatic copies of H in K_n among all possible two-colorings of the edges of K_n . An early result of Goodman [14] showed that K_3 is common. Motivated by this result, Burr and Rosta [7] conjectured that all graphs H are common, which was disproved by Sidorenko (who showed that a triangle with a pendant edge is uncommon) and Thomason [26] (who showed that K_4 is uncommon). In fact, any graph containing a K_4 is uncommon [16]. We do not even have a good guess at what a complete characterization of common graphs might look like.

In this paper, we are interested in an arithmetic analogue (over finite fields) of this notion of commonness. To motivate the setup, we observe that for a given set $A \subseteq \mathbb{F}_p^n$ satisfying $-A = A$, the additive quadruples in A (that is, $(x, y, z, w) \in A^4$ such that $x + y - z - w = 0$) correspond to 4-cycles in the Cayley graph on A (that is, the graph with vertex set indexed by \mathbb{F}_p^n where $x \sim y$ if and only if $x - y \in A$). As such, a natural arithmetic analogue of commonness would be to ask for which linear equations in \mathbb{F}_p is it true that the random two-coloring of \mathbb{F}_p^n minimizes the density of monochromatic solutions to the linear equation. We can similarly define an arithmetic analogue of Sidorenko’s conjecture. The study of such arithmetic analogues was first posed by Saad and Wolf [22]. In this paper, we are interested in the commonness of certain linear *systems* of equations. Before we formalize this setup, we begin by introducing some definitions.

Definition 1.1. We say that L is an $m \times k$ linear system if it is a collection of m linear equations L_1, \dots, L_m on k variables with integer coefficients. In other words, L is a collection of linear

Li was supported by the Trinity Studentship in Mathematics. Most of this work was completed while Li was an undergraduate at MIT. Zhao was supported in part by NSF CAREER Award DMS-2044606.

equations

$$\begin{aligned} L_1 \mathbf{x} &= a_{1,1}x_1 + \cdots + a_{1,k}x_k, \\ &\vdots \\ L_m \mathbf{x} &= a_{m,1}x_1 + \cdots + a_{m,k}x_k, \end{aligned}$$

with $\{a_{i,j} \in \mathbb{Z} : i \in [m], j \in [k]\}$.

We note that there is a one-to-one correspondence between $m \times k$ linear systems and matrices in $\mathbb{Z}^{m \times k}$; this means we can also represent L as the matrix

$$L = \begin{pmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,k} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m,1} & a_{m,2} & \cdots & a_{m,k} \end{pmatrix} \in \mathbb{Z}^{m \times k}.$$

We say that an $m \times k$ linear system L is *common over* \mathbb{F}_p if for every 2-coloring of \mathbb{F}_p^n , the number of monochromatic solutions $\mathbf{x} \in \mathbb{F}_p^n$ to $L\mathbf{x} = 0$ is at least $1 - o(1)$ times the expected number of monochromatic solutions given by a random 2-coloring, asymptotically as $n \rightarrow \infty$. Below, we will state an equivalent definition as a precise inequality.

In the following, we restrict our attention to *irredundant linear systems*. Roughly speaking, a system L is irredundant if it does not have ‘‘repeated’’ equations or variables.

Definition 1.2. Let L be a linear system. We say that L is *irredundant* if:

- (1) the rows of L are linearly independent,
- (2) no column of L is the zero vector,
- (3) the row span of L does not contain vectors of the form $(0, \dots, 0, 1, 0, \dots, 0, -1, 0, \dots, 0)$.

If L is not irredundant, we say that it is *redundant*.

Such a restriction is without loss of generality: if L is redundant, then by removing extraneous equations and variables from L we can obtain an irredundant linear system L' such that L is common if and only if L' is common. Therefore, we restrict our attention to irredundant linear systems for the rest of this paper (see also [17, Section 2] for a discussion on this). By restricting our attention to irredundant linear systems, we can also pass to a ‘‘functional’’ analogue of commonness which we introduce next.

Earlier on, we defined the commonness property of a linear system L in terms of minimizing the number of monochromatic solutions to $L\mathbf{x} = 0$ in a 2-coloring of \mathbb{F}_p^n . It turns out that there is a ‘‘functional’’ variant of this definition which is more convenient for us to work with. This is the version of the definition of common and Sidorenko systems that we now state. We refer the interested reader to [13, Section 2] for a proof of the equivalence of the functional and set definitions of the properties of being common and Sidorenko.

Definition 1.3. Let L be an irredundant $m \times k$ linear system and p be prime. For every $n \in \mathbb{N}$ and $f : \mathbb{F}_p^n \rightarrow \mathbb{R}$, define the *density function*

$$t_L(f) = \mathbb{E}_{\mathbf{x} \in \mathbb{F}_p^n : L\mathbf{x} = 0} f(x_1) \cdots f(x_k).$$

- (1) We say that L is *common over* \mathbb{F}_p if for all $n \in \mathbb{N}$ and $f : \mathbb{F}_p^n \rightarrow [0, 1]$, we have

$$t_L(f) + t_L(1 - f) \geq 2^{1-k}.$$

If this does not hold, then we say that L is *uncommon over* \mathbb{F}_p .

- (2) We say that L is *Sidorenko over* \mathbb{F}_p if for all $n \in \mathbb{N}$ and $f : \mathbb{F}_p^n \rightarrow [0, 1]$, we have

$$t_L(f) \geq (\mathbb{E}f)^k.$$

Remark 1.4. If L is Sidorenko over \mathbb{F}_p , then it is common over \mathbb{F}_p . Indeed, if L is Sidorenko, then $t_L(f) + t_L(1-f) \geq (\mathbb{E}f)^k + (1-\mathbb{E}f)^k \geq 2^{-k+1}$ by convexity.

Several recent works address the question of characterizing common and Sidorenko linear systems. When L is itself a single linear equation (i.e., $m = 1$), the *length* of L is defined to be the number of its nonzero coefficients. Fox, Pham and Zhao [13] showed that a linear equation L is Sidorenko if and only if its coefficients can be partitioned into cancelling pairs; furthermore, it is common and not Sidorenko if and only if it has odd length. Their results give a complete characterization of common and Sidorenko linear equations.

No complete characterization is known for linear systems of higher rank. Some partial results in this direction include the following: Versteegen [27] proved that 4-AP is uncommon – this establishes the arithmetic analogue of Thomason’s result on the uncommonness of K_4 . Kamčev, Liebenau and Morrison proved that if a linear system consists of Sidorenko linear equations aligned in a “tree template”, then it is Sidorenko [18]; they also proved that any linear system that has an irredundant 2×4 subsystem is uncommon [17]. Altman [1] explored the notion of local Sidorenko property and showed that any linear system of two equations, with every 2×2 minor invertible, is not Sidorenko. On the other hand, Altman [2] also established that there is a non-Sidorenko linear system that becomes common after adding sufficiently many free variables, answering a question of [22].

In this paper, we determine commonness of all 2×5 linear systems up to a small number of cases. As a byproduct, we answer [18, Question 5.2] by constructing an uncommon 2×5 linear system that contains an additive quadruple. We also prove [17, Conjecture 6.1] that all $2 \times k$ linear systems, with k even and “girth” (which we define in Definition 1.7) $k - 1$, are uncommon over \mathbb{F}_p for all sufficiently large p . Before we formally state our results, we need to introduce some definitions.

Definition 1.5. Let L be an $m \times k$ linear system on variables x_1, \dots, x_k , and L' be an $m' \times k'$ linear system on a subset of the variables $x_{i_1}, \dots, x_{i_{k'}}$. Write $\mathbf{x} = (x_1, \dots, x_k)$ and $\mathbf{x}' = (x_{i_1}, \dots, x_{i_{k'}})$.

- (1) We say that L' is a *subsystem* of L if $L\mathbf{x} = 0$ implies $L'\mathbf{x}' = 0$.
- (2) Suppose $k' = k$ (so $\mathbf{x}' = \mathbf{x}$). We say that L and L' are *isomorphic* if we can permute the columns of L' so that the obtained matrix has the same row span as that of L .

Example 1.6. The four-term arithmetic progression

$$L'(x_1, x_2, x_3, x_4) = \begin{pmatrix} 1 & -2 & 1 & 0 \\ 0 & 1 & -2 & 1 \end{pmatrix}$$

is a subsystem of the five-term arithmetic progression

$$L(x_1, x_2, x_3, x_4, x_5) = \begin{pmatrix} 1 & -2 & 1 & 0 & 0 \\ 0 & 1 & -2 & 1 & 0 \\ 0 & 0 & 1 & -2 & 1 \end{pmatrix}.$$

Finally, we recall that a graph with odd girth is not Sidorenko (see for instance [12, Theorem 3.1]). In [18], the authors defined a natural notion of *girth* for linear systems and proved a parallel result that a linear system with odd girth is not Sidorenko. We next give the definition of *girth* for a linear system.

Definition 1.7. Let $L \in \mathbb{Z}^{m \times k}$ be an $m \times k$ linear system. The *girth* of L , denoted as $s(L)$, is the minimum possible size of the support of a nonzero vector in the row span of L .

We are now ready to formally state our main results. While odd girth was enough to guarantee that a linear system L is not Sidorenko, we cannot expect it alone to give the stronger conclusion that L is uncommon: we have seen earlier that single, odd-length equations are common. But what if we restrict our attention to systems of at least two equations? Would we be able to deduce the uncommonness of such a system from its girth? Our first result gives a partial answer to

this question and shows that all two-equation linear systems with even number of variables and maximum girth are uncommon.

Theorem 1.8. *Suppose $k \geq 4$ is even. Let L be an irredundant $2 \times k$ system with $s(L) = k - 1$. Then there exists p_L such that L is uncommon over \mathbb{F}_p for all $p \geq p_L$.*

By combining the proof strategy of Theorem 1.8 and some additional ideas from [17] and [1], we obtain the following more general result, which is also a generalization of [17, Corollary 1.2].

Theorem 1.9. *Suppose $k \geq 4$ is even. Let L be a linear system, with $s(L) = k - 1$, that has an irredundant $2 \times k$ subsystem. Then there exists p_L such that L is uncommon over \mathbb{F}_p for all $p \geq p_L$.*

We remark that an independent proof of Theorems 1.8 and 1.9 was given by Altman and Liebenau [3].

Next, one can ask about the commonness of $2 \times k$ linear systems for k odd. This turns out to be much more subtle than the k even case; even for $k = 5$, there are common 2×5 linear systems with $s(L) = 3$ and $s(L) = 4$. In fact, for each 2×5 linear system that is not one of the two unknown cases, we are able to determine whether it is common over \mathbb{F}_p for all p , or it is uncommon over \mathbb{F}_p for all sufficiently large p .

Theorem 1.10. *Let L be an irredundant 2×5 linear system that is not isomorphic to*

$$\begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 1 & -1 & 3 & 0 & -3 \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 2 & -2 & 3 & 0 & -3 \end{pmatrix}.$$

(1) *If L is isomorphic to*

$$\begin{pmatrix} 1 & 0 & -1 & 2 & -2 \\ 0 & 1 & 2 & -1 & -2 \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} 1 & -1 & 1 & -1 & 0 \\ 1 & 2 & -1 & 0 & -2 \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 2 & -2 & 1 & 0 & -1 \end{pmatrix}$$

or

$$\begin{pmatrix} a & b & 0 & 0 & c \\ 0 & 0 & a & b & c \end{pmatrix} \quad \text{with } a, b, c \in \mathbb{Z} \setminus \{0\}$$

or

$$\begin{pmatrix} c & -c & d & -d & 0 \\ a & b & 0 & 0 & -a - b \end{pmatrix} \quad \text{with } a, b, a + b, c, d \in \mathbb{Z} \setminus \{0\}$$

such that $\{|a/b|, |a/(a+b)|, |b/(a+b)|\} \cap \{|c/d|, |d/c|\} \neq \emptyset$,

then L is common over \mathbb{F}_p for all primes p .

(2) *If L is not isomorphic to any of the above, then there exists p_L such that L is uncommon over \mathbb{F}_p for all $p \geq p_L$.*

The proofs of uncommonness in Theorem 1.8, Theorem 1.9 and Theorem 1.10 utilize suitably designed *Fourier templates* which we introduce in Section 3; these Fourier templates can be thought of as functions with suitably chosen Fourier coefficients that certificate uncommonness over \mathbb{F}_p for sufficiently large p . We are unable to construct a suitable Fourier template for the two unknown cases in Theorem 1.10, although we believe that our main constraints are essentially computational barriers which limit the size of the support of the Fourier templates we are able to consider. It is also plausible that the systems are actually common, although we are also unable to prove that.

Question 1.11. Are the two systems not covered by Theorem 1.10 common or uncommon?

There also remains the interesting question of classifying $2 \times k$ systems for k odd and $k \geq 7$, and it is natural to ask if there is a general way to do this classification for all odd k at once.

Question 1.12. What is the classification of the commonness of $2 \times k$ systems for $k \geq 7$ odd?

Outline. In Section 2, we introduce the Fourier analytic approach to solve problems on identifying common and Sidorenko linear systems. In Section 3, we define the notion of a Fourier template and demonstrate its utility by proving (Theorem 3.5) that if we can find an appropriate Fourier template corresponding to a given linear system L , then L is uncommon over all \mathbb{F}_p for sufficiently large p . Next, we prove Theorem 1.8 and Theorem 1.9 in Section 4 and Theorem 1.10 in Sections 5 and 6. We begin each of Sections 4 (Section 4.1) and 5 (Section 5.1) with some examples to illustrate the key proof ideas behind our construction of the relevant Fourier templates.

Acknowledgements. We thank Yunkun Zhou for providing the proof that the linear system

$$\begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 2 & -2 & 1 & 0 & -1 \end{pmatrix}$$

is common. We thank Dylan Toh for suggesting the greatly simplified proof of Fact 5.18. We thank Pablo Parrilo for providing suggestions and help on symmetry-adapted semi-definite programs. We also thank Nitya Mani and Mark Saengrungkongka for helpful discussions.

2. PRELIMINARIES

2.1. Fourier analytic approach. In this section, we discuss the Fourier analytic approach for identifying common and Sidorenko linear systems. Let $\widehat{\mathbb{F}_p^n}$ denote the set of homomorphisms from $\mathbb{F}_p^n \rightarrow \mathbb{C}^\times$. Since every element in $\widehat{\mathbb{F}_p^n}$ is of the form $x \mapsto e^{2\pi i r \cdot x/p}$, we have an isomorphism between \mathbb{F}_p^n and $\widehat{\mathbb{F}_p^n}$. For a function $f : \mathbb{F}_p^n \rightarrow \mathbb{R}$, the Fourier transform of f is the function $\widehat{f} : \widehat{\mathbb{F}_p^n} \rightarrow \mathbb{C}$ defined by

$$\widehat{f}(r) = \mathbb{E}_{x \in \mathbb{F}_p^n} f(x) \overline{r(x)}$$

with inverse Fourier transform

$$f(x) = \sum_{r \in \widehat{\mathbb{F}_p^n}} \widehat{f}(r) r(x).$$

Thus, we can rewrite the density function $t_L(f)$ as

$$t_L(f) = \mathbb{E}_{L\mathbf{x}=0} \sum_{r_1, \dots, r_k \in \widehat{\mathbb{F}_p^n}} \widehat{f}(r_1) \cdots \widehat{f}(r_k) r_1(x_1) \cdots r_k(x_k) = \sum_{r_1, \dots, r_k \in \widehat{\mathbb{F}_p^n}} \widehat{f}(r_1) \cdots \widehat{f}(r_k) \mathbb{E}_{L\mathbf{x}=0} r_1(x_1) \cdots r_k(x_k).$$

Now observe that $\mathbb{E}_{L\mathbf{x}=0} r_1(x_1) \cdots r_k(x_k) = \mathbb{E}_{L\mathbf{x}=0} (e^{2\pi i/p})^{r_1 \cdot x_1 + \cdots + r_k \cdot x_k}$ equals 1 if (r_1, \dots, r_k) is a linear combination of the row vectors of L , and equals 0 otherwise. Let $\vec{\ell}_1, \dots, \vec{\ell}_m \in \mathbb{Z}^k$ be the row vectors of L and $\vec{a}_1, \dots, \vec{a}_k \in \mathbb{Z}^m$ the column vectors of L . We know that

$$\text{span}(\vec{\ell}_1, \dots, \vec{\ell}_m) = \{(\vec{a}_1 \cdot r, \dots, \vec{a}_k \cdot r) : r \in (\widehat{\mathbb{F}_p^n})^m\}.$$

Hence we have

$$t_L(f) = \sum_{r \in (\widehat{\mathbb{F}_p^n})^m} \widehat{f}(\vec{a}_1 \cdot r) \cdots \widehat{f}(\vec{a}_k \cdot r). \quad (1)$$

In particular, when L itself is a linear equation $L\mathbf{x} = a_1x_1 + \cdots + a_kx_k$, we can rewrite the density function $t_L(f)$ as

$$t_L(f) = \sum_{r \in \widehat{\mathbb{F}_p^n}} \widehat{f}(a_1r) \cdots \widehat{f}(a_kr). \quad (2)$$

2.2. Reduction to critical sets. In this section, we define critical sets of a linear system and then introduce a reductive step in [17] that will be useful in our proof. The following two definitions can be found in [17, Section 2.2].

Definition 2.1. For every linear system L with girth $s(L)$, define

$$c(L) = \begin{cases} s(L) & s(L) \text{ even} \\ s(L) + 1 & s(L) \text{ odd.} \end{cases}$$

We say that $B \subseteq [k]$ is a *critical set* of L if

- (1) $|B| = c(L)$, and
- (2) L has a nonempty subsystem L' whose variable set is $\{x_i : i \in B\}$.

Let $\mathcal{C}(L)$ denote the collection of critical sets of L .

Definition 2.2. Let L be a linear system and $B \subseteq [k]$ be a critical set of L .

- (1) Define

$$m_B = \max\{m' : L \text{ has an } m' \times c(L) \text{ subsystem on variable set } \{x_i : i \in B\}\}.$$

- (2) Define L_B to be the (unique up to isomorphism) $m_B \times c(L)$ subsystem of L on variable set $\{x_i : i \in B\}$.

The proof of the uniqueness of L_B is standard linear algebra, and we refer the readers to [17, Lemma 2.4].

Remark 2.3. Since no equation of length $< s(L)$ can be a subsystem of L , we have the following:

- When $s(L)$ is even (so $c(L) = s(L)$), every $B \in \mathcal{C}(L)$ has $m_B = 1$, and L_B must be a length- $c(L)$ equation.
- When $s(L)$ is odd (so $c(L) = s(L) + 1$), every $B \in \mathcal{C}(L)$ has $m_B \in \{1, 2\}$, and L_B is either a length- $c(L)$ equation or a $2 \times c(L)$ linear system with girth $s(L)$.

One of the results in [17] states that to conclude a linear system L is uncommon, it sometimes suffices to study the collection $\{L_B : B \in \mathcal{C}(L)\}$.

Proposition 2.4 ([17, Theorem 3.1]). *Let L be an $m \times k$ linear system. If there exists $f : \mathbb{F}_p^n \rightarrow \mathbb{R}$ such that $\mathbb{E}f = 0$ and $\sum_{B \in \mathcal{C}(L)} t_{L_B}(f) < 0$, then L is uncommon over \mathbb{F}_p .*

Proof. Consider any $h : \mathbb{F}_p^n \rightarrow [0, 1]$ with $\mathbb{E}h = 1/2$, and define $f = h - 1/2 : \mathbb{F}_p^n \rightarrow [-1/2, 1/2]$. Expanding each term, one can check that

$$\begin{aligned} & t_L(h) + t_L(1 - h) \\ &= \mathbb{E}_{L\mathbf{x}=0}(1/2 + f(x_1)) \cdots (1/2 + f(x_k)) + \mathbb{E}_{L\mathbf{x}=0}(1/2 - f(x_1)) \cdots (1/2 - f(x_k)) \\ &= 2^{1-k} + 2 \sum_{\substack{\emptyset \neq B \subseteq [k] \\ |B| \text{ even}}} 2^{|B|-k} \mathbb{E}_{L\mathbf{x}=0} \left[\prod_{i \in B} f(x_i) \right] = 2^{1-k} + 2 \sum_{\substack{B \subseteq [k] \\ |B| \geq c(L) \text{ even}}} 2^{|B|-k} \mathbb{E}_{L\mathbf{x}=0} \left[\prod_{i \in B} f(x_i) \right], \end{aligned}$$

where the last equality follows from the fact that $\mathbb{E}f = 0$, and L does not have nonempty subsystems on at most $c(L) - 2$ variables. Thus, if there is some $f : \mathbb{F}_p^n \rightarrow [-1/2, 1/2]$ such that $\mathbb{E}f = 0$ and

$$\sum_{\substack{B \subseteq [k] \\ |B| \geq c(L) \text{ even}}} 2^{|B|-k} \mathbb{E}_{L\mathbf{x}=0} \left[\prod_{i \in B} f(x_i) \right] < 0,$$

then by Definition 1.3, L is uncommon over \mathbb{F}_p .

Observe that

$$\begin{aligned} & \sum_{\substack{B \subseteq [k] \\ |B| \geq c(L) \text{ even}}} 2^{|B|-k} \mathbb{E}_{L\mathbf{x}=0} \left[\prod_{i \in B} f(x_i) \right] \\ &= \sum_{B \in \mathcal{C}(L)} 2^{c(L)-k} \mathbb{E}_{L\mathbf{x}=0} \left[\prod_{i \in B} f(x_i) \right] + \sum_{\substack{B \subseteq [k] \\ |B| \geq c(L)+2 \text{ even}}} 2^{|B|-k} \mathbb{E}_{L\mathbf{x}=0} \left[\prod_{i \in B} f(x_i) \right]. \end{aligned}$$

Consider any critical set $B \in \mathcal{C}(L)$ and its corresponding subsystem L_B . A linear algebra calculation (see also [17, Lemma 2.6]) shows that every solution to L_B extends to the same number of solutions to L . That is, for every solution \mathbf{x}' to $L_B \mathbf{x}' = 0$, there are $(p^n)^{k-m-c(L)+m_B}$ solutions \mathbf{x} to $L\mathbf{x} = 0$ such that \mathbf{x} restricted on the variables $\{x_i : i \in B\}$ equals \mathbf{x}' . Thus we get that

$$\mathbb{E}_{L\mathbf{x}=0} \left[\prod_{i \in B} f(x_i) \right] = \mathbb{E}_{L_B \mathbf{x}'=0} \left[\prod_{i \in B} f(x_i) \right] = t_{L_B}(f).$$

Suppose there exists $\tilde{f} : \mathbb{F}_p^n \rightarrow \mathbb{R}$ with $\mathbb{E}\tilde{f} = 0$ and $\sum_{B \in \mathcal{C}(L)} t_{L_B}(\tilde{f}) < 0$. Taking $f = \epsilon \tilde{f}$ with $\epsilon > 0$ sufficiently small, we can obtain $f : \mathbb{F}_p^n \rightarrow [-1/2, 1/2]$ with

$$\begin{aligned} & \sum_{B \in \mathcal{C}(L)} 2^{c(L)-k} \mathbb{E}_{L\mathbf{x}=0} \left[\prod_{i \in B} \epsilon \tilde{f}(x_i) \right] + \sum_{\substack{B \subseteq [k] \\ |B| \geq c(L)+2 \text{ even}}} 2^{|B|-k} \mathbb{E}_{L\mathbf{x}=0} \left[\prod_{i \in B} \epsilon \tilde{f}(x_i) \right] \\ &= 2^{c(L)-k} \epsilon^{c(L)} \sum_{B \in \mathcal{C}(L)} t_{L_B}(\tilde{f}) + O(\epsilon^{c(L)+2}), \end{aligned}$$

which is negative. Hence L is uncommon over \mathbb{F}_p . \square

3. FOURIER TEMPLATE

As discussed in Section 2, for linear system L and function $f : \mathbb{F}_p \rightarrow \mathbb{R}$, we can write the solution density $t_L(f)$ in terms of the Fourier coefficients of f . In order to construct a function f that certifies uncommonness, it is often fruitful to view the problem in the Fourier space and construct a suitable $\hat{f} : \widehat{\mathbb{F}_p} \rightarrow \mathbb{C}$. We will introduce a notion of a *Fourier template* which can then be turned into some $\hat{f} : \widehat{\mathbb{F}_p} \rightarrow \mathbb{C}$ for all sufficiently large p .

Example 3.1. As an example, to show that the linear equation

$$L\mathbf{x} = x_1 + 2x_2 + 3x_3 + 4x_4$$

is uncommon over \mathbb{F}_p , we can consider the function $f : \mathbb{F}_p \rightarrow \mathbb{R}$ defined by

$$\begin{cases} \hat{f}(1) = \hat{f}(-1) = -1 \\ \hat{f}(2) = \hat{f}(-2) = \hat{f}(3) = \hat{f}(-3) = \hat{f}(4) = \hat{f}(-4) = 1 \\ \hat{f}(r) = 0 \quad \text{otherwise.} \end{cases}$$

Notice that $\mathbb{E}f = \hat{f}(0) = 0$. Moreover, for all sufficiently large p , we have

$$t_L(f) = \sum_{r \in \widehat{\mathbb{F}_p}} \hat{f}(r) \hat{f}(2r) \hat{f}(3r) \hat{f}(4r) = \hat{f}(1) \hat{f}(2) \hat{f}(3) \hat{f}(4) + \hat{f}(-1) \hat{f}(-2) \hat{f}(-3) \hat{f}(-4) = -2 < 0.$$

By taking $\epsilon > 0$ sufficiently small such that $\epsilon f : \mathbb{F}_p \rightarrow [-1/2, 1/2]$ and letting $h = 1/2 + \epsilon f$, we can obtain a function $h : \mathbb{F}_p \rightarrow [0, 1]$ such that

$$\begin{aligned} & t_L(h) + t_L(1-h) \\ &= \mathbb{E}_{L\mathbf{x}=0}[(1/2 + \epsilon f(x_1)) \cdots (1/2 + \epsilon f(x_4))] + \mathbb{E}_{L\mathbf{x}=0}[(1/2 - \epsilon f(x_4)) \cdots (1/2 - \epsilon f(x_1))] \\ &= 1/8 + 2\epsilon^4 \mathbb{E}_{L\mathbf{x}=0}[f(x_1) \cdots f(x_4)] = 1/8 + 2\epsilon^4 t_L(f) < 1/8. \end{aligned}$$

Therefore, L is uncommon over \mathbb{F}_p for all sufficiently large p .

The key step in Example 3.1 is to construct an appropriate Fourier transform $\widehat{f} : \widehat{\mathbb{F}_p} \rightarrow \mathbb{C}$ such that $t_L(f) < 0$. Since \widehat{f} has finite support, this construction works for all sufficiently large p . What we essentially did is creating a finitely supported map $g : \mathbb{Z} \rightarrow \mathbb{C}$ from the *integers*, defined by $\pm 1 \mapsto -1$ and $\pm 2, \pm 3, \pm 4 \mapsto 1$, such that the Fourier transform $\widehat{f} = g$ (which is well-defined for all sufficiently large p) gives us a function f that helps certify uncommonness. This map $g : \mathbb{Z} \rightarrow \mathbb{C}$ is what we call a 1-dimensional Fourier template.

We now formally introduce the definition of Fourier templates.

Definition 3.2. For every $m \times k$ linear system L and $d \in \mathbb{N}$, we say that $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ is a *d-dimensional Fourier template* if the following holds:

- g has finite support,
- $g(-r) = \overline{g(r)}$ for all $r \in \mathbb{Z}^d$,
- $g(0) = 0$.

Definition 3.3. Let $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ be a *d-dimensional Fourier template*, and

$$L = (\vec{a}_1 \quad \vec{a}_2 \quad \cdots \quad \vec{a}_k) = \begin{pmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,k} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m,1} & a_{m,2} & \cdots & a_{m,k} \end{pmatrix} \in \mathbb{Z}^{m \times k}$$

be an $m \times k$ linear system. (Here $\vec{a}_1, \dots, \vec{a}_k \in \mathbb{Z}^m$ are the columns of L .)

(1) Define the associated function $g_L : \mathbb{Z}^{d \times m} \rightarrow \mathbb{C}$ by

$$g_L(R) = g(R\vec{a}_1) \cdots g(R\vec{a}_k) = \prod_{i=1}^k g(R\vec{a}_i).$$

(2) Define $\sigma_L(g) = \sum_{R \in \mathbb{Z}^{d \times m}} g_L(R)$.

Remark 3.4. Notice that $\sigma_L(g)$ is well-defined as g has finite support. Moreover, since

$$\sigma_L(g) = \sum_{R \in \mathbb{Z}^{d \times m}} g_L(R) = \sum_{R \in \mathbb{Z}^{d \times m}} g_L(-R) = \sum_{R \in \mathbb{Z}^{d \times m}} \overline{g_L(R)} = \overline{\sigma_L(g)},$$

we always have $\sigma_L(g) \in \mathbb{R}$.

We now state and prove the main theorem of this section.

Theorem 3.5. *Let L be an $m \times k$ linear system. If there exists a d -dimensional Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ such that $\sum_{B \in \mathcal{C}(L)} \sigma_{L_B}(g) < 0$, then L is uncommon over \mathbb{F}_p for all sufficiently large p .*

When $d = 1$, the proof of Theorem 3.5 essentially goes as Example 3.1. When $d > 1$, the main idea is to construct an appropriate Freiman homomorphism $\gamma : \mathbb{Z}^d \rightarrow \mathbb{Z}$ that projects \mathbb{Z}^d onto a generalized arithmetic progression in \mathbb{Z} . We will then build the certificate $f : \mathbb{F}_p \rightarrow \mathbb{C}$ by setting $\widehat{f}(r) = g(\gamma^{-1}(r))$ for all $r \in \text{im } \gamma \subseteq \widehat{\mathbb{F}_p}$.

Proof. Consider any critical set $B \in \mathcal{C}(L)$. Suppose

$$L_B = (\vec{a}_1 \ \cdots \ \vec{a}_{k'}) = \begin{pmatrix} a_{1,1}x_1 & \cdots & a_{1,k'}x_{k'} \\ \vdots & \ddots & \vdots \\ a_{m',1}x_1 & \cdots & a_{m',k'}x_{k'} \end{pmatrix} \in \mathbb{Z}^{m' \times k'},$$

so that $\vec{a}_1, \dots, \vec{a}_{k'} \in \mathbb{Z}^{m'}$ are the column vectors of L_B . Recall from Remark 2.3 that we have $m' \in \{1, 2\}$ and $k' = c(L)$.

Pick matrix $Q_B \in \mathbb{Z}^{(k'-m') \times k'}$ so that the rows of L_B span the kernel of Q_B . Then for $f : \mathbb{F}_p \rightarrow \mathbb{R}$, we have (recall (1))

$$t_{L_B}(f) = \sum_{r \in (\widehat{\mathbb{F}_p})^{m'}} \widehat{f}(\vec{a}_1 \cdot r) \cdots \widehat{f}(\vec{a}_{k'} \cdot r) = \sum_{\substack{\vec{v} \in (\widehat{\mathbb{F}_p})^{k'} \\ Q_B \vec{v} = 0}} \widehat{f}(v_1) \cdots \widehat{f}(v_{k'}). \quad (3)$$

For $M \in \mathbb{N}$, let $\llbracket -M, M \rrbracket$ denote the subset $\{-M, \dots, -1, 0, 1, \dots, M\} \subseteq \mathbb{Z}$. Suppose the Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ vanishes outside $\llbracket -M, M \rrbracket^d$. Observe that, for $C \in \mathbb{N}$ sufficiently large, the projection $\gamma : \llbracket -M, M \rrbracket^d \rightarrow \mathbb{Z}$ defined by

$$\gamma(w_0, \dots, w_{d-1}) = w_0 + w_1 C + \cdots + w_{d-1} C^{d-1}$$

is injective. Moreover, the inverse $\phi := \gamma^{-1} : \text{im} \gamma \rightarrow \llbracket -M, M \rrbracket^d$ satisfies the following property: for all $B \in \mathcal{C}(L)$ and $\vec{v} \in (\text{im} \gamma)^{k'}$, we have $Q_B \vec{v} = 0 \in \mathbb{Z}^{k'-m'}$ if and only if $Q_B \phi(\vec{v}) = 0 \in \mathbb{Z}^{(k'-m') \times d}$, where

$$\phi(\vec{v}) = \begin{pmatrix} \phi(v_1) \\ \vdots \\ \phi(v_{k'}) \end{pmatrix} \in \mathbb{Z}^{k' \times d}.$$

Suppose p is sufficiently large. Construct $f : \mathbb{F}_p \rightarrow \mathbb{R}$ as follows:

- For all $-p/2 < r < p/2$ with $r \in \text{im} \gamma$, set $\widehat{f}(r) = g(\phi(r))$.
- Otherwise, set $\widehat{f}(r) = 0$.

By (3), we have

$$t_{L_B}(f) = \sum_{\substack{\vec{v} \in (\widehat{\mathbb{F}_p})^{k'} \\ Q_B \vec{v} = 0}} \widehat{f}(v_1) \cdots \widehat{f}(v_{k'}) = \sum_{\substack{\vec{v} \in (\widehat{\mathbb{F}_p})^{k'} \\ Q_B \phi(\vec{v}) = 0}} \widehat{f}(v_1) \cdots \widehat{f}(v_{k'}) = \sum_{\substack{U \in \mathbb{Z}^{k' \times d} \\ Q_B U = 0}} g(U_1) \cdots g(U_{k'}),$$

where $U_1, \dots, U_{k'} \in \mathbb{Z}^d$ are the row vectors for each $U \in \mathbb{Z}^{k' \times d}$. Since $Q_B U = 0$ if and only if

$$U = \begin{pmatrix} \vec{a}_1 \cdot r_1 & \cdots & \vec{a}_1 \cdot r_d \\ \vdots & \ddots & \vdots \\ \vec{a}_{k'} \cdot r_1 & \cdots & \vec{a}_{k'} \cdot r_d \end{pmatrix} \text{ for } r_1, \dots, r_d \in \mathbb{Z}^{m'},$$

we get that

$$\begin{aligned} t_{L_B}(f) &= \sum_{r_1, \dots, r_d \in \mathbb{Z}^{m'}} g(\vec{a}_1 \cdot r_1, \dots, \vec{a}_1 \cdot r_d) \cdots g(\vec{a}_{k'} \cdot r_1, \dots, \vec{a}_{k'} \cdot r_d) \\ &= \sum_{R \in \mathbb{Z}^{d \times m'}} g(R\vec{a}_1) \cdots g(R\vec{a}_{k'}) = \sigma_{L_B}(g). \end{aligned}$$

Therefore, if $\sum_{B \in \mathcal{C}(L)} \sigma_{L_B}(g) < 0$, then we have $\sum_{B \in \mathcal{C}(L)} t_{L_B}(f) = \sum_{B \in \mathcal{C}(L)} \sigma_{L_B}(g) < 0$ for all sufficiently large p . By Proposition 2.4, L is uncommon over \mathbb{F}_p . \square

Finally, we remark that we can join two Fourier templates g, \tilde{g} to form a new Fourier template h . Moreover, the joined Fourier template satisfies $\sigma_L(h) = \sigma_L(g)\sigma_L(\tilde{g})$.

Proposition 3.6. *Let $g : \mathbb{Z}^{d_1} \rightarrow \mathbb{C}$, $\tilde{g} : \mathbb{Z}^{d_2} \rightarrow \mathbb{C}$ be two Fourier templates. Define the map $h : \mathbb{Z}^{d_1+d_2} \rightarrow \mathbb{C}$ by*

$$h(r_1, r_2) = g(r_1)\tilde{g}(r_2) \quad \text{for all } r_1 \in \mathbb{Z}^{d_1}, r_2 \in \mathbb{Z}^{d_2}.$$

Then h is a $(d_1 + d_2)$ -dimensional Fourier template. Moreover, for every linear system L , we have

$$\sigma_L(h) = \sigma_L(g)\sigma_L(\tilde{g}).$$

Proof. One can check directly from Definition 3.2 that h is a $(d_1 + d_2)$ -dimensional Fourier template. Let

$$L = (\vec{a}_1 \quad \dots \quad \vec{a}_k) \in \mathbb{Z}^{m \times k}$$

be an $m \times k$ linear system. For matrix $R \in \mathbb{Z}^{(d_1+d_2) \times m}$, write

$$R = \begin{pmatrix} R_1 \\ R_2 \end{pmatrix}$$

where $R_1 \in \mathbb{Z}^{d_1 \times m}$ and $R_2 \in \mathbb{Z}^{d_2 \times m}$. We then have $R\vec{a}_i = (R_1\vec{a}_i, R_2\vec{a}_i)$ and $h(R\vec{a}_i) = g(R_1\vec{a}_i)\tilde{g}(R_2\vec{a}_i)$ for every $i \in [k]$, which gives

$$\begin{aligned} \sigma_L(h) &= \sum_{R \in \mathbb{Z}^{(d_1+d_2) \times m}} h_L(R) = \sum_{R \in \mathbb{Z}^{(d_1+d_2) \times m}} h(R\vec{a}_1) \cdots h(R\vec{a}_k) \\ &= \sum_{R_1 \in \mathbb{Z}^{d_1 \times m}} \sum_{R_2 \in \mathbb{Z}^{d_2 \times m}} g(R_1\vec{a}_1) \cdots g(R_1\vec{a}_k) \tilde{g}(R_2\vec{a}_1) \cdots \tilde{g}(R_2\vec{a}_k) \\ &= \sum_{R_1 \in \mathbb{Z}^{d_1 \times m}} g(R_1\vec{a}_1) \cdots g(R_1\vec{a}_k) \sum_{R_2 \in \mathbb{Z}^{d_2 \times m}} \tilde{g}(R_2\vec{a}_1) \cdots \tilde{g}(R_2\vec{a}_k) = \sigma_L(g)\sigma_L(\tilde{g}). \end{aligned}$$

□

4. PROOF OF THEOREM 1.8

4.1. Introduction. In this section, we prove Theorem 1.8 and Theorem 1.9. We prove Theorem 1.8 by showing that, for all $2 \times k$ linear systems L with $s(L) = k - 1$, there exists a Fourier template g that satisfies the condition in Theorem 3.5.

The existence of such Fourier template will be given by a probabilistic argument. Recall that for every $M \in \mathbb{N}$, we let $\llbracket -M, M \rrbracket$ denote the subset $\{-M, \dots, -1, 0, 1, \dots, M\} \subseteq \mathbb{Z}$. Define the random Fourier template $f_M : \mathbb{Z} \rightarrow \mathbb{C}$ by

$$\begin{cases} f_M(r) = \xi_r & r \in \llbracket -M, M \rrbracket \setminus \{0\} \\ f_M(r) = 0 & \text{otherwise,} \end{cases}$$

where every ξ_r is an independent random unit complex number, subject to the constraint that $\xi_r = \overline{\xi_{-r}}$.

We say that two numbers form a *cancelling pair* if they sum up to zero. The key observation is that, for all $a_1, \dots, a_t \in \mathbb{Z}$, we have

$$\mathbb{E}[f_M(a_1) \cdots f_M(a_t)] = \begin{cases} 1 & a_1, \dots, a_t \in \llbracket -M, M \rrbracket \setminus \{0\} \\ & \text{and } a_1, \dots, a_t \text{ can be partitioned into cancelling pairs} \\ 0 & \text{otherwise.} \end{cases} \quad (\dagger)$$

A result in [13] showed that any even length equation whose coefficients cannot be partitioned into cancelling pairs is uncommon over sufficiently large \mathbb{F}_p . This can be proved using a first moment argument on the above f_M : if $L = a_1x_1 + \cdots + a_kx_k$ is such an equation, then for every $r \in \mathbb{Z}$, we have $\mathbb{E}[f_M(a_1r) \cdots f_M(a_kr)] = 0$ as a_1, \dots, a_k cannot be partitioned into cancelling pairs. Thus, for sufficiently large M , we have $\mathbb{E}[\sigma_L(f_M)] = \mathbb{E}[\sum_{r \in \mathbb{Z}} f_M(a_1r) \cdots f_M(a_kr)] = 0$ while $\mathbb{P}[\sigma_L(f_M) > 0] > 0$, which means that there is a choice of f_M with $\sigma_L(f_M) < 0$.

When L is a $2 \times k$ linear system, it turns out that a first moment argument is not enough. For example, consider the linear system

$$L = \begin{pmatrix} 1 & 0 & -3 & -4 \\ 0 & 1 & 1 & 2 \end{pmatrix},$$

so that

$$\sigma_L(f_M) = \sum_{r_1, r_2 \in \mathbb{Z}} f_M(r_1) f_M(r_2) f_M(-3r_1 + r_2) f_M(-4r_1 + 2r_2).$$

By the above key observation (\dagger), we get that

$$\begin{aligned} \mathbb{E}[\sigma_L(f_M)] &= |\{(r_1, r_2) \in \mathbb{Z}^2 : r_1, r_2, -3r_1 + r_2, -4r_1 + 2r_2 \in \llbracket -M, M \rrbracket \setminus \{0\}, \\ &\quad r_1, r_2, -3r_1 + r_2, -4r_1 + 2r_2 \text{ can be partitioned into cancelling pairs}\}| \\ &= |\{(r_1, r_2) \in \mathbb{Z}^2 : r_1, r_2, -3r_1 + r_2, -4r_1 + 2r_2 \in \llbracket -M, M \rrbracket \setminus \{0\}, \\ &\quad -3r_1 + 2r_2 = 0 \\ &\quad \text{or } r_1 + r_2 = -7r_1 + 3r_2 = 0 \\ &\quad \text{or } -2r_1 + r_2 = -4r_1 + 3r_2 = 0\}| \\ &= \Theta(M) > 0. \end{aligned}$$

Therefore, to prove Theorem 1.8, a first moment argument is insufficient and we need a more careful analysis on the distribution of $\sigma_L(f_M)$.

4.2. First attempt: second moment method. The most natural next step is to try to extract more information about $\sigma_L(f_M)$ from higher moments. For the sake of concreteness, we continue to work with

$$L = \begin{pmatrix} 1 & 0 & -3 & -4 \\ 0 & 1 & 1 & 2 \end{pmatrix}.$$

First we compute the variance of $\sigma_L(f_M)$. Define

$$\begin{aligned} A_M &= \{(r_1, r_2) \in \mathbb{Z}^2 : r_1, r_2, -3r_1 + r_2, -4r_1 + r_2 \in \llbracket -M, M \rrbracket \setminus \{0\}, \\ &\quad r_1, r_2, -3r_1 + r_2, -4r_1 + r_2 \text{ cannot be partitioned into cancelling pairs}\}, \\ B_M &= \{(r_1, r_2) \in \mathbb{Z}^2 : r_1, r_2, -3r_1 + r_2, -4r_1 + r_2 \in \llbracket -M, M \rrbracket \setminus \{0\}, \\ &\quad r_1, r_2, -3r_1 + r_2, -4r_1 + r_2 \text{ can be partitioned into cancelling pairs}\}. \end{aligned}$$

Then we know that

- (1) $\sigma_L(f_M) = \sum_{(r_1, r_2) \in A_M \cup B_M} f_M(r_1) f_M(r_2) f_M(-3r_1 + r_2) f_M(-4r_1 + 2r_2)$;
- (2) If $(r_1, r_2) \in B_M$, then $f_M(r_1) f_M(r_2) f_M(-3r_1 + r_2) f_M(-4r_1 + 2r_2)$ is constant 1;
- (3) If $(r_1, r_2) \in A_M$, then $\mathbb{E}[f_M(r_1) f_M(r_2) f_M(-3r_1 + r_2) f_M(-4r_1 + 2r_2)] = 0$.

Thus, we have

$$\begin{aligned} \text{Var}[\sigma_L(f_M)] &= \mathbb{E}\left[\left(\sum_{(r_1, r_2) \in A_M} f_M(r_1) f_M(r_2) f_M(-3r_1 + r_2) f_M(-4r_1 + 2r_2)\right)^2\right] \\ &= \sum_{(r_1, r_2), (r'_1, r'_2) \in A_M} \mathbb{E}[f_M(r_1) f_M(r_2) f_M(-3r_1 + r_2) f_M(-4r_1 + 2r_2) \\ &\quad \cdot f_M(r'_1) f_M(r'_2) f_M(-3r'_1 + r'_2) f_M(-4r'_1 + 2r'_2)] \\ &\stackrel{(\dagger)}{=} |\{((r_1, r_2), (r'_1, r'_2)) \in A_M^2 : r_1, r_2, -3r_1 + r_2, -4r_1 + 2r_2, r'_1, r'_2, -3r'_1 + r'_2, -4r'_1 + 2r'_2 \\ &\quad \text{can be partitioned into cancelling pairs}\}| \\ &\geq |\{((r_1, r_2), (r'_1, r'_2)) \in A_M^2 : r'_1 = -r_1, r'_2 = -r_2\}| = \Omega(M^2). \end{aligned}$$

In particular, this implies that $\text{Var}[\sigma_L(f_M)] = \Omega(\mathbb{E}[\sigma_L(f_M)]^2)$, so the second moment argument also does not provide a sufficiently strong anticoncentration statement to give the desired conclusion.

4.3. Speculating on higher moments. However, the observation (\dagger) suggests a way to compute higher moments. In particular, the d -th moment of $\sigma_L(f_M)$ is number of d -tuples $((r_1^{(1)}, r_2^{(1)}), \dots, (r_1^{(d)}, r_2^{(d)})) \in A_M^d$ such that the union of multisets $\bigcup_{i=1}^d \{\{r_1^{(i)}, r_2^{(i)}, -3r_1^{(i)} + 2r_2^{(i)}, -4r_1^{(i)} + 2r_2^{(i)}\}\}$ can be partitioned into cancelling pairs.

When $d = 2t$ is even, this number is at least

$$\binom{2t}{2} \binom{2t-2}{2} \cdots \binom{4}{2} \binom{2}{2} A_M^t = \frac{(2t)!}{2^{t!}} (cM)^{2t}.$$

Here $\binom{2t}{2} \binom{2t-2}{2} \cdots \binom{4}{2} \binom{2}{2}$ is the number of ways to pair up $[2t]$ into t pairs $(i_1, i'_1), \dots, (i_t, i'_t)$, as in each such pairing, we can pick arbitrary $(r_1^{(i_j)}, r_2^{(i_j)}) \in A_M$ and set $(r_1^{(i'_j)}, r_2^{(i'_j)}) = -(r_1^{(i_j)}, r_2^{(i_j)})$ to meet the requirement. It is plausible to guess that this is the dominant term in the $(2t)$ -th moment, and this ansatz suggests that $\sigma_L(f_M)$ might be close to a normal random variable with mean $\Theta(M)$ and variance $\Theta(M^2)$.

However, if we want to prove this by the method of higher moments, then we need to show that the other ways to partition $\bigcup_{i=1}^d \{\{r_1^{(i)}, r_2^{(i)}, -3r_1^{(i)} + 2r_2^{(i)}, -4r_1^{(i)} + 2r_2^{(i)}\}\}$ into cancelling pairs correspond to lower order terms $o(M^d)$. There are $\frac{(4d)!}{2^{2d}(2d)!}$ ways to partition this set into pairs; although most of them cannot be a valid partition into cancelling pairs, such approach becomes computationally messy very quickly. This motivates us to find another way to prove that $\sigma_L(f_M)$ is close in distribution to a normal random variable, with information from low moments only.

4.4. Second attempt: Stein's method. As mentioned earlier, we would like to prove that $\sigma_L(f_M)$ is close in distribution to a normal random variable without appealing to high moment calculations. The method of moments fundamentally relies on the fact that a distribution is uniquely determined by its characteristic function. Another approach to distributional approximation is *Stein's method* in which we replace the characteristic function with a characterizing operator.

Lemma 4.1 (Stein's lemma). *Define the functional operator \mathcal{A} by $\mathcal{A}f(x) = f'(x) - xf(x)$. If a random variable Z is such that $\mathbb{E}\mathcal{A}f(Z) = 0$ for all absolutely continuous functions f with $\|f'\|_\infty < \infty$, then $Z \sim \mathcal{N}(0, 1)$.*

We say that \mathcal{A} is the *characterizing operator* of the standard normal. An immediate consequence of the above is that we can get quantitative distributional approximation using the characterizing operator.

Corollary 4.2 (Another form of Stein's lemma). *Let W be any random variable and $Z \sim \mathcal{N}(0, 1)$. For any family of functions \mathcal{H} , we have that*

$$\sup_{h \in \mathcal{H}} |\mathbb{E}h(W) - \mathbb{E}h(Z)| = \sup_{h \in \mathcal{H}} |\mathbb{E}[f'_h(W) - Wf_h(W)]|,$$

where f'_h is the solution to

$$f'_h(x) - xf_h(x) = h(x) - \mathbb{E}h(Z).$$

A working mathematician's form of the above corollary is the following:

Definition 4.3 (Wasserstein distance). For random variables $X \sim \mu$ and $Y \sim \nu$ defined on a probability space Ω , the *Wasserstein distance* between X, Y is defined by

$$d_{\text{Wass}}(X, Y) = \sup \left\{ \left| \int f d\mu - \int f d\nu \right| : f : \Omega \rightarrow \mathbb{R} \text{ is 1-Lipschitz} \right\}.$$

Corollary 4.4 ([21, Theorem 3.1]). *Define*

$$\mathcal{D} = \left\{ f \in \mathcal{C}^1(\mathbb{R}) : f' \text{ is absolutely continuous, } \|f\|_\infty \leq 1, \|f'\|_\infty \leq \sqrt{2/\pi}, \|f''\|_\infty \leq 2 \right\}.$$

If W is a random variable and $Z \sim \mathcal{N}(0, 1)$, then

$$d_{\text{Wass}}(W, Z) \leq \sup_{f \in \mathcal{D}} |\mathbb{E}[f'(W) - Wf(W)]|.$$

Note that in our previous example, two variables $f_M(r_1)f_M(r_2)f_M(-3r_1 + r_2)f_M(-4r_1 + 2r_2)$, $f_M(r'_1)f_M(r'_2)f_M(-3r'_1 + r'_2)f_M(-4r'_1 + 2r'_2)$ are independent unless r_1, r'_1, r_2, r'_2 satisfy certain linear relations. This is also the reason why we would expect Corollary 4.4 to be useful for us: it allows us to quantify the intuitive idea that a sum of only locally dependent random variables is approximately normal. More precisely, we have the following consequence of Corollary 4.4, which we formalize using the notion of *dependency graph*.

Definition 4.5 (Dependency graph). Given a graph $G = (V, E)$ and a collection of random variables $\{X_i\}_{i \in V}$, we say that G is a *dependency graph* for $\{X_i\}_{i \in V}$ if for any disjoint subsets $S, T \subseteq V$ with no edges between S and T , the two random vectors $(X_i)_{i \in S}$ and $(X_i)_{i \in T}$ are independent.

For a graph $G = (V, E)$ and vertex $u \in V$, we use N_u to denote the closed neighborhood of u ; that is, $N_u = \{v : uv \in E\} \cup \{u\}$.

Lemma 4.6. *Let $\{X_v\}_{v \in V}$ be a family of random variables with dependency graph $G = (V, E)$. Let $\rho = \sqrt{\text{Var}(\sum_v X_v)}$ and $W = \rho^{-1} \sum_{v \in V} X_v$. Suppose $\mathbb{E}X_v = 0$ for all $v \in V$. Then for $Z \sim \mathcal{N}(0, 1)$ and any $x \in \mathbb{R}$, we have*

$$\begin{aligned} & |\mathbb{P}[W \leq x] - \mathbb{P}[Z \leq x]|^2 \\ & \leq 2/\pi \cdot \rho^{-3} \sum_{u \in V} \left(3 \sum_{v, w \in N_u} |\mathbb{E}[X_u X_v X_w]| + 4 \sum_{v \in N_u} |\mathbb{E}[X_u X_v]| \cdot \mathbb{E} \left[\left| \sum_{w \in N_u \cup N_v} X_w \right| \right] \right). \end{aligned}$$

A proof of Lemma 4.6 (which is effectively a modified version of [4, Theorem 3.1]) is provided in Appendix A.

4.5. Proof of Theorem 1.8. Suppose $k \geq 4$ is even, and L is a $2 \times k$ linear system with $s(L) = k - 1$. To prove Theorem 1.8, by Theorem 3.5, it suffices to construct a Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ such that $\sigma_L(g) < 0$.

Suppose L takes the form

$$L = \begin{pmatrix} a_1 & \cdots & a_k \\ b_1 & \cdots & b_k \end{pmatrix}.$$

Since $s(L) = k - 1$, the column vectors $(a_1, b_1), \dots, (a_k, b_k) \in \mathbb{Z}^2$ are pairwise linearly independent. For any Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$, we have

$$\sigma_L(g) = \sum_{r_1, r_2 \in \mathbb{Z}} g(a_1 r_1 + b_1 r_2) \cdots g(a_k r_1 + b_k r_2).$$

The key idea for proving Theorem 1.8 is to combine Theorem 3.5 with a probabilistic argument.

Proposition 4.7. *Let L be a $2 \times k$ linear system with $k \geq 4$ even and $s(L) = k - 1$. For every $M \in \mathbb{N}$, define the random Fourier template $f_M : \mathbb{Z} \rightarrow \mathbb{C}$ by*

$$\begin{cases} f_M(r) = \xi_r & r \in [-M, M] \setminus \{0\} \\ f_M(r) = 0 & \text{otherwise,} \end{cases}$$

where every ξ_r is a uniformly random unit complex number, subject to $\xi_r = \overline{\xi_{-r}}$ but independent otherwise. Then $\mathbb{P}[\sigma_L(f_M) < 0] > 0$ for all sufficiently large M .

Theorem 1.8 then directly follows from Proposition 4.7 and Theorem 3.5.

Proof of Theorem 1.8. Let L be a $2 \times k$ linear system with $k \geq 4$ even and $s(L) = k - 1$. By Proposition 4.7, there exists a Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ such that $\sigma_L(g) < 0$. By Theorem 3.5, L is uncommon over \mathbb{F}_p for all sufficiently large p . \square

It remains to show Proposition 4.7.

Proof of Proposition 4.7. Fix a $2 \times k$ linear system L with $k \geq 4$ even and $s(L) = k - 1$. Choose $\lambda \in \mathbb{N}$ sufficiently large such that, for all $M \in \mathbb{N}$, if $a_1 r_1 + b_1 r_2, \dots, a_k r_1 + b_k r_2 \in \llbracket -M, M \rrbracket$, then $-\lambda M \leq r_1, r_2 \leq \lambda M$.

Let $N = \lambda M$. Since f_M is always supported on $\llbracket -M, M \rrbracket \setminus \{0\}$, we have

$$\sigma_L(f_M) = \sum_{r_1, r_2 \in \mathbb{Z}} f_M(a_1 r_1 + a_1 r_2) \cdots f_M(a_k r_1 + b_k r_2) = \sum_{r_1, r_2 \in \llbracket -N, N \rrbracket} f_M(a_1 r_1 + a_1 r_2) \cdots f_M(a_k r_1 + b_k r_2).$$

For every multiset $S \subseteq \mathbb{Z}$, define the associated random variable

$$X_S = \prod_{s \in S} f_M(s),$$

where the product goes over all $s \in S$ counting multiplicities. Note that X_S has the following properties:

- (i) $X_S X_{S'} = X_{S \cup S'}$.
- (ii)

$$\begin{cases} X_S = 1 & S \subseteq \llbracket -M, M \rrbracket \setminus \{0\}, S \text{ can be partitioned into cancelling pairs,} \\ \mathbb{E} X_S = 0 & \text{otherwise.} \end{cases}$$

- (iii) Let \mathcal{S} and \mathcal{S}' be two collections of subsets of \mathbb{Z} . If $\pm \bigcup_{S \in \mathcal{S}} S$ and $\pm \bigcup_{S' \in \mathcal{S}'} S'$ are disjoint, then the two random vectors $(X_S)_{S \in \mathcal{S}}$ and $(X_{S'})_{S' \in \mathcal{S}'}$ are independent.

For every $(r_1, r_2) \in \llbracket -N, N \rrbracket^2$, define the associated multiset

$$S_{(r_1, r_2)} = \{\{a_1 r_1 + b_1 r_2, \dots, a_k r_1 + b_k r_2\}\}.$$

We simply write $X_{(r_1, r_2)}$ to denote $X_{S_{(r_1, r_2)}}$. Thus, we have

$$\sigma_L(f_M) = \sum_{r_1, r_2 \in \llbracket -N, N \rrbracket} f_M(a_1 r_1 + a_1 r_2) \cdots f_M(a_k r_1 + b_k r_2) = \sum_{r_1, r_2 \in \llbracket -N, N \rrbracket} X_{(r_1, r_2)}.$$

Consider the partition $\llbracket -N, N \rrbracket^2 = A_N \cup B_N$ defined by

$$\begin{aligned} A_N &= \{(r_1, r_2) \in \llbracket -N, N \rrbracket^2 : S_{(r_1, r_2)} \not\subseteq \llbracket -M, M \rrbracket \setminus \{0\} \\ &\quad \text{or } S_{(r_1, r_2)} \text{ cannot be partitioned into cancelling pairs}\}, \\ B_N &= \{(r_1, r_2) \in \llbracket -N, N \rrbracket^2 : S_{(r_1, r_2)} \subseteq \llbracket -M, M \rrbracket \setminus \{0\} \\ &\quad \text{and } S_{(r_1, r_2)} \text{ can be partitioned into cancelling pairs}\}. \end{aligned}$$

For all $(r_1, r_2) \in B_N$, we always have $X_{(r_1, r_2)} = 1$. Therefore we have

$$\sigma_L(f_M) = |B_N| + \sum_{(r_1, r_2) \in A_N} X_{(r_1, r_2)}.$$

We study the distribution of $\sum_{(r_1, r_2) \in A_N} X_{(r_1, r_2)}$ using Lemma 4.6. Let G be a dependency graph for the family of variables $\{X_{(r_1, r_2)} : (r_1, r_2) \in A_N\}$, defined by

$$\begin{aligned} V(G) &= A_N, \\ E(G) &= \{\{(r_1, r_2), (r'_1, r'_2)\} : \pm S_{(r_1, r_2)} \cap \pm S_{(r'_1, r'_2)} \neq \emptyset\} \subseteq \binom{A_N}{2}. \end{aligned}$$

Recall from Definition 4.5 that $N_{(r_1, r_2)}$ denotes the closed neighborhood of (r_1, r_2) in G , which in this case equals

$$\{(r'_1, r'_2) \in A_N : \pm S_{(r_1, r_2)} \cap \pm S_{(r'_1, r'_2)} \neq \emptyset\}.$$

We will verify the following properties of $\sum_{(r_1, r_2) \in A_N} X_{(r_1, r_2)}$. In the asymptotic bounds below, we allow the constants in $O(\cdot)$ and $\Omega(\cdot)$ to depend on k , as L is fixed.

Claim 4.8.

- (1) $\mathbb{E}[\sum_{u \in A_N} X_u] = 0$, and $\text{Var}(\sum_{u \in A_N} X_u) = \Omega(N^2)$;
- (2) For every $u \in A_N$, $\sum_{v \in A_N} \mathbb{E}[X_u X_v] = O(1)$;
- (3) $\sum_{u, v, w \in A_N} \mathbb{E}[X_u X_v X_w] = O(N^2)$;
- (4) For every $u, v \in A_N$, $\sum_{w_1, w_2 \in N_u \cup N_v} \mathbb{E}[X_{w_1} \overline{X_{w_2}}] = O(N)$.

We first prove Proposition 4.7 assuming Claim 4.8. Let $Z \sim \mathcal{N}(0, 1)$ be the normal distribution and $\rho = \sqrt{\text{Var}(\sum_{u \in A_N} X_u)} = \Omega(N)$ (per Claim 4.8(1)). We know from Claim 4.8(4) that

- (5) For every $u, v \in A_N$, by applying the Cauchy–Schwarz inequality, we have

$$\begin{aligned} (\mathbb{E}[\sum_{w \in N_u \cup N_v} X_w])^2 &\leq \mathbb{E}[\sum_{w \in N_u \cup N_v} X_w]^2 = \mathbb{E}[(\sum_{w \in N_u \cup N_v} X_w)(\sum_{w \in N_u \cup N_v} \overline{X_w})] \\ &= \sum_{w_1, w_2 \in N_u \cup N_v} \mathbb{E}[X_{w_1} \overline{X_{w_2}}] = O(N). \end{aligned}$$

Let $W = \rho^{-1} \sum_{u \in A_N} X_u$. By Lemma 4.6, for $Z \sim \mathcal{N}(0, 1)$ and all $x \in \mathbb{R}$, we have

$$\begin{aligned} &\pi/2 \cdot |\mathbb{P}[W \leq x] - \mathbb{P}[Z \leq x]|^2 \\ &\leq \rho^{-3} \sum_{u \in A_N} (3 \sum_{v, w \in N_u} |\mathbb{E}[X_u X_v X_w]| + 4 \sum_{v \in N_u} |\mathbb{E}[X_u X_v]| \cdot \mathbb{E}[\sum_{w \in N_u \cup N_v} X_w]) \\ &\stackrel{(*)}{\leq} \rho^{-3} (3 \sum_{u, v, w \in A_N} \mathbb{E}[X_u X_v X_w] + 4 \sum_{u, v \in A_N} \mathbb{E}[X_u X_v] \cdot \mathbb{E}[\sum_{w \in N_u \cup N_v} X_w]) \\ &\leq O(N^{-3})(O(N^2) + O(N^{1/2})) \sum_{u, v \in A_N} \mathbb{E}[X_u X_v] \tag{1}(3)(5) \\ &\leq O(N^{-3})(O(N^2) + O(N^{5/2})) = O(N^{-1/2}), \tag{2} \end{aligned}$$

where $(*)$ is due to the fact that (recall properties (i)(ii) at the beginning of proof)

$$0 \leq \mathbb{E}[X_u X_v], \mathbb{E}[X_u X_v X_w] \leq 1 \quad \text{for all } u, v, w \in A_N.$$

Since $\sigma_L(f_M) = |B_N| + \sum_{v \in A_N} X_v$, we have

$$\mathbb{P}(\sigma_L(f_M) < 0) = \mathbb{P}(\sum_{v \in A_N} X_v < -|B_N|) = \mathbb{P}(W < -|B_N|/\rho),$$

and therefore

$$|\mathbb{P}(\sigma_L(f_M) < 0) - \mathbb{P}(Z < -|B_N|/\rho)| = O(N^{-1/4}).$$

Since $|B_N| = O(N)$ and $\rho = \Omega(N)$, we have $|B_N|/\rho = O(1)$. Thus $\mathbb{P}(Z < -|B_N|/\rho) = c > 0$, and therefore $\mathbb{P}(\sigma_L(f_M) < 0) \geq c - O(N^{-1/4}) > 0$ for all N sufficiently large. \square

It remains to verify Claim 4.8.

Proof of Claim 4.8.

(1) For every $u = (r_1, r_2) \in A_N$, since $S_{(r_1, r_2)}$ cannot be partitioned into cancelling pairs, we have $\mathbb{E}X_{(r_1, r_2)} = 0$. Moreover, we have

$$\begin{aligned} \mathbb{E}[(\sum_{u \in A_N} X_u)^2] &= \sum_{u, v \in A_N} \mathbb{E}[X_u X_v] = \sum_{u, v \in A_N} \mathbb{E}[X_{S_u \cup S_v}] \\ &= |\{(u, v) \in A_N^2 : S_u \cup S_v \subseteq \llbracket -M, M \rrbracket \setminus \{0\}, S_u \cup S_v \text{ can be partitioned into cancelling pairs}\}| \\ &\geq |\{((r_1, r_2), (r'_1, r'_2)) \in A_N^2 : S_{(r_1, r_2)} \subseteq \llbracket -M, M \rrbracket \setminus \{0\}, r'_1 = -r_1, r'_2 = -r_2\}| \\ &= \Omega(N^2). \end{aligned}$$

This gives $\text{Var}(\sum_{u \in A_N} X_u) = \Omega(N^2)$.

(2) For all $u = (r_1, r_2) \in A_N$, we have

$$\begin{aligned} \sum_{v \in A_N} \mathbb{E}[X_u X_v] &= \sum_{v \in A_N} \mathbb{E}[X_{S_u \cup S_v}] \\ &\leq |\{v \in A_N : S_u \cup S_v \text{ can be partitioned into cancelling pairs}\}|. \end{aligned}$$

Recall that S_u itself cannot be partitioned into cancelling pairs. Therefore, if $S_u \cup S_v$ can be partitioned into cancelling pairs, then $v = (r'_1, r'_2)$ must satisfy equations of the form

$$\begin{aligned} a_i r'_1 + b_i r'_2 &= -(a_i r_1 + b_i r_2) \\ a_{j'} r'_1 + b_{j'} r'_2 &= -(a_{j'} r_1 + b_{j'} r_2) \end{aligned}$$

with $i \neq j$ and $i' \neq j'$. Since $(a_{i'}, b_{i'})$ and $(a_{j'}, b_{j'})$ are *linearly independent* due to the girth condition $s(L) = k - 1$, every such pair of equations has at most one solution. Moreover, there are at most k^4 such pairs. Thus, for every $u \in A_N$, we get that

$$|\{v \in A_N : S_u \cup S_v \text{ can be partitioned into cancelling pairs}\}| = O(1).$$

Proofs of (3)(4) have similar ideas as in (2).

(3) For every triple $(u, v, w) \in A_N^3$, define a graph $H_{u, v, w}$ on the multiset $S_u \cup S_v \cup S_w$, such that $s_1 \sim s_2$ in $H_{u, v, w}$ if and only if $s_1 + s_2 = 0$. Observe that

$$\begin{aligned} \sum_{u, v, w \in A_N} \mathbb{E}[X_u X_v X_w] &= \sum_{u, v, w \in A_N} \mathbb{E}[X_{S_u \cup S_v \cup S_w}] \\ &= |\{(u, v, w) \in A_N^3 : S_u, S_v, S_w \subseteq \llbracket -M, M \rrbracket \setminus \{0\}, \\ &\quad S_u \cup S_v \cup S_w \text{ can be partitioned into cancelling pairs}\}| \\ &= |\{(u, v, w) \in A_N^3 : S_u, S_v, S_w \subseteq \llbracket -M, M \rrbracket \setminus \{0\}, \\ &\quad H_{u, v, w} \text{ has a perfect matching}\}|. \end{aligned}$$

Note that every graph $H_{u, v, w}$ has $|S_u \cup S_v \cup S_w| = 3k$ vertices. We prove that, for every perfect matching \mathcal{M} on $3k$ vertices, there are $O(N^2)$ choices of $(u, v, w) \in A_N^3$ such that $S_u \cup S_v \cup S_w \subseteq \llbracket -M, M \rrbracket \setminus \{0\}$ and $H_{u, v, w}$ contains \mathcal{M} as a subgraph. Since there are finitely many perfect matchings on $3k$ vertices, we get that

$$|\{(u, v, w) \in A_N^3 : S_u, S_v, S_w \subseteq \llbracket -M, M \rrbracket \setminus \{0\}, H_{u, v, w} \text{ has a perfect matching}\}| = O(N^2).$$

Fix a perfect matching \mathcal{M} on $3k$ vertices. Suppose $(u, v, w) \in A_N^3$ satisfies $S_u \cup S_v \cup S_w \subseteq \llbracket -M, M \rrbracket \setminus \{0\}$, and $H_{u, v, w}$ contains \mathcal{M} as a subgraph. For $U, V \subseteq S_u \cup S_v \cup S_w$, let $e_{\mathcal{M}}(U, V)$ denote the number of edges in \mathcal{M} whose vertices intersect both U and V . Since none of S_u, S_v, S_w can be partitioned into cancelling pairs, we must have

$$e_{\mathcal{M}}(S_u, S_v \cup S_w) \geq 2, \quad e_{\mathcal{M}}(S_v, S_u \cup S_w) \geq 2, \quad e_{\mathcal{M}}(S_w, S_u \cup S_v) \geq 2.$$

Up to permuting u, v, w , we have the following two cases.

Case 1. $e_{\mathcal{M}}(S_u, S_v) \geq 2$. In this case, \mathcal{M} gives $i \neq j$ and $i' \neq j'$ such that

$$a_{i'}r'_1 + b_{i'}r'_2 = -(a_i r_1 + b_i r_2), \quad a_{j'}r'_1 + b_{j'}r'_2 = -(a_j r_1 + b_j r_2)$$

for $u = (r_1, r_2)$ and $v = (r'_1, r'_2)$. Since $(a_{i'}, b_{i'})$ and $(a_{j'}, b_{j'})$ are linearly independent, $v = (r'_1, r'_2)$ is uniquely determined by $u = (r_1, r_2)$. Since $e_{\mathcal{M}}(S_w, S_u \cup S_v) \geq 2$, \mathcal{M} also gives $i'' \neq j''$ and i°, i^* such that

$$a_{i''}r''_1 + b_{i''}r''_2 = -(a_{i^\circ}r_1^\circ + b_{i^\circ}r_2^\circ), \quad a_{j''}r''_1 + b_{j''}r''_2 = -(a_{i^*}r_1^* + b_{i^*}r_2^*)$$

for $w = (r''_1, r''_2)$ and $(r_1^\circ, r_2^\circ), (r_1^*, r_2^*) \in \{(r_1, r_2), (r'_1, r'_2)\} = \{u, v\}$. Since $(a_{i''}, b_{i''})$ and $(a_{j''}, b_{j''})$ are linearly independent, $w = (r''_1, r''_2)$ is uniquely determined by $u = (r_1, r_2)$ and $v = (r'_1, r'_2)$. Therefore, the number of choices for u, v, w is at most the number of choices for $u = (r_1, r_2)$, which is $O(N^2)$.

Case 2. $e_{\mathcal{M}}(S_u, S_v) = e_{\mathcal{M}}(S_u, S_w) = e_{\mathcal{M}}(S_v, S_w) = 1$. In this case, \mathcal{M} gives $i \neq j$, $i' \neq j'$ and $i'' \neq j''$ such that

$$\begin{aligned} a_{i'}r'_1 + b_{i'}r'_2 &= -(a_i r_1 + b_i r_2), \\ a_{i''}r''_1 + b_{i''}r''_2 &= -(a_j r_1 + b_j r_2), \\ a_{j''}r''_1 + b_{j''}r''_2 &= -(a_{j'}r'_1 + b_{j'}r'_2). \end{aligned}$$

Since $S_u \cup S_v \cup S_w \subseteq \llbracket -M, M \rrbracket \setminus \{0\}$, none of the above three equations can be 0.

Now since $|S_u| = |S_v| = |S_w| = k \geq 4$, \mathcal{M} also has at least one edge within each of S_u, S_v, S_w . Therefore, \mathcal{M} in particular gives i', k', ℓ' distinct, i'', k'', ℓ'' distinct, and $i \neq j$ such that

$$\begin{aligned} a_{i'}r'_1 + b_{i'}r'_2 &= -(a_i r_1 + b_i r_2), \\ a_{i''}r''_1 + b_{i''}r''_2 &= -(a_j r_1 + b_j r_2), \\ a_{k'}r'_1 + b_{k'}r'_2 &= -(a_{\ell'}r'_1 + b_{\ell'}r'_2), \\ a_{k''}r''_1 + b_{k''}r''_2 &= -(a_{\ell''}r''_1 + b_{\ell''}r''_2). \end{aligned}$$

Since

- (i) $(a_{k'}, b_{k'})$ and $(a_{\ell'}, b_{\ell'})$ are linearly independent,
- (ii) $(a_{k''}, b_{k''})$ and $(a_{\ell''}, b_{\ell''})$ are linearly independent,
- (iii) $a_i r_1 + b_i r_2$ and $a_j r_1 + b_j r_2$ are nonzero,

we get that $v = (r'_1, r'_2)$ and $w = (r''_1, r''_2)$ are uniquely determined by $u = (r_1, r_2)$. Hence the number of choices for u, v, w is at most the number of choices for $u = (r_1, r_2)$, which is $O(N^2)$.

(4) For every $u = (r_1, r_2), v = (r'_1, r'_2) \in A_N$, we have

$$\begin{aligned} \sum_{w_1, w_2 \in N_u \cup N_v} \mathbb{E}[X_{w_1} \overline{X_{w_2}}] &= \sum_{w_1, w_2 \in N_u \cup N_v} \mathbb{E}[X_{w_1} X_{-w_2}] \\ &\leq |\{(w_1, w_2) \in (N_u \cup N_v)^2 : S_{w_1} \cup S_{-w_2} \text{ can be partitioned into cancelling pairs}\}|. \end{aligned}$$

Recall that each of S_{w_1}, S_{w_2} cannot be partitioned into cancelling pairs. Therefore, if $S_{w_1} \cup S_{-w_2}$ can be partitioned into cancelling pairs, then $w_1 = (r''_1, r''_2), w_2 = (r'''_1, r'''_2)$ must satisfy two equations

$$a_{i''}r''_1 + b_{i''}r''_2 = -(a_{i'''}r'''_1 + b_{i'''}r'''_2), \quad a_{j''}r''_1 + b_{j''}r''_2 = -(a_{j'''}r'''_1 + b_{j'''}r'''_2)$$

with $i'' \neq j''$ and $i''' \neq j'''$. Again, by linear independence of the columns of L , for fixed choices of i'', j'', i''', j''' , each one of w_1, w_2 will uniquely determine another.

We now look at the number of possible choices for w_1 . Since $w_1 \in N_u \cup N_v$, by Definition 4.5, $w_1 = (r''_1, r''_2)$ must also satisfy

$$a_{\ell}r''_1 + b_{\ell}r''_2 = a_{\ell^\circ}r_1^\circ + b_{\ell^\circ}r_2^\circ \quad \text{or} \quad a_{\ell}r''_1 + b_{\ell}r''_2 = -(a_{\ell^\circ}r_1^\circ + b_{\ell^\circ}r_2^\circ)$$

for some ℓ, ℓ° and $(r_1^\circ, r_2^\circ) \in \{(r_1, r_2), (r'_1, r'_2)\}$. Given ℓ, ℓ° and (r_1°, r_2°) , since we cannot have $a_\ell = b_\ell = 0$, the number of possible choices for w_1 is at most $O(N)$.

Overall, since there are finitely many choices for $i'', j'', i''', j''', \ell, \ell^\circ$ and $(r_1^\circ, r_2^\circ) \in \{(r_1, r_2), (r_1', r_2')\}$, we get that

$$|\{(w_1, w_2) \in (N_u \cup N_v)^2 : S_{w_1} \cup S_{-w_2} \text{ can be partitioned into cancelling pairs}\}| = O(N).$$

□

4.6. Proof of Theorem 1.9. Let L be a linear system with $c = c(L)$. Recall from Remark 2.3 that all elements in $\{L_B : B \in \mathcal{C}(L)\}$ are either length- c linear equations or $2 \times c$ systems with girth $c - 1$.

Consider the partition $\mathcal{C}(L) = \mathcal{C}_1 \cup \mathcal{C}_2$ defined by

$$\begin{aligned} \mathcal{C}_1 &= \{B \in \mathcal{C}(L) : L_B \text{ is length-}c \text{ linear equation}\}, \\ \mathcal{C}_2 &= \{B \in \mathcal{C}(L) : L_B \text{ is } 2 \times c \text{ linear system}\}. \end{aligned}$$

The following lemma is essentially due to Altman [1, Theorem 5.2], although Altman did not work in the language of Fourier templates. For the sake of completeness, we provide its proof here.

Proposition 4.9. *Let L be a linear system with $s(L)$ odd. Suppose there exists a Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ such that $\sum_{B \in \mathcal{C}_2} \sigma_{L_B}(g) < 0$. Then L is uncommon over all sufficiently large p .*

Proof. Suppose $\sum_{B \in \mathcal{C}_2} \sigma_{L_B}(g) < 0$ for some Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$. By Theorem 3.5, this gives a function $f : \mathbb{F}_p \rightarrow \mathbb{R}$ with $\mathbb{E}f = 0$ and $\sum_{B \in \mathcal{C}_2} t_{L_B}(f) < 0$.

However, to show that L is uncommon over all sufficiently large p , we should take into account subsystems in \mathcal{C}_1 as well. That is, we wish to show that there is some function $h : \mathbb{F}_p^n \rightarrow \mathbb{R}$ with $\mathbb{E}h = 0$ and $\sum_{B \in \mathcal{C}(L)} t_{L_B}(h) < 0$, so that we can utilize Proposition 2.4.

To do so, we take a “tensoring” of f with the indicator function of 0. Namely, define $h : \mathbb{F}_p^2 \rightarrow \mathbb{R}$ by $h(y, z) = f(y)\mathbf{1}_{z=0}$. Since f has mean 0, so does h . Moreover, for any $B \in \mathcal{C}(L)$, we have (with $c = c(L)$)

$$t_{L_B}(h) = \mathbb{E}_{\substack{\mathbf{x} \in (\mathbb{F}_p^2)^c \\ L_B \mathbf{x} = 0}} h(\mathbf{x}_1) \cdots h(\mathbf{x}_c) = \mathbb{E}_{\substack{y \in \mathbb{F}_p^c \\ L_B y = 0}} f(y_1) \cdots f(y_c) \cdot \mathbb{E}_{\substack{\mathbf{z} \in \mathbb{F}_p^c \\ L_B \mathbf{z} = 0}} \mathbf{1}_{z_1=0} \cdots \mathbf{1}_{z_c=0} = t_{L_B}(f)t_{L_B}(\mathbf{1}_{z=0}).$$

Observe that if L_B is a length- c equation, then it has p^{c-1} solutions $\mathbf{z} \in \mathbb{F}_p^c$; if L_B is a $2 \times c$ linear system, then it has p^{c-2} solutions $\mathbf{z} \in \mathbb{F}_p^c$. Thus, we have

$$t_{L_B}(\mathbf{1}_{z=0}) = \begin{cases} p^{1-c} & B \in \mathcal{C}_1 \\ p^{2-c} & B \in \mathcal{C}_2. \end{cases}$$

Therefore, given that $\sum_{B \in \mathcal{C}_2} t_{L_B}(f) < 0$, we have

$$\sum_{B \in \mathcal{C}(L)} t_{L_B}(h) = \sum_{B \in \mathcal{C}_1} t_{L_B}(h) + \sum_{B \in \mathcal{C}_2} t_{L_B}(h) = p^{1-c} \sum_{B \in \mathcal{C}_1} t_{L_B}(f) + p^{2-c} \sum_{B \in \mathcal{C}_2} t_{L_B}(f),$$

which is negative for p sufficiently large. Thus, by Proposition 2.4, L is uncommon over all sufficiently large p . □

We are now ready to prove Theorem 1.9.

Proof. By Proposition 4.9, it suffices to construct a Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ such that $\sum_{B \in \mathcal{C}_2} \sigma_{L_B}(g) < 0$. Suppose $\mathcal{C}_2(L) = \{B_1, \dots, B_\ell\}$, with

$$L_{B_j} = \begin{pmatrix} a_{j,1} & \cdots & a_{j,c} \\ b_{j,1} & \cdots & b_{j,c} \end{pmatrix} \quad \text{for every } j \in [\ell].$$

Define the random Fourier template $f_M : \mathbb{Z} \rightarrow \mathbb{C}$ as in Proposition 4.7. Also, choose $\lambda \in \mathbb{N}$ sufficiently large such that if $a_{j,1}r_1 + b_{j,1}r_2, \dots, a_{j,k}r_1 + b_{j,k}r_2$ lie in $[-M, M] \setminus \{0\}$ for all $j \in [\ell]$, then $-\lambda M \leq r_1, r_2 \leq \lambda M$.

For every $(r_1, r_2) \in \llbracket -N, N \rrbracket$, define the associated multisets

$$\begin{aligned} S_{(r_1, r_2)}^1 &= \{a_{1,1}r_1 + b_{1,1}r_2, \dots, a_{1,k}r_1 + b_{1,k}r_2\}, \\ &\vdots \\ S_{(r_1, r_2)}^\ell &= \{a_{\ell,1}r_1 + b_{\ell,1}r_2, \dots, a_{\ell,k}r_1 + b_{\ell,k}r_2\}. \end{aligned}$$

We then have

$$\sum_{B \in \mathcal{C}_2} \sigma_{L_B}(f_M) = \sum_{r_1, r_2 \in \llbracket -N, N \rrbracket} \sum_{j=1}^{\ell} X_{S_{(r_1, r_2)}^j}.$$

Let ρ denote the standard deviation of $\sum_{r_1, r_2 \in \llbracket -N, N \rrbracket} \sum_{j=1}^{\ell} X_{S_{(r_1, r_2)}^j}$, and define

$$W = \rho^{-1} \sum_{r_1, r_2 \in \llbracket -N, N \rrbracket} \sum_{j=1}^{\ell} X_{S_{(r_1, r_2)}^j}.$$

An argument almost identical to Proposition 4.7 shows that $\mathbb{P}\left(\sum_{B \in \mathcal{C}_2} \sigma_{L_B}(f_M) < 0\right) > 0$ for all M sufficiently large. In particular, there exists Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ such that $\sum_{B \in \mathcal{C}_2} \sigma_{L_B}(g) < 0$. Finally, by Proposition 4.9, we know that L is uncommon over \mathbb{F}_p for all p sufficiently large. \square

5. PROOF OF THEOREM 1.10

5.1. Introduction. In this section, we prove Theorem 1.10. Let L be an irredundant 2×5 linear system on variables x_1, \dots, x_5 . The bulk of Theorem 1.10 considers the case $s(L) = 4$. Under the assumption $s(L) = 4$, we know that the elements in $\{L_B : B \in \mathcal{C}(L)\}$ of L are five length-4 linear equations, supported on five different 4-subsets of $\{x_1, \dots, x_5\}$.

Let L_1, \dots, L_5 denote these five equations. By Theorem 3.5, to show that L is uncommon, it suffices to find a Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{L_i}(g) < 0$. Since each $\sigma_{L_i}(g)$ is defined by both g and the coefficients in L_i , to simplify our further discussion, we give special names to the set of coefficients in each L_i .

Setup 5.1. Suppose L is an irredundant 2×5 linear system with $s(L) = 4$. For every $i = 1, \dots, 5$, we admit the following notation:

- Let $L_i \mathbf{x} := a_{i,1}x_1 + \dots + a_{i,5}x_5$ be the unique (up to constant multiplication) equation in the row span of L , such that $a_{i,i} = 0$.
- Let $A_i = \{a_{i,j} : j \in [5], j \neq i\}$ be the multiset of nonzero coefficients in L_i .
- For every Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$, define the associated function $g_{A_i} : \mathbb{Z}^d \rightarrow \mathbb{C}$ by

$$g_{A_i}(r) := \prod_{a \in A_i} g(ar)$$

and the associated sum $\sigma_{A_i}(g)$ by

$$\sigma_{A_i}(g) = \sum_{r \in \mathbb{Z}^d} g_{A_i}(r),$$

so that $g_{A_i} = g_{L_i}$ and $\sigma_{A_i}(g) = \sigma_{L_i}(g)$ as in Definition 3.3.

Suppose we wish to show that L is uncommon, and hope to find some Fourier template g with $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$. This is easy if none of L_1, \dots, L_5 is common, as a random Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ has $\sum_{i=1}^5 \mathbb{E}[\sigma_{A_i}(g)] = 0$. However, if some equation L_{i_0} among L_1, \dots, L_5 is common, with $A_{i_0} = \{1, -1, \lambda, -\lambda\}$, then $\sigma_{A_{i_0}}(g) = \sum_r |g(r)|^2 |g(\lambda r)|^2$ must be a positive contribution to $\sum_{i=1}^5 \sigma_{A_i}(g)$. In this case, we have to carefully pick some g so that the other terms $\sigma_{A_i}(g)$, $i \neq i_0$ are “negative enough” to balance out this positive term. Now, if some L_{i_0} is an additive quadruple

$L_{i_0}\mathbf{x} = x_{j_1} + x_{j_2} - x_{j_3} - x_{j_4}$, then we have $A_{i_0} = \{1, 1, -1, -1\}$, and the sum $\sigma_{A_{i_0}}(g) = \sum_r |g(r)|^4$ would be even harder to balance out.

This motivates us to classify all the irredundant 2×5 linear systems L according to the number of common equations and additive quadruples among L_1, \dots, L_5 , and prove Theorem 1.10 under each case.

Lemma 5.2. *Let L be an irredundant 2×5 linear system, with $s(L) = 4$. Then there cannot be two additive quadruples among L_1, \dots, L_5 .*

Proof. Without loss of generality, suppose L_1 and L_2 are both additive quadruple. Up to isomorphism, L is one of

$$\begin{pmatrix} 0 & 1 & 1 & -1 & -1 \\ 1 & 0 & 1 & -1 & -1 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 1 & -1 & -1 \\ 1 & 0 & -1 & 1 & -1 \end{pmatrix}.$$

In either case, we have $s(L) \leq 3$, which is a contradiction. \square

Corollary 5.3. *Let L be an irredundant 2×5 linear system, with $s(L) = 4$. Then L satisfies exactly one of the following:*

- Case A. None of L_1, \dots, L_5 is an additive quadruple, and at least three of them are uncommon.*
- Case B. Exactly one of L_1, \dots, L_5 is an additive quadruple, and none of the others is common.*
- Case C. Exactly one of L_1, \dots, L_5 is an additive quadruple, and exactly one among the others is common but not an additive quadruple.*
- Case D. At least three of L_1, \dots, L_5 are common.*

Our strategy of finding the appropriate Fourier template g with $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$ differs in each case listed in Corollary 5.3. Before diving into the proof of Theorem 1.10, we provide some examples illustrating our main ideas in Cases A, B and C.

In Case A, mostly, we aim to pick a Fourier template g with $\sigma_{A_{i_0}}(g)$ “very negative” for some uncommon L_{i_0} , while keeping $\sum_{i \in [5], L_i \text{ common}} \sigma_{A_i}(g)$ relatively small.

Example 5.4 (Case A). Let L be the linear system

$$\begin{pmatrix} 1 & 3 & -1 & -3 & 0 \\ 2 & 3 & -3 & 0 & -2 \end{pmatrix}.$$

We then have

$$\begin{aligned} L_5\mathbf{x} &= x_1 + 3x_2 - x_3 - 3x_4 & A_5 &= \{1, 3, -1, -3\} \\ L_4\mathbf{x} &= 2x_1 + 3x_2 - 3x_3 - 2x_5 & A_4 &= \{2, 3, -3, -2\} \\ L_3\mathbf{x} &= x_1 + 6x_2 - 9x_4 + 2x_5 & A_3 &= \{1, 6, -9, 2\} \\ L_2\mathbf{x} &= x_1 - 2x_3 + 3x_4 - 2x_5 & A_2 &= \{1, -2, 3, -2\} \\ L_1\mathbf{x} &= 3x_2 + x_3 - 6x_4 + 2x_5 & A_1 &= \{3, 1, -6, 2\}. \end{aligned}$$

We wish to find a Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ such that

$$\begin{aligned} & \sum_{r \in \mathbb{Z}} (|g(r)|^2 |g(3r)|^2 + |g(2r)|^2 |g(3r)|^2 \\ & + g(r)g(6r)g(-9r)g(2r) + g(r)g(-2r)g(3r)g(-2r) + g(3r)g(r)g(-6r)g(2r)) = \sum_{i=1}^5 \sigma_{A_i}(g) < 0. \end{aligned}$$

Observe that the first two terms always have nonnegative contribution. This motivates us to find some g that makes one of the last three terms “very negative” without making the first two terms “too positive”.

For example, consider the Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ defined by

$$\begin{cases} g(2) = g(-2) = g(6) = g(-6) = g(9) = g(-9) = 1 \\ g(1) = g(-1) = -100 \\ g(r) = 0 \quad \text{otherwise.} \end{cases}$$

We then have

$$\sum_{r \in \mathbb{Z}} g(r)g(6r)g(-9r)g(2r) = g(1)g(6)g(-9)g(2) + g(-1)g(-6)g(9)g(-2) = -200,$$

while other terms

$$\begin{aligned} \sum_{r \in \mathbb{Z}} |g(r)|^2 |g(3r)|^2 &= |g(2)|^2 |g(6)|^2 + |g(-2)|^2 |g(-6)|^2 = 2, \\ \sum_{r \in \mathbb{Z}} |g(2r)|^2 |g(3r)|^2 &= |g(6)|^2 |g(9)|^2 + |g(-6)|^2 |g(-9)|^2 = 2, \\ \sum_{r \in \mathbb{Z}} g(r)g(-2r)g(3r)g(-2r) &= \sum_{r \in \mathbb{Z}} g(3r)g(r)g(-6r)g(2r) = 0 \end{aligned}$$

have small absolute value. Altogether, we get that $\sum_{i=1}^5 \sigma_{A_i}(g) = -196 < 0$.

We will discuss Case A further in Section 5.2.

In Case B, we continue to work with the Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$. However, the strategy in Case A fails in the presence of an additive quadruple, because setting $|g(r)|$ to be big for any particular $r \in \mathbb{Z}$ will make $|g(r)|^4$, the term contributed by the additive quadruple, blow-up the fastest.

As such, we need to crucially use the fact that we have at least four uncommon equations, say A_1, \dots, A_4 . In Case B, our construction is roughly in two steps:

- We construct our Fourier template to have support in some $S \subseteq \mathbb{Z}$ such that there are many nonvanishing terms in $\sum_{i=1}^4 \sum_{r \in S} g_{A_i}(r)$. We do this by taking S to be a *multiplicative grid*, illustrated in the example below.
- Next, we choose each $g(r)$ to be complex numbers with appropriate phases that align in a way to ensure that $\sum_{i=1}^4 \sigma_{A_i}(g)$ is a complex vector with negative real part.

The details of this procedure is also illustrated in the example below.

Example 5.5 (Case B). Let L be the 2×5 linear system

$$\begin{pmatrix} 1 & -1 & 1 & -1 & 0 \\ -2 & 4 & 3 & 0 & -9 \end{pmatrix}.$$

We then have

$$\begin{aligned} L_5 \mathbf{x} &= x_1 - x_2 + x_3 - x_4 & A_5 &= \{1, 1, -1, -1\} \\ L_4 \mathbf{x} &= -2x_1 + 4x_2 + 3x_3 - 9x_5 & A_4 &= \{-2, 4, 3, -9\} \\ L_3 \mathbf{x} &= 5x_1 - 7x_2 - 3x_4 + 9x_5 & A_3 &= \{5, -7, -3, 9\} \\ L_2 \mathbf{x} &= 2x_1 + 7x_3 - 4x_4 - 9x_5 & A_2 &= \{2, 7, -4, -9\} \\ L_1 \mathbf{x} &= 2x_2 + 5x_3 - 2x_4 - 9x_5 & A_1 &= \{2, 5, -2, -9\}. \end{aligned}$$

Again, we wish to find a Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ such that

$$\begin{aligned} \sum_{r \in \mathbb{Z}} (|g(r)|^4 + g(-2r)g(4r)g(3r)g(-9r) + g(5r)g(-7r)g(-3r)g(9r) \\ + g(2r)g(7r)g(-4r)g(-9r) + g(2r)g(5r)g(-2r)g(-9r)) = \sum_{i=1}^5 \sigma_{A_i}(g) < 0. \end{aligned}$$

Since L contains an additive quadruple, we cannot hope to use a construction similar to Example 5.4, as enlarging $|g(r)|$ for any particular r would make $\sum_{r \in \mathbb{Z}} |g(r)|^4$ the dominant term.

Instead, we build some Fourier template whose support is “almost closed” under multiplication by any coefficient in A_1, \dots, A_5 . Consider the *multiplicative grid* G defined by

$$G = \{2^{d_1} 3^{d_2} 5^{d_3} 7^{d_4} : d_1, \dots, d_4 \in \{0, 1, \dots\}\} \subseteq \mathbb{N}.$$

We can think of G as a four-dimensional grid, with each dimension corresponding to a prime divisor. Here we include $2, 3, 5, 7$ as they are precisely the prime divisors of elements in $A_1 \cup \dots \cup A_5$. Consequently, G has the nice property that $\pm 2G, \pm 3G, \pm 4G, \pm 5G, \pm 7G, \pm 9G \subseteq \pm G$; in other words, $\pm G$ is closed under multiplication by any coefficient in $A_1 \cup \dots \cup A_5$.

However, a Fourier template must have finite support, and therefore cannot be nonzero on the whole of $\pm G$. Therefore, we will first build some $g : \pm G \rightarrow \mathbb{C}$, and eventually replace it by its restriction on $\pm G_D$, where

$$G_D = \{2^{d_1} 3^{d_2} 5^{d_3} 7^{d_4} : d_1, \dots, d_4 \in \{0, 1, \dots, D\}\}$$

is a finite truncation of G .

Our $g : \pm G \rightarrow \mathbb{C}$ will be of the form

$$\begin{cases} g(2^{d_1} 3^{d_2} 5^{d_3} 7^{d_4}) = e((\theta_1 d_1 + \theta_2 d_2 + \theta_3 d_3 + \theta_4 d_4)t) \\ g(-2^{d_1} 3^{d_2} 5^{d_3} 7^{d_4}) = e(-(\theta_1 d_1 + \theta_2 d_2 + \theta_3 d_3 + \theta_4 d_4)t) \end{cases}$$

where $\theta_1, \dots, \theta_4 \in \mathbb{Q}$ and $t \in \mathbb{R}$. In other words, g is the composition of $e(\cdot)$ and a “linear map” from the multidimensional grid G to \mathbb{R} . This choice of g will ensure that the phases of $\sigma_{A_i}(g)$ line up in a way to ensure that $\sum_{i=2}^5 \sigma_{A_i}(g)$ is very negative.

Observe that $|g(r)| = 1$ for all $r \in \pm G$. Moreover, for all $x, y \in G$, we have $g(xy) = g(x)g(y)$ and $g(-xy) = g(-x)g(-y)$. Hence for all $r \in \pm G$, we have

$$g(-2r)g(4r)g(3r)g(-9r) = \begin{cases} g(-2)g(4)g(3)g(-9) \cdot g(r)^2 g(-r)^2 = e((\theta_1 - \theta_2)t) & r \in G \\ g(2)g(-4)g(-3)g(9) \cdot g(r)^2 g(-r)^2 = e((\theta_2 - \theta_1)t) & r \in -G. \end{cases}$$

Similarly, we have

$$\begin{aligned} g(5r)g(-7r)g(-3r)g(9r) &= \begin{cases} e((\theta_2 + \theta_3 - \theta_4)t) & r \in G \\ e((-\theta_2 - \theta_3 + \theta_4)t) & r \in -G \end{cases} \\ g(2r)g(7r)g(-4r)g(-9r) &= \begin{cases} e((-\theta_1 - 2\theta_2 + \theta_4)t) & r \in G \\ e((\theta_1 + 2\theta_2 - \theta_4)t) & r \in -G \end{cases} \\ g(2r)g(5r)g(-2r)g(-9r) &= \begin{cases} e((-\theta_2 + \theta_3)t) & r \in G \\ e((2\theta_2 - \theta_3)t) & r \in -G. \end{cases} \end{aligned}$$

Therefore, for all $r \in \pm G$, we have

$$\sum_{i=1}^5 \operatorname{Re}(g_{A_i}(r)) = 1 + \cos((\theta_1 - \theta_2)t) + \cos((\theta_2 + \theta_3 - \theta_4)t) + \cos((-\theta_1 - 2\theta_2 + \theta_4)t) + \cos((-\theta_2 + \theta_3)t).$$

Setting $\theta_1 = \theta_3 = 1$, $\theta_2 = \theta_4 = 0$ and $t = \pi$, we get that

$$\sum_{i=1}^5 \operatorname{Re}(g_{A_i}(r)) = 1 + \cos \pi + \cos \pi + \cos(-\pi) + \cos \pi = -3.$$

Finally, we switch our attention from g to its restriction on $\pm G_D$. Consider the Fourier template $g_D : \mathbb{Z} \rightarrow \mathbb{C}$ defined by

$$g_D(r) = \begin{cases} e((\theta_1 d_1 + \theta_2 d_2 + \theta_3 d_3 + \theta_4 d_4)t) & r = 2^{d_1} 3^{d_2} 5^{d_3} 7^{d_4} \in G_D \\ e(-(\theta_1 d_1 + \theta_2 d_2 + \theta_3 d_3 + \theta_4 d_4)t) & r = -2^{d_1} 3^{d_2} 5^{d_3} 7^{d_4} \in -G_D \\ 0 & \text{otherwise.} \end{cases}$$

Observe that if $r \notin \pm G_D$, then $(g_D)_{A_i}(r) = 0$ for all $i \in [5]$. Moreover, for those $r \in \pm G_D$ such that $r, 2r, 4r, 3r, 9r, 5r, 7r \in \pm G_D$, we have $\sum_{i=1}^5 \operatorname{Re}((g_D)_{A_i}(r)) = \sum_{i=1}^5 \operatorname{Re}(g_{A_i}(r)) = -3$. Since

$$\lim_{D \rightarrow \infty} \frac{|\{r \in \pm G_D : r, 2r, 4r, 3r, 9r, 5r, 7r \in G_D\}|}{|\pm G_D|} = 1,$$

we get that

$$\lim_{D \rightarrow \infty} \frac{\sum_{i=1}^5 \sigma_{A_i}(g_D)}{|\pm G_D|} = \lim_{D \rightarrow \infty} \frac{\sum_{r \in \pm G_D} \sum_{i=1}^5 \operatorname{Re}((g_D)_{A_i}(r))}{|\pm G_D|} = -3$$

(recall from Remark 3.4 that $\sigma_{A_i}(g_D)$ is always real). Taking D sufficiently large, we can obtain the desired Fourier template g_D with $\sum_{i=1}^5 \sigma_{A_i}(g_D) < 0$.

We discuss Case B further in Section 5.3.

Case B covers the case where exactly one of L_1, \dots, L_5 is additive quadruple, and the other four are uncommon. What if one of the other four becomes common? The following examples illustrates why the construction in Case B fails and what the alternate strategy is.

Example 5.6 (Case C). Let L be the 2×5 linear system

$$\begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 1 & -1 & 4 & 0 & -4 \end{pmatrix}.$$

We then have

$$\begin{aligned} L_5 \mathbf{x} &= x_1 + x_2 - x_3 - x_4 & A_5 &= \{\{1, 1, -1, -1\}\} \\ L_4 \mathbf{x} &= x_1 - x_2 + 4x_3 - 4x_5 & A_4 &= \{\{1, -1, 4, -4\}\} \\ L_3 \mathbf{x} &= 5x_1 + 3x_2 - 4x_4 - 4x_5 & A_3 &= \{\{3, 5, -4, -4\}\} \\ L_2 \mathbf{x} &= 2x_1 + 3x_3 - 1x_4 - 4x_5 & A_2 &= \{\{2, 3, -1, -4\}\} \\ L_1 \mathbf{x} &= 2x_2 - 5x_2 - 1x_4 + 4x_5 & A_1 &= \{\{2, -5, -1, 4\}\}. \end{aligned}$$

The difference between this example and Example 5.5 is that, in addition to the additive quadruple L_5 , we have another common linear equation L_4 .

Let us first try the strategy in Example 5.5. Since 2, 3, 5 are the prime divisors of at least one coefficient in L_1, \dots, L_5 , consider the multiplicative grid $G = \{2^{d_1} 3^{d_2} 5^{d_3} : d_1, d_2, d_3 \in \{0, 1, \dots\}\}$ and some $g : \pm G \rightarrow \mathbb{C}$ of the form

$$\begin{cases} g(2^{d_1} 3^{d_2} 5^{d_3}) = e((\theta_1 d_1 + \theta_2 d_2 + \theta_3 d_3)t) \\ g(-2^{d_1} 3^{d_2} 5^{d_3}) = e(-(\theta_1 d_1 + \theta_2 d_2 + \theta_3 d_3)t). \end{cases}$$

We wish to find some $\theta_1, \theta_2, \theta_3, t \in \mathbb{R}$ such that for all $r \in \pm G$, we have

$$\begin{aligned} 0 > \sum_{i=1}^5 \operatorname{Re}(g_{A_i}(r)) &= |g(r)|^4 + |g(r)|^2 |g(4r)|^2 + g(5r)g(3r)g(-4r)g(-4r) \\ &\quad + g(2r)g(3r)g(-r)g(-4r) + g(-2r)g(5r)g(r)g(-4r) \\ &= 2 + \cos((-4\theta_1 + \theta_2 + \theta_3)t) + \cos(-\theta_1 + \theta_2)t + \cos((-3\theta_1 + \theta_3)t). \end{aligned}$$

However, since there are no $\alpha, \beta \in \mathbb{R}$ with $\cos \alpha + \cos \beta + \cos(\alpha + \beta) < -2$, we cannot find solutions $\theta_1, \theta_2, \theta_3, t \in \mathbb{R}$ to the above inequality.

This motivates us to pursue a different route. Instead of letting $\sum_{i=1}^5 \operatorname{Re}(g_{A_i}(r))$ be constant for every $r \in \pm G$, we choose some $g : \pm G \rightarrow \mathbb{C}$ that is ‘‘periodic’’ in the multiplicative grid, so that $\sum_{i=1}^5 \operatorname{Re}(g_{A_i}(r))$ sums up to a negative number within every period.

We construct some $g : \pm G_L \rightarrow \mathbb{C}$ so that for all $d_1, d_2, d_3 \in \{0, 1, \dots\}$, we have $g(2^{d_1} 3^{d_2} 5^{d_3}) = g(2^{d_1+4} 3^{d_2+2} 5^{d_3+2})$. More precisely, consider

$$\begin{cases} g(2^{d_1} 3^{d_2} 5^{d_3}) = h(\operatorname{Mod}(d_1, 4), \operatorname{Mod}(d_2, 2), \operatorname{Mod}(d_3, 2)) \\ g(-2^{d_1} 3^{d_2} 5^{d_3}) = \overline{h(\operatorname{Mod}(d_1, 4), \operatorname{Mod}(d_2, 2), \operatorname{Mod}(d_3, 2))}, \end{cases}$$

where $h : \mathbb{Z}_4 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \rightarrow \mathbb{C}$ is defined by

$$\begin{array}{cccc} h(0, 0, 0) = 0.25 + 0.42i & h(1, 0, 0) = 0 & h(2, 0, 0) = -0.35 + 0.35i & h(3, 0, 0) = 1 - i \\ h(0, 0, 1) = 0.35 - 0.35i & h(1, 0, 1) = -1 - i & h(2, 0, 1) = 0.48 + 0.12i & h(3, 0, 1) = 0 \\ h(0, 1, 0) = -0.35 + 0.35i & h(1, 1, 0) = -1 - i & h(2, 1, 0) = 0.48 + 0.12i & h(3, 1, 0) = 0 \\ h(0, 1, 1) = -0.25 - 0.42i & h(1, 1, 1) = 0 & h(2, 1, 1) = 0.35 - 0.35i & h(3, 1, 1) = 1 - i. \end{array}$$

One can check that, for all $r \in G$, we have

$$\begin{aligned} &\sum_{j=0}^3 \sum_{k, \ell=0}^1 \left(\sum_{i=1}^5 g_{A_i}(2^j 3^k 5^\ell r) \right) \\ &= \sum_{\substack{j \in \mathbb{Z}_4 \\ k, \ell \in \mathbb{Z}_2}} (|h(j, k, \ell)|^4 + |h(j, k, \ell)|^2 |h(j+2, k, \ell)| + h(j, k, \ell+1)h(j, k+1, \ell) \overline{h(j+2, k, \ell)} \\ &\quad + h(i+1, j, k)h(i, j+1, k) \overline{h(i, j, k)h(i+2, j, k)} + \overline{h(i+1, j, k)h(i, j, k+1)h(i, j, k)h(i+2, j, k)}) \\ &= -0.249573 + 0.723675i. \end{aligned}$$

Similarly, for all $r \in -G$, we have

$$\sum_{j=0}^3 \sum_{k, \ell=0}^1 \left(\sum_{i=1}^5 g_{A_i}(2^j 3^k 5^\ell r) \right) = -0.249573 - 0.723675i.$$

As before, let $g_D : \mathbb{Z} \rightarrow \mathbb{C}$ be the restriction of g on the finite truncation $\pm G_D$, so that g_D is a Fourier template. Since

$$\lim_{D \rightarrow \infty} \frac{|\{r \in \pm G_D : \{1, 2, 4, 8\} \times \{1, 3\} \times \{1, 5\} \times r \subseteq \pm G_D\}|}{|\pm G_D|} = 1,$$

we get that

$$\lim_{D \rightarrow \infty} \frac{\sum_{i=1}^5 \sigma_{A_i}(g_D)}{|\pm G_D|/16} = -0.249573 < 0.$$

Taking D sufficiently large, we can obtain the desired Fourier template g_D with $\sum_{i=1}^5 \sigma_{A_i}(g_D) < 0$.

In general, to prove Theorem 1.10 under Case C, we will repeatedly use and generalize the above function h . At this point, it might not be intuitive why the values of $h(i, j, k)$ are chosen in this way, and how they can be generalized. We will discuss this further in Section 5.4, which hopefully will explain more to the readers.

The above three cases form the majority of the proof of Theorem 1.10. Case D does not have many possibilities, as each linear system under Case D can be parameterized by one single variable; we will prove Theorem 1.10 under Case D in Section 5.5. In Section 5.6, we will show Theorem 1.10 for 2×5 linear systems with girth $s(L) \leq 3$. Finally, in Section 5.7, we show commonness of the several linear systems indicated in Theorem 1.10.

Having illustrated some of our proof strategies, we now begin the proof of Theorem 1.10.

5.2. Proof of Theorem 1.10 under Case A. In this section, we prove Theorem 1.10 under Case A listed in Corollary 5.3.

Definition 5.7. Suppose L is an irredundant 2×5 system with $s(L) = 4$. Recall Setup 5.1.

- For $i \in [5]$ and $\lambda \in \mathbb{Q} \setminus \{0\}$, we say that L_i is λ -common if A_i is of the form $\{\{a, -a, \lambda a, -\lambda a\}\}$.
- For $a, b \in \mathbb{Q}$, we say that a, b are L -coincidental if there exist $\lambda \in \{\pm a/b, \pm b/a\}$ and $i \in [5]$ such that L_i is λ -common.

Suppose L is a linear system that falls under Case A listed in Corollary 5.3. We show that L is always uncommon. This follows from combining the following two statements:

- (1) (Lemma 5.8) If L falls under Case A in Corollary 5.3, then there exists A_i that cannot be partitioned into two L -coincidental pairs.
- (2) (Lemmas 5.9 and 5.10) If for some $i \in [5]$, A_i cannot be partitioned into two L -coincidental pairs, then L is uncommon.

Note that, if A_i cannot be partitioned into two L -coincidental pairs, then one of the following occurs:

- (a) there exists $a \in A_i$ that is not L -coincidental with any element in $A_i \setminus \{a\}$,
- (b) there exists $a \in A_i$ that is L -coincidental with every element in $A_i \setminus \{a\}$, while no two of the three elements in $A_i \setminus \{a\}$ are L -coincidental.

In particular, Lemma 5.9 shows that L is uncommon when (a) occurs, and Lemma 5.10 shows that L is uncommon when (b) occurs.

The detailed statements are as follows.

Lemma 5.8. *Suppose L is an irredundant 2×5 linear system that falls under Case A in Corollary 5.3. Assume Setup 5.1. Then there exists $i \in [5]$ such that A_i cannot be partitioned into two L -coincidental pairs.*

Lemma 5.9. *Suppose L is an irredundant 2×5 linear system that falls under Case A in Corollary 5.3. Assume Setup 5.1. If there exists $i \in [5]$ and $a \in A_i$ such that a is not L -coincidental with any element in $A_i \setminus \{a\}$, then there exists a Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.*

Lemma 5.10. *Suppose L is an irredundant 2×5 linear system that falls under Case A in Corollary 5.3. Assume Setup 5.1. Moreover, suppose:*

- (1) *for every $i \in [5]$ and every $a \in A_i$, there is some $b \in A_i \setminus \{a\}$ such that a, b are L -coincidental.*
- (2) *there exists $i_0 \in [5]$, $A_{i_0} = \{\{a_0, b_0, c_0, d_0\}\}$ such that:*
 - a_0 *is L -coincidental with each of b_0, c_0, d_0 ,*
 - b_0, c_0, d_0 *are pairwise not L -coincidental.*

Then there exists a Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.

The proof of Lemma 5.10 is effectively a generalization of Example 5.4.

Proof of Lemma 5.10. Consider the Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ defined by

$$\begin{cases} g(a_0) = g(-a_0) = 1 \\ g(b_0) = g(-b_0) = g(c_0) = g(-c_0) = g(d_0) = g(-d_0) = -C \\ g(r) = 0 \quad \text{otherwise.} \end{cases}$$

We then have $\sigma_{A_{i_0}}(g) = -2C^3$. Moreover, because of condition (1) in the lemma statement, there is no $i \in [5]$ such that A_i has all elements lying in $\{\pm b_0, \pm c_0, \pm d_0\}$. Therefore, for all $i \in [5]$, we either have $\sigma_{A_i}(g) = -2C^3$ or $|\sigma_{A_i}(g)| = O(C^2)$. This gives $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$ when $C > 0$ is sufficiently large. \square

The proof of Lemma 5.9 has more case discussions than Lemma 5.10, although the underlying principle is identical. Similarly, the proof of Lemma 5.8 is routine. As such, we defer the proofs of these lemmas to Appendix B.1.

5.3. Proof of Theorem 1.10 under Case B. In this section, we prove Theorem 1.10 under Case B of Corollary 5.3. In this case, we always assume the following setup:

Setup 5.11. Suppose

$$\begin{aligned} L_5 \mathbf{x} &= x_1 + x_2 - x_3 - x_4 & A_5 &= \{\{1, 1, -1, -1\}\} \\ L_4 \mathbf{x} &= ax_1 + bx_2 + cx_3 + dx_5 & A_4 &= \{\{a, b, c, d\}\} \\ L_3 \mathbf{x} &= (a+c)x_1 + (b+c)x_2 - cx_4 + dx_5 & A_3 &= \{\{a+c, b+c, -c, d\}\} \\ L_2 \mathbf{x} &= (a-b)x_1 + (b+c)x_3 + bx_4 + dx_5 & A_2 &= \{\{a-b, b+c, b, d\}\} \\ L_1 \mathbf{x} &= (b-a)x_3 + (a+c)x_3 + ax_4 + dx_5 & A_1 &= \{\{b-a, a+c, a, d\}\} \end{aligned}$$

with $a, b, c, d \in \mathbb{Z} \setminus \{0\}$, and none of L_1, \dots, L_4 is common. By replacing a, b, c, d by their negations if necessary, we may suppose that at least two of a, b, c, d are positive.

We aim to show that L is always uncommon. As in Example 5.5, the proof strategy is in two steps:

- (1) First, we construct an appropriate ‘‘multiplicative grid’’ G_L which is the support of the Fourier template g .
- (2) Next, we set the values of $g(r)$ for $r \in G_L$ to be some complex numbers so that the phases align in a way for $\sum_{i=1}^5 \sigma_{A_i}(g)$ to be negative. We will in fact choose g to be the finite truncation of some $h : \pm G_L \rightarrow \mathbb{C}$ where each h_{A_i} is periodic (in some suitable sense that we will define) on G_L . Most of the times h will look like the functions in Example 5.5, given by the composition of $e(\cdot)$ with a linear map from $G_L \rightarrow \mathbb{R}$.

As such, we begin by defining the ‘‘multiplicative grid’’ G_L associated to a 2×5 linear system L as follows.

Definition 5.12. Let L be a 2×5 linear system with $s(L) = 4$.

- Let $P_L = \{p_1, \dots, p_\ell\} = \{p : p \text{ is a prime divisor of some element in } A_1 \cup \dots \cup A_5\}$.
- Let $G_L = \{p_1^{d_1} \cdots p_\ell^{d_\ell} : d_1, \dots, d_\ell \in \{0, 1, \dots\}\}$.

For a function $h : \pm G_L \rightarrow \mathbb{C}$, $i \in [5]$ and $\vec{u} = (u_1, \dots, u_\ell) \in \mathbb{N}^\ell$, we say that h_{A_i} is \vec{u} -periodic if for all $r \in G_L$, we have

$$h_{A_i}(r) = h_{A_i}(p_1^{u_1} \cdots p_\ell^{u_\ell} \cdot r).$$

Remark 5.13. We note that while a more natural definition for \vec{u} -periodicity is $h(r) = h(p_1^{u_1} \cdots p_\ell^{u_\ell} \cdot r)$ – and some of our constructions satisfy this stronger periodicity condition – we do need the more general notion of periodicity in Definition 5.12 for the construction given in Example 5.5.

Now, we construct a function $h: \pm G_L \rightarrow \mathbb{C}$ such that h_{A_1}, \dots, h_{A_5} are \vec{u} -periodic on G_L and has the “phase cancellation” property on a period. That is,

$$\operatorname{Re} \sum_{0 \leq d_1 \leq u_1 - 1} \sum_{i=1}^5 h_{A_i}(p_1^{d_1} \dots p_\ell^{d_\ell} \cdot r) < 0. \quad (4)$$

$$\vdots$$

$$0 \leq d_\ell \leq u_\ell - 1$$

It turns out that the above is enough to guarantee the existence of a Fourier template $g: \mathbb{Z} \rightarrow \mathbb{C}$ such that $\operatorname{Re}(\sigma_{A_i}(g)) < 0$. We have seen some version of this idea in Example 5.5.

Lemma 5.14. *Let L be a 2×5 system with $s(L) = 4$ and suppose G_L is the associated multiplicative grid as in Definition 5.12. Conider $h: \pm G_L \rightarrow \mathbb{C}$ such that $h(-r) = \overline{h(r)}$ for all $r \in G_L$. Suppose h_{A_1}, \dots, h_{A_5} are \vec{u} -periodic for some $\vec{u} \in \mathbb{N}^\ell$, such that (4) holds. Then there exists a Fourier template $g: \mathbb{Z} \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.*

We defer the formal proof of Lemma 5.14 to Appendix B.2. However, the main idea is just the last part of Example 5.5. Roughly speaking, we “tessellate” enough copies of h on a sufficiently large truncation $\pm G_D$, where $G_D = \{p_1^{d_1} \dots p_\ell^{d_\ell} : d_1, \dots, d_\ell \in \{0, 1, \dots, D\}\} \subseteq G_L$. Then, we will define the Fourier template $g: \pm G_D \rightarrow \mathbb{C}$ by

$$g(r) = h(r) \quad \text{for every } r \in \pm G_D$$

(so the support of g is $\pm G_D \subseteq \mathbb{Z}$, which is finite). By choosing D large, we can ensure that the terms involved in the “wrap around”/boundary terms in $\sum_{i=1}^5 \sigma_{A_i}(g)$ is much smaller than the number of main terms coming from terms in (4), and so the overall sum stays negative.

As such, in what follows, we restrict our attention to constructing periodic functions on $\pm G_L$ that sum to a negative value in its period. Depending on the properties of A_1, \dots, A_5 , we construct h as a suitable product formed from four special “primitive functions”: h_1, h_2, h_3, h_4 .

Definition 5.15. Let A be a multiset of size 4. We say that A is *cancelling* if A is of the form $\{\{a_1, a_2, b_1, b_2\}\}$ such that $a_1, a_2, -b_1, -b_2 \in \mathbb{N}$, with $a_1 a_2 = b_1 b_2$.

We already know that $A_5 = \{\{1, 1, -1, -1\}\}$. For each A_i among A_1, A_2, A_3, A_4 , we know that A_i satisfies one of the following:

- (i) A_i has four positive elements,
- (ii) A_i has three positive and one negative elements,
- (iii) A_i has two positive and two negative elements, but is not cancelling,
- (iv) A_i is cancelling (but not an additive quadruple).

Observe that if A_i meets (i) or (ii), then we can pick very simple functions $h: \pm G_L \rightarrow \mathbb{C}$ to make h_{A_i} negative.

Observation 5.16. *There exist functions $h_1, h_2: \pm G_L \rightarrow \mathbb{C}$, defined by*

$$\begin{cases} h_1(r) = e(1/8) & r \in G_L \\ h_1(r) = e(-1/8) & r \in -G_L \end{cases} \quad \begin{cases} h_2(r) = i & r \in G_L \\ h_2(r) = -i & r \in -G_L \end{cases}$$

such that, for all 4-multisets A , $(h_1)_A$ and $(h_2)_A$ are both constant on G_L with the following values:

$$(h_1)_A = \begin{cases} -1 & A \text{ has four positive elements} \\ i \text{ or } -i & A \text{ has three positive and one negative elements} \\ 1 & A \text{ has two positive and two negative elements,} \end{cases}$$

$$(h_2)_A = \begin{cases} -1 & A \text{ has three positive and one negative elements} \\ 1 & A \text{ has two positive and two negative elements.} \end{cases}$$

However, if some A_i , $i \in [4]$ has an equal number of positive and negative elements, then we have $(h_1)_{A_i} = (h_2)_{A_i} = 1$, so new constructions are needed when this occurs.

We first discuss the case when all of A_1, \dots, A_4 meets (iii). The construction for this case generalizes that in Example 5.5.

Proposition 5.17. *Let L be a 2×5 system that falls under Case B in Corollary 5.3. Let G_L be the associated multiplicative grid as given by Definition 5.12. Suppose A_5 is an additive quadruple, and each of A_1, \dots, A_4 contains an equal number of positive and negative elements, but is not cancelling.*

Then there exists a function $h_3: \pm G_L \rightarrow \mathbb{C}$ with $h(-r) = \overline{h(r)}$ such that each $(h_3)_{A_i}$, $i \in [5]$ is constant on G_L . Furthermore, $\operatorname{Re} \sum_{i=1}^5 (h_3)_{A_i}(r) < 0$ for every $r \in \pm G_L$.

In the proof of Proposition 5.17, we will need the following fact, whose proof we put in Appendix B.2. Fact 5.18 is the driving force behind the idea of getting the phases of $h_{A_i}(r)$ to align and produce a negative real value.

Fact 5.18. *For all $\gamma_1, \gamma_2, \gamma_3, \gamma_4 \in \mathbb{Q} \setminus \{0\}$, there exists $t \in \mathbb{R}$ such that $1 + \cos(\gamma_1 t) + \cos(\gamma_2 t) + \cos(\gamma_3 t) + \cos(\gamma_4 t) < 0$.*

The proof of Proposition 5.17 generalizes what we saw in Example 5.5.

Proof of Proposition 5.17. Recall that

$$P_L = \{p_1, \dots, p_\ell\} = \{p : p \text{ is a prime divisor of some element in } A_1 \cup \dots \cup A_5\}.$$

For some $\theta_1, \dots, \theta_\ell \in \mathbb{Q}$ and $t \in \mathbb{R}$ to be chosen later, define $h_3 : \pm G_L \rightarrow \mathbb{C}$ by

$$\begin{cases} h_3(p_1^{d_1} \dots p_\ell^{d_\ell}) = e((\theta_1 d_1 + \dots + \theta_\ell d_\ell)t) \\ h_3(-p_1^{d_1} \dots p_\ell^{d_\ell}) = e(-(\theta_1 d_1 + \dots + \theta_\ell d_\ell)t). \end{cases}$$

Observe that, since each of A_1, \dots, A_5 contains two positive and two negative elements, $(h_3)_{A_1}, \dots, (h_3)_{A_5}$ are constant on G_L . In particular, we have $(h_3)_{A_5} = 1$. Because none of A_1, \dots, A_4 are cancelling, by some elementary linear algebra, we can choose $\theta_1, \dots, \theta_\ell \in \mathbb{Q}$ so that

$$\operatorname{Re} \sum_{i=1}^5 (h_3)_{A_i}(r) = 1 + \cos(\gamma_1 t) + \cos(\gamma_2 t) + \cos(\gamma_3 t) + \cos(\gamma_4 t) \quad \text{with } \gamma_1, \gamma_2, \gamma_3, \gamma_4 \in \mathbb{Q} \setminus \{0\}.$$

By Fact 5.18, we can choose $t \in \mathbb{R}$ so that $1 + \cos(\gamma_1 t) + \cos(\gamma_2 t) + \cos(\gamma_3 t) + \cos(\gamma_4 t) < 0$. This ensures that (4) holds. \square

It is clear that the construction in Proposition 5.17 fails when some A_i , $i \in [4]$ is cancelling, because then h_{A_i} becomes identically 1, and we can no longer leverage Fact 5.18 to do the phase cancellation. To handle this final case, we can leverage the discrepancy between the largest exponents of primes that divide various elements of A_i ; that is, h is a function of these largest exponents, for which we use the following notation.

Definition 5.19.

- (1) For prime number p and $r \in \mathbb{Z} \setminus \{0\}$, define $v_p(r) = \max\{d \in \{0, 1, \dots\} : p^d \mid |r|\}$.
- (2) For prime number p and multiset A , define $V_p(A) = \{\{v_p(a) : a \in A\}\}$.

Next, we demonstrate our construction through two examples.

Example 5.20. Let L be the 2×5 linear system

$$\begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 1 & -3 & -4 & 0 & 12 \end{pmatrix}.$$

We then have

$$\begin{aligned}
L_5\mathbf{x} &= x_1 + x_2 - x_3 - x_4 & A_5 &= \{\{1, 1, -1, -1\}\} \\
L_4\mathbf{x} &= x_1 - 3x_2 - 4x_3 + 12x_5 & A_4 &= \{\{1, -3, -4, 12\}\} \\
L_3\mathbf{x} &= -3x_1 - 7x_2 + 4x_4 + 12x_5 & A_3 &= \{\{-3, -7, 4, 12\}\} \\
L_2\mathbf{x} &= 4x_1 - 7x_3 - 3x_4 + 12x_5 & A_2 &= \{\{4, -7, -3, 12\}\} \\
L_1\mathbf{x} &= -4x_2 - 3x_3 + x_4 + 12x_5 & A_1 &= \{\{-4, -3, 1, 12\}\}.
\end{aligned}$$

Note that in this context we have that $G_L = \{2^{d_1}3^{d_2}7^{d_3} : d_1, d_2, d_3 \in \{0, 1, \dots\}\}$. By Lemma 5.14, to show that L is uncommon, it suffices to find a \vec{u} -periodic function $h: \pm G_L \rightarrow \mathbb{C}$ with $h(-r) = \overline{h(r)}$ such that (4) holds.

Note that $v_2(3) = 0, v_2(4) = 2, v_2(12) = 2$ and $v_3(3) = 1, v_3(4) = 0, v_3(12) = 1$. It turns out to be advantageous to consider the largest exponent of 2 and 3, which are the primes that “witness” the cancelling structure in A_4 . In particular, sending $A_i \mapsto \{(v_2(r), v_3(r)) : r \in A_i\}$ for each i , we have:

$$\begin{aligned}
A_5 &= \{\{1, 1, -1, -1\}\} \mapsto \{(0, 0), (0, 0), (0, 0), (0, 0)\} \\
A_4 &= \{\{1, -3, -4, 12\}\} \mapsto \{(0, 0), (0, 1), (2, 0), (2, 1)\} \\
A_3 &= \{\{-3, -7, 4, 12\}\} \mapsto \{(0, 1), (0, 0), (2, 0), (2, 1)\} \\
A_2 &= \{\{4, -7, -3, 12\}\} \mapsto \{(2, 0), (0, 0), (1, 0), (2, 1)\} \\
A_1 &= \{\{-4, -3, 1, 12\}\} \mapsto \{(2, 0), (0, 1), (0, 0), (2, 1)\}.
\end{aligned}$$

Consider the following function $h: \pm G_L \rightarrow \mathbb{C}$ given by

$$h(r) = \begin{cases} -1 & \text{if } \left\lfloor \frac{v_2(r)}{2} \right\rfloor \equiv v_3(r) \equiv 1 \pmod{2} \\ 1 & \text{otherwise.} \end{cases}$$

Then, since $h(2^4 \cdot 3^2 \cdot r) = h(r)$ for every r , we know that every h_{A_i} is \vec{u} -periodic with $\vec{u} = (4, 2, 1)$. Moreover, we have

$$\begin{aligned}
& \sum_{\substack{0 \leq d_1 \leq 3 \\ 0 \leq d_2 \leq 1 \\ r = 2^{d_1}3^{d_2}}} (|h(r)|^4 + h(r)h(-3r)h(-4r)h(12r) + h(-3r)h(-7r)h(4r)h(12r) \\
& \quad + h(4r)h(-7r)h(-3r)h(12r) + h(-4r)h(-3r)h(r)h(12r)) \\
&= 8 - 8 - 8 - 8 - 8 = -24 < 0.
\end{aligned}$$

Therefore, h satisfies the condition in Lemma 5.14.

Example 5.21. Let L be the 2×5 linear system

$$\begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 6 & 24 & -144 & 0 & -1 \end{pmatrix}.$$

We then have

$$\begin{aligned}
L_5\mathbf{x} &= x_1 + x_2 - x_3 - x_4 & A_5 &= \{\{1, 1, -1, -1\}\} \\
L_4\mathbf{x} &= 6x_1 + 24x_2 - 144x_3 - x_5 & A_4 &= \{\{6, 24, -144, -1\}\} \\
L_3\mathbf{x} &= -138x_1 - 120x_2 + 144x_4 - x_5 & A_3 &= \{\{-138, -120, 144, -1\}\} \\
L_2\mathbf{x} &= -18x_1 - 120x_3 + 24x_4 - x_5 & A_2 &= \{\{-18, -120, 24, -1\}\} \\
L_1\mathbf{x} &= 18x_2 - 138x_3 + 6x_4 - x_5 & A_1 &= \{\{18, -138, 6, -1\}\}
\end{aligned}$$

Note that in this context we have $G_L = \{2^{d_1}3^{d_2}5^{d_3}23^{d_4} : d_1, d_2, d_3, d_4 \in \mathbb{N}\}$. By Lemma 5.14, to show that L is uncommon, it suffices to find a \vec{u} -periodic function $h: \pm G_L \rightarrow \mathbb{C}$ such that (4) holds.

In this case, note that $v_2(6), v_2(24), v_2(144) \geq 1$ and $v_2(6) + v_2(24) = v_2(144)$, and so the cancelling structure in A_4 is “witnessed” by the prime 2. This stands in contrast to Example 5.20, where two primes were needed to “witness” this cancelling structure. At present it turns out to be advantageous to isolate the largest exponent of 2 that divide each of the elements of A_i . In particular, sending $A_i \mapsto \{\{v_2(r) : r \in A_i\}\}$ for each i , we have:

$$\begin{aligned} A_5 &= \{\{1, 1, -1, -1\}\} \mapsto \{\{0, 0, 0, 0\}\} \\ A_4 &= \{\{6, 24, -144, -1\}\} \mapsto \{\{1, 3, 4, 0\}\} \\ A_3 &= \{\{-138, -120, 144, -1\}\} \mapsto \{\{1, 3, 4, 0\}\} \\ A_2 &= \{\{-18, -120, 24, -1\}\} \mapsto \{\{1, 3, 3, 0\}\} \\ A_1 &= \{\{18, -138, 6, -1\}\} \mapsto \{\{1, 1, 1, 0\}\} \end{aligned}$$

We will choose an appropriate $\phi: \mathbb{N} \rightarrow \{\pm 1\}$ and show that $h(r) = \phi(v_2(r))$ has the desired properties. We will choose ϕ to be periodic, and then it will follow that h is periodic on G_L as well. Consider

$$\phi(v) = \begin{cases} -1 & \text{if } v \equiv 0, 1 \pmod{4}, \\ 1 & \text{if } v \equiv 2, 3 \pmod{4}. \end{cases}$$

Then, since $h(2^4 \cdot r) = h(r)$ for every r , we know that every h_{A_i} is \vec{u} -periodic with $\vec{u} = (4, 1, 1, 1)$. Moreover, for $r = 2^{d_1}$ with $d_1 = 0, 1, 2, 3$, the values of each $h_{A_i}(r)$ varies as follows:

- $h_{A_5} = 1$, as

$$\phi(0)^4 = \dots = \phi(3)^4 = 1.$$

- $h_{A_4}, h_{A_3} = -1$, as

$$\phi(1)\phi(3)\phi(0)^2 = \phi(2)\phi(0)\phi(1)^2 = \phi(3)\phi(1)\phi(2)^2 = \phi(0)\phi(2)\phi(3)^2 = -1.$$

- $h_{A_1} = h_{A_2} = 1$ when $d_1 \in \{0, 2\}$ and $h_{A_1} = h_{A_2} = -1$ when $d_1 \in \{1, 3\}$, as

$$\phi(1)\phi(0) = \phi(3)\phi(2) = 1, \quad \phi(2)\phi(1) = \phi(0)\phi(3) = -1.$$

Altogether, we have

$$\begin{aligned} &\sum_{\substack{0 \leq d_1 \leq 3 \\ r=2^{d_1}}} (|h(r)|^4 + h(6r)h(24r)h(-144r)h(r) + h(-138r)h(-120r)h(144r)h(-r) \\ &\quad + h(-18r)h(-120r)h(24r)h(-r) + h(18r)h(-138r)h(6r)h(-r)) \\ &= 4 - 4 - 4 + 0 + 0 = -4 < 0. \end{aligned}$$

Therefore, h satisfies the condition in Lemma 5.14.

The general result, whose proof we put in Appendix B.2, is the following.

Proposition 5.22. *Suppose L is an irredundant 2×5 linear system that falls under Case B in Corollary 5.3 and suppose that for some $i \in [4]$, A_i is a cancelling pair. Then there exists a Fourier template $g: \mathbb{Z} \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.*

The proof of Proposition 5.22 proceeds in two steps. First, in Corollary B.6 we make precise the notion of “witnessing the cancelling structure” in the above examples, by classifying the types of cancelling sets that can occur. Second, we handle each case in the series of Proposition B.10–Proposition B.13. The main ideas on how to handle each case generalize those in Example 5.20 and Example 5.21.

We summarize some of the properties of the above constructions in the following table.

TABLE 1. Periodic sums of $(h_j)_A$ on G_L

	h_1	h_2	h_3	h_4
A has four positive elements	-1			
A has three positive and one negative elements	$\pm i$	-1		
A has two positive and two negative elements but is not cancelling	1	1	$e^{i\gamma t}$	
A is cancelling but not additive quadruple	1	1	1	-1
Relevant results	5.16	5.16	5.17	5.22

Finally, we put together Observation 5.16, Proposition 5.17 and Proposition 5.22 to complete the proof of case B.

Proposition 5.23. *Suppose L is an irredundant 2×5 linear system that falls under Case B in Corollary 5.3. Then there exists a Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.*

Roughly speaking, depending on the structures of A_1, \dots, A_5 that appear in the linear system L – in particular, which rows in Table 1 these A_i belong to – we choose some subset of h_j as described in Table 1 such that the product of each row restricted to chosen h_j sums up to negative. Then we take the Fourier template g_j to be the finite truncation of h_j , and finally take g to be the joined Fourier template from the g_j .

In the proof of Proposition 5.23, we will need the following analogue of Fact 5.18 (also proved in Appendix B.2).

Fact 5.24. *For all $\gamma_1, \gamma_2 \in \mathbb{Q} \setminus \{0\}$, there exists $t \in \mathbb{R}$ such that $1 + \cos(\gamma_1 t) + \cos(\gamma_2 t) < 0$.*

Proof of Proposition 5.23. If at least one of A_1, \dots, A_4 is cancelling then we can conclude by Proposition 5.22. It remains to handle the case when none of A_1, A_2, A_3, A_4 is cancelling.

- (1) Suppose two of A_1, A_2, A_3, A_4 contains four positive elements and the other two contain three positive and one negative elements. Then we can apply Lemma 5.14 and Observation 5.16 with $h_1 : \pm G_L \rightarrow \mathbb{C}$.
- (2) Suppose all four of A_1, A_2, A_3, A_4 contain three positive and one negative element. Then we can apply Lemma 5.14 and Observation 5.16 with $h_2 : \pm G_L \rightarrow \mathbb{C}$.
- (3) Suppose each of A_1, \dots, A_4 contain two positive and two negative elements. Then we can apply Lemma 5.14 and Proposition 5.17 with $h_3 : \pm G_L \rightarrow \mathbb{C}$.
- (4) Suppose that two of A_1, \dots, A_4 (say A_{i_1}, A_{i_2}) contain three positive and one negative elements, and the other two contain two positive and two negative elements (say A_{i_3}, A_{i_4}). We construct two functions h_3 and $h_3 h_2$, and it will be clear in a moment why we do so.

We set h_3 to be the function that we constructed in Proposition 5.17, but with θ_i and t chosen according to Fact 5.24. More precisely, recall that

$$P_L = \{p_1, \dots, p_\ell\} = \{p : p \text{ is a prime divisor of some element in } A_1 \cup \dots \cup A_5\}.$$

For some $\theta_1, \dots, \theta_\ell \in \mathbb{Q}$ and $t \in \mathbb{R}$ to be chosen later, define $h_3 : \pm G_L \rightarrow \mathbb{C}$ by

$$\begin{cases} h_3(p_1^{d_1} \cdots p_\ell^{d_\ell}) = e((\theta_1 d_1 + \cdots + \theta_\ell d_\ell)t) \\ h_3(-p_1^{d_1} \cdots p_\ell^{d_\ell}) = e(-(\theta_1 d_1 + \cdots + \theta_\ell d_\ell)t) \end{cases}$$

Observe that $(h_3)_{A_{i_1}}, (h_3)_{A_{i_2}}, (h_3)_{A_5}$ are constant on G_L , and $(h_3)_{A_5} = 1$. Since A_{i_1} and A_{i_2} are not cancelling, we are able to choose $\theta_1, \dots, \theta_\ell \in \mathbb{Q}$ so that

$$(h_3)_{A_5} + \operatorname{Re}((h_3)_{A_{i_1}}) + \operatorname{Re}((h_3)_{A_{i_2}}) = 1 + \cos(\gamma_1 t) + \cos(\gamma_2 t), \quad \gamma_1, \gamma_2 \in \mathbb{Q} \setminus \{0\}.$$

Using Fact 5.24, we can choose $t \in \mathbb{R}$ so that $1 + \cos(\gamma_1 t) + \cos(\gamma_2 t) < 0$.

Now consider $h_3 h_2$. Since $(h_2)_{A_{i_1}}, (h_2)_{A_{i_2}}, (h_2)_{A_{i_5}} = 1$ and $(h_2)_{A_{i_3}} = (h_2)_{A_{i_4}} = -1$ (as summarized in Table 1), we have either

$$\operatorname{Re} \sum_{i=1}^5 h_3(r) < 0 \quad \text{or} \quad \operatorname{Re} \sum_{i=1}^5 h_3(r) h_2(r) < 0$$

for every $r \in \pm G_L$. Applying Lemma 5.14 gives the result. \square

5.4. Proof of Theorem 1.10 under Case C. In this section, we prove Theorem 1.10 under Case C listed in Corollary 5.3.

Suppose L falls under Case C in Corollary 5.3. We show that, if L is not isomorphic to

$$\begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 1 & -1 & 3 & 0 & -3 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 2 & -2 & 3 & 0 & -3 \end{pmatrix}$$

then L is common if it is isomorphic to

$$\begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 2 & -2 & 1 & 0 & -1 \end{pmatrix}$$

and uncommon otherwise.

Suppose L is an irredundant 2×5 linear system under Case C listed in Corollary 5.3. In this case, we always assume the following setup:

Setup 5.25. Suppose

$$\begin{aligned} L_5 \mathbf{x} &= x_1 + x_2 - x_3 - x_4 & A_5 &= \{1, 1, -1, -1\} \\ L_4 \mathbf{x} &= bx_1 - bx_2 + ax_3 - ax_5 & A_4 &= \left\{1, -1, \frac{a}{b}, -\frac{a}{b}\right\} \\ L_3 \mathbf{x} &= (a+b)x_1 + (a-b)x_2 - ax_4 - ax_5 & A_3 &= \left\{\frac{a+b}{b}, \frac{a-b}{b}, -\frac{a}{b}, -\frac{a}{b}\right\} \\ L_2 \mathbf{x} &= 2bx_1 + (a-b)x_3 - bx_4 - ax_5 & A_2 &= \left\{2, \frac{a-b}{b}, -1, -\frac{a}{b}\right\} \\ L_1 \mathbf{x} &= 2bx_2 - (a+b)x_3 - bx_4 + ax_5 & A_1 &= \left\{2, -\frac{a+b}{b}, -1, \frac{a}{b}\right\} \end{aligned}$$

with $a, b \in \mathbb{N}$ coprime, and none of L_1, L_2, L_3 is common.

In order to build some intuition for this case, we revisit Example 5.6 and reframe the construction in a manner which generalizes easily.

Example 5.26 (Take two of Example 5.6). We recall the setup. Let L be the linear system

$$\begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 1 & -1 & 4 & 0 & -4 \end{pmatrix}.$$

We then have

$$\begin{aligned} L_5 \mathbf{x} &= x_1 + x_2 - x_3 - x_4 & A_5 &= \{1, 1, -1, -1\} \\ L_4 \mathbf{x} &= x_1 - x_2 + 4x_3 - 4x_5 & A_4 &= \{1, -1, 4, -4\} \\ L_3 \mathbf{x} &= 5x_1 + 3x_2 - 4x_4 - 4x_5 & A_3 &= \{3, 5, -4, -4\} \\ L_2 \mathbf{x} &= 2x_1 + 3x_3 - 1x_4 - 4x_5 & A_2 &= \{2, 3, -1, -4\} \\ L_1 \mathbf{x} &= 2x_2 - 5x_3 - 1x_4 + 4x_5 & A_1 &= \{2, -5, -1, 4\}. \end{aligned}$$

We have the corresponding multiplicative grid

$$G_L = \{2^{d_1} 3^{d_2} 5^{d_3} : d_1, d_2, d_3 \in \{0, 1, \dots\}\},$$

and the goal is to construct a \vec{u} -periodic $h: \pm G_L \rightarrow \mathbb{C}$ with $h(-r) = \overline{h(r)}$ and the property (4):

$$\operatorname{Re} \sum_{\substack{0 \leq d_1 \leq u_1 - 1 \\ 0 \leq d_2 \leq u_2 - 1 \\ 0 \leq d_3 \leq u_3 - 1}} \sum_{i=1}^5 h_{A_i}(2^{d_1} 3^{d_2} 5^{d_3} \cdot r) < 0.$$




FIGURE 1. An illustration of the “orbits” of h . The pink/dotted arrow represents the action of multiplying by 4, the green/solid arrow represents the action of multiply by 3 and the blue/dashed arrow represents the action of multiplying by 5.

Figure 1 illustrates our construction. In particular, we chose $\vec{u} = (4, 2, 2)$, which means that $h(2^4 \cdot 3^2 \cdot 5^2 \cdot r) = h(r)$ for every $r \in G_L$. The left column denotes the value of $h(r)$ when $r = 2^i 3^j 5^k$ with $i \in \{0, 2\}$ and $j, k \in \mathbb{Z}_2$; right left column denotes the value of $h(r)$ when $r = 2^i 3^j 5^k$ with $i \in \{1, 3\}$ and $j, k \in \mathbb{Z}_2$. And as always, we take $h(-r) = \overline{h(r)}$ for every $r \in G_L$.

The key observation is that, since $h_{A_1}(r), h_{A_2}(r), h_{A_4}(r)$ are nonvanishing only if $h(r), h(4r)$ are both nonzero, we have $h_{A_1}(r) = h_{A_2}(r) = h_{A_4}(r) = 0$ if r is any of the yellow/shaded elements. This gives

$$\begin{aligned} \sum_{r \text{ yellow}} \sum_{i=1}^5 h_{A_i}(r) &= \sum_{r \text{ yellow}} (h_{A_5}(r) + h_{A_3}(r)) \\ &= 2|1 - i|^4 + 2|-1 - i|^4 + 2(-1 - i)^2(1 + i)^2 + 2(1 - i)^2(-1 + i)^2 = 0. \end{aligned}$$

We draw attention to the fact that multiplication by 3 and 5 create orbits of length two. This fact, coupled with the choice of values of h on the yellow/shaded elements, is essential in ensuring that the contributions from h_{A_3} cancels out the contributions of h_{A_5} on the yellow/shaded elements. In effect the only contribution to the sum $\sum_{\substack{0 \leq d_1 \leq 3 \\ 0 \leq d_2 \leq 1 \\ 0 \leq d_3 \leq 1}} \sum_{i=1}^5 h_{A_i}(2^{d_1} 3^{d_2} 5^{d_3})$ comes from $\sum_{i=1}^5 h_{A_i}(r)$

when r are the white elements on the left. This final part is a computation:

$$\operatorname{Re} \sum_{r \text{ white}} \sum_{i=1}^5 h_{A_i}(r) = 0.474101 + 0.474026 - 0.2401 - 0.4788 - 0.4788 = -0.249573.$$

To summarize, we can think of our construction as a combination of two components:

- A function $f: \mathbb{Z}_4 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \rightarrow \mathbb{C}$ represented by the table of values in Figure 1.
- A function $\phi: G_L \rightarrow \mathbb{Z}_4 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ with the “orbit structure” given by:
 - $\phi(4r) = \phi(r) + (2, 0, 0)$,
 - $\phi(3r) = \phi(r) + (0, 1, 0)$,
 - $\phi(5r) = \phi(r) + (0, 0, 1)$,
 - $\phi(2r) = \phi(r) + (1, 0, 0)$ when $\phi(r)_1 \in \{0, 2\}$.

The Fourier template in Example 5.6 can therefore be thought of as $h = f \circ \phi$. The combination of these two elements ensure that the yellow/shaded cells cancel in contribution to give a net zero contribution to the sum

$$\operatorname{Re} \sum_{0 \leq d_1 \leq u_1 - 1} \sum_{i=1}^5 h_{A_i}(p_1^{d_1} \dots p_\ell^{d_\ell} \cdot r),$$

$$\vdots$$

$$0 \leq d_\ell \leq u_\ell - 1$$

while the white cells contribute a negative value. Using such a framework, we can generalize our construction to handle other systems L in Case C. In the first step, we show how to extend Figure 1 to a larger table, which gives us more flexibility with constructing the function ϕ in the second step. This is the content of the following Proposition 5.27.

Proposition 5.27. *Suppose L is an irredundant 2×5 linear system that falls under Case C in Corollary 5.3. Assume Setup 5.25. If there exists $m \geq 2$ and $\phi: G_L \rightarrow \mathbb{Z}_{2m} \times \mathbb{Z}_2 \times \mathbb{Z}_2$ such that*

$$\begin{aligned} \phi\left(\frac{a}{b} \cdot r\right) &= \phi(r) + (m, 0, 0) \\ \phi\left(\frac{|a-b|}{b} \cdot r\right) &= \phi(r) + (0, 1, 0) \\ \phi\left(\frac{a+b}{b} \cdot r\right) &= \phi(r) + (0, 0, 1) \\ \phi(2r) &= \phi(r) + (1, 0, 0) \text{ when } \phi(r)_1 \in \{0, m\}, \end{aligned}$$

Then there exists a Fourier template $g: \mathbb{Z} \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.

Vaguely speaking, we can think of Proposition 5.27 as “extending” the table in Table 1 by duplicating more columns of yellow/shaded cells in Figure 1 and leaving the white cells intact.

Proof. Suppose such ϕ exists. We first consider the case $a - b > 0$. In this case, define $h = \phi \circ f: G_L \rightarrow \mathbb{C}$, where $f: \mathbb{Z}_{2m} \times \mathbb{Z}_2 \times \mathbb{Z}_2 \rightarrow \mathbb{C}$ is given by

$$\begin{array}{ll} (0, 0, 0) \mapsto 0.25 + 0.42i & (1, 0, 0), \dots, (m-1, 0, 0) \mapsto 0 \\ (0, 0, 1) \mapsto 0.35 - 0.35i & (1, 0, 1), \dots, (m-1, 0, 1) \mapsto -1 - i \\ (0, 1, 0) \mapsto -0.35 + 0.35i & (1, 1, 0), \dots, (m-1, 1, 0) \mapsto -1 - i \\ (0, 1, 1) \mapsto -0.25 - 0.42i & (1, 1, 1), \dots, (m-1, 1, 1) \mapsto 0 \\ (m, 0, 0) \mapsto -0.35 + 0.35i & (m+1, 0, 0), \dots, (2m-1, 0, 0) \mapsto 1 - i \\ (m, 0, 1) \mapsto 0.48 + 0.12i & (m+1, 0, 1), \dots, (2m-1, 0, 1) \mapsto 0 \\ (m, 1, 0) \mapsto 0.48 + 0.12i & (m+1, 1, 0), \dots, (2m-1, 1, 0) \mapsto 0 \\ (m, 1, 1) \mapsto 0.35 - 0.35i & (m+1, 1, 1), \dots, (2m-1, 1, 1) \mapsto 1 - i. \end{array}$$

Take $\vec{u} = (u_1, \dots, u_\ell) \in \mathbb{N}^\ell$ such that $\phi(p_1^{u_1} \dots p_\ell^{u_\ell} r) = \phi(r)$. Then h_{A_1}, \dots, h_{A_5} are \vec{u} -periodic. Moreover, we have

$$\begin{aligned} \sum_{\substack{0 \leq d_1 \leq u_1 - 1 \\ \dots \\ 0 \leq d_\ell \leq u_\ell - 1}} \sum_{i=1}^5 h_{A_i}(p_1^{d_1} \dots p_\ell^{d_\ell} \cdot r) = & \sum_{i,j,k \in \mathbb{Z}_{2m} \times \mathbb{Z}_2 \times \mathbb{Z}_2} |f(i, j, k)|^4 + |f(i, j, k)|^2 |f(i+m, j, k)|^2 \\ & + f(i, j, k+1) f(i, j+1, k) \overline{f(i+m, j, k)}^2 \\ & + f(i+1, j, k) f(i, j+1, k) \overline{f(i, j, k) f(i+m, j, k)} \\ & + f(i+1, j, k) \overline{f(i, j, k+1) f(i, j, k)} f(i+m, j, k). \end{aligned}$$

A quick observation is that, if $i \notin \{0, m\}$, then one of $f(i, j, k)$ and $f(i+m, j, k)$ equals 0, so the second, fourth and fifth terms above vanish. Moreover, the first and third terms sum up to

$$2|1-i|^4 + 2|-1-i|^4 + 2(-1-i)^2(1+i)^2 + 2(1-i)^2(-1+i)^2 = 0.$$

Therefore, the above sum reduces to

$$\begin{aligned} \sum_{i,j,k \in \{0,m\} \times \mathbb{Z}_2 \times \mathbb{Z}_2} & |f(i, j, k)|^4 + |f(i, j, k)|^2 |f(i+m, j, k)|^2 + f(i, j, k+1) f(i, j+1, k) \overline{f(i+m, j, k)}^2 \\ & + f(i+1, j, k) f(i, j+1, k) \overline{f(i, j, k) f(i+m, j, k)} \\ & + f(i+1, j, k) \overline{f(i, j, k+1) f(i, j, k)} f(i+m, j, k) \end{aligned}$$

which equals $-0.249573 + 0.723675i$. Hence h has property (4). By Lemma 5.14, this gives Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.

Similarly, when $a-b < 0$ (so $h(\frac{a-b}{b} \cdot r) = \overline{h(\frac{|a-b|}{b} \cdot r)}$ for $r \in G_L$), we define $h = \phi \circ f' : G_L \rightarrow \mathbb{C}$ where $f' : \mathbb{Z}_{2m} \times \mathbb{Z}_2 \times \mathbb{Z}_2 \rightarrow \mathbb{C}$ is given by

$$f'(i, j, k) = \begin{cases} f(i, j, k) & j = 0 \\ \overline{f(i, j, k)} & j = 1. \end{cases}$$

A similar argument shows that h_{A_1}, \dots, h_{A_5} are \vec{u} -periodic, with property (4). By Lemma 5.14, this gives Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$. \square

In Lemma 5.29, we show how to construct the corresponding function $\phi : \mathbb{Z}_{2m} \times \mathbb{Z}_2 \times \mathbb{Z}_2$ for all but five specific linear systems under Case C in Corollary 5.3. Combining this with Proposition 5.27 proves that all systems that fall under Case C, barring these five exceptions, are uncommon.

To do so, we first extend the function $v_p(\cdot) : \mathbb{Z} \setminus \{0\} \rightarrow \mathbb{Z}$ introduced in Definition 5.19 to one whose domain is the set of all nonzero rational numbers.

Definition 5.28. For every prime number p , define the function $v_p(\cdot) : \mathbb{Q} \setminus \{0\} \rightarrow \mathbb{Z}$ as follows:

- (1) for all $r \in \mathbb{Z} \setminus \{0\}$, define $v_p(r) = \max\{d \in \{0, 1, \dots\} : p^d \mid |r|\}$;
- (2) for all $r \in \mathbb{Q} \setminus \mathbb{Z}$, define $v_p(r) = v_p(x) - v_p(y)$, where $r = x/y$ and $x, y \in \mathbb{Z} \setminus \{0\}$.

Lemma 5.29. *Suppose L is a linear system that falls under Case C in Corollary 5.3. Then, unless L is isomorphic to one of the following five linear systems*

$$\begin{aligned} & \begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 1 & -1 & 3 & 0 & -3 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 2 & -2 & 3 & 0 & -3 \end{pmatrix} \\ & \begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 2 & -2 & 1 & 0 & -1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 3 & -3 & 1 & 0 & -1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 3 & -3 & 2 & 0 & -2 \end{pmatrix}, \end{aligned}$$

there exists $m \geq 2$ and $\phi : G_L \rightarrow \mathbb{Z}_{2m} \times \mathbb{Z}_2 \times \mathbb{Z}_2$ that meets the condition in Proposition 5.27.

Proof. We work under Setup 5.25. One can check that, if L is not isomorphic to one of the above five systems, then one of the following occurs:

- There exist odd primes p_1, p_2, p_3 such that

$$\begin{aligned} v_{p_1}(a/b) &\neq 0, & v_{p_1}(|a-b|/b) &= v_{p_1}((a+b)/b) = 0, \\ v_{p_2}(|a-b|/b) &\neq 0, & v_{p_2}(a/b) &= v_{p_2}((a+b)/b) = 0, \\ v_{p_3}((a+b)/b) &\neq 0, & v_{p_3}(a/b) &= v_{p_3}(|a-b|/b) = 0. \end{aligned}$$

In this case, we define $\phi : G_L \rightarrow \mathbb{Z}_4 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ by

$$\phi(r) = (2 \left\lfloor \frac{v_{p_1}(r)}{v_{p_1}(a/b)} \right\rfloor + v_2(r), \left\lfloor \frac{v_{p_1}(r)}{v_{p_2}(|a-b|/b)} \right\rfloor, \left\lfloor \frac{v_{p_3}(r)}{v_{p_3}((a+b)/b)} \right\rfloor),$$

so that ϕ meets the condition in Proposition 5.27.

E.g., when $a = 10$ and $b = 1$, we define

$$\phi(r) = (2v_5(r) + v_2(r), \lfloor v_3(r)/2 \rfloor, v_{11}(r)).$$

- There exist odd primes p_2, p_3 such that

$$\begin{aligned} v_{p_2}(|a-b|/b) &\neq 0, & v_{p_2}(a/b) &= v_{p_2}((a+b)/b) = 0, \\ v_{p_3}((a+b)/b) &\neq 0, & v_{p_3}(a/b) &= v_{p_3}(|a-b|/b) = 0. \end{aligned}$$

Moreover, we have $m := |v_2(a/b)| \geq 2$.

In this case, we define $\phi : G_L \rightarrow \mathbb{Z}_{2m} \times \mathbb{Z}_2 \times \mathbb{Z}_2$ by

$$\phi(r) = (v_2(r), \left\lfloor \frac{v_{p_1}(r)}{v_{p_2}(|a-b|/b)} \right\rfloor, \left\lfloor \frac{v_{p_3}(r)}{v_{p_3}((a+b)/b)} \right\rfloor),$$

so that ϕ meets the condition in Proposition 5.27.

E.g., in Example 5.6, when $a = 4$ and $b = 1$, we define

$$\phi(r) = (v_2(r), v_3(r), v_5(r)).$$

- There exist odd primes p_1, p_2 such that

$$\begin{aligned} v_{p_1}(a/b) &\neq 0, & v_{p_1}(|a-b|/b) &= v_{p_1}((a+b)/b) = 0, \\ v_{p_2}(|a-b|/b) &\neq 0, & v_{p_2}(a/b) &= v_{p_2}((a+b)/b) = 0. \end{aligned}$$

Moreover, we have $m := |v_2((a+b)/b)| \geq 2$.

In this case, we define $\phi : G_L \rightarrow \mathbb{Z}_{2m} \times \mathbb{Z}_2 \times \mathbb{Z}_2$ by

$$\phi(r) = (mv_{p_1}(r) + \text{Mod}(v_2(r), m), \left\lfloor \frac{v_{p_2}(r)}{v_{p_2}(|a-b|/b)} \right\rfloor, \left\lfloor \frac{v_2(r)}{m} \right\rfloor),$$

so that ϕ meets the condition in Proposition 5.27.

E.g., when $a = 7$ and $b = 1$, we define

$$\phi(r) = (3v_7(r) + \text{Mod}(v_2(r), 3), v_3(r), \lfloor v_2(r)/3 \rfloor).$$

- There exist odd primes p_1, p_3 such that

$$\begin{aligned} v_{p_1}(a/b) &\neq 0, & v_{p_1}(|a-b|/b) &= v_{p_1}((a+b)/b) = 0, \\ v_{p_3}((a+b)/b) &\neq 0, & v_{p_1}(a/b) &= v_{p_2}(|a-b|/b) = 0. \end{aligned}$$

Moreover, we have $m := |v_2(|a-b|/b)| \geq 2$.

In this case, we define $\phi : G_L \rightarrow \mathbb{Z}_{2m} \times \mathbb{Z}_2 \times \mathbb{Z}_2$ by

$$\phi(r) = (mv_{p_1}(r) + \text{Mod}(v_2(r), m), \left\lfloor \frac{v_2(r)}{m} \right\rfloor, \left\lfloor \frac{v_{p_3}(r)}{v_{p_3}((a+b)/b)} \right\rfloor),$$

so that ϕ meets the condition in Proposition 5.27.

E.g., when $a = 5$ and $b = 1$, we define

$$\phi(r) = (2v_5(r) + \text{Mod}(v_2(r), 2), \lfloor v_2(r)/2 \rfloor, v_3(r)).$$

- $|a - b| = 2$ and $a + b = 2^m$, with $m \geq 3$. Moreover, there exist odd primes p_1, p_2 such that

$$\begin{aligned} v_{p_1}(a) &\neq 0, & v_{p_1}(b) &= 0, \\ v_{p_2}(b) &\neq 0, & v_{p_2}(a) &= 0. \end{aligned}$$

In this case, we define $\psi : G_L \rightarrow \mathbb{Z}_{2m-2} \times \mathbb{Z}_2 \times \mathbb{Z}_2$ by

$$\psi(r) = (u(r)(m-1) + \text{Mod}(w(r) + u(r), m-1), \left\lfloor \frac{v_{p_2}(r)}{v_{p_2}(b)} \right\rfloor - u(r), \left\lfloor \frac{w(r) + u(r)}{m-1} \right\rfloor)$$

where

$$u(r) = \left\lfloor \frac{v_{p_1}(r)}{v_{p_1}(a)} \right\rfloor, \quad w(r) = v_2(r) - \left\lfloor \frac{v_{p_2}(r)}{v_{p_2}(b)} \right\rfloor.$$

Then, we take $\phi(r) = (\psi(r)_1, \psi(r)_2 - \psi(r)_3, \psi(r)_3)$.

E.g., when $a = 5$ and $b = 3$, we define

$$\psi(r) = (2v_5(r) + \text{Mod}(v_2(r) + v_5(r) - v_3(r), 2), v_3(r) - v_5(r), \left\lfloor \frac{v_2(r) + v_5(r) - v_3(r)}{2} \right\rfloor).$$

□

Finally, for the five specific linear systems not covered by Lemma 5.29, we will show in Section 5.7 that

$$\begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 2 & -2 & 1 & 0 & -1 \end{pmatrix}$$

is common, and show in the following lemma that two of the other four are uncommon (via simple periodic constructions). This finishes the proof of Theorem 1.10 under Case C.

Lemma 5.30. *The two 2×5 linear systems*

$$\begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 3 & -3 & 1 & 0 & -1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 3 & -3 & 2 & 0 & -2 \end{pmatrix}$$

are uncommon over \mathbb{F}_p for sufficiently large p .

Proof. The first linear system has

$$\begin{aligned} L_5 \mathbf{x} &= x_1 + x_2 - x_3 - x_4 & A_5 &= \{1, 1, -1, -1\} \\ L_4 \mathbf{x} &= 3x_1 - 3x_2 + x_3 - x_5 & A_4 &= \{3, -3, 1, -1\} \\ L_3 \mathbf{x} &= 4x_1 - 2x_2 - x_4 - x_5 & A_3 &= \{4, -2, -1, -1\} \\ L_2 \mathbf{x} &= 6x_1 - 2x_3 - 3x_4 - 1x_5 & A_2 &= \{6, -2, -3, -1\} \\ L_1 \mathbf{x} &= 6x_2 - 4x_3 - 3x_4 + 1x_5 & A_1 &= \{6, -4, -3, 1\} \end{aligned}$$

which gives $P_L = \{2, 3\}$ and $G_L = \{2^{d_1} 3^{d_2} : d_1, d_2 \in \{0, 1, \dots\}\}$. Define $h : \pm G_L \rightarrow \mathbb{C}$ by

$$\begin{cases} h(r) = \phi(v_2(r), v_3(r)) & r \in G_L \\ h(r) = \overline{\phi(v_2(r), v_3(r))} & r \in -G_L \end{cases}$$

where $\phi : \mathbb{Z}_2 \times \mathbb{Z}_4 \rightarrow \mathbb{C}$ is given by

$$\begin{aligned} \phi(0, 0) &= 1 & \phi(0, 1) &= 0 & \phi(0, 2) &= 0.25i & \phi(0, 3) &= 0.5i \\ \phi(1, 0) &= -1 & \phi(1, 1) &= 0 & \phi(1, 2) &= -0.25i & \phi(1, 3) &= -0.5i. \end{aligned}$$

One can then check that

$$\operatorname{Re} \sum_{\substack{0 \leq d_1 \leq 1 \\ 0 \leq d_2 \leq 3}} \sum_{i=1}^5 h_{A_i}(2^{d_1} 3^{d_2}) = -0.265625 < 0.$$

The second linear system has

$$\begin{aligned} L_5 \mathbf{x} &= x_1 + x_2 - x_3 - x_4 & A_5 &= \{1, 1, -1, -1\} \\ L_4 \mathbf{x} &= 3x_1 - 3x_2 + 2x_3 - 2x_5 & A_4 &= \{3, -3, 2, -2\} \\ L_3 \mathbf{x} &= 5x_1 - x_2 - 2x_4 - 2x_5 & A_3 &= \{5, -1, -2, -2\} \\ L_2 \mathbf{x} &= 6x_1 - x_3 - 3x_4 - 2x_5 & A_2 &= \{6, -1, -3, -2\} \\ L_1 \mathbf{x} &= 6x_2 - 5x_3 - 3x_4 + 2x_5 & A_1 &= \{6, -5, -3, 2\} \end{aligned}$$

which gives $P_L = \{2, 3, 5\}$ and $G_L = \{2^{d_1} 3^{d_2} 5^{d_3} : d_1, d_2, d_3 \in \{0, 1, \dots\}\}$. Define $h : \pm G_L \rightarrow \mathbb{C}$ by

$$\begin{cases} h(r) = \phi(v_5(r), v_3(r)) & r \in G_L \\ h(r) = \overline{\phi(v_5(r), v_3(r))} & r \in -G_L \end{cases}$$

where $\phi : \mathbb{Z}_2 \times \mathbb{Z}_4 \rightarrow \mathbb{C}$ is given by

$$\begin{aligned} \phi(0, 0) &= 1 & \phi(0, 1) &= 0 & \phi(0, 2) &= 0.25i & \phi(0, 3) &= 0.5i \\ \phi(1, 0) &= -1 & \phi(1, 1) &= 0 & \phi(1, 2) &= -0.25i & \phi(1, 3) &= -0.5i. \end{aligned}$$

One can then check that

$$\operatorname{Re} \sum_{\substack{0 \leq d_1 \leq 1 \\ 0 \leq d_2 \leq 3}} \sum_{i=1}^5 h_{A_i}(2^{d_1} 3^{d_2}) = -0.265625 < 0.$$

The result follows by Lemma 5.14. \square

5.5. Proof of Theorem 1.10 under Case D. In this section, we prove Theorem 1.10 under Case D listed in Corollary 5.3. We show that L is common if and only if it is isomorphic to

$$\begin{pmatrix} 1 & -1 & 1 & -1 & 0 \\ 1 & 2 & -1 & 0 & -2 \end{pmatrix} \text{ or } \begin{pmatrix} 1 & -1 & 1 & -1 & 0 \\ 0 & 1 & 2 & -1 & -2 \end{pmatrix}.$$

Without loss of generality, suppose

$$\begin{aligned} L_5 &= x_1 - x_2 + \alpha x_3 - \alpha x_4 \\ L_4 &= x_1 + \beta x_2 - x_3 - \beta x_5 \\ L_3 &= (\alpha + 1)x_1 + (\alpha\beta - 1)x_2 - \alpha x_4 - \alpha\beta x_5 \\ L_2 &= (\beta + 1)x_1 + (\alpha\beta - 1)x_3 - \alpha\beta x_4 - \beta x_5 \\ L_1 &= (\beta + 1)x_2 + (-1 - \alpha)x_3 + \alpha x_4 - \beta x_5 \end{aligned}$$

with $\alpha, \beta \in \mathbb{Q}$ and $\alpha, \beta > 0$, and at least one of L_1, L_2, L_3 is common. Some elementary calculations (which we defer to Appendix B.3) show that we can reduce to the following system:

$$\begin{aligned} L_5 &= x_1 - x_2 + \alpha x_3 - \alpha x_4 \\ L_4 &= x_1 + \alpha x_2 - x_3 - \alpha x_5 \\ L_3 &= (\alpha + 1)x_1 + (\alpha^2 - 1)x_2 - \alpha x_4 - \alpha^2 x_5 \\ L_2 &= (\alpha + 1)x_1 + (\alpha^2 - 1)x_3 - \alpha^2 x_4 - \alpha x_5 \\ L_1 &= (\alpha + 1)x_2 + (-1 - \alpha)x_3 + \alpha x_4 - \alpha x_5 \end{aligned} \tag{5}$$

In this case, first observe that L_2 and L_3 are uncommon, as none of $\alpha = \alpha + 1$ or $\alpha = \alpha^2 - 1$ has rational solutions. Next, observe that $\alpha \notin \{-1, 1\}$, as either $\alpha = 1$ or $\alpha = -1$ would give $s(L) < 4$. This implies that L_4 and L_5 cannot be additive quadruples.

Suppose L_1 is an additive quadruple, i.e., $\alpha = -1/2$. In this case, L is isomorphic to

$$\begin{pmatrix} 1 & -1 & 1 & -1 & 0 \\ 1 & 2 & -1 & 0 & -2 \end{pmatrix}$$

which is common (see Section 5.7 for proof of its commonness).

Suppose none of L_1, L_4, L_5 is an additive quadruple. Then the main idea is to utilize the constructions in Section 5.2, where we set the value of the Fourier template to be large on some well-chosen values. Recall that Lemma 5.9 and Lemma 5.10 give a criterion for when such a construction works. Applying this criterion, it follows that L is common only if we can partition $\{\alpha + 1, \alpha^2 - 1, -\alpha, -\alpha^2\}$ into two pairs, such that the ratio between the two numbers in each pair lies in $\{\alpha, -\alpha, \frac{\alpha}{1+\alpha}, -\frac{\alpha}{1+\alpha}\}$. By a finite case check, the only $\alpha \in \mathbb{Q} \setminus \{-1, 0, 1\}$ with this property is $\alpha = 1/2$ ¹. When $\alpha = 1/2$, L is isomorphic to

$$\begin{pmatrix} 1 & 0 & -1 & 2 & -2 \\ 0 & 1 & 2 & -1 & -2 \end{pmatrix}$$

which is common (see Section 5.7 for proof of its commonness).

5.6. Proof of Theorem 1.10 for $s(L) < 4$. In this section, we prove Theorem 1.10 for $s(L) < 4$.

Suppose L is an irredundant 2×5 linear system with $s(L) < 4$. We show that L is common over all \mathbb{F}_p if and only if it is isomorphic to a linear system of the form

$$\begin{pmatrix} a & b & 0 & 0 & c \\ 0 & 0 & a & b & c \end{pmatrix} \tag{6}$$

$$\text{or } \begin{pmatrix} 1 & -1 & \lambda & -\lambda & 0 \\ a & b & 0 & 0 & -a-b \end{pmatrix}, \quad \{|a/b|, |a/(a+b)|, |b/(a+b)|\} \cap \{|\lambda|, |\lambda|^{-1}\} \neq \emptyset. \tag{7}$$

First, we recall the following lemma from [17].

Lemma 5.31 ([17, Theorem 1.3]). *Let L be an irredundant 2×5 linear system with $s(L)$ even. If every $1 \times s(L)$ subsystem of L (i.e., every length- $s(L)$ equation that is a subsystem of L) is uncommon over \mathbb{F}_p , then L is uncommon over \mathbb{F}_p .*

Applying Lemma 5.31, we first handle a few simple cases.

If $s(L) = 2$, then since L is irredundant (i.e., not having subsystems of the form $x_i = x_j$), we know that every 1×2 subsystem of L is uncommon. By Lemma 5.31, L is uncommon.

If $s(L) = 3$ and L has a 2×4 subsystem, then by Theorem 1.1 of [17], L is uncommon.

Suppose from now that $s(L) = 3$ and L does not have a 2×4 subsystem. If L has two linearly independent 1×3 subsystems, then it is isomorphic to

$$\begin{pmatrix} a_1 & b_1 & 0 & 0 & c \\ 0 & 0 & a_2 & b_2 & c \end{pmatrix}.$$

In this case, we have $c(L) = 4$ and $\mathcal{C}(L) = \{\{1, 2, 3, 4\}\}$, i.e., the only element in $\{L_B : B \in \mathcal{C}(L)\}$ is the linear equation

$$L_5 \mathbf{x} = a_1 x_1 + b_1 x_2 - a_2 x_3 - b_2 x_4.$$

If L_5 is uncommon, then we can find a Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ with $\sigma_{L_5}(g) < 0$, so L is uncommon over sufficiently large p by Theorem 3.5. We will show in Section 5.7 that, when L_5 is common, L is common if and only if it is of the form listed in (6) and (7).

¹Verifying this by hand is straightforward. Alternatively, readers can find a Mathematica code checking this at ‘case-d.nb’ in the supplemental file.

Now suppose that L has only one 1×3 subsystem. Again, by Theorem 3.5, if all 1×4 subsystems L are uncommon, then L is uncommon as well. Hence it suffices to consider the case where L has a common 1×4 subsystem, i.e., L is isomorphic to either of the following:

$$\begin{pmatrix} 1 & -1 & \lambda & -\lambda & 0 \\ a & b & 0 & 0 & c \end{pmatrix} \quad \begin{pmatrix} 1 & -1 & \lambda & -\lambda & 0 \\ a & 0 & b & 0 & c \end{pmatrix} \quad (8)$$

with $\lambda \in \mathbb{Q}$, $\lambda > 0$ and $a, b, c \in \mathbb{Z} \setminus \{0\}$.

In the first case, the elements in $\{L_B : B \in \mathcal{C}(L)\}$ are

$$\begin{aligned} L_5 \mathbf{x} &= x_1 - x_2 + \lambda x_3 - \lambda x_4 & A_5 &= \{1, -1, \lambda, -\lambda\} \\ L_1 \mathbf{x} &= (a+b)x_2 - a\lambda x_3 + a\lambda x_4 + cx_5 & A_1 &= \{a+b, -a\lambda, a\lambda, c\} \\ L_2 \mathbf{x} &= (a+b)x_1 + b\lambda x_3 - b\lambda x_4 + cx_5 & A_2 &= \{a+b, b\lambda, -b\lambda, c\}. \end{aligned}$$

We first show that, if $a+b+c \neq 0$, then L is uncommon. We prove this by finding Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ such that $\sigma_{L_1}(g) + \sigma_{L_2}(g) + \sigma_{L_5}(g) < 0$ and applying Theorem 3.5. The construction of suitable Fourier templates uses nearly identical ideas as that for Observation 5.16 and Proposition 5.17. Indeed, we have the following cases:

- Each of A_1, A_2 has three positive and one negative elements. Consider $h : \pm G_L \rightarrow \mathbb{C}$ with

$$\begin{cases} h(r) = i & r > 0 \\ h(r) = -i & r < 0. \end{cases}$$

We know that each of $h_{A_1}, h_{A_2}, h_{A_5}$ is constant on $\pm G_L$, with $h_{A_5} = 1$ and $h_{A_1} = h_{A_2} = -1$. Hence Lemma 5.14 applies.

- Each of A_1, A_2 contains two positive and two negative elements, with $a+b+c \neq 0$. Then since A_1, A_2 are not cancelling, we can find some $h : \pm G_L \rightarrow \mathbb{C}$ as in Proposition 5.17, so that each of $\text{Re}(h_{A_1}), \text{Re}(h_{A_2}), h_{A_5}$ is constant on $\pm G_L$, and $\text{Re}(h_{A_1}) + \text{Re}(h_{A_2}) + h_{A_5} < 0$. Hence Lemma 5.14 applies.

Finally, for $a+b+c = 0$, we show in Section 5.7 that L is common if and only if it is of the form listed in (6) and (7).

Lastly, we have the second case, where the elements in $\{L_B : B \in \mathcal{C}(L)\}$ are given by

$$\begin{aligned} L_5 \mathbf{x} &= x_1 - x_2 + \lambda x_3 - \lambda x_4 & A_5 &= \{1, -1, \lambda, -\lambda\} \\ L_1 \mathbf{x} &= ax_2 + (b-a\lambda)x_3 + a\lambda x_4 + cx_5 & A_1 &= \{b-a\lambda, a, a\lambda, c\} \\ L_3 \mathbf{x} &= (a\lambda-b)x_1 + bx_2 + b\lambda x_4 + c\lambda x_5 & A_3 &= \{a\lambda-b, b, b\lambda, c\lambda\}. \end{aligned} \quad (9)$$

Notice that we overlap with the previous case if $\lambda = -1$. We handle the typical case $\lambda \neq 1$ and defer the case $\lambda = 1$ to the appendix; the construction for the case when $\lambda = 1$ uses the same idea as that in Example 5.20, Example 5.21 and the proof of Proposition 5.22.

When $\lambda \notin \{1, -1\}$, the key idea is to reduce to the setup of Lemma 5.9 and Lemma 5.10. Namely, Lemma 5.9 and Lemma 5.10 together show that if the A_i corresponding to L cannot be partitioned into two L -coincidental pairs, then L is uncommon. For the system of L above, we will show that this is the case because of parity reasons.

Suppose $\lambda \notin \{1, -1\}$. In this case, note that at least one of $|\frac{b-a\lambda}{c}|$ and $|\frac{a\lambda-b}{c\lambda}|$ is not an odd power of λ . Without loss of generality, suppose $|\frac{a\lambda-b}{c\lambda}|$ is not an odd power of λ . Note that L_5 is λ -common. Moreover, if L_1 is common, then it must be λ -common as well. This implies that $A_3 = \{a\lambda-b, b, b\lambda, c\lambda\}$ cannot be partitioned into L -coincidental pairs, so L is uncommon.

5.7. Proofs of commonality. In this section, we show the positive side of Theorem 1.10. That is, L is common if it is one of the forms listed in Theorem 1.10(1).

5.7.1. *Common 2×5 linear systems with $s(L) = 4$.* In this section, we show that the three 2×5 linear systems

$$\begin{pmatrix} 1 & -1 & 1 & -1 & 0 \\ 1 & 2 & -1 & 0 & -2 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & -1 & 2 & -2 \\ 0 & 1 & 2 & -1 & -2 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & -1 & -1 & 0 \\ 2 & -2 & 1 & 0 & -1 \end{pmatrix}$$

are common over all \mathbb{F}_p . Proofs that the first two linear systems are common can also be found at [18, Examples 4.5 and 4.6].

Let L be an irredundant 2×5 linear system with $s(L) = 4$. For $f : \mathbb{F}_p^n \rightarrow [0, 1]$ with $\mathbb{E}f = \alpha$, we have

$$\begin{aligned} t_L(f) + t_L(1-f) &= \mathbb{E}_{L\mathbf{x}=0}[f(x_1) \cdots f(x_5) + (1-f(x_1)) \cdots (1-f(x_5))] \\ &= 1 - 5\alpha + \binom{5}{2}\alpha^2 - \binom{5}{3}\alpha^3 + \sum_{i=1}^5 \mathbb{E}_{L\mathbf{x}=0} \left[\prod_{j \in [5] \setminus \{i\}} f(x_j) \right] \\ &= (1-\alpha)^5 + \alpha^5 - 5\alpha^4 + \sum_{i=1}^5 t_{L_i}(f) \stackrel{(2)}{=} (1-\alpha)^5 + \alpha^5 + \sum_{i=1}^5 \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} \prod_{a \in A_i} \widehat{f}(ar). \end{aligned}$$

Thus, if

$$\sum_{i=1}^5 \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} \prod_{a \in A_i} \widehat{f}(ar) \geq 0 \quad \text{for all } f : \mathbb{F}_p^n \rightarrow [0, 1], \quad (10)$$

we would have $t_L(f) + t_L(1-f) \geq 2^{-4}$ for all f , which implies that L is common over all \mathbb{F}_p .

We now verify that (10) holds for the three linear systems mentioned above. The first one has

$$\begin{aligned} L_5\mathbf{x} &= x_1 - x_2 + x_3 - x_4 & A_5 &= \{1, 1, -1, -1\} \\ L_4\mathbf{x} &= x_1 + 2x_2 - x_3 - 2x_5 & A_4 &= \{1, -1, 2, -2\} \\ L_3\mathbf{x} &= 2x_1 + x_2 - x_4 - 2x_5 & A_3 &= \{1, -1, 2, -2\} \\ L_2\mathbf{x} &= 3x_1 + x_3 - 2x_4 - 2x_5 & A_2 &= \{1, 3, -2, -2\} \\ L_1\mathbf{x} &= 3x_2 - 2x_3 + x_4 - 2x_5 & A_1 &= \{1, 3, -2, -2\}. \end{aligned}$$

For all $f : \mathbb{F}_p^n \rightarrow [0, 1]$, we have

$$\sum_{i=1}^5 \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} \prod_{a \in A_i} \widehat{f}(ar) = \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} 2|\widehat{f}(r)|^2|\widehat{f}(2r)|^2 + |\widehat{f}(r)|^2|\widehat{f}(3r)|^2 + 2\widehat{f}(r)\widehat{f}(2r)\widehat{f}(3r)\widehat{f}(-6r).$$

Using the Cauchy–Schwarz inequality, we get that

$$\left| \sum_r \widehat{f}(r)\widehat{f}(2r)\widehat{f}(3r)\widehat{f}(-6r) \right| \leq \sqrt{\left(\sum_r |\widehat{f}(r)|^2|\widehat{f}(2r)|^2 \right) \left(\sum_r |\widehat{f}(3r)|^2|\widehat{f}(-6r)|^2 \right)} = \sum_r |\widehat{f}(r)|^2|\widehat{f}(2r)|^2,$$

which implies that $\sum_{i=1}^5 \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} \prod_{a \in A_i} \widehat{f}(ar) \geq 0$ always.

The second linear system has

$$\begin{aligned} L_5\mathbf{x} &= x_1 - x_2 - 3x_3 + 3x_4 & A_5 &= \{1, -1, 3, -3\} \\ L_4\mathbf{x} &= x_1 + 2x_2 + 3x_3 - 6x_5 & A_4 &= \{1, 2, 3, -6\} \\ L_3\mathbf{x} &= 2x_1 + x_2 + 3x_4 - 6x_5 & A_3 &= \{1, 2, 3, -6\} \\ L_2\mathbf{x} &= x_1 - x_3 + 2x_4 - 2x_5 & A_2 &= \{1, -1, 2, -2\} \\ L_1\mathbf{x} &= x_2 + 2x_3 - x_4 - 2x_5 & A_1 &= \{1, -1, 2, -2\}. \end{aligned}$$

For all $f : \mathbb{F}_p^n \rightarrow [0, 1]$, we have

$$\sum_{i=1}^5 \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} \prod_{a \in A_i} \widehat{f}(ar) = \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} |\widehat{f}(r)|^4 + 2|\widehat{f}(r)|^2|\widehat{f}(2r)|^2 + 2\widehat{f}(r)\widehat{f}(3r)\widehat{f}(-2r)^2.$$

Using the Cauchy–Schwarz inequality twice, we get that

$$\begin{aligned} \left| \sum_r \widehat{f}(r)\widehat{f}(3r)\widehat{f}(-2r)^2 \right| &\leq \left(\sum_r |\widehat{f}(r)|^2 |\widehat{f}(-2r)|^2 \right)^{1/2} \left(\sum_r |\widehat{f}(-2r)|^2 |\widehat{f}(3r)|^2 \right)^{1/2} \\ &\leq \left(\sum_r |\widehat{f}(r)|^2 |\widehat{f}(-2r)|^2 \right)^{1/2} \left(\frac{1}{2} \left(\sum_r |\widehat{f}(-2r)|^4 + \sum_r |\widehat{f}(3r)|^4 \right) \right)^{1/2} \\ &\leq \frac{1}{2} \sum_r |\widehat{f}(r)|^2 |\widehat{f}(-2r)|^2 + \frac{1}{4} \left(\sum_r |\widehat{f}(-2r)|^4 + \sum_r |\widehat{f}(3r)|^4 \right) \\ &= \frac{1}{2} \sum_r |\widehat{f}(r)|^2 |\widehat{f}(2r)|^2 + \frac{1}{2} \sum_r |\widehat{f}(r)|^4, \end{aligned}$$

which implies that $\sum_{i=1}^5 \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} \prod_{a \in A_i} \widehat{f}(ar) \geq 0$ always.

The third linear system has

$$\begin{aligned} L_5 \mathbf{x} &= x_1 - x_2 + x_3 - x_4 & A_5 &= \{1, 1, -1, -1\} \\ L_4 \mathbf{x} &= 2x_1 - 2x_2 + x_3 - x_5 & A_4 &= \{2, -2, 1, -1\} \\ L_3 \mathbf{x} &= 3x_1 - x_2 - x_4 - x_5 & A_3 &= \{3, -1, -1, -1\} \\ L_2 \mathbf{x} &= 4x_1 - x_3 - 2x_4 - x_5 & A_2 &= \{4, -1, -2, -1\} \\ L_1 \mathbf{x} &= 4x_2 - 3x_3 - 2x_4 + x_5 & A_1 &= \{4, -3, -2, 1\}. \end{aligned}$$

For all $f : \mathbb{F}_p^n \rightarrow [0, 1]$, we have

$$\begin{aligned} &\sum_{i=1}^5 \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} \prod_{a \in A_i} \widehat{f}(ar) \\ &= \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} |\widehat{f}(r)|^4 + |\widehat{f}(r)|^2 |\widehat{f}(2r)|^2 + \widehat{f}(3r)\widehat{f}(-r)^3 + \widehat{f}(4r)\widehat{f}(-2r)\widehat{f}(-r)^2 + \widehat{f}(4r)\widehat{f}(-3r)\widehat{f}(-2r)\widehat{f}(r) \\ &= \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} \frac{3}{4} |\widehat{f}(r)|^4 + \frac{1}{4} |\widehat{f}(3r)|^4 + |\widehat{f}(2r)|^2 |\widehat{f}(4r)|^2 \\ &\quad + \operatorname{Re}(\widehat{f}(3r)\widehat{f}(-r)^3 + \widehat{f}(4r)\widehat{f}(-2r)\widehat{f}(-r)^2 + \widehat{f}(4r)\widehat{f}(-3r)\widehat{f}(-2r)\widehat{f}(r)). \end{aligned}$$

Thus, to show that L is common, it suffices to show that

$$\begin{aligned} &\frac{3}{4} |\widehat{f}(r)|^4 + \frac{1}{4} |\widehat{f}(3r)|^4 + |\widehat{f}(2r)|^2 |\widehat{f}(4r)|^2 \\ &\quad + \operatorname{Re}(\widehat{f}(3r)\widehat{f}(-r)^3 + \widehat{f}(4r)\widehat{f}(-2r)\widehat{f}(-r)^2 + \widehat{f}(4r)\widehat{f}(-3r)\widehat{f}(-2r)\widehat{f}(r)) \geq 0 \quad \text{for all } r \in \widehat{\mathbb{F}}_p^n. \end{aligned}$$

Consider any $r \in \widehat{\mathbb{F}}_p^n$. Let

$$a_1 = |\widehat{f}(r)|, \quad a_2 = |\widehat{f}(2r)|, \quad a_3 = |\widehat{f}(3r)|, \quad a_4 = |\widehat{f}(4r)|.$$

Since

$$\widehat{f}(3r)\widehat{f}(-r)^3 \cdot \widehat{f}(4r)\widehat{f}(-3r)\widehat{f}(-2r)\widehat{f}(r) = \widehat{f}(4r)\widehat{f}(-2r)\widehat{f}(-r)^2 \cdot |\widehat{f}(r)|^2 |\widehat{f}(3r)|^2,$$

letting α, β be the arguments of $\widehat{f}(3r)\widehat{f}(-r)^3$, $\widehat{f}(4r)\widehat{f}(-3r)\widehat{f}(-2r)\widehat{f}(r)$ respectively, our goal becomes showing that

$$I := \frac{3}{4}a_1^4 + \frac{1}{4}a_3^4 + a_2^2a_4^2 + a_1^3a_3 \cos(\alpha) + a_1^2a_2a_4 \cos(\alpha + \beta) + a_1a_2a_3a_4 \cos(\beta) \geq 0. \quad (11)$$

Setting $x = a_1^2$, $y = a_1a_3$ and $z = a_2a_4$, we have

$$\begin{aligned} I &\geq \frac{1}{2}a_1^4 + \frac{1}{2}a_1^2a_3^2 + a_2^2a_4^2 + a_1^3a_3 \cos(\alpha) + a_1^2a_2a_4 \cos(\alpha + \beta) + a_1a_2a_3a_4 \cos(\beta) \\ &= \frac{1}{2}(x^2 + y^2 + 2z^2 + 2xy \cos(\alpha) + 2xz \cos(\alpha + \beta) + 2yz \cos(\beta)). \end{aligned}$$

Since

$$\begin{aligned} 0 &\leq (y + z \cos(\beta) + x \cos(\alpha))^2 + (z \sin(\beta) - x \sin(\alpha))^2 \\ &= y^2 + z^2 + x^2 + 2yz \cos(\beta) + 2xy \cos(\alpha) + 2xz \cos(\alpha + \beta), \end{aligned}$$

we conclude that $I \geq 0$. This gives (11).

5.7.2. *Common 2×5 linear systems with $s(L) = 3$.* We first show that linear systems of the form

$$L = \begin{pmatrix} a & b & 0 & 0 & c \\ 0 & 0 & a & b & c \end{pmatrix} \quad a, b, c \in \mathbb{Z} \setminus \{0\}$$

are common. Define

$$\begin{aligned} L_1\mathbf{x} &= ax_1 + bx_2 + cx_5 \\ L_2\mathbf{x} &= ax_1 + bx_2 - ax_3 - bx_4. \end{aligned}$$

Consider any $f : \mathbb{F}_p^n \rightarrow [0, 1]$ with $\mathbb{E}f = \alpha \in [0, 1]$. For any $\{i, j, k\} \in \binom{[5]}{3}$, we have

$$\mathbb{E}_{L\mathbf{x}=0} f(x_i)f(x_j)f(x_k) = \begin{cases} t_{L_1}(f) & \{i, j, k\} = \{1, 2, 5\} \text{ or } \{3, 4, 5\} \\ \alpha^3 & \text{otherwise.} \end{cases}$$

For any $\{i, j, k, \ell\} \in \binom{[5]}{4}$, we have

$$\mathbb{E}_{L\mathbf{x}=0} f(x_i)f(x_j)f(x_k)f(x_\ell) = \begin{cases} t_{L_2}(f) & \{i, j, k, \ell\} = \{1, 2, 3, 4\} \\ \alpha t_{L_1}(f) & \text{otherwise.} \end{cases}$$

Thus, we have

$$\begin{aligned} t_L(f) + t_L(1-f) &= \mathbb{E}_{L\mathbf{x}=0} f(x_1) \cdots f(x_5) + (1-f(x_1)) \cdots (1-f(x_5)) \\ &= 1 - 5\alpha + 10\alpha^2 - (8\alpha^3 + 2t_{L_1}(f)) + (t_{L_2}(f) + 4\alpha t_{L_1}(f)). \end{aligned}$$

Since

$$\begin{aligned} t_{L_1}(f) &= \sum_{r \in \widehat{\mathbb{F}}_p^n} \widehat{f}(ar)\widehat{f}(br)\widehat{f}(cr) = \alpha^3 + \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} \widehat{f}(ar)\widehat{f}(br)\widehat{f}(cr), \\ t_{L_2}(f) &= \sum_{r \in \widehat{\mathbb{F}}_p^n} |\widehat{f}(ar)|^2 |\widehat{f}(br)|^2 = \alpha^4 + \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} |\widehat{f}(ar)|^2 |\widehat{f}(br)|^2, \end{aligned}$$

we get that

$$t_L(f) + t_L(1-f) = \alpha^5 + (1-\alpha)^5 + (4\alpha - 2) \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} \widehat{f}(ar)\widehat{f}(br)\widehat{f}(cr) + \sum_{r \in \widehat{\mathbb{F}}_p^n \setminus \{0\}} |\widehat{f}(ar)|^2 |\widehat{f}(br)|^2.$$

Letting

$$X = \sum_{r \in \widehat{\mathbb{F}_p^n} \setminus \{0\}} \widehat{f}(ar)\widehat{f}(br)\widehat{f}(cr), \quad Y = \sum_{r \in \widehat{\mathbb{F}_p^n} \setminus \{0\}} |\widehat{f}(ar)|^2 |\widehat{f}(br)|^2,$$

we know from Cauchy–Schwarz and Parseval that

$$X^2 \leq Y \sum_{r \in \widehat{\mathbb{F}_p^n} \setminus \{0\}} |\widehat{f}(cr)|^2 = Y \sum_{r \in \widehat{\mathbb{F}_p^n} \setminus \{0\}} |\widehat{f}(r)|^2 = Y(\mathbb{E}[|f|^2] - \alpha^2) \leq Y(\|f\|_\infty \mathbb{E}[f] - \alpha^2) \leq Y(\alpha - \alpha^2).$$

Since $0 \leq Y = t_{L_2}(f) - \alpha^4 \leq 1$, we get that

$$t_L(f) + t_L(1-f) \geq \min_{0 \leq \alpha, Y \leq 1} \alpha^5 + (1 - \alpha^5) - |4\alpha - 2| \sqrt{Y(\alpha - \alpha^2)} + Y \geq \frac{1}{16}.$$

Hence L is common.

Next, we show that linear systems of the form

$$L = \begin{pmatrix} 1 & -1 & \lambda & -\lambda & 0 \\ a & b & 0 & 0 & -a-b \end{pmatrix}, \quad \{|a/b|, |(a+b)/b|, |(a+b)/a|\} \cap \{|\lambda|, |\lambda|^{-1}\} \neq \emptyset$$

are common. Define

$$\begin{aligned} L_1 \mathbf{x} &= ax_1 + bx_2 - (a+b)x_5 \\ L_2 \mathbf{x} &= x_1 - x_2 + \lambda x_3 - \lambda x_4 \\ L_3 \mathbf{x} &= (a+b)x_1 + b\lambda x_3 - b\lambda x_4 - (a+b)x_4 \\ L_4 \mathbf{x} &= (a+b)x_2 - a\lambda x_2 + a\lambda x_3 - (a+b)x_4. \end{aligned}$$

For all $f : \mathbb{F}_p^n \rightarrow [0, 1]$ with $\mathbb{E}f = \alpha \in [0, 1]$, again we can rewrite

$$\begin{aligned} t_L(f) + t_L(1-f) &= \alpha^5 + (1 - \alpha)^5 + (2\alpha - 1) \sum_{r \in \widehat{\mathbb{F}_p^n} \setminus \{0\}} \widehat{f}(ar)\widehat{f}(br)\overline{\widehat{f}((a+b)r)} + \sum_{r \in \widehat{\mathbb{F}_p^n} \setminus \{0\}} |\widehat{f}(r)|^2 |\widehat{f}(\lambda r)|^2 \\ &\quad + \sum_{r \in \widehat{\mathbb{F}_p^n} \setminus \{0\}} |\widehat{f}(b\lambda r)|^2 |\widehat{f}((a+b)r)|^2 + \sum_{r \in \widehat{\mathbb{F}_p^n} \setminus \{0\}} |\widehat{f}(a\lambda r)|^2 |\widehat{f}((a+b)r)|^2. \end{aligned}$$

Let

$$X = \sum_{r \in \widehat{\mathbb{F}_p^n} \setminus \{0\}} \widehat{f}(ar)\widehat{f}(br)\overline{\widehat{f}((a+b)r)}, \quad Y = \sum_{r \in \widehat{\mathbb{F}_p^n} \setminus \{0\}} |\widehat{f}(r)|^2 |\widehat{f}(\lambda r)|^2.$$

Since one of $|a/b|, |a/b|^{-1}, |(a+b)/b|, |(a+b)/b|^{-1}, |(a+b)/a|, |(a+b)/a|^{-1}$ equals $|\lambda|$, we know from Cauchy–Schwarz and Parseval that

$$X^2 \leq Y \sum_{r \in \widehat{\mathbb{F}_p^n} \setminus \{0\}} |\widehat{f}(r)|^2 = Y(\mathbb{E}[|f|^2] - \alpha^2) \leq Y(\|f\|_\infty \mathbb{E}[f] - \alpha^2) \leq Y(\alpha - \alpha^2).$$

Since $0 \leq Y \leq 1$, we get that

$$t_L(f) + t_L(1-f) \geq \min_{0 \leq \alpha, Y \leq 1} \alpha^5 + (1 - \alpha^5) - |2\alpha - 1| \sqrt{Y(\alpha - \alpha^2)} + Y \geq \frac{1}{16}.$$

Hence L is common.

Finally, we show that linear systems

$$\begin{pmatrix} a & -a & 0 & 0 & c \\ 0 & 0 & b & -b & c \end{pmatrix} \quad \begin{pmatrix} 1 & -1 & \lambda & -\lambda & 0 \\ a & b & 0 & 0 & -a-b \end{pmatrix}$$

not isomorphic to (6) and (7) are uncommon over sufficiently large \mathbb{F}_p . For the first linear system, define

$$\begin{aligned} L_1\mathbf{x} &= ax_1 - ax_2 + cx_5 \\ L_2\mathbf{x} &= bx_3 - bx_4 + cx_5 \\ L_3\mathbf{x} &= ax_1 - ax_2 + bx_3 - bx_4. \end{aligned}$$

Since the linear system is not isomorphic to (6), we have $|a| \neq |b|$. Since it is not isomorphic to (7), we also have $|a/b| \notin \{|a/c|, |a/c|^{-1}\}$. For all $f : \mathbb{F}_p^n \rightarrow [0, 1]$ with $\mathbb{E}f = \alpha \in [0, 1]$, we have

$$\begin{aligned} t_L(f) + t_L(1-f) &= \alpha^5 + (1-\alpha)^5 + (2\alpha-1) \sum_{r \in \widehat{\mathbb{F}_p} \setminus \{0\}} |\widehat{f}(ar)|^2 \widehat{f}(cr) \\ &\quad + (2\alpha-1) \sum_{r \in \widehat{\mathbb{F}_p} \setminus \{0\}} |\widehat{f}(br)|^2 \widehat{f}(cr) + \sum_{r \in \widehat{\mathbb{F}_p} \setminus \{0\}} |\widehat{f}(ar)|^2 |\widehat{f}(br)|^2. \end{aligned}$$

Suppose p is sufficiently large. Define $f : \mathbb{F}_p \rightarrow [0, 1]$ by

$$\widehat{f}(0) = 0.4995 := \alpha, \quad \widehat{f}(a) = \widehat{f}(-a) = \widehat{f}(c) = \widehat{f}(-c) = 0.0834 := \beta.$$

We then have

$$\begin{aligned} t_L(f) + t_L(1-f) &\leq \alpha^5 + (1-\alpha)^5 + (2\alpha-1) \sum_{r \in \widehat{\mathbb{F}_p} \setminus \{0\}} |\widehat{f}(ar)|^2 \widehat{f}(cr) \\ &= \alpha^5 + (1-\alpha)^5 + (2\alpha-1)2\beta^3 = 0.0624995 < 1/16. \end{aligned}$$

For the second linear system, define

$$\begin{aligned} L_1\mathbf{x} &= ax_1 + bx_2 - (a+b)x_5 \\ L_2\mathbf{x} &= x_1 - x_2 + \lambda x_3 - \lambda x_4 \\ L_3\mathbf{x} &= (a+b)x_1 + b\lambda x_3 - b\lambda x_4 - (a+b)x_4 \\ L_4\mathbf{x} &= (a+b)x_2 - a\lambda x_2 + a\lambda x_3 - (a+b)x_4. \end{aligned}$$

Since the linear system is not isomorphic to (7), we know that $\lambda \notin \{1, -1\}$. Moreover, we have

$$\{|a/b|, |(a+b)/b|, |(a+b)/a|\} \cap \{|\lambda|, |\lambda|^{-1}, |(a+b)/(b\lambda)|, |b\lambda/(a+b)|, |(a+b)/(a\lambda)|, |a\lambda/(a+b)|\} = \emptyset.$$

For all $f : \mathbb{F}_p^n \rightarrow [0, 1]$ with $\mathbb{E}f = \alpha \in [0, 1]$, we have

$$\begin{aligned} t_L(f) + t_L(1-f) &= \alpha^5 + (1-\alpha)^5 + (2\alpha-1) \sum_{r \in \widehat{\mathbb{F}_p}} \widehat{f}(ar) \widehat{f}(br) \overline{\widehat{f}((a+b)r)} \\ &\quad + \sum_{r \in \widehat{\mathbb{F}_p} \setminus \{0\}} |\widehat{f}(r)|^2 |\widehat{f}(\lambda r)|^2 + |\widehat{f}((a+b)r)|^2 |\widehat{f}(b\lambda r)|^2 + |\widehat{f}((a+b)r)|^2 |\widehat{f}(a\lambda r)|^2. \end{aligned}$$

Suppose p is sufficiently large. Define $f : \mathbb{F}_p \rightarrow [0, 1]$ by

$$\widehat{f}(0) = 0.4995 := \alpha, \quad \widehat{f}(a) = \widehat{f}(-a) = \widehat{f}(b) = \widehat{f}(-b) = \widehat{f}(a+b) = \widehat{f}(-a-b) = 0.0834 := \beta.$$

We then have

$$\begin{aligned} t_L(f) &= \alpha^5 + (1-\alpha)^5 + (2\alpha-1) \sum_{r \in \widehat{\mathbb{F}_p} \setminus \{0\}} \widehat{f}(ar) \widehat{f}(br) \overline{\widehat{f}((a+b)r)} \\ &= \alpha^5 + (1-\alpha)^5 + (2\alpha-1)2\beta^3 = 0.0624995 < 1/16. \end{aligned}$$

REFERENCES

- [1] Daniel Altman, *Local aspects of the Sidorenko property for linear equations*, (2022), arXiv:2210.17493.
- [2] Daniel Altman, *On a common-extendable, non-Sidorenko linear system*, *Comb. Theory* **3** (2023), Paper No. 5, 12.
- [3] Daniel Altman and Anita Liebenau, *On the uncommonness of minimal rank-2 systems of linear equations*, (2024), arXiv:2404.18908.
- [4] P. Baldi, Y. Rinott, and C. Stein, *A normal approximation for the number of local maxima of a random function on a graph*, *Probability, statistics, and mathematics*, Academic Press, Boston, MA, 1989, pp. 59–81.
- [5] A. D. Barbour and Louis H. Y. Chen, *An introduction to Stein's method*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2005.
- [6] G. R. Blakley and Prabir Roy, *A Hölder type inequality for symmetric matrices with nonnegative entries*, *Proc. Amer. Math. Soc.* **16** (1965), 1244–1245.
- [7] Stefan A. Burr and Vera Rosta, *On the Ramsey multiplicities of graphs—problems and recent results*, *J. Graph Theory* **4** (1980), 347–361.
- [8] David Conlon, Jacob Fox, and Benny Sudakov, *An approximate version of Sidorenko's conjecture*, *Geom. Funct. Anal.* **20** (2010), 1354–1366.
- [9] David Conlon, Jeong Han Kim, Choongbum Lee, and Joonkyung Lee, *Some advances on Sidorenko's conjecture*, *J. Lond. Math. Soc. (2)* **98** (2018), 593–608.
- [10] David Conlon and Joonkyung Lee, *Finite reflection groups and graph norms*, *Adv. Math.* **315** (2017), 130–165.
- [11] David Conlon and Joonkyung Lee, *Sidorenko's conjecture for blow-ups*, *Discrete Anal.* (2021), Paper No. 2, 13.
- [12] David Conlon, Joonkyung Lee, and Alexander Sidorenko, *Extremal numbers and sidorenko's conjecture*, (2023), arXiv:2307.04588.
- [13] Jacob Fox, Huy Tuan Pham, and Yufei Zhao, *Common and Sidorenko linear equations*, *Q. J. Math.* **72** (2021), 1223–1234.
- [14] A. W. Goodman, *On sets of acquaintances and strangers at any party*, *Amer. Math. Monthly* **66** (1959), 778–783.
- [15] Hamed Hatami, *Graph norms and Sidorenko's conjecture*, *Israel J. Math.* **175** (2010), 125–150.
- [16] Chris Jagger, Pavel Štoviček, and Andrew Thomason, *Multiplicities of subgraphs*, *Combinatorica* **16** (1996), 123–141.
- [17] Nina Kamčev, Anita Liebenau, and Natasha Morrison, *On uncommon systems of equations*, (2021), arXiv:2106.08986.
- [18] Nina Kamčev, Anita Liebenau, and Natasha Morrison, *Towards a characterization of Sidorenko systems*, *Q. J. Math.* **74** (2023), 957–974.
- [19] Jeong Han Kim, Choongbum Lee, and Joonkyung Lee, *Two approaches to Sidorenko's conjecture*, *Trans. Amer. Math. Soc.* **368** (2016), 5057–5074.
- [20] László Lovász, *Subgraph densities in signed graphons and the local Simonovits-Sidorenko conjecture*, *Electron. J. Combin.* **18** (2011), Paper 127, 21.
- [21] Nathan Ross, *Fundamentals of Stein's method*, *Probab. Surv.* **8** (2011), 210–293.
- [22] A. Saad and J. Wolf, *Ramsey multiplicity of linear patterns in certain finite abelian groups*, *Q. J. Math.* **68** (2017), 125–140.
- [23] A. F. Sidorenko, *Inequalities for functionals generated by bipartite graphs*, *Diskret. Mat.* **3** (1991), 50–65.
- [24] Alexander Sidorenko, *A correlation inequality for bipartite graphs*, *Graphs Combin.* **9** (1993), 201–204.
- [25] Balazs Szegedy, *An information theoretic approach to Sidorenko's conjecture*, (2014), arXiv:1406.6738.
- [26] Andrew Thomason, *A disproof of a conjecture of Erdős in Ramsey theory*, *J. London Math. Soc. (2)* **39** (1989), 246–255.
- [27] Leo Versteegen, *Linear configurations containing 4-term arithmetic progressions are uncommon*, *J. Combin. Theory Ser. A* **200** (2023), Paper No. 105792, 38.

APPENDIX A. PROOF OF LEMMA 4.6

In this section, we prove Lemma 4.6.

Proof of Lemma 4.6. We will prove that

$$d_{\text{Wass}}(W, Z) \leq \rho^{-3} \sum_{u \in V} \left(3 \sum_{v, w \in N_u} |\mathbb{E}[X_u X_v X_w]| + 4 \sum_{v \in N_u} |\mathbb{E}[X_u X_v]| \cdot \mathbb{E} \left[\left\| \sum_{w \in N_u \cup N_v} X_w \right\| \right] \right).$$

This will imply the desired conclusion because we have the following (see for instance [5, Theorem 3.1]) for any $x \in \mathbb{R}$:

$$|\mathbb{P}[W \leq x] - \mathbb{P}[Z \leq x]| \leq \sqrt{2d_{\text{Wass}}(W, Z)/\pi}.$$

Define the set of test functions

$$\mathcal{D} = \left\{ f \in \mathcal{C}^1(\mathbb{R}) : f' \text{ absolutely continuous, } \|f\|_\infty \leq 1, \|f'\|_\infty \leq \sqrt{2/\pi}, \|f''\|_\infty \leq 2 \right\}.$$

By Corollary 4.4, it follows that

$$d_{\text{Wass}}(W, Z) \leq \sup_{f \in \mathcal{D}} |\mathbb{E}[f'(W) - Wf(W)]|.$$

Define

$$Y_v = \rho^{-1} \sum_{u \in N_v} X_u, \quad J_v = \rho^{-1} \sum_{u \notin N_v} X_u, \quad K_{vw} = \rho^{-1} \sum_{u \notin N_w \cup N_v} X_u, \quad T_{vw} = \rho^{-1} \sum_{u \in N_w \setminus N_v} X_u.$$

In particular, observe that for any choice of $v, w \in V$, we have $Y_v + K_{vw} + T_{vw} = W$.

Write $\mathbb{E}[Wf(W) - f'(W)] = \text{(I)} + \text{(II)} + \text{(III)}$ as a telescoping sum, where

$$\begin{aligned} \text{(I)} &= \mathbb{E}Wf(W) - \rho^{-1} \sum_{v \in V} \mathbb{E}X_v Y_v f'(J_v), \\ \text{(II)} &= \rho^{-1} \sum_{v \in V} \mathbb{E}X_v Y_v f'(J_v) - \rho^{-2} \sum_{v \in V} \sum_{u \in N_v} \mathbb{E}[X_u X_v] \mathbb{E}[f'(K_{uv})], \\ \text{(III)} &= \rho^{-2} \sum_{v \in V} \sum_{u \in N_v} \mathbb{E}[X_u X_v] (\mathbb{E}f'(K_{uv}) - \mathbb{E}f'(W)). \end{aligned}$$

By the triangle inequality, we have

$$|\mathbb{E}Wf(W) - \mathbb{E}f'(W)| \leq |\text{(I)}| + |\text{(II)}| + |\text{(III)}|. \quad (12)$$

In what remains, we upper bound each of the individual pieces $|\text{(I)}|$, $|\text{(II)}|$ and $|\text{(III)}|$.

For (I), we write $W = Y_v + J_v$ and by Taylor expansion about J_v to obtain

$$Wf(W) = \rho^{-1} \sum_{v \in V} \left(X_v f(J_v) + X_v Y_v f'(J_v) + \frac{1}{2} X_v Y_v^2 f''(J_1) \right),$$

where J_1 is a random variable in the interval $[J_v, W]$. It follows by the assumptions on f that

$$|\text{(I)}| \leq \rho^{-1} \sum_{v \in V} |\mathbb{E}(X_v Y_v^2)| \leq \rho^{-3} \sum_{v \in V} \sum_{u, w \in N_v} |\mathbb{E}X_u X_v X_w|.$$

For (II), fix $v \in V$ and by writing $J_v = K_{vw} + T_{vw}$, we know from Taylor expansion about K_{vw} that

$$X_v Y_v f'(J_v) = \rho^{-1} \sum_{w \in N_v} X_v X_w (f'(K_{vw}) + T_{vw} f''(J_2))$$

where J_2 is a random variable in the interval $[K_{vw}, J_v]$. By the assumptions on f , we have

$$|\text{(II)}| \leq 2\rho^{-2} \sum_{v \in V} \left| \mathbb{E} \left[X_v \sum_{w \in N_v} X_w T_{vw} \right] \right| \leq 2\rho^{-3} \sum_{v \in V} \sum_{u, w \in N_v} |\mathbb{E}X_u X_v X_w|.$$

Finally, for (III), fix $u, v \in V$. Since $K_{uv} = W - (Y_u + T_{uv})$, by the mean value theorem, we have

$$f'(K_{uv}) = f'(W) - (Y_u + T_{uv}) f''(J_3)$$

for some $J_3 \in [K_{uv}, W]$. Consequently, we have

$$|\mathbb{E}f'(K_{uv}) - \mathbb{E}f'(W)| \leq 2\mathbb{E}|Y_u + T_{uv}| \leq 2\rho^{-1} \mathbb{E} \left| \sum_{w \in N_u \cup N_v} X_w \right|.$$

Putting all the various pieces into (12) gives the desired bound. \square

APPENDIX B. DEFERRED PROOFS FROM SECTION 5

B.1. Case A.

Proof of Lemma 5.8. Suppose first that exactly one of L_1, \dots, L_5 is common. Without loss of generality, let

$$L_5 \mathbf{x} = x_1 - x_2 + \lambda x_3 - \lambda x_4, \quad \lambda \in \mathbb{Q} \setminus \{-1, 0, 1\}.$$

We use $a \sim b$ to denote that a, b are L -coincidental, i.e., $\frac{a}{b} \in \{\lambda, -\lambda, 1/\lambda, -1/\lambda\}$. Up to replacing λ by one of $-\lambda, 1/\lambda, -1/\lambda$ and permuting the variables, we have the following two cases:

- (1) For every $i = 1, \dots, 4$, letting $L_i \mathbf{x} = a_{i,1}x_1 + \dots + a_{i,5}x_5$, we have $a_{i,1} \sim a_{i,2}$ or $a_{i,3} \sim a_{i,4}$.
- (2) Letting $L_4 \mathbf{x} = a_{4,1}x_1 + a_{4,2}x_2 + a_{4,3}x_3 + a_{4,5}x_5$, we have $a_{4,1} \sim a_{4,3}$ and $a_{4,2} \sim a_{4,5}$.

Write

$$\begin{aligned} L_5 \mathbf{x} &= x_1 - x_2 + \lambda x_3 - \lambda x_4 & A_5 &= \{\{1, -1, \lambda, -\lambda\}\} \\ L_4 \mathbf{x} &= ax_1 + bx_2 + cx_3 + dx_5 & A_4 &= \{\{a, b, c, d\}\} \\ L_3 \mathbf{x} &= (\lambda a - c)x_1 + (\lambda b + c)x_2 + \lambda cx_4 + \lambda dx_5 & A_3 &= \{\{\lambda a - c, \lambda b + c, \lambda c, \lambda d\}\} \\ L_2 \mathbf{x} &= (a + b)x_1 + (\lambda b + c)x_3 - \lambda bx_4 + dx_5 & A_2 &= \{\{a + b, \lambda b + c, -\lambda b, d\}\} \\ L_1 \mathbf{x} &= (a + b)x_2 + (c - \lambda a)x_3 + \lambda ax_4 + dx_5 & A_1 &= \{\{a + b, c - \lambda a, \lambda a, d\}\}. \end{aligned}$$

Suppose (1) occurs. if each of A_1, \dots, A_4 can be partitioned into two L -coincidental pairs, then we must have

$$a \sim b, \quad c \sim d, \quad (\lambda a - c) \sim (\lambda b + c), \quad (\lambda b + c) \sim \lambda b, \quad (c - \lambda a) \sim \lambda a, \quad a + b \sim d.$$

Recall that $a \sim b$ corresponds to one of the four identities $\lambda a = b$, $\lambda a = -b$, $\lambda b = a$, $\lambda b = -a$. Thus, for the above relation to hold, a, b, c, d, λ must satisfy a system of six equations on five variables, and there are 4^6 possible such systems of equations. A computer search² yields that none of these 4^6 systems of equations has a solution with $a, b, c, d \in \mathbb{Q} \setminus \{0\}$ and $\lambda \in \mathbb{Q} \setminus \{-1, 0, 1\}$.

Suppose (2) occurs. Similarly, if each of A_1, \dots, A_4 can be partitioned into two L -coincidental pairs, then we must have

$$\begin{aligned} a \sim c, \quad b \sim d, \\ ((\lambda a - c) \sim (\lambda b + c), \quad \lambda c \sim \lambda d) \quad \text{or} \quad ((\lambda a - c) \sim \lambda d, \quad (\lambda b + c) \sim \lambda c), \\ (-\lambda b \sim (\lambda b + c), \quad (a + b) \sim d) \quad \text{or} \quad ((\lambda b + c) \sim d, \quad (a + b) \sim -\lambda b), \\ ((c - \lambda a) \sim \lambda a, \quad (a + b) \sim d) \quad \text{or} \quad ((c - \lambda a) \sim (a + b), \quad \lambda a \sim d). \end{aligned}$$

This is because $b \sim d$, so we cannot have $-\lambda b \sim d$ in A_2 . Then we either have $(\lambda b + c) \sim d$ or $(\lambda b + c) \sim -\lambda b$ in A_2 , which means that $|(\lambda b + c)/d|$ is an odd power of $|\lambda|$, so we cannot have $(\lambda b + c) \sim \lambda d$ in A_3 . Then we either have $(\lambda a - c) \sim (\lambda b + c)$ or $(\lambda a - c) \sim \lambda d$ in A_3 , which means that $|(\lambda a - c)/d|$ is an even power of $|\lambda|$, so we cannot have $(c - \lambda a) \sim d$ in A_1 . Therefore, we obtain $8 \cdot 4^6$ possible systems of eight equations on five variables. Again, using a computer search³, one can verify that none of these $8 \cdot 4^6$ systems of equations has a solution with $a, b, c, d \in \mathbb{Q} \setminus \{0\}$ and $\lambda \in \mathbb{Q} \setminus \{-1, 0, 1\}$.

²The code can be found at ‘case-a-1.nb’ in the supplemental file.

³The code can be found at ‘case-a-2.nb’ in the supplemental file.

Now suppose two of L_1, \dots, L_5 are common. Without loss of generality, suppose

$$\begin{array}{ll} L_5 \mathbf{x} = x_1 - x_2 + \lambda x_3 - \lambda x_4 & A_5 = \{\{1, -1, \lambda, -\lambda\}\} \\ L_4 \mathbf{x} = x_1 + \mu x_2 - x_3 - \mu x_5 & A_4 = \{\{1, -1, \mu, -\mu\}\} \\ L_3 \mathbf{x} = (\lambda + 1)x_1 + (\lambda\mu - 1)x_2 - \lambda x_4 - \lambda\mu x_5 & A_3 = \{\{\lambda + 1, \lambda\mu - 1, -\lambda, -\lambda\mu\}\} \\ L_2 \mathbf{x} = (\mu + 1)x_1 + (\lambda\mu - 1)x_3 - \lambda\mu x_4 - \mu x_5 & A_2 = \{\{\mu + 1, \lambda\mu - 1, -\mu, -\lambda\mu\}\} \\ L_1 \mathbf{x} = (\mu + 1)x_2 + (-\lambda - 1)x_3 + \lambda x_4 - \mu x_5 & A_1 = \{\{\mu + 1, -\lambda - 1, \lambda, -\mu\}\} \end{array}$$

with $\lambda, \mu \in \mathbb{Q} \setminus \{-1, 0, 1\}$. Since L_4 and L_5 are both common, $a \sim b$ corresponds to one of the eight identities $\lambda a = b$, $\lambda a = -b$, $\lambda b = a$, $\lambda b = -a$, $\mu a = b$, $\mu a = -b$, $\mu b = a$, $\mu b = -a$. Since A_1 can be partitioned into two cancelling pairs, we must have

$$(\mu + 1) \sim (-\lambda - 1), \quad \lambda \sim -\mu$$

or

$$(\mu + 1) \sim \lambda, \quad (-\lambda - 1) \sim -\mu$$

or

$$(\mu + 1) \sim -\mu, \quad \lambda \sim (-\lambda - 1).$$

Thus, $\lambda, \mu \in \mathbb{Q} \setminus \{-1, 0, 1\}$ must satisfy one of the $3 \cdot 8^2$ possible systems of two equations on two variables. A computer search⁴ gives that this is only possible when (λ, μ) have the following values:

$$\begin{aligned} &(1/2, -1/4), (-1/4, 1/2), (-2, -1/3), (-1/3, -2), \\ &(-2, -2/3), (-2/3, -2), (1/2, -3/4), (-3/4, 1/2), \\ &(-2, 3), (3, -2), (-2, -5), (-5, 2), (-2, -2), (\lambda, -\lambda/(1 + \lambda)). \end{aligned}$$

This is what we obtain just by assuming that A_1 can be partitioned into two cancelling pairs. One can then verify that A_2, A_3 cannot be both partitioned into L -coincidental pairs in any of these solutions. \square

Proof of Lemma 5.9. Let E denote the collection of those $i \in [5]$ that satisfies the condition in Lemma 5.9. That is, for every $i \in E$, there exists some $a \in A_i$ such that a, b are not L -coincidental for any $b \in A_i$.

Suppose there exists $i_0 \in E$ with $A_{i_0} = \{\{1, 1, 1, 1\}\}$ or $\{\{-1, -1, -1, -1\}\}$. In this case, the Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ defined by

$$\begin{cases} g(1) = e(1/8) \\ g(-1) = e(-1/8) \\ g(r) = 0 & \text{otherwise} \end{cases}$$

has $\sum_{i=1}^5 \sigma_{A_i}(g) \leq -2 < 0$.

Otherwise, suppose there exists $i_0 \in E$ with $A_{i_0} = \{\{1, 1, 1, -1\}\}$ or $\{\{1, -1, -1, -1\}\}$. In this case, the Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ defined by

$$\begin{cases} g(1) = i \\ g(-1) = -i \\ g(r) = 0 & \text{otherwise} \end{cases}$$

has $\sum_{i=1}^5 \sigma_{A_i}(g) \leq -2 < 0$.

⁴The code can be found at 'case-a-3.nb' in the supplemental file.

Otherwise, suppose there exists $i_0 \in E$ with $A_{i_0} = \{\{a, a, a, b\}\}$ or $\{\{a, a, -a, b\}\}$ with $|a| \neq |b|$. In this case, the Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ defined by

$$\begin{cases} g(a) = g(-a) = C \\ g(b) = g(-b) = -1 \\ g(r) = 0 \quad \text{otherwise} \end{cases}$$

has $\sum_{i=1}^5 \sigma_{A_i}(g) = -2C^3 + O(C^2)$, which is negative for $C > 0$ sufficiently large.

Otherwise, suppose there exists $i_0 \in E$ with $A_{i_0} = \{\{a, a, b, b\}\}$ with $|a| \neq |b|$. In this case, consider the joined Fourier template $h : \mathbb{Z}^2 \rightarrow \mathbb{C}$ defined by $h(r_1, r_2) = g(r_1)\tilde{g}(r_2)$, where $g, \tilde{g} : \mathbb{Z} \rightarrow \mathbb{C}$ are 1-dimensional Fourier templates given by

$$\begin{cases} g(a) = g(b) = e(1/8) \\ g(-a) = g(-b) = e(-1/8) \\ g(r) = 0 \quad \text{otherwise} \end{cases} \quad \begin{cases} \tilde{g}(a) = \tilde{g}(-b) = e(1/8) \\ \tilde{g}(-a) = \tilde{g}(b) = e(-1/8) \\ \tilde{g}(r) = 0 \quad \text{otherwise.} \end{cases}$$

Observe that, since a, b are not L -coincidental, there is no $i \in [5]$ with $A_i = \{\{a, -a, b, -b\}\}$. Thus, $\sigma_{A_i}(h) = \sigma_{A_i}(g)\sigma_{A_i}(\tilde{g}) \neq 0$ if and only if A_i equals one of $\pm S_1, \pm S_2, \pm S_3, \pm S_4$, defined by

$$S_1 = \{\{a, a, b, b\}\}, \quad S_2 = \{\{a, -a, b, b\}\}, \quad S_3 = \{\{a, a, b, -b\}\}, \quad S_4 = \{\{a, a, -b, -b\}\}.$$

One can check that

$$\sigma_{S_1}(g) = g(a)^2g(b)^2 + g(-a)^2g(-b)^2 = -2, \quad \sigma_{S_1}(\tilde{g}) = \tilde{g}(a)^2\tilde{g}(b)^2 + \tilde{g}(-a)^2\tilde{g}(-b)^2 = 2,$$

and similarly

$$\begin{array}{ccc} \sigma_{S_2}(g) = 0 & \sigma_{S_3}(g) = 0 & \sigma_{S_4}(g) = 2 \\ \sigma_{S_2}(\tilde{g}) = 0 & \sigma_{S_3}(\tilde{g}) = 0 & \sigma_{S_4}(\tilde{g}) = -2. \end{array}$$

Therefore, we have $\sigma_{A_i}(h) \leq 0$ for all $i \in [5]$. Since $A_{i_0} = S_1$ with $\sigma_{A_{i_0}}(h) = \sigma_{A_{i_0}}(g)\sigma_{A_{i_0}}(\tilde{g}) = -4$, we get that $\sum_{i=1}^5 \sigma_{A_i}(h) < 0$.

Otherwise, suppose there exists $i_0 \in E$ with $A_{i_0} = \{\{a, a, b, -b\}\}$ with $|a| \neq |b|$. In this case, consider the joined Fourier template $h : \mathbb{Z}^2 \rightarrow \mathbb{C}$ defined by $h(r_1, r_2) = g(r_1)\tilde{g}(r_2)$, where $g, \tilde{g} : \mathbb{Z} \rightarrow \mathbb{C}$ are 1-dimensional Fourier templates given by

$$\begin{cases} g(a) = g(-a) = 1 \\ g(b) = i, \quad g(-b) = -i \\ g(r) = 0 \quad \text{otherwise} \end{cases} \quad \begin{cases} \tilde{g}(b) = \tilde{g}(-b) = 1 \\ \tilde{g}(a) = i, \quad \tilde{g}(-a) = -i \\ \tilde{g}(r) = 0 \quad \text{otherwise.} \end{cases}$$

Again, since a, b are not L -coincidental, there is no $i \in [5]$ with $A_i = \{\{a, -a, b, -b\}\}$. Thus, $\sigma_{A_i}(h) = \sigma_{A_i}(g)\sigma_{A_i}(\tilde{g}) \neq 0$ if and only if A_i is one of $\pm S_2, \pm S_3$ defined above. One can check that

$$\sigma_{S_2}(g) = 2 \quad \sigma_{S_3}(g) = -2 \quad \sigma_{S_2}(\tilde{g}) = -2 \quad \sigma_{S_3}(\tilde{g}) = 2.$$

Therefore, we have $\sigma_{A_i}(h) \leq 0$ for all $i \in [5]$. Since $A_{i_0} = S_3$ with $\sigma_{A_{i_0}}(h) = \sigma_{A_{i_0}}(g)\sigma_{A_{i_0}}(\tilde{g}) = -4$, we get that $\sum_{i=1}^5 \sigma_{A_i}(h) < 0$.

Otherwise, suppose there exists $i_0 \in E$ with $A_{i_0} = \{\{a, a, b, c\}\}$ or $\{\{a, -a, b, c\}\}$, $|a|, |b|, |c|$ distinct, and a is not L -coincidental with b or c . In this case, the Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ defined by

$$\begin{cases} g(a) = g(-a) = C \\ g(b) = g(-b) = 1 \\ g(c) = g(-c) = -1 \\ g(r) = 0 \quad \text{otherwise} \end{cases}$$

has $\sum_{i=1}^5 \sigma_{A_i}(g) = -2C^2 + O(C)$, which is negative for $C > 0$ sufficiently large.

Finally, suppose for all $i \in E$ and all $a \in A_i$ not L -coincidental with any element in A_i , we have $|A_i \cap \{a, -a\}| = 1$. In this case, pick any $i_0 \in E$ and suppose $A_{i_0} = \{\{a_0, b_0, c_0, d_0\}\}$, such that a_0 is not L -coincidental with any of b_0, c_0, d_0 . Consider the Fourier template $g : \mathbb{Z} \rightarrow \mathbb{C}$ defined by

$$\begin{cases} g(a_0) = g(-a_0) = -C \\ g(b_0) = g(-b_0) = g(c_0) = g(-c_0) = g(d_0) = g(-d_0) = 1 \\ g(r) = 0 \quad \text{otherwise.} \end{cases}$$

Then we have $\sigma_{A_{i_0}}(g) = -2C$. Moreover, for all $i \in [5] \setminus \{i_0\}$, we either have $\sigma_{A_i}(g) = -2C$ or have $|\sigma_{A_i}(g)| = O(1)$. Hence for sufficiently large $C > 0$, we have $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$. \square

B.2. Case B. We first prove Lemma 5.14. To do so, we introduce the notion of ‘‘periodic sum’’ for h_{A_i} on the multiplicative grid G_L .

Definition B.1. Suppose $h : \pm G_L \rightarrow \mathbb{C}$ is function such that h_{A_1}, \dots, h_{A_5} are \vec{u} -periodic (recall Definition 5.12). Define

$$\sigma(h_{A_i}, \vec{u}) = \sum_{\substack{0 \leq d_1 \leq u_1 - 1 \\ \vdots \\ 0 \leq d_\ell \leq u_\ell - 1}} h_{A_i}(p_1^{d_1} \cdots p_\ell^{d_\ell} \cdot r).$$

Proof of Lemma 5.14. For $D > 0$, define $G_D = \{p_1^{d_1} \cdots p_\ell^{d_\ell} : d_1, \dots, d_\ell \in \{0, 1, \dots, D\}\}$ to be a finite truncation of G_L , and $g : \pm G_D \rightarrow \mathbb{C}$ by $g(r) = h(r)$. Observe that, if $r \in \pm G_D$ satisfies

$$\{ap_1^{d_1} \cdots p_\ell^{d_\ell} r : a \in A_1 \cup \cdots \cup A_5, 0 \leq d_1 < u_1, \dots, 0 \leq d_\ell < u_\ell\} \subseteq \pm G_D, \quad (*)$$

then we will have

$$\sum_{0 \leq d_1 < u_1} \cdots \sum_{0 \leq d_\ell < u_\ell} g_{A_i}(p_1^{d_1} \cdots p_\ell^{d_\ell} r) = \sigma_{A_i}(h, \vec{u}).$$

As $D \rightarrow \infty$, we have

$$\frac{|\{r \in \mathbb{Z} : g_{A_i} \neq 0 \text{ for some } i \in [5]\}|}{|\{r \in \pm G_D : r \text{ satisfies } (*)\}|} \rightarrow 1.$$

This captures the idea that the fraction of ‘‘boundary terms’’ become negligible and the main contribution comes from the internal portion of the grid. That is, we have

$$1 = \lim_{D \rightarrow \infty} \frac{\sum_{i=1}^5 \sigma_{A_i}(g)}{\sum_{i=1}^5 \sum_{r \in \pm G_D \text{ satisfies } (*)} g_{A_i}(r)} = \lim_{D \rightarrow \infty} \frac{\sum_{i=1}^5 \sigma_{A_i}(g)}{\sum_{i=1}^5 2\text{Re}(\sigma_{A_i}(h, \vec{u})) \cdot D^\ell / u_1 \cdots u_\ell}.$$

Therefore, given $\sum_{i=1}^5 \text{Re}(\sigma_{A_i}(h, \vec{u})) < 0$, we know that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$ for D sufficiently large. \square

Proof of Fact 5.24. For $\gamma_1, \gamma_2 \in \mathbb{Q} \setminus \{0\}$, we wish to find $t \in \mathbb{R}$ such that $1 + \cos(\gamma_1 t) + \cos(\gamma_2 t) < 0$. Without loss of generality, suppose $\gamma_1, \gamma_2 > 0$. The statement clearly holds for $\gamma_1 = \gamma_2$, so suppose $\gamma_1 < \gamma_2$.

If $\gamma_1 > \gamma_2/2$, we can take $t = \pi/\gamma_2$, so that $\gamma_2 t = \pi$ and $\pi/2 < \gamma_1 t < \pi$.

If $\gamma_2/3 \leq \gamma_1 < \gamma_2/2$, we can take $t = \frac{8\pi}{3\gamma_2}$, so that $\gamma_2 t = \frac{8\pi}{3}$ and $\frac{8\pi}{9} \leq \gamma_1 t < \frac{4\pi}{3}$.

If $\gamma_1 < \gamma_2/3$, then $(\frac{2\pi}{\gamma_1} \cdot \frac{2\pi}{3}, \frac{\gamma_2}{\gamma_1} \cdot \frac{4\pi}{3})$ has length $> 2\pi$, so we can find t such that $\gamma_1 t, \gamma_2 t \in (\frac{2\pi}{3}, \frac{4\pi}{3})$.

Finally, if $\gamma_2 = 2\gamma_1$, we can take $t = 1.4\pi/\gamma_1$, so that $1 + \cos(1.4\pi) + \cos(2.8\pi) < 0$. \square

Proof of Fact 5.18. Without loss of generality, assume $0 < \gamma_1 \leq \gamma_2 \leq \gamma_3 \leq \gamma_4$. Suppose for the sake of contradiction that $f(t) = 1 + \cos(\gamma_1 t) + \cos(\gamma_2 t) + \cos(\gamma_3 t) + \cos(\gamma_4 t) \geq 0$ for all $t \in \mathbb{R}$. In particular, we have that $f(t_k) \geq 0$ for all $k \in \mathbb{Z}$, with $t_k := \frac{(2k+1)\pi}{\gamma_4}$. This choice of t_k ensures that $\cos(\gamma_4 t_k) = -1$, so we have $f(t_k) = \cos(\gamma_1 t_k) + \cos(\gamma_2 t_k) + \cos(\gamma_3 t_k) \geq 0$.

Observe that for any $m > 0$, we have

$$\sum_{k=0}^{\gamma_4-1} \cos(mt_k) = \sum_{k=0}^{\gamma_4-1} \cos\left(\frac{m\pi}{\gamma_4} + \frac{2km\pi}{\gamma_4}\right) = \begin{cases} 0 & \text{if } \gamma_4 \nmid m \\ (-1)^{m/\gamma_4} & \text{if } \gamma_4 \mid m. \end{cases}$$

By the maximality of γ_4 and non-negativity of $f(\cdot)$, it follows that

$$0 \leq \sum_{k=0}^{\gamma_4-1} f(t_k) = \sum_{j=1}^3 \sum_{k=0}^{\gamma_4-1} \cos(\gamma_j t_k) = -\gamma_4 |\{i \in [3] : \gamma_i = \gamma_4\}| \leq 0.$$

Consequently equality holds in the above chain of inequalities, which implies that we have:

- $f(t_k) = 0$ for all $0 \leq k \leq \gamma_4 - 1$,
- $0 < \gamma_1, \gamma_2, \gamma_3 < \gamma_4$.

Thus, for all $i, j \in [3]$, we have $|\gamma_i - \gamma_j| < \gamma_4$ and $0 < \gamma_i + \gamma_j < 2\gamma_4$. This gives

$$\begin{aligned} 0 &= \sum_{k=0}^{\gamma_4-1} f(t_k)^2 = \sum_{i,j=1}^3 \sum_{k=0}^{\gamma_4-1} \cos(\gamma_i t_k) \cos(\gamma_j t_k) = \frac{1}{2} \sum_{i,j=1}^3 \sum_{k=0}^{\gamma_4-1} (\cos((\gamma_i + \gamma_j)t_k) + \cos((\gamma_i - \gamma_j)t_k)) \\ &= \frac{\gamma_4}{2} (|\{(i, j) : i, j \in [3], \gamma_i = \gamma_j\}| - |\{(i, j) : i, j \in [3], \gamma_i + \gamma_j = \gamma_4\}|). \end{aligned}$$

Since $|\{(i, j) : i, j \in [3], \gamma_i = \gamma_j\}| \geq 3$, we must have $|\{(i, j) : i, j \in [3], \gamma_i + \gamma_j = \gamma_4\}| \geq 3$, which implies that $\gamma_1 + \gamma_3 = 2\gamma_2 = \gamma_4$. In particular, we have

$$\cos(\gamma_3\pi/\gamma_4) = \cos(\pi - \gamma_1\pi/\gamma_4) = -\cos(\gamma_1\pi/\gamma_4), \quad \cos(\gamma_2\pi/\gamma_4) = \cos(\pi/2) = 0.$$

Thus, at $t_0 = \pi/\gamma_4$, we have

- $f(t_0) = \cos(\gamma_1 t_0) + \cos(\gamma_2 t_0) + \cos(\gamma_3 t_0) = 0$, and
- $f'(t_0) = -\gamma_1 \sin(\gamma_1 t_0) - \gamma_2 \sin(\gamma_2 t_0) - \gamma_3 \sin(\gamma_3 t_0) < 0$ since $0 < \gamma_1 t_0, \gamma_2 t_0, \gamma_3 t_0 < \gamma_4 t_0 = \pi$.

It follows that $f(t_0 + \epsilon) < 0$ for sufficiently small $\epsilon > 0$, which is a contradiction as desired. \square

In the remainder of this section we prove Proposition 5.22. In order to handle the more intricate cases when some of the A_i are cancelling, we need to introduce more notation.

Definition B.2. For primes $p \neq \tilde{p}$ and multiset A , define $V_{p, \tilde{p}}(A) = \{(v_p(a), v_{\tilde{p}}(a)) : a \in A\}$.

Example B.3. Suppose $A = \{-1, 2, 6, -12\}$. Then

- $V_2(A) = \{(0, 1, 1, 2)\}$, $V_3(A) = \{(0, 0, 1, 1)\}$,
- $V_{2,3}(A) = \{(0, 0), (1, 0), (1, 1), (2, 1)\}$.

Observation B.4. Consider $a, a' \in \mathbb{Z} \setminus \{0\}$ and prime number p .

If $v_p(a) \neq v_p(a')$, then $v_p(a + a') = v_p(a - a') = \min\{v_p(a), v_p(a')\}$.

If $v_p(a) = v_p(a')$, then $v_p(a + a'), v_p(a - a') \geq v_p(a)$.

Recall that in Case B, we assume (Setup 5.11) that L is of the form

$$\begin{aligned} L_5 \mathbf{x} &= x_1 + x_2 - x_3 - x_4 & A_5 &= \{(1, 1, -1, -1)\} \\ L_4 \mathbf{x} &= ax_1 + bx_2 + cx_3 + dx_5 & A_4 &= \{(a, b, c, d)\} \\ L_3 \mathbf{x} &= (a + c)x_1 + (b + c)x_2 - cx_4 + dx_5 & A_3 &= \{(a + c, b + c, -c, d)\} \\ L_2 \mathbf{x} &= (a - b)x_1 + (b + c)x_3 + bx_4 + dx_5 & A_2 &= \{(a - b, b + c, b, d)\} \\ L_1 \mathbf{x} &= (b - a)x_3 + (a + c)x_3 + ax_4 + dx_5 & A_1 &= \{(b - a, a + c, a, d)\} \end{aligned}$$

with $a, b, c, d \in \mathbb{Z} \setminus \{0\}$, and none of L_1, \dots, L_4 is common. By replacing a, b, c, d by their negations if necessary, we may suppose that at least two of a, b, c, d are positive.

Lemma B.5. *Suppose L is an irredundant 2×5 linear system that falls under Case B in Corollary 5.3. Assume Setup 5.11. Then there cannot be three cancelling sets among A_1, \dots, A_4 .*

Proof. By contradiction, suppose three among these sets are cancelling. Up to isomorphism, we can suppose A_2, A_3, A_4 are cancelling and $a > 0$. We then have four different cases dependings on the coefficients in L_1, \dots, L_5 :

- (1) $a > 0, b > 0, c < 0, d < 0, a + c > 0, b + c < 0, a - b > 0,$
 $ab = cd, -c(a + c) = (b + c)d, b(a - b) = d(b + c);$
- (2) $a > 0, b > 0, c < 0, d < 0, a + c < 0, b + c > 0, a - b < 0,$
 $ab = cd, -c(b + c) = (a + c)d, b(b + c) = d(a - b);$
- (3) $a > 0, c > 0, b < 0, d < 0, a + c > 0, b + c > 0, a - b > 0,$
 $ac = bd, (a + c)(b + c) = -cd, (a - b)(b + c) = bd;$
- (4) $a > 0, d > 0, b < 0, c < 0, a + c < 0, b + c < 0, a - b > 0,$
 $ad = bc, (a + c)(b + c) = -cd, d(a - b) = b(b + c).$

One can then check by a direct computation that none of these cases has real solutions. \square

Corollary B.6. *Suppose L is an irredundant 2×5 linear system that falls under Case B in Corollary 5.3. Assume Setup 5.11. Then up to isomorphism, at least one of the following occurs:*

- (1) A_4 is cancelling and $V_p(A_4) = \{0, v_1, v_2, v_1 + v_2\}$ with $0 < v_1 < v_2$ for some prime p .
- (2) A_4 is cancelling and $V_p(A_4) = \{0, v, v, 2v\}$ with $v > 0$ for some prime p .
- (3) A_4 is cancelling and $V_{p, \tilde{p}}(A_4) = \{(0, 0), (v_1, 0), (0, v_2), (v_1, v_2)\}$ with $v_1, v_2 > 0$ for some primes $p \neq \tilde{p}$. Moreover, none of A_1, A_2, A_3 is cancelling.
- (4) A_4 is cancelling and $V_{p, \tilde{p}}(A_4) = \{(0, 0), (v_1, 0), (0, v_2), (v_1, v_2)\}$ with $v_1, v_2 > 0$ for some primes $p \neq \tilde{p}$. Moreover, there is exactly one $i \in [3]$ such that the following two conditions hold:
 - (i) A_i is cancelling, and
 - (ii) $V_{p', \tilde{p}'}(A_i) = \{(0, 0), (w_1, 0), (0, w_2), (w_1, w_2)\}$ with $w_1, w_2 > 0$ for some primes $p' \neq \tilde{p}'$.

We now state a reduction step which says that, when $h : \pm G_L \rightarrow \mathbb{C}$ satisfies certain conditions, we can combine it with constructions in Observation 5.16 and Proposition 5.17 appropriately, to obtain Fourier template g with $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.

Definition B.7. Let A be a multiset of size 4. We say that A is (P2) if it contains two positive and two negative elements, and is (P3) if it contains three positive and one negative elements.

Proposition B.8. *Suppose L is an irredundant 2×5 linear system that falls under Case B in Corollary 5.3. Assume Setup 5.11. Moreover, suppose A_4 is cancelling, and one of the following holds:*

- (1) One of A_1, A_2, A_3 is cancelling, and there exists $h : \pm G_L \rightarrow \mathbb{C}$ such that h_{A_1}, \dots, h_{A_5} are \vec{u} -periodic, with $\sum_{i \in [5], A_i \text{ cancelling}} \sigma(h_{A_i}, \vec{u}) < 0$.
- (2) None of A_1, A_2, A_3 is cancelling, and there exists $h : \pm G_L \rightarrow \mathbb{C}$ such that h_{A_1}, \dots, h_{A_5} are \vec{u} -periodic, with $\sigma(h_{A_5}, \vec{u}) + \sigma(h_{A_4}, \vec{u}) = 0$ and $|\{i \in [3] : \text{Re}(\sigma(h_{A_i}, \vec{u})) \neq 0\}| \in \{1, 3\}$.

Then there exists a Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.

Remark B.9. The caveat here is that, if for some $h : \pm G_L \rightarrow \mathbb{C}$ we have $\sigma(h_{A_5}, \vec{u}) + \sigma(h_{A_4}, \vec{u}) = 0$ and $|\{i \in [3] : \text{Re}(\sigma(h_{A_i}, \vec{u})) \neq 0\}| = 2$, we might not be able to produce, based on this, a Fourier template g with $\text{Re} \sum_{i=1}^5 \sigma(h_{A_i}, \vec{u})$ strictly less than 0. (For example, $1 - 1 + xe^{i\gamma t} - xe^{i\gamma t} + 0 = 0$ no matter what value t takes.)

Proof of Proposition B.8. We work under Setup 5.11. One can check that, since A_4 is cancelling, either A_1, \dots, A_4 are all (P2), or two of them are (P2) and the other two are (P3).

Recall that the function h_2 defined in Observation 5.16 satisfies

$$(h_2)_{A_i} = \begin{cases} 1 & A_i \text{ is (P2)} \\ -1 & A_i \text{ is (P3)}. \end{cases}$$

The function h_3 (with $\theta_1, \dots, \theta_\ell, t$ to be determined) defined in Proposition 5.17 satisfies

$$(h_3)_{A_i} = \begin{cases} 1 & A_i \text{ is cancelling} \\ e^{i\gamma t}, \gamma \neq 0 & A_i \text{ is (P2) but not cancelling.} \end{cases}$$

Suppose condition (1) in Proposition B.8 occurs. Then we can pick $\theta_1, \dots, \theta_\ell, t$ so that the corresponding h_3 has

$$\operatorname{Re} \sum_{\substack{A_i \text{ is (P2) but} \\ \text{not cancelling}}} \sigma((hh_3)_{A_i}, \vec{u}) \leq 0,$$

which then gives

$$\operatorname{Re} \sum_{A_i \text{ is (P2)}} \sigma((hh_3)_{A_i}, \vec{u}) < 0.$$

Let g be the finite truncation of hh_3 (recall Lemma 5.14) and g_2 be the finite truncation of h_2 . Then, by Proposition 3.6, we have either $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$ or $\sum_{i=1}^5 \sigma_{A_i}(gg_2) < 0$.

Suppose condition (2) in Proposition B.8 occurs. If exactly one $i_0 \in [3]$ has $\operatorname{Re}(\sigma(h_{A_{i_0}}, \vec{u})) \neq 0$, and this A_{i_0} is (P3), then letting g be the finite truncation of h and g_2 be the finite truncation of h_2 , we have either $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$ or $\sum_{i=1}^5 \sigma_{A_i}(gg_2) < 0$.

Otherwise, we have

$$|\{i \in [3] : \operatorname{Re}(\sigma(h_{A_i}, \vec{u})) \neq 0, A_i \text{ is (P2)}\}| \in \{1, 3\},$$

so we can pick $\theta_1, \dots, \theta_\ell, t$ such that

$$\operatorname{Re} \sum_{A_i \text{ is (P2)}} \sigma((hh_3)_{A_i}, \vec{u}) < 0.$$

Again, letting g be the finite truncation of hh_3 and g_2 be the finite truncation of h_2 , we have either $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$ or $\sum_{i=1}^5 \sigma_{A_i}(gg_2) < 0$. \square

We will show in Propositions B.10 to B.13 that, in each of the four cases listed in Corollary B.6, there exists some Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.

Proposition B.10. *Suppose L is an irredundant 2×5 linear system that falls under Case B in Corollary 5.3. Assume Setup 5.11. Moreover, suppose A_4 is cancelling and*

$$V_{p_1}(A_4) = \{\{0, v_1, v_2, v_1 + v_2\}\}, \quad 0 < v_1 < v_2 < v_1 + v_2.$$

Then there exists a Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.

Proof. Suppose $\gcd(v_1, v_2) = 1$. By Observation B.4 and the fact that $V_{p_1}(A_4) = \{\{0, v_1, v_2, v_1 + v_2\}\}$, one can check that $\{V_{p_1}(A_1), V_{p_1}(A_2), V_{p_1}(A_3)\}$ must be one of the following:

- $\{\{\{0, 0, 0, v_1 + v_2\}\}, \{\{0, v_1, v_1, v_1 + v_2\}\}, \{\{0, v_1, v_2, v_1 + v_2\}\}\}$,
- $\{\{\{0, 0, 0, v_1\}\}, \{\{0, v_2, v_2, v_1\}\}, \{\{0, v_2, v_1 + v_2, v_1\}\}\}$,
- $\{\{\{0, 0, 0, v_2\}\}, \{\{0, v_1, v_1, v_2\}\}, \{\{0, v_1, v_2, v_1 + v_2\}\}\}$,
- $\{\{\{0, v_1, v_1, v_1\}\}, \{\{0, v_1, v_2, v_2\}\}, \{\{v_1, v_2, v_1 + v_2, 0\}\}\}$.

Let

$$\mathcal{V} = \{\{\{0, 0, 0, v_1\}\}, \{\{0, 0, 0, v_2\}\}, \{\{0, 0, 0, v_1 + v_2\}\}, \{\{0, v_1, v_1, v_1\}\}, \\ \{\{0, v_1, v_1, v_2\}\}, \{\{0, v_1, v_1, v_1 + v_2\}\}, \{\{0, v_1, v_2, v_2\}\}, \{\{0, v_1, v_2, v_1 + v_2\}\}\}$$

denote the collection of all 4-multisets that appeared in the above possibilities. Also, let

$$W = V_{p_1}(A_4) = \{\{0, v_1, v_2, v_1 + v_2\}\}.$$

Suppose first that one of v_1, v_2 is even. Call this even number u_e . Define $\phi : \{0, 1, \dots\} \rightarrow \mathbb{R}$ by

$$\begin{cases} \phi(v) = -1 & v \equiv 0, 2, \dots, u_e - 2 \pmod{2u_e} \\ \phi(v) = 1 & \text{otherwise} \end{cases}$$

and $h = \phi \circ v_{p_1} : \pm G_L \rightarrow \mathbb{R}$. Let $\vec{u} = (2u_e, 1, \dots, 1)$. Then h_{A_1}, \dots, h_{A_5} are \vec{u} -periodic, with

$$\sigma(h_{A_i}, \vec{u}) = \begin{cases} 2u_e & A_i = A_5 \\ -2u_e & V_{p_1}(A_i) = W \\ 0 & V_{p_1}(A_i) \in \mathcal{V} \setminus \{W\}. \end{cases}$$

Since exactly two elements $i \in [4]$ have $V_{p_1}(A_i) = W$, we get that $\sum_{i=1}^5 \sigma(h_{A_i}, \vec{u}) = -2u_e < 0$, so Lemma 5.14 applies.

Suppose v_1, v_2 are both odd. In this case, we build $h = \phi \circ v_{p_1} : \pm G_L \rightarrow \mathbb{C}$ that satisfies Proposition B.8. If $v_1 + v_2$ is a multiple of 4, define $u_e = v_1 + v_2$ and $w_e = v_2 - v_1$; if $v_1 - v_2$ is a multiple of 4, define $u_e = v_2 - v_1$ and $w_e = v_1 + v_2$.

Note that w_e is a multiple of 2 but not 4, and $\gcd(w_e/2, u_e/2) = 1$. This means that w_e has order $u_e/2$ in \mathbb{Z}_{u_e} , so we can define $\phi : \{0, 1, \dots\} \rightarrow \mathbb{R}$ by

$$\begin{cases} \phi(v) = -1 & v \equiv 0, 1, 2w_e, 2w_e + 1, \dots, -2w_e, -2w_e + 1 \pmod{u_e} \\ \phi(v) = 1 & \text{otherwise.} \end{cases}$$

Let $\vec{u} = (u_e, 1, \dots, 1)$. Then h_{A_1}, \dots, h_{A_5} are \vec{u} -periodic, with

$$\sigma(h_{A_i}, \vec{u}) = \begin{cases} 2u_e & A_i = A_5 \\ -2u_e & V_{p_1}(A_i) = \{\{0, v_1, v_2, v_1 + v_2\}\} \\ 2u_e \text{ or } -2u_e & V_{p_1}(A_i) \in \{\{\{0, 0, 0, v_1 + v_2\}\}, \{\{0, v_1, v_1, v_1 + v_2\}\}\} \\ 0 & \text{otherwise.} \end{cases}$$

This implies the condition of Proposition B.8, so we are done.

In general, for $\gcd(v_1, v_2) = w > 1$, up to replacing $\phi(v)$ by $\phi(\lfloor v/w \rfloor)$ and the period $\vec{u} = (u_1, 1, \dots, 1)$ by $(u_1 w_1, 1, \dots, 1)$, we can still use the previous constructions. \square

Proposition B.11. *Suppose L is an irredundant 2×5 linear system that falls under Case B in Corollary 5.3. Assume Setup 5.11. Moreover, suppose A_4 is cancelling and*

$$V_{p_1}(A_4) = \{\{0, v, v, 2v\}\}, \quad v > 0.$$

Then there exists a Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.

Proof. We first assume that $v = 1$, i.e., $V_{p_1}(A_4) = \{\{0, 1, 1, 2\}\}$. Using Observation B.4, one can check that $\{V_{p_1}(A_1), V_{p_1}(A_2), V_{p_1}(A_3)\}$ must be one of the following form:

- $\{\{\{0, 0, 0, 2\}\}, \{\{0, 1, x, 2\}\}, \{\{0, 1, x, 2\}\}\}$ for some $x \geq 1$,
- $\{\{\{0, 0, 0, 1\}\}, \{\{0, 1, 1, 1\}\}, \{\{0, 1, 2, 1\}\}\}$,
- $\{\{\{1, x, 2, 0\}\}, \{\{1, x, 2, 0\}\}, \{\{1, 1, 2, 0\}\}\}$ for some $x \geq 1$.

Suppose A_i is cancelling for some $i \in [3]$. Then we have $V_{p_1}(A_i) = \{\{0, 1, x, 2\}\}$ with $x = 1$ or 3 . If $x = 3$, then we can apply Proposition B.10 to find the desired g . If $x = 1$, define $h : \pm G_L \rightarrow \mathbb{C}$ by

$$\begin{cases} h(r) = i & v_{p_1}(r) \text{ odd, } r > 0 \\ h(r) = -i & v_{p_1}(r) \text{ odd, } r < 0 \\ h(r) = 1 & v_{p_1}(r) \text{ even} \end{cases}$$

so that for every $i \in [5]$, h_{A_i} is \vec{u} -periodic with $\vec{u} = (2, 1, \dots, 1)$. Moreover, we have

$$\sigma(h_{A_i}, \vec{u}) = \begin{cases} 2 & A_i = A_5 \\ -2 & i \in [4], A_i \text{ is cancelling.} \end{cases}$$

This gives $\sum_{A_i \text{ cancelling}} \sigma(h_{A_i}, \vec{u}) \leq -2 < 0$, so Proposition B.8 applies.

Now suppose that none of A_1, A_2, A_3 is cancelling. In this case, define $h : \pm G_L \rightarrow \mathbb{C}$ by

$$\begin{cases} h(r) = e(1/8) & v_{p_1}(r) \text{ odd, } r > 0 \text{ or } v_{p_1}(r) \text{ even, } r < 0 \\ h(r) = e(-1/8) & v_{p_1}(r) \text{ odd, } r < 0 \text{ or } v_{p_1}(r) \text{ even, } r > 0. \end{cases}$$

One can check that

- $h_{A_5} = 1$ and $h_{A_4} = -1$ on $\pm G_L$;
- if A_i is (P3), and $V_{p_1}(A_i)$ contains an odd number of even elements, then h_{A_i} is a constant function that is either 1 or -1 ;
- if A_i is (P2), and $V_{p_1}(A_i)$ contains an even number of even elements, then h_{A_i} is a constant function that is either 1 or -1 ;
- otherwise, $\text{Re}(\sigma(h_{A_i}, \vec{u})) = 0$.

This implies the condition of Proposition B.8(2), so we are done.

In general, suppose $v > 1$. If all elements in $V_{p_1}(A_1) \cup V_{p_1}(A_2) \cup V_{p_1}(A_3)$ are divisible by v , then we can use the above constructions up to replacing $v_{p_1}(r)$ by $\lfloor v_{p_1}(r)/v \rfloor$. Thus, it suffices to assume that $\{V_{p_1}(A_1), V_{p_1}(A_2), V_{p_1}(A_3)\}$ equals $\{\{(0, 0, 0, 2v)\}, \{(0, v, x, 2v)\}, \{(0, v, x, 2v)\}\}$ or $\{\{(v, x, 2v, 0)\}, \{(v, x, 2v, 0)\}, \{(v, v, 2v, 0)\}\}$ with $v \nmid x$. Define $h : \pm G \rightarrow \mathbb{C}$ by

$$\begin{cases} h(r) = 1 & \lfloor v_{p_1}(r)/(2v) \rfloor \text{ even} \\ h(r) = -1 & \lfloor v_{p_1}(r)/(2v) \rfloor \text{ odd.} \end{cases}$$

In this case, one can check that

- $h_{A_5} = 1$ and $h_{A_4} = -1$ on $\pm G_L$;
- $h_{A_1}, h_{A_2}, h_{A_3}$ are \vec{u} -periodic with $\vec{u} = (4v, 1, \dots, 1)$;
- $\sigma(h_{A_1}, \vec{u}), \sigma(h_{A_2}, \vec{u}), \sigma(h_{A_3}, \vec{u})$ are all nonzero.

Again, this implies the condition of Proposition B.8(2), so we are done. \square

Proposition B.12. *Suppose L is an irredundant 2×5 linear system that falls under Case B in Corollary 5.3. Assume Setup 5.11. Moreover, suppose A_4 is cancelling, none of A_1, A_2, A_3 is cancelling, and*

$$V_{p_1, p_2}(A_4) = \{(0, 0), (v_1, 0), (0, v_2), (v_1, v_2)\}, \quad v_1, v_2 > 0.$$

Then there exists a Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.

Proof. Using Observation B.4, one can check that $\{V_{p_1, p_2}(A_1), V_{p_1, p_2}(A_2), V_{p_1, p_2}(A_3)\}$ must be one of the following form:

- $\{(v_1, 0), (0, 0), (x, 0), (0, 0)\}, \{(0, 0), (0, v_2), (0, y), (0, 0)\}, \{(x, 0), (0, y), (v_1, v_2), (0, 0)\}$
- $\{(0, 0), (x, 0), (0, y), (v_1, v_2)\}, \{(x, 0), (0, v_2), (0, 0), (v_1, v_2)\}, \{(0, y), (0, 0), (v_1, 0), (v_1, v_2)\}$
- $\{(0, 0), (x, 0), (0, 0), (v_1, 0)\}, \{(x, 0), (0, v_2), (0, y), (v_1, 0)\}, \{(0, 0), (0, y), (v_1, v_2), (v_1, 0)\}$
- $\{(0, 0), (0, y), (0, 0), (0, v_2)\}, \{(0, y), (v_1, 0), (x, 0), (0, v_2)\}, \{(0, 0), (x, 0), (v_1, v_2), (0, v_2)\}$

where $x, y \geq 0$.

The proof idea is similar to Proposition B.11. Suppose first that $v_1 \mid x$ and $v_2 \mid y$. In this case, define $h : \pm G_L \rightarrow \mathbb{C}$ by

$$\begin{cases} h(r) = e(1/8) & \begin{cases} \lfloor v_{p_1}(r)/v_1 \rfloor \equiv \lfloor v_{p_2}(r)/v_2 \rfloor \pmod{2}, r > 0 \\ \text{or } \lfloor v_{p_1}(r)/v_1 \rfloor \not\equiv \lfloor v_{p_2}(r)/v_2 \rfloor \pmod{2}, r < 0 \end{cases} \\ h(r) = e(-1/8) & \begin{cases} \lfloor v_{p_1}(r)/v_1 \rfloor \not\equiv \lfloor v_{p_2}(r)/v_2 \rfloor \pmod{2}, r > 0 \\ \text{or } \lfloor v_{p_1}(r)/v_1 \rfloor \equiv \lfloor v_{p_2}(r)/v_2 \rfloor \pmod{2}, r < 0. \end{cases} \end{cases}$$

In this case, one can check that

- $h_{A_5} = 1$ and $h_{A_4} = -1$;
- if A_i is cancelling for some $i \in [3]$, then $A_i = A_4$ and $h_{A_i} = h_{A_4} = -1$;
- if A_i is (P3) and $V_{p_1}(A_i)$ contains an odd number of elements in $\{(0, 0), (v_1, v_2)\} \pmod{(2v_1, 2v_2)}$, then h_{A_i} is a constant function that is either 1 or -1 ;
- if A_i is (P2) and $V_{p_1}(A_i)$ contains an even number of elements in $\{(0, 0), (v_1, v_2)\} \pmod{(2v_1, 2v_2)}$, then h_{A_i} is a constant function that is either 1 or -1 ;
- otherwise, $\text{Re}(\sigma(h_{A_i}, \vec{u})) = 0$.

Thus, in each of the above possibilities of $\{V_{p_1, p_2}(A_1), V_{p_1, p_2}(A_2), V_{p_1, p_2}(A_3)\}$, the condition of Proposition B.8 holds.

Suppose now that $2x \mid v_1$ and $2y \mid v_2$; moreover, either $x \equiv v_1/2 \pmod{v_1}$ or $y \equiv v_2/2 \pmod{v_2}$. In this case, define $h : \pm G_L \rightarrow \mathbb{C}$ by $h = h_1 h_2$ or $h = h_1 h_3$, where

$$\begin{cases} h_1(r) = 1 & \lfloor v_{p_1}(r)/v_1 \rfloor \equiv \lfloor v_{p_2}(r)/v_2 \rfloor \equiv 0 \pmod{2} \\ h_1(r) = -1 & \text{otherwise} \end{cases}$$

$$\begin{cases} h_2(r) = 1 & \lfloor v_{p_1}(r)/x \rfloor + \lfloor v_{p_2}(r)/y \rfloor \equiv 0 \pmod{2} \\ h_2(r) = i & \lfloor v_{p_1}(r)/x \rfloor + \lfloor v_{p_2}(r)/y \rfloor \equiv 1 \pmod{2}, r > 0 \\ h_2(r) = -i & \lfloor v_{p_1}(r)/x \rfloor + \lfloor v_{p_2}(r)/y \rfloor \equiv 1 \pmod{2}, r < 0 \end{cases}$$

$$\begin{cases} h_2(r) = 1 & \lfloor v_{p_1}(r)/x \rfloor + \lfloor v_{p_2}(r)/y \rfloor \equiv 1 \pmod{2} \\ h_2(r) = i & \lfloor v_{p_1}(r)/x \rfloor + \lfloor v_{p_2}(r)/y \rfloor \equiv 0 \pmod{2}, r > 0 \\ h_2(r) = -i & \lfloor v_{p_1}(r)/x \rfloor + \lfloor v_{p_2}(r)/y \rfloor \equiv 0 \pmod{2}, r < 0. \end{cases}$$

In this case, one can check that

- $h_{A_5} = 1$ and $h_{A_4} = -1$;
- By picking $h \in \{h_1 h_2, h_1 h_3\}$ appropriately, we have $|\{i \in [3] : \text{Re}(\sigma(h_{A_i}, \vec{u})) \neq 0\}| \in \{1, 3\}$.

Finally, suppose we have $2v_1 \nmid y$ or $2v_2 \nmid y$. Set

$$\begin{cases} h(r) = -1 & \lfloor v_{p_1}(r)/v_1 \rfloor, \lfloor v_{p_2}(r)/v_2 \rfloor \text{ both odd} \\ h(r) = 1 & \text{otherwise,} \end{cases}$$

so that h_{A_1}, \dots, h_{A_5} are \vec{u} -periodic with period $\vec{u} = (2v_1, 2v_2, 1, \dots, 1)$. Moreover, we have $\sigma(h_{A_5}, \vec{u}) = 4v_1 v_2$, $\sigma(h_{A_4}, \vec{u}) = -4v_1 v_2$, and $\sigma(h_{A_3}, \vec{u}), \sigma(h_{A_2}, \vec{u}), \sigma(h_{A_1}, \vec{u})$ are all nonzero. Thus the condition of Proposition B.8 always holds, so we are done. \square

Proposition B.13. *Suppose L is an irredundant 2×5 linear system that falls under Case B in Corollary 5.3. Assume Setup 5.11. Moreover, suppose*

- A_4 is cancelling and $V_{p_1, p_2}(A_4) = \{(0, 0), (v_1, 0), (0, v_2), (v_1, v_2)\}$ with $v_1, v_2 > 0$.
- A_{i_0} is cancelling for some $i_0 \in [3]$, and $V_{p_{j_1}, p_{j_2}}(A_i) = \{(0, 0), (w_1, 0), (0, w_2), (w_1, w_2)\}$ with $w_1, w_2 > 0$.
- A_i is not cancelling for all $i \in [3] \setminus \{i_0\}$.

Then there exists a Fourier template $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ such that $\sum_{i=1}^5 \sigma_{A_i}(g) < 0$.

Proof. Suppose first that $p_1 = p_{j_1}$ and $p_2 = p_{j_2}$. In this case, we must have $v_1 = w_1$ and $v_2 = w_2$. Consider $h : \pm G_L \rightarrow \mathbb{C}$ defined by

$$\begin{cases} h(r) = -1 & [v_{p_1}(r)/v_1] \equiv [v_{p_2}(r)/v_2] \equiv 0 \pmod{2} \\ h(r) = 1 & \text{otherwise} \end{cases}$$

so that $h_{A_4}, h_{A_{i_0}}$ are the constant -1 function on $\pm G_L$. This implies Proposition B.8(1).

Now suppose $\{p_1, p_2\} \neq \{p_{j_1}, p_{j_2}\}$. In this case, since A_4, A_{i_0} are cancelling, $V_{p_1, p_2}(A_{i_0})$ and $V_{p_{j_1}, p_{j_2}}(A_4)$ can be partitioned into cancelling pairs. Consider functions $h, h' : \pm G \rightarrow \mathbb{C}$ defined by

$$\begin{cases} h(r) = -1 & [v_{p_1}(r)/v_1] \equiv [v_{p_2}(r)/v_2] \equiv 0 \pmod{2} \\ h(r) = 1 & \text{otherwise,} \\ h'(r) = -1 & [v_{p_{j_1}}(r)/w_1] \equiv [v_{p_{j_2}}(r)/w_2] \equiv 0 \pmod{2} \\ h'(r) = 1 & \text{otherwise.} \end{cases}$$

We already know that h_{A_4} and $h'_{A_{i_0}}$ are the constant -1 function. Moreover, since $V_{p_1, p_2}(A_{i_0})$ and $V_{p_{j_1}, p_{j_2}}(A_4)$ can be partitioned into cancelling pairs, we additionally know that h'_{A_4} and $h_{A_{i_0}}$ are the constant 1 function. Therefore, $(hh')_{A_4}$ and $(hh')_{A_{i_0}}$ are the constant -1 function. Again, this implies the condition of Proposition B.8(1), so we are done. \square

B.3. Case D. In this subsection, we give the reduction to (5). We begin by recalling the setup. Recall that we had the following system L ,

$$\begin{aligned} L_5 &= x_1 - x_2 + \alpha x_3 - \alpha x_4 \\ L_4 &= x_1 + \beta x_2 - x_3 - \beta x_5 \\ L_3 &= (\alpha + 1)x_1 + (\alpha\beta - 1)x_2 - \alpha x_4 - \alpha\beta x_5 \\ L_2 &= (\beta + 1)x_1 + (\alpha\beta - 1)x_3 - \alpha\beta x_4 - \beta x_5 \\ L_1 &= (\beta + 1)x_2 + (-1 - \alpha)x_3 + \alpha x_4 - \beta x_5 \end{aligned}$$

with $\alpha, \beta \in \mathbb{Q}$ and $\alpha, \beta > 0$, and at least one of L_1, L_2, L_3 is common. We then have one of the following:

- (1) L_3 is common, which means that $(\alpha + 1)(\alpha\beta - 1) = \alpha \cdot \alpha\beta$;
- (2) L_2 is common, which means that $(\beta + 1)(\alpha\beta - 1) = \beta \cdot \alpha\beta$;
- (3) L_1 is common, which means that $\alpha(\beta + 1) = \beta(\alpha + 1)$.

Note that the first two cases are the same up to switching α and β . The first case implies that $\alpha\beta = \alpha + 1$, which gives

$$\begin{aligned} L_5 &= x_1 - x_2 + \alpha x_3 - \alpha x_4 \\ L_4 &= x_1 + \frac{\alpha + 1}{\alpha} \cdot x_2 - x_3 - \frac{\alpha + 1}{\alpha} \cdot x_5. \end{aligned}$$

The third case implies that $\alpha = \beta$, which gives

$$\begin{aligned} L_5 &= x_1 - x_2 + \alpha x_3 - \alpha x_4 \\ L_4 &= x_1 + \alpha x_2 - x_3 - \alpha x_5. \end{aligned}$$

From here, we can see that these two cases are the same up to sending $\alpha \mapsto \frac{\alpha+1}{\alpha}$. Therefore, we may restrict our attention to the third case.

B.4. Case E.

Proof that (9) is uncommon when $\lambda = 1$. Recall that the setup is as follows: the elements in $\{L_B : B \in \mathcal{C}(L)\}$ are

$$\begin{aligned} L_5 &= x_1 - x_2 + x_3 - x_4 & A_5 &= \{1, -1, 1, -1\} \\ L_1 &= ax_2 + (b-a)x_3 + ax_4 + cx_5 & A_1 &= \{b-a, a, a, c\} \\ L_3 &= (a-b)x_1 + bx_2 + bx_4 + cx_5 & A_3 &= \{a-b, b, b, c\}. \end{aligned}$$

Suppose first that $|a-b| \neq |c|$. Take prime p such that $v := |v_p(a-b) - v_p(c)| > 0$. Then the map $h : \pm G_L \rightarrow \mathbb{C}$ with

$$\begin{cases} h(r) = 1 & [v_p(r)/v] \equiv 0 \pmod{2} \\ h(r) = -1 & [v_p(r)/v] \equiv 1 \pmod{2} \end{cases}$$

gives $\operatorname{Re}(h_{A_1}) + \operatorname{Re}(h_{A_3}) + \operatorname{Re}(h_{A_5}) = 1 - 1 - 1 = -1$.

Now suppose $|a-b| = |c|$. Without loss of generality, let $b-a = c$, so we have $A_5 = \{1, 1, -1, -1\}$, $A_1 = \{a, a, c, c\}$, $A_3 = \{a+c, a+c, c, -c\}$. If a, c are of the same sign, then we can choose $z \in \mathbb{C}$ with $|z| = 1$ and $\operatorname{Re}(z^4 + z^2 + 1) < 0$, and set $h : \pm G_L \rightarrow \mathbb{C}$ by

$$\begin{cases} h(r) = z & r > 0 \\ h(r) = \bar{z} & r < 0, \end{cases}$$

so that $\operatorname{Re}(h_{A_5}), \operatorname{Re}(h_{A_1}), \operatorname{Re}(h_{A_3})$ are constant on $\pm G_L$, with $\operatorname{Re}(h_{A_5}) + \operatorname{Re}(h_{A_1}) + \operatorname{Re}(h_{A_3}) = \operatorname{Re}(z^4 + z^2 + 1) < 0$.

If a, c are of different signs, then there exists some prime p that divides exactly one of $|a|$ and $|c|$. Suppose $\{v_p(a), v_p(c)\} = \{v, 0\}$ with $v \in \mathbb{N}$. Again, we choose $z \in \mathbb{C}$ with $|z| = 1$ and $\operatorname{Re}(z^4 + z^2 + 1) < 0$. Set $h : \pm G_L \rightarrow \mathbb{C}$ by

$$h(r) = \begin{cases} z & [v_p(r)/v] \text{ is even, } r > 0 \\ \bar{z} & [v_p(r)/v] \text{ is odd, } r > 0 \\ \bar{z} & [v_p(r)/v] \text{ is even, } r < 0 \\ z & [v_p(r)/v] \text{ is odd, } r < 0. \end{cases}$$

One can check that $\operatorname{Re}(h_{A_5}), \operatorname{Re}(h_{A_1}), \operatorname{Re}(h_{A_3})$ are constant on $\pm G_L$, with $\operatorname{Re}(h_{A_5}) + \operatorname{Re}(h_{A_1}) + \operatorname{Re}(h_{A_3}) = \operatorname{Re}(z^4 + z^2 + 1) < 0$. \square

DONG: DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA 02138, USA.
Email address: ddong@math.harvard.edu

LI: DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS (CENTRE FOR MATHEMATICAL SCIENCES), UNIVERSITY OF CAMBRIDGE, CAMBRIDGE CB30WB, ENGLAND, UK.
Email address: aqli@stanford.edu

ZHAO: DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA.
Email address: yufeiz@mit.edu