

# Walks, infinite series and spectral radius of graphs

Wenqian Zhang

School of Mathematics and Statistics, Shandong University of Technology

Zibo, Shandong 255000, P.R. China

## Abstract

For a graph  $G$ , the spectral radius  $\rho(G)$  of  $G$  is the largest eigenvalue of its adjacency matrix. In this paper, we seek the relationship between  $\rho(G)$  and the walks of the subgraphs of  $G$ . Especially, if  $G$  contains a complete multi-partite graph as a spanning subgraph, we give a formula for  $\rho(G)$  by using an infinite series on walks of the subgraphs of  $G$ . These results are useful for the current popular spectral extremal problem.

**Keywords:** spectral radius; spectral extrema; walk; infinite series.

**2020 Mathematics Subject Classification:** 05C50.

## 1 Introduction

All graphs considered in this paper are finite, undirected and without parallel edges or loops. For a graph  $G$ , let  $\overline{G}$  be its complement. The vertex set and edge set of  $G$  are denoted by  $V(G)$  and  $E(G)$ , respectively. For a vertex  $u$ , let  $d_G(u)$  be its degree. For two vertices  $u$  and  $v$ , we say  $u \sim v$  if they are adjacent in  $G$ . For a certain integer  $n$ , let  $K_n$  and  $K_{1,n-1}$  be the complete graph and the star of order  $n$ , respectively. For  $\ell \geq 2$  vertex-disjoint graphs  $G_1, G_2, \dots, G_\ell$ , let  $\cup_{1 \leq i \leq \ell} G_i$  be the disjoint union of them. Let  $G_1 \vee G_2$  be the join of  $G_1$  and  $G_2$ . For any terminology used but not defined here, one may refer to [2].

Let  $G$  be a graph with vertices  $v_1, v_2, \dots, v_n$ . The *adjacency matrix* of  $G$  is an  $n \times n$  matrix  $(a_{ij})$ , where  $a_{ij} = 1$  if  $v_i \sim v_j$ , and  $a_{ij} = 0$  otherwise. The *spectral radius*  $\rho(G)$  of  $G$  is the largest eigenvalue of its adjacency matrix. By Perron–Frobenius theorem,  $\rho(G)$  has a non-negative eigenvector. Any non-negative eigenvector corresponding to  $\rho(G)$  is called a *Perron vector* of  $G$ . If  $G$  is connected, any Perron vector of  $G$  has positive entries.

For a family  $\mathcal{H}$  of graphs, a graph  $G$  is call  $\mathcal{H}$ -free if  $G$  does not contain any one in  $\mathcal{H}$  as a subgraph. Let  $\text{EX}(n, \mathcal{H})$  be the set of  $\mathcal{H}$ -free graphs of order  $n$  with the maximum

---

E-mails: zhangwq@pku.edu.cn

number of edges, and let  $\text{SPEX}(n, \mathcal{H})$  be the set of  $\mathcal{H}$ -free graphs of order  $n$  with the maximum spectral radius. For  $r \geq 2, n \geq r$ , let  $T_{n,r}$  be the Turán graph of order  $n$  with  $r$  parts. Turán's famous theorem states that  $\text{EX}(n, \{K_{r+1}\}) = \{T_{n,r}\}$ . Nikiforov [17] proves that  $\text{SPEX}(n, \{K_{r+1}\}) = \{T_{n,r}\}$ . In 2010, Nikiforov [16] formally presented a spectral version of Turán-type problem: what is the maximum spectral radius of an  $\mathcal{H}$ -free graph of order  $n$ ? In recent years, this problem has become very popular (for example, see [1, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15, 18, 19]). In particular, Wang, Kang and Xue [18] proved the following nice result, which is conjectured by Cioabă, Desai and Tait [5].

**Theorem 1.1** (Wang, Kang and Xue [18]) *For  $r \geq 2$  and sufficiently large  $n$ , let  $\mathcal{H}$  be a finite family of graphs such that all graphs in  $\text{EX}(n, \mathcal{H})$  are obtained from  $T_{n,r}$  by embedding  $k$  edges into its  $r$  parts, where  $k$  is a bounded integer. Then  $\text{SPEX}(n, \mathcal{H}) \subseteq \text{EX}(n, \mathcal{H})$ .*

In general,  $\text{SPEX}(n, \mathcal{H})$  is a proper subset of  $\text{EX}(n, \mathcal{H})$  in Theorem 1.1. To precisely characterize the graphs in  $\text{SPEX}(n, \mathcal{H})$ , we need to develop new techniques.

Let  $G$  be a graph. For an integer  $\ell \geq 1$ , a *walk* of length  $\ell$  in  $G$  is an ordered sequence of vertices  $v_0, v_1, \dots, v_\ell$ , such that  $v_i \sim v_{i+1}$  for any  $0 \leq i \leq \ell - 1$ . The walk is called an  $\ell$ -walk, and the vertex  $v_0$  is called the starting vertex of the walk. For a vertex  $u$  of  $G$ , let  $w_G^\ell(u)$  be the number of  $\ell$ -walks starting at  $u$  in  $G$ . Let  $W^\ell(G) = \sum_{v \in V(G)} w_G^\ell(v)$ . Clearly,  $w_G^1(u) = d_G(u)$  and

$$w_G^{\ell+1}(u) = \sum_{v \in V(G), v \sim u} w_G^\ell(v)$$

for any  $\ell \geq 1$ . Thus  $W^1(G) = 2|E(G)|$ , and

$$W^2(G) = \sum_{u \in V(G)} \sum_{v \in V(G), v \sim u} d_G(v) = \sum_{v \in V(G)} d_G^2(v).$$

Let  $G_1$  and  $G_2$  be two graphs (probably of different orders). We say that  $G_1$  is strictly *walk-preferable* to  $G_2$  (denoted by  $G_1 \succ G_2$ ), if there is an integer  $\ell \geq 1$  such that  $W^\ell(G_1) > W^\ell(G_2)$  and  $W^i(G_1) = W^i(G_2)$  for any  $1 \leq i \leq \ell - 1$ ;  $G_1$  is *walk-equivalent* to  $G_2$  (denoted by  $G_1 \equiv G_2$ ), if  $W^i(G_1) = W^i(G_2)$  for any  $i \geq 1$ . We say  $G_1$  is *walk-preferable* to  $G_2$  (denoted by  $G_1 \succeq G_2$ ), if  $G_1 \succ G_2$  or  $G_1 \equiv G_2$ . We also write  $G_2 \prec G_1$  or  $G_2 \preceq G_1$ , if  $G_1 \succ G_2$  or  $G_1 \succeq G_2$ . Let  $\mathcal{G}$  be a set of graphs. Let  $\text{SPEX}(\mathcal{G})$  denote the set of graphs with the maximum spectral radius in  $\mathcal{G}$ . Define  $\text{EX}^1(\mathcal{G}) = \{G \in \mathcal{G} \mid W^1(G) \geq W^1(G') \text{ for any } G' \in \mathcal{G}\}$ , and for any  $\ell \geq 2$  define  $\text{EX}^\ell(\mathcal{G}) = \{G \in \text{EX}^{\ell-1}(\mathcal{G}) \mid W^\ell(G) \geq W^\ell(G') \text{ for any } G' \in \text{EX}^{\ell-1}(\mathcal{G})\}$ . Clearly,  $\text{EX}^1(\mathcal{G})$  is precisely the set of graphs in  $\mathcal{G}$  with the maximum number of edges, and  $\text{EX}^{i+1}(\mathcal{G}) \subseteq \text{EX}^i(\mathcal{G})$  for any  $i \geq 1$ . Let  $\text{EX}^\infty(\mathcal{G}) = \bigcap_{1 \leq i \leq \infty} \text{EX}^i(\mathcal{G})$ . The graphs in  $\text{EX}^\infty(\mathcal{G})$  are called the most walk-preferable graphs in  $\mathcal{G}$ .

The following Theorem 1.2 and Theorem 1.3 reveal the relationship between the spectral radius of graphs and walks of their subgraphs.

**Theorem 1.2** *Let  $G$  be a connected graph of order  $n$ , and let  $S$  be a subset of  $V(G)$  with  $1 \leq |S| < n$ . Assume that  $T$  is a set of some isolated vertices of  $G - S$ , such that each vertex in  $T$  is adjacent to each vertex in  $S$  in  $G$ . Let  $H_1$  and  $H_2$  be two graphs with vertex set  $T$ . For any  $1 \leq i \leq 2$ , let  $G_i$  be the graph obtained from  $G$  by embedding the edges of  $H_i$  into  $T$ . When  $\rho(G)$  is sufficiently large (compared with  $|T|$ ), we have the following conclusions.*

- (i) *If  $H_1 \equiv H_2$ , then  $\rho(G_1) = \rho(G_2)$ .*
- (ii) *If  $H_1 \succ H_2$ , then  $\rho(G_1) > \rho(G_2)$ .*
- (iii) *If  $H_1 \prec H_2$ , then  $\rho(G_1) < \rho(G_2)$ .*

**Theorem 1.3** *For an integer  $r \geq 2$ , let  $K_{n_1, n_2, \dots, n_r}$  be the complete  $r$ -partite graph of order  $n$  with parts  $V_1, V_2, \dots, V_r$ , where  $n_i = |V_i| \geq 1$  for  $1 \leq i \leq r$  and  $\sum_{1 \leq i \leq r} n_i = n$ . For each  $1 \leq i \leq r$ , let  $H_i$  be a graph with vertex set  $V(H_i) \subseteq V_i$  and maximum degree at most  $\Delta$ . Let  $G$  be the graph obtained from  $K_{n_1, n_2, \dots, n_r}$  by embedding the edges of  $H_i$  into  $V_i$  for all  $1 \leq i \leq r$ . Let  $\rho = \rho(G)$ . If  $\rho > \Delta$ , then*

$$\sum_{1 \leq s \leq r} \frac{1}{1 + \frac{n_s}{\rho} + \sum_{i=1}^{\infty} \frac{W^i(H_s)}{\rho^{i+1}}} = r - 1.$$

To characterize the graphs in  $\text{SPEX}(n, \mathcal{H})$  in Theorem 1.1, we first introduce some notations. For  $r \geq 2$  and sufficiently large  $n$ , let  $T(n, r)$  be the Turán graph of order  $n$  with  $r$ -parts  $V_1, V_2, \dots, V_r$ , where  $n_i = |V_i|$  for  $1 \leq i \leq r$  and  $\lfloor \frac{n}{r} \rfloor = n_1 \leq n_2 \leq \dots \leq n_r = \lceil \frac{n}{r} \rceil$ . For a fixed integer  $t \geq 1$ , let  $\mathcal{T}_{n,r,t}$  be the set of all the graphs obtained from  $T(n, r)$  by embedding  $t$  edges into the  $r$ -parts  $V_1, V_2, \dots, V_r$ . Given a graph  $G \in \mathcal{T}_{n,r,t}$ , let  $H_i^G$  (probably empty) be the subgraph of  $G$  induced by the edges embedded into  $V_i$  for each  $1 \leq i \leq r$ . Clearly,  $H_1^G, H_2^G, \dots, H_r^G$  are vertex-disjoint. Let  $H^G = \cup_{1 \leq i \leq r} H_i^G$ . Then  $|E(H^G)| = \sum_{1 \leq i \leq r} |E(H_i^G)| = t$ .

For a subset  $\mathcal{G}$  of  $\mathcal{T}_{n,r,t}$ , let  $\mathcal{G}_j = \{G \in \mathcal{G} \mid H^G = \cup_{1 \leq i \leq r} H_i^G = H_j^G\}$  for any  $1 \leq j \leq r$ . Define  $\mathcal{H}^{\mathcal{G}} = \{H^G \mid G \in \mathcal{G}\}$  and  $\mathcal{H}^{\mathcal{G}_j} = \{H^G \mid G \in \mathcal{G}_j\}$  (if  $\mathcal{G}_j$  is not empty) for  $1 \leq j \leq r$ . The subset  $\mathcal{G}$  is called a *normal set* of  $\mathcal{T}_{n,r,t}$ , if it satisfies the property: for any  $1 \leq i \neq j \leq r$  and any  $G \in \mathcal{G}_i$ , there is a  $G' \in \mathcal{G}_j$  such that  $H^{G'} = H_j^{G'}$  is a copy of  $H^G = H_i^G$ .

**Theorem 1.4** *For  $t \geq 1, r \geq 2$  and sufficiently large  $n$ , let  $\mathcal{G}$  be a normal subset of  $\mathcal{T}_{n,r,t}$ . Let  $\mathcal{G}_i, \mathcal{H}^{\mathcal{G}}$  and  $\mathcal{H}^{\mathcal{G}_i}$  be defined as above for any  $1 \leq i \leq r$ . If  $\{G \in \mathcal{G} \mid H^G \in \text{EX}^2(\mathcal{H}^{\mathcal{G}})\} \subseteq \cup_{1 \leq i \leq r} \mathcal{G}_i$ , then*

- (i)  $\text{SPEX}(\mathcal{G}) = \cup_{i, n_i = \lfloor \frac{n}{r} \rfloor} \text{SPEX}(\mathcal{G}_i)$ ;

(ii) for any  $1 \leq i \leq r$  and any graph  $G \in \mathcal{G}_i$ ,  $G \in \text{SPEX}(\mathcal{G}_i)$  if and only if  $H^G \in \text{EX}^\infty(\mathcal{H}^{\mathcal{G}_i})$ .

Considering Theorem 1.2 and Theorem 1.4, it is interesting to study the following problem:

**Problem.** For a specified family  $\mathcal{H}$  of graphs, characterize the graphs in  $\text{EX}^\infty(\mathcal{H})$ .

Recall that  $W^2(G) = \sum_{v \in V(G)} d_G^2(v)$  for any graph  $G$ . The following result is taken from Lemma 2.4 of [10] (it is trivial for  $m = 1$ ).

**Lemma 1.5** (Ismailescu and Stefanica [10]) *Let  $\mathcal{M}_{n,m}$  be the set of all the graphs of order  $n$  with  $m$  edges, where  $m \geq 1$  and  $n \geq m+2$ . Then  $\text{EX}^2(\mathcal{M}_{n,m}) = \{K_{1,3} \cup \overline{K_{n-4}}, K_3 \cup \overline{K_{n-3}}\}$  for  $m = 3$ , and  $\text{EX}^2(\mathcal{M}_{n,m}) = \{K_{1,m} \cup \overline{K_{n-1-m}}\}$  otherwise.*

It is easy to see that  $W^3(K_3) = 24 > 18 = W^3(K_{1,3})$ . Since a graph of order less than  $n$  can be viewed as a graph of order  $n$  by adding some isolated vertices, the following result can be deduced from Lemma 1.5 directly (by letting  $n = 2m + 2$ ).

**Corollary 1.6** *Let  $\mathcal{M}_m$  be the set of graphs with  $m$  edges and without isolated vertices, where  $m \geq 1$ . Then we have the following conclusions.*

- (i)  $\text{EX}^2(\mathcal{M}_m) = \{K_{1,3}, K_3\}$  for  $m = 3$ , and  $\text{EX}^2(\mathcal{M}_m) = \{K_{1,m}\}$  otherwise
- (ii)  $\text{EX}^\infty(\mathcal{M}_m) = \text{EX}^3(\mathcal{M}_m) = \{K_3\}$  for  $m = 3$ , and  $\text{EX}^\infty(\mathcal{M}_m) = \text{EX}^3(\mathcal{M}_m) = \{K_{1,m}\}$  otherwise.

For a graph  $G$ , the *maximum average degree* of  $G$  is the maximum value of  $\frac{2|E(H)|}{|H|}$ , where  $H$  is an induced subgraph of  $G$  with  $|H| \geq 1$ . For  $k \geq 1$  and  $n \geq 2k + 1$ , Let  $\mathcal{N}_{n,k}$  be the set of graphs of order  $n$  with maximum average degree at most  $2k$ . Zhang [20] proved that any graph  $G \in \text{SPEX}(\mathcal{N}_{n,k})$  is obtained from  $K_k \vee \overline{K_{n-k}}$  by embedding  $\frac{k(k+1)}{2}$  edges into the part  $V(\overline{K_{n-k}})$ . When  $k$  is given and  $n$  is sufficiently large, from Theorem 1.2 together with Corollary 1.6, we see that such  $G$  is unique, of which the subgraph induced by the  $\frac{k(k+1)}{2}$  edges is  $K_3$  for  $k = 2$ , and is  $K_{1, \frac{k(k+1)}{2}}$  otherwise. (Note that, we can assume that the  $\frac{k(k+1)}{2}$  edges are embedded into a fixed subset of size  $k(k+1)$  which is viewed as  $T$  in Theorem 1.2.)

A graph  $F$  is called *color-critical* if there is an edge  $e$  whose deletion induces a subgraph with less chromatic number than  $F$ . As in [9], for  $k \geq 2$  color-critical graphs with chromatic number  $r+1 \geq 3$ :  $F_1, F_2, \dots, F_k$ , let  $\mathcal{G}(F_1, F_2, \dots, F_k)$  denote the family of graphs which consists of  $k$  edge-disjoint copies of  $F_1, F_2, \dots, F_k$ . Fang, Tait and Zhai [9] proved that for sufficiently large  $n$ ,  $\text{EX}(n, \mathcal{G}(F_1, F_2, \dots, F_k))$  is precisely  $\mathcal{T}_{n,r,k-1}$  defined as above. From Theorem 1.1 we see  $\text{SPEX}(n, \mathcal{G}(F_1, F_2, \dots, F_k)) \subseteq \text{EX}(n, \mathcal{G}(F_1, F_2, \dots, F_k)) = \mathcal{T}_{n,r,k-1}$ .

The authors in [9] also devoted themselves to characterize the graphs with the maximum spectral radius in  $\mathcal{T}_{n,r,k-1}$  (this result is given by Lin, Zhai and Zhai [14] for the case  $F_1 = F_2 = \dots = F_k = K_3$ ). In detail, they [9] proved the following Corollary 1.7, which is directly deduced from Theorem 1.4 together with Corollary 1.6. (The condition in Theorem 1.4 holds for  $\mathcal{G} = \mathcal{T}_{n,r,k-1}$ . Clearly,  $\mathcal{T}_{n,r,k-1}$  itself is a normal subset of  $\mathcal{T}_{n,r,k-1}$ . Moreover, the condition

$$\{G \in \mathcal{T}_{n,r,k-1} \mid H^G \in \text{EX}^2(\mathcal{H}^{\mathcal{T}_{n,r,k-1}})\} \subseteq \cup_{1 \leq i \leq r} \mathcal{G}_i$$

holds, since the one or two graphs in  $\text{EX}^2(\mathcal{H}^{\mathcal{T}_{n,r,k-1}})$  are connected by Corollary 1.6.)

**Corollary 1.7** ([9, 14]) *For  $k \geq 2, r \geq 2$  and sufficiently large  $n$ , let  $\mathcal{T}_{n,r,k-1}$  be defined as above. Then each graph in  $\text{SPEX}(\mathcal{T}_{n,r,k-1})$  is obtained from  $T_{n,r}$  by embedding  $k-1$  edges into its one part of size  $\lfloor \frac{n}{r} \rfloor$ , of which the subgraph induced by the  $k-1$  edges is  $K_3$  for  $k=4$  and is  $K_{1,k-1}$  otherwise.*

## 2 Proofs of Theorem 1.2 and Theorem 1.3

The following Lemma 2.1 is taken from Theorem 8.1.1 of [2].

**Lemma 2.1** ([2]) *If  $H$  is a subgraph of a connected graph  $G$ , then  $\rho(H) \leq \rho(G)$ , with equality if and only if  $H = G$ .*

**Lemma 2.2** *Let  $G$  be a connected graph of order  $n$ , and let  $S$  be a subset of  $V(G)$  with  $1 \leq |S| < n$ . Assume that  $H$  is a component of  $G - S$  with maximum degree at most  $\Delta$ , such that each vertex of  $H$  is adjacent to each vertex in  $S$ . Let  $\rho = \rho(G)$  and let  $\mathbf{x} = (x_v)_{v \in V(G)}$  be the Perron vector of  $G$  such that  $\sum_{v \in S} x_v = \rho$ . If  $\rho > \Delta$ , then for any  $u \in T$  we have*

$$x_u = 1 + \sum_{i=1}^{\infty} \frac{w_H^i(u)}{\rho^i}.$$

**Proof:** Denote  $V(H) = T$ . Recall that  $\sum_{v \in S} x_v = \rho$ . For any  $u \in T$ , we have

$$\rho x_u = \left( \sum_{v \in S} x_v \right) + \sum_{v \sim u, v \in T} x_v = \rho + \sum_{v \sim u, v \in T} x_v.$$

This implies that  $x_u = 1 + \frac{\sum_{v \sim u, v \in T} x_v}{\rho}$ .

**Claim 1.** For any vertex  $u$  in  $T$ , we have  $x_u \geq 1 + \sum_{1 \leq i \leq \ell} \frac{w_H^i(u)}{\rho^i}$  for any integer  $\ell \geq 1$ .

**Proof of Claim 1.** From  $x_u = 1 + \frac{\sum_{v \sim u, v \in T} x_v}{\rho}$  we see that  $x_u \geq 1$  for any  $u \in T$ . Thus, for any  $v \in T$  and  $v \sim u$  we have  $x_v \geq 1$ . By  $x_u = 1 + \frac{\sum_{v \sim u, v \in T} x_v}{\rho}$  we have  $x_u \geq 1 + \frac{w_H^1(u)}{\rho}$

for any  $u \in T$ . Thus Claim 1 holds for  $\ell = 1$ . Assume that Claim 1 holds for  $\ell = j \geq 1$ . That is  $x_u \geq 1 + \sum_{1 \leq i \leq j} \frac{w_H^i(u)}{\rho^i}$  for any  $u \in T$ . Now we consider the case  $\ell = j + 1$ . Let  $u$  be a vertex in  $T$ . Note that  $x_v \geq 1 + \sum_{1 \leq i \leq j} \frac{w_H^i(v)}{\rho^i}$  for any  $v \in T$  and  $v \sim u$  by induction on  $\ell = j$ . By  $x_u = 1 + \frac{\sum_{v \sim u, v \in T} x_v}{\rho}$ , we have

$$x_u \geq 1 + \frac{\sum_{v \sim u, v \in T} (1 + \sum_{1 \leq i \leq j} \frac{w_H^i(v)}{\rho^i})}{\rho} = 1 + \sum_{1 \leq i \leq j+1} \frac{w_H^i(u)}{\rho^i}.$$

So, Claim 1 holds for  $\ell = j + 1$ . By induction on  $\ell$ , we complete the proof of Claim 1.  $\square$

**Claim 2.** For any vertex  $u$  in  $T$ , we have  $x_u \leq (1 + \sum_{1 \leq i \leq \ell} \frac{w_H^i(u)}{\rho^i}) + \frac{\Delta}{\rho - \Delta} \frac{w_H^\ell(u)}{\rho^\ell}$  for any integer  $\ell \geq 1$ .

**Proof of Claim 2.** Let  $u^*$  be a vertex in  $T$  such that  $x_{u^*} = \max_{v \in T} x_v$ . Recall that  $\rho > \Delta$ . From  $x_{u^*} = 1 + \frac{\sum_{v \sim u^*, v \in T} x_v}{\rho} \leq 1 + \frac{\Delta \cdot x_{u^*}}{\rho}$  we see that  $x_{u^*} \leq \frac{\rho}{\rho - \Delta}$ .

For any vertex  $u \in T$ , from  $x_u = 1 + \frac{\sum_{v \sim u, v \in T} x_v}{\rho}$  we obtain that

$$x_u \leq 1 + \frac{w_H^1(u) \cdot x_{u^*}}{\rho} \leq (1 + \frac{w_H^1(u)}{\rho}) + \frac{\Delta}{\rho - \Delta} \frac{w_H^1(u)}{\rho}.$$

Thus Claim 2 holds for  $\ell = 1$ . Assume that Claim 2 holds for  $\ell = j \geq 1$ . That is  $x_u \leq (1 + \sum_{1 \leq i \leq j} \frac{w_H^i(u)}{\rho^i}) + \frac{\Delta}{\rho - \Delta} \frac{w_H^j(u)}{\rho^j}$  for any  $u \in T$ . Now we consider the case  $\ell = j + 1$ . Let  $u \in T$ . Note that  $x_v \leq (1 + \sum_{1 \leq i \leq j} \frac{w_H^i(v)}{\rho^i}) + \frac{\Delta}{\rho - \Delta} \frac{w_H^j(v)}{\rho^j}$  for any  $v \in T$  and  $v \sim u$  by induction on  $\ell = j$ . By  $x_u = 1 + \frac{\sum_{v \sim u, v \in T} x_v}{\rho}$ , we have

$$x_u \leq 1 + \frac{\sum_{v \sim u, v \in T} ((1 + \sum_{1 \leq i \leq j} \frac{w_H^i(v)}{\rho^i}) + \frac{\Delta}{\rho - \Delta} \frac{w_H^j(v)}{\rho^j})}{\rho} = (1 + \sum_{1 \leq i \leq j+1} \frac{w_H^i(u)}{\rho^i}) + \frac{\Delta}{\rho - \Delta} \frac{w_H^{j+1}(u)}{\rho^{j+1}}.$$

So, Claim 2 holds for  $\ell = j + 1$ . By induction on  $\ell$ , we complete the proof of Claim 2.  $\square$

Since  $\rho > \Delta$ , we see that the infinite series  $\sum_{i=1}^{\infty} (\frac{\Delta}{\rho})^i$  exists and  $(\frac{\Delta}{\rho})^\ell \rightarrow 0$  for  $\ell \rightarrow \infty$ . Let  $u \in T$ . For any  $\ell \geq 1$ , clearly,  $w_H^\ell(u) \leq \Delta^\ell$ . Thus  $\sum_{i=1}^{\infty} \frac{w_H^i(u)}{\rho^i}$  exists and  $\frac{w_H^\ell(u)}{\rho^\ell} \rightarrow 0$  for  $\ell \rightarrow \infty$ . By Claim 1 and Claim 2 we have

$$1 + \sum_{1 \leq i \leq \ell} \frac{w_H^i(u)}{\rho^i} \leq x_u \leq (1 + \sum_{1 \leq i \leq \ell} \frac{w_H^i(u)}{\rho^i}) + \frac{\Delta}{\rho - \Delta} \frac{w_H^\ell(u)}{\rho^\ell},$$

for any integer  $\ell \geq 1$ . By letting  $\ell \rightarrow \infty$  we obtain that  $x_u = 1 + \sum_{i=1}^{\infty} \frac{w_H^i(u)}{\rho^i}$ . This completes the proof.  $\square$

**Proof of Theorem 1.2.** Clearly, both  $G_1$  and  $G_2$  are connected. Let  $\rho_1 = \rho(G_1)$  and  $\rho_2 = \rho(G_2)$ . Then  $\rho_1, \rho_2 \geq \rho(G)$  by Lemma 2.1. Let  $\mathbf{x} = (x_v)_{v \in V(G_1)}$  be a Perron vector

of  $G_1$  such that  $\sum_{v \in S} x_v = \rho_1$ . Let  $\mathbf{y} = (y_v)_{v \in V(G_2)}$  be a Perron vector of  $G_2$  such that  $\sum_{v \in S} x_v = \rho_2$ .

**Claim 1.**  $(\rho_1 - \rho_2) \sum_{v \in V(G)} x_v y_v = \sum_{\ell=0}^{\infty} \left( \frac{W^{\ell+1}(H_1)}{\rho_1^\ell} - \frac{W^{\ell+1}(H_2)}{\rho_2^\ell} \right)$ .

**Proof of Claim 1.** Clearly, the only distinct edges of  $G_1$  and  $G_2$  are inside  $T$ . Let  $A(G_1)$  and  $A(G_2)$  be the adjacency matrices of  $G_1$  and  $G_2$ , respectively. Let  $\mathbf{x}'$  be the transpose of  $\mathbf{x}$ . For any  $u \in T$ , we write  $w_{H_i}^0(u) = 1$  for any  $1 \leq i \leq 2$ . Thus  $x_u = \sum_{i=0}^{\infty} \frac{w_{H_1}^i(u)}{\rho_1^i}$  and  $y_u = \sum_{i=0}^{\infty} \frac{w_{H_2}^i(u)}{\rho_2^i}$  by Lemma 2.2. Then

$$\begin{aligned}
& (\rho_1 - \rho_2) \sum_{v \in V(G)} x_v y_v \\
&= \mathbf{x}' (A(G_1) - A(G_2)) \mathbf{y} \\
&= \left( \sum_{uv \in E(H_1)} (x_u y_v + x_v y_u) \right) - \left( \sum_{uv \in E(H_2)} (x_u y_v + x_v y_u) \right) \\
&= \left( \sum_{u \in T} y_u \sum_{v \in T, vu \in E(H_1)} x_v \right) - \left( \sum_{u \in T} x_u \sum_{v \in T, vu \in E(H_2)} y_v \right) \\
&= \left( \sum_{u \in T} \left( \sum_{i=0}^{\infty} \frac{w_{H_2}^i(u)}{\rho_2^i} \right) \left( \sum_{v \in T, vu \in E(H_1)} \sum_{i=0}^{\infty} \frac{w_{H_1}^i(v)}{\rho_1^i} \right) \right) - \left( \sum_{u \in T} \left( \sum_{i=0}^{\infty} \frac{w_{H_1}^i(u)}{\rho_1^i} \right) \left( \sum_{v \in T, vu \in E(H_2)} \sum_{i=0}^{\infty} \frac{w_{H_2}^i(v)}{\rho_2^i} \right) \right) \\
&= \left( \sum_{u \in T} \left( \sum_{i=0}^{\infty} \frac{w_{H_2}^i(u)}{\rho_2^i} \right) \left( \sum_{i=0}^{\infty} \frac{w_{H_1}^{i+1}(u)}{\rho_1^i} \right) \right) - \left( \sum_{u \in T} \left( \sum_{i=0}^{\infty} \frac{w_{H_1}^i(u)}{\rho_1^i} \right) \left( \sum_{i=0}^{\infty} \frac{w_{H_2}^{i+1}(u)}{\rho_2^i} \right) \right) \\
&= \left( \sum_{u \in T} \sum_{\ell=0}^{\infty} \sum_{i, j \geq 0, i+j=\ell} \frac{w_{H_1}^{i+1}(u) w_{H_2}^j(u)}{\rho_1^i \rho_2^j} \right) - \left( \sum_{u \in T} \sum_{\ell=0}^{\infty} \sum_{i, j \geq 0, i+j=\ell} \frac{w_{H_1}^i(u) w_{H_2}^{j+1}(u)}{\rho_1^i \rho_2^j} \right) \\
&= \sum_{u \in T} \sum_{\ell=0}^{\infty} \left( \frac{w_{H_1}^{\ell+1}(u) w_{H_2}^0(u)}{\rho_1^\ell \rho_2^0} - \frac{w_{H_1}^0(u) w_{H_2}^{\ell+1}(u)}{\rho_1^0 \rho_2^\ell} \right) \\
&= \sum_{\ell=0}^{\infty} \left( \frac{W^{\ell+1}(H_1)}{\rho_1^\ell} - \frac{W^{\ell+1}(H_2)}{\rho_2^\ell} \right).
\end{aligned}$$

This finishes the proof of Claim 1.  $\square$

Clearly, there are at most  $2^{\frac{1}{2}|T|^2}$  graphs with vertex set  $T$ . Thus there are at most  $\frac{1}{2}(2^{\frac{1}{2}|T|^2})^2 = 2^{|T|^2-1}$  pairs of graphs  $Q_1, Q_2$  with vertex set  $T$  such that  $Q_1 \succ Q_2$ . Let  $\ell_{Q_1, Q_2}$  be the integer such that  $W^{\ell_{Q_1, Q_2}}(Q_1) > W^{\ell_{Q_1, Q_2}}(Q_2)$  and  $W^i(Q_1) = W^i(Q_2)$  for any  $1 \leq i \leq \ell_{Q_1, Q_2} - 1$ . Let  $C = \max_{Q_1 \succ Q_2, V(Q_1) = V(Q_2) = T} \ell_{Q_1, Q_2}$ . Note that  $C$  is only related to  $|T|$ . When  $\rho(G)$  is sufficiently large, we can assume  $\rho(G) > |T| - 1 + (|T| - 1)^{C+1} |T|$  in the following discussion.

(i) Let  $H_1 \equiv H_2$ . Then  $W^i(H_1) = W^i(H_2)$  for any  $i \geq 1$ . Without loss of generality,

assume  $\rho_1 \geq \rho_2$ . Then by Claim 1 we have

$$(\rho_1 - \rho_2) \sum_{v \in V(G)} x_v y_v = \sum_{\ell=0}^{\infty} \left( \frac{W^{\ell+1}(H_1)}{\rho_1^\ell} - \frac{W^{\ell+1}(H_2)}{\rho_2^\ell} \right) \leq 0.$$

It follows that  $\rho_1 \leq \rho_2$ . Consequently,  $\rho_1 = \rho_2$ .

(ii) Let  $H_1 \succ H_2$ . There is an integer  $1 \leq k \leq C$  (defined as above), such that  $W^k(H_1) > W^k(H_2)$  and  $W^i(H_1) = W^i(H_2)$  for any  $1 \leq i \leq k-1$ . We shall prove  $\rho_1 > \rho_2$ . Suppose  $\rho_1 \leq \rho_2$  by contradiction. Then by Claim 1 we have

$$\begin{aligned} (\rho_1 - \rho_2) \sum_{v \in V(G)} x_v y_v &= \sum_{\ell=0}^{\infty} \left( \frac{W^{\ell+1}(H_1)}{\rho_1^\ell} - \frac{W^{\ell+1}(H_2)}{\rho_2^\ell} \right) \\ &\geq \sum_{\ell=0}^{\infty} \left( \frac{W^{\ell+1}(H_1)}{\rho_2^\ell} - \frac{W^{\ell+1}(H_2)}{\rho_2^\ell} \right) \\ &= \sum_{\ell=k-1}^{\infty} \left( \frac{W^{\ell+1}(H_1)}{\rho_2^\ell} - \frac{W^{\ell+1}(H_2)}{\rho_2^\ell} \right) \\ &\geq \frac{1}{\rho_2^{k-1}} - \sum_{\ell=k}^{\infty} \frac{W^{\ell+1}(H_2)}{\rho_2^\ell} \\ &\geq \frac{1}{\rho_2^{k-1}} - \sum_{\ell=k}^{\infty} \frac{(|T| - 1)^{\ell+1} |T|}{\rho_2^\ell} \\ &= \frac{1}{\rho_2^{k-1}(\rho_2 - |T| + 1)} (\rho_2 - |T| + 1 - (|T| - 1)^{k+1} |T|) \\ &\geq \frac{1}{\rho_2^{k-1}(\rho_2 - |T| + 1)} (\rho_2 - |T| + 1 - (|T| - 1)^{C+1} |T|) > 0. \end{aligned}$$

It follows that  $\rho_1 > \rho_2$ , which contradicts the assumption  $\rho_1 \leq \rho_2$ . Hence we have  $\rho_1 > \rho_2$ .

(iii) The proof is very similar to (ii). This completes the proof.  $\square$

**Proof of Theorem 1.3.** Let  $\mathbf{x} = (x_v)_{v \in V(G)}$  be the Perron vector of  $G$  such that  $\sum_{v \in V(G)} x_v = 1$ . For each  $1 \leq s \leq r$ , let  $\mathbf{y}^s = \frac{\rho}{1 - \sum_{v \in V_s} x_v} \mathbf{x}$ . Then  $\mathbf{y}^s = (y_v^s)_{v \in V(G)}$  is a Perron vector of  $G$  such that

$$\sum_{v \in V(G) - V_s} y_v^s = \frac{\rho}{1 - \sum_{v \in V_s} x_v} \sum_{v \in V(G) - V_s} x_v = \rho.$$

By Lemma 2.2, we have

$$y_u^s = 1 + \sum_{i=1}^{\infty} \frac{w_{H_s}^i(u)}{\rho^i},$$

for any  $u \in V_s$ . It follows that

$$x_u = \frac{1 - \sum_{v \in V_s} x_v}{\rho} \left( 1 + \sum_{i=1}^{\infty} \frac{w_{H_s}^i(u)}{\rho^i} \right),$$



for any  $u \in V_s$ . Thus

$$\sum_{u \in V_s} x_u = \frac{1 - \sum_{v \in V_s} x_v}{\rho} \sum_{u \in V_s} \left(1 + \sum_{i=1}^{\infty} \frac{w_{H_s}^i(u)}{\rho^i}\right) = \frac{1 - \sum_{u \in V_s} x_u}{\rho} \left(n_s + \sum_{i=1}^{\infty} \frac{W^i(H_s)}{\rho^i}\right).$$

It follows that

$$\sum_{u \in V_s} x_u = 1 - \frac{1}{1 + \frac{n_s}{\rho} + \sum_{i=1}^{\infty} \frac{W^i(H_s)}{\rho^{i+1}}}.$$

Then

$$1 = \sum_{1 \leq s \leq r} \sum_{u \in V_s} x_u = r - \sum_{1 \leq s \leq r} \frac{1}{1 + \frac{n_s}{\rho} + \sum_{i=1}^{\infty} \frac{W^i(H_s)}{\rho^{i+1}}},$$

implying that

$$\sum_{1 \leq s \leq r} \frac{1}{1 + \frac{n_s}{\rho} + \sum_{i=1}^{\infty} \frac{W^i(H_s)}{\rho^{i+1}}} = r - 1.$$

This completes the proof.  $\square$

### 3 Proof of Theorem 1.4

In this section, we prove Theorem 1.4 by using Theorem 1.2 and Theorem 1.3. For  $t \geq 1, r \geq 2$  and sufficiently large  $n$ , let  $\mathcal{G}$  be a normal subset of  $\mathcal{T}_{n,r,t}$ . Let  $\mathcal{G}_i, \mathcal{H}^{\mathcal{G}}$  and  $\mathcal{H}^{\mathcal{G}_i}$  for  $1 \leq i \leq r$  be defined as in the introduction.

**Proof of Theorem 1.4.** For any  $G \in \mathcal{G}$ , we have  $\rho(G) \geq \rho(T(n, r))$  by Lemma 2.1, and  $\rho(G) \leq \rho(T(n, r)) + \rho(H^{\mathcal{G}})$  by Courant-Weyl inequalities (see Theorem 1.3.15 of [2]). As we know,  $\rho(H^{\mathcal{G}}) \leq \sqrt{2|E(H^{\mathcal{G}})|} = \sqrt{2t}$ . Thus  $\rho(T(n, r)) \leq \rho(G) \leq \rho(T(n, r)) + \sqrt{2t}$ . Note that  $\frac{\rho(T(n, r))}{n} \rightarrow 1 - \frac{1}{r}$  for  $n \rightarrow \infty$ . Thus for any  $1 \leq s \leq r$ , we have  $\frac{n_s}{\rho(G)} \rightarrow \frac{\frac{1}{r}}{1 - \frac{1}{r}} = \frac{1}{r-1}$  for  $n \rightarrow \infty$ . Let  $\rho = \rho(G)$ . Then by Theorem 1.3 we have

$$\sum_{1 \leq s \leq r} \frac{1}{1 + \frac{n_s}{\rho} + \sum_{i=1}^{\infty} \frac{W^i(H_s^{\mathcal{G}})}{\rho^{i+1}}} = r - 1.$$

Let

$$f(G, x) = \sum_{1 \leq s \leq r} \frac{1}{1 + \frac{n_s}{x} + \sum_{i=1}^{\infty} \frac{W^i(H_s^{\mathcal{G}})}{x^{i+1}}},$$

where  $x > 0$  is large enough such that  $\sum_{i=1}^{\infty} \frac{W^i(H_s^{\mathcal{G}})}{x^{i+1}}$  exists for  $1 \leq s \leq r$ . Clearly,  $f(G, x)$  is strictly increasing with respect to  $x$ . Thus  $\rho$  is the largest root of  $f(G, x) = r - 1$ . Note that  $\frac{1}{1+x} = 1 + \sum_{i \geq 1} (-x)^i$  for  $0 \leq x < 1$ . Thus for  $1 \leq s \leq r$ , (noting that  $W^i(H_s^{\mathcal{G}}) \leq 2t \cdot t^i$

for any  $i \geq 1$ ) we have

$$\begin{aligned}
& \frac{1}{1 + \frac{n_s}{\rho} + \sum_{i=1}^{\infty} \frac{W^i(H_s^G)}{\rho^{i+1}}} \\
&= \frac{1}{1 + \frac{n_s}{\rho}} \frac{1}{1 + \frac{1}{1 + \frac{n_s}{\rho}} \sum_{i=1}^{\infty} \frac{W^i(H_s^G)}{\rho^{i+1}}} \\
&= \frac{1}{1 + \frac{n_s}{\rho}} \left( 1 - \frac{1}{1 + \frac{n_s}{\rho}} \frac{W^1(H_s^G)}{\rho^2} - \frac{1}{1 + \frac{n_s}{\rho}} \frac{W^2(H_s^G)}{\rho^3} + \Theta\left(\frac{1}{\rho^4}\right) \right).
\end{aligned}$$

Then

$$f(G, x) = \left( \sum_{1 \leq s \leq r} \left( \frac{1}{1 + \frac{n_s}{x}} - \frac{1}{(1 + \frac{n_s}{x})^2} \frac{W^1(H_s^G)}{x^2} - \frac{1}{(1 + \frac{n_s}{x})^2} \frac{W^2(H_s^G)}{x^3} \right) \right) + \Theta\left(\frac{1}{x^4}\right).$$

Without loss of generality, assume  $G \in \text{SPEX}(\mathcal{G})$ . Let  $G' \in \mathcal{G}$  such that  $H^{G'} \in \text{EX}^2(\mathcal{H}^{\mathcal{G}})$ . Then  $G' \in \cup_{1 \leq i \leq r} \mathcal{G}_i$  by the condition. Since  $\mathcal{G}$  is a normal subset of  $\mathcal{T}_{n,r,t}$ , we can assume that  $G' \in \mathcal{G}_1$ . Note that  $\rho = \rho(G) \geq \rho(G')$ . Since  $f(G', x)$  is strictly increasing with respect to  $x$  and  $f(G', \rho(G')) = r - 1$ , we have  $f(G', \rho) \geq r - 1$ . It follows that  $f(G', \rho) \geq f(G, \rho)$  as  $f(G, \rho) = r - 1$ . Thus  $\rho^3(f(G', \rho) - f(G, \rho)) \geq 0$ . On the other hand, we have

$$\begin{aligned}
& \rho^3(f(G', \rho) - f(G, \rho)) \\
&= \rho^3 \left( \sum_{1 \leq s \leq r} \left( \frac{1}{1 + \frac{n_s}{\rho}} - \frac{1}{(1 + \frac{n_s}{\rho})^2} \frac{W^1(H_s^{G'})}{\rho^2} - \frac{1}{(1 + \frac{n_s}{\rho})^2} \frac{W^2(H_s^{G'})}{\rho^3} \right) \right) + \Theta\left(\frac{1}{\rho}\right) \\
&\quad - \rho^3 \left( \sum_{1 \leq s \leq r} \left( \frac{1}{1 + \frac{n_s}{\rho}} - \frac{1}{(1 + \frac{n_s}{\rho})^2} \frac{W^1(H_s^G)}{\rho^2} - \frac{1}{(1 + \frac{n_s}{\rho})^2} \frac{W^2(H_s^G)}{\rho^3} \right) \right) - \Theta\left(\frac{1}{\rho}\right) \\
&= \left( \rho \sum_{1 \leq s \leq r} \frac{W^1(H_s^G) - W^1(H_s^{G'})}{(1 + \frac{n_s}{\rho})^2} \right) + \left( \sum_{1 \leq s \leq r} \frac{W^2(H_s^G) - W^2(H_s^{G'})}{(1 + \frac{n_s}{\rho})^2} \right) + \Theta\left(\frac{1}{\rho}\right).
\end{aligned}$$

Since  $G' \in \mathcal{G}_1$ , we have that  $H^{G'} = H_1^{G'}$ , i.e.,  $H_i^{G'}$  is empty for  $2 \leq i \leq r$ . Thus  $W^1(H_1^{G'}) = \sum_{1 \leq s \leq r} W^1(H_s^G) = 2t$ , implying that  $\sum_{1 \leq s \leq r} \frac{W^1(H_s^G) - W^1(H_s^{G'})}{(1 + \frac{n_s}{\rho})^2} \leq 0$  (recall  $n_1 = \lfloor \frac{n}{r} \rfloor$ ). From  $\rho^3(f(G', \rho) - f(G, \rho)) \geq 0$ , we obtain that

$$\left( \sum_{1 \leq s \leq r} \frac{W^2(H_s^G) - W^2(H_s^{G'})}{(1 + \frac{n_s}{\rho})^2} \right) + \Theta\left(\frac{1}{\rho}\right) \geq 0.$$

Note that  $W^2(H_1^{G'}) \geq \sum_{1 \leq s \leq r} W^2(H_s^G) = W^2(H^G)$  as  $H^{G'} \in \text{EX}^2(\mathcal{H}^{\mathcal{G}})$ . If  $W^2(H_1^{G'}) > \sum_{1 \leq s \leq r} W^2(H_s^G)$ , then for  $n \rightarrow \infty$ ,

$$\sum_{1 \leq s \leq r} \frac{W^2(H_s^G) - W^2(H_s^{G'})}{(1 + \frac{n_s}{\rho})^2} \leq -\frac{1}{(1 + \frac{n_1}{\rho})^2} \rightarrow -\frac{1}{(1 + \frac{1}{r-1})^2}.$$

This contradicts that  $(\sum_{1 \leq s \leq r} \frac{W^2(H_s^G) - W^2(H_s^{G'})}{(1 + \frac{n_s}{\rho})^2}) + \Theta(\frac{1}{\rho}) \geq 0$ . Hence

$$W^2(H_1^{G'}) = \sum_{1 \leq s \leq r} W^2(H_s^G) = W^2(H^G).$$

Recall that  $H^{G'} \in \text{EX}^2(\mathcal{H}^{\mathcal{G}})$ . Thus  $H^G \in \text{EX}^2(\mathcal{H}^{\mathcal{G}})$ , implying that  $G \in \cup_{1 \leq i \leq r} \mathcal{G}_i$  by the condition of the theorem.

Suppose that  $G \in \mathcal{G}_j$  for some  $1 \leq j \leq r$  and  $n_j = \lceil \frac{n}{r} \rceil > \lfloor \frac{n}{r} \rfloor$ . Then for  $n \rightarrow \infty$ ,

$$\begin{aligned} \rho \sum_{1 \leq s \leq r} \frac{W^1(H_s^G) - W^1(H_s^{G'})}{(1 + \frac{n_s}{\rho})^2} &= t\rho \left( \frac{1}{(1 + \frac{n_j}{\rho})^2} - \frac{1}{(1 + \frac{n_1}{\rho})^2} \right) = \frac{-t(2 + \frac{n_j}{\rho} + \frac{n_1}{\rho})}{(1 + \frac{n_j}{\rho})^2 (1 + \frac{n_1}{\rho})^2} \\ &\rightarrow \frac{-t(2 + \frac{2}{r-1})}{(1 - \frac{1}{r-1})^4}. \end{aligned}$$

Clearly,  $\sum_{1 \leq s \leq r} \frac{W^2(H_s^G) - W^2(H_s^{G'})}{(1 + \frac{n_s}{\rho})^2} \leq 0$ . Thus for large  $n$ ,

$$\left( \rho \sum_{1 \leq s \leq r} \frac{W^1(H_s^G) - W^1(H_s^{G'})}{(1 + \frac{n_s}{\rho})^2} \right) + \left( \sum_{1 \leq s \leq r} \frac{W^2(H_s^G) - W^2(H_s^{G'})}{(1 + \frac{n_s}{\rho})^2} \right) + \Theta\left(\frac{1}{\rho}\right) < 0.$$

Recall that  $\rho^3(f(G', \rho) - f(G, \rho)) \geq 0$ , i.e.,

$$\left( \rho \sum_{1 \leq s \leq r} \frac{W^1(H_s^G) - W^1(H_s^{G'})}{(1 + \frac{n_s}{\rho})^2} \right) + \left( \sum_{1 \leq s \leq r} \frac{W^2(H_s^G) - W^2(H_s^{G'})}{(1 + \frac{n_s}{\rho})^2} \right) + \Theta\left(\frac{1}{\rho}\right) \geq 0.$$

This is a contradiction. Hence  $G \in \mathcal{G}_j$  for some  $1 \leq j \leq r$  and  $n_j = \lfloor \frac{n}{r} \rfloor$ . Thus we obtain

$$\text{SPEX}(\mathcal{G}) = \cup_{i, n_i = \lfloor \frac{n}{r} \rfloor} \text{SPEX}(\mathcal{G}_i).$$

For a graph  $G'' \in \mathcal{G}_i$  with  $1 \leq i \leq r$ , note that  $H^{G''} = H_i^{G''}$ . Since  $H_i^{G''}$  has  $t$  edges, we can assume that the edges of  $H_i^{G''}$  are embedded into a fixed subset of size  $2t$  in  $V_i$  (this fixed subset is viewed as  $T$  in Theorem 1.2). By Theorem 1.2, we see that  $G'' \in \text{SPEX}(\mathcal{G}_i)$  if and only if  $H^{G''} = H_i^{G''} \in \text{EX}^\infty(\mathcal{H}^{\mathcal{G}_i})$ . This completes the proof.  $\square$

### Declaration of competing interest

There is no conflict of interest.

### Data availability statement

No data was used for the research described in the article.

## References

- [1] J. Byrne, D.N. Desai and M. Tait, A general theorem in spectral extremal graph theory, arXiv:2401.07266v1.
- [2] D. Cvetković, P. Rowlinson and S. Simić, An Introduction to the Theory of Graph Spectra, Cambridge University Press, Cambridge, 2010.
- [3] S. Cioabă, L. Feng, M. Tait and X. Zhang, The maximum spectral radius of graphs without friendship subgraphs, *Electron. J. Combin.* 27 (2020) #4.22.
- [4] S. Cioabă, D. Desai and M. Tait, The spectral even cycle problem, arXiv:2205.00990v1.
- [5] S. Cioabă, D. Desai and M. Tait, The spectral radius of graphs with no odd wheels, *European J. Combin.* 99 (2022) 103420.
- [6] S. Cioabă, D. Desai and M. Tait, A spectral Erdős-Sós theorem, *SIAM J. Discrete Math.* 37 (2023) 2228-2239.
- [7] D. Desai, L. Kang, Y. Li, Z. Ni, M. Tait and J. Wang, Spectral extremal graphs for intersecting cliques, *Linear Algebra Appl.* 644 (2022) 234-258.
- [8] L. Fang, H. Lin, J. Shu and Z. Zhang, Spectral extremal results on trees, arXiv:2401.05786v2.
- [9] L. Fang, M. Tait and M. Zhai, Turán numbers for non-bipartite graphs and applications to spectral extremal problems, arXiv:2404.09069v1.
- [10] D. Ismailescu and D. Stefanica, Minimizer graphs for a class of extremal problems, *J. Graph Theory* 39 (2002) 230-240.
- [11] X. Lei and S. Li, Spectral extremal problem on disjoint color-critical graphs, *Electron. J. Combin.* 31 (2024) # 1.25.
- [12] Y. Li, and Y. Peng, Refinement on spectral Turán's theorem, *SIAM J. Discrete Math.* 37 (2023) 2462-2485.
- [13] Y. Li, and Y. Peng, The spectral radius of graphs with no intersecting odd cycles, *Discrete Math.* 345 (2022) 112907.
- [14] H. Lin, M. Zhai and Y. Zhao, Spectral radius, edge-disjoint cycles and cycles of the same length, *Electron. J. Combin.* 29 (2022) # 2.1.

- [15] Z. Ni, J. Wang and L. Kang, Spectral extremal graphs for disjoint cliques, *Electron. J. Combin.* 30 (2023) #1.20.
- [16] V. Nikiforov, The spectral radius of graphs without paths and cycles of specified length, *Linear Algebra Appl.* 432 (2010) 2243-2256.
- [17] V. Nikiforov, Bounds on graph eigenvalues II, *Linear Algebra Appl.* 427 (2007) 183-189.
- [18] J. Wang, L. Kang and Y. Xue, On a conjecture of spectral extremal problems, *J. Combin. Theory Ser. B* 159 (2023) 20-41.
- [19] M. Zhai and H. Lin, Spectral extrema of graphs: Forbidden hexagon, *Discrete Math.* 343 (2020) 112028.
- [20] W. Zhang, The spectral radius, maximum average degree and consecutive lengths of graphs, *Graphs Combin.* 40 (2024) 32.