

# ON A LEVEL ANALOG OF SELBERG'S RESULT ON $S(t)$

QINGFENG SUN AND HUI WANG

ABSTRACT. Let  $S(t, f) = \pi^{-1} \arg L(1/2 + it, f)$ , where  $f$  is a holomorphic Hecke cusp form of weight 2 and prime level  $q$ . In this paper, we establish an asymptotic formula for the moments of  $S(t, f)$  without assuming the GRH.

## 1. INTRODUCTION

For  $T > 0$ , let  $N(T)$  be the number of zeros  $\rho = \beta + i\gamma$  of the Riemann zeta function  $\zeta(s)$  in the rectangle  $0 < \beta < 1$ ,  $0 < \gamma \leq T$ . In particular, Davenport [2, Chapter 15] gives that the approximate formula of  $N(T)$  is

$$N(T) = \frac{T}{2\pi} \log \frac{T}{2\pi} - \frac{T}{2\pi} + \frac{7}{8} + S(T) + O(T^{-1}),$$

where

$$S(T) = \frac{1}{\pi} \arg \zeta\left(\frac{1}{2} + iT\right).$$

The argument of  $\zeta(1/2 + iT)$  is defined as the continuous variation along the horizontal segment from  $\infty + iT$  to  $1/2 + iT$  starting at  $\infty + iT$  with the value 0. Understanding the complicated behavior of  $S(t)$  is an interesting topic in analytic number theory, and a series of results have emerged in recent years due to Selberg's pioneering work.

In [12, 13], Selberg showed that for any  $n \in \mathbb{N}$  one has

$$\int_T^{2T} S(t)^{2n} dt = \frac{(2n)!}{n!(2\pi)^{2n}} T (\log \log T)^n + O(T (\log \log T)^{n-1/2}), \quad (1.1)$$

which can imply, on average sense,  $|S(t)|$  has order of magnitude  $\sqrt{\log \log T}$ . The same method can be applied to get a similar result relating to Dirichlet  $L$ -functions  $L(s, \chi)$ , where  $\chi$  is the primitive Dirichlet character of modulus  $q$ . More precisely, for prime modulus  $q$  and  $t > 0$ , Selberg [14] showed that

$$\sum_{\chi \bmod q}^* S(t, \chi)^{2n} = \frac{(2n)!}{n!(2\pi)^{2n}} q (\log \log q)^n + O(q (\log \log q)^{n-1/2}), \quad (1.2)$$

where  $*$  denotes the summation goes through the primitive characters  $\chi(\bmod q)$ , and  $S(t, \chi)$  is defined by

$$S(t, \chi) = \frac{1}{\pi} \arg L\left(\frac{1}{2} + it, \chi\right).$$

---

2020 *Mathematics Subject Classification.* 11F12, 11F66, 11F72.

*Key words and phrases.* Moments, holomorphic Hecke cusp forms, Selberg's limit theorem.

Q. Sun was partially supported by the National Natural Science Foundation of China (Grant Nos. 11871306 and 12031008).

Note that the counterparts of (1.1) and (1.2) for the odd-order moments can also be obtained by making a minor adaptation in Selberg's techniques.

For higher rank  $L$ -functions, there are some analogues of (1.2) in the spectral and weight aspects, respectively. For example, under Generalized Riemann Hypothesis (GRH), Hejhal and Luo [4] (resp. Liu and Liu [10]) obtained an asymptotic formula for the spectral moments of  $S_F(t) = \pi^{-1} \arg L(\frac{1}{2} + it, F)$  for each fixed  $t > 0$ , where  $F$  is the Hecke–Maass cusp form for  $\mathrm{SL}_2(\mathbb{Z})$  (resp.  $\mathrm{SL}_3(\mathbb{Z})$ ) with eigenvalue  $\lambda_F = 1/4 + t_F^2$  (resp. the spectral parameter  $\nu_F = (\nu_{F,1}, \nu_{F,2}, \nu_{F,3})$ ). Recently, when  $f$  is a holomorphic Hecke cusp form of  $\mathrm{SL}_2(\mathbb{Z})$  with weight  $k$ , Liu and Shim [11] unconditionally establish the asymptotic formula for the moments of  $S(t, f) = \frac{1}{\pi} \arg L(\frac{1}{2} + it, f)$ , by means of the weighted version of the zero-density estimate obtained by Hough [5] in the weight aspect.

In this paper, we are devoted to get an unconditional analog for  $\mathrm{GL}_2$  in the level aspect without assuming the GRH. Assume  $q$  is a fixed large prime. Let  $S_2(q)$  be the space of holomorphic cusp forms of weight 2 and level  $q$  equipped with the Petersson inner product

$$\langle f, g \rangle = \int_{\Gamma_0(q) \backslash \mathbb{H}} f(z) \overline{g(z)} \frac{dx dy}{y^2}.$$

Here  $\Gamma_0(q) \backslash \mathbb{H}$  is a fundamental domain of the action of the Hecke congruence subgroup  $\Gamma_0(q)$  on the upper-half plane  $\mathbb{H}$ . Let  $H_2(q)$  be a basis of  $S_2(q)$  that are eigenfunctions of all the Hecke operators and have the first Fourier coefficient  $\lambda_f(1) = 1$ . For each  $f \in H_2(q)$ , let

$$f(z) = \sum_{n=1}^{\infty} n^{1/2} \lambda_f(n) e(nz), \quad z \in \mathbb{H}$$

be its Fourier expansion. The Fourier coefficients  $\lambda_f(n)$  are real and satisfy the Deligne's bound (in particular, this is due to Eichler–Shimura [15] in the case of weight 2)

$$|\lambda_f(n)| \leq \tau(n), \tag{1.3}$$

with  $\tau(n)$  being the divisor function. For  $\mathrm{Re}(s) > 1$ , the Hecke  $L$ -function associated to  $f \in H_2(q)$  is defined by

$$L(s, f) = \sum_{n=1}^{\infty} \frac{\lambda_f(n)}{n^s} = \prod_p (1 - \lambda_f(p)p^{-s} + \varepsilon_q(p)p^{-2s})^{-1},$$

where  $\varepsilon_q(\cdot)$  is the principal Dirichlet character modulo  $q$ . It can be analytically continued to the whole complex plane and has the completed  $L$ -function

$$\Lambda(s, f) = \left( \frac{\sqrt{q}}{2\pi} \right)^s \Gamma\left(s + \frac{1}{2}\right) L(s, f),$$

which satisfies the functional equation

$$\Lambda(s, f) = \epsilon_f \Lambda(1 - s, f),$$

where  $\epsilon_f = q^{1/2} \lambda_f(q) \in \{\pm 1\}$  is the sign of the functional equation. Moreover, we have the following multiplicative property of the coefficients  $\lambda_f(n)$ ,

$$\lambda_f(m) \lambda_f(n) = \sum_{d|(m,n)} \varepsilon_q(d) \lambda_f\left(\frac{mn}{d^2}\right), \quad \text{for } m, n \geq 1. \tag{1.4}$$

If  $p$  is a prime  $\neq q$ , we write  $1 - \lambda_f(p)Y + Y^2 = (1 - \alpha_f(p)Y)(1 - \beta_f(p)Y)$ , so

$$\lambda_f(p) = \alpha_f(p) + \beta_f(p).$$

The bound (1.3) is equivalent (for  $n$  coprime with the level) with the assertion that  $|\alpha_f(p)| = 1$  for all  $p \neq q$ . For  $p = q$ , the  $p$ -factor of  $L(s, f)$  is of degree at most 1, and we let  $\alpha_f(p) = \lambda_f(p)$ , which is shown to be of modulus at most 1, and  $\beta_f(p) = 0$ .

Now we define  $S(t, f)$  by

$$S(t, f) := \frac{1}{\pi} \arg L(1/2 + it, f),$$

where the argument  $\arg L(1/2 + it, f)$  is obtained by continuous variation along the line  $\text{Im}(s) = t$  from  $\sigma = +\infty$  to  $\sigma = 1/2$ . Our main result is the following theorem.

**Theorem 1.1.** *Let  $t > 0$  and  $n \in \mathbb{N}$  be given. For sufficiently large prime number  $p$ , we have*

$$\sum_{f \in H_2(q)} \frac{S(t, f)^n}{4\pi \langle f, f \rangle} = C_n (\log \log q)^{n/2} + O_{t,n}((\log \log q)^{(n-1)/2}),$$

where  $C_n$  is defined by

$$C_n = \begin{cases} \frac{n!}{(n/2)!(2\pi)^n}, & \text{if } n \text{ is even,} \\ 0, & \text{if } n \text{ is odd.} \end{cases}$$

**Remark 1.** For a holomorphic Hecke cusp form  $f$  with weight 2 and level prime number  $q$ , Theorem 1.1 and Lemma 2.3 indicate that  $|S(t, f)|$  on average has order of magnitude  $\sqrt{\log \log q}$ .

It is another interesting problem to consider the following probability measure  $\mu_q$  on  $\mathbb{R}$ , defined by

$$\mu_q(E) := \left( \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \mathbf{1}_E \left( \frac{S(t, f)}{\sqrt{\log \log q}} \right) \right) / \left( \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \right),$$

where  $\mathbf{1}_E$  is the characteristic function on a Borel measurable set  $E$  in  $\mathbb{R}$ . As a consequence of Theorem 1.1, we obtain the following weighted central limit theorem.

**Theorem 1.2.** *As  $q \rightarrow \infty$ , the probability measure  $\mu_q$  converges to the Gaussian distribution of mean 0 and variance  $(2\pi^2)^{-1}$ ; that is*

$$\lim_{q \rightarrow \infty} \mu_q([a, b]) = \sqrt{\pi} \int_a^b \exp(-\pi^2 \xi^2) d\xi$$

for any  $a < b$ .

**Notation.** Throughout the paper,  $\varepsilon$  and  $A$  are arbitrarily small and arbitrarily large positive real numbers, which may be different at each occurrence. As usual,  $e(z) = \exp(2\pi iz) = e^{2\pi iz}$ . We write  $f = O(g)$  or  $f \ll g$  to mean  $|f| \leq cg$  for some unspecified positive constant  $c$ . The sign  $\bar{a}$  denotes the multiplicative inverse of  $a \pmod{r}$ .

## 2. PRELIMINARIES

We firstly collect some classical results as the beginning of this section.

**Lemma 2.1.** *Let  $x \geq 2$ .*

(1) *There exist constants  $c_1$  and  $c_2$ , such that*

$$\sum_{p \leq x} \frac{1}{p} = \log \log x + c_1 + O(e^{c_2 \sqrt{\log x}}). \quad (2.1)$$

(2) *We have*

$$\sum_{p \leq x} \frac{\log p}{p} = \log x + O(1),$$

and

$$\sum_{p \leq x} \frac{(\log p)^2}{p} \ll (\log x)^2. \quad (2.2)$$

The Petersson trace formula is given by the following basic orthogonality relation on  $H_2(q)$  (see e.g., Kowalski–Michel [8]).

**Lemma 2.2.** *Let  $m, n \geq 1$ . Then*

$$\sum_{f \in H_2(q)} \omega_f \cdot \lambda_f(m) \lambda_f(n) = \delta_{m,n} - 2\pi \sum_{c \equiv 0 \pmod{q}} \frac{S(m, n; c)}{c} J_1 \left( \frac{4\pi \sqrt{mn}}{c} \right),$$

where  $J_1$  is the Bessel function and  $S(m, n; c)$  is a classical Kloosterman sum,

$$\omega_f := \frac{1}{4\pi \langle f, f \rangle} = \frac{\zeta(2)}{|H_2(q)|} \cdot \frac{1}{L(1, \text{sym}^2 f)} \cdot \left( 1 + O \left( \frac{(\log q)^4}{q} \right) \right)$$

is the harmonic weight of  $f$ .

We record here some properties of Bessel functions will be used. The Bessel function of the first kind,  $J_\nu(x)$  with  $\nu \in \mathbb{C}$ , has Taylor series about zero given by (see [3, pg. 918, Eq. 8.440])

$$J_\nu(x) = \left( \frac{x}{2} \right)^\nu \sum_{m=0}^{\infty} \frac{(-1)^m}{m! \Gamma(\nu + 1 + m)} \left( \frac{x}{2} \right)^{2m}.$$

Specializing to  $\nu = k - 1$  with  $k \geq 1$ , we have the following bound for  $x > 0$ ,

$$J_{k-1}(x) \ll_k \begin{cases} x^{k-1}, & \text{if } x \leq 1, \\ x^{-1/2}, & \text{if } x > 1. \end{cases} \quad (2.3)$$

By a general bound for Kloosterman sums (see e.g., [7, (2.13)])

$$|S(m, n; c)| \leq (m, n, c) \min \left\{ \frac{c}{(n, c)}, \frac{c}{(m, c)} \right\}^{\frac{1}{2}} \tau(c),$$

and the bound  $J_1(x) \ll x$  for  $J$ -Bessel function, we derive (see e.g., [7, Corollary 2.3])

**Lemma 2.3.** *For any integers  $m, n \geq 1$  with  $6\pi\sqrt{mn} \leq q$ , we have*

$$\sum_{f \in H_2(q)} \omega_f \cdot \lambda_f(m) \lambda_f(n) = \delta_{m,n} + O \left( q^{-3/2} \frac{(m, n, q)(mn)^{1/2}}{((m, q) + (n, q))^{1/2}} (\tau(m, n)) \right).$$

We will need the following weighted version of zero-density estimate which was established by Kowalski–Michel [9, Theorem 4].

**Lemma 2.4.** *Let  $q$  be a prime number and*

$$N(f; \sigma, T) := \#\{\rho = \beta + i\gamma \mid L(\rho, f) = 0, \sigma < \beta, |\gamma| < T\}.$$

*There exists an absolute constant  $A > 0$  such that for any real number  $T$  with  $T \geq \frac{1}{\log q}$ , for any  $\frac{1}{2} + \frac{1}{\log q} \leq \sigma \leq 1$ , and for any  $c$ ,  $0 < c < \frac{1}{2}$ , one has*

$$\sum_{f \in H_2(q)} \omega_f \cdot N(f; \sigma, T) \ll (1 + T)^A q^{-2c(\sigma-1/2)} (\log q) T,$$

where the implied constant depends only on  $c$ .

Finally, we need the lemma of the theory of moments in probability theory (see, for example, [1, Theorem 30.2]).

**Lemma 2.5.** *Suppose that the distribution of random variable  $X$  is determined by its moments, that the  $X_n$  have moments of all orders, and that  $\lim_n E[X_n^r] = E[X^r]$  for  $r = 1, 2, \dots$ . Then  $X_n \Rightarrow X$ .*

### 3. AN APPROXIMATION OF $S(t, f)$

In this section we will prove several lemmas and derive an approximation of  $S(t, f)$ . We denote by  $\rho = \beta + i\gamma$  a typical zero of  $L(s, f)$  inside the critical strip, i.e.,  $0 < \beta < 1$ . For  $\text{Re}(s) > 1$ , we have the Dirichlet series expansion for the logarithmic derivative of  $L(s, f)$ ,

$$-\frac{L'}{L}(s, f) = \sum_{n=1}^{\infty} \frac{\Lambda(n) C_f(n)}{n^s}, \quad (3.1)$$

where  $\Lambda(n)$  denotes the von Mangoldt function, and

$$C_f(n) = \begin{cases} \alpha_f(p)^m + \beta_f(p)^m, & \text{if } n = p^m, \\ 0, & \text{otherwise.} \end{cases} \quad (3.2)$$

**Lemma 3.1.** *Let  $x > 1$ . For  $s \neq \rho$ , and  $s \neq -m - \frac{1}{2}$ , we have the following identity*

$$\begin{aligned} \frac{L'}{L}(s, f) = & - \sum_{n \leq x^3} \frac{C_f(n) \Lambda_x(n)}{n^s} - \frac{1}{\log^2 x} \sum_{\rho} \frac{x^{\rho-s} (1 - x^{\rho-s})^2}{(\rho - s)^3} \\ & - \frac{1}{\log^2 x} \sum_{m=0}^{\infty} \frac{x^{-m-\frac{1}{2}-s} (1 - x^{-m-\frac{1}{2}-s})^2}{(-m - \frac{1}{2} - s)^3}, \end{aligned}$$

where

$$\Lambda_x(n) = \begin{cases} \Lambda(n), & \text{if } n \leq x, \\ \Lambda(n) \frac{\log^2(x^3/n) - 2 \log^2(x^2/n)}{2 \log^2 x}, & \text{if } x \leq n \leq x^2, \\ \Lambda(n) \frac{\log^2(x^3/n)}{2 \log^2 x}, & \text{if } x^2 \leq n \leq x^3, \\ 0, & \text{if } n \geq x^3. \end{cases}$$

*Proof.* Recall the discontinuous integral

$$\frac{1}{2\pi i} \int_{(\alpha)} \frac{y^s}{s^3} ds = \begin{cases} \frac{\log^2 y}{2}, & \text{if } y \geq 1, \\ 0, & \text{if } 0 < y \leq 1 \end{cases}$$

for  $\alpha > 0$ . It follows from (3.1) and (3.2) that

$$-\log^2 x \sum_{n=1}^{\infty} \frac{C_f(n)\Lambda_x(n)}{n^s} = \frac{1}{2\pi i} \int_{(\alpha)} \frac{x^u(1-x^u)^2}{u^3} \frac{L'}{L}(s+u, f) du,$$

where  $\alpha = \max\{2, 1 + \operatorname{Re}(s)\}$ . By moving the line of integration all way to the left, we pick up the residues at  $u = 0$ ,  $u = \rho - s$  and  $u = -m - \frac{1}{2} - s$  ( $m = 0, 1, 2, \dots$ ), and deduce that

$$\begin{aligned} & \frac{1}{2\pi i} \int_{(\alpha)} \frac{x^u(1-x^u)^2}{u^3} \frac{L'}{L}(s+u, f) du \\ &= \frac{L'}{L}(s, f) \log^2 x + \sum_{\rho} \frac{x^{\rho-s}(1-x^{\rho-s})^2}{(\rho-s)^3} + \sum_{m=0}^{\infty} \frac{x^{-m-\frac{1}{2}-s}(1-x^{-m-\frac{1}{2}-s})^2}{(-m-\frac{1}{2}-s)^3}. \end{aligned}$$

Thus we complete the proof of lemma.  $\square$

**Lemma 3.2.** For  $s = \sigma + it$ ,  $s' = \sigma' + it'$  such that  $1/2 \leq \sigma, \sigma' \leq 10$ ,  $s \neq \rho$ ,  $s' \neq \rho$ , we have

$$\operatorname{Im} \left( \frac{L'}{L}(s, f) - \frac{L'}{L}(s', f) \right) = \operatorname{Im} \sum_{\rho} \left( \frac{1}{s-\rho} - \frac{1}{s'-\rho} \right) + O(1),$$

and

$$\operatorname{Re} \frac{L'}{L}(s, f) = \sum_{\rho} \frac{\sigma - \beta}{(\sigma - \beta)^2 + (t - \gamma)^2} + O(\log(|t| + q)). \quad (3.3)$$

*Proof.* By the Hadamard factorization theorem of the entire function  $\Lambda(s, f)$ , we have

$$\frac{L'}{L}(s, f) = b(f) + \sum_{\rho} \left( \frac{1}{s-\rho} + \frac{1}{\rho} \right) - \frac{\Gamma'(s+1/2)}{\Gamma(s+1/2)} - \frac{1}{2} \log q + \log(2\pi),$$

for some  $b \in \mathbb{C}$  with  $\operatorname{Re}(b(f)) = -\operatorname{Re} \sum_{\rho} \frac{1}{\rho}$  (see [6, Proposition 5.7 (3)]). Note that for  $z \notin -\mathbb{N}$ ,

$$\frac{\Gamma'}{\Gamma}(z) = -\gamma + \sum_{k \geq 1} \left( \frac{1}{k} - \frac{1}{k-1+z} \right), \quad (3.4)$$

where  $\gamma$  is the Euler constant. Using (3.4), we get that for  $\frac{1}{4} \leq u$ ,

$$\operatorname{Im} \frac{\Gamma'}{\Gamma}(u+iv) = \sum_{k \geq 1} \frac{v}{(k+u-1)^2 + v^2} \ll 1.$$

And for  $|\arg z| \leq \pi - \varepsilon$ ,

$$\frac{\Gamma'}{\Gamma}(z) = \log z - \frac{1}{2z} + O_{\varepsilon} \left( \frac{1}{|z|^2} \right),$$

thus we complete the proof of lemma.  $\square$

Let  $x \geq 4$ , we define

$$\sigma_x = \sigma_{x,f} = \sigma_{x,f,t} := \frac{1}{2} + 2 \max \left\{ \left| \beta - \frac{1}{2} \right|, \frac{5}{\log x} \right\}, \quad (3.5)$$

where  $\rho = \beta + i\gamma$  runs through the zeros of  $L(s, f)$  for which

$$|t - \gamma| \leq \frac{x^{3|\beta-1/2|}}{\log x}. \quad (3.6)$$

**Lemma 3.3.** *Let  $x \geq 4$ . For  $\sigma \geq \sigma_x$ , we have*

$$\frac{L'}{L}(\sigma + it, f) = - \sum_{n \leq x^3} \frac{C_f(n)\Lambda_x(n)}{n^{\sigma+it}} + O \left( x^{1/4-\sigma/2} \left| \sum_{n \leq x^3} \frac{C_f(n)\Lambda_x(n)}{n^{\sigma_x+it}} \right| \right) + O \left( x^{1/4-\sigma/2} \log(|t| + q) \right),$$

and

$$\sum_{\rho} \frac{\sigma_x - 1/2}{(\sigma_x - \beta)^2 + (t - \gamma)^2} = O \left( \left| \sum_{n \leq x^3} \frac{C_f(n)\Lambda_x(n)}{n^{\sigma_x+it}} \right| \right) + O(\log(|t| + q)).$$

*Proof.* This proof follows [11, Lemma 4.3] closely, and we give the details here for completeness. By (3.3), we have

$$\operatorname{Re} \frac{L'}{L}(\sigma_x + it, f) = \sum_{\rho} \frac{\sigma_x - \beta}{(\sigma_x - \beta)^2 + (t - \gamma)^2} + O(\log(|t| + q)). \quad (3.7)$$

Moreover, if  $\rho = \beta + i\gamma$  is zero of  $L(s, f)$ , then  $(1 - \beta) + i\gamma$  is also a zero of  $L(s, f)$ . Thus we have

$$\begin{aligned} & \sum_{\rho} \frac{\sigma_x - \beta}{(\sigma_x - \beta)^2 + (t - \gamma)^2} \\ &= \frac{1}{2} \left( \sum_{\rho} \frac{\sigma_x - \beta}{(\sigma_x - \beta)^2 + (t - \gamma)^2} + \sum_{\rho} \frac{\sigma_x - (1 - \beta)}{(\sigma_x - (1 - \beta))^2 + (t - \gamma)^2} \right) \\ &= \left( \sigma_x - \frac{1}{2} \right) \sum_{\rho} \frac{(\sigma_x - 1/2)^2 - (\beta - 1/2)^2 + (t - \gamma)^2}{((\sigma_x - \beta)^2 + (t - \gamma)^2)((\sigma_x - 1 + \beta)^2 + (t - \gamma)^2)}. \end{aligned} \quad (3.8)$$

If  $|\beta - 1/2| \leq \frac{\sigma_x - 1/2}{2}$ , then

$$\begin{aligned} (\sigma_x - 1/2)^2 - (\beta - 1/2)^2 &\geq \frac{1}{2}((\sigma_x - 1/2)^2 + (\beta - 1/2)^2) \\ &= \frac{1}{4}((\sigma_x - \beta)^2 + (\sigma_x - 1 + \beta)^2). \end{aligned}$$

Thus,

$$(\sigma_x - 1/2)^2 - (\beta - 1/2)^2 + (t - \gamma)^2 \geq \frac{1}{4}((\sigma_x - 1 + \beta)^2 + (t - \gamma)^2). \quad (3.9)$$

If  $|\beta - 1/2| > \frac{\sigma_x - 1/2}{2}$ , then by (3.5) and (3.6) we have

$$|t - \gamma| > \frac{x^{3|\beta-1/2|}}{\log x} > 3|\beta - 1/2|.$$

Thus,

$$\begin{aligned}
& (\sigma_x - 1/2)^2 - (\beta - 1/2)^2 + (t - \gamma)^2 \\
&= ((\sigma_x - 1/2)^2 + (\beta - 1/2)^2) + (t - \gamma)^2 - 2(\beta - 1/2)^2 \\
&\geq \frac{1}{2}((\sigma_x - \beta)^2 + (\sigma_x - 1 + \beta)^2) + \frac{7}{9}(t - \gamma)^2 \\
&\geq \frac{1}{4}((\sigma_x - 1 + \beta)^2 + (t - \gamma)^2).
\end{aligned} \tag{3.10}$$

Combining (3.9), (3.10) and (3.8), we have

$$\sum_{\rho} \frac{\sigma_x - \beta}{(\sigma_x - \beta)^2 + (t - \gamma)^2} \geq \frac{1}{4}(\sigma_x - 1/2) \sum_{\rho} \frac{1}{((\sigma_x - \beta)^2 + (t - \gamma)^2)}.$$

Using this bound and (3.7), we obtain

$$\sum_{\rho} \frac{1}{((\sigma_x - \beta)^2 + (t - \gamma)^2)} \leq \frac{4}{\sigma_x - 1/2} \left| \frac{L'}{L}(\sigma_x + it, f) \right| + O\left(\frac{\log(|t| + q)}{\sigma_x - 1/2}\right). \tag{3.11}$$

On the other hand, by Lemma 3.1, we have

$$\begin{aligned}
\frac{L'}{L}(\sigma + it, f) &= - \sum_{n \leq x^3} \frac{C_f(n) \Lambda_x(n)}{n^{\sigma+it}} + \frac{w(x, \sigma, t)}{\log^2 x} \sum_{\rho} \frac{x^{\beta-\sigma}(1 + x^{\beta-\sigma})^2}{((\sigma - \beta)^2 + (t - \gamma)^2)^{3/2}} \\
&\quad + O\left(\frac{x^{-1/2-\sigma}}{\log^2 x}\right),
\end{aligned} \tag{3.12}$$

with  $|w(x, \sigma, t)| \leq 1$ . Next we claim that

$$\frac{x^{\beta-\sigma}(1 + x^{\beta-\sigma})^2}{((\sigma - \beta)^2 + (t - \gamma)^2)^{3/2}} \leq 2 \log x \frac{x^{1/4-\sigma/2}}{(\beta - \sigma_x)^2 + (t - \gamma)^2}. \tag{3.13}$$

If  $\beta \leq \frac{\sigma_x + 1/2}{2}$ , then we have

$$\begin{aligned}
\frac{x^{\beta-\sigma}(1 + x^{\beta-\sigma})^2}{((\sigma - \beta)^2 + (t - \gamma)^2)^{3/2}} &\leq \frac{4x^{1/4-\sigma/2}}{(\sigma_x - \beta)((\sigma_x - \beta)^2 + (t - \gamma)^2)} \\
&\leq \frac{8}{\sigma_x - 1/2} \frac{x^{1/4-\sigma/2}}{((\sigma_x - \beta)^2 + (t - \gamma)^2)} \\
&\leq \frac{4}{5} \log x \frac{x^{1/4-\sigma/2}}{((\sigma_x - \beta)^2 + (t - \gamma)^2)}.
\end{aligned}$$

If  $\beta > \frac{\sigma_x + 1/2}{2}$ , then we have

$$|t - \gamma| > \frac{x^{3|\beta-1/2|}}{\log x} > 3|\beta - 1/2| \geq 3|\beta - \sigma_x|.$$

Thus,  $(t - \gamma)^2 > \frac{8}{9}((\beta - \sigma_x)^2 + (t - \gamma)^2)$ . Hence

$$\begin{aligned} \frac{x^{\beta-\sigma}(1+x^{\beta-\sigma})^2}{((\sigma-\beta)^2+(t-\gamma)^2)^{3/2}} &\leq \frac{x^{\beta-\sigma}(1+x^{\beta-1/2})^2}{|t-\gamma|(t-\gamma)^2} \leq \frac{\log x}{x^{3|\beta-1/2|}} \frac{9}{8} \frac{x^{\beta-\sigma}(1+x^{\beta-1/2})^2}{(\beta-\sigma_x)^2+(t-\gamma)^2} \\ &= \frac{9}{8}(\log x)(1+x^{-(\beta-1/2)})^2 \frac{x^{1/2-\sigma}}{(\sigma_x-\beta)^2+(t-\gamma)^2} \\ &\leq \frac{9}{8}(\log x)(1+e^{-5})^2 \frac{x^{1/2-\sigma}}{(\sigma_x-\beta)^2+(t-\gamma)^2} \\ &< 2(\log x) \frac{x^{1/2-\sigma}}{(\sigma_x-\beta)^2+(t-\gamma)^2}. \end{aligned}$$

Using (3.11) and (3.13), we have

$$\begin{aligned} &\sum_{\rho} \frac{x^{\beta-\sigma}(1+x^{\beta-\sigma})^2}{((\sigma-\beta)^2+(t-\gamma)^2)^{3/2}} \\ &\leq 8 \log x \frac{x^{1/4-\sigma/2}}{\sigma_x-1/2} \left| \frac{L'}{L}(\sigma_x+it, f) \right| + O\left( \frac{(\log x)x^{1/4-\sigma/2} \log(|t|+q)}{\sigma_x-1/2} \right) \\ &\leq \frac{4}{5}(\log x)^2 x^{1/4-\sigma/2} \left| \frac{L'}{L}(\sigma_x+it, f) \right| + O\left( (\log x)^2 x^{1/4-\sigma/2} \log(|t|+q) \right). \end{aligned}$$

Inserting this bound into (3.12), we have

$$\begin{aligned} \frac{L'}{L}(\sigma+it, f) &= - \sum_{n \leq x^3} \frac{C_f(n)\Lambda_x(n)}{n^{\sigma+it}} + \frac{4}{5} w'(x, \sigma, t) x^{1/4-\sigma/2} \left| \frac{L'}{L}(\sigma_x+it, f) \right| \\ &\quad + O\left( x^{1/4-\sigma/2} \log(|t|+q) \right), \end{aligned} \tag{3.14}$$

with  $|w'(x, \sigma, t)| \leq 1$ . By taking  $\sigma = \sigma_x$ ,

$$\left( 1 - \frac{4}{5} w'(x, \sigma_x, t) x^{1/4-\sigma_x/2} \right) \frac{L'}{L}(\sigma_x+it, f) = O\left( \left| \sum_{n \leq x^3} \frac{C_f(n)\Lambda_x(n)}{n^{\sigma_x+it}} \right| \right) + O\left( x^{1/4-\sigma_x/2} \log(|t|+q) \right).$$

Since  $|1 - \frac{4}{5} w'(x, \sigma_x, t) x^{1/4-\sigma_x/2}| \geq 1 - |\frac{4}{5} w'(x, \sigma_x, t) x^{1/4-\sigma_x/2}| \geq \frac{1}{5}$ , we have

$$\frac{L'}{L}(\sigma_x+it, f) = O\left( \left| \sum_{n \leq x^3} \frac{C_f(n)\Lambda_x(n)}{n^{\sigma_x+it}} \right| \right) + O\left( x^{1/4-\sigma_x/2} \log(|t|+q) \right). \tag{3.15}$$

Inserting (3.15) into (3.14) and (3.11), respectively, we complete the proof of lemma.  $\square$

The following theorem provides an approximation of  $S(t, f)$ .

**Theorem 3.4.** *For  $t \neq 0$ , and  $x \geq 4$ , we have*

$$\begin{aligned} S(t, f) &= \frac{1}{\pi} \operatorname{Im} \sum_{n \leq x^3} \frac{C_f(n)\Lambda_x(n)}{n^{\sigma_x+it} \log n} + O\left( (\sigma_x-1/2) \left| \sum_{n \leq x^3} \frac{C_f(n)\Lambda_x(n)}{n^{\sigma_x+it}} \right| \right) \\ &\quad + O\left( (\sigma_x-1/2) \log(|t|+q) \right), \end{aligned}$$

where  $\sigma_x$  is defined by (3.5).

*Proof.* We have

$$\begin{aligned}
\pi S(t, f) &= - \int_{1/2}^{\infty} \operatorname{Im} \frac{L'}{L}(\sigma + it, f) d\sigma \\
&= - \int_{\sigma_x}^{\infty} \operatorname{Im} \frac{L'}{L}(\sigma + it, f) d\sigma - (\sigma_x - 1/2) \operatorname{Im} \frac{L'}{L}(\sigma_x + it, f) \\
&\quad + \int_{1/2}^{\sigma_x} \operatorname{Im} \left( \frac{L'}{L}(\sigma_x + it, f) - \frac{L'}{L}(\sigma + it, f) \right) d\sigma \\
&=: J_1 + J_2 + J_3.
\end{aligned}$$

By Lemma 3.3, we have

$$\begin{aligned}
J_1 &= - \int_{\sigma_x}^{\infty} \operatorname{Im} \frac{L'}{L}(\sigma + it, f) d\sigma \\
&= \int_{\sigma_x}^{\infty} \operatorname{Im} \sum_{n \leq x^3} \frac{C_f(n) \Lambda_x(n)}{n^{\sigma+it}} d\sigma + O \left( \left| \sum_{n \leq x^3} \frac{C_f(n) \Lambda_x(n)}{n^{\sigma_x+it}} \right| \int_{\sigma_x}^{\infty} x^{1/4-\sigma/2} d\sigma \right) \\
&\quad + O \left( \log(|t| + q) \int_{\sigma_x}^{\infty} x^{1/4-\sigma/2} d\sigma \right) \\
&= \operatorname{Im} \sum_{n \leq x^3} \frac{C_f(n) \Lambda_x(n)}{n^{\sigma_x+it} \log n} + O \left( \frac{1}{\log x} \left| \sum_{n \leq x^3} \frac{C_f(n) \Lambda_x(n)}{n^{\sigma_x+it}} \right| \right) + O \left( \frac{\log(|t| + q)}{\log x} \right).
\end{aligned}$$

Taking  $\sigma = \sigma_x$  in Lemma 3.3, we have

$$\begin{aligned}
J_2 &\leq (\sigma_x - 1/2) \left| \frac{L'}{L}(\sigma_x + it, f) \right| \\
&\ll (\sigma_x - 1/2) \left| \sum_{n \leq x^3} \frac{C_f(n) \Lambda_x(n)}{n^{\sigma_x+it}} \right| + (\sigma_x - 1/2) \log(|t| + q).
\end{aligned}$$

We apply Lemma 3.2 to bound  $J_3$  and get

$$\begin{aligned}
&\operatorname{Im} \left( \frac{L'}{L}(\sigma_x + it, f) - \frac{L'}{L}(\sigma + it, f) \right) \\
&= \sum_{\rho} \frac{(\sigma_x - \sigma)(\sigma + \sigma_x - 2\beta)(t - \gamma)}{((\sigma_x - \beta)^2 + (t - \gamma)^2)((\sigma - \beta)^2 + (t - \gamma)^2)} + O(1).
\end{aligned}$$

Hence

$$\begin{aligned}
|J_3| &\leq \sum_{\rho} \int_{1/2}^{\sigma_x} \frac{(\sigma_x - \sigma) |\sigma + \sigma_x - 2\beta| |t - \gamma|}{((\sigma_x - \beta)^2 + (t - \gamma)^2)((\sigma - \beta)^2 + (t - \gamma)^2)} d\sigma + O(\sigma_x - 1/2) \\
&\leq \sum_{\rho} \frac{\sigma_x - 1/2}{(\sigma_x - \beta)^2 + (t - \gamma)^2} \int_{1/2}^{\sigma_x} \frac{|\sigma + \sigma_x - 2\beta| |t - \gamma|}{(\sigma - \beta)^2 + (t - \gamma)^2} d\sigma + O(\sigma_x - 1/2).
\end{aligned}$$

If  $|\beta - 1/2| \leq \frac{1}{2}(\sigma_x - 1/2)$ , then for  $1/2 \leq \sigma \leq \sigma_x$ ,

$$\begin{aligned}
|\sigma + \sigma_x - 2\beta| &= |(\sigma - 1/2) + (\sigma_x - 1/2) - 2(\beta - 1/2)| \\
&\leq |\sigma - 1/2| + |\sigma_x - 1/2| + 2|\beta - 1/2| \leq 3|\sigma_x - 1/2|.
\end{aligned}$$

Thus,

$$\begin{aligned} \int_{1/2}^{\sigma_x} \frac{|\sigma + \sigma_x - 2\beta||t - \gamma|}{(\sigma - \beta)^2 + (t - \gamma)^2} d\sigma &\leq 3(\sigma_x - 1/2) \int_{-\infty}^{\infty} \frac{|t - \gamma|}{(\sigma - \beta)^2 + (t - \gamma)^2} d\sigma \\ &\leq 10(\sigma_x - 1/2). \end{aligned}$$

If  $|\beta - 1/2| > \frac{1}{2}(\sigma_x - 1/2)$ , then

$$|t - \gamma| > \frac{x^{3|\beta - 1/2|}}{\log x} > 3|\beta - 1/2|$$

and for  $1/2 \leq \sigma \leq \sigma_x$ ,

$$|\sigma + \sigma_x - 2\beta| \leq |\sigma - 1/2| + |\sigma_x - 1/2| + 2|\beta - 1/2| \leq 3|\sigma_x - 1/2| \leq 6|\beta - 1/2|.$$

Thus,

$$\begin{aligned} \int_{1/2}^{\sigma_x} \frac{|\sigma + \sigma_x - 2\beta||t - \gamma|}{(\sigma - \beta)^2 + (t - \gamma)^2} d\sigma &< \int_{1/2}^{\sigma_x} \frac{|\sigma + \sigma_x - 2\beta|}{|t - \gamma|} d\sigma \\ &\leq \int_{1/2}^{\sigma_x} \frac{6|\beta - 1/2|}{3|\beta - 1/2|} d\sigma = 2(\sigma_x - 1/2). \end{aligned}$$

By Lemma 3.3, we obtain

$$\begin{aligned} |J_3| &\leq 10(\sigma_x - 1/2) \sum_{\rho} \frac{\sigma_x - 1/2}{(\sigma_x - \beta)^2 + (t - \gamma)^2} + O(\sigma_x - 1/2) \\ &= O\left((\sigma_x - 1/2) \left| \sum_{n \leq x^3} \frac{C_f(n)\Lambda_x(n)}{n^{\sigma_x + it}} \right| \right) + O((\sigma_x - 1/2) \log(|t| + q)). \end{aligned}$$

Thus we complete the proof of lemma.  $\square$

**Theorem 3.5.** *Under the GRH, we have*

$$S(t, f) \ll \frac{\log(|t| + q)}{\log \log(|t| + q)}.$$

*Proof.* We have  $\sigma_x = \frac{1}{2} + \frac{10}{\log x}$  by mean of the GRH. By (1.3),

$$|C_f(p^m)\Lambda_x(p^m)| \ll \log p.$$

Thus,

$$\left| \operatorname{Im} \sum_{n \leq x^3} \frac{C_f(n)\Lambda_x(n)}{n^{\sigma_x + it} \log n} \right| \ll \sum_{p \leq x^3} p^{-1/2} \ll \frac{x^{3/2}}{\log x},$$

and

$$(\sigma_x - 1/2) \left| \sum_{n \leq x^3} \frac{C_f(n)\Lambda_x(n)}{n^{\sigma_x + it}} \right| \ll \frac{1}{\log x} \sum_{p \leq x^3} p^{-1/2} \log p \ll \frac{x^{3/2}}{\log x}.$$

Then the theorem follows by taking  $x = (\log(|t| + q))^{2/3}$  in Theorem 3.4.  $\square$

## 4. PROOF OF THEOREMS 1.1 AND 1.2

**Lemma 4.1.** *Let  $t > 0$  be given. For prime number  $q$  sufficiently large and  $x = q^{\delta/3}$  with  $0 < \delta < \frac{6c}{8n+3A+3}$ , we have*

$$\sum_{f \in H_2(q)} \frac{(\sigma_{x,f} - 1/2)^{4n} x^{4n(\sigma_{x,f} - 1/2)}}{4\pi \langle f, f \rangle} \ll_{t,n,\delta,A} \frac{1}{(\log q)^{4n}},$$

where  $\sigma_{x,f}$  is defined by (3.5),  $c$  and  $A$  are as in Lemma 2.4.

*Proof.* By the definition of  $\sigma_{x,f}$ ,

$$\begin{aligned} & (\sigma_{x,f} - 1/2)^{4n} x^{4n(\sigma_{x,f} - 1/2)} \\ & \leq \left( \frac{10}{\log x} \right)^{4n} x^{40n/\log x + 2^{4n+1}} \sum_{\substack{\beta > \frac{1}{2} + \frac{5}{\log x} \\ |t-\gamma| \leq \frac{x^{3|\beta-1/2|}}{\log x}}} (\beta - 1/2)^{4n} x^{8n(\beta-1/2)}. \end{aligned}$$

On the other hand,

$$\begin{aligned} & \sum_{\substack{\beta > \frac{1}{2} + \frac{5}{\log x} \\ |t-\gamma| \leq \frac{x^{3|\beta-1/2|}}{\log x}}} (\beta - 1/2)^{4n} x^{8n(\beta-1/2)} \\ & \leq \sum_{j=5}^{\frac{1}{2}[\log x]} \left( \frac{j+1}{\log x} \right)^{4n} x^{8n \frac{j+1}{\log x}} \sum_{\substack{\frac{1}{2} + \frac{j}{\log x} < \beta \leq \frac{1}{2} + \frac{j+1}{\log x} \\ |t-\gamma| \leq \frac{x^{3|\beta-1/2|}}{\log x}}} 1 \\ & \leq \frac{1}{(\log x)^{4n}} \sum_{j=5}^{\frac{1}{2}[\log x]} (j+1)^{4n} e^{8n(j+1)} N \left( f; \frac{1}{2} + \frac{j}{\log x}, |t| + \frac{e^{3(j+1)}}{\log x} \right). \end{aligned}$$

By Lemma 2.4, there exists an absolute constant  $A > 0$  such that

$$\begin{aligned} & \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \sum_{\substack{\beta > \frac{1}{2} + \frac{5}{\log x} \\ |t-\gamma| \leq \frac{x^{3|\beta-1/2|}}{\log x}}} (\beta - 1/2)^{4n} x^{8n(\beta-1/2)} \\ & \leq \frac{1}{(\log x)^{4n}} \sum_{j=5}^{\frac{1}{2}[\log x]} (j+1)^{4n} e^{8n(j+1)} \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} N \left( f; \frac{1}{2} + \frac{j}{\log x}, |t| + \frac{e^{3(j+1)}}{\log x} \right) \\ & \ll \frac{1}{(\log x)^{4n}} \sum_{j=5}^{\infty} (j+1)^{4n} e^{8n(j+1)} \left( 1 + |t| + \frac{e^{3(j+1)}}{\log x} \right)^A q^{-2c \frac{j}{\log x}} \left( |t| + \frac{e^{3(j+1)}}{\log x} \right) \log q \\ & \ll_{t,n,A,\delta} \frac{\log q}{(\log x)^{4n+A+1}} \sum_{j=5}^{\infty} (j+1)^{4n} e^{(8n+3A+3-\frac{6c}{\delta})j} \\ & \ll_{t,n,A,\delta} \frac{1}{(\log q)^{4n}} \end{aligned}$$

provided that  $0 < \delta < \frac{6c}{8n+3A+3}$ . In addition, by Lemma 2.3,

$$\sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \left( \frac{10}{\log x} \right)^{4n} x^{40n/\log x} \ll \frac{1}{(\log x)^{4n}} \ll \frac{1}{(\log q)^{4n}}.$$

Thus we complete the proof of lemma.  $\square$

For a positive parameter  $x$  (to be chosen later), let

$$M(t, f) := \frac{1}{\pi} \operatorname{Im} \sum_{p \leq x^3} \frac{C_f(p)}{p^{1/2+it}}, \quad \text{and} \quad R(t, f) := S(t, f) - M(t, f),$$

where  $C_f(p) = \alpha_f(p) + \beta_f(p) = \lambda_f(p)$ .

**Lemma 4.2.** *We have*

$$\begin{aligned} R(t, f) &= O \left( \left| \operatorname{Im} \sum_{p \leq x^3} \frac{C_f(p)(\Lambda_x(p) - \Lambda(p))}{p^{1/2+it} \log p} \right| \right) + O \left( \left| \operatorname{Im} \sum_{p \leq x^{3/2}} \frac{C_f(p^2)\Lambda_x(p^2)}{p^{1+2it} \log p} \right| \right) \\ &\quad + O \left( (\sigma_x - 1/2)x^{\sigma_x-1/2} \int_{1/2}^{\infty} x^{1/2-\sigma} \left| \sum_{p \leq x^3} \frac{C_f(p)\Lambda_x(p) \log(xp)}{p^{\sigma+it}} \right| d\sigma \right) \\ &\quad + O((\sigma_x - 1/2) \log(|t| + q)) + O(1). \end{aligned}$$

*Proof.* By Theorem 3.4,

$$\begin{aligned} R(t, f) &= S(t, f) - M(t, f) \\ &= \frac{1}{\pi} \operatorname{Im} \sum_{p \leq x^3} \frac{C_f(p)(\Lambda_x(p)p^{1/2-\sigma_x} - \Lambda(p))}{p^{1/2+it} \log p} + \frac{1}{\pi} \operatorname{Im} \sum_{m=2}^{\infty} \sum_{p^m \leq x^3} \frac{C_f(p^m)\Lambda_x(p^m)}{p^{m(\sigma_x+it)} \log p^m} \\ &\quad + O \left( (\sigma_x - 1/2) \left| \sum_{m=1}^{\infty} \sum_{p^m \leq x^3} \frac{C_f(p^m)\Lambda_x(p^m)}{p^{m(\sigma_x+it)}} \right| \right) + O((\sigma_x - 1/2) \log(|t| + q)). \end{aligned}$$

Using the bound (1.3), we deduce that

$$\sum_{m=3}^{\infty} \sum_{p^m \leq x^3} \frac{C_f(p^m)\Lambda_x(p^m)}{p^{m(\sigma_x+it)}} = O(1), \quad \text{and} \quad \sum_{m=3}^{\infty} \sum_{p^m \leq x^3} \frac{C_f(p^m)\Lambda_x(p^m)}{p^{m(\sigma_x+it)} \log p} = O(1).$$

Note that

$$\begin{aligned} (\sigma_x - 1/2) \left| \sum_{p \leq x^{3/2}} \frac{C_f(p^2)\Lambda_x(p^2)}{p^{2(\sigma_x+it)}} \right| &\ll (\sigma_x - 1/2) \sum_{p \leq x^{3/2}} \frac{\log p}{p} \\ &\ll (\sigma_x - 1/2) \log x \ll (\sigma_x - 1/2) \log q. \end{aligned}$$

Thus,

$$\begin{aligned}
R(t, f) &= \frac{1}{\pi} \operatorname{Im} \sum_{p \leq x^3} \frac{C_f(p)(\Lambda_x(p) - \Lambda(p))}{p^{1/2+it} \log p} - \frac{1}{\pi} \operatorname{Im} \sum_{p \leq x^3} \frac{C_f(p)\Lambda_x(p)(1 - p^{1/2-\sigma_x})}{p^{1/2+it} \log p} \\
&\quad + \frac{1}{\pi} \operatorname{Im} \sum_{p \leq x^{3/2}} \frac{C_f(p^2)\Lambda_x(p^2)}{p^{1+2it} \log p^2} + \frac{1}{\pi} \operatorname{Im} \sum_{p \leq x^{3/2}} \frac{C_f(p^2)\Lambda_x(p^2)(p^{1-2\sigma_x} - 1)}{p^{1+2it} \log p^2} \\
&\quad + O\left((\sigma_x - 1/2) \left| \sum_{p \leq x^3} \frac{C_f(p)\Lambda_x(p)}{p^{\sigma_x+it}} \right| \right) + O((\sigma_x - 1/2) \log(|t| + q)) + O(1).
\end{aligned}$$

Note that for  $1/2 \leq a \leq \sigma_x$ ,

$$\begin{aligned}
\left| \sum_{p \leq x^3} \frac{C_f(p)\Lambda_x(p)}{p^{a+it}} \right| &= x^{a-1/2} \left| \int_a^\infty x^{1/2-\sigma} \sum_{p \leq x^3} \frac{C_f(p)\Lambda_x(p) \log(xp)}{p^{\sigma+it}} d\sigma \right| \\
&\leq x^{\sigma_x-1/2} \int_{1/2}^\infty x^{1/2-\sigma} \left| \sum_{p \leq x^3} \frac{C_f(p)\Lambda_x(p) \log(xp)}{p^{\sigma+it}} \right| d\sigma.
\end{aligned}$$

Thus,

$$\begin{aligned}
\left| \sum_{p \leq x^3} \frac{C_f(p)\Lambda_x(p)(1 - p^{1/2-\sigma_x})}{p^{1/2+it} \log p} \right| &= \left| \int_{1/2}^{\sigma_x} \sum_{p \leq x^3} \frac{C_f(p)\Lambda_x(p)}{p^{a+it}} da \right| \\
&\leq (\sigma_x - 1/2) x^{\sigma_x-1/2} \int_{1/2}^\infty x^{1/2-\sigma} \left| \sum_{p \leq x^3} \frac{C_f(p)\Lambda_x(p) \log(xp)}{p^{\sigma+it}} \right| d\sigma,
\end{aligned}$$

and

$$(\sigma_x - 1/2) \left| \sum_{p \leq x^3} \frac{C_f(p)\Lambda_x(p)}{p^{\sigma_x+it}} \right| \leq (\sigma_x - 1/2) x^{\sigma_x-1/2} \int_{1/2}^\infty x^{1/2-\sigma} \left| \sum_{p \leq x^3} \frac{C_f(p)\Lambda_x(p) \log(xp)}{p^{\sigma+it}} \right| d\sigma.$$

Moreover,

$$\begin{aligned}
\left| \sum_{p \leq x^{3/2}} \frac{C_f(p^2)\Lambda_x(p^2)(p^{1-2\sigma_x} - 1)}{p^{1+2it} \log p} \right| &\ll \sum_{p \leq x^{3/2}} \frac{1}{p} (1 - p^{1-2\sigma_x}) \ll \sum_{p \leq x^{3/2}} (\sigma_x - 1/2) \frac{\log p}{p} \\
&\ll (\sigma_x - 1/2) \log q.
\end{aligned}$$

Thus we complete the proof of lemma.  $\square$

**Proposition 4.3.** *Let  $t > 0$  be given. For prime number  $q$  sufficiently large and  $x = q^{\delta/3}$  with sufficiently small  $\delta > 0$ , we have*

$$\sum_{f \in H_2(q)} \frac{M(t, f)^n}{4\pi \langle f, f \rangle} = C_n (\log \log q)^{n/2} + O_{t,n}((\log \log q)^{n/2-1}), \quad (4.1)$$

and

$$\sum_{f \in H_2(q)} \frac{|R(t, f)|^{2n}}{4\pi \langle f, f \rangle} = O_{t,n}(1). \quad (4.2)$$

Here  $C_n$  is defined by

$$C_n = \begin{cases} \frac{n!}{(n/2)!(2\pi)^n}, & \text{if } n \text{ is even,} \\ 0, & \text{if } n \text{ is odd.} \end{cases}$$

*Proof.* Recall that

$$M(t, f) = \frac{1}{\pi} \operatorname{Im} \sum_{p \leq x^3} \frac{C_f(p)}{p^{1/2+it}} = \frac{-i}{2\pi} \left( \sum_{p \leq x^3} \frac{\lambda_f(p)}{p^{1/2+it}} - \sum_{p \leq x^3} \frac{\lambda_f(p)}{p^{1/2-it}} \right).$$

Set  $x = q^{\delta/3}$  for a suitably small  $\delta > 0$  (to be specified later). Thus we have

$$M(t, f)^n = \frac{(-i)^n}{(2\pi)^n} \left( \sum_{p \leq q^\delta} \frac{\lambda_f(p)}{p^{1/2+it}} - \sum_{p \leq q^\delta} \frac{\lambda_f(p)}{p^{1/2-it}} \right)^n.$$

A general term in the expansion of the above has the form

$$\frac{\lambda_f(p_1)^{m(p_1)+n(p_1)}}{p_1^{m(p_1)(1/2+it)} (-1)^{n(p_1)} p_1^{n(p_1)(1/2-it)}} \times \cdots \times \frac{\lambda_f(p_r)^{m(p_r)+n(p_r)}}{p_r^{m(p_r)(1/2+it)} (-1)^{n(p_r)} p_r^{n(p_r)(1/2-it)}}, \quad (4.3)$$

where  $p_1 < p_2 < \cdots < p_r \leq q^\delta$ ,  $m(p_j) + n(p_j) \geq 1$ , and  $\sum_{j=1}^r (m(p_j) + n(p_j)) = n$ .

Now we discuss the contribution from the general term (4.3) in the following cases.

Case (I): In the general term (4.3),  $m(p_{j_0}) \not\equiv n(p_{j_0}) \pmod{2}$  for some  $j_0$ . With the Hecke relation (1.4), the numerator of this general term (4.3) can be written as a sum whose terms are all of the form  $\lambda_f(m)\lambda_f(n)$  where  $m \neq n$ . Since for these terms  $\delta_{m,n} = 0$ , we see from Lemma 2.3 that the contribution from the general term (4.3) to the moment (4.1) is

$$\begin{aligned} & \ll_n \sum_{p_1 < p_2 < \cdots < p_r \leq q^\delta} p_1^{-\frac{1}{2}(m(p_1)+n(p_1))} \cdots p_r^{-\frac{1}{2}(m(p_r)+n(p_r))} q^{-\frac{3}{2} + \frac{\delta n}{2}(1+\varepsilon)} \\ & \ll_n \left( \sum_{p \leq q^\delta} p^{-\frac{1}{2}} \right)^r q^{-\frac{3}{2} + \frac{\delta n}{2}(1+\varepsilon)} \ll_n q^{-\frac{3}{2} + \frac{\delta n}{2}(2+\varepsilon)}, \end{aligned}$$

which is negligible by taking  $0 < \delta < \frac{1}{n}$ .

Case (II): In the general term (4.3),  $m(p_j) \equiv n(p_j) \pmod{2}$  for all  $j$ , and  $m(p_{j_0}) + n(p_{j_0}) \geq 4$  for some  $j_0$ . In this case,  $n$  is even and  $n > 2r$ , so  $r < \frac{n}{2}$ . By using Hecke relations (1.4), one sees that it is possible to have resulting terms  $\lambda_f(m)\lambda_f(n)$  with  $m = n$ . Using Lemma 2.3, the contribution to the moment (4.1) such terms is bounded by the diagonal term of size  $\asymp 1$ . The contribution from the off-diagonal terms is clearly negligible compared to 1, given that  $0 < \delta < \frac{1}{n}$ . Combining the bound (1.3) and (2.1), we have the contribution from this general

term to the moment (4.1) is

$$\begin{aligned} & \ll_n \sum_{p_1 < p_2 < \dots < p_r \leq q^\delta} p_1^{-\frac{1}{2}(m(p_1)+n(p_1))} \dots p_r^{-\frac{1}{2}(m(p_r)+n(p_r))} \\ & \ll_n \left( \sum_{p \leq q^\delta} p^{-1} \right)^r \ll_n (\log \log q)^{\frac{n}{2}-1}. \end{aligned}$$

It remains to discuss the following: In the general term (4.3),  $m(p_j) \equiv n(p_j) \pmod{2}$  and  $m(p_j) + n(p_j) = 2$  for all  $j$ . Clearly  $n$  must be even, say  $n = 2m$  and so  $r = m$ .

Case (III):  $m(p_{j_0}) = 2$  or  $n(p_{j_0}) = 2$  for some  $j_0$ . We deduce that the contribution from these terms to (4.1) is at most  $O((\log \log q)^{\frac{n}{2}-1})$  by using (1.3) and the convergence of  $\sum_p p^{-1-it}$  for  $t \neq 0$ .

Case (IV):  $m(p_j) = n(p_j) = 1$  for all  $j$ . The contribution from these terms to (4.1) is

$$\sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \binom{2m}{m} (m!) (m!) \frac{(-1)^m (-i)^{2m}}{(2\pi)^{2m}} \sum_{p_1 < p_2 < \dots < p_m \leq q^\delta} \frac{\lambda_f(p_1)^2 \dots \lambda_f(p_m)^2}{p_1 \dots p_m}.$$

By the Hecke relations, we have  $\lambda_f(p_1)^2 \dots \lambda_f(p_m)^2 = \lambda_f(p_1 \dots p_m) \lambda_f(p_1 \dots p_m)$ . Applying Lemma 2.3, we have the above equals

$$\begin{aligned} & \frac{(2m)!}{(2\pi)^{2m}} \sum_{p_1 < p_2 < \dots < p_m \leq q^\delta} \frac{1}{p_1 \dots p_m} \cdot \left( 1 + O\left( q^{-3/2} (p_1 \dots p_m) (\tau(p_1 \dots p_m)) \right) \right) \\ & = \frac{(2m)!}{m! (2\pi)^{2m}} \sum_{\substack{p_1, p_2, \dots, p_m \leq q^\delta \\ p_j \text{ distinct}}} \frac{1}{p_1 \dots p_m} + O\left( q^{-3/2} \sum_{\substack{p_1, p_2, \dots, p_m \leq q^\delta \\ p_j \text{ distinct}}} 1 \right) \\ & = \frac{(2m)!}{m! (2\pi)^{2m}} (\log \log q)^m + O_m((\log \log q)^{m-1}). \end{aligned}$$

Now the asymptotic formula (4.1) follows from Cases (I)–(IV).

Next we are ready to prove (4.2). Recall that  $\sigma_x = \sigma_{x,f}$  depending on  $f$ . By Lemma 4.2,

$$\begin{aligned} & \sum_{f \in H_2(q)} \frac{|R(t, f)|^{2n}}{4\pi \langle f, f \rangle} \\ & \ll \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \left| \operatorname{Im} \sum_{p \leq x^3} \frac{C_f(p) (\Lambda_x(p) - \Lambda(p))}{p^{1/2+it} \log p} \right|^{2n} + \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \left| \operatorname{Im} \sum_{p \leq x^{3/2}} \frac{C_f(p^2) \Lambda_x(p^2)}{p^{1+2it} \log p} \right|^{2n} \\ & + \sum_{f \in H_2(q)} \frac{(\sigma_x - 1/2)^{2n} x^{2n(\sigma_x - 1/2)}}{4\pi \langle f, f \rangle} \left( \int_{1/2}^{\infty} x^{1/2-\sigma} \left| \sum_{p \leq x^3} \frac{C_f(p) \Lambda_x(p) \log(xp)}{p^{\sigma+it}} \right| d\sigma \right)^{2n} \\ & + \sum_{f \in H_2(q)} \frac{(\sigma_x - 1/2)^{2n}}{4\pi \langle f, f \rangle} (\log(|t| + q))^{2n}. \end{aligned} \tag{4.4}$$

Since

$$|\Lambda_x(p) - \Lambda(p)| = O\left(\frac{\log^3 p}{\log^2 x}\right) \quad \text{and} \quad C_f(p^2) = \lambda_f(p^2) - \varepsilon_q(p),$$

the first two terms is of  $O(1)$  by the argument as (4.1). The last term is of  $O(1)$  by Lemma 4.1. For the third term, it follows from Cauchy's inequality that

$$\begin{aligned} & \sum_{f \in H_2(q)} \frac{(\sigma_x - 1/2)^{2n} x^{2n(\sigma_x - 1/2)}}{4\pi \langle f, f \rangle} \left( \int_{1/2}^{\infty} x^{1/2 - \sigma} \left| \sum_{p \leq x^3} \frac{C_f(p) \Lambda_x(p) \log(xp)}{p^{\sigma + it}} \right| d\sigma \right)^{2n} \\ & \leq \left( \sum_{f \in H_2(q)} \frac{(\sigma_x - 1/2)^{4n} x^{4n(\sigma_x - 1/2)}}{4\pi \langle f, f \rangle} \right)^{1/2} \\ & \times \left( \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \left( \int_{1/2}^{\infty} x^{1/2 - \sigma} \left| \sum_{p \leq x^3} \frac{C_f(p) \Lambda_x(p) \log(xp)}{p^{\sigma + it}} \right| d\sigma \right)^{4n} \right)^{1/2}. \end{aligned}$$

By Hölder's inequality with the exponents  $4n/(4n - 1)$  and  $4n$ ,

$$\begin{aligned} & \left( \int_{1/2}^{\infty} x^{1/2 - \sigma} \left| \sum_{p \leq x^3} \frac{C_f(p) \Lambda_x(p) \log(xp)}{p^{\sigma + it}} \right| d\sigma \right)^{4n} \\ & \leq \left( \int_{1/2}^{\infty} x^{1/2 - \sigma} d\sigma \right)^{4n-1} \left( \int_{1/2}^{\infty} x^{1/2 - \sigma} \left| \sum_{p \leq x^3} \frac{C_f(p) \Lambda_x(p) \log(xp)}{p^{\sigma + it}} \right|^{4n} d\sigma \right) \\ & = \frac{1}{(\log x)^{4n-1}} \int_{1/2}^{\infty} x^{1/2 - \sigma} \left| \sum_{p \leq x^3} \frac{C_f(p) \Lambda_x(p) \log(xp)}{p^{\sigma + it}} \right|^{4n} d\sigma. \end{aligned}$$

Moreover, using Lemma 2.3 and the bound (2.2), we have

$$\begin{aligned} & \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \left| \sum_{p \leq x^3} \frac{C_f(p) \Lambda_x(p) \log(xp)}{p^{\sigma + it}} \right|^{4n} \\ & \ll \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \left( \sum_{p \leq x^3} \frac{|\Lambda_x(p) \log(xp) p^{1/2 - \sigma}|^2}{p} \right)^{2n} \\ & \ll \left( \sum_{p \leq x^3} \frac{|\Lambda_x(p) \log(xp) p^{1/2 - \sigma}|^2}{p} \right)^{2n} \ll \left( \sum_{p \leq x^3} \frac{(\log p)^2 (\log x)^2}{p} \right)^{2n} \ll (\log x)^{8n}. \end{aligned}$$

Thus,

$$\begin{aligned} & \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \left( \int_{1/2}^{\infty} x^{1/2-\sigma} \left| \sum_{p \leq x^3} \frac{C_f(p) \Lambda_x(p) \log(xp)}{p^{\sigma+it}} \right| d\sigma \right)^{4n} \\ & \ll \frac{1}{(\log x)^{4n-1}} \int_{1/2}^{\infty} x^{1/2-\sigma} (\log x)^{8n} d\sigma \ll (\log x)^{4n}. \end{aligned}$$

Using this bound and Lemma 4.1, we have the third term in (4.4) is of  $O(1)$ . Thus we complete the proof of proposition.  $\square$

**Proof of Theorem 1.1.** By the binomial theorem, we have

$$S(t, f)^n = M(t, f)^n + O_n \left( \sum_{\ell=1}^{\infty} |M(t, f)|^{n-\ell} |R(t, f)|^{\ell} \right).$$

For  $1 \leq \ell \leq n$ , we apply the generalized Hölder's inequality and Proposition 4.3, and get

$$\begin{aligned} & \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} |M(t, f)|^{n-\ell} |R(t, f)|^{\ell} \\ & \ll \left( \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \right)^{\frac{1}{2}} \left( \sum_{f \in H_2(q)} \frac{|M(t, f)|^{2n}}{4\pi \langle f, f \rangle} \right)^{\frac{n-\ell}{2n}} \left( \sum_{f \in H_2(q)} \frac{|R(t, f)|^{2n}}{4\pi \langle f, f \rangle} \right)^{\frac{\ell}{2n}} \\ & \ll_{t,n} (\log \log q)^{(n-\ell)/2} \ll_{t,n} (\log \log q)^{(n-1)/2}. \end{aligned}$$

**Proof of Theorem 1.2.** The  $n$ -th moment of  $\mu_q$  is

$$\int_{\mathbb{R}} \xi^n d\mu_q(\xi) = \left( \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \left( \frac{S(t, f)}{\sqrt{\log \log q}} \right)^n \right) / \left( \sum_{f \in H_2(q)} \frac{1}{4\pi \langle f, f \rangle} \right).$$

Using Theorem 1.1 and Lemma 2.3, we obtain that for all  $n$ ,

$$\lim_{q \rightarrow \infty} \int_{\mathbb{R}} \xi^n d\mu_q(\xi) = C_n = \sqrt{\pi} \int_{\mathbb{R}} \xi^n \exp(-\pi^2 \xi^2) d\xi.$$

By the theory of moments in probability theory (see Lemma 2.5), we complete the proof of theorem.

## REFERENCES

1. P. Billingsley, *Probability and measure*. (English summary) Third edition. Wiley Series in Probability and Mathematical Statistics. A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1995. xiv+593 pp.
2. H. Davenport, *Multiplicative number theory*. Second edition. Revised by Hugh L. Montgomery. Graduate Texts in Mathematics, 74. Springer-Verlag, New York-Berlin, 1980. xiii+177 pp.
3. I. Gradshteyn and I. Ryzhik, *Table of integrals, series, and products*. Translated from the Russian. Translation edited and with a preface by Alan Jeffrey and Daniel Zwillinger. With one CD-ROM (Windows, Macintosh and UNIX). Seventh edition. Elsevier/Academic Press, Amsterdam, 2007. xlviii+1171 pp.
4. D. A. Hejhal and W. Luo, *On a spectral analog of Selberg's result on  $S(T)$* . Internat. Math. Res. Notices 1997, no. 3, 135–151.
5. B. Hough, *Zero-density estimate for modular form  $L$ -functions in weight aspect*. Acta Arith. **154** (2012), no. 2, 187–216.

6. H. Iwaniec and E. Kowalski, *Analytic number theory*. American Mathematical Society Colloquium Publications, 53. American Mathematical Society, Providence, RI, 2004. xii+615 pp.
7. H. Iwaniec, W. Luo and P. Sarnak, *Low lying zeros of families of L-functions*. Inst. Hautes Études Sci. Publ. Math. No. **91** (2000), 55–131.
8. E. Kowalski and P. Michel, *The analytic rank of  $J_0(q)$  and zeros of automorphic L-functions*. Duke Math. J. **100** (1999), no. 3, 503–542.
9. E. Kowalski and P. Michel, *Explicit upper bound for the (analytic) rank of  $J_0(q)$* . Israel J. Math. **120** (2000), part A, 179–204.
10. S. C. Liu and S. Liu, *A  $GL_3$  analog of Selberg’s result on  $S(t)$* . Ramanujan J. **56** (2021), no. 1, 163–181.
11. S. C. Liu and J. Shim, *Moments of  $S(t, f)$  associated with holomorphic Hecke cusp forms*. Taiwanese J. Math. **26** (2022), no. 3, 463–482.
12. A. Selberg, *On the remainder in the formula for  $N(T)$ , the number of zeros of  $\zeta(s)$  in the strip  $0 < t < T$* . Avh. Norske Vid.-Akad. Oslo I 1944 (1944), no. 1, 27 pp.
13. A. Selberg, *Contributions to the theory of Dirichlet’s L-functions*. Skr. Norske Vid.-Akad. Oslo I 1946 (1946), no. 3, 62 pp.
14. A. Selberg, *Contributions to the theory of the Riemann zeta-function*. Arch. Math. Naturvid. **48** (1946), no. 5, 89–155.
15. G. Shimura, *Introduction to the arithmetic theory of automorphic functions*. Kanô Memorial Lectures, No. 1. Publications of the Mathematical Society of Japan, No. 11. Iwanami Shoten Publishers, Tokyo; Princeton University Press, Princeton, NJ, 1971. xiv+267 pp.

SCHOOL OF MATHEMATICS AND STATISTICS, SHANDONG UNIVERSITY, WEIHAI, WEIHAI, SHANDONG 264209, CHINA

*Email address:* `qfsun@sdu.edu.cn`

SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN, SHANDONG 250100, CHINA

*Email address:* `wh0315@mail.sdu.edu.cn`