Provable Benefit of Annealed Langevin Monte Carlo for Non-log-concave Sampling

Wei Guo, Molei Tao, Yongxin Chen Georgia Institute of Technology {wei.guo,mtao,yongchen}@gatech.edu

Abstract

We address the outstanding problem of sampling from an unnormalized density that may be non-log-concave and multimodal. To enhance the performance of simple Markov chain Monte Carlo (MCMC) methods, techniques of annealing type have been widely used. However, quantitative theoretical guarantees of these techniques are under-explored. This study takes a first step toward providing a non-asymptotic analysis of annealed MCMC. Specifically, we establish, for the first time, an oracle complexity of $\widetilde{O}\left(\frac{d\beta^2 A^2}{\varepsilon^6}\right)$ for simple annealed Langevin Monte Carlo algorithm to achieve ε^2 accuracy in Kullback-Leibler divergence to the target distribution $\pi \propto e^{-V}$ on \mathbb{R}^d with β -smooth potential V. Here, \mathcal{A} represents the action of a curve of probability measures interpolating the target distribution π and a readily sampleable distribution.

1 Introduction

We study the task of efficient sampling from a probability distribution $\pi \propto e^{-V}$ on \mathbb{R}^d . This fundamental problem is pivotal across various fields including computational statistics [42, 50, 5, 48], Bayesian inference [30], statistical physics [46, 37], and finance [17], and has been extensively studied in the literature [12]. The most common approach to this problem is Markov Chain Monte Carlo (MCMC), among which Langevin Monte Carlo (LMC) [20, 61, 13, 44] is a particularly popular choice. LMC can be understood as a time-discretization of a diffusion process, known as Langevin diffusion (LD), whose stationary distribution is the target distribution π , and has been attractive partly due to its robust performance despite conceptual simplicity.

Although LMC and its variants converge rapidly when the target distribution π is strongly logconcave or satisfies isoperimetric inequalities such as the log-Sobolev inequality (LSI) [20, 61, 13], its effectiveness diminishes when dealing with target distributions that are not strongly log-concave or are multimodal, such as mixtures of Gaussians. In such scenarios, the sampler often becomes confined to a single mode, severely limiting its ability to explore the entire distribution effectively. This results in significant challenges in transitioning between modes, which can dramatically increase the mixing time, making it exponential in dimension d. Such limitations highlight the need for enhanced MCMC methodologies that can efficiently navigate the complex landscapes of multimodal distributions, thereby improving convergence rates and overall sampling efficiency.

To address the challenges posed by multimodality, techniques around the notion of annealing have been widely employed [29, 45]. The general philosophy involves constructing a sequence of intermediate distributions $\pi_0, \pi_1, ..., \pi_M$ that bridge the gap between an easily samplable distribution π_0 (e.g., Gaussian or Dirac-like), and the target distribution $\pi_M = \pi$. The process starts with sampling from π_0 and progressively samples from each subsequent distribution until π_M is reached. When π_i and π_{i+1} are close enough, approximate samples from π_i can serve as a warm start for sampling from π_{i+1} , thereby facilitating this transition. Employing LMC within this framework gives rise to what is known as the annealed LMC algorithm, which is the focus of our study. Despite its empirical success [54, 55, 69, 68], a thorough theoretical understanding of annealed LMC, particularly its non-asymptotic complexity bounds, remains elusive.

In this work, we take a first step toward providing a non-asymptotic analysis of annealed MCMC. By leveraging the Girsanov theorem to quantify the differences between the sampling dynamics and a reference process, we establish an upper bound of the error of annealed MCMC by the summation of two terms. One term is equal to the ratio of an action integral in the Wasserstein geometry induced by optimal transport and the duration of the process and diminishes when the annealing is sufficiently slow. Another term quantifies the discretization error in implementation. Our approach challenges the traditional point of view that annealed MCMC is a cascade of warm start. Analysis based on this viewpoint still requires log-concavity or isoperimetry. Instead, our theoretical analysis of the annealed Langevin-based MCMC advances the field by eliminating the reliance on assumptions related to log-concavity or isoperimetry. Our strategy represents a paradigm shift in analyzing annealed MCMC. Even though a fully polynomial complexity bound has not been achieved yet. We hope our strategy can inspire future work towards this ambitious goal.

Our key technical contributions are summarized as follows.

- 1. We discover a novel strategy to analyze the non-asymptotic complexity bound of annealed MCMC algorithms that can circumvent the need for log-concavity or isoperimetry.
- 2. In Section 4, we investigate the annealed LD, which involves running LD with a dynamically changing target distribution and derive a surprising bound on the time required to simulate the SDE for achieving ε^2 -accuracy in KL divergence.
- 3. Building on the insights from the analysis of the continuous dynamic and accounting for discretization errors, we establish a non-asymptotic oracle complexity bound for annealed LMC in Section 5, which is applicable to a broad range of annealing schemes.

The quantitative results are summarized and compared to other sampling algorithms in Table 1. Notably, our approach operates under the least stringent assumptions and exhibits the most favorable ε -dependence among all isoperimetry-free sampling methods.

Algorithm	Isoperimetric Assumptions	Other Assumptions	Criterion	Complexity
LMC [61]	C-LSI	Potential smooth	$\varepsilon^2, \operatorname{KL}\left(\cdot \ \pi\right)$	$\widetilde{O}\left(C^2d\varepsilon^{-2}\right)$
PS [22]	C-LSI	Potential smooth	ε, TV	$\widetilde{O}\left(Cd^{1/2}\log\varepsilon^{-1}\right)$
STLMC [28]	/	Translated mixture of a well-conditioned distribution	ε, TV	$O\left(\operatorname{poly}\left(d,\varepsilon^{-1}\right)\right)$
RDMC [35]	/	Potential smooth, nearly convex at ∞	ε, TV	$O\left(\operatorname{poly}\left(d\right) e^{\operatorname{poly}\left(\varepsilon^{-1}\right)}\right)$
ZOD-MC [32]	/	Potential growing at most quadratically	ε , TV+W2	$\mathrm{e}^{\widetilde{O}(d)O\left(\log\varepsilon^{-1} ight)}$
ALMC (ours)	/	Potential smooth	$\varepsilon^2, \operatorname{KL}(\pi \ \cdot)$	$\widetilde{O}\left(d\mathcal{A}(d)^2\varepsilon^{-6}\right)$

Table 1: Comparison of oracle complexities in terms of d, ε , and LSI constant for sampling from $\pi \propto e^{-V}$. The notation "poly(·)" denotes a polynomial function of the indicated parameters.

Related works and comparison. We provide a brief overview of the literature, mainly focusing on the algorithms for non-log-concave sampling and their theoretical analysis.

• <u>Samplers based on tempering</u>. The fundamental concept of tempering involves sampling the system at various temperatures simultaneously: at higher temparatures, the distribution flattens, allowing particles to easily transition between modes, while at lower temperatures, particles can more effectively explore local structures. In simulated tempering [43, 64], the system's temperature is randomly switched, while in parallel tempering (also known as replica exchange) [57, 38], the temperatures of two particles are swapped according to a

specific rule. However, quantitative theoretical results for tempering are limited, and the existing results (e.g., [27, 28, 19]) apply primarily to certain special classes of non-log-concave distributions.

- Samplers based on general diffusions. Inspired by score-based diffusion models [33, 53, 56], recent advances have introduced sampling methods that reverse the Ornstein-Uhlenbeck process, as detailed in [35, 34, 32]. These samplers exhibit reduced sensitivity to isoperimetric conditions, but rely on estimating score functions (gradients of log-density) via importance sampling, which poses significant challenges in high-dimensional settings. Concurrently, studies such as [66, 59, 60, 49] have employed neural networks to approximate unknown drift terms, enabling an SDE to transport a readily sampleable distribution to the target distribution. This approach has shown excellent performance in handling complex distributions, albeit at the expense of significant computational resources required for neural network training. In contrast, annealed LMC runs on a known interpolation of probability distributions, thus simplifying sampling by obviating the need for intensive score estimation or neural network training.
- Non-asymptotic analysis for non-log-concave sampling. Drawing upon the stationarypoint analysis in non-convex optimization, the seminal work [4] characterizes the convergence of non-log-concave sampling via Fisher divergence. Subsequently, [11] applies this methodology to examine the local mixing of LMC. However, Fisher divergence is a relatively weak criterion compared to more commonly employed metrics such as totalvariational distance or Wasserstein distances. In contrast, our study provides a convergence guarantee in terms of KL divergence, which implies convergence in total-variation distance and offers a stronger result.

Notations and definitions. For $a, b \in \mathbb{R}$, we define $\llbracket a, b \rrbracket := [a, b] \cap \mathbb{Z}$, $a \wedge b := \min(a, b)$, and $a \vee b := \max(a, b)$. For a, b > 0, the notations $a \leq b$, a = O(b), and $b = \Omega(a)$ indicate that $a \leq Cb$ for some universal constant C > 0, and the notations $a \approx b$ and $a = \Theta(b)$ stand for both a = O(b) and b = O(a). $\widetilde{O}(\cdot)$ hides logarithmic dependence in $O(\cdot)$. A function $U \in C^2(\mathbb{R}^d)$ is $\alpha(> 0)$ -strongly-convex if $\nabla^2 U \succeq \alpha I$, and is $\beta(> 0)$ -smooth if $-\beta I \preceq \nabla^2 U \preceq \beta I$. The total-variation (TV) distance is defined as $\operatorname{TV}(\mu, \nu) = \sup_{A \subset \mathbb{R}^d} |\mu(A) - \nu(A)|$, and the Kullback-Leibler (KL) divergence is defined as $\operatorname{KL}(\mu \| \nu) = \mathbb{E}_{\mu} \left[\log \frac{d\mu}{d\nu} \right]$. $\|\cdot\|$ represents the ℓ^2 norm on \mathbb{R}^d . For $f : \mathbb{R}^d \to \mathbb{R}^{d'}$ and a probability measure μ on \mathbb{R}^d , $\|f\|_{L^2(\mu)} := \left(\int \|f\|^2 \,\mathrm{d}\mu\right)^{1/2}$, and the second-order moment of μ is defined as $\mathbb{E}_{\mu} \left[\|\cdot\|^2 \right]$.

2 Preliminaries

2.1 Stochastic Differential Equations and Girsanov Theorem

A stochastic differential equation (SDE) $X = (X_t)_{t \in [0,T]}$ is a stochastic process on $\Omega = C([0,T]; \mathbb{R}^d)$, the space of continuous functions from [0,T] to \mathbb{R}^d . The dynamics of X are typically represented by the equation $dX_t = b(X_t, t)dt + \sigma(X_t, t)dB_t, t \in [0,T]$, where $(B_t)_{t \in [0,T]}$ is a standard Brownian motion in \mathbb{R}^d . The **path measure** of X, denoted \mathbb{P}^X , characterizes the distribution of X over Ω and is defined by $\mathbb{P}^X(A) = \mathbb{P}(X \in A)$ for all measurable subset A of Ω . The following lemma, as a corollary of the Girsanov theorem [36, 58, 41], provides a methodology for computing the KL divergence between two path measures and serves as a crucial technical tool in our proof.

Lemma 1. Assume we have the following two SDEs on Ω :

$$\mathrm{d}X_t = a_t(X_t)\mathrm{d}t + \sqrt{2}\mathrm{d}B_t, \ X_0 \sim \mu; \qquad \mathrm{d}Y_t = b_t(Y_t)\mathrm{d}t + \sqrt{2}\mathrm{d}B_t, \ Y_0 \sim \nu.$$

Let \mathbb{P}^X and \mathbb{P}^Y denote the path measures of X and Y, respectively. Then

$$\mathrm{KL}\left(\mathbb{P}^{X} \left\|\mathbb{P}^{Y}\right) = \mathrm{KL}\left(\mu\|\nu\right) + \frac{1}{4} \mathbb{E}_{X \sim \mathbb{P}^{X}}\left[\int_{0}^{T} \left\|a_{t}(X_{t}) - b_{t}(X_{t})\right\|^{2} \mathrm{d}t\right].$$

2.2 Langevin Diffusion and Langevin Monte Carlo

The Langevin diffusion (LD) with target distribution $\pi \propto e^{-V}$ is the solution to the SDE

$$dX_t = -\nabla V(X_t)dt + \sqrt{2}dB_t, \ t \in [0,\infty); \ X_0 \sim \mu_0.$$
⁽¹⁾

It is well-known that under mild conditions, π is the unique stationary distribution this SDE, and when π has good regularity properties, the marginal distribution of X_t converges to π as $t \to +\infty$, so we can sample from π by simulating Equation (1) for a long time. However, in most of the cases, LD is intractable to simulate exactly, and the Euler-Maruyama discretization of Equation (1) leads to the Langevin Monte Carlo (LMC) algorithm. LMC with step size h > 0 and target distribution $\pi \propto e^{-V}$ is a Markov chain $\{X_{kh}\}_{k=0,1,...}$ constructed by iterating the following update rule:

$$X_{(k+1)h} = X_{kh} - h\nabla V(X_{kh}) + \sqrt{2}(B_{(k+1)h} - B_{kh}), \ k = 0, 1, ...; \ X_0 \sim \mu_0,$$
(2)

where $\{B_{(k+1)h} - B_{kh}\}_{k=0,1,\ldots} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0,hI).$

2.3 Wasserstein Distance and Curves of Probability Measures

We briefly introduce several fundamental concepts in optimal transport, and direct readers to authoritative textbooks [62, 63, 2, 1] for an in-depth exploration.

For two probability measures μ, ν on \mathbb{R}^d with finite second-order moments, the **Wasserstein-2** (W2) **distance** between μ and ν is defined as

$$W_2(\mu,\nu) = \inf_{\gamma \in \Pi(\mu,\nu)} \left(\int \|x-y\|^2 \gamma(\mathrm{d} x,\mathrm{d} y) \right)^{\frac{1}{2}},$$

where $\Pi(\mu, \nu)$ is the set of all couplings of (μ, ν) , i.e., probability measure γ on $\mathbb{R}^d \times \mathbb{R}^d$ with $\gamma(A \times \mathbb{R}^d) = \mu(A)$ and $\gamma(\mathbb{R}^d \times A) = \nu(A)$, for all measurable set $A \subset \mathbb{R}^d$.

Given a vector field $v = (v_t : \mathbb{R}^d \to \mathbb{R}^d)_{t \in [a,b]}$ and a curve of probability measures $\rho = (\rho_t)_{t \in [a,b]}$ on \mathbb{R}^d with finite second-order moments, we say that v generates ρ if the continuity equation $\partial_t \rho_t + \nabla \cdot (\rho_t v_t) = 0, t \in [a, b]$ holds. The **metric derivative** of ρ at $t \in [a, b]$ is defined as

$$\left|\dot{\rho}\right|_{t} := \lim_{\delta \to 0} \frac{W_{2}(\rho_{t+\delta}, \rho_{t})}{\left|\delta\right|}$$

which can be interpreted as the "speed" of this curve. If $|\dot{\rho}|_t$ exists and is finite for all $t \in [a, b]$, we say that ρ is **absolutely continuous (AC)**. The metric derivative and the continuity equation are closely related by the following important fact (see [2, Theorem 8.3.1]):

Lemma 2. For an AC curve of probability measures $(\rho_t)_{t \in [a,b]}$, any vector field $(v_t)_{t \in [a,b]}$ that generates $(\rho_t)_{t \in [a,b]}$ satisfies $|\dot{\rho}|_t \leq ||v_t||_{L^2(\rho_t)}$ for all $t \in [a,b]$. Moreover, the equality is reachable.

Finally, define the **action** of an AC curve of probability measures $(\rho_t)_{t \in [a,b]}$ as $\mathcal{A}(\rho) := \int_a^b |\dot{\rho}|_t^2 dt$.

2.4 Isoperimetric Inequalities

A probability measure π on \mathbb{R}^d satisfies a **log-Sobolev inequality (LSI)** with constant C, or C-LSI, if for all $f \in C^1(\mathbb{R}^d)$ with $\mathbb{E}_{\pi}[f^2] > 0$,

$$\mathbb{E}_{\pi}\left[f^{2}\log\frac{f^{2}}{\mathbb{E}_{\pi}\left[f^{2}\right]}\right] \leq 2C \mathbb{E}_{\pi}\left[\left\|\nabla f\right\|^{2}\right].$$

It is worth noting that α -strongly-log-concave distributions satisfy $\frac{1}{\alpha}$ -LSI (Bakry-Émery theorem, [3]). When $\pi \propto e^{-V}$ satisfies *C*-LSI and the potential *V* is β -smooth, it is established in [61] that the LD converges exponentially fast in KL divergence, while the LMC also converges exponentially with a bias that vanishes when the step size approaches 0.

3 Problem Setup

In this paper, we consider the following mild assumption on the target distribution $\pi \propto e^{-V}$ on \mathbb{R}^d , which is widely used in the field of sampling (see, e.g., [20, 61, 4, 13]):

Assumption 1. The potential V is β -smooth, and there exists a global minimizer x_* of V such that $||x_*|| \leq R$. Moreover, π has finite second-order moment.

Recall that the rationale behind annealing involves a gradual transition from a distribution that is easy to sample from to the more complex target distribution, which is crucial to our algorithm. Throughout this paper, we define the following curve of probability measures on \mathbb{R}^d :

$$\pi_{\theta} \propto \exp\left(-\eta(\theta)V - \frac{\lambda(\theta)}{2} \left\|\cdot\right\|^{2}\right), \ \theta \in [0, 1].$$
(3)

We use the term **annealing schedule** to refer to the functions $\eta(\cdot)$ and $\lambda(\cdot)$. They need to be differentiable and monotone, satisfying the boundary conditions $\eta_0 = \eta(0) \nearrow \eta(1) = 1$ and $\lambda_0 = \lambda(0) \searrow \lambda(1) = 0$. The values of $\eta_0 \in [0, 1]$ and $\lambda_0 \in (1, +\infty)$ will be determined later. This flexible interpolation scheme includes many general cases. For example, [6] and [26] used the schedule $\eta(\cdot) \equiv 1$, while [45] used the schedule $\lambda(\theta) = c(1 - \eta(\theta))$. The key motivation for this interpolation is that when $\theta = 0$, the quadratic term predominates, making the potential of π_0 strongly-convex with a moderate condition number, thus π_0 is easy to sample from; on the other hand, when $\theta = 1$, π_1 is just the target distribution π . As θ gradually increases from 0 to 1, the readily sampleable distribution π_0 slowly transform into the target distribution π_1 .

In Lemma 5, we prove that π_{θ} also has finite second-order moment, so the W2 distance between π_{θ} 's makes sense. We further make a mild assumption on the absolute continuity of the curve:

Assumption 2. The curve $(\pi_{\theta})_{\theta \in [0,1]}$ is AC with finite action $\mathcal{A}_{\eta,\lambda}(\pi) := \int_0^1 |\dot{\pi}|_{\theta}^2 d\theta$.

To facilitate easy sampling from π_0 , we consider two choices of the parameters η_0 and λ_0 , which have the following complexity guarantees.

Lemma 3. When $\eta_0 = 0$, $\pi_0 = \mathcal{N}\left(0, \frac{1}{\lambda_0}I\right)$ can be sampled directly; when $\eta_0 \in (0, 1]$, we choose $\lambda_0 = \eta_0 d\beta$, so that under Assumption 1, it takes $O\left(1 \lor \log \frac{\eta_0 \beta R^2}{d^2}\right) = \widetilde{O}(1)$ queries to the oracle of V and ∇V in expectation to precisely sample from π_0 via rejection sampling.

See Appendix A for the rejection sampling algorithm as well as the full proof of this lemma. The parameter R reflects prior knowledge about global minimizer(s) of the potential function V. Unless it is exceptionally large, indicating scarce prior information about the global minimizer(s) of V, this $\tilde{O}(1)$ complexity is negligible compared to the overall complexity of sampling. In particular, when the exact location of a global minimizer of V is known, we can adjust the potential V so that 0 becomes a global minimizer, thereby reducing the complexity to O(1).

Equipped with this foundational setup, we now proceed to introduce the annealed LD and annealed LMC algorithms.

4 Analysis of Annealed Langevin Diffusion

To elucidate the concept of annealing more clearly, we first consider the **annealed Langevin dif**fusion (ALD) algorithm, which samples from the $\pi \propto e^{-V}$ by running LD with a dynamically changing target distribution. For the purpose of this discussion, let us assume that we can exactly simulate *any* SDE with known drift and diffusion terms.

Fix an annealing schedule $\eta(\cdot), \lambda(\cdot)$ and choose a sufficiently long time T. We define a reparametrized curve of probability measures $(\tilde{\pi}_t := \pi_{t/T})_{t \in [0,T]}$. Starting with an initial sample $X_0 \sim \pi_0 = \tilde{\pi}_0$, we run the SDE

$$dX_t = \nabla \log \tilde{\pi}_t(X_t) dt + \sqrt{2} dB_t, \ t \in [0, T],$$
(4)

and ultimately output $X_T \sim \nu^{ALD}$ as an approximate sample from the target distribution π . Intuitively, when $\tilde{\pi}_t$ is changing slowly, the distribution of X_t should closely resemble $\tilde{\pi}_t$, leading to an output distribution ν^{ALD} that approximates the target distribution. This turns out to be true, as is confirmed by the following theorem, which provides a convergence guarantee for the ALD process.

Theorem 1. When choosing
$$T = \frac{A_{\eta,\lambda}(\pi)}{4\varepsilon^2}$$
, it follows that $\operatorname{KL}(\pi \| \nu^{\operatorname{ALD}}) \leq \varepsilon^2$.

Proof. Let \mathbb{Q} be the path measure of ALD (Equation (4)) initialized at $X_0 \sim \tilde{\pi}_0$, and define \mathbb{P} as the path measure corresponding to the following reference SDE:

$$dX_t = (\nabla \log \widetilde{\pi}_t + v_t)(X_t)dt + \sqrt{2}dB_t, \ X_0 \sim \widetilde{\pi}_0, \ t \in [0, T].$$
(5)

The vector field $v = (v_t)_{t \in [0,T]}$ is designed such that $X_t \sim \tilde{\pi}_t$ for all $t \in [0,T]$. According to the Fokker-Planck equation, v must satisfy the PDE

$$\partial_t \widetilde{\pi}_t = -\nabla \cdot \left(\widetilde{\pi}_t \left(\nabla \log \widetilde{\pi}_t + v_t \right) \right) + \Delta \widetilde{\pi}_t = -\nabla \cdot \left(\widetilde{\pi}_t v_t \right), \ t \in [0, T],$$

which means that v generates $(\tilde{\pi}_t := \pi_{t/T})_{t \in [0,T]}$. We can compute $\mathrm{KL}(\mathbb{P}||\mathbb{Q})$ using Lemma 1:

$$\operatorname{KL}\left(\mathbb{P}\|\mathbb{Q}\right) = \frac{1}{4} \operatorname{\mathbb{E}}_{\mathbb{P}}\left[\int_{0}^{T} \left\|v_{t}(X_{t})\right\|^{2} \mathrm{d}t\right] = \frac{1}{4} \int_{0}^{T} \left\|v_{t}\right\|_{L^{2}(\widetilde{\pi}_{t})}^{2} \mathrm{d}t$$

Leveraging Lemma 2, among all vector fields v that generate $(\tilde{\pi}_t := \pi_{t/T})_{t \in [0,T]}$, we can choose the one that minimizes $||v_t||_{L^2(\tilde{\pi}_t)}$, thereby making $||v_t||_{L^2(\tilde{\pi}_t)} = |\dot{\tilde{\pi}}|_t$, the metric derivative. With the reparameterization $\tilde{\pi}_t = \pi_{t/T}$, we have the following relation by chain rule:

$$\left|\dot{\pi}\right|_{t} = \lim_{\delta \to 0} \frac{W_2(\widetilde{\pi}_{t+\delta}, \widetilde{\pi}_t)}{|\delta|} = \lim_{\delta \to 0} \frac{W_2(\pi_{(t+\delta)/T}, \pi_{t/T})}{T |\delta/T|} = \frac{1}{T} |\dot{\pi}|_{t/T}$$

Employing the change-of-variable formula leads to

$$\mathrm{KL}\left(\mathbb{P}\|\mathbb{Q}\right) = \frac{1}{4} \int_0^T \left|\dot{\pi}\right|_t^2 \mathrm{d}t = \frac{1}{4T} \int_0^1 |\dot{\pi}|_\theta^2 \mathrm{d}\theta = \frac{\mathcal{A}_{\eta,\lambda}(\pi)}{4T}.$$

Finally, using the data-processing inequality ([12, Theorem 1.5.3]), with $T = \frac{A_{\eta,\lambda}(\pi)}{4\epsilon^2}$,

$$\mathrm{KL}\left(\pi \left\|\nu^{\mathrm{ALD}}\right) = \mathrm{KL}\left(\mathbb{P}_{T} \|\mathbb{Q}_{T}\right) \leq \mathrm{KL}\left(\mathbb{P} \|\mathbb{Q}\right) = \varepsilon^{2},$$

where \mathbb{P}_T and \mathbb{Q}_T stand for the marginal distributions of \mathbb{P} and \mathbb{Q} at time T, respectively.

Although we adopt a specific parametrization of the interpolating distribution (Equation (3)), which is widely used in practice, the results in Theorem 1 are applicable to any curve of probability measures $(\pi_{\theta})_{\theta \in [0,1]}$ that bridge π_0 and $\pi_1 = \pi$. This applicability extends as long as the closed forms of the drift and diffusion terms in the SDE, which precisely follow the path $(\pi_{\theta})_{\theta \in [0,1]}$, are known.

Let us delve deeper into the mechanics of the ALD. Although at time t the SDE (Equation (4)) targets the distribution $\tilde{\pi}_t$, the distribution of X_t does not precisely match $\tilde{\pi}_t$. Nevertheless, by choosing a sufficiently long time T, we actually move on the curve $(\pi_{\theta})_{\theta \in [0,1]}$ sufficiently slowly, thus minimizing the discrepancy between the path measure of $(X_t)_{t \in [0,T]}$ and the reference curve $(\tilde{\pi}_t)_{t \in [0,T]}$. By applying data-processing inequality, we can upper bound the error between the marginal distributions at time T by the joint distributions of the two paths. In essence, moving more slowly, sampling more precisely.

Our analysis predominantly addresses the *global* error across the entire curve of probability measures, rather than focusing solely on the *local* error at time T. This approach is inspired by [18] (see also [12, Section 4.4]) and [9], and stands in contrast to the traditional isoperimetry-based analyses of LD (e.g., [61, 13, 4]), which emphasize the decay of the KL divergence from the distribution of X_t to the target distribution and require LSI to bound the time derivative of this quantity. Notably, the total time T needed to run the SDE depends solely on the action of the curve $(\pi_{\theta})_{\theta \in [0,1]}$, obviating the need for assumptions related to log-concavity or isoperimetry.

We also remark that the ALD plays a significant role in another important subject known as nonequilibrium stochastic thermodynamics [52]. Recently, a refinement of the fluctuation theorem was discovered in [10, 24], stating that the irreversible entropy production in a stochastic thermodynamic system is equal to the ratio of a similar action integral as in Theorem 1 and the duration of the process T, resembling Theorem 1.

5 Analysis of Annealed Langevin Monte Carlo

It is crucial to recognize that the ALD (Equation (4)) cannot be precisely simulated in practice. Transitioning from LD to LMC introduces additional errors due to discretization. This section will present a detailed convergence analysis for the **annealed Langevin Monte Carlo (ALMC)** algorithm, which is implementable in practical scenarios.

A straightforward yet non-optimal method to discretize Equation (4) involves employing the Euler-Maruyama scheme, i.e.,

$$X_{t+\Delta t} \approx X_t + \nabla \log \widetilde{\pi}_t(X_t) \Delta t + \sqrt{2} (B_{t+\Delta t} - B_t), \ 0 \le t < t + \Delta t \le T.$$

However, considering that $\nabla \log \tilde{\pi}_t(x) = -\eta \left(\frac{t}{T}\right) \nabla V(x) - \lambda \left(\frac{t}{T}\right) x$, the integral of the linear term can be computed in closed form, so we can use the exponential-integrator scheme [65, 67] to further reduce the discretization error.

Given the total time T, we define a sequence of points $0 = \theta_0 < \theta_1 < ... < \theta_M = 1$, and set $T_{\ell} = T\theta_{\ell}, h_{\ell} = T(\theta_{\ell} - \theta_{\ell-1})$. The exponential-integrator scheme is then expressed as

$$dX_t = \left(-\eta\left(\frac{t}{T}\right)\nabla V(X_{t-}) - \lambda\left(\frac{t}{T}\right)X_t\right)dt + \sqrt{2}dB_t, \ t \in [0,T], \ X_0 \sim \pi_0,$$
(6)

where $t_{-} := T_{\ell-1}$ when $t \in [T_{\ell-1}, T_{\ell}), \ell \in [1, M]$. The explicit update rule is detailed in Algorithm 1, with x_{ℓ} denoting $X_{T_{\ell}}$, and the derivation of Equation (7) is presented in Appendix B.1. Notably, replacing $\nabla V(X_{t_{-}})$ with $\nabla V(X_{t})$ recovers the ALD (Equation (4)), and setting $\eta \equiv 1$ and $\lambda \equiv 0$ reduces to the classical LMC iterations.

Algorithm 1: Annealed LMC Algorithm

Input: Target distribution $\pi \propto e^{-V}$, total time *T*, annealing schedule $\eta(\cdot)$ and $\lambda(\cdot)$, discrete points $\theta_0, ..., \theta_M$. For $0 \le \theta \le \theta' \le 1$ define $\Lambda_{\alpha}(\theta', \theta) = \exp\left(-T \int_{-\infty}^{\theta'} \chi(x) dx\right)$

$$H(\theta',\theta) = T \int_{\theta}^{\theta'} \eta(u) \Lambda_0(u,\theta') du, \text{ and } \Lambda_1(\theta',\theta) = \sqrt{2T \int_{\theta}^{\theta'} \Lambda_0^2(u,\theta') du};$$

2 Obtain a sample $x_0 \sim \pi_0$ using rejection sampling (Algorithm 2);

- 3 for $\ell = 1, 2, ..., M$ do
- 4 Sample an independent Gaussian noise $\xi_{\ell} \sim \mathcal{N}(0, I)$;
- 5 Update

$$x_{\ell} = \Lambda_0(\theta_{\ell}, \theta_{\ell-1}) x_{\ell-1} - H(\theta_{\ell}, \theta_{\ell-1}) \nabla V(x_{\ell-1}) + \Lambda_1(\theta_{\ell}, \theta_{\ell-1}) \xi_{\ell}.$$
(7)

6 end

Output:
$$x_M \sim \nu^{\text{ALMC}}$$
, an approximate sample from π .

We illustrate the ALMC algorithm in Figure 1. The underlying intuition behind this algorithm is that by setting a sufficiently long total time T, the trajectory of the continuous dynamic (i.e., annealed LD) approaches the reference trajectory closely, as established in Theorem 1. Additionally, by adopting sufficiently small step sizes $h_1, ..., h_M$, the discretization error can be substantially reduced. Unlike traditional annealed LMC methods, which require multiple LMC update steps for each intermediate distribution $\pi_1, ..., \pi_M$, our approach views the annealed LMC as a discretization of a continuous-time process. Consequently, it is adequate to perform *a single step* of LMC towards each intermediate distribution $\tilde{\pi}_{T_1}, ..., \tilde{\pi}_{T_M}$, provided that T is sufficiently large and the step sizes are appropriately small.

The subsequent theorem provides a convergence guarantee for the annealed LMC algorithm, with the complete proof detailed in Appendix B.2.

Theorem 2. Under Assumptions 1 and 2, Algorithm 1 can generate a distribution ν^{ALMC} satisfying $\text{KL}(\pi \| \nu^{\text{ALMC}}) \leq \varepsilon^2$ within

$$\widetilde{O}\left(\frac{d\beta^2 \mathcal{A}_{\eta,\lambda}(\pi)^2}{\varepsilon^6}\right)$$

calls to the oracle of V and ∇V in expectation.




Figure 1: Illustration of the ALMC algorithm. The ℓ -th green arrow, proceeding from left to right, represents one step of LMC towards $\tilde{\pi}_{T_{\ell}}$ with step size h_{ℓ} , while each red arrow corresponds to the application of the same transition kernel, initialized at $\tilde{\pi}_{T_{\ell-1}}$ on the reference trajectory \mathbb{P} , which is depicted in purple. To evaluate KL ($\mathbb{P} || \mathbb{Q}$), the Girsanov theorem implies that we only need to bound the aggregate KL divergence across each small interval (i.e., the sum of the blue "distances").

Sketch of Proof. Let \mathbb{Q} be the path measure of ALMC (Equation (4)), whose marginal distribution at time T is the output distribution ν^{ALMC} . Again, let \mathbb{P} denote the reference path measure of Equation (5) used in the proof of Theorem 1, in which the same vector field $(v_t)_{t \in [0,T]}$ ensures that $X_t \sim \tilde{\pi}_t$ under \mathbb{P} for all $t \in [0,T]$.

Applying Girsanov theorem (Lemma 1) and carefully dealing with the discretization error, we can upper bound $\text{KL}(\mathbb{P}||\mathbb{Q})$ by

$$\operatorname{KL}\left(\mathbb{P}\|\mathbb{Q}\right) \lesssim \sum_{\ell=1}^{M} \left(\frac{1+\eta(\theta_{\ell})^{2}\beta^{2}h_{\ell}^{2}}{T} \int_{\theta_{\ell-1}}^{\theta_{\ell}} |\dot{\pi}|_{\theta}^{2} \mathrm{d}\theta + \eta(\theta_{\ell})^{2}\beta^{2} dh_{\ell}^{2} \left(1+h_{\ell}\left(\beta\eta(\theta_{\ell})+\lambda(\theta_{\ell-1})\right)\right)\right).$$

The first summation is governed by the total time T, which pertains to the convergence of the continuous dynamic (i.e., ALD). Setting $T \simeq \frac{A_{\eta,\lambda}(\pi)}{\varepsilon^2}$ ensures that the first summation remains $O(\varepsilon^2)$, provided that the step size h_ℓ is sufficiently small. The second summation addresses the discretization error, and it suffices to determine the appropriate value of θ_ℓ to minimize M, the total number of calls to the oracle of ∇V for discretizing the SDE. Combining M with the complexity of sampling from π_0 determines the overall complexity of the algorithm.

Once again, our analysis relies on bounding the global error between two path measures by Girsanov theorem. The annealing schedule $\eta(\cdot)$ and $\lambda(\cdot)$ influence the complexity exclusively through the action $\mathcal{A}_{\eta,\lambda}(\pi)$. Identifying the optimal annealing schedule to minimize this action remains a significant challenge, and we propose this as an area for future research.

We also note that our Assumption 1 encompasses strongly-log-concave target distributions. For sampling from these well-conditioned distributions via LMC, the complexity required to achieve ε -in TV distance scales as $\tilde{O}(\varepsilon^{-2})$ [12]. However, using Pinsker inequality $KL \ge 2TV^2$, our complexity to meet the same error criterion is $O(\varepsilon^{-6})$, indicating a significantly higher computational demand. Refining the parameter dependence in our algorithm to reduce this complexity remains a key area for future work.

Finally, we present a conjecture concerning the action $\mathcal{A}_{\eta,\lambda}(\pi)$, leaving its proof or disproof as an open question for future research. Following this conjecture, we demonstrate its applicability to a specific class of non-log-concave distributions in the subsequent example, with the detailed proof provided in Appendix C.

Conjecture 1. Under Assumption 1, the action $\mathcal{A}_{\eta,\lambda}(\pi)$ is bounded above by a polynomial function of the problem parameters, including d, β , R, $\mathbb{E}_{\pi} \left[\left\| \cdot \right\|^2 \right] \eta_0$, λ_0 , etc.

Example 1. Consider a mixture of Gaussian distributions in \mathbb{R}^d defined by $\pi = \sum_{i=1}^N p_i \mathcal{N}(y_i, \beta^{-1}I)$, where the weights $p_i > 0$, $\sum_{i=1}^N p_i = 1$, and $||y_i|| = r$ for all $i \in [\![1, N]\!]$. Consequently, the potential $V = -\log \pi$ is B-smooth, where $B = \beta(4r^2\beta + 1)$. With an annealing schedule defined by $\eta(\cdot) \equiv 1$ and $\lambda(\theta) = dB(1-\theta)^{\gamma}$ for some $1 \leq \gamma = O(1)$, it follows that $\mathcal{A}_{\eta,\lambda}(\pi) = O\left(d(r^2\beta + 1)\left(r^2 + \frac{d}{\beta}\right)\right)$.

To demonstrate the superiority of our theoretical results, let us consider a simplified scenario where N = 2, $y_1 = -y_2$, and $r^2 \gg \frac{1}{\beta}$. We derive from Example 1 and Pinsker inequality that the total complexity required to obtain a sample that is ε -accurate in TV is $\widetilde{O}\left(d^3\beta^2r^4(r^4\beta^2\vee d^2)\varepsilon^{-6}\right)$. In contrast, studies such as [51, 8, 19] indicate that the LSI constant of π is $\Omega\left(e^{\Theta(\beta r^2)}\right)$, so existing bounds in Table 1 suggest that LMC, to achieve the same accuracy, can only provide an exponential complexity guarantee of $\widetilde{O}\left(e^{\Theta(\beta r^2)}d\varepsilon^{-2}\right)$.

6 Conclusions and Future Work

In this paper, we have explored the complexity of annealed LMC for sampling from a non-logconcave probability measure without relying on log-concavity or isoperimetry. Central to our analysis are the Girsanov theorem and optimal transport techniques, thus circumventing the need for isoperimetric inequalities. While our proof primarily focuses on the annealing scheme as described in Equation (3), there is potential for adaptation to more general interpolations. Further exploration of these applications will be a focus of our future research. Technically, our proof methodology could be expanded to include a broader range of target distributions beyond those with smooth potentials, such as those with Hölder-continuous gradients [7, 21, 40, 47, 23, 22], or even heavy-tailed target distributions [44, 31]. Furthermore, eliminating the necessity of assuming global minimizers of the potential function would enhance the practical utility of our algorithm. Theoretically, proving **Conjecture 1** or obtaining a polynomial bound of $A_{\eta,\lambda}(\pi)$ under less restrictive conditions would further validate that our complexity bounds are genuinely polynomial. Finally, while this work emphasizes the upper bounds for non-log-concave sampling, an intriguing future avenue would be to verify the tightness of these bounds and explore potential lower bounds for this problem [16, 15, 14].

References

- Luigi Ambrosio, Elia Brué, and Daniele Semola. Lectures on optimal transport. Vol. 130. UNITEXT. Springer Cham, 2021. DOI: 10.1007/978-3-030-72162-6. URL: https://link.springer.com/book/10.1007/978-3-030-72162-6.
- [2] Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. Gradient Flows: In Metric Spaces and in the Space of Probability Measures. 2nd ed. Lectures in Mathematics. ETH Zürich. Birkhäuser Basel, 2008. DOI: 10.1007/978-3-7643-8722-8.
- [3] Dominique Bakry, Ivan Gentil, and Michel Ledoux. Analysis and geometry of Markov diffusion operators. 1st ed. Vol. 103. Grundlehren der mathematischen Wissenschaften. Springer Cham, 2014. DOI: 10.1007/978-3-319-00227-9.
- [4] Krishna Balasubramanian et al. "Towards a Theory of Non-Log-Concave Sampling: First-Order Stationarity Guarantees for Langevin Monte Carlo". In: *Proceedings of Thirty Fifth Conference on Learning Theory*. Ed. by Po-Ling Loh and Maxim Raginsky. Vol. 178. Proceedings of Machine Learning Research. PMLR, Feb. 2022, pp. 2896–2923. URL: https://proceedings.mlr.press/v178/balasubramanian22a.html.
- [5] Steve Brooks et al. Handbook of Markov chain Monte Carlo. CRC press, 2011.
- [6] Nicolas Brosse, Alain Durmus, and Éric Moulines. "Normalizing constants of log-concave densities". In: *Electronic Journal of Statistics* 12.1 (2018), pp. 851–889. DOI: 10.1214/18-EJS1411. URL: https://doi.org/10.1214/18-EJS1411.
- [7] Niladri Chatterji et al. "Langevin Monte Carlo without smoothness". In: Proceedings of the Twenty Third International Conference on Artificial Intelligence and Statistics. Ed. by Silvia Chiappa and Roberto Calandra. Vol. 108. Proceedings of Machine Learning Research. PMLR, 26–28 Aug 2020, pp. 1716– 1726. URL: https://proceedings.mlr.press/v108/chatterji20a.html.
- [8] Hong-Bin Chen, Sinho Chewi, and Jonathan Niles-Weed. "Dimension-free log-Sobolev inequalities for mixture distributions". In: *Journal of Functional Analysis* 281.11 (2021), p. 109236. ISSN: 0022-1236. DOI: https://doi.org/10.1016/j.jfa.2021.109236. URL: https://www.sciencedirect.com/science/article/pii/S0022123621003189.
- [9] Sitan Chen et al. "Sampling is as easy as learning the score: theory for diffusion models with minimal data assumptions". In: *The Eleventh International Conference on Learning Representations*. 2023. URL: https://openreview.net/forum?id=zyLVMgsZOU_.

- [10] Yongxin Chen, Tryphon T. Georgiou, and Allen Tannenbaum. "Stochastic Control and Nonequilibrium Thermodynamics: Fundamental Limits". In: *IEEE Transactions on Automatic Control* 65.7 (2020), pp. 2979–2991. DOI: 10.1109/TAC.2019.2939625.
- [11] Xiang Cheng et al. "Fast Conditional Mixing of MCMC Algorithms for Non-logconcave Distributions". In: Advances in Neural Information Processing Systems. Ed. by A. Oh et al. Vol. 36. Curran Associates, Inc., 2023, pp. 13374–13394. URL: https://proceedings.neurips.cc/paper_files/paper/2023/file/2b00b3331bd0f5fbfdd966ac06338f6d-Paper-Conf
- [12] Sinho Chewi. Log-Concave Sampling. Book draft, in preparation, 2024. URL: https://chewisinho.github.io.
- [13] Sinho Chewi et al. "Analysis of Langevin Monte Carlo from Poincaré to Log-Sobolev". In: Proceedings of Thirty Fifth Conference on Learning Theory. Ed. by Po-Ling Loh and Maxim Raginsky. Vol. 178. Proceedings of Machine Learning Research. PMLR, Feb. 2022, pp. 1–2. URL: https://proceedings.mlr.press/v178/chewi22a.html.
- [14] Sinho Chewi et al. "Fisher information lower bounds for sampling". In: Proceedings of The 34th International Conference on Algorithmic Learning Theory. Ed. by Shipra Agrawal and Francesco Orabona. Vol. 201. Proceedings of Machine Learning Research. PMLR, 20 Feb-23 Feb 2023, pp. 375-410. URL: https://proceedings.mlr.press/v201/chewi23b.html.
- [15] Sinho Chewi et al. "Query lower bounds for log-concave sampling". In: 2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS). 2023, pp. 2139–2148. DOI: 10.1109/F0CS57990.2023.00131.
- [16] Sinho Chewi et al. "The query complexity of sampling from strongly log-concave distributions in one dimension". In: *Proceedings of Thirty Fifth Conference on Learning Theory*. Ed. by Po-Ling Loh and Maxim Raginsky. Vol. 178. Proceedings of Machine Learning Research. PMLR, Feb. 2022, pp. 2041– 2059. URL: https://proceedings.mlr.press/v178/chewi22b.html.
- [17] John S. Dagpunar. Simulation and Monte Carlo: With Applications in Finance and MCMC. John Wiley & Sons, Ltd, 2007. DOI: 10.1002/9780470061336.
- [18] Arnak Dalalyan and Alexandre B. Tsybakov. "Sparse regression learning by aggregation and Langevin Monte-Carlo". In: Journal of Computer and System Sciences 78.5 (2012). JCSS Special Issue: Cloud Computing 2011, pp. 1423-1443. ISSN: 0022-0000. DOI: https://doi.org/10.1016/j.jcss.2011.12.023. URL: https://www.sciencedirect.com/science/article/pii/S0022000012000220.
- [19] Jing Dong and Xin T. Tong. "Spectral gap of replica exchange Langevin diffusion on mixture distributions". In: *Stochastic Processes and their Applications* 151 (2022), pp. 451– 489. ISSN: 0304-4149. DOI: https://doi.org/10.1016/j.spa.2022.06.006. URL: https://www.sciencedirect.com/science/article/pii/S0304414922001314.
- [20] Alain Durmus, Szymon Majewski, and Błażej Miasojedow. "Analysis of Langevin Monte Carlo via Convex Optimization". In: Journal of Machine Learning Research 20.73 (2019), pp. 1–46. URL: http://jmlr.org/papers/v20/18-173.html.
- [21] Murat A Erdogdu and Rasa Hosseinzadeh. "On the Convergence of Langevin Monte Carlo: The Interplay between Tail Growth and Smoothness". In: *Proceedings of Thirty Fourth Conference on Learning Theory*. Ed. by Mikhail Belkin and Samory Kpotufe. Vol. 134. Proceedings of Machine Learning Research. PMLR, 15–19 Aug 2021, pp. 1776–1822. URL: https://proceedings.mlr.press/v134/erdogdu21a.html.
- [22] Jiaojiao Fan, Bo Yuan, and Yongxin Chen. "Improved dimension dependence of a proximal algorithm for sampling". In: *Proceedings of Thirty Sixth Conference on Learning Theory*. Ed. by Gergely Neu and Lorenzo Rosasco. Vol. 195. Proceedings of Machine Learning Research. PMLR, Dec. 2023, pp. 1473– 1521. URL: https://proceedings.mlr.press/v195/fan23a.html.
- [23] Jiaojiao Fan et al. "Nesterov Smoothing for Sampling Without Smoothness". In: 2023 62nd IEEE Conference on Decision and Control (CDC). 2023, pp. 5313–5318. DOI: 10.1109/CDC49753.2023.10383529.
- [24] Rui Fu et al. "Maximal power output of a stochastic thermodynamic engine". In: *Automatica* 123 (2021), p. 109366.
- [25] Guillaume Garrigos and Robert M Gower. "Handbook of convergence theorems for (stochastic) gradient methods". In: *arXiv preprint arXiv:2301.11235* (2023).
- [26] Rong Ge, Holden Lee, and Jianfeng Lu. "Estimating Normalizing Constants for Log-Concave Distributions: Algorithms and Lower Bounds". In: *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing.* STOC 2020. Chicago, IL, USA: Association for Computing Machinery, 2020, pp. 579–586. ISBN: 9781450369794. DOI: 10.1145/3357713.3384289. URL: https://doi.org/10.1145/3357713.3384289.

- [27] Rong Ge, Holden Lee, and Andrej Risteski. "Beyond Log-concavity: Provable Guarantees for Sampling Multi-modal Distributions using Simulated Tempering Langevin Monte Carlo". In: Advances in Neural Information Processing Systems. Ed. by S. Bengio et al. Vol. 31. Curran Associates, Inc., 2018. URL: https://proceedings.neurips.cc/paper_files/paper/2018/file/c6ede20e6f597abf4b3f6bb30cee16c7-Paper.pdf.
- [28] Rong Ge, Holden Lee, and Andrej Risteski. "Simulated tempering Langevin Monte Carlo II: An improved proof using soft Markov chain decomposition". In: *arXiv preprint arXiv:1812.00793* (2018).
- [29] Saul Brian Gelfand, Sanjoy K Mitter, et al. "On sampling methods and annealing algorithms". In: technical report (1990).
- [30] Andrew Gelman et al. *Bayesian data analysis*. 3rd ed. Chapman and Hall/CRC, 2013.
- [31] Ye He, Krishnakumar Balasubramanian, and Murat A. Erdogdu. "An Analysis of Transformed Unadjusted Langevin Algorithm for Heavy-Tailed Sampling". In: *IEEE Transactions on Information Theory* 70.1 (2024), pp. 571–593. DOI: 10.1109/TIT.2023.3318152.
- [32] Ye He, Kevin Rojas, and Molei Tao. "Zeroth-Order Sampling Methods for Non-Log-Concave Distributions: Alleviating Metastability by Denoising Diffusion". In: *arXiv preprint arXiv:2402.17886* (2024).
- [33] Jonathan Ho, Ajay Jain, and Pieter Abbeel. "Denoising diffusion probabilistic models". In: Advances in Neural Information Processing Systems 33 (2020), pp. 6840–6851.
- [34] Xunpeng Huang et al. "Faster Sampling without Isoperimetry via Diffusion-based Monte Carlo". In: *arXiv preprint arXiv:2401.06325* (2024).
- [35] Xunpeng Huang et al. "Reverse Diffusion Monte Carlo". In: *The Twelfth International Conference on Learning Representations*. 2024. URL: https://openreview.net/forum?id=kIPEyMSdFV.
- [36] Ioannis Karatzas and Steven E. Shreve. *Brownian Motion and Stochastic Calculus*. 2nd ed. Graduate Texts in Mathematics. Springer New York, NY, 1991. DOI: 10.1007/978-1-4612-0949-2.
- [37] David Landau and Kurt Binder. A Guide to Monte Carlo Simulations in Statistical Physics. Cambridge University Press, 2021. DOI: 10.1017/9781108780346.
- [38] Holden Lee and Zeyu Shen. "Improved Bound for Mixing Time of Parallel Tempering". In: *arXiv* preprint arXiv:2304.01303 (2023).
- [39] Jiaming Liang and Yongxin Chen. "A Proximal Algorithm for Sampling". In: *Transactions on Machine Learning Research* (2023). ISSN: 2835-8856. URL: https://openreview.net/forum?id=CkXOwlhf27.
- [40] Jiaming Liang and Yongxin Chen. "A Proximal Algorithm for Sampling from Non-Smooth Potentials". In: 2022 Winter Simulation Conference (WSC). 2022, pp. 3229–3240. DOI: 10.1109/WSC57314.2022.10015293.
- [41] Robert S. Liptser and Albert N. Shiryaev. Statistics of Random Processes. I. General Theory. 2nd ed. Stochastic Modelling and Applied Probability. Springer Berlin, Heidelberg, 2013. DOI: 10.1007/978-3-662-13043-8.
- [42] Jun S. Liu. Monte Carlo strategies in scientific computing. Springer Series in Statistics. Springer New York, NY, 2004. DOI: 10.1007/978-0-387-76371-2.
- [43] Enzo Marinari and Giorgio Parisi. "Simulated Tempering: A New Monte Carlo Scheme". In: Europhysics Letters 19.6 (July 1992), p. 451. DOI: 10.1209/0295-5075/19/6/002. URL: https://dx.doi.org/10.1209/0295-5075/19/6/002.
- [44] Alireza Mousavi-Hosseini et al. "Towards a Complete Analysis of Langevin Monte Carlo: Beyond Poincaré Inequality". In: Proceedings of Thirty Sixth Conference on Learning Theory. Ed. by Gergely Neu and Lorenzo Rosasco. Vol. 195. Proceedings of Machine Learning Research. PMLR, Dec. 2023, pp. 1–35. URL: https://proceedings.mlr.press/v195/mousavi-hosseini23a.html.
- [45] Radford M. Neal. "Annealed importance sampling". In: Statistics and Computing 11.2 (Apr. 2001), pp. 125–139. ISSN: 1573-1375. DOI: 10.1023/A:1008923215028. URL: https://doi.org/10.1023/A:1008923215028.
- [46] Mark E. J. Newman and Gerard T. Barkema. Monte Carlo Methods in Statistical Physics. Oxford University Press, Feb. 1999. ISBN: 9780198517962. DOI: 10.1093/oso/9780198517962.001.0001. URL: https://doi.org/10.1093/oso/9780198517962.001.0001.
- [47] Dao Nguyen, Xin Dang, and Yixin Chen. "Unadjusted Langevin Algorithm Potentials". for Non-convex Weakly Smooth In: Communications in Math-10.1007/s40304-023-00350-w. ematics and **Statistics** (2023). DOI: URL: https://link.springer.com/article/10.1007/s40304-023-00350-w.
- [48] Art B. Owen. Monte Carlo theory, methods and examples. 2013. URL: https://artowen.su.domains/mc/.
- [49] Lorenz Richter and Julius Berner. "Improved sampling via learned diffusions". In: The Twelfth International Conference on Learning Representations. 2024. URL: https://openreview.net/forum?id=h4pNROsO06.
- [50] Christian P. Robert and George Casella. *Monte Carlo Statistical Methods*. 2nd ed. Springer Texts in Statistics. Springer New York, NY, 2004. DOI: 10.1007/978-1-4757-4145-2.

- [51] André Schlichting. "Poincaré and Log-Sobolev Inequalities for Mixtures". In: Entropy 21.1 (2019). ISSN: 1099-4300. DOI: 10.3390/e21010089. URL: https://www.mdpi.com/1099-4300/21/1/89.
- [52] Udo Seifert. "Stochastic thermodynamics, fluctuation theorems and molecular machines". In: *Reports on progress in physics* 75.12 (2012), p. 126001.
- [53] Jiaming Song, Chenlin Meng, and Stefano Ermon. "Denoising Diffusion Implicit Models". In: International Conference on Learning Representations. 2021. URL: https://openreview.net/forum?id=St1giarCHLP.
- [54] Yang Song and Stefano Ermon. "Generative Modeling by Estimating Gradients of the Data Distribution". In: Advances in Neural Information Processing Systems. Ed. by H. Wallach et al. Vol. 32. Curran Associates, Inc., 2019. URL: https://proceedings.neurips.cc/paper_files/paper/2019/file/3001ef257407d5a371a96dcd947c7d93-Paper.pdf.
- [55] Yang Song and Stefano Ermon. "Improved Techniques for Training Score-Based Generative Models". In: Advances in Neural Information Processing Systems. Ed. by H. Larochelle et al. Vol. 33. Curran Associates, Inc., 2020, pp. 12438-12448. URL: https://proceedings.neurips.cc/paper_files/paper/2020/file/92c3b916311a5517d9290576e3ea37ad-Paper.pdf.
- [56] Yang Song et al. "Score-Based Generative Modeling through Stochastic Differential Equations". In: International Conference on Learning Representations. 2021. URL: https://openreview.net/forum?id=PxTIG12RRHS.
- [57] Robert H. Swendsen and Jian-Sheng Wang. "Replica Monte Carlo Simulation of Spin-Glasses". In: *Phys. Rev. Lett.* 57 (21 Nov. 1986), pp. 2607–2609. DOI: 10.1103/PhysRevLett.57.2607. URL: https://link.aps.org/doi/10.1103/PhysRevLett.57.2607.
- [58] Ali Süleyman Üstünel and Moshe Zakai. *Transformation of Measure on Wiener Space*. 1st ed. Springer Monographs in Mathematics. Springer Berlin, Heidelberg, 2013. DOI: 10.1007/978-3-662-13225-8.
- [59] Francisco Vargas, Will Sussman Grathwohl, and Arnaud Doucet. "Denoising Diffusion Samplers". In: *The Eleventh International Conference on Learning Representations*. 2023. URL: https://openreview.net/forum?id=8pvnfTAbu1f.
- [60] Francisco Vargas et al. "Transport meets Variational Inference: Controlled Monte Carlo Diffusions". In: The Twelfth International Conference on Learning Representations. 2024. URL: https://openreview.net/forum?id=PP1rudnxiW.
- [61] Santosh S. Vempala and Andre Wibisono. "Rapid Convergence of the Unadjusted Langevin Algorithm: Isoperimetry Suffices". In: Advances in Neural Information Processing Systems. Ed. by H. Wallach et al. Vol. 32. Curran Associates, Inc., 2019. URL: https://proceedings.neurips.cc/paper_files/paper/2019/file/65a99bb7a3115fdede20da98b08a370f-Paper.pdf.
- [62] Cédric Villani. Optimal Transport: Old and New. 1st ed. Grundlehren der mathematischen Wissenschaften. Springer Berlin, Heidelberg, 2008. DOI: 10.1007/978-3-540-71050-9. URL: https://link.springer.com/book/10.1007/978-3-540-71050-9.
- [63] Cédric Villani. Topics in optimal transportation. Vol. 58. American Mathematical Society, 2021.
- [64] Dawn B. Woodard, Scott C. Schmidler, and Mark Huber. "Conditions for rapid mixing of parallel and simulated tempering on multimodal distributions". In: *The Annals of Applied Probability* 19.2 (2009), pp. 617–640. DOI: 10.1214/08-AAP555. URL: https://doi.org/10.1214/08-AAP555.
- [65] Qinsheng Zhang and Yongxin Chen. "Fast Sampling of Diffusion Models with Exponential Integrator". In: The Eleventh International Conference on Learning Representations. 2023. URL: https://openreview.net/forum?id=Loek7hfb46P.
- [66] Qinsheng Zhang and Yongxin Chen. "Path Integral Sampler: A Stochastic Control Approach For Sampling". In: International Conference on Learning Representations. 2022. URL: https://openreview.net/forum?id=_uCb2ynRu7Y.
- [67] Qinsheng Zhang, Molei Tao, and Yongxin Chen. "gDDIM: Generalized denoising diffusion implicit models". In: *The Eleventh International Conference on Learning Representations*. 2023. URL: https://openreview.net/forum?id=1hKE9qjvz-.
- [68] Nicolas Zilberstein, Ashutosh Sabharwal, and Santiago Segarra. "Solving Linear Inverse Problems Using Higher-Order Annealed Langevin Diffusion". In: *IEEE Transactions on Signal Processing* 72 (2024), pp. 492–505. DOI: 10.1109/TSP.2023.3348202.
- [69] Nicolas Zilberstein et al. "Annealed Langevin Dynamics for Massive MIMO Detection". In: IEEE Transactions on Wireless Communications 22.6 (2023), pp. 3762–3776. DOI: 10.1109/TWC.2022.3221057.

A Proof of Lemma 3

Our proof is inspired by [39].

We only consider the nontrivial case $\eta_0 \in (0, 1]$. The potential of π_0 is $V_0 := \eta_0 V + \frac{\lambda_0}{2} \|\cdot\|^2$, which is $(\lambda_0 - \eta_0\beta)$ -strongly-convex and $(\lambda_0 + \eta_0\beta)$ -smooth. Note that for any fixed point $x' \in \mathbb{R}^d$,

$$\pi_0(x) \propto \exp(-V_0(x)) \le \exp\left(-V_0(x') - \langle \nabla V_0(x'), x - x' \rangle - \frac{\lambda_0 - \eta_0 \beta}{2} \|x - x'\|^2\right).$$

The right-hand side is the unnormalized density of $\pi'_0 := \mathcal{N}\left(x' - \frac{\nabla V_0(x')}{\lambda_0 - \eta_0 \beta}, \frac{1}{\lambda_0 - \eta_0 \beta}I\right)$. The rejection sampling algorithm is as follows: sample $X \sim \pi'_0$, and accept X as a sample from π_0 with probability

$$\exp\left(-V_0(X) + V_0(x') + \langle \nabla V_0(x'), X - x' \rangle + \frac{\lambda_0 - \eta_0 \beta}{2} \|X - x'\|^2\right) \in (0, 1].$$

By [12, Theorem 7.1.1], the number of queries to the oracle of V until acceptance follows a geometric distribution with mean

$$\frac{\int \exp\left(-V_0(x') - \langle \nabla V_0(x'), x - x' \rangle - \frac{\lambda_0 - \eta_0 \beta}{2} \|x - x'\|^2\right) dx}{\int \exp\left(-V_0(x)\right) dx} \\
\leq \frac{\int \exp\left(-V_0(x') - \langle \nabla V_0(x'), x - x' \rangle - \frac{\lambda_0 - \eta_0 \beta}{2} \|x - x'\|^2\right) dx}{\int \exp\left(-V_0(x') - \langle \nabla V_0(x'), x - x' \rangle - \frac{\lambda_0 + \eta_0 \beta}{2} \|x - x'\|^2\right) dx} \\
= \frac{\int \exp\left(-\langle \nabla V_0(x'), x \rangle - \frac{\lambda_0 - \eta_0 \beta}{2} \|x\|^2\right) dx}{\int \exp\left(-\langle \nabla V_0(x'), x \rangle - \frac{\lambda_0 + \eta_0 \beta}{2} \|x\|^2\right) dx} \\
= \frac{\int \exp\left(-\frac{\lambda_0 - \eta_0 \beta}{2} \|x + \frac{\nabla V_0(x')}{\lambda_0 - \eta_0 \beta}\|^2 + \frac{\|\nabla V_0(x')\|^2}{2(\lambda_0 - \eta_0 \beta)}\right) dx} \\
= \left(\frac{\lambda_0 + \eta_0 \beta}{\lambda_0 - \eta_0 \beta}\right)^{\frac{d}{2}} \exp\left(\frac{\eta_0 \beta}{\lambda_0^2 - \eta_0^2 \beta^2} \|\nabla V_0(x')\|^2\right) \\
\leq \exp\left(\frac{\eta_0 \beta d}{\lambda_0 - \eta_0 \beta}\right) \exp\left(\frac{\eta_0 \beta}{\lambda_0^2 - \eta_0^2 \beta^2} \|\nabla V_0(x')\|^2\right)$$

We choose $\lambda_0 = \eta_0 \beta d$ such that $\exp\left(\frac{\eta_0 \beta d}{\lambda_0 - \eta_0 \beta}\right) \lesssim 1$. With this λ_0 , $\exp\left(\frac{\eta_0 \beta}{\lambda_0^2 - \eta_0^2 \beta^2} \|\nabla V_0(x')\|^2\right)$ is also O(1) as long as $\|\nabla V_0(x')\|^2 \lesssim \eta_0 \beta d^2$.

Let x'' be the global minimizer of the strongly convex potential function V_0 , which satisfies

$$0 = \nabla V_0(x'') = \eta_0 \nabla V(x'') + \lambda_0 x'' = \eta_0 \nabla V(x'') + \eta_0 \beta dx''.$$

Given the smoothness of V_0 ,

$$\|\nabla V_0(x')\|^2 = \|\nabla V_0(x') - \nabla V_0(x'')\|^2 \le (\eta_0 \beta d + \eta_0 \beta)^2 \|x' - x''\|^2.$$

Therefore, to guarantee $\|\nabla V_0(x')\|^2 \lesssim \eta_0 \beta d^2$, it suffices to find an x' such that $\|x' - x''\| \lesssim \frac{1}{\sqrt{\eta_0 \beta}}$. We first derive an upper bound of $\|x''\|$ under Assumption 1. Since x_* is a global minimizer of V,

$$\beta d \|x''\| = \|\nabla V(x'')\| = \|\nabla V(x'') - \nabla V(x_*)\|$$
$$\leq \beta \|x'' - x_*\| \leq \beta (\|x''\| + R),$$
$$\implies \|x''\| \leq \frac{R}{d-1} \lesssim \frac{R}{d}.$$

When R = 0, i.e., 0 is a known global minimizer of both V and V_0 , we can directly set x' = 0; otherwise, we need to run optimization algorithms to find such an x'. According to standard results in convex optimization (see, e.g., [25, Theorem 3.6]), running gradient descent on function V_0 with step size $\frac{1}{\lambda_0 + \eta_0 \beta}$ yields the following convergence rate: starting from $x_0 = 0$, the *t*-th iterate x_t satisfies

$$\|x_t - x''\|^2 \le \left(1 - \frac{\lambda_0 - \eta_0 \beta}{\lambda_0 + \eta_0 \beta}\right)^t \|0 - x''\|^2 \lesssim \left(\frac{2}{d}\right)^t \frac{R^2}{d^2} \le \frac{R^2}{e^t d^2},$$

where the last inequality holds when $d \ge 6$. Thus, $\log \frac{\eta_0 \beta R^2}{d^2} + O(1)$ iterations are sufficient to find a desired x'. We summarize the rejection sampling algorithm in Algorithm 2. In conclusion, precisely sampling from π_0 requires

$$O\left(1 \vee \log \frac{\eta_0 \beta R^2}{d^2}\right) = \widetilde{O}\left(1\right)$$

calls to the oracle of V and ∇V in expectation.

Algorithm 2: Rejection Sampling for π_0 . Input: $\pi \propto \exp(-V), \eta_0 \in (0, 1], \lambda_0 = \eta_0 \beta d$. 1 Let $V_0 := \eta_0 V + \frac{\lambda_0}{2} \|\cdot\|^2$; 2 Use gradient descent to find an x' that is $O\left(\frac{1}{\sqrt{\eta_0\beta}}\right)$ -close to the global minimizer of V_0 ; 3 Let $\pi'_0 := \mathcal{N}\left(x' - \frac{\nabla V_0(x')}{\lambda_0 - \eta_0\beta}, \frac{1}{\lambda_0 - \eta_0\beta}I\right)$; 4 while *True* do 5 Sample $X \sim \pi'_0$ and $U \in [0, 1]$ independently; 6 Accept X as a sample of π_0 when $U \leq \exp\left(-V_0(X) + V_0(x') + \langle \nabla V_0(x'), X - x' \rangle + \frac{\lambda_0 - \eta_0\beta}{2} \|X - x'\|^2\right)$. 7 end Output: $X \sim \pi_0$.

B Proofs for Annealed LMC

B.1 Proof of Equation (7)

Define $\Lambda(t) := \int_0^t \lambda\left(\frac{\tau}{T}\right) d\tau$, whose derivative is $\Lambda'(t) = \lambda\left(\frac{t}{T}\right)$. By Itô's formula, we have

$$d\left(e^{A(t)}X_{t}\right) = e^{A(t)}\left(\lambda\left(\frac{t}{T}\right)X_{t}dt + dX_{t}\right) = e^{A(t)}\left(-\eta\left(\frac{t}{T}\right)\nabla V(X_{t-})dt + \sqrt{2}dB_{t}\right)$$

Integrating over $t \in [T_{\ell-1}, T_{\ell})$ (note that in this case $t_{-} = T_{\ell-1}$), we obtain

$$e^{A(T_{\ell})}X_{T_{\ell}} - e^{A(T_{\ell-1})}X_{T_{\ell-1}} = -\left(\int_{T_{\ell-1}}^{T_{\ell}} \eta\left(\frac{t}{T}\right)e^{A(t)}dt\right)\nabla V(X_{T_{\ell-1}}) + \sqrt{2}\int_{T_{\ell-1}}^{T_{\ell}} e^{A(t)}dB_t,$$

i.e.,

$$X_{T_{\ell}} = e^{-(\Lambda(T_{\ell}) - \Lambda(T_{\ell-1}))} X_{T_{\ell-1}} - \left(\int_{T_{\ell-1}}^{T_{\ell}} \eta\left(\frac{t}{T}\right) e^{-(\Lambda(T_{\ell}) - \Lambda(t))} dt \right) \nabla V(X_{T_{\ell-1}}) + \sqrt{2} \int_{T_{\ell-1}}^{T_{\ell}} e^{-(\Lambda(T_{\ell}) - \Lambda(t))} dB_t.$$

Since

$$\Lambda(T_{\ell}) - \Lambda(t) = \int_{t}^{T_{\ell}} \lambda\left(\frac{\tau}{T}\right) \mathrm{d}\tau = T \int_{t/T}^{T_{\ell}/T} \lambda(u) \mathrm{d}u,$$

by defining $\Lambda_0(\theta',\theta) = \exp\left(-T\int_{\theta}^{\theta'}\lambda(u)du\right)$, we have $e^{-(\Lambda(T_\ell)-\Lambda(T_{\ell-1}))} = \Lambda_0(\theta_\ell,\theta_{\ell-1})$. Similarly, we can show that

$$\int_{T_{\ell-1}}^{T_{\ell}} \eta\left(\frac{t}{T}\right) e^{-(\Lambda(T_{\ell}) - \Lambda(t))} dt = \int_{T_{\ell-1}}^{T_{\ell}} \eta\left(\frac{t}{T}\right) \Lambda_0\left(\frac{T_{\ell}}{T}, \frac{t}{T}\right) dt = T \int_{\theta_{\ell-1}}^{\theta_{\ell}} \eta(u) \Lambda_0(\theta_{\ell}, u) du,$$

and $\sqrt{2} \int_{T_{\ell-1}}^{T_{\ell}} e^{-(\Lambda(T_{\ell}) - \Lambda(t))} dB_t$ is a zero-mean Gaussian random vector with covariance

$$2\int_{T_{\ell-1}}^{T_{\ell}} e^{-2(\Lambda(T_{\ell})-\Lambda(t))} dt \cdot I = 2\int_{T_{\ell-1}}^{T_{\ell}} \Lambda_0^2\left(\frac{T_{\ell}}{T}, \frac{t}{T}\right) dt \cdot I = 2T\int_{\theta_{\ell-1}}^{\theta_{\ell}} \Lambda_0^2(\theta_{\ell}, u) du \cdot I$$

B.2 Proof of Theorem 2

We denote the path measure of ALMC (Equation (6)) by \mathbb{Q} . Then, \mathbb{Q}_T , the marginal distribution of X_T , serves as the output distribution ν^{ALMC} . Similar to the methodology in the proof of Theorem 1, we use \mathbb{P} to denote the reference path measure of Equation (5), in which the vector field $(v_t)_{t \in [0,T]}$ generates the cure of probability distributions $(\tilde{\pi}_t)_{t \in [0,T]}$.

Using the data-processing inequality, it suffices to demonstrate that $\text{KL}(\mathbb{P}||\mathbb{Q}) \leq \varepsilon^2$. By Girsanov theorem (Lemma 1) and triangle inequality, we have

$$\operatorname{KL}\left(\mathbb{P}\|\mathbb{Q}\right) = \frac{1}{4} \int_{0}^{T} \mathbb{E}_{\mathbb{P}}\left[\left\|\eta\left(\frac{t}{T}\right)\left(\nabla V(X_{t}) - \nabla V(X_{t_{-}})\right) - v_{t}(X_{t})\right\|^{2}\right] \mathrm{d}t$$
$$\lesssim \int_{0}^{T} \mathbb{E}_{\mathbb{P}}\left[\eta\left(\frac{t}{T}\right)^{2}\left\|\nabla V(X_{t}) - \nabla V(X_{t_{-}})\right\|^{2} + \left\|v_{t}(X_{t})\right\|^{2}\right] \mathrm{d}t$$
$$\leq \sum_{\ell=1}^{M} \eta\left(\frac{T_{\ell}}{T}\right)^{2} \beta^{2} \int_{T_{\ell-1}}^{T_{\ell}} \mathbb{E}_{\mathbb{P}}\left[\left\|X_{t} - X_{t_{-}}\right\|^{2}\right] \mathrm{d}t + \int_{0}^{T} \left\|v_{t}\right\|_{L^{2}(\tilde{\pi}_{t})}^{2} \mathrm{d}t$$

The last inequality arises from the smoothness of V and the increasing property of $\eta(\cdot)$. To bound $\mathbb{E}_{\mathbb{P}}\left[\left\|X_t - X_{t_-}\right\|^2\right]$, note that under \mathbb{P} , for $t \in [T_{\ell-1}, T_{\ell})$, we have

$$X_{t} - X_{t_{-}} = \int_{T_{\ell-1}}^{t} \left(\nabla \log \tilde{\pi}_{\tau} + v_{\tau} \right) (X_{\tau}) \mathrm{d}\tau + \sqrt{2} (B_{t} - B_{T_{\ell-1}})$$

Thanks to the fact that $X_t \sim \tilde{\pi}_t$ under \mathbb{P} ,

$$\begin{split} \mathbb{E}_{\mathbb{P}}\left[\left\|X_{t}-X_{t-}\right\|^{2}\right] &\lesssim \mathbb{E}_{\mathbb{P}}\left[\left\|\int_{T_{\ell-1}}^{t}\left(\nabla\log\tilde{\pi}_{\tau}+v_{\tau}\right)\left(X_{\tau}\right)\mathrm{d}\tau\right\|^{2}\right] + \mathbb{E}\left[\left\|\sqrt{2}(B_{t}-B_{T_{\ell-1}})\right\|^{2}\right] \\ &\lesssim \left(t-T_{\ell-1}\right)\int_{T_{\ell-1}}^{t}\mathbb{E}_{\mathbb{P}}\left[\left\|\left(\nabla\log\tilde{\pi}_{\tau}+v_{\tau}\right)\left(X_{\tau}\right)\right\|^{2}\right]\mathrm{d}\tau + d(t-T_{\ell-1}) \\ &\lesssim \left(t-T_{\ell-1}\right)\int_{T_{\ell-1}}^{t}\left(\left\|\nabla\log\tilde{\pi}_{\tau}\right\|^{2}_{L^{2}(\tilde{\pi}_{\tau})} + \left\|v_{\tau}\right\|^{2}_{L^{2}(\tilde{\pi}_{\tau})}\right)\mathrm{d}\tau + d(t-T_{\ell-1}) \\ &\lesssim h_{\ell}\int_{T_{\ell-1}}^{T_{\ell}}\left(\left\|\nabla\log\tilde{\pi}_{\tau}\right\|^{2}_{L^{2}(\tilde{\pi}_{\tau})} + \left\|v_{\tau}\right\|^{2}_{L^{2}(\tilde{\pi}_{\tau})}\right)\mathrm{d}\tau + dh_{\ell}. \end{split}$$

The second inequality arises from the application of the Cauchy-Schwarz inequality, and the last inequality is due to the definition $h_{\ell} = T_{\ell} - T_{\ell-1}$. Taking integral over $t \in [T_{\ell-1}, T_{\ell}]$,

$$\int_{T_{\ell-1}}^{T_{\ell}} \mathbb{E}\left[\left\|X_t - X_{t_-}\right\|^2\right] \mathrm{d}t \lesssim h_{\ell}^2 \int_{T_{\ell-1}}^{T_{\ell}} \left(\left\|\nabla\log\widetilde{\pi}_t\right\|_{L^2(\widetilde{\pi}_t)}^2 + \left\|v_t\right\|_{L^2(\widetilde{\pi}_t)}^2\right) \mathrm{d}t + dh_{\ell}^2.$$

Recall that the potential of $\tilde{\pi}_t$ is $\left(\eta\left(\frac{t}{T}\right)\beta + \lambda\left(\frac{t}{T}\right)\right)$ -smooth. By Lemma 4 and the monotonicity of $\eta(\cdot)$ and $\lambda(\cdot)$, we have

$$\begin{split} \int_{T_{\ell-1}}^{T_{\ell}} \|\nabla \log \widetilde{\pi}_t\|_{L^2(\widetilde{\pi}_t)}^2 \, \mathrm{d}t &\leq \int_{T_{\ell-1}}^{T_{\ell}} d\left(\eta\left(\frac{t}{T}\right)\beta + \lambda\left(\frac{t}{T}\right)\right) \, \mathrm{d}t \\ &\leq dh_\ell \left(\beta\eta\left(\frac{T_\ell}{T}\right) + \lambda\left(\frac{T_{\ell-1}}{T}\right)\right) \\ &= dh_\ell \left(\beta\eta\left(\theta_\ell\right) + \lambda\left(\theta_{\ell-1}\right)\right). \end{split}$$

Therefore, $\mathrm{KL}\left(\mathbb{P}\|\mathbb{Q}\right)$ is upper bounded by

$$\lesssim \sum_{\ell=1}^{M} \eta \left(\theta_{\ell}\right)^{2} \beta^{2} \int_{T_{\ell-1}}^{T_{\ell}} \mathbb{E}_{\mathbb{P}} \left[\left\| X_{t} - X_{t-} \right\|^{2} \right] \mathrm{d}t + \int_{0}^{T} \left\| v_{t} \right\|_{L^{2}(\widetilde{\pi}_{t})}^{2} \mathrm{d}t \\ \lesssim \sum_{\ell=1}^{M} \eta \left(\theta_{\ell}\right)^{2} \beta^{2} \left(dh_{\ell}^{3} \left(\beta \eta \left(\theta_{\ell}\right) + \lambda \left(\theta_{\ell-1}\right)\right) + h_{\ell}^{2} \int_{T_{\ell-1}}^{T_{\ell}} \left\| v_{t} \right\|_{L^{2}(\widetilde{\pi}_{t})}^{2} \mathrm{d}t + dh_{\ell}^{2} \right) + \int_{0}^{T} \left\| v_{t} \right\|_{L^{2}(\widetilde{\pi}_{t})}^{2} \mathrm{d}t \\ = \sum_{\ell=1}^{M} \left(\left(1 + \eta (\theta_{\ell})^{2} \beta^{2} h_{\ell}^{2} \right) \int_{T_{\ell-1}}^{T_{\ell}} \left\| v_{t} \right\|_{L^{2}(\widetilde{\pi}_{t})}^{2} \mathrm{d}t + \eta (\theta_{\ell})^{2} \beta^{2} dh_{\ell}^{2} \left(1 + h_{\ell} \left(\beta \eta (\theta_{\ell}) + \lambda (\theta_{\ell-1})\right) \right) \right).$$

For the remaining integral, given that $(v_t)_{t \in [0,T]}$ generates $(\tilde{\pi}_t)_{t \in [0,T]}$, according to Lemma 2, we may choose v_t such that $||v_t||_{L^2(\tilde{\pi}_t)} = \left|\dot{\tilde{\pi}}\right|_t$. Thus,

$$\int_{T_{\ell-1}}^{T_{\ell}} \|v_t\|_{L^2(\tilde{\pi}_t)}^2 \,\mathrm{d}t = \int_{T_{\ell-1}}^{T_{\ell}} \left|\dot{\tilde{\pi}}\right|_t^2 \,\mathrm{d}t = \int_{T_{\ell-1}}^{T_{\ell}} \frac{1}{T^2} \,|\dot{\pi}|_{t/T}^2 \,\mathrm{d}t = \frac{1}{T} \int_{\theta_{\ell-1}}^{\theta_{\ell}} |\dot{\pi}|_{\theta}^2 \,\mathrm{d}\theta,$$

through a change-of-variable analogous to that used in the proof of Theorem 1. Therefore,

$$\operatorname{KL}\left(\mathbb{P}\|\mathbb{Q}\right) \lesssim \sum_{\ell=1}^{M} \left(\frac{1 + \eta(\theta_{\ell})^{2} \beta^{2} h_{\ell}^{2}}{T} \int_{\theta_{\ell-1}}^{\theta_{\ell}} |\dot{\pi}|_{\theta}^{2} \mathrm{d}\theta + \eta(\theta_{\ell})^{2} \beta^{2} dh_{\ell}^{2} \left(1 + h_{\ell} \left(\beta \eta(\theta_{\ell}) + \lambda(\theta_{\ell-1})\right)\right) \right).$$

Assume $h_{\ell} \lesssim \frac{1}{\beta d}$ (which will be verified later), so we can further simplify the above expression to

$$\operatorname{KL}\left(\mathbb{P}\|\mathbb{Q}\right) \lesssim \sum_{\ell=1}^{M} \left(\frac{1}{T} \int_{\theta_{\ell-1}}^{\theta_{\ell}} |\dot{\pi}|_{\theta}^{2} \,\mathrm{d}\theta + \eta(\theta_{\ell})^{2} \beta^{2} dh_{\ell}^{2}\right)$$
$$= \frac{\mathcal{A}_{\eta,\lambda}(\pi)}{T} + \beta^{2} d\sum_{\ell=1}^{M} \eta(\theta_{\ell})^{2} h_{\ell}^{2}$$
$$= \frac{\mathcal{A}_{\eta,\lambda}(\pi)}{T} + \beta^{2} d\sum_{\ell=1}^{M} \eta(\theta_{\ell})^{2} T^{2} (\theta_{\ell} - \theta_{\ell-1})^{2}$$

To bound the above expression by ε^2 , we first select $T \approx \frac{A_{\eta,\lambda}(\pi)}{\varepsilon^2}$, mirroring the total time T required for running annealed LD as specified in Theorem 1. This guarantees that the continuous dynamics closely approximate the reference path measure. Given that $\eta(\cdot) \leq 1$, it remains only to ensure

$$\beta^2 d \sum_{\ell=1}^M \eta(\theta_\ell)^2 T^2 (\theta_\ell - \theta_{\ell-1})^2 \le \beta^2 d \frac{\mathcal{A}_{\eta,\lambda}(\pi)^2}{\varepsilon^4} \sum_{\ell=1}^M (\theta_\ell - \theta_{\ell-1})^2 \lesssim \varepsilon^2,$$

which is equivalent to

$$\sum_{\ell=1}^{M} (\theta_{\ell} - \theta_{\ell-1})^2 \lesssim \frac{\varepsilon^6}{d\beta^2 \mathcal{A}_{\eta,\lambda}(\pi)^2}.$$
(8)

To minimize M, we apply Cauchy-Schwarz inequality:

$$\left(\sum_{\ell=1}^{M} 1\right) \left(\sum_{\ell=1}^{M} (\theta_{\ell} - \theta_{\ell-1})^2\right) \ge \left(\sum_{\ell=1}^{M} (\theta_{\ell} - \theta_{\ell-1})\right)^2 = 1,$$

which establishes a lower bound for M. The equality is achieved when $\theta_{\ell} - \theta_{\ell-1} = \frac{1}{M}$ for all $\ell \in [\![1, M]\!]$. Thus, selecting

$$M \asymp \frac{d\beta^2 \mathcal{A}_{\eta,\lambda}(\pi)^2}{\varepsilon^6}$$

satisfies the constraint given in Equation (8). In this case, the step size $h_{\ell} = \frac{T}{M} \approx \frac{\varepsilon^4}{d\beta^2} \lesssim \frac{1}{\beta d}$. Combining this with the $\widetilde{O}(1)$ complexity for sampling from π_0 , we have completed the proof.

C Proof of Example 1

The smoothness of V comes from [11, Lemma 4].

Note that $\pi(x) \propto \sum_{i=1}^{N} p_i \exp\left(-\frac{\beta}{2} \left\|x - y_i\right\|^2\right)$, and define

$$\begin{aligned} \widehat{\pi}_{\lambda}(x) &\propto \pi(x) \exp\left(-\frac{\lambda}{2} \|x\|^{2}\right) \\ &= \sum_{i=1}^{N} p_{i} \exp\left(-\frac{\lambda\beta}{2(\lambda+\beta)} \|y_{i}\|^{2} - \frac{\lambda+\beta}{2} \left\|x - \frac{\beta}{\lambda+\beta}y_{i}\right\|^{2}\right) \\ &\propto \sum_{i=1}^{N} p_{i} \exp\left(-\frac{\lambda+\beta}{2} \left\|x - \frac{\beta}{\lambda+\beta}y_{i}\right\|^{2}\right) \\ &= \sum_{i=1}^{N} p_{i} \mathcal{N}\left(\frac{\beta}{\lambda+\beta}y_{i}, \frac{1}{\lambda+\beta}I\right). \end{aligned}$$

We use coupling method to upper bound $W_2^2(\widehat{\pi}_{\lambda}, \widehat{\pi}_{\lambda+\delta})$. We first sample I with distribution $\mathbb{P}\left(I=i\right)=p_i, i\in \llbracket 1,N \rrbracket$, and then independently sample $\eta \sim \mathcal{N}\left(0,I\right)$. Then,

$$X := \frac{\beta}{\lambda + \beta} y_I + \frac{1}{\sqrt{\lambda + \beta}} \eta \sim \hat{\pi}_{\lambda},$$
$$Y := \frac{\beta}{\lambda + \delta + \beta} y_I + \frac{1}{\sqrt{\lambda + \delta + \beta}} \eta \sim \hat{\pi}_{\lambda + \delta}.$$

By definition of W2 distance, we have

$$\begin{split} W_2^2(\widehat{\pi}_{\lambda}, \widehat{\pi}_{\lambda+\delta}) &\leq \mathbb{E}\left[\|X - Y\|^2 \right] \\ &= \mathbb{E}_I \left[\mathbb{E}_{\eta} \left[\left\| \left(\frac{\beta}{\lambda+\beta} - \frac{\beta}{\lambda+\delta+\beta} \right) y_I + \left(\frac{1}{\sqrt{\lambda+\beta}} - \frac{1}{\sqrt{\lambda+\delta+\beta}} \right) \eta \right\|^2 \right] \right] \\ &= \mathbb{E}_I \left[\left(\frac{\beta}{\lambda+\beta} - \frac{\beta}{\lambda+\delta+\beta} \right)^2 \|y_I\|^2 + \left(\frac{1}{\sqrt{\lambda+\beta}} - \frac{1}{\sqrt{\lambda+\delta+\beta}} \right)^2 d \right] \\ &= \left(\frac{\beta}{\lambda+\beta} - \frac{\beta}{\lambda+\delta+\beta} \right)^2 r^2 + \left(\frac{1}{\sqrt{\lambda+\beta}} - \frac{1}{\sqrt{\lambda+\delta+\beta}} \right)^2 d. \end{split}$$

This implies

$$\left|\hat{\pi}\right|_{\lambda}^{2} = \lim_{\delta \to 0} \frac{W_{2}^{2}(\hat{\pi}_{\lambda}, \hat{\pi}_{\lambda+\delta})}{\delta^{2}} \leq \frac{\beta^{2}r^{2}}{(\lambda+\beta)^{4}} + \frac{d}{4(\lambda+\beta)^{3}}.$$

By time reparameterization $\pi_{\theta} = \hat{\pi}_{\lambda(\theta)}, |\dot{\pi}|_{\theta} = \left|\dot{\pi}\right|_{\lambda(\theta)} |\dot{\lambda}(\theta)|$. With $\lambda(\theta) = \lambda_0 (1-\theta)^{\gamma}, |\dot{\lambda}(\theta)| = |\gamma \lambda_0 (1-\theta)^{\gamma-1}| \le \gamma \lambda_0 \lesssim \lambda_0$. Therefore,

$$\begin{split} \mathcal{A}_{\eta,\lambda}(\pi) &= \int_0^1 |\dot{\pi}|_{\theta}^2 \,\mathrm{d}\theta = \int_0^1 \left|\dot{\pi}\right|_{\lambda(\theta)}^2 \left|\dot{\lambda}(\theta)\right|^2 \mathrm{d}\theta \\ &\lesssim \lambda_0 \int_0^1 \left|\dot{\pi}\right|_{\lambda(\theta)}^2 \left|\dot{\lambda}(\theta)\right| \mathrm{d}\theta = \lambda_0 \int_0^{\lambda_0} \left|\dot{\pi}\right|_{\lambda}^2 \mathrm{d}\lambda \\ &= \lambda_0 \int_0^{\lambda_0} \left(\frac{\beta^2 r^2}{(\lambda+\beta)^4} + \frac{d}{4(\lambda+\beta)^3}\right) \mathrm{d}\lambda \\ &\lesssim \lambda_0 \left(\beta^2 r^2 \frac{1}{\beta^3} + d\frac{1}{\beta^2}\right) \\ &\lesssim d\beta (r^2\beta + 1) \left(\frac{r^2}{\beta} + \frac{d}{\beta^2}\right) \\ &= d(r^2\beta + 1) \left(r^2 + \frac{d}{\beta}\right). \end{split}$$

It follows from the proof that as long as $\max_{\theta \in [0,1]} |\dot{\lambda}(\theta)|$ is bounded by a polynomial function of β , d and r, so is the action $\mathcal{A}_{\eta,\lambda}(\pi)$.

D Supplementary Lemmas

Lemma 4 ([12, Lemma 4.E.1]). Consider a probability measure $\mu \propto e^{-U}$ on \mathbb{R}^d . If $\nabla^2 U \preceq \beta I$ for some $\beta > 0$, then $\mathbb{E}_{\mu} \left[\|\nabla U\|^2 \right] \leq \beta d$.

Lemma 5. Under Assumption 1, π_{θ} defined in Equation (3) has finite second-order moment when $\eta(\theta) \in [0, 1]$.

Proof. When $\eta(\theta) = 1$, $\pi_{\theta} \propto \exp\left(-V - \frac{\lambda(\theta)}{2} \|\cdot\|^2\right) \leq \exp\left(-V\right)$, and the claim is straightforward. Otherwise, by convexity of $u \mapsto e^{-u}$, we have

$$\exp\left(-\eta(\theta)V - \frac{\lambda(\theta)}{2} \left\|\cdot\right\|^{2}\right) = \exp\left(-\eta(\theta)V - (1 - \eta(\theta))\frac{\lambda(\theta)}{2(1 - \eta(\theta))} \left\|\cdot\right\|^{2}\right)$$
$$\leq \eta(\theta)\exp\left(-V\right) + (1 - \eta(\theta))\exp\left(-\frac{\lambda(\theta)}{2(1 - \eta(\theta))} \left\|\cdot\right\|^{2}\right).$$

Multiplying both sides by $\|\cdot\|^2$ and taking integral over \mathbb{R}^d , we see that $\mathbb{E}_{\pi_\theta}\left[\|\cdot\|^2\right] < +\infty$.