

# Fermion-Vortex Interactions in Axion Electrodynamics

Saurav Kantha

*S. N. Bose National Centre for Basic Sciences,*

*JD Block, Sector 3, Salt Lake,*

*Kolkata 700106, WB, INDIA*

*and*

*Laboratoire de Physique Théorique,*

*Université Paul Sabatier, 31400 Toulouse, France\**

Amitabha Lahiri<sup>†</sup>

*S. N. Bose National Centre for Basic Sciences,*

*JD Block, Sector 3, Salt Lake, Kolkata 700106, WB, INDIA*

(Dated: July 26, 2024)

## Abstract

A relativistic action for scalar condensate-fermion mixture is considered where both the scalar boson and the fermion fields are coupled to a  $U(1)$  gauge field. The dynamics of the gauge field is governed by a linear combination of the Maxwell term, and the Lorentz invariant  $\mathbf{E} \cdot \mathbf{B}$  term with a constant coefficient  $\theta$ . We obtain an effective action describing an emergent fermion-fermion interaction and fermion-vortex tube interaction by using the particle-string duality, and find that the  $\theta$  term can significantly affect the interaction of fermions and vortices. We also perform a dimensional reduction to show a  $\theta$  dependent flux attachment to the itinerant fermions.

---

\* saurav.kantha@irsamc.ups-tlse.fr

† amitabha@bose.res.in

## I. INTRODUCTION

The interaction of charged currents with rank-1 gauge fields is a well understood concept in physics, both in high-energy and condensed matter sectors. In relativistic physics, the coupling goes by the name of minimal coupling, whereas in condensed matter physics, the gauge fields are introduced on the lattice via Peierls substitution [1]. In both cases, the gauge invariance is ensured by a local phase change of the states as the gauge field  $A$  goes to  $A + d\phi$  under the action of the underlying gauge group.

However, as we move one rank higher in the gauge field i.e. from a rank-1 gauge field  $A_\mu$  to a rank-2 gauge field  $B_{\mu\nu}$ , a gauge-invariant local coupling of fermions to the gauge field is no longer possible. A gauge-invariant, but nonlocal, interaction between fermions and an antisymmetric rank-2 tensor gauge field was recently found in the context of topological mass generation [2]. In this interaction, the gauge field  $B_{\mu\nu}$  couples to the fermion current  $j^\mu$  via the nonlocal term  $B_{\mu\nu}\tilde{J}^{\mu\nu}$ , where  $\tilde{J}^{\mu\nu} = \epsilon^{\mu\nu\rho\lambda}\frac{1}{\square}\partial_\rho j_\lambda$  is a nonlocal current because of the appearance of the inverse of the four-dimensional wave operator  $\square \equiv \partial_\mu\partial^\mu$ . For the purposes of calculations, we can think of  $\frac{1}{\square}$  as a Green function, or in momentum space as  $-\frac{1}{k^2}$ .

The consequences of such a coupling have not been explored very much. It has been known for a long time that the rank-2 antisymmetric gauge field  $B_{\mu\nu}$  mediates interactions between strings [3]. They have in the recent past also found applications in the context of various field theoretic dualities [4] and in condensed matter physics, in the description of bosonic topological insulators [5–8].

In this work, the presence of charged fermions gives rise to the nonlocal interaction. The non-relativistic limit of the nonlocal current  $\tilde{J}^{\mu\nu}$  includes the spin density current of the fermions, so  $B_{\mu\nu}$  can be thought of as a gauge field for spin. It was shown recently that such couplings could result in interesting physics such as a linear attractive potential between fermions [9], and attachment of magnetic flux to fermions resulting in anyonic statistics [10]. The flux-attached fermions in such a system are essentially bound states of topological vortices and electrons. Bound states between topological textures and electrons, specifically skyrmion-electron bound states have been recently studied in [11–13]. The field theoretical formalism used in this paper to study vortex-electron interactions can be adapted to the study of such systems.

There is another aspect of gauge fields, which has received a lot of attention in the past two decades, particularly in the context of Topological Insulators (TIs) and Topological Superconductors (TSCs). This aspect is the topological action of the gauge field [14–24]. In 3+1 dimensions, there are two possible dynamical terms of a rank-1 gauge field. One of them is the usual Maxwell term  $-\frac{1}{4}F^{\mu\nu}F_{\mu\nu}$  and the other one is the so called axion term or the  $\theta$ -term, usually written as  $\frac{\theta}{32\pi^2}\epsilon_{\mu\nu\lambda\rho}F^{\mu\nu}F^{\lambda\rho}$  which is topological, meaning that it does not affect the local dynamics of the gauge field. In particular, the importance of the axion term has been highlighted in [14] where the authors have established that the axion term is responsible for the quantized electromagnetic response of time-reversal invariant topological insulators in 3 spatial dimensions. A recent work [4] also explores the dualities between these two the field theories describing 3+1 dimensional TIs and TSCs and their parallels to the particle-vortex duality [25]. The axion term has also been found to be important systems like semimetals [26–31].

In this paper, we will be interested in systems which include features of both of the aforementioned elements, i.e. a rank-2 gauge field  $B_{\mu\nu}$  and the axion term for a rank-1 gauge field. One example of such a system is a construction involving two interfaces, proposed in a recent work by Nogueira et al. [32, 33] for the detection of the Witten effect [34]. In their work, the authors consider a superconductor-topological insulator (TI)-superconductor junction with vortex flux tubes within the two type II superconductors end on the TI surface where there is a change in the value of axion parameter  $\theta$ , leading to a Josephson-Witten effect.

To this end, we will begin with a theory of a mixture of bosons and fermions in the background of a linear combination of the Maxwell and axion terms for a rank-1 gauge field, where the coupling parameter  $\theta$  of the axion term is taken to be a constant. When the axion parameter  $\theta$  is space-time independent, the axion term in the action is a boundary term and therefore one expects it to not play a role in local interactions. In this work, we show that in the presence of flux-tubes in the background,  $\theta$  does affect local interactions.

Theories involving mixtures of bosons and fermions have been proposed in various contexts including high- $T_c$  superconductivity [35–38] to BCS-BEC crossover [39, 40]. In terms of the appearance of flux tubes, our construction bears some resemblance to the Friedberg-Lee model [41, 42], proposed as a description of high- $T_c$  superconductivity. We consider the Abelian Higgs model with the presence additional itinerant fermions minimally coupled to

the  $U(1)$  gauge field. In this theory, vortex flux tubes appear as a consequence of a  $U(1)$  symmetry breaking of the bosonic field. In the symmetry broken phase, the theory can be dualized, using a 3+1 dimensional adaptation of the particle vortex duality, into another theory comprised of flux tubes and a rank-2 gauge field mediating the flux tube-flux tube interaction. The effective interactions between fermions and flux tubes can then be studied by integrating out first the rank-1 gauge field and then the rank-2 gauge field from the partition function describing the system. We do this systematically by bringing the Lagrangian to linear order in the rank-1 gauge field  $A_\mu$ . This is achieved by multiplying the partition function of the system by a Gaussian path integral comprised of the Maxwell term  $F_{\mu\nu}$ , its dual and a new rank-2 antisymmetric field  $\chi_{\mu\nu}$ . After linearizing the Lagrangian, the gauge field  $A_\mu$  is integrated out and the antisymmetric field  $\chi_{\mu\nu}$  is traded away for various nonlocal gauge invariant couplings between the gauge field  $B_{\mu\nu}$  and the fermion current. The gauge field is then integrated out to get the effective interactions. Finally, we perform a dimensional reduction by compactifying the length of the system in the  $z$  direction to obtain a planar action and the resultant interactions in an effective 2+1 dimensional system.

The paper is organized as follows. In section II, we start with the partition function of the system described in terms of a rank-1 gauge field, a scalar condensate and itinerant fermions and show its dual transformation to another partition function in terms of flux tubes and the Kalb-Ramond gauge field [3] along with the usual rank-1 gauge field  $A_\mu$ . In section III, we linearize the Lagrangian in the rank-1 gauge field to obtain a partition function in terms of the rank-1 gauge field and integrate it out to obtain a partition function with the rank-2 gauge fields, fermions and flux tubes. In section IV, the gauge field  $B_{\mu\nu}$  is integrated out to obtain the relevant interactions where it is shown that the axion parameter  $\theta$  appears in all local interactions. In the subsequent section we perform a dimensional reduction to an effective 2+1 dimensional system and demonstrate flux attachment to fermions and finally conclude with a summary and outlook.

## II. THE MODEL

We consider a modified version of the Abelian Higgs model , which includes the parity violating  $\theta$  term or the axion term  $\theta\epsilon^{\mu\nu\alpha\beta}F_{\mu\nu}F_{\alpha\beta}$  alongside the usual Maxwell term  $F_{\mu\nu}F^{\mu\nu}$ . Additionally, we also consider the presence of Dirac fermions in the system which are also

minimally coupled to the gauge field  $A_\mu$ . We work with 3+1 dimensional relativistic quantum field theory, even though the objects and systems we consider are all non-relativistic. There are two important reasons for this approach. One is that the procedure of dualization, which changes the effective variables from the scalar field to the flux tubes and the anti-symmetric tensor gauge field, works best in four dimensions. The other is that electron spin appears naturally in the theory and we do not need to put in an interaction by hand. The coupling of electron spin to the anti-symmetric gauge field then appears automatically from the dualization.

To this end, we start with the Lagrangian

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{\theta}{32\pi^2}\epsilon^{\mu\nu\alpha\beta}F_{\mu\nu}F_{\alpha\beta} + \frac{1}{2}(D_\mu\phi)^\dagger(D^\mu\phi) - V(|\phi|^2) \\ & + \bar{\psi}(i\gamma^\mu\partial_\mu - m)\psi - eA_\mu\bar{\psi}\gamma^\mu\psi, \end{aligned} \quad (1)$$

where  $D_\mu = \partial_\mu + iqA_\mu$ .

We focus on the broken symmetry phase in which  $D_\mu\phi = 0$  and there are flux strings carrying quantized magnetic flux [43]

$$\oint_C dx^\mu A_\mu = -\frac{2n\pi}{q}, \quad (2)$$

where  $n \in \mathbb{Z}$  is the winding number. The description of  $\phi$  in this phase is of the form

$$\phi = vf(|\mathbf{r} - \mathbf{r}_0|) \exp(i\chi(\mathbf{r} - \mathbf{r}_0)), \quad (3)$$

where  $v$  is the vacuum expectation value of the scalar condensate  $\phi$ , while  $f$  is a function of the distance of a point  $\mathbf{r}$  from the core of the flux tube  $\mathbf{r}_0$  with the property that as  $\mathbf{r} \rightarrow \mathbf{r}_0$ ,  $f \rightarrow 0$  and as  $\mathbf{r} \rightarrow \infty$ , we have  $f \rightarrow 1$ . For  $\phi$  to admit a topological solution, its phase  $\chi$  must be a multivalued function. Hence we can split  $\chi$  into a well behaved single valued regular part  $\chi_r$  and a singular part  $\chi_s$  as  $\chi = \chi_r + \chi_s$ . The regular part of the phase  $\chi_r$  can be integrated out by the introduction of an auxiliary field [44–46], to arrive at the dual

Lagrangian away from the core

$$\begin{aligned}
Z = & \int (\mathcal{D}A_\mu \mathcal{D}B_{\mu\nu} \mathcal{D}\bar{\psi} \mathcal{D}\psi \mathcal{D}\chi_s) \\
& \exp \left[ i \int d^4x \left( -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{\theta}{32\pi^2} \epsilon_{\mu\nu\alpha\beta} F^{\mu\nu} F^{\alpha\beta} \right. \right. \\
& \quad + \frac{1}{12v^2} H^{\mu\nu\lambda} H_{\mu\nu\lambda} - \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} B_{\alpha\beta} \partial_\mu \partial_\nu \chi_s \\
& \quad \left. \left. - \frac{q}{2} \epsilon^{\mu\nu\alpha\beta} H_{\nu\alpha\beta} A_\mu + \bar{\psi} (i\gamma^\mu \partial_\mu - m) \psi - e A_\mu \bar{\psi} \gamma^\mu \psi \right) \right], \tag{4}
\end{aligned}$$

where we have written  $H_{\nu\alpha\beta} = \frac{1}{3}(\partial_\nu B_{\alpha\beta} + \partial_\alpha B_{\beta\nu} + \partial_\beta B_{\nu\alpha})$ .

Since the scalar field  $\phi$  vanishes at the core of the flux tube, derivatives on the singular part of its phase  $\chi_s$  are non-commuting. The commutator  $\partial_{[\mu} \partial_{\nu]} \chi_s$  is the vorticity of the flux tube, which can also be represented by the two form field  $\Sigma^{\alpha\beta} = \epsilon^{\mu\nu\alpha\beta} \partial_\mu \partial_\nu \chi_s$ . This two-form also represents the world-sheet of the vortex tubes in  $3 + 1$  dimensions,

$$\Sigma^{\mu\nu} = \int d\sigma_{\rho\lambda} \delta(x - X),$$

where  $X^\mu$  are the coordinates of the tube and  $d\sigma_{\mu\nu}$  is the world-sheet surface element. The integral of the vorticity along a closed loop  $C$  which contains the core of the flux tube is non-vanishing.

### III. DUALIZATION AND EMERGENT COUPLINGS

We start by linearizing the gauge field  $A_\mu$  in the Lagrangian by introducing an anti-symmetric tensor  $\chi_{\mu\nu}$ . To this end, we multiply the partition function of Eq. (4) by the following Gaussian functional

$$Z_\chi = \int D\chi_{\mu\nu} \exp \left( -i \int d^4x (\alpha F^{\mu\nu} + \frac{\beta}{2} \epsilon_{\mu\nu\alpha\beta} F^{\alpha\beta} + \frac{1}{2} \chi_{\mu\nu})^2 \right), \tag{5}$$

and set

$$\alpha^2 - \beta^2 = -\frac{1}{4}, \tag{6}$$

$$\alpha\beta = -\frac{\theta}{32\pi^2}. \tag{7}$$

With this choice of  $\alpha$  and  $\beta$ , the product is the partition function

$$\begin{aligned}
Z_{dual} = & \int (\mathcal{D}\chi_{\mu\nu} \mathcal{D}A_\mu \mathcal{D}B_{\mu\nu} \mathcal{D}\bar{\psi} \mathcal{D}\psi \mathcal{D}\chi_s) \\
& \exp \left[ i \int d^4x \left( -\frac{1}{4} \chi_{\mu\nu} \chi^{\mu\nu} - \alpha F^{\mu\nu} \chi_{\mu\nu} - \frac{\beta}{2} \epsilon_{\mu\nu\alpha\beta} \chi^{\alpha\beta} F^{\mu\nu} \right. \right. \\
& \quad \left. \left. + \frac{1}{12v^2} H^{\mu\nu\lambda} H_{\mu\nu\lambda} - \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} B_{\alpha\beta} \partial_\mu \partial_\nu \chi_s \right. \right. \\
& \quad \left. \left. - \frac{q}{2} \epsilon^{\mu\nu\alpha\beta} \partial_\nu B_{\alpha\beta} A_\mu + \bar{\psi} (i\gamma^\mu \partial_\mu - m) \psi - e A_\mu \bar{\psi} \gamma^\mu \psi \right) \right]. \quad (8)
\end{aligned}$$

The Lagrangian in this partition function is linear in the gauge field  $A_\mu$ , which can be integrated out to produce a  $\delta$ -functional:

$$\begin{aligned}
Z_{dual} = & \int (\mathcal{D}\chi_{\mu\nu} \mathcal{D}B_{\mu\nu} \mathcal{D}\bar{\psi} \mathcal{D}\psi \mathcal{D}\Sigma_{\mu\nu} \mathcal{D}\chi_{\mu\nu}) \delta[C^\mu] \exp \left[ i \int d^4x \left( -\frac{1}{4} \chi_{\mu\nu} \chi^{\mu\nu} \right. \right. \\
& \quad \left. \left. + \frac{1}{12v^2} H_{\mu\nu\lambda} H^{\mu\nu\lambda} - \frac{1}{2} B_{\mu\nu} \Sigma^{\mu\nu} + \bar{\psi} (i\gamma^\mu \partial_\mu - m) \psi \right) \right], \quad (9)
\end{aligned}$$

where

$$C^\mu = -2\alpha \partial_\nu \chi^{\mu\nu} - \beta \epsilon^{\mu\nu\alpha\beta} \partial_\nu \chi_{\alpha\beta} - \frac{q}{2} \epsilon^{\mu\nu\alpha\beta} \partial_\nu B_{\alpha\beta} - e \bar{\psi} \gamma^\mu \psi.$$

To satisfy the  $\delta$  function constraint, we set

$$2\alpha \chi_{\mu\nu} + \beta \epsilon_{\mu\nu\alpha\beta} \chi^{\alpha\beta} = -\frac{q}{2} \epsilon_{\mu\nu\alpha\beta} B^{\alpha\beta} + e J_{\mu\nu}, \quad (10)$$

where we have defined the nonlocal current  $J_{\mu\nu} = \frac{1}{\square} \partial_{[\mu} j_{\nu]}$  in terms of the fermion current  $j^\mu \equiv \bar{\psi} \gamma^\mu \psi$ .

We contract Eq. (10) with  $\epsilon^{\mu\nu\rho\sigma}$  and solve for  $\chi_{\mu\nu}$  to get

$$\chi_{\mu\nu} = \frac{\alpha}{2(\alpha^2 + \beta^2)} \left( -q \tilde{B}_{\mu\nu} + e J_{\mu\nu} \right) - \frac{\beta}{2(\alpha^2 + \beta^2)} \left( q B_{\mu\nu} + e \tilde{J}_{\mu\nu} \right), \quad (11)$$

where we have defined

$$\tilde{B}_{\mu\nu} \equiv \frac{1}{2} \epsilon_{\mu\nu\alpha\beta} B^{\alpha\beta}, \quad (12)$$

$$\tilde{J}_{\mu\nu} \equiv \frac{1}{2} \epsilon_{\mu\nu\alpha\beta} J^{\alpha\beta}. \quad (13)$$

Substituting  $\chi_{\mu\nu}$  from Eq. (11) back into the partition function (9), we get

$$\begin{aligned}
Z_{\text{dual}} = & \int (\mathcal{D}\chi_{\mu\nu} \mathcal{D}B_{\mu\nu} \mathcal{D}\bar{\psi} \mathcal{D}\psi \mathcal{D}\Sigma_{\mu\nu} \mathcal{D}\chi_{\mu\nu}) \\
& \exp \left[ i \int d^4x \left[ -\frac{1}{4(\alpha^2 + \beta^2)^2} \left[ \frac{1}{16} (q^2 v^2 B_{\mu\nu} B^{\mu\nu} - e^2 J_{\mu\nu} J^{\mu\nu}) \right. \right. \right. \\
& \quad \left. \left. + \frac{qve}{8} \tilde{B}_{\mu\nu} J^{\mu\nu} - \frac{\alpha\beta}{2} \left( -q^2 v^2 B_{\mu\nu} \tilde{B}^{\mu\nu} + e^2 J_{\mu\nu} \tilde{J}^{\mu\nu} + qve B_{\mu\nu} J^{\mu\nu} \right) \right] \right. \\
& \quad \left. \left. + \frac{1}{12} H_{\mu\nu\lambda} H^{\mu\nu\lambda} - \frac{v}{2} B_{\mu\nu} \Sigma^{\mu\nu} + \bar{\psi} (i\gamma^\mu \partial_\mu - m) \psi \right] \right], \tag{14}
\end{aligned}$$

where we have substituted  $\alpha^2 - \beta^2 = -\frac{1}{4}$  and rescaled  $B_{\mu\nu} \rightarrow v B_{\mu\nu}$ .

We choose to keep the coefficient  $(\alpha^2 + \beta^2)^{-2}$  intact for notational simplicity and only replace it in terms of  $\theta$  in the end. The Lagrangian presented above is the dual counterpart to the one described in Eq. (1) and describes a system consisting of flux tubes whose world-sheets are described by the tensor  $\Sigma^{\mu\nu}$ . These flux tubes are coupled to the two-form gauge field  $B_{\mu\nu}$  which mediates the interaction between two flux tubes. Additionally,  $B_{\mu\nu}$  is also coupled to a non-local fermion current described by the antisymmetric tensor  $J^{\mu\nu}$ . This kind of coupling was first described in [2] and leads to interactions between the flux tubes and fermions, mediated by the two-form gauge field.

#### IV. EFFECTIVE INTERACTIONS IN THE DUAL THEORY

Let us now calculate the effective interactions as described by the dual Lagrangian in Eq. (14). To get the effective interaction between flux tubes, between fermions, or between fermions and flux tubes, we integrate out the two-form gauge field  $B_{\mu\nu}$  from the path integral. Since the dual Lagrangian is quadratic in  $B_{\mu\nu}$ , this integral can be done exactly. In addition, since  $B_{\mu\nu}$  is massive in the dual Lagrangian, a gauge fixing term not necessary. We can then segregate the part of the partition function which depends on  $B_{\mu\nu}$  and write it as

$$\begin{aligned}
Z_{\text{dual}}^B = & \int \mathcal{D}B_{\mu\nu} \exp \left[ i \int d^4x \left[ \frac{1}{12} H_{\mu\nu\lambda} H^{\mu\nu\lambda} + \frac{1}{4(\alpha^2 + \beta^2)^2} \left( -\frac{q^2 v^2}{16} B_{\mu\nu} B^{\mu\nu} - \frac{qve}{8} B_{\mu\nu} \tilde{J}^{\mu\nu} \right. \right. \right. \\
& \quad \left. \left. + \frac{\theta}{32\pi^2} \left( \frac{q^2 v^2}{2} B_{\mu\nu} \tilde{B}^{\mu\nu} - \frac{1}{2} qve B_{\mu\nu} J^{\mu\nu} \right) \right) - \frac{v}{2} B_{\mu\nu} \Sigma^{\mu\nu} \right], \tag{15}
\end{aligned}$$



which can be recast in the form

$$Z_{dual}^B = \int \mathcal{D}B_{\mu\nu} \exp \left[ i \int d^4x \left[ -\frac{1}{4} B_{\mu\nu} K^{\mu\nu\alpha\beta} B_{\alpha\beta} - \frac{1}{2} B_{\mu\nu} P^{\mu\nu} \right] \right], \quad (16)$$

where

$$K^{\mu\nu\alpha\beta} = \frac{1}{2} \left( \square + M^2 \right) g^{\mu[\alpha} g^{\beta]\nu} - \frac{\theta}{8\pi^2} M^2 \epsilon^{\mu\nu\alpha\beta} B_{\alpha\beta} + \frac{1}{2} \left( g^{\nu[\alpha} g^{\beta]\sigma} \partial_\sigma \partial^\mu - \mu \leftrightarrow \nu \right), \quad (17)$$

and

$$P^{\mu\nu} = \left( \frac{qve}{16(\alpha^2 + \beta^2)^2} \tilde{J}^{\mu\nu} + \frac{qve\theta}{128\pi^2(\alpha^2 + \beta^2)^2} J^{\mu\nu} + v\Sigma^{\mu\nu} \right). \quad (18)$$

We have also written

$$M^2 = \frac{q^2 v^2}{16(\alpha^2 + \beta^2)^2} = \frac{q^2 v^2}{\left(1 + \frac{\theta^2}{16\pi^4}\right)}. \quad (19)$$

Integrating out  $B_{\mu\nu}$  from Eq. (16) we get the following expression for the effective interaction Lagrangian,

$$\mathcal{L}_{dual}^B = \int \frac{d^4k}{(2\pi)^4} \left( -\frac{1}{4} P^{\mu\nu}(-k) G_{\mu\nu\alpha\beta}(k) P^{\alpha\beta}(k) \right), \quad (20)$$

where  $P^{\alpha\beta}(k)$  is the Fourier transformation of Eq. (18) and  $G_{\mu\nu\alpha\beta}(k)$  is the inverse of the Fourier transformed  $K^{\mu\nu\alpha\beta}$  such that

$$K^{\mu\nu\alpha\beta}(k) G_{\mu\nu\alpha'\beta'}(k) = \delta_{[\alpha}^{\mu} \delta_{\beta']}^{\nu]}. \quad (21)$$

The propagator is easily calculated,

$$G_{\mu\nu\alpha\beta}(k) = \frac{1}{(-k^2 + M^2)^2 + \frac{\theta^2 M^4}{16\pi^4}} \left( (-k^2 + M^2) g_{\mu[\alpha} g_{\beta]\nu} - \frac{\theta}{4\pi^2} M^2 \epsilon_{\mu\nu\alpha\beta} \right) \quad (22)$$

$$- \left( 1 - \frac{k^2}{M^2} \right) \left( g_{\nu[\alpha} g_{\beta]\sigma} k^\sigma k_\mu - \mu \leftrightarrow \nu \right), \quad (23)$$

where we have used Eq. (7) to substitute for the value of  $\alpha\beta$ .

From Eqs. (18), (20), and (23), we can calculate the effective interactions, which can be split into three parts as follows:

$$\begin{aligned} \mathcal{L}_{dual}^B = & \int \frac{d^4k}{(2\pi)^4} \left( \frac{qve}{16(\alpha^2 + \beta^2)^2} \tilde{J}^{\mu\nu}(-k) + \frac{qve\theta}{128\pi^2(\alpha^2 + \beta^2)^2} J^{\mu\nu}(-k) \right) G_{\mu\nu\alpha\beta}(k) \\ & \times \left( \frac{qve}{16(\alpha^2 + \beta^2)^2} \tilde{J}^{\alpha\beta}(k) + \frac{qve\theta}{128\pi^2(\alpha^2 + \beta^2)^2} J^{\alpha\beta}(k) \right) \\ & + v^2 \int \frac{d^4k}{(2\pi)^4} \Sigma^{\mu\nu}(-k) G_{\mu\nu\alpha\beta}(k) \Sigma^{\alpha\beta}(k) + \dots \end{aligned} \quad (24)$$

The first integral in the above equation corresponds to the effective fermion-fermion interaction. The second integral describes the interaction between flux tubes while the ellipses represent the  $J - \Sigma$  interaction terms.

Evaluating the first integral, we get

$$\frac{e^2}{4(\alpha^2 + \beta^2)^2} \int \frac{d^4k}{(2\pi)^4} \left[ j_\mu(-k) \left[ \frac{M^2}{(-k^2 + M^2)^2 + \frac{\theta^2 M^4}{4\pi^4}} \frac{1}{k^2} \left( k^2 - M^2 + \frac{\theta^2}{128\pi^4} (k^2 + 7M^2) \right) \right] j^\mu(k) \right]. \quad (25)$$

The above expression gives the effective interaction between two fermions in the system. Clearly, in the presence of a non-zero  $\theta$  term, the interaction is different from that of a Coulombic or a Yukawa interaction. In addition, it is clear that  $\theta$  also contributes to the mass of the  $B$  field. It is also interesting to note that  $\theta$  and  $M$  appear together in this expression. Thus the effect of the axion term on fermionic interactions depends on the scale at which condensation occurs, as well as the charge of the condensate. It is important to note here that the axion term contributes to the dynamics of fermions in the bulk despite being a topological term and it also shifts the pole of the propagator. The position of the pole in the complex plane now depends on the value of  $\theta$ . Additionally, as  $\theta \rightarrow 0$ , the resulting effective interaction between fermions reduces to

$$\int \frac{d^4k}{(2\pi)^4} \frac{e^2}{2(\alpha^4 + \beta^4)} \left[ j_\mu(-k) \left( \frac{1}{k^2 - M^2} - \frac{1}{k^2} \right) j^\mu(k) \right], \quad (26)$$

which agrees with the result obtained in [47] where the  $\theta$  term was not considered.

We will now focus on the second integral in Eq. (24), which is

$$2v^2 \int \frac{d^4k}{(2\pi)^4} \Sigma_{\mu\nu}(-k) \left[ \frac{1}{(-k^2 + M^2)^2 + \frac{\theta^2 M^4}{16\pi^4}} \left( (-k^2 + M^2) \Sigma^{\mu\nu}(k) - \frac{\theta M^2}{2\pi^2} \tilde{\Sigma}^{\mu\nu}(k) \right) \right]. \quad (27)$$

As stated before, this term gives the effective interaction between two flux tubes in the system. It can be seen that the effect of the axion term is two-fold. Firstly, it has shifted the pole of the propagator of the gauge field  $B_{\mu\nu}$ . And secondly, there is a new kind of interaction between the world-sheet tensor  $\Sigma_{\mu\nu}$  and its dual which is generated purely because of the axion term.

To obtain the fermion-flux tubes interactions, we evaluate the terms represented by the

ellipses in Eq. (24), remembering the gauge invariance condition  $\partial_\mu \Sigma^{\mu\nu} = 0$ , to get

$$\frac{4eM^2}{q} \int \frac{d^4k}{(2\pi)^4} j_\nu(k) \frac{1}{(-k^2 + M^2)^2 + \frac{\theta^2 M^4}{16\pi^4}} \left[ -\frac{k_\mu}{k^2} \left( (-k^2 + M^2) \left( 1 + \frac{\theta^2 M^2}{8\pi^2} \right) \right) \right] \tilde{\Sigma}^{\mu\nu}(-k). \quad (28)$$

It can be seen again that the effect of the axion term is to change the interaction between the fermion current  $j_\mu$  and the dual of flux tube given by  $\Sigma^{\mu\nu}$ . The interaction between these two fields in the absence of the  $\theta$  term was obtained in [47]. We note that all of these interactions, fermion-fermion, flux tube-flux tube and the fermion-flux tube have the common feature of having a term proportional to  $\theta M^2$  which means that the axion term has an observable effect when there is a charged massive scalar condensate in the background.

## V. DIMENSIONAL REDUCTION AND EFFECTIVE (2+1)D THEORY

In the previous sections we have shown how the presence of a non-zero  $\theta$  term affects local interactions in the presence of a massive scalar condensate. It was shown in [9] that with for vanishing  $\theta$ , one obtains magnetic flux attachment to electrons present in the system when the theory is dimensionally reduced to 2+1 dimensions. Motivated by this, we would like to investigate if the  $\theta$  term also plays a role in flux attachment.

We follow the steps outlined in [10] to perform a dimensional reduction of our system. We assume that the system has a small extension  $L$  along the third direction. In addition, we also assume that all the fields in the Lagrangian are independent of the third coordinate  $z$ . The resulting theory in the small  $L$  limit will be the effective dual theory in  $(2+1)$  dimensions. We begin by segregating the  $(0, 1, 2)$  components from the 3 component of all the fields in the Lagrangian

$$\begin{aligned} -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} &= -\frac{1}{4} F_{ab} F^{ab} + \partial_a \phi \partial^a \phi, \quad (29) \\ \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi - eA_\mu j^\mu &= \bar{\psi}(i\gamma^a \partial_a - m)\psi - eA_a j^a - e\phi j^3, \\ \frac{1}{12} H^{\mu\nu\lambda} H_{\mu\nu\lambda} &= -\frac{1}{4} H^{ab} H_{ab} + \frac{1}{12} H^{abc} H_{abc}, \\ B^{\mu\nu} B_{\mu\nu} &= 2B_a B^a + B_{ab} B^{ab} \quad (30) \end{aligned}$$

where  $\epsilon_{abc} = \epsilon_{abc3}$  and  $\phi = A_z$ . Additionally, the wave operator  $\square = \partial_\mu \partial^\mu$  now takes the form  $\Delta = \partial_a \partial^a$ . The Latin indices  $a, b, c, \dots$  represent the  $(2+1)$ -dimensional components and take the values  $0, 1, 2$ . Substituting the above expressions into the dual partition

function, we get

$$\begin{aligned}
Z_{\text{dual}} = & \int (\mathcal{D}\chi_{\mu\nu} \mathcal{D}B_{\mu\nu} \mathcal{D}\bar{\psi} \mathcal{D}\psi \mathcal{D}\Sigma_{\mu\nu} \mathcal{D}\chi_{\mu\nu} \mathcal{D}A_\mu) \\
& \exp \left[ i \int d^4x \left[ -\frac{1}{4} F_{ab} F^{ab} + \partial_a \phi \partial^a \phi + \bar{\psi} (i\gamma^a \partial_a - m) \psi - \kappa \tilde{e} A_a j^a - \kappa \tilde{e} \phi j^3 \right] \right. \\
& + \frac{1}{4(\alpha^2 + \beta^2)^2} \left[ -\frac{q^2 v^2}{16} (2B_a B^a + B_{ab} B^{ab}) - \frac{qve}{4} \left( \tilde{B}_a \frac{1}{\Delta} \partial^a j^3 + \frac{1}{2} \epsilon_{abc} B^a \partial^b \frac{1}{\Delta} j^c \right) \right. \\
& + \left. \left. \frac{\theta}{32\pi^2} \left( +4q^2 v^2 \tilde{B}_c B^c + 4qve \left( \partial^a B_a \frac{1}{\Delta} j^3 - \partial^a B_{ab} \frac{1}{\Delta} j^b \right) \right) \right] \right. \\
& \left. - \frac{1}{4} H^{ab} H_{ab} + \frac{1}{12} H^{abc} H_{abc} - v B_a \Sigma^a - \frac{v}{2} B_{ab} \Sigma^{ab} \right], \tag{31}
\end{aligned}$$

where we have used the following relation to restore the photon field  $A_\mu$  in Eq. (14):

$$\int \mathcal{D}A_\mu \exp \left( i \int d^4x \left( -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + A_\mu K^\mu \right) \right) \propto \int \exp \left( -\frac{i}{2} \int d^4x \left( K_\mu \frac{1}{\square} K^\mu \right) \right). \tag{32}$$

We consider the special case when the fermion current is confined to the  $X - Y$  plane, and set  $j_3 = 0$ , to get

$$\begin{aligned}
Z_{\text{dual}} \Big|_{j_3=0} = & \int (\mathcal{D}B_{ab} \mathcal{D}B_a \mathcal{D}\bar{\psi} \mathcal{D}\psi \mathcal{D}\Sigma_{ab} \mathcal{D}\Sigma_a \mathcal{D}A_a) \\
& \exp \left[ i \int d^4x \left[ -\frac{1}{4} F_{ab} F^{ab} + \partial_a \phi \partial^a \phi + \bar{\psi} (i\gamma^a \partial_a - m) \psi - \kappa \tilde{e} A_a j^a \right. \right. \\
& - \frac{M^2}{4} (2B_a B^a + B_{ab} B^{ab}) - \frac{qv\kappa\tilde{e}}{2} \epsilon_{abc} B^a \partial^b \frac{1}{\Delta} j^c \\
& + \frac{\theta M^2}{2\pi^2} \tilde{B}_c B^c - \frac{2\theta\kappa\tilde{e}}{\pi^2} \partial^a B_{ab} \frac{1}{\Delta} j^b \\
& \left. \left. - \frac{1}{4} H^{ab} H_{ab} + \frac{1}{12} H^{abc} H_{abc} - v B_a \Sigma^a - \frac{v}{2} B_{ab} \Sigma^{ab} \right] \right], \tag{33}
\end{aligned}$$

where  $\kappa = L^{-\frac{1}{2}}$  and the ‘‘screened’’ charge  $\tilde{e}$  is defined as

$$\tilde{e} = \frac{e}{16(\alpha^2 + \beta^2)} = \frac{Me}{2qv}. \tag{34}$$

We wish to integrate out the  $B_a$  and the  $B_{ab}$  fields from the partition function to obtain the effective electrodynamics on the thin layer. To this end, we integrate the two fields separately, treating the mixed term  $\frac{\theta M^2}{2\pi^2} \tilde{B}_c B^c$  perturbatively. We begin by writing out the

terms in the partition function containing the relevant fields

$$\begin{aligned}
Z_{\text{dual}}^B \Big|_{j_3=0} &= \int (\mathcal{D}B_{ab} \mathcal{D}B_a) \\
&\exp \left[ i \int d^3x \left( -\frac{1}{4} H^{ab} H_{ab} - \frac{M^2}{2} B_a B^a - \frac{qv\kappa\tilde{\epsilon}}{2} \epsilon_{abc} B^a \partial^b \frac{1}{\Delta} j^c - v B_a \Sigma^a \right) \right. \\
&\quad + \left( \frac{1}{12} H^{abc} H_{abc} - \frac{M^2}{2} B_{ab} B^{ab} - \frac{2\theta\kappa\tilde{\epsilon}}{\pi^2} B_{ab} \partial^a \frac{1}{\Delta} j^b + B_{ab} \Sigma^{ab} \right) \\
&\quad \left. + \frac{\theta M^2}{2\pi^2} \tilde{B}_c B^c \right] \tag{35}
\end{aligned}$$

where  $\kappa = 1/\sqrt{L}$  takes into account the rescaling of charge upon going to a smaller dimension [10]. The propagators for  $B_a$  and  $B_{ab}$  are respectively

$$\Delta_{ab}^0 = \frac{1}{-k^2 + M^2} \left( g_{ab} - \frac{k_a k_b}{M^2} \right) \tag{36}$$

and

$$\Delta_{ab,cd}^0 = \frac{1}{-k^2 + M^2} \left[ (g_{ac} g_{bd} - g_{ad} g_{bc}) - \frac{g_{ac} k_b k_d - g_{ad} k_b k_c}{M^2} \right]. \tag{37}$$

The total propagator including the mixed term can be calculated from an infinite sum [48],

$$\Delta_{ab} = \Delta_{ab}^0 + \Delta_{ac}^0 \Theta_{cd} \Delta_{db}^0 + \Delta_{ac}^0 \Theta_{cd} \Delta_{de}^0 \Theta_{ef} \Delta_{fb}^0 + \dots \tag{38}$$

where

$$\Theta_{ab} = V_{a,cd} \Delta_{cd,ef}^0 V_{ef,b}, \tag{39}$$

with  $V_{ab,c} = V_{c,ab} = \frac{\theta M^2}{2\pi^2} \epsilon_{abc}$ . The total propagator then is

$$\Delta_{ab} = -\frac{g_{ab} - \frac{k_a k_b}{M^2}}{k^2 - (M^2 + \mu^2)}, \tag{40}$$

where

$$\mu^2 = \frac{\theta^2 M^2}{8\pi^4}. \tag{41}$$

Having found the effective propagator for  $B_a$ , we can write down the vortex-fermion and the fermion-fermion interaction by integrating out the  $B$  fields from the partition function of Eq. (35),

$$Z_{\text{dual}}^B \Big|_{j_3=0} \propto \int \exp \left[ -\frac{i}{2} \int d^3x d^3y \left( P_a(x) \Delta^{ab}(x-y) P_b(y) \right) \right], \tag{42}$$

where we have written

$$P_a(x) = -\frac{qv\kappa\tilde{e}}{2}\epsilon_{abc}\partial^b\frac{1}{\Delta}j^c - v\Sigma_a \equiv -\frac{qv\kappa\tilde{e}}{2}\tilde{j}_a - v\Sigma_a. \quad (43)$$

The vorticity  $\Sigma^a$  can be split into two parts, one coming from the singular part of the scalar field  $\phi$  that we started with in Eq. (4) and another coming from any external magnetic field that is applied, hence  $\Sigma^a = \epsilon^{abc}\partial_b\partial_c\chi^s + \Sigma_e^a \equiv \epsilon^{abc}\partial_b S_c + \Sigma_e^a$ . The partition function in Eq. (42) can then be rewritten as

$$\int \exp \left[ -\frac{i}{2} \int d^3x d^3y \left( \frac{q^2 v^2 \kappa \tilde{e}^2}{4} \tilde{j}^a \frac{1}{\Delta + (M^2 + \mu^2)} \tilde{j}_a + qv^2 \kappa \tilde{e} S^d \frac{1}{\Delta + (M^2 + \mu^2)} \epsilon_{bcd} \partial^c \tilde{j}^b \right. \right. \\ \left. \left. + \frac{v^2}{2} G_{ab} \frac{1}{\Delta^2 + (M^2 + \mu^2)} G^{ab} + v^2 \Sigma_e^a \frac{1}{\Delta^2 + (M^2 + \mu^2)} \Sigma_{ea} \right. \right. \\ \left. \left. + v^2 \epsilon^{abc} G_{bc} \frac{1}{\Delta^2 + (M^2 + \mu^2)} \Sigma_{ea} + qv^2 \kappa \tilde{e} \Sigma_e^a \frac{1}{\Delta^2 + (M^2 + \mu^2)} \tilde{j}_a \right) \right], \quad (44)$$

where we have written  $G_{ab} = \partial_a S_b - \partial_b S_a$ . Collecting the terms with the field  $S^a$ , and using the fact that at low energies  $\frac{1}{\Delta^2 + (M^2 + \mu^2)} \approx \frac{1}{M^2 + \mu^2}$ , we get for the vortex interaction

$$\int \exp \left[ -\frac{i}{2} \int d^3x \left[ qv^2 \kappa \tilde{e} S^d \frac{1}{(M^2 + \mu^2)} \epsilon_{bcd} \partial^c \tilde{j}^b + \frac{v^2}{2} G_{ab} \frac{1}{(M^2 + \mu^2)} G^{ab} \right. \right. \\ \left. \left. + v^2 \epsilon^{abc} G_{bc} \frac{1}{(M^2 + \mu^2)} \Sigma_{ea} \right] \right]. \quad (45)$$

We rescale  $S^a \rightarrow \frac{v}{\sqrt{M^2 + \mu^2}} S^a$  and extract the Lagrangian from the above partition function as

$$\mathcal{L}_S = -\frac{1}{4} G_{ab} G^{ab} + S_c \left( \frac{qv\kappa\tilde{e}}{\sqrt{M^2 + \mu^2}} j^c + \frac{v}{\sqrt{M^2 + \mu^2}} \epsilon^{abc} \partial_a \Sigma_{eb} \right). \quad (46)$$

Following [10], we can say that the equation of motion derived from the above Lagrangian shows that the number of vortices present in the system, including those due to the external magnetic field, is exactly the same as the charge,

$$\epsilon^{bc} \partial_b S_c + \frac{v}{q\sqrt{M^2 + \mu^2}} \Sigma_{e0} = -\frac{v\kappa\tilde{e}}{\sqrt{M^2 + \mu^2}} \tilde{j}_0. \quad (47)$$

## VI. SUMMARY AND OUTLOOK

In this paper, we have considered vortex strings in an Abelian Higgs model coupled to  $\theta$ -electrodynamics. In addition, we have taken the electromagnetic field to be also coupled to charged fermions. Treating this system as a four-dimensional relativistic field theory,

we have dualized the fields to write the path integral in terms of a rank-2 tensor field  $B_{\mu\nu}$  which couples to the itinerant fermions via a nonlocal interaction term. The emergence of the 2-form gauge field  $B_{\mu\nu}$  is not unexpected, as it is well known that such a field mediates the interaction between vortex strings. The nonlocal interaction, although less well known, emerges in a straightforward manner from the path integral of the model. Using these, we calculate the effective interaction between fermions, between strings, and between fermions and strings. These may be relevant to type-II superconductors with itinerant fermions.

We find that the parameter  $\theta$  appears in all of these interactions, modifying the pole of the propagator, as well as the overall strength of the interaction. Since the axion term is a total divergence, it is not expected to affect local physics. It is thus a surprise that, as we have shown in this paper, the presence of vortex strings and itinerant fermions in the system brings the effect of the axion term down to local dynamics. This is our main result. Intuitively, we can understand this by realizing that the existence of the string creates a circle at infinity from which the total divergence term gets a nontrivial contribution for the path integral. However, we are not aware of any method of seeing this contribution in local dynamics other than the dualization procedure used in this paper. This result also justifies our use of 3+1-dimensional relativistic quantum field theory to analyze this system, even though we may be thinking of its application in superconductors, as the dualization is specific to 3+1-dimensional relativistic field theories.

We have also considered a dimensionally reduced version of our system, corresponding to all the fields being restricted to a thin layer. We know that in the absence of the axion term, the reduced system shows that charges are attached to the vortex strings in a kind of flux attachment. Our calculation with the axion term finds the same result — the density of vortices is exactly the same the density of the (dual) charge. It should be possible, using the same techniques, to calculate the effective interactions between vortices in thin slabs of type-II superconductors when an axion term is present. We leave these and similar questions for future work.

## VII. ACKNOWLEDGEMENTS

S.K. thanks Department of Science and Technology, India for Junior Research Fellowship, and Shantonu Mukherjee for helpful discussions.

- 
- [1] R. Peierls, Zur theorie des diamagnetismus von leitungselektronen, Zeitschrift für Physik **80**, 763 (1933).
  - [2] I. D. Choudhury, M. C. Diamantini, G. Guarnaccia, and et al., 4d topological mass by gauging spin, Journal of High Energy Physics **2015**, 81 (2015).
  - [3] M. Kalb and P. Ramond, Classical direct interstring action, Phys. Rev. D **9**, 2273 (1974).
  - [4] J. Murugan and H. Nastase, A 4d duality web (2021), arXiv:2103.12667 [hep-th].
  - [5] G. Y. Cho and J. E. Moore, Topological bf field theory description of topological insulators, Annals of Physics **326**, 1515 (2011).
  - [6] A. Vishwanath and T. Senthil, Physics of three-dimensional bosonic topological insulators: Surface-deconfined criticality and quantized magnetoelectric effect, Phys. Rev. X **3**, 011016 (2013).
  - [7] B. Moy, H. Goldman, R. Sohal, and E. Fradkin, Theory of oblique topological insulators, SciPost Phys. **14**, 023 (2023).
  - [8] A. Tiwari, X. Chen, T. Neupert, L. H. Santos, S. Ryu, C. Chamon, and C. Mudry, Topological bf theory of the quantum hydrodynamics of incompressible polar fluids, Physical Review B **90**, 235118 (2014).
  - [9] S. Mukherjee and A. Lahiri, Spin gauge theory, duality and fermion pairing, Journal of High Energy Physics **2022**, 68 (2022).
  - [10] S. Mukherjee and A. Lahiri, Spin-flux attachment by dimensional reduction of vortices, Nuclear Physics B **986**, 116050 (2023).
  - [11] N. Davier and R. Ramazashvili, Electron states bound to a texture in a néel antiferromagnet, Phys. Rev. B **109**, 224425 (2024).
  - [12] N. Davier and R. Ramazashvili, Texture-induced spin-orbit coupling and skyrmion-electron bound states in a néel antiferromagnet, Phys. Rev. B **107**, 014406 (2023).



- [13] F. A. Gómez Albarracín, H. D. Rosales, M. Udagawa, P. Pujol, and L. D. C. Jaubert, From chiral spin liquids to skyrmion fluids and crystals, and their interplay with itinerant electrons, *Phys. Rev. B* **109**, 064426 (2024).
- [14] X.-L. Qi, T. L. Hughes, and S.-C. Zhang, Topological field theory of time-reversal invariant insulators, *Phys. Rev. B* **78**, 195424 (2008).
- [15] L. Wu, M. Salehi, N. Koirala, J. Moon, S. Oh, and N. P. Armitage, Quantized faraday and kerr rotation and axion electrodynamics of a 3d topological insulator, *Science* **354**, 1124 (2016), <https://www.science.org/doi/pdf/10.1126/science.aaf5541>.
- [16] A. Sekine and K. Nomura, Axion electrodynamics in topological materials, *Journal of Applied Physics* **129**, 141101 (2021).
- [17] Y.-L. Lee, H. C. Park, J. Ihm, and Y.-W. Son, Manifestation of axion electrodynamics through magnetic ordering on edges of a topological insulator, *Proceedings of the National Academy of Sciences* **112**, 11514 (2015), <https://www.pnas.org/doi/pdf/10.1073/pnas.1515664112>.
- [18] K. Shiozaki and S. Fujimoto, Dynamical axion in topological superconductors and superfluids, *Phys. Rev. B* **89**, 054506 (2014).
- [19] A. Martín-Ruiz, M. Cambiaso, and L. F. Urrutia, Axion electrodynamics in magnetoelectric media, in *Chirality, Magnetism and Magnetoelectricity: Separate Phenomena and Joint Effects in Metamaterials* edited by E. Kamenetskii (Springer International Publishing, Cham, 2021) pp. 459–492.
- [20] K. M. Fijalkowski, N. Liu, M. Hartl, M. Winnerlein, P. Mandal, A. Coschizza, A. Fothergill, S. Grauer, S. Schreyeck, K. Brunner, M. Greiter, R. Thomale, C. Gould, and L. W. Molenkamp, Any axion insulator must be a bulk three-dimensional topological insulator, *Phys. Rev. B* **103**, 235111 (2021).
- [21] J. Wang, B. Lian, and S.-C. Zhang, Dynamical axion field in a magnetic topological insulator superlattice, *Phys. Rev. B* **93**, 045115 (2016).
- [22] A. Chan, T. L. Hughes, S. Ryu, and E. Fradkin, Effective field theories for topological insulators by functional bosonization, *Phys. Rev. B* **87**, 085132 (2013).
- [23] J. Schultz, F. S. Nogueira, B. Büchner, J. van den Brink, and A. Lubk, Axion Mie theory of electron energy loss spectroscopy in topological insulators, *SciPost Phys. Core* **4**, 023 (2021).
- [24] S. Ryu, J. E. Moore, and A. W. W. Ludwig, Electromagnetic and gravitational responses and anomalies in topological insulators and superconductors, *Phys. Rev. B* **85**, 045104 (2012).

- [25] J. Murugan and H. Nastase, Particle-vortex duality in topological insulators and superconductors, *Journal of High Energy Physics* **2017**, 159 (2017).
- [26] Y. Chen, S. Wu, and A. A. Burkov, Axion response in weyl semimetals, *Phys. Rev. B* **88**, 125105 (2013).
- [27] J. Gooth, B. Bradlyn, S. Honnali, *et al.*, Axionic charge-density wave in the weyl semimetal (TaSe<sub>4</sub>)<sub>2</sub>i, *Nature* **575**, 315 (2019).
- [28] D. Schmeltzer, The hall conductivity in weyl semimetals as a tool to detect the axion field, *Annals of Physics* **420**, 168237 (2020).
- [29] E. Mottola, A. V. Sadofyev, and A. Stergiou, Axions and superfluidity in weyl semimetals, *Phys. Rev. B* **109**, 134512 (2024).
- [30] M. M. Vazifeh and M. Franz, Electromagnetic response of weyl semimetals, *Phys. Rev. Lett.* **111**, 027201 (2013).
- [31] Z. Wang and S.-C. Zhang, Chiral anomaly, charge density waves, and axion strings from weyl semimetals, *Phys. Rev. B* **87**, 161107 (2013).
- [32] F. S. Nogueira, Z. Nussinov, and J. van den Brink, Josephson currents induced by the witten effect, *Phys. Rev. Lett.* **117**, 167002 (2016).
- [33] F. S. Nogueira, Z. Nussinov, and J. van den Brink, Duality of a compact topological superconductor model and the witten effect, *Phys. Rev. D* **94**, 085003 (2016).
- [34] E. Witten, Dyons of charge  $e\theta/2\pi$ , *Physics Letters B* **86**, 283 (1979).
- [35] P. Salas, M. Fortes, M. Solís, and F. Sevilla, Specific heat of underdoped cuprate superconductors from a phenomenological layered boson-fermion model, *Physica C: Superconductivity and its Applications* **524**, 37 (2016).
- [36] G. Pawłowski, R. Micnas, and S. Robaszkiewicz, Effects of disorder on superconductivity of systems with coexisting itinerant electrons and local pairs, *Phys. Rev. B* **81**, 064514 (2010).
- [37] R. Micnas, S. Robaszkiewicz, and A. Bussmann-Holder, Anisotropic superconductivity in systems with coexisting electrons and local pairs, *Phys. Rev. B* **66**, 104516 (2002).
- [38] J. Ranninger and J. Robin, The boson-fermion model of high- $T_c$  superconductivity. doping dependence, *Physica C: Superconductivity* **253**, 279 (1995).
- [39] J. Deng, A. Schmitt, and Q. Wang, Relativistic bcs-bec crossover in a boson-fermion model, *Phys. Rev. D* **76**, 034013 (2007).

- [40] M. M. Maška and N. Trivedi, Temperature-driven bcs-bec crossover and cooper-paired metallic phase in coupled boson-fermion systems, *Phys. Rev. B* **102**, 144506 (2020).
- [41] R. Friedberg and T. D. Lee, Gap energy and long-range order in the boson-fermion model of superconductivity, *Phys. Rev. B* **40**, 6745 (1989).
- [42] R. Friedberg, T. D. Lee, and H. C. Ren, Coherence length and vortex filament in the boson-fermion model of superconductivity, *Phys. Rev. B* **42**, 4122 (1990).
- [43] H. Nielsen and P. Olesen, Vortex-line models for dual strings, *Nuclear Physics B* **61**, 45 (1973).
- [44] M. Mathur and H. S. Sharatchandra, Manifestly lorentz-covariant local quantum field theory of dyons via dual matter fields, *Phys. Rev. Lett.* **66**, 3097 (1991).
- [45] K. Lee, Dual formulation of cosmic strings and vortices, *Phys. Rev. D* **48**, 2493 (1993).
- [46] E. T. Akhmedov, M. N. Chernodub, M. I. Polikarpov, and M. A. Zubkov, Quantum theory of strings in an abelian higgs model, *Phys. Rev. D* **53**, 2087 (1996).
- [47] S. Mukherjee and A. Lahiri, Emergent vortex–electron interaction from dualization, *Annals of Physics* **418**, 168167 (2020).
- [48] T. J. Allen, M. J. Bowick, and A. Lahiri, Topological mass generation in (3+1)-dimensions, *Mod. Phys. Lett. A* **6**, 559 (1991).