Evaluating the Effectiveness of Public Policies on COVID-19 Containment: A PSM-DID Approach

Zihan Wang

Department of Statistics and Data Science, Tsinghua University, Beijing 100084, China

Abstract

The implementation of public policies is crucial in controlling the spread of COVID-19. However, the effectiveness of different policies can vary across different aspects of epidemic containment. Identifying the most effective policies is essential for providing informed recommendations for pandemic control. This paper examines the relationship between various public policy responses and their impact on COVID-19 containment. Using the propensity score matching-difference in differences (PSM-DID) model to address endogeneity, we analyze the causal significance of each policy on epidemic control. Our analysis reveals that that policies related to vaccine delivery, debt relief, and the cancellation of public events are the most effective measures. These findings provide key insights for policymakers, highlighting the importance of focusing on specific, high-impact measures in managing public health crises.

Keywords: Causal inference; Change point detection; COVID-19; PSM-DID model; Public policy.

1 Introduction

The COVID-19 pandemic has profoundly impacted global life and economies since 2020, marking it as the most widespread and severe health crisis in the past century. Unlike the pandemics of the early 20th century, such as the influenza outbreak, the modern world faces new challenges due to increased global interconnectivity and changes in human lifestyles. These factors complicate the prevention and control of epidemics in our era. However, advancements in governance, biomedicine, and epidemiology have provided societies with more sophisticated tools to combat such crises. Countries worldwide have implemented a variety of public policies to curb the pandemic's spread, including lockdowns, medical interventions, and economic measures. The widespread rollout of vaccines, supported by biomedical advancements, has also become a crucial element in controlling the virus. Each nation's public health strategy is shaped by factors such as population density, age demographics, ethnic diversity, and healthcare infrastructure, and these strategies often evolve as the pandemic progresses. The simultaneous implementation of multiple policies further complicates the analysis of any single policy's effectiveness in controlling the epidemic.

Numerous studies have examined the impact of public policies on COVID-19 containment using a variety of statistical models. For instance, Page-Tan and Corbin (2021) employed propensity score matching to explore the relationship between state and local policies and COVID-19 mortality rates. Sun et al. (2022) analyzed the effects of intervention timeliness, stringency, and duration on cumulative infections using counterfactual estimates. Yang and Yang (2021) applied the difference-in-differences method to assess how firm-level volatility responded to the COVID-19 shock through the lens of economic policy uncertainty. In another study, Zhang et al. (2021) utilized structural equation modeling to provide scientifically grounded evidence for designing more effective COVID-19 policies in the transport and public health sectors. Dzator et al. (2022) employed a panel data model to examine the effects of government policy stringency and handwashing measures.

Other approaches include the use of negative binomial regression (Gaskin et al., 2021), latent factor models (Chen et al., 2022), and logistic regression (Liu et al., 2021) to study the impact

of policies on daily life. Linear regression (Sylvester, 2021) and time series models (Pelagatti and Maranzano, 2021) have been used to predict COVID-19 cases. Spatial modeling has also been employed; for example, Guliyev (2020) explored the spatial impact of various geographic factors on COVID-19, Jiang et al. (2021) provided insights on mobility restriction policies, and Zhang and Lin (2021) used a generalized linear model for variable selection. Additionally, James et al. (2021) developed an infectious disease model to investigate the effects of public policy interventions.

In this paper, we explore the impact of public policies on the progression of the COVID-19 pandemic from multiple perspectives. By employing the widely used PSM-DID model in causal inference, we account for differences in national conditions across countries and analyze the effects of nearly 20 public policies, including lockdown measures, healthcare interventions, and economic policies, on the pandemic's trajectory. Our goal is to identify which policies are effective and which may have limited impact. Overall, this study approaches the analysis of public policy effects on the pandemic through the lens of causal inference.

The rest of the paper is organized as follows. Section 2 provides a detailed description of the data used in this study, including the specific definitions of each public policy variable, and presents visualizations of changes in the epidemic reproduction rate and virus variants across different countries. In Section 3, we explain the application of the PSM-DID model to control for confounding factors between countries and assess the significance of each policy. Section 4 discusses the empirical results and their corresponding policy implications. Finally, Section 5 gives a brief conclusion of this study.

2 Data

In this section, we will give the sources, descriptive statistics and exploratory data analysis of the data used in the statistical modeling later.

2.1 Data source and variable descriptions

To obtain the policy response intensity of governments in the past two years to deal with COVID-19, we use a popular public dataset called the Oxford COVID-19 Government Response Tracker (OxCGRT). The OxCGRT (Hale et al., 2021) systematically collects information on several different common policy responses governments have taken, records these policies on a scale to reflect the extent of government action, and aggregates these scores into a suite of policy indices. Here, we choose 8 containment and closure policies (C), 2 economic policies (E), and 6 health system policies (H). Then, in order to measure the effect of policies, we use new confirmed cases of (7-day smoothed) per million people as the dependent variable, which is collected by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University. Arroyo-Marioli et al. (2021) gave the real-time estimate of the effective reproduction rate (R) of COVID-19, which is used to detect the change points in Section 2.3. In terms of country and date selection, we only retain the data of 38 European countries with not too small population and relatively complete data from March 15, 2020 to November 30, 2021. All the variable descriptions and basic statistics are shown in Table 1. For detailed codebook, see https: //github.com/OxCGRT/covid-policy-tracker. The GitHub repository, which includes the code and data for the empirical result reproduction of this paper, can be founded in https: //github.com/Wang-ZH-Stat/COVID-19.

Variable	Description	Mean	Sd.	Min.	Max.	#n
C1	Closings of schools and universities		0.87	0	3	23729
C2	Closings of workplaces	1.67	0.76	0	3	23732
C3	Canceling public events	1.52	0.61	0	2	23732
C4	Limits on gatherings	3.16	1.10	0	4	23732
C5	Closing of public transport	0.43	0.59	0	2	23717
C6	Requirements of staying at home	0.88	0.82	0	3	23707
C7	Restrictions on internal movement		0.84	0	2	23662
C8	Restrictions on international travel	2.61	0.87	0	4	23730
E1	If the government is providing direct cash pay-	1.55	0.66	0	2	23724
	ments to people who lose jobs					
E2	If the government is freezing financial obliga-	1.24	0.76	0	2	23717
	tions for households					
H1	Presence of public info campaigns	1.96	0.21	0	2	23680
H2	Policies on who has access to testing	2.33	0.69	0	3	23709
H3	Policies on contact tracing after a positive di-	1.52	0.62	0	2	23625
	agnosis					
H6	Policies on the use of facial coverings	2.26	1.11	0	4	23690
m H7	Policies for vaccine delivery	2.00	2.10	0	5	23709
H8	Policies for protecting elderly people	1.88	0.95	0	3	23693
GRI	Overall government response index (all indica-	58.27	11.91	6.77	89.69	23740
	tors)					
SI	Stringency index (all C indicators, plus H1)	55.18	16.67	8.33	100.0	23742
CHI	Containment and health index (all C and H	57.54	11.99	7.74	90.00	23741
	indicators)					
ESI	Economic support index (all E indicators)	63.46	27.26	0	100	23723
NCSM	New confirmed cases of (7-day smoothed) per	182.23	238.34	21.63	2078.77	23788
	1,000,000 people					
R	Estimate of the effective reproduction rate	1.08	0.31	0.09	3.69	23668

Table 1: Variable descriptions and basic statistics.

5

2.2 Exploratory data analysis

In this section, we will show the preliminary results of visualizations of the data. In Figure 1, we illustrate the correlation between policy responses and their respective trends over time. We can find that there are always strong positive correlations within containment and closure policies, economic policies and health system policies, while there might be negatively correlated between containment and closure policies and health system policies. For containment and closure policies (denoted as "C"), the policy intensity varies from top to bottom, which is determined by the degree of epidemic spread. Economic policies and health system policies.




Figure 1: The correlation between policy responses and their respective trends over time. The data in the right panel averages 38 countries.

As for the comparison at the country level, Figure 2 shows the new cases smoothed per million in different countries. Here we select the ten most populous countries, and daily data are centralized in these countries to clearly show the performance of these countries in epidemic containment. We can see that the UK, Romania and Spain played important roles in epidemic outbreak, while Netherlands and Poland performed relatively well.




Figure 2: The steam plot of new cases smoothed per million in different countries. The new cases data are centralized for each date.

Figure 3 gives the trend of proportion of different variants over time in six main countries. We can find that, before January 2021, the variants spread in these countries are different, including B.1.177, B.1.160 and B.1.258. From January 2021 to July 2021, the variant "alpha" originated from the UK accounted for a dominant proportion in all countries while since July 2021, the variant "delta" originated from India become the mainstream.






Figure 3: The trend of proportion of different variants over time in different countries. The data is collected from GISAID (Elbe and Buckland-Merrett, 2017).

2.3 Detection of epidemic outbreak point

In order to test the effect of policy response, we need to focus on a period of time before and after the epidemic outbreak. Here we consider the sequential and multiple change point detection for the effective reproduction rate. Since the reproduction rate data is obviously doesn't follow the normal distribution, we use the Mann-Whitney test statistic (Ross et al., 2011) to detect location shifts in a stream with a (possibly unknown) non-Gaussian distribution. We only retain the 6 change points of effective reproduction rate in the rising stage, and the results are shown in Figure 4. Then observing the position of these change points on the smoothed confirmed new cases curve, we find three main epidemic outbreak points, including "2020-09-14", "2021-02-12" and "2021-10-04". In the following sections, we mainly measure the effect the policy responses based on the epidemic date near these three points. Specifically, 30 days before and 30 days after the epidemic outbreak points will be studied.




Figure 4: The results of sequential and multiple change point detection using the Mann-Whitney test statistic. Around the change points, the smoothed confirmed new cases present a piecewise linear trend.

3 Methodology

In this section, we will introduce the most popular technique in empirical research to evaluate policy effects, which is propensity score matching-difference in differences (PSM-DID) model (Hirano et al., 2003). This technique can be decomposed into two separated procedures. In this section, we illustrate the PSM procedure and DID procedure respectively, using the data of policy response C3 (canceling public events) and smoothed new confirmed cases near October 4, 2021 as an example.

3.1 **PSM** procedure

Propensity score matching (PSM), which was introduced by Rosenbaum and Rubin (1983), is a statistical matching technique that attempts to reduce the bias due to confounding variables. The possibility of bias arises because a difference in the treatment outcome between treated and untreated groups may be caused by a factor that predicts treatment rather than the treatment itself. For each covariate, randomization implies that treatment-groups will be balanced on average. Unfortunately, for observational studies, the assignment of treatments to research subjects is typically not random. Matching attempts to reduce the treatment assignment bias, and mimic randomization, by creating a sample of units that received the treatment that is comparable on all observed covariates to a sample of units that did not receive the treatment.

The covariates used to match countries are shown in Table 2. They are some macro factors that may affect the policy-making of government when the epidemic outbreaks.

In our illustration example (policy response C3, epidemic outbreak point "2021-10-04"), 38 countries are divided into the treatment group and the control group according to whether the value of C3 at date "2021-10-04" is greater than 1, and we get 10 countries in the control group and 28 in the treatment group. We run a logistic regression and use the estimated probability of a given country coming from the treatment group as the propensity score. Then, we perform the optimal pair matching (Hansen and Klopfer, 2006). Each country in the control group has been matched to a similar country from the treatment group. The distribution of propensity scores and the differences of covariates between control group and treatment group before and after matching are shown in Figure 5.

Variable	Description	Source	
population	Population (latest available values)	United Nations	
population_density	Number of people divided by land area	World Bank	
aged_65_older	Share of the population that is 65 years	World Bank	
	and older		
gdp_per_capita	Gross domestic product at purchasing	World Bank	
	power parity		
cardiovasc_death_rate	Death rate from cardiovascular disease	Global Burden of Diseas	
	in 2017	Collaborative Network	
diabetes_prevalence	Diabetes prevalence ($\%$ of population	World Bank	
	aged 20 to 79) in 2017		
hospital_beds_per_thousand	Hospital beds per 1,000 people	OECD	
life_expectancy	Life expectancy at birth in 2019	United Nations	
human_development_index	A composite index measuring average	United Nations	
	achievement in three basic dimensions		
	of human development		

Table 2: Macro covariates used to match countries.




Figure 5: The distribution of propensity scores, see subplots (a) and (b), and the comparison of covariates balance before and after matching, see subplots (c) and (d).

3.2 DID procedure

Difference-in-difference (DID) evaluates the effect of a treatment on an outcome by comparing the average change over time in the outcome variable for the treatment group to the average change over time for the control group. In contrast to a time-series estimate of the treatment effect on units or a cross-section estimate of the treatment effect, DID uses panel data to measure the differences, between the treatment and control group, of the changes in the outcome variable that occur over time. DID is one of the four most commonly used methods for policy evaluation. The other three methods include matching, synthetic control and regression discontinuity, see Imbens and Wooldridge (2009) for details.

There are many assumptions related to the DID model, and one of the most important is the parallel trend assumption. If individuals in the treatment group did not receive a treatment, which is counterfactual, the trends of response variable from treatment group and control group should be parallel. When this assumption is violated, we need to use PSM to reduce the endogeneity of the model. We only need to include the matched units to run a DID model.

In our illustration example, there are 10 units in both groups. By Figure 4, the smoothed new confirmed cases seems to be piecewise linear before and after the outbreak epidemic point. For a given policy response, consider the piecewise linear regression

$$Y_{it} = \beta_0 + \beta_1 X_t + \beta_2 P_{it} + \beta_3 (X_t - X_{t_0}) D_t + \beta_4 (D_t - D_{t_0}) D_t P_{it} + \varepsilon_t,$$
(1)

where Y_{it} is the smoothed new confirmed cases of country *i*, X_t is the time variable taking the same values as *t* from 1 to 60, t_0 is the policy response time (in our case, $t_0 = 30$), P_{it} is a binary variable taking 0 if the country *i* is in the control group and taking 1 if in the treatment group, D_t is a dummy variable taking 0 when $1 \le t \le 30$ and taking 1 when $31 \le t \le 60$, and ε_t is the error term. Here, the negative significance of $\hat{\beta}_4$ can be regarded as the evidence of significant policy response effect. To get obtain more informative measurement, we define the containment ratio (CR) as

$$CR = \frac{\max(-\beta_4, 0)}{\beta_1 + \beta_3} \times 100\%,$$
(2)

which illustrates the proportion of the slope of smoothed new confirmed cases after the outbreak point with time decreases by carrying out one given policy. The larger the value of CR, the more effective the policy is. In our example, $\hat{\beta}_4 = -3.4422$ with *p*-value taking 0.0041, and $\widehat{CR} = 52.31\%$. Canceling public events has a significant containment effect on the spread of the epidemic. The fitted lines of the control group, the treatment group and the counterfactual treatment group are shown in Figure 6.




Figure 6: The fitted lines of the control group, the treatment group and the counterfactual treatment group.

4 Results

In this section, we give the evaluation the effectiveness of most policy responses by PSM-DID technique. We use the similar procedures as those in Section 3 on every policy response and three epidemic outbreak points. The results are shown in Table 3. Some policy response have been omitted, since we get three meaningless '/', implying that the intensity of this policy does not differ much among these European countries at the same date. We still divided 38 countries into control group and treatment group according to the intensity of policy response, and the division criteria are consistent at three time epidemic outbreak points.

Table 3: The estimations of PSM-DID effects and containment ratios. * means $0.01 \le p < 0.05$, ** means $0.001 \le p < 0.01$ and *** p < 0.001. In every cell of the table, the first row gives the estimation of β_4 in model (1), the second row gives the standard deviation, and the third row gives the estimation of containment ratio. '/' means that there are less than 3 countries in the control group or treatment group.

Policy	2020-09-14	2021-02-12	2021-10-04	Policy	2020-09-14	2021-02-12	2021-10-04
C1	0.1451	-0.4832	4.4789		-1.294***	-6.1438***	-3.4422**
	(0.2426)	(1.2224)	(1.6829)	C3	(0.2629)	(1.5246)	(1.1956)
	0%	11.07%	0%		35.89%	100.99%	52.31%
C4	-0.6398*		-0.3187			-2.6323**	-11.02**
	(0.2517)	/	(1.0109)	C6	/	(0.9514)	(1.5776)
	22.66%		3.97%			45.99%	93.11%
E1	1.7113***	0.2987	-0.0969		0.2083	-6.446***	-0.2798
	(0.2499)	(1.2112)	(1.0466)	E2	(0.2545)	(1.0998)	(1.1292)
	0%	0%	1.18%		0% 109.12%		3.44%
H2	-1.6034***	-3.0598**	0.2454			-2.3722*	-5.5568***
	(0.2928)	(1.1526)	(1.8826)	m H7	/	(1.1726)	(1.4866)
	38.99%	61.4%	0			48.6%	63.73%
H8	-0.3310	0.1783	-4.0268***				
	(0.2709)	(1.3543)	(1.1184)				
	10.75%	0%	36.5%				

For containment and closure policies, closings of schools and universities (C1) and limits on gatherings (C4) have poor effectiveness, and canceling public events (C3) and requirements of staying at home (C6) are significantly effect. One possible explanation can be that canceling public events and requirements of staying at home are more operable and supervised than limits on gatherings, and closings of schools and universities has little effect on containing the epidemic. For economic policies, debt and contract relief (E2) is more effective than income support (E1). Even E1 is significantly positive at "2020-09-14". One possible explanation can be that providing more cash support to idle people will encourage them to gather for fun, while debt and contract relief can ensure that people are not forced to go out to make a living.

For health system policies, testing policy (H2) is much effective in the first two epidemic outbreak points, and protection of elderly people (H8) becomes more important in recent days. Vaccination policy (H7) is always effect, which is reasonable. With the normalization of the epidemic, the attitudes of people towards testing has become more passive. Perhaps in the future, the government can pay more attention to the protection of elderly people and carry out more refined policies.

Based on the containment ratios, the four most effective policy responses are canceling public events, requirements of staying at home, testing policy and vaccination policy.

5 Conclusions

In this paper, we examine the impact of public policies on epidemic containment using causal inference techniques. The PSM-DID model, which mimics randomization, is employed to assess the effectiveness of various policy responses. Our analysis reveals that for containment measures, canceling public events and enforcing stay-at-home requirements are more effective than limiting gatherings. In terms of economic policies, debt relief proves to be more impactful than direct cash support. For health system policies, combining vaccination efforts with protections for the elderly can lead to more effective outcomes.

Despite the insights provided by the causal inference method, some limitations exist. First, the PSM-DID model assumes that units are uncorrelated, but there may be spatial dependencies among European countries that warrant consideration of spatial models. Second, since countries often implement multiple policies simultaneously, analyzing each policy in isolation may not fully capture their combined effects. Integrated policy indices, such as the GRI, could offer a more comprehensive alternative. Lastly, studying multiple countries and policies concurrently presents challenges due to numerous confounding factors. The synthetic control method proposed by Abadie and Gardeazabal (2003) could be a useful approach for evaluating the effects of multiple policies within a single country.

References

- Abadie, A. and Gardeazabal, J. (2003). The economic costs of conflict: A case study of the basque country. *American Economic Review*, 93(1):113–132.
- Arroyo-Marioli, F., Bullano, F., Kucinskas, S., and Rondón-Moreno, C. (2021). Tracking R of COVID-19: A new real-time estimation using the kalman filter. *PloS One*, 16(1):e0244474.
- Chen, C., Feng, T., and Gu, X. (2022). Role of latent factors and public policies in travel decisions under COVID-19 pandemic: Findings of a hybrid choice model. *Sustainable Cities and Society*, 78:103601.
- Dzator, J., Acheampong, A. O., Dzator, M., Paolucci, F., Yawe, B. L., Asmah, E. E., Andoh, F. K., Kabagenyi, A., and Gillespie, J. (2022). Policy stringency, handwashing and COVID-19 cases: Evidence from global dataset. *Health Policy and Technology*, 11(2):100574.
- Elbe, S. and Buckland-Merrett, G. (2017). Data, disease and diplomacy: GISAID's innovative contribution to global health. *Global Challenges*, 1(1):33–46.
- Gaskin, D. J., Zare, H., and Delarmente, B. A. (2021). Geographic disparities in COVID-19 infections and deaths: The role of transportation. *Transport Policy*, 102:35–46.
- Guliyev, H. (2020). Determining the spatial effects of COVID-19 using the spatial panel data model. *Spatial Statistics*, 38:100443.
- Hale, T., Angrist, N., Goldszmidt, R., Kira, B., Petherick, A., Phillips, T., Webster, S., Cameron-Blake, E., Hallas, L., Majumdar, S., et al. (2021). A global panel database of pandemic policies (Oxford COVID-19 government response tracker). *Nature Human Behaviour*, 5(4):529–538.
- Hansen, B. B. and Klopfer, S. O. (2006). Optimal full matching and related designs via network flows. Journal of Computational and Graphical Statistics, 15(3):609–627.
- Hirano, K., Imbens, G. W., and Ridder, G. (2003). Efficient estimation of average treatment effects using the estimated propensity score. *Econometrica*, 71(4):1161–1189.
- Imbens, G. W. and Wooldridge, J. M. (2009). Recent developments in the econometrics of program evaluation. Journal of Economic Literature, 47(1):5–86.
- James, L. P., Salomon, J. A., Buckee, C. O., and Menzies, N. A. (2021). The use and misuse of mathematical modeling for infectious disease policymaking: lessons for the COVID-19 pandemic. *Medical Decision Making*, 41(4):379–385.
- Jiang, P., Fu, X., Van Fan, Y., Klemeš, J. J., Chen, P., Ma, S., and Zhang, W. (2021). Spatial-temporal potential exposure risk analytics and urban sustainability impacts related to COVID-19 mitigation: A perspective from car mobility behaviour. *Journal of Cleaner Production*, 279:123673.

- Liu, R., Zhang, Y., Nicholas, S., Leng, A., Maitland, E., and Wang, J. (2021). COVID-19 vaccination willingness among chinese adults under the free vaccination policy. *Vaccines*, 9(3):292.
- Page-Tan, C. and Corbin, T. B. (2021). Protective policies for all? an analysis of Covid-19 deaths and protective policies among low-, medium-, and high-vulnerability groups. *Disas*ters, 45:S119–S145.
- Pelagatti, M. and Maranzano, P. (2021). Assessing the effectiveness of the italian risk-zones policy during the second wave of COVID-19. *Health Policy*, 125(9):1188–1199.
- Rosenbaum, P. R. and Rubin, D. B. (1983). The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70(1):41–55.
- Ross, G. J., Adams, N. M., Tasoulis, D. K., and Hand, D. J. (2011). A nonparametric change point model for streaming data. *Technometrics*, 53(4):379–389.
- Sun, J., Zheng, Y., Liang, W., Yang, Z., Zeng, Z., Li, T., Luo, J., Ng, M. T. A., He, J., and Zhong, N. (2022). Quantifying the effect of public activity intervention policies on COVID-19 pandemic containment using epidemiologic data from 145 countries. *Value in Health*, 25(5):699–708.
- Sylvester, S. M. (2021). COVID-19 and motivated reasoning: The influence of knowledge on covid-related policy and health behavior. Social Science Quarterly, 102(5):2341–2359.
- Yang, J. and Yang, C. (2021). Economic policy uncertainty, COVID-19 lockdown, and firm-level volatility: Evidence from China. *Pacific-Basin Finance Journal*, 68:101597.
- Zhang, J., Zhang, R., Ding, H., Li, S., Liu, R., Ma, S., Zhai, B., Kashima, S., and Hayashi, Y. (2021). Effects of transport-related COVID-19 policy measures: A case study of six developed countries. *Transport Policy*, 110:37–57.
- Zhang, T. and Lin, G. (2021). Generalized k-means in GLMs with applications to the outbreak of COVID-19 in the United States. *Computational Statistics & Data Analysis*, 159:107217.