STRANGE PARTICLE PRODUCTION IN NUCLEAR COLLISIONS AT CERN-NA49

D. BARNA for the NA49 Collaboration

Central Research Institute for Physics, Konkoly-Thege út 29-33, H-1121, Budapest, Hungary

In order to quantify isospin effects in the comparison of elementary to nuclear collisions, p-p and n-p interactions have been studied in the NA49 experiment in addition to p-Pb and Pb-Pb reactions. Together with first measurements of cascade hyperon and Ω production in p-p collisions, isospin-corrected K/π ratios and cascade yields are discussed. The $\overline{\Omega}^+/\Omega^-$ ratio in p-p collisions is found to be less than 0.5 with 95% confidence level.

1 Introduction

NA49 is a general purpose large acceptance hadron spectrometer[1]. It's main tracking devices are 4 large volume TPCs, which make particle identification possible. The experiment has collected data on a wide range of hadronic interactions (p-p, d-p, π -p, p-Pb, π -Pb, Pb-Pb). These data are used to make a systematic study and understand the complex A-A interactions.

2 Isospin Effects on the K and π Production

In order to trace the signatures of new physics, heavy ion collisions are usually compared to elementary (p-p or p-A) collisions. Beyond the possible 'interesting' differences there is a 'trivial' difference however: heavy ions are composed of neutrons and protons (for Pb the mixture is 60% n, 40% p). In order to study differences between proton and neutron fragmentation, NA49 has collected data with a deuteron beam incident upon a proton target with $p_{\text{beam}}=158~\text{GeV/nucleon}$. By tagging on a spectator proton with $p\approx 158~\text{GeV}$ one can select neutron-proton collisions.




Figure 1: Pion (left) and kaon (right) invariant p_T distributions in p-p and n-p collisions (preliminary)




Figure 2: (a): the $R^+ = K^+/\pi^+$ and $R^- = K^-/\pi^-$ ratio in pp and np collisions. (b) and (c): the R^+ and R^- ratio in pPb and PbPb collisions, compared to the same ratio in pp collisions. The estimated isospin-effect was removed from the PbPb results, see text for details (preliminary results)




Figure 3: Rapidity distributions of Λ , $\overline{\Lambda}$, Ξ and $\overline{\Xi}$ particles in p+p collisions

Charged kaon and pion yields were determined via dE/dx measurements. Figure 1 shows the transverse momentum distributions in p-p and n-p collisions in the projectile (n or p) hemisphere at fixed Feynman-x. The flip of positive and negative pions and the unchanged kaon yields, when replacing the proton projectile by a neutron projectile, are clearly visible. Consequently the $R^+ = K^+/\pi^+$ ratio is higher, the $R^- = K^-/\pi^-$ ratio is lower in n than in p fragmentation (fig. 2a). A properly weighted average (40%p + 60%n fragmentation) is used then to estimate the magnitude of isospin effects in Pb-Pb collisions. Figures 2 b.) and c.) show the $R_{\rm pPb,\ PbPb}^{\pm}/R_{\rm pp}^{\pm}$ double ratios as a function of Feynman-x, after the elimination of the isospin effect from the Pb-Pb results. Note that above $x_F = 0.05$ the p-Pb and Pb-Pb results agree within errors. The deviation close to $x_F = 0$ is due to the target feed-over: target nucleons in p-Pb and Pb-Pb behave differently. They suffer 1 (in p-Pb) or several collisions (in Pb-Pb).

3 Strange Baryon Production

3.1 Strangeness Enhancement

Strange and multistrange baryons are reconstructed via their weak decay, using two independent reconstruction methods. The acceptance extends over a wide rapidity range (± 1 unit around midrapidity), and covers the full p_T range. Rapidity distributions of Λ , $\overline{\Lambda}$, Ξ and $\overline{\Xi}$ are shown in figure 3. The Λ ($\overline{\Lambda}$) results have been feeddown-corrected with respect to the $\Xi^-, \Xi^0 \to \Lambda$ ($\overline{\Xi}^+, \overline{\Xi}^0 \to \overline{\Lambda}$) channel. For this the measured Ξ^- and $\overline{\Xi}^+$ yields were used. The Ξ^0 ($\overline{\Xi}^0$) yields were estimated as $\Xi^0 = 1.5 \cdot \Xi^-$ and $\overline{\Xi}^0 = 0.66 \cdot \overline{\Xi}^+$.

Strangeness enhancement in heavy ion collisions is often defined as the increase of the




Figure 4: Ξ^- (left panel) and Ξ^+ (right panel) enhancement: (a) and (c) – without isospin correction, (b) and (d) – with isospin correction. The Pb-Pb point is an average of WA97[2] and NA49[3] results.

strangeness/number-of-participants ratio compared to p-p collisions. Strangeness enhancement can be attributed to the presence of the Quark Gluon Plasma, or the higher number of collisions suffered by an average participant. Assuming the latter, p-A data can be used to make a study of enhancement as a function of the number of collisions ν .

Midrapidity yields are contributed to by both the target (T) and the projectile (P). In the symmetric p-p collisions both contributions are therefore $T_{pp} = P_{pp} = \frac{1}{2} \cdot \frac{dN}{dy} \Big|_{pp}^{y=0}$. In the asymmetric p-A collisions no enhancement per nucleon is assumed on the target side (since each target nucleon suffers only 1 collision). The target contribution (T_{pA}) is therefore assumed to be the target contribution derived from the p-p results (T_{pp}) , scaled by the number of colliding target nuclei (ν) . All the enhancement (E_{ν}) is attributed to the projectile, suffering ν collisions: $\frac{dN}{dy}\Big|_{pA}^{y=0} = \nu \cdot \frac{1}{2} \frac{dN}{dy}\Big|_{pp}^{y=0} + E_{\nu} \cdot \frac{1}{2} \frac{dN}{dy}\Big|_{pp}^{y=0}$. The enhancement in the symmetric A-A collisions is defined as $\frac{dN}{dy}\Big|_{AA}^{y=0} = E_{\nu} \cdot \frac{N_W}{2} \cdot \frac{dN}{dy}\Big|_{pp}^{y=0}$, where N_W is the number of wounded nucleons. Panels (a) and (c) of figure 4 show the enhancement E_{ν} for Ξ^- and Ξ^+ in p-Be, p-Pb and Pb-Pb collisions as a function of ν (which is the average number of collisions suffered by the projectile for p-A collisions, and by an average nucleon for A-A collisions). Enhancement factors for Ξ^- are larger than for Ξ^+ . The enhancement factors for both the baryon and the antibaryon are of the same order in p-Pb and Pb-Pb collisions.

Since Ξ production from a neutron or a proton can be different, isospin effects can also play a role in strangeness enhancement. In order to take this into account, we redefined strangeness enhancement, with this isospin effect detached:

$$\frac{dN}{dy}\Big|_{pA}^{y=0} = \nu \cdot \alpha \cdot \frac{1}{2} \cdot \frac{dN}{dy}\Big|_{pp}^{y=0} + E_{\nu} \cdot \frac{1}{2} \cdot \frac{dN}{dy}\Big|_{pp}^{y=0} \tag{1}$$

$$\frac{dN}{dy}\Big|_{AA}^{y=0} = E_{\nu} \cdot \frac{N_W}{2} \cdot \alpha \cdot \frac{dN}{dy}\Big|_{pp}^{y=0}$$
(2)

where the α factor accounts for the increase or decrease of the Ξ yield due to the proton-neutron composition of the lead nucleus. This could be derived by comparing an appropriately weighted average of Ξ yields from proton (40%) and neutron (60%) induced reactions to the Ξ yield in proton-induced reactions. Our limited statistics of n+p data did not allow a measurement of Ξ production, therefore the α factor for Ξ^- and $\overline{\Xi}^+$ could only be estimated: we used the factors measured[4] for proton and antiproton: $\alpha_{\Xi^-} = \alpha_{\overline{p}} = 1.3$ and $\alpha_{\overline{\Xi}^+} = \alpha_p = 0.7$. The enhancements as defined in equations 1 and 2 are shown in the panels (b) and (d) of figure 4 for Ξ^- and $\overline{\Xi}^+$,




Figure 5: Omega mass spectra in p-p and Pb-Pb collisions

respectively. After taking into account the isospin effects the enhancement factors for Ξ^- and $\overline{\Xi}^+$ approached each other, and p-Pb and Pb-Pb results are still of the same order.

3.2 Ω Production in p-p Collisions

 Ω^- signals have been observed in p-p collisions for the first time. Panels (a) and (b) of figure 5 show the Ω^- and $\overline{\Omega}^+$ mass spectra in p-p collisions; panel (c) shows the $\Omega^- + \overline{\Omega}^+$ mass spectrum in Pb-Pb collisions. From the absence of the $\overline{\Omega}^+$ signal in p-p events we estimate the $\overline{\Omega}^+/\Omega^-$ ratio to be less than 0.5 with a confidence level of 95%.

4 Conclusions

Results obtained from p-p and n-p collisions indicate a sizeable isospin-dependence of the K/π ratio. Using these data to eliminate isospin effects from Pb-Pb results the K/π ratio in Pb-Pb and p-Pb collisions in the forward hemisphere agree within errors.

The final Ξ and feeddown-corrected Λ results in p-p and Pb-Pb collisions at the NA49 experiment are obtained. The Ξ -enhancements at midrapidity in p-A and A-A collisions with respect to p-p collisions are of the same order. An increased statistics should make the measurement of Λ and Ξ particles in n-p collisions possible in the near future. In the present study the isospin effect on Ξ production in p-Pb and Pb-Pb collisions was estimated using the factors determined for protons and antiprotons from p-p and n-p collisions. Isospin-corrected enhancement factors of Ξ^- and $\overline{\Xi}^+$ approach each other.

An Ω^- signal has been observed in p-p collisions for the first time. The $\overline{\Omega}^+/\Omega^-$ ratio is estimated to be less than 0.5 with a confidence level of 95%. Ω production has also been observed in Pb-Pb collisions in NA49.

Acknowledgements

This work was partially supported by the Hungarian Scientific Research Fund (OTKA) under the contracts F034707 and T0032293.

References

- 1. S. V. Afanasiev et al, Nucl. Instrum. Methods A430, 210 (1999)
- 2. F. Antinori et al, Nucl. Phys. A661, 130c (1999)
- 3. S.V. Afanasiev et al., hep-ex/0202037
- 4. S. V. Afanasev et al., "Progress report and beam request for 2002", CERN-SPSC-2002-008