

Soft QCD Results from ATLAS and CMS

Claire Gwenlan*, on behalf of the ATLAS and CMS Collaborations

University of Oxford

E-mail: c.gwenlan1@physics.ox.ac.uk

The ATLAS and CMS Collaborations at the LHC have measured a wide range of properties of soft particle production. These include charged particle differential distributions, forward-backward and azimuthal correlations, azimuthal ordering of hadrons, event shapes and identified particle production. Many of the measurements can be used to develop and tune Monte Carlo models. The inclusive and diffractive cross sections have been measured using vertex counting, particle tags, and rapidity gaps. In addition, the mean transverse energy flow and its density is measured out to forward rapidities providing a sensitive probe of models of the underlying event and QCD radiation.

*The European Physical Society Conference on High Energy Physics -EPS-HEP2013
18-24 July 2013
Stockholm, Sweden*

*Speaker.



1. Introduction

Soft QCD physics (low momentum transfer, strong force interactions) dominates the total cross section at hadron colliders. Since its description involves non-perturbative QCD it is not well understood, and we must rely on phenomenological models. Improving our understanding and description of such physics is not only of interest in its own right, but is also needed for improved measurements of high-momentum transfer observables, since soft QCD processes contribute to the overall final state, influencing, for example, measurements of inclusive jet cross sections, missing transverse momentum, and isolation cuts. Measurements of total cross sections, particle yields and kinematic distributions are essential for understanding the physics of hadron production, including the relative roles of soft and hard contributions, and for testing, and tuning, our existing Monte Carlo (MC) models, in order to provide the best description of data. These proceedings provide an overview of selected results, from the wealth of soft QCD measurements now available from the ATLAS [1] and CMS [2] Collaborations at the Large Hadron Collider (LHC).

2. Results

2.1 Inclusive Cross Section Measurements

The total hadronic cross section can be separated into elastic and inelastic components. The LHC experiments are sensitive to a large fraction of the inelastic cross section, where at least one proton loses a fraction of its energy. The inelastic cross section can, itself, be separated into diffractive and non-diffractive processes. Diffractive events are characterised by a large rapidity “gap” (i.e. absence of activity) due to the exchange of a colour singlet object. Diffractive processes have contributions from single diffractive ($pp \rightarrow pX$), double diffractive ($pp \rightarrow XY$) and (a smaller proportion of) central diffractive ($pp \rightarrow pXp$) events.

ATLAS and CMS have both measured the total inelastic cross section [3, 4], making minimal experimental requirements to ensure an inelastic collision has occurred. For ATLAS, two hits in the Minimum Bias Trigger Scintillators (MBTS) are required while, for CMS, events in which ≥ 5 GeV is deposited in the forward hadronic calorimeter, are accepted. The measurements are restricted to the kinematic region $\xi = (M_X^2)/s > 5 \times 10^{-6}$ (where M_X is the (highest) invariant mass of particles, selected on one side of the largest rapidity gap in the event), in order to maintain a high efficiency for the minimum bias triggers. The measurements from ATLAS and CMS agree well with each other, as shown in Fig. 1 (black triangle and red circle).




Figure 1: ATLAS and CMS measurements of the inelastic cross section compared to predictions from several MC models, as labelled. Plot from [4].




Figure 2: SD (left) and DD (right) cross sections as a function of $\xi = M_X^2/s$ compared to PYTHIA6, PYTHIA8-4C and PYTHIA8-MBR predictions. Plots from [7].

41 CMS have also used an alternative technique, counting the number of pileup vertices in an
 42 event [4]. This method relies on the assumption that the number of inelastic pp interactions (n), in
 43 a given bunch crossing, follows a Poisson distribution, $P(n) = \frac{(L\sigma_{\text{inel}})^n}{n!} e^{-L\sigma_{\text{inel}}}$. The data is plotted
 44 in bins of instantaneous luminosity, L , for each measured value of n , and fitted with the Poisson
 45 distribution to extract values for the total inelastic cross section, σ_{inel} . The final results are obtained
 46 by averaging the values extracted from each luminosity bin, and are presented as a function of the
 47 minimum number of tracks used to build the vertices (1, 2 or 3), as shown in Fig. 1 (red squares).

48 The measurements are compared to a variety of MC models: PYTHIA6 and 8 (which give very
 49 similar predictions, independent of tune); PYTHIA8-MBR, with an alternative model of diffrac-
 50 tion; and a second class of models based on the same Regge-Gribov phenomenology, but with
 51 different implementations of the model ingredients: PHOJET, and three MC models commonly
 52 used in cosmic ray physics (EPOS, QGSJET and SIBYLL). The ATLAS and CMS cross section
 53 measurements are generally well described by QGSJET01, QGSJET II-04 and PYTHIA8-MBR.
 54 Other models tested tend to overestimate the data. Several models, such as PYTHIA6 and 8, are
 55 able to reproduce the TOTEM measurement [5] of $\sigma_{\text{inel}}^{\text{total}}$, suggesting these models overestimate the
 56 contribution from high mass diffraction and underestimate the low mass component.

57 ATLAS and CMS have also performed measurements of diffractive cross sections [6, 7]. Both
 58 collaborations have measured the total inelastic cross section as a function of $\Delta\eta_F$ (the largest empty
 59 pseudorapidity interval, measured from the edge of the detector). The results show the dominance
 60 of non-diffractive events at small $\Delta\eta_F$, and the diffractive plateau at large values of $\Delta\eta_F$. Varying
 61 the transverse momentum (p_T) threshold requirements on the particles entering the measurement,
 62 shows that for high p_T , non-diffractive events dominate even out to large values of $\Delta\eta_F$ showing that
 63 inclusive events are dominated by low- p_T production. PYTHIA8 tune 4C and the PYTHIA8-MBR
 64 model give a reasonable description of the data.

65 CMS have further discriminated between single- (SD) and double-diffractive (DD) events [7]
 66 based on topology. Using the CASTOR very forward calorimeter (which covers the pseudorapidity
 67 region $\eta = -6.6 - -5.2$), SD- and DD-enhanced events have been selected. Activity was required
 68 in the central detector and, in addition, activity in CASTOR was either vetoed or required to select
 69 SD- and DD-enhanced events, respectively. The results show that PYTHIA8-MBR [8], a Regge-
 70 based model with hadronisation tuned to describe diffractive masses at low energies, and including




Figure 3: Inclusive charged-particle jet p_T spectrum at high multiplicity ($110 < N_{ch} < 140$), compared to various MC models. Plot from [10].




Figure 4: E_T density compared to various MC models and tunes for the inclusive selection. The filled band represents the total uncertainty on the unfolded data. Plot from [11].

an implementation of the central diffractive process, gives the best description of the data. In particular, the SD-enhanced distribution (Fig 2, left) shows a different trend between the standard PYTHIA (Schuler-Sjöstrand) model of diffraction, and PYTHIA8-MBR. DD cross sections were also measured using only the central detector, by requiring a central rapidity gap, $\Delta\eta > 3$.

2.2 Particle Production

ATLAS and CMS have performed a wealth of measurements of particle production, including charged particle differential distributions, transverse energy flow, event shapes, particle correlations, and identified particle production. Comparison of such measurements with predictions can highlight areas of discrepancy, and the data can be used to further our understanding and improve the description by the models.

CMS have measured inclusive charged particle production [9] for two different topologies: inclusive and non-single-diffractive (NSD) events. The TOTEM T2 telescopes (triple Gas Electron Multipliers, covering $|\eta| = 5.3 - 6.5$, $p_T > 40$ MeV), were used to select the events. For the inclusive sample, tracks in T2 were required in either the forward or backward hemispheres, while for the NSD-enhanced sample, tracks were required in both hemispheres. Measurements were performed in two kinematic regions ($p_T > 100$ MeV, $p_T > 1$ GeV) in order to study both soft and hard scattering. The data were compared to a variety of models and tunes: PYTHIA6 Z2*, PYTHIA8 4C, Herwig++ EE3C, EPOS and QGSJET II-04. The level of agreement with the data varies with both p_T and topology. For the inclusive sample, the PYTHIA6 Z2* tune provides the best description of the data, for both kinematic regions. Most other models overestimate the data for the inclusive topology, though Herwig++ underestimates the measurement for $p_T > 1$ GeV. For the NSD-enhanced data, no model tested gives a good description of the data. Discrepancies are observed for both topologies, and both kinematic regions, of up to 15%.

CMS have also measured the properties of soft and hard particles as a function of multiplicity [10]. Charged particle tracks were classified as either lying inside (intra-jet) or outside jets, and various properties, such as multiplicity and p_T distributions of the charged particles, jet rates and




Figure 5: Normalised distribution of the transverse sphericity (left); mean value of the transverse sphericity as a function of $\sum p_{\text{T}}$ (right). Plots from [12].

97 jet shapes were measured. Deficiencies in the description of the data were observed, especially at
 98 high multiplicities. In particular, the PYTHIA6 Z2* and PYTHIA8 4C models predict too many
 99 jets, and the jets are too hard. Herwig++ shows the opposite trend, producing too few jets, that are
 100 too soft (see, for example, Fig. 3). The MCs bracket the data, perhaps providing an indication of
 101 where the models could be improved.

102 ATLAS have performed a measurement of the transverse energy flow, and its density, using
 103 both charged and neutral particles [11]. The measurement uses the full acceptance of the ATLAS
 104 calorimeters, and has been performed for both inclusive (minimum bias) events and in dijet events
 105 in the region transverse to the highest- p_{T} jet ($60^\circ < |\phi| < 120^\circ$), where the distributions are sensitive
 106 to the underlying event. The results show that in the central region the data is reasonably well
 107 described by recent MC tunes, while in the forward region the models underestimate the data (see,
 108 for example, Fig 4, showing the E_{T} density for the inclusive selection). Note that the PYTHIA6
 109 DW tune uses only pre-LHC data, and Herwig++ UE7-2 is optimised for the description of the
 110 underlying event, so the discrepancies observed with these models also in the central region is not
 111 necessarily surprising. Varying the proportion of diffractive events has little effect on the shape
 112 of the distributions. Changing the proton PDFs from the default for the analysis, CTEQ6L1, to
 113 MSTW08LO (which has a larger gluon density at high- and low- x) provides a better description of
 114 the energy flow in the forward region. However, the prediction remains $\sim 15\%$ below the data at
 115 very forward pseudorapidities.

116 ATLAS have also performed measurements of charged particle event shapes, which were up-
 117 dated for this conference [12]. Event shape variables, such as transverse thrust and transverse
 118 sphericity, describe the geometric properties of energy flow in the final state. The measurements
 119 show that in inclusive events, which are dominated by low- p_{T} production, there is a prevalence of
 120 spherical (isotropic) events (see e.g. Fig 5, left). As the p_{T} of the leading (highest- p_{T}) particle is
 121 increased the events become more jet-like (i.e. the peak of the distribution shifts towards higher
 122 values of transverse thrust and lower values of transverse sphericity). The mean value of the event
 123 shape variables show an increase as a function of $\sum p_{\text{T}}$, up to a maximum, and then fall off at very
 124 high $\sum p_{\text{T}}$ (Fig 5, right). The distributions are generally best described by the PYTHIA6 Z2* tune.
 125 However, the mean values of the event shape variables, as a function of $\sum p_{\text{T}}$, show that all models
 126 underestimate the data at large $\sum p_{\text{T}}$.




Figure 6: Forward-backward multiplicity correlations in symmetrically opposite η intervals, compared to a selection of MC models. Plot from [13].




Figure 7: Power spectrum $S_\eta(\xi)$, an observable sensitive to the azimuthal ordering of hadrons, compared to various MC models using conventional hadronisation algorithms. Plot from [14].

127 ATLAS have made a number of recent measurements of charged particle correlations. For
 128 example, the so-called forward-backward correlations [13] in which the multiplicity and $\sum p_T$ cor-
 129 relations are measured in symmetrically opposite η regions. The correlation observable is the
 130 normalised covariance between the two distributions relative to the mean¹. Measuring such
 131 correlations as a function of η separation, and particle- p_T , probes both hard and soft contributions,
 132 characterised by short-range and long-range correlations, respectively. The results show that the
 133 measured correlations can be adequately described by recent MC tunes, which include other LHC
 134 data. In particular, the PYTHIA6 AMBT2B tune, which includes this same dataset, though uses
 135 different observables, gives a good description of the data. Other recent measurements include
 136 charged particle azimuthal distributions [13], azimuthal ordering of hadrons [14] and two-particle
 137 correlations [15]. Results show that current MCs have difficulty in describing all features of the
 138 available correlation data, tending to over- or underestimate the strength of the correlations. One
 139 particular example, is the power spectrum $S_\eta(\xi)$ (an observable sensitive to the azimuthal ordering
 140 of hadrons), as shown in Fig. 7, demonstrating the poor description by current models. It has been
 141 suggested that alternative hadronisation models might be required in order to describe these data.

142 Both ATLAS and CMS have produced recent measurements of identified particle produc-
 143 tion [16, 17]. ATLAS have measured the spectra of kaons and lambdas, and CMS have studied
 144 pion, kaon and proton production. Many features are well described by recent MC tunes. For
 145 example, Fig. 8 (left) shows the p_T spectra of pions, kaons and protons as a function of p_T , demon-
 146 strating that the PYTHIA6 Z2* tune describes the data reasonably well except, perhaps, for protons
 147 at low- p_T . However, inadequacies in the description of certain features are observed, especially in
 148 strange particle production. This is apparent in the kaon-to-pion ratios as a function of p_T (as
 149 shown in Fig 8 (right)), and in lambda distributions at high- p_T . Discrepancies are also observed
 150 in the description of proton $\langle p_T \rangle$ at high multiplicities, and as a function of \sqrt{s} , where current
 151 models tend to underestimate the data.

¹For example, the forward-backward multiplicity correlation is defined as: $\rho_{FB}^n = \frac{\sum x_{F(B)}^n}{N \sigma_F^n \sigma_B^n}$ where $x_{F(B)}$ indicates the deviation of forward (backward) multiplicities from their mean, $\sigma_{F(B)}$ is the standard deviation of forward (backward) distributions about their mean, and the sum is over N events.




Figure 8: Transverse momentum distributions of identified (positively) charged pions, kaons and protons (left). Ratio of particle yields as a function of transverse momentum (right). Plots from [17]

152 3. Summary

153 The LHC provides a rich laboratory for studying soft QCD. ATLAS and CMS have provided a
 154 wealth of data on event characteristics, particle properties and correlations at the hadron level. Mea-
 155 surements of inelastic cross sections have been made by the experiments using different methods.
 156 While many aspects of particle production are adequately described deficiencies are still apparent -
 157 most especially at high multiplicities, in the forward region, in certain particle correlation measure-
 158 ments, and in strange particle production. The measurements presented here can be used to further
 159 our understanding and improve MC models and tuning in the future.

160 References

- 161 [1] ATLAS Coll., 2008 JINST 3 S08003.
 162 [2] CMS Coll., 2008 JINST 3 S08004.
 163 [3] ATLAS Coll., Nature Commun. 2 (2011) 463.
 164 [4] CMS Coll., Phys. Lett. B 722 (2013) 5.
 165 [5] TOTEM Coll., Eurphys. Lett. 96 (2011) 21002.
 166 [6] ATLAS Coll., Eur. Phys. J. C72 (2012) 1926.
 167 [7] CMS Coll., CMS PAS FSQ-12-005, <https://cds.cern.ch/record/1547898>.
 168 [8] R. Ciesielski and K. Goulianos, arXiv:1205.1446.
 169 [9] CMS Coll., CMS PAS FSQ-12-026, <https://cds.cern.ch/record/1546365>.
 170 [10] CMS Coll., CMS PAS FSQ-12-022, <https://cds.cern.ch/record/1543202>.
 171 [11] ATLAS Coll., JHEP 11 (2012) 033.
 172 [12] ATLAS Coll., Phys. Rev. D88 (2013) 032004.
 173 [13] ATLAS Coll., JHEP 1207 (2012) 019.
 174 [14] ATLAS Coll., Phys. Rev. D86 (2012) 052005.
 175 [15] ATLAS Coll., JHEP 1205 (2012) 157.
 176 [16] ATLAS Coll., Phys. Rev. D85 (2012) 012001.
 177 [17] CMS Coll., Eur. Phys. J. C72 (2012) 2164.